Species-specific variation in mitochondrial genome tandem repeat

polymorphisms in hares (*Lepus* spp., Lagomorpha, Leporidae)

provides insight into their evolution

Riikka Tapanainen[1], Koit Aasumets[1], Zsófia Fekete[1,2], Steffi Goffart[1], Eric Dufour[3¶], &

Jaakko L. O. Pohjoismäki[1*¶]

[1]University of Eastern Finland, Department of Environmental and Biological Sciences –

Joensuu, Finland

[2]Hungarian University of Agriculture and Life Sciences, Institute of Genetics and

Biotechnology, Gödöllő, Hungary

[3]Mitochondrial Bioenergetics and Metabolism, Faculty of Medicine and Health Technology,

FI-33014 Tampere University, Finland

*Corresponding author

¶These authors contributed equally

Short title: Species-specific variation of mtDNA tandem repeats in hares

Abstract

The non-coding regions of some vertebrate mitochondrial genomes contain tandem repeat

elements, whose copy number can vary between species, populations and even individuals.

Both short (~20 bp) and long (130–160 bp) tandem repeats of variable lengths are present in

the mitochondrial genomes of hares, rabbits and pika (Lagomorpha), while none can be found

in related mammalian orders. In the presented study, we provide functional annotation for

mountain hare (*Lepus timidus*) and brown hare (*L. europaeus*) mitochondrial DNA (mtDNAs)

non-coding regions, together with a species as well as population-level analysis of tandem

repeat variation. While the mountain hare short tandem repeats (SRs) consist of two randomly

repeating but highly conserved 10 bp sequence motifs, brown hares have only one type of 10

bp motif, which shows more sequence variation within the species. The one repeat motif SRs

seem unique to brown hares, as the other members of the genus *Lepus* have two motifs.

Similarly, the long tandem repeats (LRs) show both sequence and copy number differences

between the species. Mountain hares have between four and seven repeats, with median value

of five and the brown hares between five and nine, with median value of six repeats.

Interestingly, mountain hare mtDNA introgressed into brown hares through hybridization

exhibit an intermediate LR copy number distribution. In cultured cells, the transfer of mountain

hare mtDNA into mtDNA-less brown hare cells typically result in a gain or loss of one to two

LR copies, the new LR haplotype being maintained in heteroplasmy with the old. The

reciprocal transfer of brown hare mtDNA had no effect on the LRs. We propose that the tandem

repeats are unlikely to have regulatory roles in hare mitochondrial genomes, but rather

represent selfish genetic elements, whose maintenance is influenced by the cellular context.

Introduction

Mitochondria are essential cellular organelles best known for production of ATP through

process called oxidative phosphorylation (OXPHOS) [1]. They are also a central hub for most

metabolic pathways and play specialized functions such as the biosynthesis of iron-sulfur

clusters, steroids, fatty acid oxidation, calcium homeostasis, programmed cell death, and more.

Due to their origin as a free-living prokaryote, mitochondria have their own small circular DNA

genome (mitochondrial DNA, mtDNA), which has undergone reductive evolution associated

with gene transfer to the nucleus [2]. In animals, mtDNA is typically less than 20 kb in size

and contains genes for 13 polypeptides of the OXPHOS complexes as well as two

mitochondrial ribosomal RNAs and 22 tRNAs required for their translation [3]. The two

strands of the mtDNA have very different gene density, with most of the genes being encoded

on the so-called heavy strand (H-strand). The designation stems from its high guanine content,

giving it a higher mass compared to the complementary light-strand (L-strand), as visualized

in alkaline CsCl density gradient centrifugation [4]. Besides the highly compact coding region,

the mitochondrial genome also has a non-coding region (NCR), sometimes called the control

region due to the presence of control elements required for DNA transcription and replication.

The length and organization of the NCR is very variable across the animal kingdom,

contributing most to the mtDNA length variation between the species [3].

In vertebrates the mtDNA non-coding region houses curious features, such as the

displacement loop (D-loop), which is generated by replication initiating from the main

replication origin (OriH: Origin of Heavy strand replication) and terminating prematurely at

the "termination associated sequence" (TAS) [5]. As a result, a triple-stranded DNA structure

is formed, including the 600–700 nt long nascent H-strand fragment, the 7S DNA, which

remains hybridized to its L-strand template. While the human and mouse mitochondrial

genomes, probably the best known mtDNAs, show considerable economy in their NCR

organization, its size can be variable in other vertebrates due to tandem repeat arrays [6]. Of

these, short tandem repeats of around 10 bp are an universal feature for the insect mitochondrial

genomes [7], also present in vertebrate species, including mammals [8]. Longer, 100-200 bp

repeats are found in certain species of fish [9], birds [10-12] and some mammals, such as rabbits

and hares [13]. The repeat number can vary from two [9] to more than one hundred [14] per

mtDNA molecule. The presence of mitochondrial tandem repeats is likely overlooked as they

can be easily lost during the sequencing and the assembly of short sequence reads, such as the

ones obtained from Illumina sequencing platforms.

The tandem repeats on the NCR of the European rabbit (*Oryctolagus cuniculus*)

mitochondrial genome were noted already in the 1980s [15] and later work showed that these

repeats are widespread also in other mitochondrial genomes of Lagomorpha [13]. There are

two types of repeated motifs in rabbits, 20 bp short repeats (SR) and 153 bp long repeats (LR).

The SR arrays are expanded or contracted relatively dynamically, with the copy number of the

units varying between three to 19. Consequently, the SR arrays can occur in heteroplasmy, i.e.

mitochondrial DNAs with different repeat lengths coexisting in the same cell or tissue. An

elegant experiment where rabbit SRs were cloned and maintained in a bacterial plasmid

demonstrated that slipped-strand mispairing during replication is the main mechanism to

explain the dynamic state of the repeat array [16]. The experiments demonstrated that SR

insertions are more common than deletions in arrays with less than 10 copies with the opposite

being true for longer tracts, resulting the SR lengths to oscillate around an optimal value.

Similar to the SRs, the LRs also have a high mutation rate (10^{-2} per animal per

generation), resulting in NCR length variation between generations but also within individuals,

manifesting as heteroplasmy as well as mosaicism [13]. In rabbits, LR tend to be present as

arrays of five repeats with shorter and longer variants being rarer. While the mtDNA haplotype

has no influence of the array length, oddly, some organs such as gonads maintain longer arrays

than other tissues [13]. Due to their length, the expansion or deletion of LRs through replication

slippage-mispairing is more complicated than for the SRs. In the nucleus, the copy number

changes of similar long satellite repeats, including expansions and contractions of rDNA arrays

[17], have been proposed to occur through break induced recombination [18]. Interestingly, a

similar copy-choice recombination mechanism involving strand invasion of the parental

molecule by a free 3′-end of a newly synthetized DNA strand at regions of short sequence

homologies has been suggested as a mechanism for mtDNA deletion formation in humans [19,

20]. Non-allelic recombination or end joining could in principle also explain the observed LR

length variation.

LR tandem repeats have been found in all Lagomorphs from pikas (*Ochotona* spp.) to

hares (*Lepus* spp., *Sylvilagus* spp.) [13]. All LRs have a 20 bp strictly conserved sequence,

which has been proposed to be a binding site for a regulatory factor, such as TFAM. In fact, the

rabbit LR has been suggested to contain both L-strand (LSP) and H-strand (HSP) promoters

for mtDNA transcription [21]. If this is the case, variation in LR length could influence primer

synthesis by MTRPOL at these promoters [22], thereby modulating also the replication

initiation.

Our group has been interested in hybridization between the mountain hare (*Lepus*

timidus) and brown hare (*L. europaeus*) at the northern contact zone of these species in Finland.

Despite being separated by three million years of evolution [23], the hybrids are fertile,

resulting in gene flow across the species barrier [24, 25]. This gene flow is biased towards the

brown hare, which seems to benefit from the hybridization by obtaining locally adapted alleles

from the mountain hare [26]. Furthermore, the hybridization results in frequent introgression

of mountain hare mtDNA into the brown hare population, encompassing up to fifth of the

individuals in certain regions in Finland [25]. When sequencing the mitochondrial genomes of

our mountain hare and brown hare cell lines [27], we recently noted a species difference in the

length of LRs in the mtDNA non-coding region. A population sampling of 151 mountain hares

and 148 brown hares with conspecific mtDNA confirmed that brown hares maintain more LR

copies (median = 6) than mountain hares (median = 5), with no correlation with their

geographic origin. Interestingly when introgressed into brown hares, the mountain hare

mtDNA presented frequent heteroplasmy and gained "brown-hare like" LRs (median = 6

copies). To further test how the nuclear background can influence LRs, we generated cybrids

of mountain hare and brown hare fibroblasts with nuclear DNA from one species and mtDNA

from another. The repeat region length changed in three cybrid cell lines out of six, regardless

of their genetic background, with the new LR variant maintained in heteroplasmy. In one case,

the LR number of brown hare mtDNA was increased from six to seven in cells with mountain

hare nucleus, a state which is very rare in the natural population of mountain hares. We

conclude that while the nuclear genetic background certainly plays a role in the LR

maintenance, environmental factors and differences in cell biology between species also

contribute to the observed variation. We also discuss the potential relation of the NCR repeat

elements to the regulation of mtDNA.

Materials and Methods

Sampling and DNA isolation

The mountain hare (*Lepus timidus*) and brown hare (*L. europaeus*) specimens presented here

are from a larger biobank of 1,202 hare samples collected in 2012–2016 across Finland [24,

25]. The origin and generation of the four mountain hare (LT1, LT4, LT5, LT6) and four brown

hare (LE1, LE2, LE3, LE4) fibroblast cell lines has been described elsewhere [27]. Most

samples are from hunted animals, with a subset from specimens found dead from the wild. The

sampling had minimal impact on the local populations and pose no threat on the habitats. The

hunting followed the regional hunting seasons and legislation (Metsästyslaki [Hunting law]

1993/615/5§). The sampling adhered to the ARRIVE guidelines and also no ethical assessment

is required. All sampling occurred in Finland and did not involve International Trade in

Endangered Species of Wild Fauna and Flora (CITES) or other export of specimens, as defined

by the Convention on Biological Diversity (CBD). The DNA was isolated from frozen ear

muscle biopsies using the pegGOLD blood & tissue DNA mini kit (VWR) and following the

protocol provided by the manufacturer.

Mitochondrial DNA sequencing

The primers used to amplify the hare mtDNA were as follows:

Le93F: TTGTTTTGTAGCAAGTTTACACATGC

Le184R: GCTTAATACCTGCTCCTCTTGATCTA

Le1580F: TTAAACCCATAGTTGGCCTAAAAGC

Le1635R: TTGAGCTTTAACGCTTTCTTAATTGA

Le3045F: AGGCGTATTATTTATCCTAGCAACCT

Le3175R: CCTCATAAGAAATGGTCTGTGCGA

Le3921F: CCCCCTAATCTTTTCCATCATCCTAT

Le4482R: TCATCCTATATGGGCAATTGAGGAAT

Le4689F: AGGCTTTATTCCAAAGTGAATTATTATTCA

Le5417R: AGGCTCCAAATAAAAGGTAGAGAGTT

Le6696F: ATACCGTCTCATCAATAGGCTCCTTC

Le6756R: ATAAAGATTATTACTATTACAGCGGTTAGA

Le8603F: AGCCTATATCTACATGATAATACTTAATGA

Le8698R: CGGATAAGGCCCCGGTAAGTGG

Le10552F: TTGAAGCAACACTAATCCCTACACTA

Le10613R: TCGTTCTGTTTGATTACCTCATCGT

Le11301F: ACCATTAACCTTCTAGGAGAGCTTCT

Le11807R: AGGATAATGATTGAGACGGCTATTGA

Le12407F: GTCTAATCCTAGCTGCTACAGGTAAG

Le12791R: GAGCATAAAAGAGTATAGCTTTGAA

Le14204F: ATTGTTAACCACTCTCTAATCGACCT

Le14514R: CCAATGTTTCAGGTTTCTAGGTAAGT

Lt16056F: TGGGGTATGCTTGGACTCAAC

Le16119R: TCGTCTACAATAAGTGCACCGG

In total, 12 separate reactions were prepared to cover the mitochondria genome:

1.   Lt16056F + Le184R: 1871 bp

2.   Le93F + Le1635R: 1543 bp

3.   Le1580F + Le3175R: 1596 bp

4.   Le3045F + Le4482R: 1438 bp

5.   Le3921F + Le5417R: 1497 bp

6.   Le4689F + Le6756R: 2068 bp

7.   Le6696F + Le8698R: 2003 bp

8.   Le8603F + Le10613R: 2011 bp

9.   Le10552F + Le11807R: 1256 bp

10.   Le11301F + Le12791R: 1491 bp

11.   Le12407F + Le14514R: 2108 bp

12.   Le14204F + Le16119R: 1916 bp

(Expected fragment size based on the published *Lepus europaeus* mtDNA sequence from

Sweden [NC_004028.1]).

The fragments were amplified from total DNA preparations using a PCR program with a 1 min

94 °C denaturing step, followed by 35 cycles of 94 °C for 15 s, 56 °C for 15 s and 72 °C for 2

min and a final 3 min elongation step at 72 °C. The obtained products were gel purified using

the GeneJET gel extraction kit (Thermo Fisher Scientific™) and sent for sequencing using

Illumina MiniSeq™ at the Genome Center of Eastern Finland.

The sequence of the non-coding region-containing PCR fragment (Lt16056F + Lt184R) was

further validated by Sanger sequencing, applying also the following additional sequencing

primer:

Le101F: TATAAATTCCTGCCAAACCCCAAAAA

Mitochondrial DNA assembly and annotation

Mitochondrial DNA was assembled using the MitoZ pipeline[28] and the best assembly was

selected by comparing the results of the pipeline's outputs with different kmer options. The

final assembly was done using the un options --clade Chordata –fastq_read_length 150, --

requiring_taxa Chordata --genetic_code 2 --kmers_megahit 21 29 39 59 79 99 119 141. The

pipeline included the tools fastp [29] for cleaning the raw data, MEGAHIT [30] for assembly,

after which sequences were filtered using HMMER [31] to ensure the correct taxa and the

completeness of protein-coding genes. Annotation was performed using TBLASTN [32],

GeneWise [33] and MiTFi [34]. The annotation of the non-coding region (NCR) as well as the

illustration of the mitochondrial genome was done with Geneious® 10.2.6 (Biomatters.

Available from https://www.geneious.com). The functional loci on the NCR were identified

based on the similarity with the human (NC_012920) and mouse (FJ374652) NCR sequences.

Genotyping of the long repeat (LR) region

The LR length genotyping of the hare samples was performed by PCR using the following

primers that bind the flanking regions of the repeat run (Fig. 1A, B):

Lepus timidus mtDNA

LtLR-F: AGAACCGTGACATAGCACTTACTTTC

LtLR-R: TAACATATTGGTGTTAGAATGTTTTTAGT

L. europaeus mtDNA

LeLR-F: TATAAATTCCTGCCAAACCCCAAAAA

LeLR-R: GCTTAATACCTGCTCCTCTTGATCTA

The used PCR program had an initial denaturation step of 94°C for 2 min, followed by 35

cycles of 94°C for 20 s., 59°C for 20 s. and 72°C for 1:30 min, with final elongation at 72°C

for 5:00 min, using AccuStart II™ PCR SuperMix (Quantabio). The PCR products were

separated over 1 % TAE agarose gel. In general, the Le primer pair works also for mountain

hares, but did not reliably amplify certain population samples.

Analyses of the SRs

Core SR sequences and their flanking regions of the Finnish *Lepus timidus* Finland and *Lepus*

europaeus were manually retrieved from the whole mitochondrial genome sequences. All

possible core dimers (1 for LE type, 4 for LT) were then used as query sequence for a basic

nucleotide BLAST search. Results were filtered to only include mitochondrial genomes

sequences from defined geographic isolates (LE query) or from one genome per *Lepus* species

(LT queries). For LT queries; *Lepus tibetanus*: MN539746.1, *L. tolai*: MN539744.1, *L. arcticus*:

NC_044769.1, *L. sinensis*: NC_025316.1, *L. coreanus*: NC_024259.1, *L. granatensis*:

NC_024042.1, *L. othus*: KJ397608.1, *L. corsicanus*: KJ397606.1, *L. capensis*: NC_015841.1

and *L. yarkandensis*: MN539747.1 mitochondrial genomes were selected. *L. townsendii* was

excluded due to poor quality sequencing of its SR region. For the LE query, *L. europaeus*

Poland 3: KY211034.1, Poland 2: KY211033.1, Poland 1: KY211032.1, Germany 2:

KY211031.1, Germany 1: KY211030.1, Greece 4: KY211029.1, Greece 3: KY211028.1,

Greece 1: KY211026.1, Cyprus 4: KY211025.1, Cyprus 3: KY211024.1, Cyprus 2:

KY211023.1, Cyprus 1: KY211022.1, and Turkey 1: KY211021.1 were selected. LE Greece 2

isolate was excluded as is SR region was identical to that of Greece 3 isolate. Rabbit

(*Oryctolagus cuniculus*) reference genome mtDNA (NC_001913.1) was used for a comparison.

The proportion of mutated core repeats was compared using Fisher exact test. Because of its

unique SR features, *L. yarkandensis* was excluded from the statistical analyses. Sequence

uncertainties in the repeats of *L. otus* (1) and *L. corsicanus* (3) were all considered as mutated

for the Fisher exact test, the mutation frequency was nevertheless significantly higher in LE-

type of SRs. To visualize the frequency of mutation in function of the repeat location, the repeat

positions were normalized to the number of repeats in the SR leading to an x-axis ranging from

0 ("beginning" of the SR region) to 1 ("end" of the SR region).

Generation of the cybrid cell lines

Cybrid cells were generated using the chemical enucleation method [35]. In brief, we first

generated recipient mtDNA-less ρ0 cells from immortalized LT1, LT4, LT6, LE1, LE2 and LE3

fibroblast cell lines by culturing them in the presence of 100 μM ddC, until mtDNA was not

detected anymore on a Southern blot. The nucleus of the mtDNA donor cells was killed with

Actinomycin D, with the effective concentration determined separately for each cell line. The

optimal concentration for most cell lines was 1 μg/ml for 18h, except for LT1 (1 μg/ml for 15h)

and LE2 (2 μg/ml for 28h). The cybrid fusions were conducted in both directions (same cell

line either a recipient or a donor of mtDNA) for the following cell line pairs: LE1♂ x LT4♀,

LE2♀ x LT1♂ and LE3♀ x LT6♂. For the control cybrids with mtDNA from the same species

but a different cell line than the nucleus, we used LE1♂(nucleus) x LE2♀(mtDNA) and LT4♀

(mtDNA) x LT6♂ (nucleus) combinations. The fusion was performed by growing the mtDNA

donor cells on 6-well plates and treating them with Actinomycin D. After replacing the

treatment medium with fresh DMEM (10% FBS) containing 1 million ρ0 cells were added to

the well for 3h. The cells were then washed 3 × with fresh DMEM before addition of 45%

polyethylene glycol (PEG, MW: 1450) for 60s. The cells were then washed again 3 × with

DMEM + 10% DMSO and 1 × with DMEM. Finally, the cells were given DMEM + 10% FBS

+ P/S. The cybrids were grown until they reached confluency, at which stage half of the cells

in a well were collected for genotyping. DNA was extracted with Quick-DNA Miniprep Kit

(Zymo Research).

As the donor and recipient cells differed by their sex, PCR-RFLP of *ZFX* and *ZFY* loci

[36] was used to genotype the cybrid nucleus and *CYTB* PCR-RFLP their mtDNA [37]. Each

original cybrid pool was genotyped and promising looking pools were subcloned on 96-wells.

These clonal cell lines were rechecked for the expected nuclear and mtDNA haplotypes,

obtaining the final cybrid cell lines. The ρ0 and cybrid cells were maintained in high glucose

(4.5 g/l) DMEM, supplemented with 10% FBS and 50 μg/ml uridine. The cybrid cells were

otherwise grown the same, but without added uridin to select for functional mitochondrial

DNA.

Agarose gel electrophoresis and Southern hybridization

Mitochondrial DNA from LE and LT parental and cybrid cell lines was isolated using

phenol:chloroform extraction and ethanol precipitation. 2 μg of uncut mitochondrial DNA was

separated over a 0.4 % agarose gel in 1x TBE buffer at 25 V for overnight at room temperature.

To facilitate the transfer of mtDNA onto nylon membrane by capillary transfer the gel was

subjected to acid depurination followed by denaturation to create suitable hybridization targets.

Transferred DNA was fixed to the blot by baking 80 °C for 2 h. Southern blot was probed using

(α-32P)-dCTP labelled PCR probe spanning hare mtDNA nucleotides 16840-88 or 17077-88

at 65 °C for overnight. Signals were captured on a phosphor screen and detected using

phosphorimager (GE Healthcare).

 Statistical tests

The RNAfold web server (http://rna.tbi.univie.ac.at/cgi-bin/RNAWebSuite/RNAfold.cgi) for

predicting secondary structures of single stranded RNA or DNA sequences was used to screen

for hairpin formation in the mitochondrial non-coding region and the QGRS mapper [38] to

identify putative Quadruplex forming G-Rich Sequences.

To test whether there was a geographical pattern in the LR length variation among

Finnish hares, a spatial autocorrelation analysis was used applying the Moran's I test with sf

and spdep packages in R (https://CRAN.R-project.org/).

Results

The structure of the mitochondrial DNA non-coding region in hares

We amplified the mitochondrial genomes from four mountain hare (LT1, LT4, LT5, LT6) and

four brown hare (LE1, LE2, LE3, LE4) cell lines as 12 overlapping PCR products, which were

subsequently sequenced using Illumina Miniseq™. When assembling the sequences, we noted

that the obtained sequence for the NCR was shorter than expected from the PCR product

lengths. Closer inspection revealed that this was because the short sequence reads (2×150 bp)

over the LR region were assembled as a single repeat element due to their redundancy. The

PCR products containing NCR elements were therefore reanalyzed using Sanger sequenced to

clarify LR copy number and finalized the assemblies (Table 1, Fig. 1). Interestingly, our

mountain hare as well as brown hare mitochondrial genomes are smaller (Table 1) than the

previously published mountain hare mtDNA from a Finnish specimen (NC_024040.1) (17,755

bp) or the brown hare mtDNA from Sweden (NC_004028.1). The size difference is explained

by the length of the non-coding region (NCR), where our LT cell lines have five and LE cell

lines six long tandem repeats (LRs) compared to seven in both previous mitochondrial

genomes.

The length of the NCR is 2,038 bp in LT1 (Fig. 2A) and 2,102 bp in LE1 (Fig. 2B). It

contains recognizable features such as a rather conserved region between the termination

associated sequence (TAS) and the cluster of short tandem repeats (SR), which by length and

location could correspond the D-loop region containing the 7S DNA. The SRs and the LRs are

separated by a variably long sequence with conserved sequence blocks (CSBs) 2 and 3, thought

to be relevant for the control of mtDNA gene expression and replication priming [39].

However, if replication is primed at or near the CSBs, the primer 3´-ends are relatively far away

from the known 5´-ends of DNA, considered to correspond the replication origin and start of

the 7S DNA [40]. In fact, Southern blot analysis of this region can only detect the 7S DNA

when using a probe that extends downstream of the SRs (Fig. 2C). Also, the size of the 7S does

not differ between the cell lines despites the difference in the length of the NCR-region. Of

note, we could confirm that Lagomorphs do not seem to have any CSB1[13, 21]. The CSB2

differs by a couple of indels between the species, the CSB3 seems highly conserved (Fig. 3A).

Hare SR elements consist of 10 bp core motifs

At first glance, the mountain hare (LT) SRs appear as a 20 bp repeat element similar in length

to the rabbits SRs (Fig. 3A). However, these are not consistent, and part of the SR arrays

showed repetition of only half of the sequence. In closer inspection, the element consists of

two 10 bp motifs, TAC<u>G</u>C<u>G</u>CACG, and TAC<u>A</u>C<u>C</u>CACG, which differ by the two underlined

nucleotides. In contrast, the brown hare (LE) SRs contain only one type of 10 bp core motif

(TAC<u>A</u>C<u>G</u>CACG), which differs from both LT motifs by one of the variable nucleotides (Fig.

3B). Interestingly, the 20 bp rabbit SR motif contains the 7 first nucleotides of the brown hare

(LE) SR motif spliced together with the last 8 nucleotides of LT SR motif1, separated by an

unique 5 bp (GTACG) spacer sequence (Fig. 3B).

While the LE-type is present only in brown hares, the two LT-types occur in all other

hare species that we analyzed (Table 2), with the two core elements alternating variably in the

SR arrays. Curiously, also the cape hare (*Lepus capensis*), a close relative of brown hare [23],

has LT-type SRs. *Lepus yarkandensis* constitute another curiosity, with randomly alternating

core 1 and 2 LT-type SR motifs being separated by di-nucleotides (TT or TC). Overall, the LT-

type SRs show considerably less variation than the LE-types (Fig. 3C, 3.35%; $p < 0.001$,

Fisher's exact test), despite being spread across many species. The copy number of the LT core

repeat sequences varied from 32 in our LT cell lines and other *L. timidus, coreanus, granatensis,*

otus and *corsicanus* isolates to 16 in *L. tibetanus* (Table 2). The copy number of the LE type,

present only in brown hares, ranged from 28 to 32.

As repetitive sequences are likely to form secondary structures, we were interested to

see whether the SR sequences of the two hares can form hairpins when single-stranded. This is

particularly interesting as hairpin formation at the second most prominent replication origin on

mitochondria, the light-strand origin (OriL) located at the WANCY tRNA cluster, is proposed

to be required for L-strand replication priming [41, 42]. A rather strong hairpin was predicted

for the H-strand SRs from the mountain hares but not from the brown hares, nor on the L-strand

SR sequences of either species (Fig. 4A–C). As for other structural features, the entire non-

coding region contained only a single sequence stretch at the 5′-end of the 7S DNA predicted

to form a strong guanine quadruplex (G4) (Fig. 4D).

LR length variation in Finnish hares

The LRs of the two species are similar in length (Fig. 5A), 139 bp in the mountain hare and

140 bp in the brown hare, differing at 21 nucleotide locations and containing the same 20 bp

conserved sequence element as previously reported for all Lagomorpha [13]. As also noted

earlier, the LR copy number shows variation between and even within individuals (Fig. 5B, C).

Although PCR tends to cause the amplification of short repeat sequences as an artefact, which

are evident as a faint ladder of PCR products of varying repeat content, the common repeat

haplotype was readily detectable as a strong band in all population samples and systematically

confirmed the sequencing results.

While the mechanism of dynamic length variation in the SRs has been shown to result

from replication slippage [16], the factors influencing the LR array length are less clear. As

they are also known to have species differences [6], we were interested to see whether the LR

array length would show population-specific variation. The most common length of LR was

five repeats in Finnish mountain hares, with six repeats being the next abundant (Fig. 6A). The

shortest observed repeat was four and longest seven, which was represented only by one

sample. Curiously, the only previously published mountain hare mtDNA from Finland

(NC_024040.1) also has seven repeats. Heteroplasmy was detected in five samples (3 %). No

geographic pattern in the LR length variation was observed among the 151 mountain hares

across Finland (Moran I statistic standard deviate = 8.09×10^{-10}, p-value = 0.5).

In contrast to mountain hares, the most common LR-region length in brown hares was

six repeats, with seven and eight repeats being more common than five (Fig. 6A). The longest

observed LR-region had nine repeats, while shorter than five repeats occurred only in

heteroplasmic samples (Fig. 5C). In the heteroplasmic samples, the shortest repeat sequence

length was three copies. Overall, brown hares showed more frequent heteroplasmy (12 %, *p* =

0.0045, Fisher exact test). Interestingly, brown hares with introgressed mountain hare mtDNA

showed an intermediate LR-region length distribution mtDNA (Fig. 6A). Although they had

the same median length of six repeats as brown hares with conspecific mtDNA, five repeats

were much more common (21% vs. 2%). Also a notable proportion (19 %) of the introgressed

samples were heteroplasmic, although the difference to the other brown hares was not

significant (*p* = 0.3077, Fisher exact test). Again, no correlation between the LR length and

geography was observed in brown hares with or without introgressed mtDNA. Only a few

mountain hares with brown hare mtDNA were present in our collection [25] and therefore were

not included in the analysis.

Transfer of mtDNA between cells causes instability in the LR copy number

Our LT and LE fibroblasts had respectively five and six LR repeats. If the repeat length is an

autonomous property of mitochondria, we expect it to remain stable despite the cohabitation

with a different nuclear genome. If it is controlled by the nuclear genome we expect it to be

rapidly converted to the copy number corresponding to the nuclear donor. While no change in

the LR copy number was observed when mtDNA was transferred from mountain hare to brown

hare cells (Fig. 6B), the six-copies LR array length of brown hare mtDNA increased to seven

in LT4, remained the same in LT1 and decreased to five in LT6 nuclear background (Fig. 6C).

Curiously, novel LR copy numbers of four and three appeared also in control cybrids with LT4

mtDNA in LT6 nuclear background (Fig. 6B), with no change observed in the brown hare

control cybrids (Fig. 6C). In all cases the new LR array haplotypes existed in heteroplasmy

with the original haplotype.

Discussion

When sequencing the mitochondrial DNAs as a part of the basic characterization of our hare

fibroblast cell lines [27], we noted a discrepancy between the expected genome size and the

assemblies obtained from short-read Illumina data. Closer inspection revealed that this was

because of misassembly of the part of the non-coding region with known long tandem repeats

(LRs). We corrected the assemblies with Sanger sequencing and provide here the fully

annotated mitochondrial genomes for four mountain hares and four brown hares from Finland

(Table 1, Figs. 1–2). Although tandem repeats are a known elements of the non-coding region

in many vertebrate species, including lagomorphs, they remain an overlooked aspects of

mitochondrial genome variation, largely due to their absence in humans and other well-studied

model organisms. In addition, and as in our case initially, they are likely to be lost when

assembling mitochondrial genome sequencing data obtained using short-read sequencing

platforms. For this reason, a lot of hare mitochondrial genome assemblies available in GenBank

have only one LR-element and many of the SR-array sequences might have issues as well. It

is likely that the same problem applies to the mitochondrial genome assemblies from other

organisms, meaning that the frequency and diversity of such elements in vertebrates is probably

underestimated. Recent development of long-molecule sequencing technologies can help to

complete our understanding of the repeat element variation, especially as the full-length

mitochondrial genomes can be assembled reliably from whole-genome sequencing data [43].

Not much is known about the function and biological significance of the repeat

elements in the non-coding region of mitochondrial genomes. Initial studies proposed that the

Lagomorph LRs are involved in the regulation of transcription and replication initiation [13,

21]. However, there are other conserved regulatory elements on the NCR (Fig. 2), which are

known to function in e.g. mtDNA replication priming in other mammals [39]. Furthermore, the

7S DNA, resulting from prematurely terminated replication, does not contain sequences

between LR- and SR-regions (Fig. 2C), indicating that the replication is likely to start at the

SR, as suggested by some authors [44].

Interestingly, the SR array length can change rather dynamically in rabbits [13, 16, 21]

and their copy number is highly variable (from three to 19). However, in hares the nature of

the repeating sequences made the delimitation of the actual repeat element difficult, as different

types of varying lengths repeats can be identified (Fig. 3A). Like in rabbits, a 20 bp SR element

can be identified in mountain hares. However, this element consists in fact of two separate

motifs (Fig. 3B), which can occur alone as the repeating element in the SR array. In brown

hares the SR arrays consist only of one core 10 bp motif, which differs from both mountain

hare motif types by a single nucleotide (Fig. 3B). If only these canonical sequences are counted,

their copy number in the different hare species varied between 16 and 32 (Table 2). Considering

the evolution of the SRs, it is interesting that the LT and LE core motif ends for also the majority

of the 20 bp rabbit, interrupted by five differing nucleotides GTACG. It is noteworthy that these

can be converted by C>T and A>G transitions to form a LT motif 2 + LE motif 20 bp consensus

sequence block (Fig. 3B). It is likely that such transition mutations can occur repeatedly during

evolution, while the G>C transversion, separating the LT motif 2 and 3′part of the rabbit motif

from the rest (Fig. 3B), is unlikely as a recurrent point mutation. Alternatively, it is possible

that the LE motif and LT motif 2 are recombinants of two ancestral repeat types, which are

present (in modified state) in rabbits. Overall, the remarkable conservation of the core LT-type

SR repeats in hares (*Lepus*) is intriguing (Fig. 3C). This suggests that their origin predates the

separation of these species and highlights the unique situation of *L. europaeus* SRs, whose core

sequence is not shared by the other *Lepus* species, including its close relative the cape hare (*L.*

capensis) [23]. Nevertheless, the identical length of the SRs core sequences (10 bp) and their

similarity in all *Lepus* species (Fig. 3B, only a single nucleotide difference *separates L.*

europaeus core sequence from each of two SR cores found in other *Lepus* species) can be only

explained by common ancestry or mechanism of origin among all species in the genus. The

sequence homogeny of the SR runs within a species is quite likely maintained by constant

contraction and reamplification of the repeat motifs, resulting in their rapid drift.

Three more observations are worth mentioning concerning the SRs region. (a) In all

species and independently of the SR core, the repeats located at the 3′ end and in the center of

the SR are the least conserved (Fig. 3C). Similar asymmetry is known to result from the

unidirectional gene conversion associated with some mitochondrial mobile elements or with

mating type switching in yeast [45, 46], which includes a double-strand break followed by

erosion of the ends that generates nucleotide variation after repair. (b) Amongst the 48 core

sequence variants in LE-type and LT-type of repeats, and even though the SR core sequences

show very strong homology, no LT-type core variant could be observed in species with LE-

type core repeats and *vice versa*. This is curious, as it suggests that a unique SR motif

amplification event has taken place specifically in the brown hare evolutionary lineage.

Sequencing mitochondrial SRs from more *Lepus* species and isolates would be needed to allow

drawing further hypotheses from this observation. (c) One species *L. yarkandensis* presents a

unique SR motive derived constituted of LT motif 1 + LT motif 2 + a two-nucleotide (TT or

TC) spacer. Similar short footprint signatures have been associated to the excision of some

transposons [47] as well as non-homologous end joining repair mechanisms [48].

The biological significance or functional roles of the SRs remain enigmatic. It is plausible

that the SRs could form secondary structures when being single-stranded, which could promote

priming by MTRPOL, as occurs during the initiation of L-strand replication at OriL [42].

Interestingly, hairpin formation is predicted to be stronger for the H-strand (Fig. 4), which –

following the OriL priming mechanism – would indicate a L-strand origin. Similarly, the

reported variation in the SR copy number [13, 21] and their proneness to mutate through strand-

slippage [16], are counterintuitive features for important genomic functions such as the

regulatory control of DNA-replication.

Like the short repeats, the long repeats contain a core (20 bp) sequence segment (Fig.

4A), which is conserved in all lagomorphs [13, 21]. Coincidentally, this is the same length as

the duplex SR motifs, although the sequence bears no resemblance to them and has no obvious

sequence features, such as palindromes. If it represents a regulatory element capable of binding

protein factors such as TFAM, as proposed [21], the existence of multiple such sites on mtDNA

is somewhat reminiscent of the initiator titration model of bacterial replication initiation [49,

50]. In *E. coli* for example, the initiation factor DnaA is titrated away from the replication origin

oriC binding sites by providing dispersed competing recognition sites dispersed elsewhere on

the genome. Replication of the genome causes the duplication of these sites, binding more DnaA and thereby reducing its availability to initiate new rounds of replication. The obvious problem for this mechanism to operate in the regulation of hare mtDNA replication initiation is the variability in the copy number of the LRs (Fig. 4B, C), where having three to nine copies of competitive binding sites per genome would effectively negate any accurate regulation, although this could be balanced by difference in mtDNA copy number and TFAM molecules. Unlike the dispersed DnaA sites, the LRs are clustered together and situated so that their doubling would occur only at the end of the replication. Thus, they are unlikely to prevent re-initiation before termination. Still, they could control replication by titrating TFAM in function of the number of mtDNA copies (and therefore SRs repeats) already present in the mitochondria. However, as with the SRs, if these elements have such an important role in genome regulation, why would they be present in some evolutionary lineages but not in others? For example, rodents (Rodentia) – a sister order of Lagomorpha – as well as most other mammals do not have such repetitive sequence arrays on their mtDNAs.

The species difference in the LR length distribution between brown hare and mountain hare (Fig. 6A) is interesting, as it tells that their variation between individuals is not random. In fact, similar differences in mtDNA tandem repeat lengths are known from other closely related species [6, 10, 11, 51]. As the species difference is fundamentally a genetic difference, one might expect that the LR copy number would then also be governed by the nuclear genome. However, the fact that LR copy number can vary between the tissues of the same individual in rabbits and that certain tissues are more prone to have longer arrays than others [13], suggests a more complicated mechanism, where the nuclear genetic background is only one variable. As different tissues have different metabolic requirements, manifesting as differences in mitochondrial activity and mtDNA maintenance strategies [52], it is plausible that the internal environment in mitochondria also indirectly effects the LR-region length through factors such

as temperature, oxidative stress and replication/repair protein availability. In fact, we have previously noted that the mountain hare fibroblasts maintain higher mitochondrial membrane potential than the brown hare cells [27], as well as have other differences in their oxidative metabolism. In this case, the energetic state of the cells can be influenced by variables such as metabolic adaptations, tissue type, ageing and stress, causing also the observed variation in the LR copy numbers. If the LRs are generated or lost as a result of strand-slippage during replication, as seems to be the case with the SRs [16], changes in the mitochondrial internal environment could increase the probability of such events, resulting in LR copy number change. Similarly, factors such as mitochondrial temperature [53], could favor certain repeat lengths over others and stabilize the LR copy number variation. Overall, it is curious that both the SR core sequence shows less conservation (Fig. 3C) and the LR arrays show more length variation in brown hares than in mountain hares.

Interestingly, our experiments with the cybrids showed mixed results (Fig. 6B). First, the effect on LR copy number was asymmetric and only presented by cells with mountain hare nucleus, regardless of the mtDNA origin. Also, the change in the LR array length in was not consistent. While in one cybrid line the brown hare mtDNA lost a LR copy, obtaining the common copy number of five for a mountain hare mtDNA, in another the copy number was increased to seven, a LR array length not presented by any of the parental cell lines. In contrast, shorter than parental mountain hare LR arrays appeared in the control cybrid with LT6 nucleus and LT4 mtDNA. Although the sample size is small, it is interesting that the observed effect is consistent with the frequent introgression of mountain hare mtDNA into brown hare population but not vice versa [25, 54], suggesting that mtDNA maintenance is perhaps more permissive or resilient in cells with brown hare nucleus. Alternatively, the mountain hare cells could be more sensitive to the stress caused by the process of generating cybrids. In all cases, the new

haplotype appeared in heteroplasmy with the old, showing that they arise as a spontaneous

rearrangement of the existing LRs and increase in abundancy through drift or selection.

As a conclusion, although being a decades old discovery, repeat elements on the non-

coding region remain an overlooked feature of Lagomorph mitochondrial genomes. We

recognized conserved 10 bp core sequences in the short repeats, with two different motifs

present in most hare species and one in brown hares. Interestingly, SR core element copies

showed sequence variation in brown hares, but not in other hare species. We also found that

while the long repeat arrays are dynamic, their presentation is not random, as the different

species show different length distributions. Considering their absence from most mammalian

species, as well as the lack of clear biological role in mtDNA maintenance and expression, it

is likely that both repeat types have risen and are maintained as selfish genetic elements. It is

unlikely that these elements are mobile and they are certainly too small to encode functional

gene products, their reproduction has to be related to mtDNA replication mechanisms. Why

such elements have not been lost from otherwise compact mitochondrial genomes and how

species-specific copy number variation is maintained, warrants for further research.

Acknowledgements

559 Dr Seppo Helisalmi and Dr Joose Raivo from the Gene diagnostics lab at Genome Center of

560 Eastern Finland are thanked for providing the mtDNA sequencing service. We thank Ms Anita

561 Kervinen for her valuable laboratory assistance in preparing the mtDNA PCRs for the

562 sequencing. This research is produced by the xHARES consortium, funded by the Academy of

563 Finland (grant no. 329264 for ED and JLOP).

References

1. Spinelli JB, Haigis MC. The multifaceted contributions of mitochondria to cellular metabolism. Nat Cell Biol. 2018;20(7):745-54. Epub 2018/06/29. doi: 10.1038/s41556-018-0124-1. PubMed PMID: 29950572; PubMed Central PMCID: PMCPMC6541229.

2. Adams KL, Palmer JD. Evolution of mitochondrial gene content: gene loss and transfer to the nucleus. Mol Phylogenet Evol. 2003;29(3):380-95. Epub 2003/11/15. doi: 10.1016/s1055-7903(03)00194-5. PubMed PMID: 14615181.

3. Boore JL. Animal mitochondrial genomes. Nucleic Acids Res. 1999;27(8):1767-80. Epub 1999/04/02. doi: 10.1093/nar/27.8.1767. PubMed PMID: 10101183; PubMed Central PMCID: PMCPMC148383.

4. Kasamatsu H, Robberson DL, Vinograd J. A novel closed-circular mitochondrial DNA with properties of a replicating intermediate. Proc Natl Acad Sci U S A. 1971;68(9):2252-7. Epub 1971/09/01. doi: 10.1073/pnas.68.9.2252. PubMed PMID: 5289384; PubMed Central PMCID: PMCPMC389395.

5. Jemt E, Persson O, Shi Y, Mehmedovic M, Uhler JP, Davila Lopez M, et al. Regulation of DNA replication at the end of the mitochondrial D-loop involves the helicase TWINKLE and a conserved sequence element. Nucleic Acids Res. 2015;43(19):9262-75. Epub 2015/08/09. doi: 10.1093/nar/gkv804. PubMed PMID: 26253742; PubMed Central PMCID: PMCPMC4627069.

6. Lunt DH, Whipple LE, Hyman BC. Mitochondrial DNA variable number tandem repeats (VNTRs): utility and problems in molecular ecology. Mol Ecol. 1998;7(11):1441-55. Epub 1998/11/20. doi: 10.1046/j.1365-294x.1998.00495.x. PubMed PMID: 9819900.

7. Solignac M, Monnerot M, Mounolou JC. Concerted Evolution of Sequence Repeats in Drosophila Mitochondrial-DNA. J Mol Evol. 1986;24(1-2):53-60. doi: Doi 10.1007/Bf02099951. PubMed PMID: WOS:A1986F784400006.

8.   Savolainen P, Arvestad L, Lundeberg J. mtDNA tandem repeats in domestic dogs and

wolves: mutation mechanism studied by analysis of the sequence of imperfect repeats.

Mol   Biol   Evol.   2000;17(4):474-88.   Epub   2000/03/31.   doi:

10.1093/oxfordjournals.molbev.a026328. PubMed PMID: 10742040.

9.   Wang TY, Tzeng CS, Teng HY, Chang T. Phylogeography and identification of a 187-bp-

long duplication within the mitochondrial control region of

(Teleostei: Balitoridae).   Zool   Stud.   2007;46(5):569-82.   PubMed   PMID:

WOS:000250416200004.

10.   Mundy NI, Helbig AJ. Origin and evolution of tandem repeats in the mitochondrial DNA

control region of shrikes (Lanius spp.). J Mol Evol. 2004;59(2):250-7. Epub 2004/10/16.

doi: 10.1007/s00239-004-2619-6. PubMed PMID: 15486698.

11.   Omote K, Nishida C, Dick MH, Masuda R. Limited phylogenetic distribution of a long

tandem-repeat cluster in the mitochondrial control region in Bubo (Aves, Strigidae) and

cluster variation in Blakiston's fish owl (Bubo blakistoni). Mol Phylogenet Evol.

2013;66(3):889-97.   Epub   2012/12/06.   doi: 10.1016/j.ympev.2012.11.015.   PubMed

PMID: 23211719.

12.   Wang X, Liu N, Zhang H, Yang XJ, Huang Y, Lei F. Extreme variation in patterns of

tandem repeats in mitochondrial control region of yellow-browed tits (Sylviparus

modestus, Paridae). Sci Rep. 2015;5:13227. Epub 2015/08/20. doi: 10.1038/srep13227.

PubMed PMID: 26288099; PubMed Central PMCID: PMCPMC4541255.

13.   Casane D, Dennebouy N, de Rochambeau H, Mounolou JC, Monnerot M. Nonneutral

evolution of tandem repeats in the mitochondrial DNA control region of lagomorphs.

Mol   Biol   Evol.   1997;14(8):779-89.   Epub   1997/08/01.   doi:

10.1093/oxfordjournals.molbev.a025818. PubMed PMID: 9254915.

14.   Hoelzel AR. Evolution by DNA turnover in the control region of vertebrate mitochondrial

614   DNA. Current Opinions in Genetics and Development. 1993;3:891-5.

615   15.   Ennafaa H, Monnerot M, Elgaaied A, Mounolou JC. Rabbit Mitochondrial-DNA -

616   Preliminary Comparison between Some Domestic and Wild Animals. Genet Sel Evol.

617   1987;19(3):279-88.   doi: DOI   10.1051/gse:19870301.   PubMed   PMID:

618   WOS:A1987K245500001.

619   16.   Pfeuty A, Gueride M, Lecellier G. Expansion/contraction of mammalian mitochondrial

620   DNA repeats in Escherichia coli mimics the mitochondrial heteroplasmy. J Mol Biol.

621   2001;314(4):709-16. Epub 2001/12/06. doi: 10.1006/jmbi.2001.5160. PubMed PMID:

622   11733991.

623   17.   Kobayashi T, Ganley AR. Recombination regulation by transcription-induced cohesin

624   dissociation in rDNA repeats. Science. 2005;309(5740):1581-4. Epub 2005/09/06. doi:

625   10.1126/science.1116102. PubMed PMID: 16141077.

626   18.   Thakur J, Packiaraj J, Henikoff S. Sequence, Chromatin and Evolution of Satellite DNA.

627   Int J Mol Sci. 2021;22(9). Epub 2021/05/01. doi: 10.3390/ijms22094309. PubMed

628   PMID: 33919233; PubMed Central PMCID: PMCPMC8122249.

629   19.   Phillips AF, Millet AR, Tigano M, Dubois SM, Crimmins H, Babin L, et al. Single-

630   Molecule Analysis of mtDNA Replication Uncovers the Basis of the Common Deletion.

631   Mol Cell. 2017;65(3):527-38 e6. Epub 2017/01/24. doi: 10.1016/j.molcel.2016.12.014.

632   PubMed PMID: 28111015.

633   20.   Persson O, Muthukumar Y, Basu S, Jenninger L, Uhler JP, Berglund AK, et al. Copy-

634   choice recombination during mitochondrial L-strand synthesis causes DNA deletions.

635   Nat Commun. 2019;10(1):759. Epub 2019/02/17. doi: 10.1038/s41467-019-08673-5.

636   PubMed PMID: 30770810; PubMed Central PMCID: PMCPMC6377680.

637 21. Dufresne C, Mignotte F, Gueride M. The presence of tandem repeats and the initiation of

638   replication in rabbit mitochondrial DNA. Eur J Biochem. 1996;235(3):593-600. doi: DOI

639   10.1111/j.1432-1033.1996.00593.x. PubMed PMID: WOS:A1996TU39700019.

640 22. Kuhl I, Miranda M, Posse V, Milenkovic D, Mourier A, Siira SJ, et al. POLRMT regulates

641   the switch between replication primer formation and gene expression of mammalian

642   mtDNA. Sci Adv. 2016;2(8):e1600963. Epub 2016/08/18. doi: 10.1126/sciadv.1600963.

643   PubMed PMID: 27532055; PubMed Central PMCID: PMCPMC4975551.

644 23. Ferreira MS, Jones MR, Callahan CM, Farelo L, Tolesa Z, Suchentrunk F, et al. The

645   Legacy of Recurrent Introgression during the Radiation of Hares. Syst Biol.

646   2021;70(3):593-607. Epub 2020/12/03. doi: 10.1093/sysbio/syaa088. PubMed PMID:

647   33263746; PubMed Central PMCID: PMCPMC8048390.

648 24. Levanen R, Thulin CG, Spong G, Pohjoismaki JLO. Widespread introgression of

649   mountain hare genes into Fennoscandian brown hare populations. Plos One. 2018;13(1).

650   doi: ARTN e0191790

651 10.1371/journal.pone.0191790. PubMed PMID: WOS:000423416600098.

652 25. Levanen R, Kunnasranta M, Pohjoismaki J. Mitochondrial DNA introgression at the

653   northern edge of the brown hare (Lepus europaeus) range. Annales Zoologici Fennici.

654   2018;55(1-3):15-24. PubMed PMID: WOS:000431406800002.

655 26. Pohjoismaki JLO, Michell C, Levanen R, Smith S. Hybridization with mountain hares

656   increases the functional allelic repertoire in brown hares. Sci Rep. 2021;11(1):15771.

657   Epub 2021/08/06. doi: 10.1038/s41598-021-95357-0. PubMed PMID: 34349207;

658   PubMed Central PMCID: PMCPMC8338973.

659 27. Gaertner K, Michell C, Tapanainen R, Goffart S, Saari S, Soininmaki M, et al. Molecular

660   phenotyping uncovers differences in basic housekeeping functions among closely related

species of hares (Lepus spp., Lagomorpha: Leporidae). Mol Ecol. 2023;32(15):4097-117.

Epub 2022/11/03. doi: 10.1111/mec.16755. PubMed PMID: 36320183.

28. Meng G, Li Y, Yang C, Liu S. MitoZ: a toolkit for animal mitochondrial genome

assembly, annotation and visualization. Nucleic Acids Res. 2019;47(11):e63. Epub

2019/03/14. doi: 10.1093/nar/gkz173. PubMed PMID: 30864657; PubMed Central

PMCID: PMCPMC6582343.

29. Chen SF, Zhou YQ, Chen YR, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor.

Bioinformatics. 2018;34(17):884-90. doi: 10.1093/bioinformatics/bty560. PubMed

PMID: WOS:000444317200035.

30. Li D, Liu CM, Luo R, Sadakane K, Lam TW. MEGAHIT: an ultra-fast single-node

solution for large and complex metagenomics assembly via succinct de Bruijn graph.

Bioinformatics.   2015;31(10):1674-6.   Epub   2015/01/23.   doi:

10.1093/bioinformatics/btv033. PubMed PMID: 25609793.

31. Wheeler TJ, Eddy SR. nhmmer: DNA homology search with profile HMMs.

Bioinformatics.   2013;29(19):2487-9.   Epub   2013/07/12.   doi:

10.1093/bioinformatics/btt403. PubMed PMID: 23842809; PubMed Central PMCID:

PMCPMC3777106.

32. Gertz EM, Yu YK, Agarwala R, Schaffer AA, Altschul SF. Composition-based statistics

and translated nucleotide searches: improving the TBLASTN module of BLAST. BMC

Biol. 2006;4:41. Epub 2006/12/13. doi: 10.1186/1741-7007-4-41. PubMed PMID:

17156431; PubMed Central PMCID: PMCPMC1779365.

33. Birney E, Clamp M, Durbin R. GeneWise and Genomewise. Genome Res.

2004;14(5):988-95. Epub 2004/05/05. doi: 10.1101/gr.1865504. PubMed PMID:

15123596; PubMed Central PMCID: PMCPMC479130.

34.   Juhling F, Putz J, Bernt M, Donath A, Middendorf M, Florentz C, Stadler PF. Improved

systematic tRNA gene annotation allows new insights into the evolution of mitochondrial

tRNA structures and into the mechanisms of mitochondrial genome rearrangements.

Nucleic Acids Res. 2012;40(7):2833-45. Epub 2011/12/06. doi: 10.1093/nar/gkr1131.

PubMed PMID: 22139921; PubMed Central PMCID: PMCPMC3326299.

35.   Bayona-Bafaluy MP, Manfredi G, Moraes CT. A chemical enucleation method for the

transfer of mitochondrial DNA to $\rho°$ cells -: art. no. e98. Nucleic Acids Research.

2003;31(16). doi: ARTN e98

10.1093/nar/gng100. PubMed PMID: WOS:000184783000045.

36.   Fontanesi L, Tazzoli M, Pecchioli E, Hauffe HC, Robinson TJ, Russo V. Sexing European

rabbits (Oryctolagus cuniculus), European brown hares (Lepus europaeus) and mountain

hares (Lepus timidus) with ZFX and ZFY loci. Mol Ecol Resour. 2008;8(6):1294-6. doi:

10.1111/j.1755-0998.2008.02167.x. PubMed PMID: WOS:000260528700022.

37.   Melo-Ferreira J, Boursot P, Suchentrunk F, Ferrand N, Alves PC. Invasion from the cold

past: extensive introgression of mountain hare (Lepus timidus) mitochondrial DNA into

three other hare species in northern Iberia. Mol Ecol. 2005;14(8):2459-64. Epub

2005/06/23. doi: 10.1111/j.1365-294X.2005.02599.x. PubMed PMID: 15969727.

38.   Kikin O, D'Antonio L, Bagga PS. QGRS Mapper: a web-based server for predicting G-

quadruplexes in nucleotide sequences. Nucleic Acids Res. 2006;34(Web Server

issue):W676-82. Epub 2006/07/18. doi: 10.1093/nar/gkl253. PubMed PMID: 16845096;

PubMed Central PMCID: PMCPMC1538864.

39.   Pham XH, Farge G, Shi Y, Gaspari M, Gustafsson CM, Falkenberg M. Conserved

sequence box II directs transcription termination and primer formation in mitochondria.

J Biol Chem. 2006;281(34):24647-52. Epub 2006/06/23. doi: 10.1074/jbc.M602429200.

PubMed PMID: 16790426.

40.   Pohjoismaki JLO, Forslund JME, Goffart S, Torregrosa-Munumer R, Wanrooij S. Known

Unknowns of Mammalian Mitochondrial DNA Maintenance. Bioessays.

2018;40(9):e1800102. Epub 2018/07/13. doi: 10.1002/bies.201800102. PubMed PMID:

29999547.

41.   Wanrooij S, Fuste JM, Farge G, Shi Y, Gustafsson CM, Falkenberg M. Human

mitochondrial RNA polymerase primes lagging-strand DNA synthesis in vitro. Proc Natl

Acad   Sci   U   S   A.   2008;105(32):11122-7.   Epub   2008/08/08.   doi:

10.1073/pnas.0805399105. PubMed PMID: 18685103; PubMed Central PMCID:

PMCPMC2516254.

42.   Fuste JM, Wanrooij S, Jemt E, Granycome CE, Cluett TJ, Shi Y, et al. Mitochondrial

RNA polymerase is needed for activation of the origin of light-strand DNA replication.

Mol Cell. 2010;37(1):67-78. Epub 2010/02/05. doi: 10.1016/j.molcel.2009.12.021.

PubMed PMID: 20129056.

43.   Uliano-Silva M, Ferreira J, Krasheninnikova K, Darwin Tree of Life C, Formenti G,

Abueg L, et al. MitoHiFi: a python pipeline for mitochondrial genome assembly from

PacBio high fidelity reads. BMC Bioinformatics. 2023;24(1):288. Epub 2023/07/19. doi:

10.1186/s12859-023-05385-y. PubMed PMID: 37464285; PubMed Central PMCID:

PMCPMC10354987.

44.   Melo-Ferreira J, Vilela J, Fonseca MM, da Fonseca RR, Boursot P, Alves PC. The elusive

nature of adaptive mitochondrial DNA evolution of an arctic lineage prone to frequent

introgression. Genome   Biol   Evol.   2014;6(4):886-96.   Epub   2014/04/04.   doi:

10.1093/gbe/evu059. PubMed   PMID: 24696399;   PubMed   Central   PMCID:

PMCPMC4007550.

45.   Stoddard BL. Homing endonucleases from mobile group I introns: discovery to genome

734   engineering. Mob DNA. 2014;5(1):7. Epub 2014/03/05. doi: 10.1186/1759-8753-5-7.

735   PubMed PMID: 24589358; PubMed Central PMCID: PMCPMC3943268.

736   46.   Richard GF, Kerrest A, Dujon B. Comparative genomics and molecular dynamics of

737   DNA repeats in eukaryotes. Microbiol Mol Biol Rev. 2008;72(4):686-727. Epub

738   2008/12/05. doi: 10.1128/MMBR.00011-08. PubMed PMID: 19052325; PubMed

739   Central PMCID: PMCPMC2593564.

740   47.   Skipper KA, Andersen PR, Sharma N, Mikkelsen JG. DNA transposon-based gene

741   vehicles - scenes from an evolutionary drive. J Biomed Sci. 2013;20(1):92. Epub

742   20131209. doi: 10.1186/1423-0127-20-92. PubMed PMID: 24320156; PubMed Central

743   PMCID: PMCPMC3878927.

744   48.   Yant SR, Kay MA. Nonhomologous-end-joining factors regulate DNA repair fidelity

745   during Sleeping Beauty element transposition in mammalian cells. Mol Cell Biol.

746   2003;23(23):8505-18.   doi: 10.1128/MCB.23.23.8505-8518.2003.   PubMed   PMID:

747   14612396; PubMed Central PMCID: PMCPMC262663.

748   49.   Ogawa T, Yamada Y, Kuroda T, Kishi T, Moriya S. The datA locus predominantly

749   contributes to the initiator titration mechanism in the control of replication initiation in

750   Escherichia   coli.   Mol   Microbiol.   2002;44(5):1367-75.   Epub   2002/06/19.   doi:

751   10.1046/j.1365-2958.2002.02969.x. PubMed PMID: 12068813.

752   50.   Hansen FG, Christensen BB, Atlung T. The initiator titration model: computer simulation

753   of chromosome and minichromosome control. Res Microbiol. 1991;142(2-3):161-7.

754   Epub 1991/02/01. doi: 10.1016/0923-2508(91)90025-6. PubMed PMID: 1925015.

755   51.   Hernández MA, Campos F, Gutiérrez-Corchero F, Amezcua A. Identification of Lanius

756   species and subspecies using tandem repeats in the mitochondrial DNA control region.

757   Ibis. 2004;146:227-30.

52.   Herbers E, Kekalainen NJ, Hangas A, Pohjoismaki JL, Goffart S. Tissue specific

differences in mitochondrial DNA maintenance and expression. Mitochondrion.

2019;44:85-92. Epub 2018/01/18. doi: 10.1016/j.mito.2018.01.004. PubMed PMID:

29339192.

53.   Chretien D, Benit P, Ha HH, Keipert S, El-Khoury R, Chang YT, et al. Mitochondria are

physiologically maintained at close to 50 degrees C. PLoS Biol. 2018;16(1):e2003992.

Epub 2018/01/26. doi: 10.1371/journal.pbio.2003992. PubMed PMID: 29370167;

PubMed Central PMCID: PMCPMC5784887.

54.   Thulin C-G, Tegelström H. Biased geographical distribution of mitochondrial DNA that

passed the species barrier from mountain hares to brown hares (genus Lepus): an effect

of genetic incompatibility and mating behaviour? J Zool. 2002;258:299-306. doi:

10.1017/S0952836902001425. PubMed PMID: WOS:000179553600006.

Tables

Table 1. The mitochondrial genomes from mountain hare (LT) and brown hare (LE) cell lines sequenced in this study.

Cell line	Species	Country	Municipality	Size	GenBank
LT1	*Lepus timidus*	Finland	Ilomantsi	17,482 bp	OR915850
LT4	*Lepus timidus*	Finland	Vesilahti	17,483 bp	OR939641
LT5	*Lepus timidus*	Finland	Outokumpu	17,481 bp	OR939643
LT6	*Lepus timidus*	Finland	Ruokolahti	17,471 bp	OR939642
LE1	*Lepus europaeus*	Finland	Liperi	17,540 bp	OR915849
LE2	*Lepus europaeus*	Finland	Outokumpu	17,546 bp	OR876275
LE3	*Lepus europaeus*	Finland	Kontiolahti	17,541 bp	OR939639
LE4	*Lepus europaeus*	Finland	Vesilahti	17,578 bp	OR939640

774   **Table 2.** The short repeat core element types and copy number variation among hares. Clade

775   indicates whether the species is more related to mountain hare (LT) or brown hare (LE). Core

776   element copy numbers represent only the analyzed mitochondrial genomes (see materials and

777   methods) and not a population study. LT+ for *Lepus yarkandensis* indicates a unique dimer

778   insertion in the classical LT-type SR (see text).

Species	Species clade	SR type	Core element copy number
Lepus timidus	LT	LT	32
Lepus tibetanus	LT	LT	16
Lepus tolai	LT	LT	17
Lepus arcticus	LT	LT	23
Lepus sinensis	LT	LT	28
Lepus coreanus	LT	LT	32
Lepus granatensis	LT	LT	32
Lepus otus	LT	LT	32
Lepus corsicanus	LT	LT	32
Lepus yarkandensis	LT	LT+	21
Lepus capensis	LE	LT	29
Lepus europaeus	LE	LE	28–32

781 **Figures**

783 **Fig. 1.** Schematic illustration and gene maps of hare mitochondrial genomes. (A) The

784 mitochondrial DNA from LT1 mountain hare (*Lepus timidus*) and (B) LE1 brown hare (*L.*

785 *europaeus*) fibroblasts. The binding sites for the long tandem repeat (LR) genotyping primers

786 (LR-F, LR-R) are indicated. See Fig. 2 for the details of the non-coding region.

Fig. 2. Sequence features of hare mitochondrial DNA non-coding region (NCR). (A) The NCR

of LT1 cells and (B) LE1 cells. The direction of transcription and replication, as well as probe

locations for the Southern blot are indicated. Note that the H-strand replication is the leading-

strand, initiated first and primed by the L-strand transcript. Key: trnP = tRNA-proline; TAS =

termination associated sequence; SR = short tandem repeat; CSB2 and 3 = conserved sequence

block 2 and 3; LR = long tandem repeat; trnaF = tRNA-phenylalanine. (C) A Southern blot of

uncut mtDNA from mountain hare (LT) and brown hare cells (LE), probed for the region

spanning nts 16,840–88 (left panel) or 17,077–88 (right panel). See the probe locations in (B).

 Right panel is overexposed to demonstrate the absence of 7S signal. Key: cat = catenated

 mtDNA forms, oc = open circles; lin = linear mtDNA; sc = supercoiled circles; 7S = 7S DNA.

 Fig. 3. Comparison of the sequence differences between the brown hare (*L. europaeus*) and

 mountain hare (*L. timidus*) mtDNA non-coding regions between the assumed start of the 7S

 DNA and the long tandem repeats. (A) The short tandem repeats (SRs) form a continuously

 repeating array of a single motif in brown hare, while the mountain hares show random

mixtures of two different repeat motifs. Repeating sequences and their direction are marked

with red bars. Mountain hare CSB2 also has more Cs flanking the central TA-pair, whereas the

CSB3 sequence is identical between the species. Note both SNP and indel variation between

the two species. (B) The 10 bp conserved motif sequences in the hare SRs. LE-type is only

seen in brown hares whereas LT-types are present in all hares analyzed here. Note the G-A

transition (red) and G-C transversion (blue) between the three core sequences. The 20 bp rabbit

SR motif is shown in two parts as comparison. Identical sequences in bold. Note the similarity

of the two rabbit sequence halves with the brown hare repeat motif. (C) Sequence conservation

of the core SR sequence motifs in hares. The LT-type shows remarkable conservation across

species, whereas LE-type shows more variation within one species.

Fig. 4. Structural features of the hare mitochondrial non-coding region. (A) Despite the

difference in the repeat element length and their overlap, there are only a few SNPs between

the short repeat elements (SR) from the brown hare (*europaeus*) and mountain hare (*timidus*).

(B) Predicted mountain hare and brown hare SR H-strand secondary structures. Note the longer

and stronger hairpin formation in mountain hare. (C) Predicted secondary structures on the SR

L-strand. (D) A putative G-quadruplex sequence close to the 5´-end of the 7S DNA (nts 17,078–

17,107 in LE1). G-duplets are highlighted.

Fig. 5. Long tandem repeats (LR) in hares. (A) Sequence comparison of LRs from brown hares

824   and mountain hares, highlighting the 20 bp conserved sequence present in all known

825   Lagomorpha. (B) An example of LR copy number variation in mountain hares. Arrowheads

826   highlight LR ladder generated as a PCR artefact. (C) Examples of LR copy number variants

827   present in brown hares. Size difference compared to (B) due to different PCR primers.

Fig. 6. Long tandem repeat (LR) copy number variation in Finnish hares. (A) The geographical distribution of LR copy number variants and their frequencies in Finnish mountain hares (*Lepus timidus*) and brown hares (*L. europaeus*) with conspecific or introgressed mtDNA. (B) LR copy numbers in the original mountain hare fibroblasts and xenomitochondrial cybrids with mtDNA from these cells. (C) The same for the parental and cybrid cells with brown hare mtDNA. LT = *Lepus timidus*; LE = *L. europaeus*. Asterisk indicates control cybrids with nucleus and mtDNA from the same species, but different parental cell lines.