Distinct neural bases of subcomponents of the attentional blink

Authors & Affiliations:

Swagata Halder[1]*, Deepak Velgapuni Raya[1], Devarajan Sridharan[1,2]*

[1]Centre for Neuroscience, Indian Institute of Science, Bangalore, India, and

[2]Computer Science and Automation, Indian Institute of Science, Bangalore, India

*Correspondence to:

Swagata Halder,
Graduate Program in Neuroscience,
Centre for Neuroscience,
Indian Institute of Science,
C. V. Raman Avenue,
Bangalore 560012, India
E-mail: swagata@iisc.ac.in

Devarajan Sridharan,
Associate Professor and
DST Swarna Jayanti Fellow,
Centre for Neuroscience,
Indian Institute of Science,
C. V. Raman Avenue,
Bangalore 560012, India
Phone: +91 802 293 3434
E-mail: sridhar@iisc.ac.in

Abstract

The attentional blink reflects a ubiquitous bottleneck with selecting and processing the second of two targets that occur in close temporal proximity. An extensive literature has examined the attention blink as a unitary phenomenon. As a result, which specific component of attention – perceptual sensitivity, choice bias or both – are compromised during the attentional blink, and their respective neural bases, remains unknown. Here, we address this question with a multialternative task and novel signal detection model, which decouples sensitivity from bias effects. We find that the attentional blink impairs specifically one component of attention – sensitivity – while leaving the other component – bias – unaffected. Distinct neural markers of the attentional blink mapped on to distinct subcomponents of the sensitivity deficits. Parieto-occipital N2p and P3 potential amplitudes characterized target detection deficits whereas long-range high-beta band (20-30 Hz) coherence between frontoparietal electrodes signalled target discrimination deficits. We synthesized these results with representational geometry analysis. The analysis revealed that detection and discrimination deficits were encoded along separable neural dimensions, whose configural distances robustly correlated with the neural markers of each. Overall, these findings provide detailed insights into the subcomponents of the attentional blink, and reveal dissociable neural bases underlying its detection and discrimination bottlenecks.

Lay summary

In daily life, our attention switches rapidly between different objects. For example, when driving, we may shift focus from a billboard on the roadside to a pedestrian in front, in quick succession. Yet, our ability to process the second object is severely compromised especially when it appears soon after the first: an impairment called the "attentional blink". In previous work, the attentional blink has been studied essentially as a "monolithic" (indivisible)

phenomenon. We design a behavioral model to divide the attentional blink into sub-components and show that the blink affects only one specific component (sensitivity). We also identify key neural markers for deficits associated with this component. Our findings may aid in understanding the neural origins of attention deficit disorders.

Keywords

attentional deficit; signal detection theory; sensitivity and bias; event-related potentials; long-range synchronization; neural population geometry

Introduction

Attention is a remarkable cognitive capacity that enables us to process relevant

information and filter out irrelevant information to guide behavior. Yet, attention is surprisingly

capacity limited[1]. This limitation is clearly revealed when we seek to perform multiple tasks

simultaneously or to tackle multiple goals in rapid succession[2]. A prime example of this

capacity limitation is the phenomenon of the attentional blink. When multiple targets are

presented sequentially – in a rapid serial visual stream – individuals are often unable to

accurately detect and identify the second of two targets, particularly when it is presented in

close temporal proximity (200-500 ms) to the first[3] (Fig. 1, left). The inability to process the

second target has been hypothesized to arise from various factors, including an inherent delay

in reallocating attention from the first target (T1) to the second target (T2)[4,5], a transient

bottleneck in working memory while processing the first target[6–10], or an inevitable

consequence of suppressing distractors that follow the first target in the visual stream[11–13].

The effect of the attentional blink on the processing of the second target is well studied.

In particular, previous studies have investigated the stage at which attentional blink affects

T2's processing (early or late) [14–17] and the neural basis of this effect, including the specific

brain regions involved[15,18–20]. Several theoretical frameworks characterize a sequence of

phases of the attentional blink, including target selection based on relevance, detection,

feature processing, and encoding into working memory[9,21]. Overall, there is little support for

attentional blink deficits at an early, sensory encoding[14] stage; by contrast, the vast majority

of literature suggests that T2's processing is affected at a late stage[8,10]. Consistent with these

behavioral results, scalp electroencephalography (EEG) studies have reported partial or

complete suppression of late event-related potential (ERP) components, particularly those

linked to attentional engagement (P2, N2, N2pc or VAN) [15,22–25], working memory (P3) [20,26–30]

or semantic processing (N400)[31]; early sensory components (P1/N1) are virtually

unaffected[20,24] (reviewed in detail in Zivony and Lamy, 2022[32]). These "late" effects are

hypothesized to be mediated by diverse functional regions within a frontoparietal network, both

based on correlational and causal evidence[33–35]. Activity in the parietal, lateral prefrontal and

anterior cingulate cortex, as measured with functional MRI (fMRI) was shown to decrease

when T2 was not detected (missed), even though stimulus-evoked activity in early visual areas

(e.g., V1) was relatively intact[36]. Long-range synchronization of EEG oscillations across fronto-

parietal regions in the beta-band (13-18 Hz) is impaired during the attentional blink[37]. In

addition, converging evidence from transcranial magnetic stimulation (TMS) studies suggest

a potentially role for the posterior parietal cortex in the attentional blink[34,35]: T2 detection and

identification performance were enhanced after the application of transcranial magnetic

stimulation (TMS) over the bilateral parietal cortex.

Despite this extensive literature, many previous studies have studied the attentional

blink as a unitary phenomenon. While some theoretical models[9,21,32] and experimental

studies[23,38] have explored distinct mechanisms underlying the attentional blink, several

fundamental questions about its distinct component mechanisms remain unanswered. One

key question concerns the precise nature of the "late" attentional blink bottleneck. The

attentional blink effect is typically quantified as a decrease in the proportion of correct T2

responses at short, relative to long, inter-target lags[3]. Yet, such an impairment can arise from

one of at least two processes – a reduction in the fidelity of T2's perceptual representation, a

deficit in decision-making based on T2's representation, or both. Signal detection theory[39] is

a widely applied psychophysical framework whose parameters – sensitivity (d') and bias (or

criterion, c) – quantify these perceptual and decisional components, respectively.

Distinguishing between sensitivity and criterion effects is crucial because a change in either

of these parameters can produce a change in the proportion of correct responses[40,41]. A lower

proportion of correct T2 detections may reflect not only a lower detection d' at short lags but

also a sub-optimal choice criterion corresponding, for instance, to a conservative detection

bias [42] (Fig. 1, right, top). Importantly, such criterion effects need not be uniform across inter-

target lags: the lag on each trial could be inferred based on various factors, such as trial length,

allowing participants to adopt different choice criteria for the different lags prior to making a

response.

Yet, whether the attentional blink induces primarily d' deficits, primarily bias deficits, or

a mixture of both, is largely unknown, and the limited evidence available is controversial[24,43].

This is perhaps because previous studies typically employed simple target detection or

identification tasks, in conjunction with conventional, one-dimensional signal detection

models[24,43], recent literature suggest that such tasks and models do not suffice to reliably

distinguish between sensitivity and bias components of attention[44–46]. A complete

understanding of the attentional blink requires dissociating sensitivity from bias deficits, with

appropriate signal detection models, and identifying their respective neural correlates.

A second question concerns the precise nature of attentional blink deficits. Is the

attentional blink a deficit with detecting T2 (Fig. 1, right, top) or discriminating T2's features,

or both (Fig. 1, right, bottom)? It is evident, from first principles, that distinct neural

computations and regions must mediate target detection versus discrimination. For example,

detecting an oriented grating in noise can be achieved with a simple, local visual cortex

computation: first, by aggregating responses of orientation-tuned visual neurons and then, by

testing if this activity exceeds a pre-set threshold[47]. By contrast, discriminating the orientation

of a target grating involves an arguably more nuanced computation; one that requires

comparing the relative activities of neurons tuned to different orientations relative to some

predefined axis (e.g., vertical meridian)[47]. This latter computation would also involve higher

cortical areas for maintaining the feature discrimination rule in working memory, such as the

prefrontal cortex[48–50]. Moreover, each of these computations could be affected by attention

differently[51].

Nonetheless, conventional attentional blink tasks often conflate these computations.

In a typical attentional blink task, participants must make identification judgments on T2[3,6]. For

example, participants may be asked to identify T2 based on its shape (e.g., a specific letter

among numbers) or its category (e.g., a face among non-faces). Yet, an identification deficit

during the attentional blink could arise from at least one of two sources[3,4,9,25,37] First, the blink

could produce a detection deficit, when the participant fails to reliably detect the occurrence

of T2 (Fig 1, Top Right). Alternatively, the blink could produce a discrimination deficit, when

the participant detects T2, but is unable to reliably discriminate its features (Fig 1, Bottom

Right). A combination of both of these deficits is also possible. To determine the precise neural

underpinnings of the attentional blink deficits, these two deficits must be dissociated, and their

respective neural correlates identified.

To tease apart these distinct subcomponents of the attentional blink, we developed a

multialternative task that involved concurrent detection of T2 and discrimination of its features

(Fig. 2A). To analyze multialternative behavioural responses, we developed a novel two-

dimensional signal detection model that decoupled, and separately quantified sensitivity and

bias deficits during the blink. With concurrent EEG recordings we analyzed key neural markers

– a local neural marker (event-related potentials) and a global neural marker (oscillatory

frontoparietal synchronization) – to test if these would map onto distinct subcomponents of the

attention blink (detection and discrimination deficits, respectively). We synthesized these

results with representational geometry analysis to understand the neural representations

underlying subcomponents of attentional blink-induced deficits. Our results reveal a double

dissociation between the neural and behavioral bases of detection and discrimination

bottlenecks underlying the attentional blink.

Results

The attentional blink produces both detection and discrimination deficits

Participants (n=24) performed an attentional blink task involving target detection and orientation discrimination (Fig. 2A, see also Methods). In a stream of serially flashed stimuli (100 ms each), the first target (T1) was an oriented grating (radius: 2 degrees of visual angle or dva) of low spatial frequency (0.6 cycles per degree or cpd), whereas the second target (T2) was either an oriented grating of comparatively higher spatial frequency (1.8 cpd, 67% of trials) or a blank (33 % of trials). T2 stimuli of high contrast (100%) and low contrast (titrated to individual threshold, see Methods) were interleaved with equal (50%) probability across trials. Targets were interspersed with plaid distractors, and inter-target onset intervals (T1-T2 "lag"-s) were drawn from a geometric distribution (100 ms to 900 ms; see Methods).

Unlike conventional attentional blink tasks[3,6,20], a key element of novelty in our task design was the following: for T1 participants provided a 2-alternative orientation judgement whereas for T2 they provided a 3-alternative, concurrent detection and identification judgement (Fig. 2A). Specifically, at the end of each trial, participants first indicated whether they perceived T1's orientation to be closer to the cardinal (0°, ±90°) or the diagonal axes (±45°) with two distinct button-press responses (two-alternative forced choice or 2-AFC) (Fig. 2A, penultimate panel from the right). Then, participants reported whether they had detected T2, and if so, whether its orientation was clockwise or counterclockwise of vertical, using 3 distinct button-press responses (three-alternative forced choice or 3-AFC) (Fig. 2A, rightmost). The 3-AFC decision enabled us to decouple the effect of the attentional blink on detection and discrimination performance.

First, we estimated detection and discrimination accuracies for T2 judgements across different T1-T2 lags; these analyses were performed only on trials in which participants provided accurate T1 responses (denoted as "T2|T1" in the figures). Detection accuracies were calculated based on the proportion of trials in which T2 was correctly detected (Methods).

Briefly, we computed the average proportion of hits, misidentifications, and correct rejections;

misidentifications were included because, although incorrectly identified, the target was

nevertheless correctly detected. In contrast, discrimination accuracies were derived from T2

present trials, based on the proportion of correct identifications alone (Methods). T2 detection

accuracies were markedly lower at the short (100 ms and 300 ms), as compared to the long,

inter-target lags (700 and 900 ms) (Fig. 2B); this induced a significant detection deficit,

measured as the difference in detection accuracies between the short and the long lags

(detection deficit = -8.9 ± 1.6%, mean ± s.e.m., z=-4.14, p<0.001, Wilcoxon signed rank test,

BF>10^2; data pooled across T2 contrasts). Similarly, T2 discrimination accuracies were also

significantly lower at the short lags compared to the long lags (discrimination deficit = -13.5 ±

1.8%, z=-4.25, p <0.001, BF>10^2) (Fig. 2C). In other words, the attentional blink induced both

a significant detection and a discrimination deficit for the processing of the second target.

Next, we tested if the magnitude of either the detection or the discrimination deficit

would vary depending on T2 stimulus contrast. For this we performed a two-way ANOVA with

either the detection or the discrimination accuracy as the response variable and lags (short or

long) and contrast (high/HC or low/LC) as independent factors. While we found a main effect

of both lag (detection: F(1,23)=29.8, p<0.001, BF>10^3, discrimination: F(1,23)=54.1, p<0.001,

BF>10^3) and contrast (detection: F(1,23)=21.02, p<0.001, BF>10^2, discrimination: F(1,23)

=13.75, p=0.001, BF=1.2), we found no significant interaction effect between lag and contrast

(detection: F(1,23)=1.92, p=0.113, BF=0.49, discrimination: F(1,23) = 0.93, p=0.450,

BF=0.40). In other words, attentional blink induced both a detection and discrimination deficit

regardless of stimulus contrast.

Finally, in addition to analyzing deficits with overall accuracies, we also analyzed blink-

induced effects on individual behavioral responses in the 3x3 stimulus-response contingency

table for the 3-AFC task (Fig. 2D). These responses fall under five categories – hits, correct

rejections, misidentifications, false alarms, and misses (Methods). Again, a two-way ANOVA

revealed a main effect of inter-target lag for all five response types (p<0.05; BF>10, SI Table

S1): essentially all correct response proportions (hits, correct rejections) decreased, and incorrect response proportions (false alarms, misses and misidentifications) increased, at short relative to long inter-target lags. A main effect of T2 contrast was observed for hit and miss proportions ($p<0.001$) but not for misidentification proportions ($p=0.090$) (SI Table S1); note that, by definition, false alarms and correct rejection proportions are independent of T2 contrast.

Overall, the attentional blink affected both components of identification – detection and discrimination – in terms of overall accuracy and individual psychometric measures. Whereas detection and discrimination accuracies varied with T2 stimuli's contrast, the strengths of detection and discrimination deficits induced by the blink did not depend on contrast.

Attentional blink selectively impairs sensitivity, but not the bias, subcomponent.

While the attentional blink induced deficits with both T2 detection and discrimination, such deficits could arise from either sensitivity or bias/criterion mechanisms. These two possibilities are illustrated in Figure 1, using a one-dimensional, signal detection theory (1-D SDT) model. In the first case (Fig. 1, right, top, upper row), a deficit in T2 detection occurs because of a deterioration in T2's signal evidence strength, which manifests behaviorally as lower perceptual sensitivity. In the second case (Fig. 1, right, top, lower row), the detection deficit occurs because of a more conservative evidence threshold for reporting the presence of T2, which manifests, behaviorally, as a higher criterion, or a lower bias. Either (or both) of these effects could induce an accuracy deficit (Fig.1, left). Similarly, discrimination deficits can occur either because of impaired sensitivity (Fig. 1, right, bottom, upper row) or a sub-optimal bias (Fig. 1, right, bottom, lower row). We sought to distinguish the precise source – sensitivity versus bias – of deficits induced by the attentional blink.

To address this question we developed a novel, two-dimensional signal detection model to analyze participants' T2 judgments in the 3-AFC task (see Discussion). Note that such multialternative responses cannot be correctly analyzed with a combination of one-

dimensional SDT models; the reasons are discussed in detail in previous studies[41,45] (see also

Discussion). Briefly, the three-alternative decision is modeled in a 2-dimensional decision

space: evidence for the presence (or absence) of T2 represented along the abscissa and

evidence for clockwise or counterclockwise orientations for T2 represented along the ordinate

(Fig. 3A); we term these the "detection" (y-axis) and "discrimination" (x-axis) axes,

respectively. On each trial, a bivariate decision variable (ψ) encodes T2's features, with ψ's

component along the detection and discrimination axes representing evidence for T2's

presence (ψ_{det}) and T2's orientation (ψ_{dis}), respectively. The distribution of ψ, across trials, is

modeled as a bivariate, isotropic Gaussian whose mean varies with T2's configuration across

each of five conditions (Fig. 3A-B; absent, LC/CW, LC/CCW, HC/CW, HC/CCW); thus, the

model incorporates one noise distribution (T2 absent), centered at the origin, and four signal

distributions. The means of the signal distributions along the detection and discrimination axes

represent the participant's sensitivity for detecting T2 (d'_{det}) and discriminating its orientation

(d'_{dis}), respectively. An inverted-Y shaped decision surface divides the decision space into 3

decision zones, one corresponding to each type of response: T2 clockwise (upper right), T2

counterclockwise (upper left) or T2 absent (lower). The decision surface is parameterized by

a detection threshold (t_{det}) and a discrimination criterion (c_{dis}); these parameters determine the

placement of the decision surface along the detection and discrimination axes, respectively.

In addition, the angle between the lower arms of the decision surface (β, Fig. 3B) is estimated

as a free parameter of the model. The model was fit to the 3x3 stimulus-response contingency

table derived from participants' 3-AFC responses (Methods) with maximum likelihood

estimation.

We fit five variants of the model with different constraints on the parameters based on

distinct sets of plausible assumptions: these included constraints on the parameters across

lags, as well as across low and high contrast T2 stimuli (Methods; SI Fig. S1A). Model

comparison analysis – based both on the Akaike Information Criterion (AIC)[52] and the

Bayesian information criterion (BIC)[52] – identified a model in which discrimination sensitivity

(d'$_{dis}$) was equal for high and low contrast T2 stimuli for each inter-target lag, and discrimination criterion (c$_{dis}$) and angle (β) were equal across lags (Model III, SI Fig. S1A, C-D). Goodness-of-fit p-values (Methods) were generally high for this model, (median >0.95; SI Fig. S1B) indicating satisfactory model fits to data.

The attentional blink significantly impaired both detection and discrimination sensitivity. T2 detection d' was significantly lower at short, compared to long, inter-target lags (d'$_{det}$ deficit = -0.94 ± 0.17, z=-3.80, p<0.001, signrank test; BF>10^3, data pooled across T2 contrasts) (Fig 3C). Similarly, T2 discrimination d' was also significantly lower at the short lags compared to the long lags (d'$_{dis}$ deficit = -0.61 ± 0.06, z=-4.25, p<0.001; BF>10^3) (Fig 3D). A two-way ANOVA with inter-target lag and T2 contrast as independent factors revealed a main effect of lag on both d'$_{det}$ (F(1,23)=30.3, p<0.001, BF>10^3) and d'$_{dis}$ (F(1,23)=100.3, p<0.001, BF>10^3). Yet, we found no significant interaction effect between lag and contrast for d'$_{det}$ (F(1,23)=2.3, p=0.141, BF=0.44). A similar interaction analysis could not be performed for d'$_{dis}$ as this parameter was constrained to be equal across contrasts, based on model selection analysis.

By contrast, attentional blink produced no systematic effect on either detection or discrimination criteria. Even though the detection threshold (t$_{det}$) was higher at short, compared to long, inter-target lags (t$_{det}$ deficit = -0.59 ± 0.17, z=-3.11, p<0.001, BF=14), the detection criterion (c$_{det}$) – the conventional SDT measure of bias (Methods) – was not statistically significantly different across lags (c$_{det}$ deficit = -0.16 ± 0.16, z=-0.28, p= 0.775, BF=0.34) (Fig 3E). Moreover, model selection analysis had already identified a model in which the discrimination criterion (c$_{dis}$) was constrained to be invariant across lags (Fig 3F), obviating analyses of lag effects on this parameter. Nonetheless, estimating this criterion – even with an unconstrained model – revealed substantial evidence against a statistically significant blink effect (c$_{dis}$ deficit = 0.02 ± 0.04, z=-0.485, p=0.627, BF=0.23).

Finally, we tested whether detection and discrimination deficits were likely to be mediated by common or dissociable processes. For this, first, we correlated the magnitude of the blink-induced detection and discrimination sensitivity deficits – the d' modulation index (MI)

– across participants (Methods). Detection and discrimination d' deficits were not statistically

significantly correlated (r=0.39, p=0.059); Bayes factor analysis revealed no clear evidence

for or against a correlation between these subcomponent deficits (BF=1.18) (SI Fig. S2, left).

Similar results were obtained upon correlating accuracy deficits also (SI Fig. S2, right). In

general, detection d' deficits varied widely even among individuals exhibiting a narrow range

of discrimination deficits, and vice versa (SI Fig. S2A-B). Second, we tested whether the

strength of detection and discrimination accuracy deficits would co-vary within each

participant. For this, we divided task blocks into those with the highest and lowest detection

accuracy MIs and compared the magnitude of the discrimination accuracy MI across these

blocks (Methods). Discrimination accuracy deficits were not statistically significantly different

between high and low detection accuracy deficit blocks (z=1.97, p=0.067), and the Bayes

factor revealed no strong evidence for or against such a difference (BF=1.42) (Fig. 3G).

In summary, we developed a novel, two-dimensional signal detection model to

simultaneously quantify blink-induced effects on T2's detection and discrimination. The

attentional blink affected both detection and discrimination sensitivity, across T2 stimuli of high

and low contrasts. Yet, no measurable effect occurred on detection or discrimination criteria.

In other words, performance deficits induced by the attentional blink could be attributed entirely

to sensitivity, rather than criterion, effects. Moreover, detection and discrimination d' deficits

were not significantly correlated both within and across participants, with no clear evidence for

or against a correlation, based on the Bayes factor. Next, we investigated electrophysiological

correlates of these behavioral deficits.

Dissociable electrophysiological markers of detection versus discrimination deficits

We sought to identify electrophysiological correlates of detection and discrimination

deficits during the attentional blink. As a first step, we examined the effect of established

signatures of the attentional blink, including event-related potential (ERP) magnitude[15,20] and

long-range synchronization[37]. EEG data were acquired from a subset (n=18/24) of

participants, while they were tested on the behavioral paradigm (Methods); high and low

contrast T2 trials were pooled to estimate reliable ERPs (Methods). To prevent confounding

neural correlates of detection and discrimination d' deficits, we quantified the partial correlation

(r_p) between the amplitudes of each EEG metric and either detection d' (d'_{det}) or discrimination

d' (d'_{dis}) across lags, while controlling for the value of the other parameter (d'_{dis} or d'_{det},

respectively) (Methods).

First, we quantified the change in peak amplitude for different ERP components

conventionally associated with the attentional blink: parietal P1[20,24], frontocentral P2[15,53],

occipital N2p[15,20,24,30] and parietal P3[15,18,54,55] (Figs. 4A-B, left). Among these ERP components,

the N2p component and the P2 component were both significantly suppressed during the blink

(Δamplitude, short-lag – long-lag: N2p=-0.47 ± 0.12 μV, z=-3.20, p=0.003, BF=40.0, P2=-0.19

± 0.07 μV, z=-2.54, p=0.021, BF=4.83, signed rank test) (Fig. 4A, right). Similarly, the parietal

P3 also showed a significant blink-induced suppression (P3= -0.45 ± 0.09μV, z=-3.59, p <

0.001, BF>10^2) (Fig. 4B, right). By contrast, the P1, an early sensory component, showed no

statistically significant blink-induced modulation (P1= 0.25 ± 0.16μV, z = 1.19, p=0.231, BF =

0.651) (SI Fig. S3). Results from various studies support each of these findings[20]. In addition,

we did not observe a statistically significant N1 component in the long lag trials (p=0.207, one-

tailed signed rank test for amplitude less than zero); the Bayes factor (BF=1.35) revealed no

clear evidence for an N1 component.

Next, we tested whether distinct ERP components correlated specifically with detection

versus the discrimination deficits induced by the attentional blink. Detection d' correlated both

with the N2p and P3 amplitudes (Fig. 4C and 4E) (partial correlation, N2p: r_p=0.34 p<0.001,

P3: r_p=0.30, p<0.001, permutation test) (Fig. 4D and 4F, top). However, discrimination d'

correlated with neither of these components (N2p: r_p=-0.01, p=0.970; P3: r_p=0.14, p=0.748)

(Fig. 4D-F, bottom). Yet, despite its amplitude being significantly modulated by the blink, the

frontocentral P2 component did not correlate with either detection d' (r_p=0.05, p=0.999) or

discrimination d' (r_p=0.23, p=0.120). Similarly, the P1 component did not correlate significantly

with either detection or discrimination d' (full set of results in SI Table S2). In other words, two

key late ERP components (N2p and P3) correlated with detection d' deficits, but not with

discrimination d' deficits, each after controlling for the variance explained by the other variable.

Finally, we tested whether long-range synchronization across the brain correlated

specifically with blink-induced detection or discrimination deficits. Following previous

reports[29,37] we investigated synchronization in the beta-band (13-30 Hz) between frontal and

parietal electrodes using spectral coherence (Methods). We observed a strong and sustained

decrease in fronto-parietal coherence during the attentional blink particularly, in the high-beta

(20-30 Hz) band (Fig. 5A-B, ΔPLV=PLV$_{short\text{-}lag}$–PLV$_{long\text{-}lag}$). Hemisphere-wise analysis revealed

a marked and significant reduction in fronto-parietal high-beta coherence over the left

hemisphere, in a cluster from 0 to 300 ms post T2 onset (permutation test, p=0.038, corrected;

cluster-forming threshold p<0.05) (Fig. 5C); by contrast, no discernible effects occurred over

the right hemispheric fronto-parietal electrodes (p>0.05) (Fig. 5D). Furthermore, we tested

whether high-beta coherence varied with detection d' or discrimination d', lag-wise, with partial

correlations (Methods). Discrimination d', but not detection d', significantly correlated with left

fronto-parietal coherence in the high-beta band (d'$_{dis}$: r_p=0.22, p=0.018; d'$_{det}$: r_p=-0.05,

p=0.316) (Fig. 5E-F).

We also examined correlations between detection d' or discrimination d' and low-beta

(13-19 Hz) coherence, as well as with beta coherence over right hemispheric fronto-parietal

electrodes but found no significant results; the full set of correlations is shown in SI Table S3.

To summarize, we observed a remarkable double dissociation between

electrophysiological markers of detection and discrimination deficits induced by the attentional

blink: whereas N2p and P3 suppression signaled blink-induced detection d' deficits, reduction

in high-beta phase synchronization between left fronto-parietal electrodes signaled

discrimination d' deficits.

Detection and discrimination deficits map to distinct neural dimensions

Recent studies suggest that an analysis of neural population geometry may provide mechanistic insights into diverse cognitive phenomena[56,57]. Inspired by these findings, we tested whether behavioral detection and discrimination deficits could be mapped to distinct neural dimensions. For this, we quantified the pairwise Euclidean distance, between the centroids of multivariate activity patterns produced by each class of T2 stimuli (clockwise, counterclockwise or no grating) in the posterior electrodes, separately for each inter-target lag (Methods). With these distances, we identified i) a "detection" dimension (η_{det}), a neural dimension that encoded the presence versus absence of T2 and ii) a "discrimination" dimension, (η_{dis}) encoded T2's orientation (clockwise versus counterclockwise) (Methods) (Figs. 6A-B, left). Neural inter-class distances ($||\eta||$) along both the detection and discrimination dimensions decreased significantly during the blink (short lag-long lag: $\Delta||\eta_{det}||$ = -1.30 $\pm$ 0.70, z=-3.68, p=0.006, BF=20; $\Delta||\eta_{dis}||$ = -1.23 $\pm$ 0.42, z=-3.54, p<0.001, BF>10^2) (Figs. 6C-D). In other words, neural representations encoding the presence of the target, as well as those encoding the target's features, became more overlapping – or less differentiated – during the attentional blink.

We tested whether the blink-induced collapse along the detection and discrimination neural dimensions would have distinct consequences for behavior. Neural inter-class distance along the detection dimension, $||\eta_{det}||$, revealed a significant partial correlation with detection d' across lags (r_p=0.27, p=0.018), but was not correlated with discrimination d' (r_p=0.16; p=0.122) (Fig. 6E-F). Conversely, neural inter-class distance along the discrimination dimension, $||\eta_{dis}||$, was significantly correlated with discrimination sensitivity (r_p=0.42, p<0.001) but not with detection sensitivity (r_p=0.20, p=0.122) (Fig. 6G-H). These results suggest a clear, double dissociation between the neural dimensions that underlie the detection and discrimination deficits induced by the attentional blink.

Next, we tested if inter-class distances along the detection and discrimination

dimensions correlated with the magnitude of blink-associated ERPs – the N2p or the P3. The

amplitude of the occipital N2p and parietal P3 were both significantly correlated with detection

distance, $||\eta_{det}||$ (N2p: $r_p=0.31$, $p=0.020$; P3: $r_p=0.46$, $p<0.001$) (Fig. 7A-B and SI Fig. S4A-B),

but not with discrimination distance, $||\eta_{dis}||$ (N2p: $r_p=0.07$, $p=0.330$; P3: $r_p=0.15$, $p=0.160$).

These results further confirm that the N2p and P3 ERP components indicate detection, but

not discrimination, bottlenecks underlying the attentional blink.

Finally, we tested whether long-range synchronization between fronto-parietal cortex

also varied with distances along these neural dimensions. In particular, we tested if left fronto-

parietal coherence correlated with distances along either the detection or the discrimination

neural dimension (η_{det} or η_{dis}, respectively). In line with our earlier findings, left frontoparietal

coherence in the high-beta band correlated significantly and specifically with discrimination

distance, $||\eta_{dis}||$ ($r_p=0.26$, $p=0.039$) (Fig. 7C-D), but not with detection distance, $||\eta_{det}||$ ($r_p=0.07$,

$p=0.255$). No such correlation occurred for either right frontoparietal coherence, or for

coherence in the low-beta band (SI Table S4). In other words, left hemispheric fronto-parietal

synchronization in the high-beta band provides a key marker for the discrimination, but not

detection, bottleneck underlying the attentional blink.

Taken together, these results indicate that the attentional blink impaired sensitivity both

for detecting T2 and discriminating its features, leaving criteria unaffected. Yet, neural markers

signaling detection and discrimination deficits were distinct. Detection deficits were signaled

by specific ERP components, namely the occipitoparietal N2p and the parietal P3. On the

other hand, discrimination deficits were signaled by long-range high-beta synchronization

between left hemispheric frontal and parietal cortex. Additionally, distinct neural dimensions

(η_{det} and η_{dis}) evidenced representational collapse associated with detection and

discrimination deficits and correlated systematically with neural markers of each type of deficit

(ERPs, and beta coherence, respectively). These results reveal a clear dissociation between

the behavioral and neural bases of detection and discrimination bottlenecks underlying the

attentional blink.

Discussion

When two successive targets appear in close temporal proximity, behavioural

judgments about the second target are compromised: a phenomenon termed the attentional

blink[3]. We show that attentional blink induces dissociable bottlenecks with both detecting the

second target and discriminating its features. Each of these bottlenecks affects the fidelity of

the second target's perceptual representations, but neither affects downstream criteria for

decisions. Moreover, these bottlenecks map on to distinct neural markers, indicating putatively

dissociable mechanisms mediating detection and discrimination deficits.

It is increasingly clear that detection and discrimination are separable processes, each

mediated by distinct neural mechanisms. Behaviorally, accurately identifying the first target,

versus merely detecting it, produces stronger deficits with identifying the second target[58].

Moreover, dissociable mechanisms have been reported to mediate object detection and

discrimination in visual adaptation contexts[59]. Neurally, shape detection and identification

judgements produce activations in non-overlapping clusters in various brain regions in the

visual cortex, inferior parietal cortex, and the medial frontal lobe[60]. Similarly, occipital ERPs

associated with conscious awareness also show clear differences between detection and

discrimination. For instance, an early posterior negative component (200-300 ms) was

significantly modulated in amplitude by success in detection, but not in identification[61]. The

closely related visual awareness negativity (VAN) was substantially stronger at the detection,

compared to the discrimination, threshold[62].

Furthermore, a significant body of previous work has reported dissociable behavioural

and neural mechanisms underlying attention's effects on target detection versus

discrimination. Behavioral studies have reported distinct effects on target detection versus

discrimination in both endogenous[63] and exogenous[64] attention tasks. Neurally, improved

detection of attended targets is accompanied by enhanced neuronal firing[65] or higher ERP

amplitudes[66]. On the other hand, improved discrimination of target features may be achieved

by selectively facilitating the activity of feature selective neurons[67], especially those that carry

maximally discriminative information about the target's features[68]. In addition, attention-

induced decorrelation of neuronal firing is hypothesized to systematically aid stimulus

discriminability[69], especially under conditions in which signal correlations are high among pairs

of neurons[70]. Furthermore, cortical network states become more distinct across object

representations, potentially facilitating target discrimination, in endogenous attention tasks[71].

Hence, there was no reason to expect, *a priori,* that impairments induced by the attentional

blink on target detection and discrimination would be correlated.

445   In line with this hypothesis, we discovered that the attentional blink induced dissociable

detection and discrimination deficits. There was no statistically significant correlation between

these two types of deficits within and across participants and evidence for such a correlation

was weak, at best. Unlike previous target identification designs that conflated attentional

blink's effect on detection versus discrimination performance[3,4,9,25,37], our 3-AFC task, and

associated signal detection model enabled quantifying each of these deficits separately and

identifying a double dissociation between their respective neural correlates. Our dissociation

of the attentional blink into distinct subcomponents is complementary to recent studies, which

examined whether the attentional blink reflects an all-or-none phenomenon[72,73]. For example,

the T2 deficit induced by the attentional blink can be either all-or-none or graded, depending

on whether T1 and T2 judgements involve distinct or common features, respectively[72]. While

a graded change in precision could reflect sensitivity effects, an all-or-none change in guess

rates – without a concomitant change in precision – may reflect a criterion increase

(conservative detection bias) effect. Future experiments that incorporate a three-alternative

response, with concurrent detection and discrimination, along with key task elements of these

earlier studies, may further help resolve these findings. The occipital N2p and parietal P3

amplitudes were modulated strongly by the attentional blink and correlated systematically and

selectively with detection deficits. Conversely, left hemispheric fronto-parietal high-beta

coherence was, again, modulated significantly by the attentional blink and strongly predictive,

specifically, of discrimination deficits. By contrast, neither the early occipital P1 nor the

frontocentral P2 systematically varied with either detection or discrimination deficits, although

the latter ERP was significantly diminished during the attentional blink.

Interestingly, we found no evidence indicating that these two computations (detection

and discrimination) were sequential; in fact, the modulation of beta coherence occurred almost

immediately after T2 onset, and lasted well afterwards (>400 ms from T2 onset) (Fig. 5A-B)

suggesting that an analysis of T2's features proceeded in parallel with its detection and

consolidation. We also modeled detection and discrimination as concurrent computations in

our SDT model (Fig. 3A-B). Previous work suggests that while object detection and

categorization processes proceed in parallel, detection and identification processes occur

sequentially[74]. Our results are in line with this literature, if we consider T2's discrimination

judgement – clockwise versus counterclockwise of vertical – to be a categorization, rather than

an identification judgement. Moreover, this earlier study[74] observed significant trial-wise

correlations between detection and categorization responses, suggesting that the two

processes involve the operation of the same perceptual filters ("analyzers"). Our study, on the

other hand, reports distinct neural bases for detection and discrimination computations. Yet,

the two sets of findings are not mutually contradictory.

In many conventional attentional blink tasks[3,20,25] , complex visual stimuli, like letters,

must be detected among a stream of background distractors with closely similar features, such

as digits. In this case, target detection would require the operation of shape-selective

perceptual filters for feature analysis. These same shape-selective filters would be involved

also for discriminating between distinct, but related target stimuli (e.g., two designated

candidate letters). In our task, target gratings needed to be distinguished in a stream of plainly

distinct background distractors (plaids), whereas the discrimination judgement involved

analysis of grating orientation. As a result, our task design likely precludes the need for the

same perceptual filters in the detection and the discrimination judgements. Absent this

common feature analysis, our results suggest distinct electrophysiological correlates for the

detection and discrimination of targets.

These results extend, and further advance, a significant literature on the attentional

blink effects on ERPs[15,18,20,24,25,29,32,53,54,75]. Several previous studies have shown that the

occipital N2 component exhibits a higher amplitude on short lag trials in which T2 was

successfully detected, versus not[15,20,24,29]. Other studies have shown that the attentional blink

also affects the amplitude of N2pc, a posterior, contralateralized N2 component[53–55]; a

decrease in the amplitude of the N2pc has been hypothesized to index impaired attentional

gating of T2. Similarly, previous studies have shown either partial or complete suppression of

P2 during the attentional blink.[15,25,53] Yet, these studies reported no change in P2 amplitude

between successfully and unsuccessfully detected targets, in line with our own findings. Taken

together with our own findings, these results establish the N2 as key marker for successful

detection of T2, achieved by robust attentional gating and filtering out of irrelevant temporal

(plaid) distractors.

The effect of the attentional blink on the P3 component has been consistently observed

in many studies, and its functional significance has been thoroughly investigated. Many

studies have reported a systematic reduction in P3 amplitude and latency in both detection

and identification tasks[15,20,24,29,53,55]. For centrally presented stimuli, the P3 recorded over

parietal cortex is generally linked to high-level cognitive functions, such as consolidation into

working memory[76]. In the present study, we observed that the parietal P3 amplitude was

correlated selectively with detection, rather than discrimination deficits. This suggests that the

P3 deficit indexes a specific bottleneck with encoding and consolidating T2 into working

memory, rather than an inability to reliably maintain its features. In this regard, a recent study[22]

measured ERP correlates of the perceptual awareness of the T2 stimulus whose relevance

was uncertain at the time of its presentation. In contrast to earlier work, this study observed

no change in P3b amplitude across seen (detected) and unseen targets. Taken together with

this study, our findings suggest that rather than indexing visual awareness, the P3 may index

detection, but only when information about the second target, or a decision about its

appearance, needs to be maintained in working memory. Additional experiments, involving

targets of uncertain relevance, along with our behavioral analysis framework, may help further

evaluate this hypothesis.

In contrast to the detection deficits associated with ERP amplitude suppression,

discrimination deficits were clearly correlated with disrupted high beta coherence in the

frontoparietal electrodes. Although previous studies have shown clear modulations in beta-

band coherence during the blink epoch[30,37], the tasks employed in these studies (e.g., letter

identification) could not distinguish between detection and discrimination deficits. To our

knowledge, ours is the first study to show that modulation of frontoparietal beta synchrony

correlates specifically with deficits associated with discrimination performance. This finding is

in line with previous literature, which suggests that frontoparietal beta synchrony plays a key

role in visual categorization, a function that relies critically on object feature analysis[77]. Beta-

band synchrony between the PFC and PPC may carry information about task relevant

categories[78]: mechanistically, the oscillations may serve to selectively communicate task

relevant features between PFC and PPC[78]. As a result, impairments in beta synchrony may

hinder the selection of relevant features leading to a selective discrimination deficit but not

detection deficit, as observed in our task. Moreover, previous studies have reported a

disruption in beta-band coherence between the left frontal and right parietal cortex[30,37]. Yet,

we observed significant left lateralization of frontoparietal beta-band coherence during the

attentional blink (Fig. 5B). One possibility is that this lateralization arises from beta oscillations

in the left motor cortex signaling right-handed responses[79]. We discount this hypothesis for

multiple reasons. First, the left-lateralization of the high-beta coherence occurred closely

following T2 onset, many hundreds of milliseconds before the response epoch. Second, the

beta coherence deficit correlated specifically with only one kind of perceptual deficit

(discrimination, but not detection). Finally, in our tasks, participants reported T2's orientation

with a bimanual response – the left index finger to report counter-clockwise orientations and the right index finger to report clockwise orientations. In other words, motor-activity linked beta oscillations were highly unlikely to be the source of the lateralized coherence deficits observed in our study.

Both successful target detection and discrimination improve significantly with stimulus contrast.[80] We had, therefore, expected to observe stronger detection and discrimination d' deficits for low contrast, as compared to high contrast targets. Surprisingly, we found no significant effect of contrast on either type of deficit (Figs. 2A-B). In other words, high (100%) contrast T2 stimuli were also strongly susceptible to the detection and discrimination bottlenecks associated with the attentional blink. Thus, despite a clear contrast-dependent encoding of T2 in early sensory cortex, the attentional blink produced a significant deficit with downstream processing, even for targets of high contrast. While at odds with some earlier work, which suggest an early-stage perceptual bottleneck [81–83], these results are largely consistent with findings from the majority of previous studies [3,7,9,11,19,20,81,84,85] which suggest a late-stage bottleneck.

But what specific neural computation underlies this late bottleneck? Previous research has documented three key computational stages mediating perceptual decisions: encoding of sensory evidence, formation of a decision variable, and application of a decision rule when formulating the choice[86]. Our results indicate that the attentional blink impacts, specifically, the second stage, associated with transforming T2's sensory evidence into a decision variable. In fact, our results indicate that the attentional blink did not affect the final, decisional stage, at all: criteria, for both detection and discrimination judgments, were essentially identical across inter-target lags.

These results appear at odds with earlier work, suggesting that the attentional blink produced both sensitivity and criterion effects in a detection task[24,43]. Lasaponara et al[24] reported a decrease in detection bias (likelihood ratio, or LR bias) with lag, whereas Caetta et

al.[43] reported an increase in detection criterion with lag. Our study, on the other hand, reports

no change in detection criterion with lag. Our results are potentially consistent with those of

Lasaponara et al. Given that d' increases with lag, a constant (positive) detection criterion

would mathematically yield a decreasing likelihood ratio bias with lag (LR bias = exp (-d' * c)).

In other words, a decrease in LR bias with lag could occur simply because of the increase in

detection d' without a concomitant change in detection criterion. On the other hand, a major

difference between Caetta et al's study[43] and ours is that Caetta et al modeled a fixed

(unvarying) detection threshold across inter-target lags, yielding changes in detection criteria;

the latter (detection criteria), and not thresholds, are the conventional SDT measure of bias[43].

Their choice of unvarying thresholds was based on the assumption that participants could not

predict the inter-lag of the trial in advance when different lags are interleaved across trials,

and, therefore, could not adopt different detection thresholds across lags. By contrast, we

considered it more reasonable that participants could modulate their detection thresholds

across lags, given that decisions about T2 were made at the end of each trial at which time

the participant would be well aware of the inter-target lag for that trial. To explore this

hypothesis further, we tested various models including these two models: one with a fixed

detection threshold and the other with a variable detection threshold across lags. The results

clearly supported a model with varying thresholds and confirmed a constant detection criterion

(bias) across lags in our data.

Taken together, these results suggest the following, testable schema (Fig. 8). First,

smaller N2p and P3 amplitudes may reflect deficits in the activation of neural populations

involved in gating and consolidating, respectively, the T2 stimulus into working memory.

Second, reduced fronto-parietal high-beta coherence may index deficits in a top-down

mechanism that enables discriminating T2's features; the latter computation relies on accurate

discrimination of clockwise versus counter-clockwise target orientations. Future research with

causal interventions like tACS or TMS could test this mechanistic schema of specific brain

regions or oscillatory patterns in mediating distinct aspects of the attentional blink. More

broadly, these findings contribute to our understanding of the relationship between attention

and perception and may be relevant for designing rational interventions for neurological

disorders that produce attention deficits.

Materials and Methods

Participants

Twenty-four healthy individuals (9 females; age range: 20-28 years) participated in the study. All participants had normal or corrected to normal vision with no known history of neurological disorders. Among these, scalp electroencephalography (EEG) recordings were acquired concurrently with behavior for eighteen participants. Experimental procedures were approved by the Institute Human Ethics Committee, Indian Institute of Science, Bangalore. Participants provided informed, written consent prior to participating in the experiments.

Behavioral Data Acquisition

Apparatus and Stimuli. Participants were tested with a typical Attentional Blink (AB task) protocol involving a rapid serial visual presentation (RSVP) stream (frame rate: 10 stimuli per second; 70 ms on, 30 ms off). Stimulus presentation and data acquisition were programmed with Psychtoolbox[87] and MATLAB 2017b MathWorks Inc. (2017), Natick, MA.). Participants were seated with their head resting on a custom chin rest in a dimly lit room, with their eyes positioned 60 cm away from a contrast-calibrated visual display (24" LCD monitor, 100 Hz refresh rate, BenQ Corp.). Responses were recorded with an RB-840 response box (Cedrus Inc). Participants were instructed to maintain gaze fixation at a central dot throughout the experiment and to avoid eye blinking during the stimulus presentation. Fixation, eye blinks and eye movements were monitored monocularly at 1000 Hz, with an infrared-based eye tracker (Eyelink, SR Research).

Task procedure. Each trial commenced with gaze fixation on a central dot (diameter: 0.5° of visual angle/dva, color: yellow) in the middle of a gray (35 cd/m^2 maximum luminance) screen. After 500 ms, task-irrelevant distractors – full contrast plaids (diameter: 4 dva) – appeared in the center of the screen in an RSVP stream (100 ms per stimulus). The plaid comprised of two superposed, square wave gratings (spatial frequency: 1.8 cpd) oriented at 45° and 135°

relative to the vertical, displayed within a circular mask (diameter: 4 dva). The phase of the

plaid stimuli underwent a 180° phase inversion (full white to full black, and vice versa) across

successive frames of the RSVP presentation. Plaids were encircled by a circular placeholder

(diameter: 4.1 dva, thickness: 0.1 dva). The fixation dot and the placeholder were present on

the screen throughout the trial.

Following a variable interval (300-1300 ms, geometrically distributed) of plaid

presentation, the first target stimulus (T1, oriented square-wave grating, spatial frequency: 0.6

cpd) appeared in the RSVP stream (Fig. 1A). Following another variable interval, the second

target stimulus (T2, oriented square-wave grating, 1.8 cpd), higher spatial frequency than T1,

was presented in the RSVP stream. The interval between T1 and T2 onsets was geometrically

distributed, and pseudorandomly selected from one of 5 possible "inter-target" lags (100, 300,

500, 700 or 900 ms); plaids, identical with those described previously, were presented in the

interleaving frames. The T2 grating appeared on only two-thirds of the trials ("T2 present"

trials). On the remaining third of the trials, randomly interleaved with the T2-present trials, no

T2 was presented ("no T2" trials); rather the plaid was replaced with a gray filled circle (0%

contrast) inside the circular placeholder. In addition, on 50% of the trials, T2 gratings were

presented at full contrast, whereas on the remaining 50% of the trials (randomly interleaved)

T2 grating were presented at a contrast staircased for each participant individually (see next).

T1 orientations, as well as T2 orientations and contrasts, were proportionately

counterbalanced across all inter-target lags. Finally, plaids were presented for a set of six

frames (600 ms) and participants were probed for response.

At the end of each trial, participants provided two responses, one for each of T1 and

T2, in succession (Fig. 1A). First, participants indicated whether the T1 grating's orientation

was closer to the cardinal (0°, 90°) or to the diagonal (45°, 135°) axes with one of two button

presses (2-alternative forced choice/2-AFC, Fig. 1A). Second, participants indicated whether

the T2 grating's orientation was clockwise of vertical (CW), counter-clockwise of vertical

(CCW) or if it was not detected at all (None), with one of three distinct button presses (3-AFC,

Fig. 1A).

<u>Training and testing.</u> Two behavioral sessions were conducted on two consecutive days. The

first session (on Day 1) comprised of training (30 minutes) followed by a staircasing session

(~1.5 hours). The training session typically comprised 2-4 blocks of trials with 84 trials each.

Only the longest inter-target lags (700 and 900 ms) were included; short-lag trials were

excluded both from this, and the subsequent staircasing, sessions based on previous reports,

which suggest that practice effects can mitigate the magnitude of the attentional blink at short-

lags[88]. At the end of each trial participants received on-screen feedback regarding their

success or failure for each target. For this session, T1 and T2 were both presented at full

contrast; T1 was presented with a ±2° deviation from cardinal or diagonal orientations, pseudo

randomly counterbalanced among the 4 different options, whereas T2 was presented at ±10°

deviation from the vertical. Note that smaller (larger) angular deviations render the

cardinal/diagonal (CW/CCW) judgement with the T1 (T2) grating easier. Following this,

participants performed a second, staircasing session (~400 trials). In this session, T1's angular

deviation from the cardinal/diagonal orientations was staircased to achieve 87.5%

discrimination accuracy for T1, whereas T2's contrast was concurrently staircased to achieve

75% discrimination accuracy for T2. T1's angular deviation and T2's contrast determined from

this session were used for the main experiment on Day 2; while the former angle was used for

all T1 trials, the latter contrast was employed in the low-contrast T2 trials alone.

On Day 2, participants were re-familiarized with the task for ~150-200 trials before the

main experiment commenced. Before testing, participants were provided with a 30-minute

break. The main experiment comprised trials with all 5 inter-target lags (see above) and

comprised 3 blocks of 312 trials each (total of 936 trials per participant). No feedback was

provided after each trial. Each block was sub-divided into 4 mini-blocks of 78 trials, and

participants were provided short breaks (~5 min) after each mini-block. Data from the staircase

and training sessions were excluded from subsequent analyses.

<u>Eye-tracking and data exclusion.</u> Throughout each experiment participants' fixation was

monitored and gaze position was recorded and stored for offline analysis. Trials in which an

eye blink occurred any time before the response period were excluded from the final analyses.

In addition, trials in which the gaze positions deviated from the central pedestal ±50 ms before

or after the presentation of either the T1 or the T2 gratings (radius: 2 dva from fixation dot)

were excluded from further analysis. Following these exclusions, the median trial rejection rate

was 2.1% [0.8-3.4%] (median, 95% confidence interval) across participants.

Behavioral data analysis

<u>Measuring attentional blink effects on psychometric quantities.</u> We quantified the effect of the

attentional blink on behavioral detection and discrimination psychometrics for the T2 grating.

All of the analyses were performed including only trials in which participants made correct

responses for the T1 judgment. Our 3-alternative task for T2 judgements, and the associated

behavioral model (see next), enabled quantifying psychophysical parameters linked to both

detection and discrimination within a single task and analysis framework. Participants' 3-AFC

responses for the T2 grating were collated into five 3x3 stimulus-response contingency tables,

separately for each of the 5 inter-target lags, and each T2 contrast (high, low). The rows of

each contingency table represent the three stimulus events (T2 CW, T2 CCW or no T2) and

the columns represent the participants' choices among the three alternatives (Fig. 2C). Each

table comprises five responses types – i) Hits (H, elements along the main diagonal, except

the bottom right) wherein the subject correctly identifies the orientation of T2 as CW or CCW,

ii) Misses (M, last column, except the bottom right), wherein the subject reported "no T2" on

trials in which T2 was actually presented, iii) False alarms (FA, last row, except the bottom

right), wherein incorrectly reported the presence of T2 (as CW or CCW) when no T2 was

presented, iv) Correct Rejections (CR, bottom right), wherein the subject accurately reported

the absence of T2, and v) Misidentifications (MI, all other entries), wherein the subject

incorrectly reported T2's orientation (CW grating as CCW, or vice versa).

Based on these psychometric measures, we computed detection and discrimination

accuracies as follows. Detection accuracies were computed as the average proportion of the

hits, misidentification and correct rejection responses; misidentifications were included

because not missing the target reflected accurate detection. By contrast, discrimination

accuracies were computed based on the average proportion of the two correct identifications

(hits) on T2 present trials alone. We performed 2-way ANOVAs on both detection and

discrimination accuracies with the inter-target lag (5 values) and T2 contrast independent

factors. We found main effects of both lag ($F_{(4,92)}=18.81$, $p<0.001$) and contrast

($F_{(1,92)}=21.78$, $p<0.001$) on detection accuracy, but no interaction effect between lag and

contrast ($F_{(4,92)}=1.92$, $p=0.113$). Similarly, we found main effects of both lag ($F_{(4,92)}=25.08$,

$p<0.001$) and contrast ($F_{(1,92)}=16.58$, $p<0.001$) on discrimination accuracy, but no interaction

effect between lag and contrast ($F_{(4,92)}=0.93$, $p=0.450$). Post-hoc tests based on Tukey's

HSD revealed a significant difference in discrimination accuracies between the two shortest

lags (100 ms and 300 ms) and the two longest lags (700 and 900 ms) for both low and high

contrast targets, and for both detection and discrimination accuracies ($p<0.01$). But they

revealed no significant difference between the two shortest lags ($p>0.25$) or the two longest

lags ($p>0.40$) for either target contrast or for either accuracy type. As a result, for subsequent

analyses, we pooled together the "short lag" (100 ms and 300 ms) and the "long lag" (700 ms

and 900 ms) trials. We quantified the effect of the attentional blink on each of the psychometric

measures as well as detection and discrimination accuracies by comparing their respective,

average values between the short lag and long lag trials, separately for the high and low T2

contrasts.

<u>Measuring attentional blink effects on psychophysical parameters.</u> We estimated

psychophysical parameters by fitting the 3-AFC contingency tables, described above, with a

novel signal detection theory (SDT) model. Observers' decisions were modeled in a two-

dimensional decision space to fit T2 detection and discrimination performance simultaneously;

the model was inspired by the recently published m-ADC model[45] widely used for the analysis

of multialternative attention tasks.

i) Signal detection model description: On each trial, T2 stimulus' features were encoded with

a bivariate Gaussian decision variable (**ψ**). The x-axis of this decision space represents

evidence for T2's orientation (CW or CCW), whereas the y-axis represents evidence for T2's

presence. In other words, a larger signal magnitude along the x-axis (either positive or

negative) favors better discrimination whereas a higher signal magnitude along the y-axis

favors more reliable detection. The mean of the bivariate decision variable's value along the

x-axis and y-axis is parameterized by two perceptual parameters, indexing perceptual

discriminability ($\pm d'_{dis}$) for discriminating T2's orientation, and perceptual sensitivity (d'_{det}) for

detecting T2's presence, respectively. On the T2 absent trials **ψ** is drawn from a distribution

centered at the origin ($d'_{dis} = d'_{det} = 0$). On T2 present trials, the decision variable **ψ** is drawn

from one of 4 distributions (Fig. 3A), depending on T2's contrast (Fig. 3A; high, low) or its

orientation (Fig. 3A; CW, CCW).

The participant makes one of 3 choices – T2 CW, T2 CCW or no T2 – based on a Y-

shaped decision surface (Fig. 3B) that divides the decision space into 3 zones (red, blue, or

gray shaded regions, respectively). This decision surface is parameterized by three decisional

parameters: i) a detection criterion (c_{det}) that determines the displacement of this surface along

the vertical direction (y-axis), ii) a discrimination criterion (c_{dis}) that determines the

displacement of this surface along the horizontal direction (x-axis) and iii) an angle (β) between

the two arms of the Y (Fig. 3B). The subject's decision on each trial is based on the decision

zone into which **ψ** fell on that trial. Note that the proportion of each type of response for each

stimulus event type CW, CCW or no T2 in the stimulus-response contingency table can be

mapped to the integral of the conditional density of the decision variable within the decision

zone for the respective T2 event and response type. These integrals were evaluated

numerically (*integral2* function in Matlab).

Parameters (d', c, β) were estimated with a maximum likelihood estimation approach[45].

Typically, parameters were fit for each lag, independently, unless specified otherwise (see

next). Goodness of fit was assessed using a randomisation test based on the chi-squared

statistic[45]; $p < 0.05$ was taken as evidence of significant deviation of model fits from

experimental data.

ii) Model comparison analysis: We fit observers' response proportions to five models, each

with a set of progressively stronger constraints (decreasing model complexity).

Model I. This model was the least constrained and comprised 35 parameters. 20 perceptual

parameters d'_{det} and d'_{dis} were estimated individually for five lags and separately for two T2

contrasts. Similarly, 15 decisional parameters c_{det}, c_{dis} and β were estimated individually for

each of the 5 lags, separately (15 c and β parameters); because the high and low T2 contrasts

trials were randomly interleaved within each block, with no apriori information about which

type of target contrast would occur, we modeled these decisional parameters as being

identical across high and low T2 contrast trials.

Model II. This model was identical to Model I except that β was constrained to be identical

across lags (20 d', 10c, 1 β). Therefore, this model comprised 31 parameters.

Model III. This model was identical with Model II, except that additional constraints were

imposed on discrimination-related parameters: i) d'_{dis} was constrained to be identical across

contrasts and ii) c_{dis} was constrained to be identical across lags. This model comprised 22

parameters.

Model IV. This model was also identical with Model II, except that additional constraints were

imposed on detection-related parameters: i) d'_{det} was constrained to be identical across

contrasts and ii) c_{det} was constrained to be identical across lags. This model also comprised

22 parameters.

Model V. This model was the most constrained of all and comprised all constraints in Models

II-IV. This model comprised 13 parameters.

Formal model comparison was performed with the Akaike Information Criterion (AIC)

and Bayesian Information Criterion (BIC)[52]. These scores represent a trade-off between model

complexity (the number of model parameters) and goodness-of-fit (based on the log-likelihood

function); a lower AIC/BIC score represents a better candidate model[52]. Based on this

analysis, the best model was determined to be Model 3 (see Results, section on "*Detection*

and discrimination deficits reflect sensitivity, rather than criterion, effects").

iii) Quantifying the attentional blink effect: To quantify the effect of the attentional blink on each

of the estimated parameters, the average value of the parameter at the shortest two lags (100

and 300 ms, SL) was subtracted from the average value of the parameters at the longest two

lags (700 and 900 ms, LL). e.g, attentional blink effect on detection sensitivity was computed

as: $\Delta d'_{det} = d'^{LL}_{det} - d'^{SL}_{det}$. This computation was performed for each T2 contrast separately In

some cases (Fig. 3G, SI Fig. S2), the deficit was quantified with a modulation index. e.g. MI-

$d'_{det} = (d'^{LL}_{det} - d'^{SL}_{det}.)./(d'^{LL}_{det} + d'^{SL}_{det}.)$ Because, in the best model selected, c_{dis} and β were

constrained to be equal across lags no blink effect was calculated for these parameters. Error

bars were computed as the standard error of mean across participants.

EEG data acquisition and preprocessing

<u>Acquisition.</u> EEG data were recorded with a dense array of scalp electrodes (128 channels,

EGI's Geodesic Inc USA), a Net Amps 400 amplifier, and EGI's Net Station software (version

5.2.0.2, Oregon, USA). During acquisition, data were referenced to the Cz electrode, and

stored for offline analyses. The EEG cap contains a few additional electrodes that are placed

around the orbits of the eye and enable monitoring an EOG signal. Channel impedances were

monitored and maintained at <50 kΩ during the experiment. Data were sampled at 1000 Hz

and stored offline for analysis.

Preprocessing. The data were pre-processed with functions from the EEGLAB toolbox (v.

14.0.0), Chronux toolbox (v. 2.12.03) as well as custom scripts developed in Matlab. First, the

data were downsampled to 250 Hz by decimation and bandpass filtered between 0.5 to 18 Hz

(FIR filter, *designfilt* function in Matlab). Next, a bandstop filter from 9-11 Hz was applied to

remove the 10 Hz oscillations evoked by the RSVP presentation. Channel traces were then

re-referenced to the average signal across channels. We then epoched the data into trials and

applied SCADS (Statistical Control of Artifacts in Dense Array EEG/MEG Studies[89]) to identify

bad epochs and artifact contaminated channels. SCADS detects artifacts based on three

measures: maximum amplitude over time, standard deviation over time, and first derivative

(gradient) over time. Any electrode or trial exhibiting values outside the specified boundaries

for these measures was excluded. The boundaries were defined as $M \pm n \ast \lambda$, where M is the

grand median across electrodes and trials for each of the three measures, and λ is the root

mean square (RMS) of the deviation of medians across sensors relative to the grand median.

We set n to 3, allowing data within three boundaries to be retained. The percentage of

electrodes per participant rejected was 6.3 ± 0.43% (mean ± s.e.m. across participants),

whereas the percentage of trials rejected per electrode and participant was 3.4 ± 0.33% (mean

± s.e.m.).

Bad channels were linearly interpolated using data from the four nearest neighbour electrodes.

Epochs were demeaned based on a baseline computed in fixation time window (250 ms) at

the beginning of each the trial and then quadratically detrended.

EEG data analysis

Event Related Potential (ERP) analysis. ERP estimation for this attentional blink task followed

a procedure outlined in (Sergent et al.,2005)[20]. First trials were sorted based on inter-target

lags (100, 300, 500, 700 and 900 ms). This yielded an average of (200±13, 171±9.71, 145 ±

7.54, 117 ± 5.43, 87 ± 4.51) (mean ± s.e.m. across participants) trials for each of the 5 lags,

respectively. Trials from the longest two lags (700 and 900 ms) were pooled to provide

sufficient trials for robust ERP estimation at these lags. Then, EEG traces were epoched from

-300 ms before to +700 ms after either T1 onset or T2 onset and averaged across trials to

estimate T1-evoked and T2-evoked ERPs, respectively. Because of the periodic RSVP

presentation these ERPs included strong, plaid-evoked responses, which occurred time-

locked to both T1 and T2. To remove this contribution, we subtracted ERPs associated with

T2-absent trials (correct rejection trials only) from ERPs associated with T2-present trials (as

in[15,20]); this subtraction was performed separately for each inter-target lag. ERP amplitudes

were quantified by measuring peak values within specific time windows corresponding to

distinct components.

We analyzed five ERP components, routinely studied as neural markers in attentional

blink tasks: i) P1 and N1 components in posterior, lateral electrodes (P7 and P8) between 40

to 140 ms and 90 to 160 ms, respectively, post T2 onset[90], ii) N2p component in

occipitoparietal electrodes (O1, O2, Oz, P3, P4, Pz and surrounding PO electrodes) between

150 to 300 ms post T2 onset[18,30,54,55], iii) P2 component in frontocentral electrodes

(C3,C4,Cz,F3,F4,Fz) between 150 to 300 ms post T2 onset[15,53], iv) P3 component in parietal

electrodes (P3,P4,Pz) between 300 to 550 ms post T2 onset[15,20,24,30]. We quantified the

change in ERP amplitudes during the attentional blink (see Methods section on "Statistical

analyses") and also performed partial correlations between the ERP amplitudes and

behavioral metrics (see Methods section on "Partial correlations between neural and

behavioral metrics"). All of the key ERP analyses were repeated without the 9-11 Hz bandstop

filter, and yielded results virtually identical to those reported in the main text and in SI Table

S2.

<u>Analysis of long-range synchronization.</u> We also quantified long-range synchronization

between the frontal and parietal cortex, an established neural marker of the attentional

blink[30,37]. For this we quantified the averaged spectral coherence between each pairs of

electrodes in the frontal (F3, F4, Fz), and parietal (P3, P4, Pz) cortex (9 pairs), respectively.

These analyses were performed with the data bandpass filtered between 0.1-40 Hz (FIR filter)

and without a 9-11 Hz bandstop filter. Coherence measures the degree of functional

connectivity at different frequencies between two regions and is quantified as the ratio of the

squared of the cross spectral density between the signals normalized by the product of the

individual signals' auto spectral densities.

$$876 \qquad C_{xy}(f) \ = \ \frac{|G_{xy}(f)|^2}{|G_{xx}(f)| \cdot |G_{yy}(f)|}$$

where $G_{xy}(f)$ denotes the cross spectral density between signals x and y whereas

$G_{xx}(f)$ and $G_{yy}(f)$ represent their, respective, auto spectral densities. We computed time

frequency representations for coherence (a "coherogram") with the *coherencyc* function in the

Matlab Chronux (v2.11) toolbox[91] with a time-half bandwidth product of 3 and 5 tapers

(TxW$_{\frac{1}{2}}$=3, K=5). Coherence values were computed in sliding windows of 300 ms duration with

a 50 ms stride computed from -200 ms before to +600 ms following T2 onset. Coherence

values were divisively normalized frequency-wise, on a per-trial basis, with coherence

computed in a baseline window from 0-300 ms prior to T2 onset. As with the ERPs, to remove

the contribution of T1-evoked responses, coherence values at each time point and frequency

of the T2-absent (correct rejection) trials were subtracted from the coherence values of the

T2-present trials; this subtraction was performed for each inter-target lag separately. To

quantify the effect of the attentional blink on the coherogram, we subtracted the coherograms

estimated from the short lag trials (100 and 300 ms) from the corresponding coherograms

from the long lag trials (700 and 900 ms). Subsequently, we conducted a cluster-based

permutation test on this difference coherogram to identify time periods and frequency bands

significantly modulated by the attentional blink (cluster forming threshold p<0.05). For

subsequent, partial correlation analyses of coherence with behavioral metrics and neural

distances (see next), we focused on a 300 ms time period (0-300 ms following T2 onset) and

high-beta frequency band (20-30 Hz) identified by the cluster-based permutation test (Fig. 5A-

C).

<u>Analysis of neural dimensions corresponding to detection and discrimination deficits</u>. To

understand how neural representations of the second target (T2) correlated with behavioral

deficits associated with the blink, we constructed distinct dimensions associated with the

detection and discrimination of T2. For this, mean EEG traces were computed from 33 bilateral

occipito-parietal electrodes (Fig. 6C-D) to form a multivariate activity representation (vector)

at each point in time. These time-varying vectors were subtracted against activity vectors on

T2 absent (correct rejection) trials, at each corresponding timepoint and across corresponding

inter-target lags, as before, to remove the contribution of the T1-evoked component.

We define the detection dimension (η_{det}) as the vector mean of each of the T2-present

activity vectors measured relative to T2-absent activity (Fig. 6A); this dimension reflects the

change in average EEG representation between T2 present and T2 absent trials. On the other

hand, the discrimination dimension (η_{dis}) represents the vector difference between the two T2-

present activity vectors (Fig. 6B); this dimension reflects the change in EEG representations

between the T2 CW and T2 CCW trials. Each of these dimensions were computed separately

for each inter-target lag; the magnitude (L2 norm) of each vector ($\|\eta_{det}\|$, $\|\eta_{dis}\|$) represents a

scalar quantifying the separability (inverse of the overlap) of the neural representations along

the detection or discrimination dimensions, for the respective lag. Note that these vector

magnitudes are equivalently computed as the Euclidean distance between the endpoints of

the mean vector representing each class of stimuli ("inter-class distance", Fig. 6C-D, left). To

mitigate biases arising from different numbers of trials across the different lags, we employed

bootstrap resampling: for each inter-target lag we randomly sampled a number of trials

corresponding to the lag with the minimum trial count with replacement and computed the

mean distances across 10 bootstrap estimates. For quantifying the effect of the attentional

blink and for correlation analyses, average inter-class neural distances were computed in a

window from 300 to 600 ms following T2 onset (Fig. 6C-D, right). These were then plotted for

each lag as a proportion of the inter-class distance for the largest lag (900 ms).

<u>Partial correlations between neural and behavioral metrics.</u> To quantify the neural correlates

of blink-induced detection and discrimination deficits, we performed bivariate correlations

between neural metrics (ERPs, inter-class distances and coherences) and the behavioral d'

for both tasks. The data were pooled across all lags and participants, and neural and

behavioral metrics were normalized for each participant by dividing each measure (neural or

behavioral) by the sum of that (respective) measure across all lags. Because the detection d'

and discrimination d' in our task were weakly correlated, we performed partial correlations,

rather than Pearson's correlations, to identify the specific neural correlate of each; these are

reported as "r_p" in the Results. To assess statistical significance of these coefficients, we

performed a non-parametric permutation test by shuffling the labels across lags independently

for each measure and participant and computing a null distribution over 1000 such random

permutations; the p-value reported is the proportion of r_p values in the null distribution that

exceeded the partial correlation coefficient observed in the real data. Given our a priori

hypothesis about the directionality of correlations, unless stated otherwise we report one-sided

p-values for these correlations; for instance, we expect a positive correlation between P3 ERP

amplitude and d' values with increasing inter-target lags. Partial correlations between neural

metrics (e.g., between inter-class distance and coherence, 7C-D) were also computed with a

similar procedure. Due to multiple comparisons, all p-values derived from permutation tests in

partial correlation were adjusted using the Bonferroni-Holm correction test.

Statistical analyses

For assessing the difference between short-lag and long-lag trials on psychometric or

psychophysical metrics (hit, miss, miss identification, false alarm and correct rejections, or d'

and c) or neural measures (e.g., ERP amplitudes, coherences, neural distances), non-

parametric Wilcoxon-signed rank tests were employed (*signrank* function in Matlab). Unless

stated otherwise, p-values in Wilcoxon-signed rank tests were computed using a normal

approximation (z-statistic); both the p-value and corresponding z-statistic are reported. All

tests were conducted as two-sided tests, unless otherwise specified. Additionally, we present

the Bayes factor (BF), which is a statistical metric comparing the likelihood of the alternative

hypothesis to that of the null hypothesis based on the observed data. The Bayes factor was

computed using the JASP toolbox[92]; all BFs were two-sided unless stated otherwise.

Goodness-of-fit of the model was assessed using a randomization test based on the χ^2-

statistic; the procedure is described in detail in previous work[45,93–96]. A small p-value (e.g.,

p<0.05) indicates that the model fit deviated significantly from the observations. Significance

testing for the attentional blink effect on sensitivity (d') and criterion (c) parameters was

performed with a 2-way ANOVA test (*anovan* function in Matlab). We employed robust (bend)

correlations (Robust Correlation Toolbox, v2. for conventional correlation analyses, and the

parcorr functon in Matlab for partial correlation analyses. Bonferroni-Holm correction tests

were employed through the *bonf_holm* function in MATLAB.

Data and code availability: Data and custom Matlab code to replicate the experimental

findings as well as the computational modeling results are available at the following link:

https://osf.io/gpzbe/?view_only=1140e43e0732451da64c7c9beaaea3d7

Acknowledgments

This research was supported by a Department of Biotechnology-Wellcome Trust India Alliance Intermediate fellowship, DST Swarna-Jayanti fellowship, a Department of Biotechnology-Indian Institute of Science Partnership Program grant, and an India-Trento Program for Advanced Research grant (all to DS).

Author Contributions

DS conceptualized the study. SH performed the experiments. SH and DR contributed novel analytic tools and analyzed the data. DS wrote the paper with inputs from all authors.

Declaration of Interest

Devarajan Sridharan is a research consultant at Google.

References

992   1.   Desimone, R. & Duncan, J. Neural Mechanisms of Selective Visual Attention. *Annu*
993   *Rev Neurosci* **18**, 193–222 (1995).
994   2.   Kanwisher Nancy.G and Potter. C. Mary. Repetition Blindness: Levels of Processing.
995   *J Exp Psychol* **16**, 30–47 (1990).
996   3.   Raymond, J. E., Shapiro, K. L. & Arnell, K. M. Temporary suppression of visual
997   processing in an RSVP task: an attentional blink? *J Exp Psychol Hum Percept Perform*
998   **18**, 849–60 (1992).
999   4.   Nieuwenstein, M. R., Potter, M. C. & Theeuwes, J. Unmasking the attentional blink. *J*
1000   *Exp Psychol Hum Percept Perform* **35**, 159–169 (2009).
1001   5.   Vul, E., Hanus, D. & Kanwisher, N. Delay of selective attention during the attentional
1002   blink. *Vision Res* **48**, 1902–1909 (2008).
1003   6.   Shapiro, K. L., Raymond, J. E. & Arnell, K. M. Attention to visual pattern information
1004   produces the attentional blink in rapid serial visual presentation. *J Exp Psychol Hum*
1005   *Percept Perform* **20**, 357–371 (1994).
1006   7.   Shapiro, K. L., Caldwell, J. & Sorensen, R. E. Personal names and the attentional
1007   blink: A visual 'cocktail party' effect. *J Exp Psychol Hum Percept Perform* **23**, 504–
1008   514 (1997).
1009   8.   Martens, S., Wolters, G. & van Raamsdonk, M. Blinks of the mind: Memory effects of
1010   attentional processes. *J Exp Psychol Hum Percept Perform* **28**, 1275–1287 (2002).
1011   9.   Chun, M. M. & Potter, M. C. A two-stage model for multiple target detection in rapid
1012   serial visual presentation. *J Exp Psychol Hum Percept Perform* **21**, 109–127 (1995).
1013   10.   Potter, M. C. *et al.* Bidirectional semantic priming in the attentional blink. *Psychon*
1014   *Bull Rev* **12**, 460–465 (2005).
1015   11.   Di Lollo, V., Kawahara, J., Shahab Ghorashi, S. M. & Enns, J. T. The attentional
1016   blink: Resource depletion or temporary loss of control? *Psychol Res* **69**, 191–200
1017   (2005).
1018   12.   Nieuwenstein, M. R. & Potter, M. C. Temporal Limits of Selection and Memory
1019   Encoding. *Psychol Sci* **17**, 471–475 (2006).
1020   13.   Kawahara, J.-I., Kumada, T. & di Lollo, V. The attentional blink is governed by a
1021   temporary loss of control. *Psychon Bull Rev* **13**, 886–890 (2006).
1022   14.   Zivony, A., Shanny, S. & Lamy, D. Perceptual Processing is Not Spared During the
1023   Attentional Blink. *J Cogn* **1**, 18 (2018).
1024   15.   Vogel, E. K., Luck, S. J. & Shapiro, K. L. Electrophysiological evidence for a
1025   postperceptual locus of suppression during the attentional blink. *J Exp Psychol Hum*
1026   *Percept Perform* **24**, 1656–1674 (1998).
1027   16.   Nieuwenstein, M. R., Chun, M. M., van der Lubbe, R. H. J. & Hooge, I. T. C. Delayed
1028   Attentional Engagement in the Attentional Blink. *J Exp Psychol Hum Percept Perform*
1029   **31**, 1463–1475 (2005).
1030   17.   Jolicœur, P. & Dell'Acqua, R. The Demonstration of Short-Term Consolidation. *Cogn*
1031   *Psychol* **36**, 138–202 (1998).
1032   18.   Dell'Acqua, R., Sessa, P., Jolicœur, P. & Robitaille, N. Spatial attention freezes during
1033   the attention blink. *Psychophysiology* **43**, 394–400 (2006).
1034   19.   Luck, S. J., Vogel, E. K. & Shapiro, K. L. Word meanings can be accessed but not
1035   reported during the attentional blink. *Nature* **383**, 616–618 (1996).
1036   20.   Sergent, C., Baillet, S. & Dehaene, S. Timing of the brain events underlying access to
1037   consciousness during the attentional blink. *Nat Neurosci* **8**, 1391–400 (2005).

21. Eimer, M. The neural basis of attentional control in visual search. *Trends Cogn Sci* **18**,
526–35 (2014).
22. Dellert, T. *et al.* Neural correlates of consciousness in an attentional blink paradigm
with uncertain target relevance. *Neuroimage* **264**, 119679 (2022).
23. Eiserbeck, A., Enge, A., Rabovsky, M. & Abdel Rahman, R. Distrust before first
sight? Examining knowledge- and appearance-based effects of trustworthiness on the
visual consciousness of faces. *Conscious Cogn* **117**, 103629 (2024).
24. Lasaponara, S., Dragone, A., Lecce, F., Di Russo, F. & Doricchi, F. The "serendipitous
brain": Low expectancy and timing uncertainty of conscious events improve awareness
of unconscious ones (evidence from the Attentional Blink). *Cortex* **71**, 15–33 (2015).
25. Vogel, E. K. & Luck, S. J. Delayed working memory consolidation during the
attentional blink. *Psychon Bull Rev* **9**, 739–743 (2002).
26. Dell'Acqua, R. *et al.* The Attentional Blink Impairs Detection and Delays Encoding of
Visual Information: Evidence from Human Electrophysiology. *J Cogn Neurosci* **27**,
720–735 (2015).
27. Chennu, S., Craston, P., Wyble, B. & Bowman, H. Attention Increases the Temporal
Precision of Conscious Perception: Verifying the Neural-ST2 Model. *PLoS Comput
Biol* **5**, e1000576 (2009).
28. Meijs, E. L., Slagter, H. A., de Lange, F. P. & van Gaal, S. Dynamic Interactions
between Top–Down Expectations and Conscious Awareness. *The Journal of
Neuroscience* **38**, 2318–2327 (2018).
29. Kranczioch, C., Debener, S. & Engel, A. K. Event-related potential correlates of the
attentional blink phenomenon. *Cognitive Brain Research* **17**, 177–187 (2003).
30. Kranczioch, C., Debener, S., Maye, A. & Engel, A. K. Temporal dynamics of access to
consciousness in the attentional blink. *Neuroimage* **37**, 947–955 (2007).
31. Batterink, L., Karns, C. M. & Neville, H. Dissociable Mechanisms Supporting
Awareness: The P300 and Gamma in a Linguistic Attentional Blink Task. *Cerebral
Cortex* **22**, 2733–2744 (2012).
32. Zivony, A. & Lamy, D. What processes are disrupted during the attentional blink? An
integrative review of event-related potential research. *Psychon Bull Rev* **29**, 394–414
(2022).
33. Marois, R., Chun, M. M. & Gore, J. C. Neural Correlates of the Attentional Blink.
Neuron **28**, 299–308 (2000).
34. Cooper, A. C. G., Humphreys, G. W., Hulleman, J., Praamstra, P. & Georgeson, M.
Transcranial magnetic stimulation to right parietal cortex modifies the attentional
blink. *Exp Brain Res* **155**, 24–29 (2004).
35. Kihara, K. *et al.* The role of left and right intraparietal sulcus in the attentional blink: a
transcranial magnetic stimulation study. *Exp Brain Res* **178**, 135–140 (2007).
36. Williams, M. A., Visser, T. A. W., Cunnington, R. & Mattingley, J. B. Attenuation of
Neural Responses in Primary Visual Cortex during the Attentional Blink. *The Journal
of Neuroscience* **28**, 9890–9894 (2008).
37. Gross, J. *et al.* Modulation of long-range neural synchrony reflects temporal
limitations of visual attention in humans. *Proceedings of the National Academy of
Sciences* **101**, 13050–13055 (2004).
38. Harris, J. A., McMahon, A. R. & Woldorff, M. G. Disruption of Visual Awareness
during the Attentional Blink Is Reflected by Selective Disruption of Late-stage Neural
Processing. *J Cogn Neurosci* **25**, 1863–1874 (2013).
39. Green, D. M. , & Swets, J. A. *Signal Detection Theory and Psychophysics.* (1966).

40.   Luo, T. Z. & Maunsell, J. H. R. Neuronal Modulations in Visual Cortex Are
Associated with Only One of Multiple Components of Attention. *Neuron* **86**, 1182–8
(2015).

41.   Sridharan, D., Steinmetz, N. A., Moore, T. & Knudsen, E. I. Does the Superior
Colliculus Control Perceptual Sensitivity or Choice Bias during Attention? Evidence
from a Multialternative Decision Framework. *The Journal of Neuroscience* **37**, 480–
511 (2017).

42.   Asplund, C. L., Fougnie, D., Zughni, S., Martin, J. W. & Marois, R. The Attentional
Blink Reveals the Probabilistic Nature of Discrete Conscious Perception. *Psychol Sci*
25, 824–831 (2014).

43.   Caetta Florent, G. A. Upshifted decision criteria in attentional blink and repetition
blindness. *Visual Cognition* **18**, 413–433 (2010).

44.   DeCarlo, L. T. On a signal detection approach to -alternative forced choice with bias,
with maximum likelihood and Bayesian approaches to estimation. *J Math Psychol* **56**,
196–207 (2012).

45.   Sridharan, D., Steinmetz, N. A., Moore, T. & Knudsen, E. I. Distinguishing bias from
sensitivity effects in multialternative detection tasks. *J Vis* **14**, 16–16 (2014).

46.   Hautus, M. J., Macmillan, N. A. & Creelman, C. D. *Detection Theory*. (Routledge,
New York, 2021). doi:10.4324/9781003203636.

47.   Parker, A. J. & Newsome, W. T. SENSE AND THE SINGLE NEURON: Probing the
Physiology of Perception. *Annu Rev Neurosci* **21**, 227–277 (1998).

48.   Funahashi, S. Working Memory in the Prefrontal Cortex. *Brain Sci* **7**, 49 (2017).

49.   Zhang, Y., Bi, K., Li, J., Wang, Y. & Fang, F. Dyadic visual perceptual learning on
orientation discrimination. *Current Biology* **33**, 2407-2416.e4 (2023).

50.   Dupont, P. *et al.* Regions in the human brain activated by simultaneous orientation
discrimination: a study with positron emission tomography. *European Journal of*
Neuroscience **10**, 3689–3699 (1998).

51.   Maunsell, J. H. R. Neuronal Mechanisms of Visual Attention. *Annu Rev Vis Sci* **1**,
373–391 (2015).

52.   Vrieze, S. I. Model selection and psychological theory: A discussion of the differences
between the Akaike information criterion (AIC) and the Bayesian information criterion
(BIC). *Psychol Methods* **17**, 228–243 (2012).

53.   Akyürek, E. G., Leszczyński, M. & Schubö, A. The temporal locus of the interaction
between working memory consolidation and the attentional blink. *Psychophysiology*
47, no-no (2010).

54.   Jolicœur, P., Sessa, P., Dell'Acqua, R. & Robitaille, N. Attentional control and capture
in the attentional blink paradigm: Evidence from human electrophysiology. *European*
Journal of Cognitive Psychology **18**, 560–578 (2006).

55.   Jolicœur, P., Sessa, P., Dell'Acqua, R. & Robitaille, N. On the control of visual spatial
attention: evidence from human electrophysiology. *Psychol Res* **70**, 414–424 (2006).

56.   Sheahan, H., Luyckx, F., Nelli, S., Teupe, C. & Summerfield, C. Neural state space
alignment for magnitude generalization in humans and recurrent networks. *Neuron*
109, 1214-1226.e8 (2021).

57.   Chung, S. & Abbott, L. F. Neural population geometry: An approach for
understanding biological and artificial neural networks. *Curr Opin Neurobiol* **70**, 137–
144 (2021).

58.   Broadbent, D. E. & Broadbent, M. H. From detection to identification: response to
multiple targets in rapid serial visual presentation. *Percept Psychophys* **42**, 105–13
(1987).

59.   Hillis, J. M. & Brainard, D. H. Distinct Mechanisms Mediate Visual Detection and
Identification. *Current Biology* **17**, 1714–1719 (2007).
60.   Straube, S. & Fahle, M. Visual detection and identification are not the same: Evidence
from psychophysics and fMRI. *Brain Cogn* **75**, 29–38 (2011).
61.   Koivisto, M., Grassini, S., Salminen-Vaparanta, N. & Revonsuo, A. Different
Electrophysiological Correlates of Visual Awareness for Detection and Identification.
J Cogn Neurosci **29**, 1621–1631 (2017).
62.   Wiens, S., Andersson, A. & Gravenfors, J. Neural electrophysiological correlates of
detection and identification awareness. *Cogn Affect Behav Neurosci* **23**, 1303–1321
(2023).
63.   Correa, Á., Lupiáñez, J., Milliken, B. & Tudela, P. Endogenous temporal orienting of
attention in detection and discrimination tasks. *Percept Psychophys* **66**, 264–278
(2004).
64.   Lubbe, R. H. J. Van der, Vogel, R. O. & Postma, A. Different Effects of Exogenous
Cues in a Visual Detection and Discrimination Task: Delayed Attention Withdrawal
and/or Speeded Motor Inhibition? *J Cogn Neurosci* **17**, 1829–1840 (2005).
65.   Treue, S. & Maunsell, J. H. R. Attentional modulation of visual motion processing in
cortical areas MT and MST. *Nature* **382**, 539–541 (1996).
66.   Luck, S. J. *et al.* Effects of spatial cuing on luminance detectability: Psychophysical
and electrophysiological evidence for early selection. *J Exp Psychol Hum Percept*
Perform **20**, 887–904 (1994).
67.   McAdams, C. J. & Maunsell, J. H. R. Effects of Attention on Orientation-Tuning
Functions of Single Neurons in Macaque Cortical Area V4. *The Journal of*
Neuroscience **19**, 431–441 (1999).
68.   Navalpakkam, V. & Itti, L. Search Goal Tunes Visual Features Optimally. *Neuron* **53**,
605–617 (2007).
69.   Kanashiro, T., Ocker, G. K., Cohen, M. R. & Doiron, B. Attentional modulation of
neuronal variability in circuit models of cortex. *Elife* **6**, (2017).
70.   Averbeck, B. B., Latham, P. E. & Pouget, A. Neural correlations, population coding
and computation. *Nat Rev Neurosci* **7**, 358–366 (2006).
71.   Rotermund, D., Taylor, K., Ernst, U. A., Kreiter, A. K. & Pawelzik, K. R. Attention
Improves Object Representation in Visual Cortical Field Potentials. *The Journal of*
Neuroscience **29**, 10120–10130 (2009).
72.   Sy, J. L., Miao, H.-Y., Marois, R. & Tong, F. Conscious perception can be both graded
and discrete. *J Exp Psychol Gen* **150**, 1461–1475 (2021).
73.   Karabay, A. *et al.* Two faces of perceptual awareness during the attentional blink:
Gradual and discrete. *J Exp Psychol Gen* **151**, 1520–1541 (2022).
74.   Grill-Spector, K. & Kanwisher, N. Visual Recognition. *Psychol Sci* **16**, 152–160
(2005).
75.   Martens, S., Elmallah, K., London, R. & Johnson, A. Cuing and stimulus probability
effects on the P3 and the AB. *Acta Psychol (Amst)* **123**, 204–218 (2006).
76.   Reed, C. L. *et al.* Selective contributions of executive function ability to the P3.
International Journal of Psychophysiology **176**, 54–61 (2022).
77.   Freedman, D. J., Riesenhuber, M., Poggio, T. & Miller, E. K. Categorical
Representation of Visual Stimuli in the Primate Prefrontal Cortex. *Science (1979)* **291**,
312–316 (2001).
78.   Antzoulatos, E. G. & Miller, E. K. Synchronous beta rhythms of frontoparietal
networks support only behaviorally relevant representations. *Elife* **5**, (2016).
79.   Murthy, V. N. & Fetz, E. E. Coherent 25- to 35-Hz oscillations in the sensorimotor
cortex of awake behaving monkeys. *Proc Natl Acad Sci U S A* **89**, 5670–4 (1992).

80.   Anton-Erxleben, K. & Carrasco, M. Attentional enhancement of spatial resolution:
linking behavioural and neurophysiological evidence. *Nat Rev Neurosci* **14**, 188–200
(2013).
81.   Fook K.Chua. The effect of target contrast on the attentional blink. *Percept*
Psychophys **5**, 770–788 (2005).
82.   Dux, P. E. & Marois, R. The attentional blink: A review of data and theory. *Atten*
Percept Psychophys **71**, 1683–1700 (2009).
83.   Simione, L. *et al.* ViSA: A neurodynamic model for visuo-spatial working memory,
attentional blink, and conscious access. *Psychol Rev* **119**, 745–769 (2012).
84.   Jolicoeur, P. Concurrent response-selection demands modulate the attentional blink. . *J*
Exp Psychol Hum Percept Perform **25**, 1097 (1999).
85.   Jolicoeur, P. Modulation of the attentional blink by on-line response selection:
Evidence from speeded and unspeeded Task1 decisions. *Mem Cognit* **26**, 1014–1032
(1998).
86.   Gold, J. I. & Shadlen, M. N. The neural basis of decision making. *Annu Rev Neurosci*
30, 535–74 (2007).
87.   M Kleiner, D. B. D. P. A. I. R. M. C. B. What's new in psychtoolbox-3. *Perception* **36**,
1–16 (2007).
88.   Nakatani, C., Baijal, S. & van Leeeuwen, C. Curbing the attentional blink: Practice
keeps the mind's eye open. *Neurocomputing* **84**, 13–22 (2012).
89.   Junghofer, M., Elbert, T., Tucker, D. M. & Rockstroh, B. Statistical control of artifacts
in dense array EEG/MEG studies. *Psychophysiology* **37**, 523–532 (2000).
90.   Chmielewski, W. X., Mückschel, M., Dippel, G. & Beste, C. Concurrent information
affects response inhibition processes via the modulation of theta oscillations in
cognitive control networks. *Brain Struct Funct* **221**, 3949–3961 (2016).
91.   Bokil, H., Andrews, P., Kulkarni, J. E., Mehta, S. & Mitra, P. P. Chronux: A platform
for analyzing neural signals. *J Neurosci Methods* **192**, 146–151 (2010).
92.   JASP Team. JASP (version 0.19.0.) [Computer Software]. Preprint at (2022).
93.   Mesulam, M.-M. Spatial attention and neglect: parietal, frontal and cingulate
contributions to the mental representation and attentional targeting of salient
extrapersonal events. *Philos Trans R Soc Lond B Biol Sci* **354**, 1325–1346 (1999).
94.   Banerjee, S., Grover, S., Ganesh, S. & Sridharan, D. Sensory and decisional
components of endogenous attention are dissociable. *J Neurophysiol* **122**, 1538–1554
(2019).
95.   Sreenivasan, V. & Sridharan, D. Subcortical connectivity correlates selectively with
attention's effects on spatial choice bias. *Proceedings of the National Academy of*
Sciences **116**, 19711–19716 (2019).
96.   Sagar, V., Sengupta, R. & Sridharan, D. Dissociable sensitivity and bias mechanisms
mediate behavioral effects of exogenous attention. *Sci Rep* **9**, 12657 (2019).

1224

1225

1226

1227

1228

1229

1230

Figures and Legends

1232

1233

1234

Figure 1. Decoupling behavioral components and neural bases of the attentional blink.

(Left) Identifying the precise origin of accuracy deficits induced by the attentional blink. T2 identification accuracies at different inter-target (T1-T2) lags, in a conventional attentional blink task. x-axis: inter-target lag in milliseconds; y-axis: T2 identification accuracy (%). Red horizontal line: asymptotic T2 identification accuracy for long inter-target lags; red vertical arrows: accuracy deficit with T2 identification for short inter-target lags (attentional blink).

(Right, top) The identification deficit could reflect impaired detection of T2's presence which, in turn, could arise either from a detection sensitivity deficit (upper row) or a detection bias (criterion) deficit (lower row). Gray and white Gaussians: decision variable distributions corresponding to signal (target present) and noise (target absent), respectively. Black vertical line: criterion for deciding between target present and absent.

(Right, bottom) The identification deficit could also reflect impaired discrimination of T2's features (e.g. orientation), which, again, could arise either from a discrimination sensitivity deficit (upper row) or a discrimination bias (criterion) deficit (lower row). Purple and orange Gaussians: decision variable distributions corresponding to a counterclockwise (CW) and

clockwise (CCW) gratings, respectively. Black vertical line: criterion for deciding between

target features (clockwise and counterclockwise orientation).

*Brain schematics (**rightmost column**): Distinct neural markers of each subcomponent –*

detection (top) or discrimination (bottom) -- of attentional blink deficits.

Figure 2. Novel task design to distinguish subcomponents of the attentional blink.

A. *Schematic of the attentional blink task. Stimuli were presented in a rapid serial visual presentation (RSVP) paradigm at a 10 Hz rate (70 ms onset, 30 ms offset). Following fixation, plaid gratings appeared for a variable interval (200-1200 ms, geometrically distributed), followed by the first target (T1): a low spatial frequency grating (100 ms). After this a series of plaid gratings appeared for variable intervals (100, 300, 500, 700, and 900 ms; geometric distribution) followed by the appearance of the second target (T2): a high*

spatial frequency grating (100 ms). Following T2, plaid gratings were presented for a fixed interval (600 ms). Finally, in the response epoch, participants reported T1's orientation as being closer to the cardinal or diagonal axes (two-alternative), and then reported T2's orientation as being clockwise or counterclockwise of vertical, or absent (three-alternative). All plaids were encircled by a circular placeholder. The fixation dot and the placeholder were present on the screen throughout the trial.

B. Psychometric function of accuracy (% correct) for T2 detection with increasing inter target (T1-T2) lags, for trials in which T1 was reported correctly (n=24 participants). Filled circles and solid lines: average accuracy for high contrast T2 gratings; open circles and dashed lines: average accuracy for low contrast T2 gratings. Error bars: s.e.m. Asterisks: significance levels for comparing accuracies between short (100 and 300 ms) and long (700 and 900 ms) lag trials; solid and dashed brackets: comparisons for high and low contrast gratings respectively. *p<0.05, **p<0.01, ***p<0.001 and n.s.: not significant.

C. Same as in panel B, but showing the psychometric function of accuracy for T2 discrimination with increasing inter target (T1-T2) lags (n=24). Other conventions are the same as in panel B, except that markers and lines are depicted in orange colour.

D. Stimulus-response contingency table for the 3-alternative T2 decision. Rows represent the three possible T2 stimulus events: clockwise orientation (CW, orange), counterclockwise orientation (CCW, purple) or absent (none, gray). Columns represent three possible choices: clockwise (CW), counterclockwise (CCW) or absent (none). The table depicts the nine stimulus-response contingencies: two each of hit rates (H), misidentification rates (MI), miss rates (M), false alarm rates (FA) – one for each orientation (CW/CCW) – and one correct rejection rate (CR).

E. Same as in panel B, but showing psychometric function of average hit rates.

F. Same as in panel B, but showing psychometric function of average misidentification rates.

G. Same as in panel B, but showing psychometric function of average miss rates.

(E-G). Other conventions are the same as in panel B except that markers and lines are denoted in black color.

1314

1315

Figure 3. A novel psychophysical model decouples attentional blink subcomponents.

A. Signal detection model for estimating T2 sensitivity and bias. The decision variable is a

bivariate Gaussian (Ψ) whose components, Ψ_det (ordinate) and Ψ_dis (abscissa), encode

sensory evidence for stimulus presence and orientation, respectively. Black circle: Ψ

distribution for T2 absent trials (noise distribution), is centered on the origin. Orange and

purple circles: Ψ distributions for clockwise and counterclockwise T2 orientations,

respectively (signal distributions). Solid and dashed outlines: High and low contrast T2,

respectively. Vertical gray and horizontal orange lines (double headed arrows): detection

sensitivity (d'_det) and discrimination sensitivity (d'_dis) for high contrast T2, respectively.

Dashed gray lines: Signal mean projections onto the detection and discrimination axes.

B. Decision surface with linear decision boundaries (thick black lines) demarcates 3 decision

zones, one for each potential 3AFC choice: Clockwise T2 (CW, orange shading),

counterclockwise T2 (CCW, purple shading), no T2 (None, gray shading). The decision

surface is parameterized by: i) a discrimination criterion (c_dis), governing the horizontal

position of the decision surface (horizontal double arrowhead), ii) a detection threshold (t_{det}), governing the vertical position of the decision surface (vertical double arrowhead), and iii) the angle between the two oblique decision boundaries (β). Other conventions are the same as in panel A.

C. Psychophysical function of detection sensitivity (d'_{det}) with increasing inter target (T1-T2) lags. Other conventions are the same as in Figure 2A.

D. Same as in panel C but showing the psychophysical function of discrimination sensitivity (d'_{dis}). Model selection yielded identical psychophysical functions for high and low contrast T2 (Methods). Other conventions are the same as in Figure 2B.

E. Same as in panel C but showing the psychophysical function of detection criterion (c_{det}). Dashed horizontal line: $c_{det}=0$.

F. Discrimination criterion (c_{dis}). Model selection constrained c_{dis} to be equal across lags (Methods). Other conventions are the same as in panels D-E.

G. Modulation index (MI) of the discrimination blink for low (open plot) and high (filled plot) detection blink MI blocks. Box plots, center line: median; box limits: upper and lower quartiles; whiskers: 1.5x the interquartile range. Violin plots: kernel density estimates.

(C-G) Asterisks: *$p<0.05$, **$p<0.01$, ***$p<0.001$; and n.s.: not significant.

1358

1359

Figure 4. N2p and P3 event related potentials signal detection sensitivity deficits.

A. *(Left) The event related potential (ERP, n=18 participants) showing the N2p component in occipitoparietal electrodes (inset), locked to T2 onset (dashed vertical line). Bright green, dull green, dark green, light gray and black traces: Average ERPs for the five inter-target lags – 100, 300, 500, 700 and 900 ms – respectively. Shading: s.e.m. Gray vertical shading: Time epoch for N2p amplitude quantification. ERPs were computed by subtracting the average ERPs on correct rejection trials (Methods). (Right) Violin plots showing the distribution of the peak N2p amplitudes across participants separately for the short (green: 100+300 ms) and the long (gray: 700+900 ms) lag trials. Asterisks: *p<0.05, **p<0.01, and ***p<0.001. Other conventions are the same as in Figure 3G.*

B. *(Left) Same as in panel A (left) but showing the P3 ERP component in the parietal*

electrodes (inset). (Right) Same as in panel A (left) but showing the quantification of the

P3 ERP. Other conventions are the same as in panel A.

C. *Variation of detection sensitivity (d'_{det}; y-axis) with N2p peak amplitude (x-axis) across*

inter-target lags (circles). r and p denote the robust correlation coefficient and permutation

test p-value, respectively (Methods). Dashed line: linear fit. Error bars: s.e.m.

D. *(Top) Partial correlation between N2p peak amplitude (x-axis, amplitude residual) and*

detection sensitivity (y-axis, d'_{det} residual) while controlling for the discrimination sensitivity

(d'_{dis}). Each of the five shapes – filled triangle, diamond, square, pentagon, circle –

represents one inter-target lag (legend). r_p and p denote the partial correlation coefficient

and permutation test p-value, respectively (Methods). Solid line: Linear fit; dashed curves:

95% confidence intervals. (Bottom) Same as in the top panel but showing partial

correlation between N2p peak amplitude (x-axis, amplitude residual) and discrimination

sensitivity (y-axis, d'_{dis} residual) while controlling for detection sensitivity (d'_{det}).

E. *Same as in panel C but showing variation of discrimination sensitivity (d'_{dis}, y-axis) with P3*

peak amplitude (x-axis) across inter-target lags.

F. *Same as in panel D but showing the partial correlation of P3 peak amplitude with d'_{det} while*

controlling for d'_{dis} (top), or, conversely, with d'_{dis} while controlling for d'_{det} (bottom).

(E-F) Other conventions are the same as in panels C-D, respectively.

1390

1391

1392

1393

1394

1395

1396

1397

1398

1402

Figure 5. High-beta fronto-parietal coherence signals discrimination sensitivity deficits.

A. *Coherence between frontal and parietal electrodes (Methods) as a function of time (x-axis) and frequency (y-axis) locked to T2 onset (t=0) (n=18 participants), normalized frequency-wise by a baseline mean (gray horizontal bar), and computed by subtracting the average coherograms for correct rejection trials. Cooler colors: higher coherence. Dashed horizontal line: low-beta (13 to 19 Hz) and high-beta (20 to 30 Hz) sub-bands. (Left and middle) Coherograms for the short (100+300 ms) and long (700+900 ms) lag trials,*

respectively. *(Right) Difference coherogram. Bold black outline: statistically significant coherence difference between short and long lag trials (cluster forming threshold p<0.05).*

B. *Bilateral fronto-parietal coherence values in the high-beta band across participants, separately for the short (100+300 ms, green) and long (700+900 ms, gray) lag trials. Topoplot: schematic of electrode pairs (red circles and black arrows) used for computing bilateral frontoparietal coherence (Methods). Other conventions are the same as in Figure 3G.*

C. *(Left) Same as in panel A (right panel) but showing the difference coherogram for the left hemispheric frontoparietal electrodes. Other conventions are the same as in panel A. (Right) Same as in panel B but showing high-beta coherence values over the left hemispheric fronto-parietal electrodes. Other conventions are the same as in panel B and Figure 3G.*

D. *(Left and right) Same as in panel C but showing the difference coherogram for the right hemispheric fronto-parietal electrodes. Other conventions are the same as in panel C. (B-D). Asterisks: *p<0.05, **p<0.01, and ***p<0.001*

E. *Same as Figure 4E but showing the variation of discrimination sensitivity (d'_{dis}, y-axis) with left fronto-parietal high-beta coherence (x-axis) across inter-target lags. Other conventions are the same as in Figure 4E.*

F. *Same as in Figure 4F but showing the partial correlation of high-beta left fronto-parietal coherence with d'_{det} while controlling for d'_{dis} (top), or, conversely, with d'_{dis} while controlling for d'_{det} (bottom). Other conventions are the same as in Figure 4F.*

Figure 6. Distinct neural dimensions encode detection and discrimination bottlenecks.

A. Schematic showing the "detection" dimension, η_{det} (y-axis): a linear dimension in a multidimensional electrode space along which neural activity encodes the presence versus absence of T2 (Methods). Gray, orange and purple dot clusters: distribution of EEG activity for T2 absent, T2 clockwise (CW) and counterclockwise (CCW) trials, respectively. Darker (lighter) shades denote longer (shorter) inter-target lags.

B. *Same as in panel A but showing the "discrimination" dimension, η_{dis} (x-axis): a linear dimension along which neural activity encodes the orientation of T2 (clockwise versus counterclockwise of vertical).*

C. *(Left) Time evolution of the average inter-class distance along the detection dimension ($\|\eta_{det}\|$) locked to T2 onset (n=18 participants). Values at each time point are plotted relative to the longest lag (900 ms). Bright green, dark green and black traces: short (100+300 ms) intermediate (500 ms) and long (700 ms) lags, respectively. Shading: s.e.m. Dashed vertical line: T2 onset. Gray shading: Time epoch for neural distance quantification and statistical comparison. (Right) Distribution of η_{det} inter-class distances across participants separately for the short (green, 100+300 ms) and the long (gray, 700+900 ms) lag trials. Asterisks: *p<0.05, **p<0.01, and ***p<0.001. Other conventions are the same as in Figure 3G.*

D. *Same as in panel C but showing the time evolution of the average neural inter-class distance along the discrimination dimension ($\|\eta_{dis}\|$) (left), and the distribution of neural distances along the discrimination dimension (right). Other conventions are the same as in panel C.*

E. *Same as in Figure 4C, but showing variation of detection sensitivity (d'$_{det}$, y-axis) with neural distance along the detection dimension ($\|\eta_{det}\|$, x-axis) across inter-target lags.*

F. *Same as in Figure 4D but showing the partial correlation of $\|\eta_{det}\|$ with d'$_{det}$ while controlling for d'$_{dis}$ (top), or, conversely, with d'$_{dis}$ while controlling for d'$_{det}$ (bottom).*

G. *Same as in Figure 4E, but showing variation of discrimination sensitivity (d'$_{dis}$, y-axis) with neural distance along the discrimination dimension ($\|\eta_{dis}\|$, x-axis) across inter-target lags.*

H. *Same as in Figure 4F but showing the partial correlation of $\|\eta_{dis}\|$ with d'$_{det}$ while controlling for d'$_{dis}$ (top), or, conversely, with d'$_{dis}$ while controlling for d'$_{det}$ (bottom).*

1472

Figure 7. Neural dimensions of detection and discrimination deficits correlate with their

respective EEG markers.

A. Same as in Figure 4C, but showing variation of the neural distance along the detection

dimension ($\|\eta_{det}\|$, y-axis) with P3 amplitude (d'_{det}, x-axis) across inter-target lags.

B. Same as in Figure 4D but showing the partial correlation of the P3 amplitude with $\|\eta_{det}\|$

while controlling for $\|\eta_{dis}\|$ (top), or, conversely, with $\|\eta_{dis}\|$ while controlling for $\|\eta_{det}\|$

(bottom).

(A-B) Other conventions are the same as in Figures 4C-D.

C. Same as in Figure 4E, but showing variation of the neural distance along the discrimination

dimension ($\|\eta_{dis}\|$, y-axis) with left high beta frontoparietal coherence (x-axis) across the 5

inter-target lags.

D. Same as in Figure 4F but showing the partial correlation of the left high beta frontoparietal

coherence with $\|\eta_{det}\|$ while controlling for $\|\eta_{dis}\|$ (top), or, conversely, with $\|\eta_{dis}\|$ while

controlling for $\|\eta_{det}\|$ (bottom).

(C-D) Other conventions are the same as in Figures 4E-F.

1488

1489

1490

Figure 8: Distinct neural bases of subcomponents of the attentional blink

The attentional blink selectively impairs a specific component of attention – perceptual

sensitivity (d') – and produces both detection (top, left) and discrimination (top, right) deficits.

Detection d' deficits – deficits with distinguishing the presence versus absence of the second

target (T2) – are correlated with reduced amplitudes of N2p and P3 ERPs (gray shading, left

top). They are also accompanied by a representational collapse along the detection

dimension (gray shading, left bottom). By contrast, discrimination d' deficits – deficits with

discriminating T2's orientation – are evidenced by reduced left fronto-parietal beta

coherence (red shading, left top) and a representational collapse along the discrimination

dimension (red shading, left bottom).

1503

1504

1505

1506

Supplementary Figures and Tables

A

Model #	# Model parameters	Constraints
Model I	35	No constraints
Model II	31	Constant beta (lags)
Model III	22	Constant beta (lags), Constant C_{dis} (lags and contrast), Constant d'_{dis} (contrast)
Model IV	22	Constant beta (lags), Constant C_{det} (lags and contrast), Constant d'_{det} (contrast)
Model V	13	Constant beta (lags), Constant C_{det} (lags and contrast), Constant d'_{det} (contrast), Constant C_{dis} (lags and contrast), Constant d'_{dis} (contrast)

B

C

1508

Supplementary Figure S1. Model comparison analysis.

A. Summary of the 5 models used in the analysis (see Methods for details).

B. Distribution of goodness of fit (p-values) with a randomization test based on the chi-squared statistic for all five models (model I – V, n=24 participants). Other conventions for the violin plot same as in Figure 3G (main text).

C. Distribution of Akaike information criterion (AIC) values for all five models. Other conventions are the same as in panel B.

D. Same as panel C but showing the distribution of Bayesian information criterion (BIC) values for all five models. Other conventions are the same as in panel C.

1519

1520

Supplementary Figure S2. Correlation between detection and discrimination blink.

A. Correlation between the modulation indices (MI) of detection and discrimination sensitivity.

The MI is calculated as MI-d'_{det} = (d'^{LL}_{det} - d'^{SL}_{det})/ (d'^{LL}_{det} + d'^{SL}_{det}) (SL: short lag, LL: long

lag; Methods)**.** Gray circles: individual participants. Dashed line: Linear fit. r and p values:

correlation coefficient and its significance level based on robust correlations (Methods).

B. Same as in panel A except showing the correlation between MIs for detection and

discrimination accuracy. Other conventions are the same as in panel A.

1528

1529

1530

1531

1532

1533

Supplementary Figure S3. P1 ERP component in parietal electrodes.

(Left) The P1 event related potential (ERP) in the bilateral parietal electrodes (inset, topoplot),

locked to T2 grating onset (n=18 participants). Bright green, dull green, dark green, light gray

and black traces: Average ERPs for the five inter-target lags (100, 300, 500, 700 and 900 ms),

respectively. Shading: s.e.m. Gray vertical bar: Time epoch considered for quantifying the P1

component amplitude. Dashed vertical black line: T2 onset. Here, and elsewhere, ERPs were

computed by subtracting the average ERPs for correct rejection trials (Methods). (Right) Violin

plots showing the distribution of the peak P1 amplitudes across participants separately for the

short (green; 100+300 ms) and the long (gray, 700+900 ms) lag conditions. Asterisks denote

significance differences: *p<0.05, **p<0.01, and ***p<0.001. Other conventions for the violin

plot are the same as in Figure 3G.

1545

Supplementary Figure S4. Linking neural dimensions with neural markers of detection deficits – N2p peak amplitude.

A. Variation of the neural distance along the detection dimension ($\|\eta_{det}\|$, y-axis) with N2p amplitude (d'_{det}, x-axis) across the 5 inter-target lags (circles in distinct shades of gray). Dashed curve: linear fit. Error bars: s.e.m along the respective axis.

B. (Top) Partial correlation of the N2p peak amplitude (x-axis, amplitude residual) with $\|\eta_{det}\|$ (y-axis, detection distance residual) while controlling for $\|\eta_{dis}\|$. The five shapes – filled triangle, diamond, square, pentagon, circle – represent the five inter-target lags – 100, 300, 500, 700 and 900 ms – respectively (data pooled across n=18 participants). r_p and p denote the partial correlation value and p-value calculated with a permutation test (Methods). Solid line: Linear fit; black dashed curves: 95% confidence intervals. (Bottom) Same as in the top panel but showing the partial correlation between N2p peak amplitude (x-axis, amplitude residual) with $\|\eta_{dis}\|$ (y-axis, discrimination distance residual) while controlling for $\|\eta_{det}\|$. Other conventions are the same as in the top panel.

1564 **Supplementary Table S1. Attentional blink effect on psychometric measures.**

1565 2-way ANOVA analysis with the psychometric measures as dependent variables and lags and

1566 contrasts as independent factors.

Psychometric measures	Analysis of Variance (ANOVA)		
	Main effect of lag (F, p, BF)	Main effect of contrast (F, p, BF)	Interaction effect (F, p, BF)
Hit	**55.78, <0.001, >10³**	**35.52, <0.001, >10²**	2.23, 0.149 (n.s.), 0.538
Miss	**10.96, 0.003, 20**	**21.6, <0.001, >10²**	2.31, 0.142 (n.s), 0.597
Misidentification	**42.45, <0.001, >10³**	3.18, 0.090 (n.s.), 0.367	0.02, 0.889 (n.s), 0.276
FA	**6.9, 0.015, >10³**	N/A	N/A
CR	**6.9, 0.015, >10³**	N/A	N/A

1567

1568

Supplementary Table S2. Partial correlations of ERP amplitudes with detection and

discrimination sensitivity.

Magnitude of blink induced deficit in ERP amplitudes, and partial correlation between ERP

amplitudes, and detection (discrimination) d', while controlling for the confounding effect of

discrimination (detection) d' (Methods). **Bold** entries correspond to significant partial

correlations.

1575

ERP components	Blink deficit (mean ± s.e.m)	Partial correlation with sensitivity (d')	
		Partial correlation with detection d'	Partial correlation with discrimination d'
P1 (parietal)	0.25 ± 0.16, p = 0.231	r_p = -0.06 p = 0.935	r_p = -0.11 p = 0.816
P2 (frontocentral)	0.19 ± 0.07, p = 0.021	r_p = 0.05 p = 0.999	r_p = 0.23 p = 0.120
N2p (occipitoparietal)	**-0.47 ± 0.12, p = 0.003**	$\mathbf{r_p}$ **= -0.34 p < 0.001**	r_p = -0.01 p = 0.970
P3 (parietal)	**0.45 ± 0.09, p < 0.001**	$\mathbf{r_p}$ **= 0.30 p < 0.001**	r_p = 0.14 p = 0.748

1576

1577

Supplementary Table S3. Partial correlation of fronto-parietal coherence with detection

and discrimination sensitivity.

Partial correlation between left and right frontoparietal beta coherence and detection and

discrimination sensitivity. Other conventions are the same as in SI Table S2.

Hemisphere	Beta band	Partial correlation with sensitivity (d')	
		Partial correlation with detection d'	Partial correlation with discrimination d'
Left fronto-parietal	High beta (20-30 Hz)	r_p = -0.05 p = 0.316	**r_p = 0.22 p = 0.018**
	Low beta (13-19 Hz)	r_p = 0.05 p = 0.728	r_p = -0.03 p = 0.728
Right fronto-parietal	High beta (20-30 Hz)	r_p = -0.01 p = 0.470	r_p = 0.21 p = 0.1
	Low beta (13-19 Hz)	r_p = -0.04 p = 0.680	r_p = 0.05 p = 0.682

Supplementary Table S4. Partial correlation between neural measures and neural distances in the detection and discrimination dimensions.

Partial correlation between neural measures (N2P and P3 ERP amplitudes and left frontoparietal high-beta coherence) and neural distances in detection (discrimination) dimension while controlling for the confounding effect of the neural distance in the discrimination (detection) dimension.

Neural measure	Partial correlation with neural distance (η)	
	Neural distance in the detection dimension ($\|\eta_{det}\|$)	Neural distance in the discrimination dimension ($\|\eta_{dis}\|$)
N2p amplitude	$r_p = -0.31, p = 0.02$	$r_p = 0.07, p = 0.330$
P3 amplitude	$r_p = 0.46, p < 0.001$	$r_p = 0.15, p = 0.160$
Left fronto-parietal high-beta (20-30 Hz) coherence	$r_p = 0.07, p = 0.255$	$r_p = 0.26, p = 0.039$