Impaired central pattern generators due to abnormal EPHA4 signaling leads to idiopathic scoliosis

Authors:

Lianlei Wang[1,2,3,4#], Sen Zhao[1,5#], Xinyu Yang[1,4#], Pengfei Zheng[6#], Wen Wen[1,2,3], Kexin Xu[1,2,3], Xi Cheng[1,2,3], Qing Li[1,2,3], Anas M. Khanshour[7], Yoshinao Koike[8,9], Junjun Liu[6], Xin Fan[10], Nao Otomo[8,9], Zefu Chen[1,2,3], Yaqi Li[1,2,3], Lulu Li[11], Haibo Xie[6], Panpan Zhu[6], Xiaoxin Li[1,2,12], Yuchen Niu[1,2,12], Shengru Wang[1,2,3], Sen Liu[1,2,3], Suomao Yuan[4], Chikashi Terao[8], Ziquan Li[1,2,3], Shaoke Chen[10], Xiuli Zhao[13], Pengfei Liu[5,14], Jennifer E. Posey[14], Zhihong Wu[1,2,3,12], Guixing Qiu[1,2,3,12], DISCO study group (Deciphering Disorders Involving Scoliosis & COmorbidities), Shiro Ikegawa[9], James R. Lupski[15,16,17], Jonathan J. Rios[7,18,19], Carol A. Wise[7,18,19], Terry Jianguo Zhang[1,2,3*], Chengtian Zhao[6*], Nan Wu[1,2,3*]

Affiliations:

[1]State Key Laboratory of Complex Severe and Rare Diseases, Department of Orthopedic Surgery, Peking Union Medical College Hospital, Peking Union Medical College and Chinese Academy of Medical Sciences; Beijing, 100730, China.

[2]Beijing Key Laboratory for Genetic Research of Skeletal Deformity; Beijing, 100730, China.

[3]Key Laboratory of Big Data for Spinal Deformities, Chinese Academy of Medical Sciences; Beijing, 100730, China.

[4]Department of Orthopedic Surgery, Qilu Hospital of Shandong University, Cheeloo College of Medicine, Shandong University; Jinan, Shandong, 250012, China.

[5]Department of Molecular and Human Genetics, Baylor College of Medicine; Houston, TX, 77030, USA.

[6]Institute of Evolution & Marine Biodiversity, College of Marine Life Science, Ocean University of China; Qingdao, 266003, China.

[7]Center for Pediatric Bone Biology and Translational Research, Scottish Rite for Children; Dallas, TX, 75219, USA.

[8]Laboratory for Statistical and Translational Genetics, RIKEN Center for Integrative Medical Sciences; Yokohama, Kanagawa, 230-0045, Japan.

[9]Laboratory for Bone and Joint Diseases, RIKEN Center for Integrative Medical Sciences; Minato-ku, Tokyo, 108-8639, Japan.

[10]Department of Pediatric Endocrine and Metabolism, Maternal and Child Health Hospital of Guangxi Zhuang Autonomous Region; Nanning, Guangxi, 530012, China.

[11]Department of Newborn Screening Center, Beijing Obstetrics and Gynecology Hospital, Capital Medical University, Beijing Maternal and Child Health Care Hospital; Beijing, 100000, China.

[12]Department of Central Laboratory, Peking Union Medical College Hospital, Peking Union Medical College and Chinese Academy of Medical Sciences; Beijing, 100730, China.

[13]Department of Medical Genetics, Institute of Basic Medical Sciences, Chinese Academy of Medical Sciences & Peking Union Medical College; Beijing, 100730, China.

[14]Baylor Genetics; Houston, TX, 77021, USA.

[15]Departments of Pediatrics, Texas Children's Hospital and Baylor College of Medicine; Houston, TX, 77030, USA.

[16]Texas Children's Hospital; Houston, TX, 77030, USA.

[17]Human Genome Sequencing Center, Baylor College of Medicine; Houston, TX, 77030, USA.

[18]Department of Orthopaedics, University of Texas Southwestern Medical Center; Dallas, TX, 75390, USA.

[19]McDermott Center for Human Growth and Development, University of Texas Southwestern Medical Center; Dallas, TX, 75390, USA.

# These authors contributed equally to this work.

*Corresponding authors:

Nan Wu, M.D., E-mail: dr.wunan@pumch.cn

Chengtian Zhao, Ph.D., E-mail: chengtian_zhao@ouc.edu.cn

Terry Jianguo Zhang M.D., E-mail: zhangjianguo@pumch.cn

Author contributions

Conceptualization: LW, SZ, CZ, NW

Methodology: LW, SZ, XY, PZ, WW, XC, AK, YK, NO, YL, CT, PL, CZ, NW

Investigation: LW, SZ, XY, PZ , WW, XC, AK, YK, JL, XF, NO, YL, LL, HX, XL, SL, CT, SC, XZ, PL, SI, CZ, NW

Visualization: LW, SZ, XY, PZ , PZ, HX, JL, CZ, NW

Funding acquisition: LW, SW, TJZ, GQ, ZW, JR, CZ, NW

Project administration: LW, SZ, YN, ZW, GQ, TJZ, CZ, NW

Supervision: LW, SZ, SY, YN, ZW,GQ, TJZ, CZ, NW

Writing – original draft: LW, SZ, XY, JL, CZ, NW

70   Writing – review & editing: LW, SZ, XY, PZ, WW, KX, XC, QL, AK, YK, JL,
71 XF, NO, YL, LL, HX, PZ, XL, YN, SW, SL, CT, ZL, SC, XZ, PL, JEP, ZW, GQ, SI,
72 JRL, CAW, TJZ, CZ, NW

Conflict-of-interest statement

J.R.L has stock ownership in 23andMe, is a paid consultant for Regeneron
Pharmaceuticals and Novartis, and is a co-inventor on multiple United States and
European patents related to molecular diagnostics for inherited neuropathies, eye
diseases and bacterial genomic fingerprinting. The Department of Molecular and
Human Genetics at Baylor College of Medicine derives revenue from the
chromosomal microarray analysis (CMA by aCGH and/or SNP arrays), clinical
exome sequencing (cES) and whole-genome sequencing (WGS) offered in the
Baylor Genetics (BG) Laboratory (http://bmgl.com).

This file includes:

Main Text
Figures 1 to 8
Table 1
Figures S1 to S7
Tables S1 to S2, S4 to S7
Movie legends S1 to S8

Other Supplementary Materials for this manuscript include the following:

91   Tables S3
92   Movies S1 to S8

Abstract

Idiopathic scoliosis (IS) is the most common form of spinal deformity with unclear pathogenesis. In this study, we firstly reanalyzed the loci associated with IS, drawing upon previous studies. Subsequently, we mapped these loci to candidate genes using either location-based or function-based strategies. To further substantiate our findings, we verified the enrichment of variants within these candidate genes across several large IS cohorts encompassing Chinese, East Asian, and European populations. Consequently, we identified variants in the *EPHA4* gene as compelling candidates for IS. To confirm their pathogenicity, we generated zebrafish mutants of *epha4a*. Remarkably, the zebrafish *epha4a* mutants exhibited pronounced scoliosis during later stages of development, effectively recapitulating the IS phenotype. We observed that the *epha4a* mutants displayed defects in left-right coordination during locomotion, which arose from disorganized neural activation in these mutants. Our subsequent experiments indicated that the disruption of the central pattern generator (CPG) network, characterized by abnormal axon guidance of spinal cord interneurons, contributed to the disorganization observed in the mutants. Moreover, when knocked down *efnb3b*, the ligand for Epha4a, we observed similar CPG defects and disrupted left-right locomotion. These findings strongly suggested that ephrin B3-Epha4 signaling is vital for the proper functioning of CPGs, and defects in this pathway could lead to scoliosis in zebrafish. Furthermore, we identified two cases of IS in *NGEF*, a downstream molecule in the *EPHA4* pathway. Collectively, our data provide compelling evidence that neural patterning impairments and disruptions in CPGs may underlie the pathogenesis of IS.

Keywords: Idiopathic scoliosis (IS), Pathogenesis, *EPHA4* (*Ephrin type-A receptor 4*), Central pattern generators (CPGs).

Introduction

Idiopathic scoliosis (IS) is the most common form of spinal deformity, affecting about 2.5% of the global population (***Hresko 2013, Luk et al. 2010***). IS may have long-term physical and mental health consequences, such as cosmetic deformity, cardiopulmonary impairment, and even disability (***Weinstein et al. 2003***). All these consequences can severely reduce the quality of life. Early intervention with conservative treatments, such as braces, can control scoliosis progression and reduce the need for surgical intervention (***Weinstein et al. 2013***). However, IS often remains undiagnosed until malformation is evident, emphasizing the importance of risk-prediction measurements in medical management.

Genetic factors are thought to play a significant role in the development of IS, while only a few genes have been associated with the condition to date (***Miller 2007, Cheng et al. 2015***). Several single-nucleotide polymorphisms (SNPs) associated with susceptibility to IS have been identified through genome-wide association studies (GWASs), including SNPs linked to genes such as *LBX1*, *GPR126*, and *BNC2*. Notably, knockdown or overexpression of zebrafish homologs of these IS-associated genes have yielded body axis defects (***Kou et al. 2019, Zhu et al. 2015, Guo et al. 2016, Kou et al. 2013, Ogura et al. 2015***). In addition to common SNPs, rare variants with larger effect sizes or causing rare Mendelian disease traits also contribute to IS. Rare variants in *FBN1*, *FBN2,* and other extracellular matrix genes are associated with severe IS (***Buchan et al. 2014, Haller et al. 2016***). Linkage analysis of familial IS suggests that it follows autosomal dominant (AD), X-linked dominant, and multifactorial patterns of inheritance (***Miller et al. 2005***). Variants in *CHD7* and *AKAP2* are also implicated in the pathogenesis of Mendelian forms of IS (***Gao et al. 2007, Li et al. 2016***). Finally, centriolar protein POC5 and planar cell polarity protein Vang-like protein 1 (VANGL1) were also shown to be associated with IS. Notably, all these genes participate in a wide range of biological processes, the mechanisms underlying IS progression still require further investigation.

Consequently, there is currently no consensus on the etiology of IS (***Tang et al.***

2021).

In this study, we developed a novel pipeline that combines SNP-to-gene

mapping and rare variant association analysis. Using this approach, we analyzed a

large Chinese population with IS and further validated our findings in East Asian

and European populations. Through these analyses, we identified *EPHA4* as a novel

gene associated with IS. To confirm the pathogenicity of this gene, we used a

zebrafish model and found that impaired *EPHA4* pathway components and resulting

defects in central pattern generators (CPGs) are essential factors in the pathogenesis

of IS. Additionally, we searched for candidate genes related to EPHA4 signaling

pathways and identified mutations in the *NGEF* among IS patients. Overall, our data

suggest that the impairment of the *EPHA4* pathway and central pattern generators

plays a previously unknown role in the development and progression of IS.

Results

Enrichment analysis of rare variants in IS candidate genes

To identify candidate genes associated with IS, we conducted a comprehensive literature review of SNPs that are linked to IS from 14 published GWASs. This led us to obtain forty-one SNPs that showed genome-wide significance ($P < 5 \times 10^{-8}$) as detailed in Table S1. We then employed positional mapping, expression quantitative trait locus (eQTL) mapping, or chromatin interaction mapping to identify 156 candidate genes that could be potentially linked to these SNPs. Further analysis of rare variants for the 156 candidate genes using exome data from 411 Chinese probands and 3,800 unrelated Chinese controls identified *EPHA4* as the only significant gene (P=0.045, OR=4.09) (Table S2). EPHA4 is a member of the Eph family of receptor tyrosine kinases which play a vital role in the development of the nervous system (***Kania and Klein 2016***). We identified three rare variants in *EPHA4* from three IS patients, including one splicing-donor variant and two missense variants (Table S2). The flowchart of the entire pipeline is shown in Fig. S1.

Inheritance pattern and functional analyses of variants in *EPHA4*

We characterized the inheritance pattern of variants in *EPHA4* using either trio sequencing data or Sanger sequencing for the parents. None of the parents being tested had scoliosis based on a clinical screening test. Of the three rare variants in *EPHA4*, two (NM_004438.3: c.1443+1G>C and c.2546G>A [p.Cys849Tyr]) arose *de novo* (Fig. 1A and Table 1). The heterozygous splicing variant (c.1443+1G>C) was identified in a 15-year-old female IS patient (SCO2003P0846) (Fig. 1B). Scoliosis in this patient was diagnosed at 13 years of age, with a main curve Cobb angle of 60°. The *in vitro* minigene assay showed that the c.1443+1G>C variant introduced a new splicing site, resulting in a 36-bp in-frame deletion in exon 6 (Fig. 1G, 1I, S2A). The *EPHA4* heterozygous missense variant (c.2546G>A, p.Cys849Tyr), identified in a 13-year-old female IS patient (SCO2003P2146) (Fig.

1C), is located in the protein tyrosine kinase domain of EPHA4 protein (Fig. 1A). The onset of scoliosis in this patient was at 11 years of age with a 70° main curve Cobb angle. As EPHA4 plays a crucial role in the phosphorylation of CDK5, which subsequently activates the downstream pathways (*Fu et al. 2007*), we conducted western blotting to quantify the phosphorylation level of CDK5. Our results revealed that the missense variant resulted in a decreased phosphorylation level of CDK5 (p=0.015), suggesting a partial loss-of-function (LoF) of *EPHA4* (Fig. 1F).

We next investigated *de novo* non-coding variants of *EPHA4* via whole genome sequencing (WGS) in 116 trio families with IS. A *de novo* heterozygous *EPHA4* intronic variant (c.1318+10344A>G) was identified in a 16-year-old female patient (SCO2003P2080) (Fig. 1D and Table 1). She developed scoliosis at 11 years of age with a 70° main curve Cobb angle. This variant is predicted to affect the branch point of the fifth intron of *EPHA4* (Fig. S2B). We performed nested PCR to show that this variant induced exon 5 skipping, resulting in a 339-bp in-frame deletion (Fig. 1H,1J).

Next, we employed an in-house gene matching approach under the framework of DISCO study, which identified a 6-year-old patient (SCO2003P3202) who had been previously diagnosed with Waardenburg syndrome caused by a 4.46 Mb *de novo* deletion at 2q35-q36.2 (*Li et al. 2015*) (Table 1). Intriguingly, the patient also presented mild scoliosis (Fig. 1E). The deletion included the entire *PAX3* gene, which was responsible for the Waardenburg syndrome phenotype, and 36 neighboring genes, including *EPHA4* (Fig. 1K). As scoliosis is not typically associated with Waardenburg syndrome caused solely by *PAX3* pathogenic variants (*Tassabehji et al. 1993*), we hypothesize that the deletion of *EPHA4* may be responsible for the IS phenotype in this patient.

Notably, the GWAS signal which we mapped to *EPHA4* (rs13398147) (*Zhu et al. 2015*) represents a significant eQTL in esophagus and colon tissues, with the T allele associated with decreased expression of *EPHA4*. In our East Asian GWAS

cohort of 6,449 adolescent IS patients and 158,068 controls, we identified another two eQTLs in *EPHA4* associated with decreased expression of *EPHA4* in brain tissue (Table S3). In the same GWAS cohort, common SNPs in *EPHA4*, after aggregation, also showed significant enrichment (P=0.023) in IS patients versus controls. Taken together, the convergence between rare and common variants of *EPHA4* that lead to LoF or hypomorphic effects highlights the pivotal role of *EPHA4* in the pathogenesis of IS.

IS-like phenotypes in zebrafish *epha4* mutants

To investigate the role of *EPHA4* in scoliosis, we utilized zebrafish as a model system due to its versatile nature in modeling adolescent IS (***Grimes et al. 2016, Bagnat and Gray 2020, Xie et al. 2022, Boswell and Ciruna 2017***). Zebrafish have two homologs of *EPHA4*, *epha4a* and *epha4b*. Using CRISPR-Cas9, we established a stable *epha4a* zebrafish mutant line with a 63-bp deletion in exon 3, which introduced a stop codon and resulted in a truncated protein (Fig. 2A-B). The homozygous *epha4a* mutant larvae had no apparent defects in either notochord or body axis development (Fig. S3A). While in consistent with previous reports, the hindbrain rhombomeric boundaries were found to be defective in both the *epha4a* mutants and morphants (Fig. S3B) (***Cayuso et al. 2019, Cooke, Kemp and Moens 2005***). Interestingly, more than 75% adult mutants showed mild scoliosis (88 of 116), and some mutants exhibited severe scoliotic phenotype (4 of 116) (Fig. 2C-D, Movie S1&S2). Remarkably, some heterozygous adult mutants also developed mild scoliosis (14 of 95), whereas none of the wild-type fish showed any signs of scoliosis (0 of 76) (Fig. 2C-D). Similarly, we further generated the *epha4b* mutants with a 25-bp deletion in exon 3, which resulted in a frameshift mutation (Fig. S4A). The homozygous *epha4b* zebrafish mutants also developed mild scoliosis (28 of 43) (Fig. S4B-C). Intriguingly, both *epha4a* and *epha4b* mutants exhibited early onset scoliosis starting from around 20 days post fertilization (Fig. S4D), a stage similar to that of IS patients. Collectively, these data suggest that mutations in Epha4 proteins

242   are linked to the scoliotic phenotype in zebrafish.

Abnormal left-right swimming pattern in the absence of Epha4a

While the *epha4a* mutants seemed to be grossly normal during the larvae stages,

they developed spinal curvature gradually during later development. We decided to

investigate whether these mutants exhibited any abnormalities in their behavior at

the larvae stages using EthoVision XT software. By monitoring the swimming

behavior of 8 days post-fertilization (dpf) zebrafish larvae, we found the motion

distance and swimming velocity were significantly decreased in *epha4a* mutants (Fig.

3A-B). In addition, we observed a remarkable difference in the relative turning angle

and angular velocity between these two groups. The wild-type group changed their

swimming direction randomly, showing a relative angle around 0 degrees (Fig.

3C-D). In contrast, the *epha4a* mutants favored turning to one side of their directions

(Fig 3C-D). Moreover, the absolute turn angles and turning speed (angular velocity)

were significantly higher in the mutants (Fig. 3E-F).

Next, we compared the swimming behavior after startle response between

wild-type and *epha4a* mutants. We used a needle to touch the head or the tail of 5

days post-fertilization (dpf) zebrafish larvae to stimulate swimming behavior. In

sibling controls, the larvae responded to the tactile stimulation and swam away

quickly (Movie S3). Conversely, the *epha4a* mutants failed to respond to the initial

stimulation, and the swimming pattern was defective with an abnormal bending

pattern (37 of 62 from tail-stimulation group and 34 of 77 from head-stimulation

group) (Fig. 3G, Movie S4). We further performed more comprehensive analysis by

high-speed video microscopy. After tactile stimulation, wild-type larvae displayed a

high-speed C-bend turn followed by a weaker counterbend turn after several

milliseconds (Fig. 3H-I). This rhythmic left/right swimming pattern ensures that the

fish swim away from the frightening stimulus. The turning angles of the control

larvae after stimulation changed with a sinusoidal wave pattern (Fig. 3J). In contrast,

this pattern was dramatically different in *epha4a* mutants (Fig. 3I,K). Of note,

although the C-bend turning angles were similar between *epha4a* mutants and

control siblings, the turning angles of the counterbend decreased significantly (Fig.

3L), implying a left/right coordination defect. Taken together, both regular

swimming and tactile stimulation analyses suggested that the left-right coordination

swimming pattern is compromised in the absence of Epha4a.

275   Defects of left-right coordination due to abnormal CPG in the absence of

276   EphA4

The coordinated left-right locomotion of zebrafish larvae relies on the

synchronized contraction of muscle fibers, a process regulated by motor neurons

situated on each side of the fish. To explore this intricate mechanism, we utilized a

Tg(*elavl3*:GAL4; *UAS*: GCaMP) double transgene, allowing the expression of a

genetically encoded calcium sensor in all neurons. We observed the rhythmic

activation of calcium signaling in motor neurons located within the spinal cord (Fig.

4A, Movie S5). In wild-type larvae, the calcium signals exhibited an alternating

pattern between the left and right sides of the body, whereas this coordinated pattern

was disrupted in *epha4a* morphants (Fig. 4 A, B vs A', B', Movie S6). Additionally,

we found that the activation frequency of motor neurons on the left and right sides

was comparable in wild-type larvae, but significantly different in the absence of

Epha4a (Fig. 4C-D).

One well-established concept regarding left-right coordination involves the

presence of central pattern generators (CPGs), which are regulated by interneuron

circuitry within the spinal cord. We further examined the axon guidance of

interneurons in *epha4a* mutants. First, we investigated the commissural trajectories

of reticulospinal (RS) interneurons. In control larvae, the large Mauthner neurons,

along with other RS neurons, were symmetrically positioned on each side of the

midline and projected their axons to the contralateral sides (Fig. 5A). These

bilaterally projected axons typically crossed at the midline and subsequently

synapsed on motoneurons of the opposite sides, contributing to the generation of

spinal cord neural circuits (**Hale et al. 2016**). However, in *epha4a* mutant larvae, we

observed an abnormal pattern in axonal projections. Specifically, the mutant axons

failed to traverse the midline and instead extended ipsilaterally (Fig. 5A). In addition,

the distance between Mauthner neurons (rhombomere 4) and rhombomere 7 was

significantly decreased and rhombomere 5 was scarcely visible in the mutant larvae

(Fig. 5A-B). Furthermore, the sites of axon crossing between two Mauthner neurons

tended to deviate to one side of the midline in the mutants (Fig. 5C-D').

Cerebrospinal fluid-contacting neurons (CSF-cNs) represent a unique type of

interneuron responsible for modulating the V0 and V2a interneurons, which are

integral components of locomotor CPGs (**Fidelin et al. 2015, Wu et al. 2021,**

Talpalar et al. 2013). In wild-type larvae, we observed that the ascending axons of

these neurons projected either to the right or left side from the midline, as visualized

using Tg(*pkd2l1*:GAL4;*UAS*:Kaede) double transgenic larvae (Fig. 6A). However, in

the absence of Epha4a, the projection of these neurons exhibited notable

disorganization, with numerous axons crossing the midline from one side of the

trunk. This disorganized pattern was observed in both heterozygotic and

homozygotic mutants (Fig 6A-B). Additionally, we employed an optogenetic

approach to activate these CSF-cNs, utilizing the Tg(Gal4^{s1020t}; *UAS*:ChR2) double

transgene. Following optical stimulation, we observed robust tail oscillations as

previously described (**Wyart et al. 2009**). In wild-type larvae, these tail oscillations

exhibited a symmetrical left-right beating pattern(Fig. 6C, D, Movie S7). However, a

striking disruption of this symmetry was observed in *epha4a* morphants, as they

consistently beat towards one side of the trunk following optical stimulation(Fig. 6C',

D' and E, Movie S8). Collectively, these findings demonstrate that the left-right

coordination deficiencies observed in *epha4a* mutants arise from abnormal neural

circuit formation, which consequently disrupts the integrity of the CPGs.

Ephrin B3-EphA4 signaling regulates interneuron axon extension

To further explore the role of Epha4a during interneuron axon extension, we examined the expression of *epha4a* during early zebrafish embryonic development. Whole-mount in situ hybridization results showed that both *epha4a* and *epha4b* were abundantly expressed in the zebrafish spinal cord (Fig. S5A). We plotted the expression of these two genes using a published single-cell transcriptome data, which showed that *epha4a* and *epha4b* were both expressed in interneurons such as serotonergic interneurons as well as V0 and V2a interneurons, suggesting a role for *epha4* in interneuron function (Fig. S5B) (*Cavone et al. 2021*). Notably, the expression of *efnb3b*, encoding the ligand of Epha4, was highly enriched in the midline floor plate cells (Fig. S5B).

Ephrins, through interacting with Eph receptors, play a critical role in repulsive axon guidance during neural development (*Flanagan and Vanderhaeghen 1998, Egea and Klein 2007*). We further analyzed axon guidance in *efnb3b* morphants. Similar to those of *epha4a* mutants, the *efnb3b* morphants also displayed axon guidance defects, as well as abnormal startle response and uncoordinated calcium activation (Fig. S6A-E). In addition, morphants larvae also displayed left-right oscillations defects after optogenetic stimulation (Fig. S6D-E).

Candidate variants in the *EPHA4*-related genes

Our zebrafish studies suggested that EPHA4 signaling is crucial for interneuron axon guidance, hence the formation of functional CPGs. Next, we further asked whether mutation of other components of the EPHA4 signaling can result in IS in humans. By searching for rare variants in *EPHA4*-related genes (Fig. S7) (*Szklarczyk et al. 2019*), we identified heterozygous *de novo* start-loss variant (c.1A>G, p.Met1?) in *NGEF* in a 17-year-old male (SCO2003P3332) (Fig. 7A-B, Table 1), whose scoliosis was diagnosed at 15 years of age with a main curve Cobb angle of 60°. *NGEF* encodes the neuronal guanine nucleotide exchange factor Ephexin that differentially affects the activity of GTPases RHOA, RAC1, and CDC42. The activation of Ephexin is triggered by ephrin through EPHA4 (*Shamah*

et al. 2001).

Strikingly, in a European-ancestry IS cohort, we further identified a dominant

missense variant (c.857G>A, p.Ala286Val) in *NGEF* in a quad family with three

affected members (Fig. 7C, Table 1). The proband (II:2, TSRHC01) with onset of

scoliosis at 14 years of age has a 58° main curve Cobb angle. This variant, which is

located in the RhoGEF domain of Ephexin, is predicted to be highly deleterious

(CADD=29.6).

Altogether, our results suggest that defects of the CPGs owing to abnormal

EPHA4 signaling maybe one of the crucial factors responsible for IS.

Discussion

IS is a disease with diverse causes, and the underlying mechanisms can vary even among patients with similar scoliotic phenotypes. Previous genetic studies have highlighted the significance of the extracellular matrix (ECM) in maintaining the balance of axial bone and supporting soft tissues in the spine, thus playing a crucial role in IS development (**Haller et al. 2016**). For example, the top SNP associated with IS maps to *LBX1*, an essential molecule for ECM maintenance and bone homeostasis (**Takahashi et al. 2011**). Additionally, genetic loci in muscle development-related genes have also been associated with the onset of scoliosis, emphasizing the intricate interaction between bones and muscles (**Ogura et al. 2015**). However, it is worth noting that the variants in these genes explain only a small fraction of the overall heritability of IS. Consequently, it is imperative to establish connections between the extensive genetic findings and biological mechanisms that can elucidate the etiological landscape of IS.

In this study, we mapped 41 significant genome-wide loci to functional genes through positional mapping and functional mapping such as eQTL. Then we determined the enrichment in patient cohorts of rare variants in these genes, which may have greater impacts compared with common SNPs. This approach revealed the convergence of SNPs and rare variants in *EPHA4* that are enriched in patients with IS. We also identified additional high-impact variants in *NGEF*, which is involved in the *EPHA4* pathway.

Our studies using zebrafish have revealed that deficiency of Epha4 can lead to the development of scoliosis. Interestingly, we observed that even heterozygotic *epha4a* mutants displayed mild scoliosis (Fig. 2C), which is consistent with the occurrence of this condition in scoliosis patients. Further analysis involving behavior and imaging demonstrated that the absence of Epha4a resulted in defective left-right coordination. This coordination is crucially governed by Central Pattern Generators

(CPGs), which generate rhythmic patterns of neural activity to coordinate limb

movements on both sides of the body (*Talpalar et al. 2013, Marder and Bucher*

2001, Kiehn 2006). Previous studies have reported the involvement of ephrin and its

receptors in axon guidance during the maturation of neural circuits, including CPGs

(*Iwasato et al. 2007, Borgius et al. 2014, Kullander et al. 2003, Andersson et al.*

2012). In line with this, we observed disrupted axon guidance of the interneurons,

which are integral components of CPGs, in *epha4a* mutants. This finding suggests

that CPG malfunction is the primary factor underlying the scoliosis observed in these

mutants. It is highly likely that the lack of coordinated left-right locomotion

generates imbalanced mechanical forces on the spine, gradually leading to spinal

curvature during later stages of development (Fig. 8).

Thus, our data provided a novel biological mechanism of IS, i.e., the

impairment of neural patterning and CPG. Previous studies have provided clues on

the role of CPGs in IS. Patients with adolescent IS showed asymmetric trunk

movement during gait, as characterized by increased relative forward rotation of the

right upper body in relation to the pelvis (*Kramers-de Quervain et al. 2004, Nishida*

et al. 2017). An electromyography (EMG) study also showed asymmetric activation

of paraspinal muscles between the convex and concave sides at the scoliosis curve

apex (*Shimode, Ryouji and Kozo 2003*). In a child with a strong family history of IS,

asymmetric hyperactivity was observed by EMG months before scoliosis was

evident (*Valentino et al. 1985*). These left-right locomotor coordination

abnormalities indicated the maldevelopment of CPGs as potential cause of IS. The

CPG asymmetry may induce an imbalance in trunk muscle strength, resulting in

asymmetric rib drooping. This leads to an abnormal vertebral rotation and then the

onset of IS. Moreover, in a companion study, Wang et al. identified a number of rare

variants in *SLC6A9*, which encodes glycine transporter 1 (GLYT1), in familial and

sporadic adolescent IS cases. The *slc6a9* mutant zebrafish also exhibited

discoordination of spinal neural activities with pronounced lateral spinal curvature,

recapitulating the human IS phenotype (data unpublished). Taken together, we

propose that the dysfunction of CPGs would cause an imbalance in the motor drive

from the spinal cord and the asymmetric transversospinalis muscle pull, eventually

producing a scoliotic curve (Fig. 8).

In summary, we showed that both common and rare variants in the *EPHA4*

pathway contribute to the genetic architecture of IS. The dysfunction of the *EPHA4*

pathway causes IS through the impairment of neural patterning and CPGs.

Materials and Methods

Mapping of candidate genes utilizing previous association studies

Through a systematic literature review, we identified IS-associated SNPs reported in GWASs and meta-analyses of GWAS. The literature search was carried out using MEDLINE (via Pubmed.gov) and Web of Science (via Clarivate Analytics) and was limited to English-language articles published from January 1980 to October 2020. The following keywords were combined to perform the search: 'idiopathic scoliosis' AND 'GWAS' OR 'SNP' OR 'single nucleotide polymorphism' OR 'variant' (Table S4). The inclusion/exclusion criteria of abstracts are provided in Table S5. After screening the titles and abstracts, we obtained from the full-text articles the rsID, chromosome, and position for SNPs with a threshold for genome-wide significance of $P < 5.0 \times 10^{-8}$. For SNPs identified in multiple studies, we recorded the lowest P-value.

SNPs reported in previous studies were first pruned using Functional Mapping and Annotation (FUMA, v1.4.1, https://fuma.ctglab.nl/) (***Watanabe et al. 2017***). Significant SNPs were considered independent at $r^2 < 0.6$. All known SNPs (available in the 1000 Genomes reference panel, https://www.internationalgenome.org/) were included for further gene mapping if they were in a linkage disequilibrium (LD) block ($r^2 \geq 0.6$) with a significant independent SNP. Three SNP-to-gene strategies were used:

1) For positional mapping of significant independent SNPs, we used annotations obtained from ANNOVAR (http://annovar.openbioinformatics.org). A 10-kb maximum distance was applied for intergenic SNPs.

2) For the eQTL mapping, we mapped significant independent SNPs and SNPs in an LD block to eQTLs acorss 44 GTEx tissue types (release V8) (***Consortium 2020***). SNP-gene pairs with a false discovery rate ≤ 0.05 were considered significant.

3) For chromatin interaction mapping, we overlapped the significant independent

451   SNPs and SNPs in an LD block with one end of significantly interacting regions

452   in all optional tissue/cell types. Genes were mapped if their promoter regions

453   overlapped with another end of the significant interactions. The promoter region

454   was defined as the region from −250 bp to +500 bp relative to the transcription

455   start site.

Cohort description

The Peking Union Medical College Hospital (PUMCH) cohort: The PUMCH

cohort comprised 411 unrelated Chinese patients with severe IS (Cobb angle ≥ 40°)

who underwent spinal surgery in the PUMCH between October 2017 and March

2022 as part of the Deciphering disorders Involving Scoliosis and COmorbidities

(DISCO) study (http://www.discostudy.org/). The clinical diagnosis was confirmed

using standing full-spine radiographs, three-dimensional computed tomography, and

magnetic resonance imaging. These patients did not show any congenital or

neuromuscular defect at the time of recruitment. The control cohort consisted of

3,800 individuals without observable scoliosis and with exome sequencing or

genome sequencing performed at PUMCH for clinical or research purposes.

Individuals with vertebrae malformation or congenital developmental defects were

excluded. A patient with a 2q35-36.2 deletion, including the *EPHA4* gene, was also

recruited through in-house gene matching of the DISCO study (***Li et al. 2015***). This

patient was subsequently evaluated for scoliosis by examination and radiography.

The East Asian cohort: The East Asian cohort includes totally 6,449 IS patients

and 158,068 controls from four independent datasets (Japanese dataset 1 (***Kou et al.***

2013, Takahashi et al. 2011): 1,261 cases, 15,019 controls; Japanese dataset 2

(***Ogura et al. 2015***): 878 cases, 21,334 controls; Japanese dataset 3 (***Kou et al. 2019***):

3,333 cases, 119,630 controls; Hong Kong dataset: 977 cases, 2,085 controls). The

inclusion criteria for IS subjects was as same as our previous studies (***Kou et al.***

2019, Fan et al. 2012).

The Texas Scottish Rite Hospital for Children (TSRHC) cohort: We used a
replication cohort of European-ancestry patients with IS from the TSRHC. Cases
considered for inclusion in the study met criteria for a positive diagnosis of IS:
lateral deviation from the midline greater than 15 degrees as measured by the Cobb
angle method from standing spinal radiographs, axial rotation toward the side of the
deviation and exclusion of relevant co-existing diagnoses.

Blood sample collection

In the PUMCH cohort, genomic DNA samples were extracted from peripheral
blood leukocytes of each subject using a QIAamp DNA Blood Mini Kit (Qiagen,
Hilden, Germany), according to the manufacturer's protocol. Purified DNA was
qualified by Nanodrop 2000 (Thermo Fisher Scientific, Waltham, MA, USA) and
quantified by Qubit 3.0 using the dsDNA HS Assay Kit (Life Technologies, Carlsbad,
CA, USA). DNA samples were stored at 4°C until used.

In the East Asian and TSRHC cohort, genomic DNA was extracted from
peripheral blood or saliva using a standard protocol.

DNA sequencing and variant calling

In the PUMCH cohort, whole exome sequencing (WES) or whole genome
sequencing (WGS) was performed on peripheral blood DNA from all individuals and
available family members (Table S6). A SureSelect Human All Exon V6+UTR r2 core
design (91 Mb, Agilent) was used for exon capture. The exomes were then sequenced
on an Illumina HiSeq 4000 (Illumina, San Diego, CA, USA) according to the
manufacturer's instructions. For WGS, sequencing libraries were prepared using the
KAPA Hyper Prep kit (KAPA Biosystems, Kusatsu, Japan) with an optimized
manufacturer's protocol. We performed multiplex sequencing using an Illumina
HiSeq X-Ten sequencer (Illumina, San Diego, CA, USA). Variant calling and
annotation were done in-house using the Peking Union Medical College Hospital
Pipeline (PUMP) described previously (***Zhao et al. 2021, Chen et al. 2021***).

In the TSRHC cohort, subjects in 162 adolescent IS families were sequenced as part of the Gabriella Miller Kids First Pediatric Research Consortium (GMKF) at The HudsonAlpha Institute for Biotechnology (Huntsville, AL). In summary, DNA was normalized, sheared and then ligated to Illumina paired-end adaptors. The purified ligated DNA was amplified and exome sequencing was performed on the Illumina HiSeq X platform. The sample's sequences were aligned into GRCh38 and genotypes were joint-called per each family by the GMKF's Data Resource Center (DRC) at Children's Hospital of Philadelphia following GATK best practices as detailed here (*Mukhopadhyay et al. 2020*). GMKF DRC's alignment and joint genotyping pipelines are open source and made available to the public via GitHub (https://github.com/kids-first/kf-alignment-workflow and https://github.com/kids-first/kf-jointgenotyping-workflow).

Rare variant association analysis of the candidate genes

To determine the contribution of rare variants in GWAS candidate genes to IS, we analyzed the gene-based mutational burden for 156 candidate genes. Only ultra-rare variants with a gnomAD population-max allele frequency ≤ 0.01% and a cohort allele count ≤ 3 were analyzed. Each variant was assigned a weight, and the mutational burden of a given gene was defined as the maximum weight value among all ultra-rare variants carried by the individual. The SAIGE-GENE+ package was used to determine the weighted mutational burden test for each gene (*Zhou et al. 2022*).

Weighting criteria for the weighted burden analysis were developed according to the variant types and in silico results. The LoF variants (canonical splicing variants, nonsense variants, or variants that cause frameshift, stop-gain, or start-loss) were calculated together with the protein-altering variants, including missense variants and in-frame indels. Each variant was assigned a weight range from 0-1 and the mutational burden for a given gene was defined as the maximum weight value among all ultra-rare variants carried by the individual. The LoF variants annotated as

'high confidence' by the loss-of-function transcript effect estimator (LOFTEE) (*Karczewski et al. 2020*) were assigned a weight value of 1.0. The LoF variants annotated as 'low confidence' or unlabeled by LOFTEE, the non-canonical splicing variants with a SpliceAI score > 0.5, and the missense variants with a rare exome variant ensemble learner (REVEL) (*Ioannidis et al. 2016*) score of 0.8 were assigned a weight value of 0.8. The in-frame insertions/deletions (indels) with a Combined Annotation Dependent Depletion (CADD) (*Kircher et al. 2014*) score > 20 and the missense variants with a REVEL score > 0.6 and ≤ 0.8 were assigned a weight value of 0.6. The in-frame indels with a CADD score > 10 and ≤ 20 and the missense variants with a REVEL score > 0.4 and ≤ 0.6 were assigned a weight value of 0.4. The in-frame indels and the missense variants with a REVEL score > 0.2 and ≤ 0.4 were assigned a weight value of 0.2. The remaining missense variants were assigned a weight value of 0.

Sanger sequencing of familial participants

Sanger sequencing of familial participants was performed to determine the origin of variants in *EPHA4* and *NGEF*. All LoF variants and protein-altering variants identified in familial participants were validated. Variant-encoding gene regions were amplified by PCR from genomic DNA obtained from probands, as well as from parents for trios, to determine the origin of the variants. The amplicons were purified using an Axygen AP-GX-50 kit (Corning, NY, USA) and sequenced by Sanger sequencing on an ABI 3730xl instrument (Thermo Fisher Scientific, Waltham, MA, USA).

Minigene assay

The splicing variant (*EPHA4:* c.1443+1G>C) was characterized by a minigene assay. Genomic DNA from the heterozygous patient was amplified by PCR using a high-fidelity DNA polymerase. Amplicons included exon 6, intron 6, exon 7, intron 7, and exon 8 of the *EPHA4* gene. PCR products were cloned into the vector via the

restriction sites BamHI and MluI for pCAS2, which is based on the pcDNA3.1

plasmid (Thermo Fisher Scientific, Waltham, MA, USA). Clones with wild-type or

mutant genomic inserts were selected and verified by sequencing the cloned DNA

fragments. The recombinant plasmids were transfected into 293T cells using

LipofectamineTM 3000 reagent (Thermo Fisher Scientific, Waltham, MA, USA).

For RT-qPCR, total RNA was isolated from the transfected cells using Trizol reagent

(Thermo Fisher Scientific, Waltham, MA, USA and reverse transcription was

performed using the GoScript™ Reverse Transcription System (Promega, Madison,

MI, USA). PCR amplification was performed using the pCAS2-RT-F and

pCAS2-RT-R primers, and the products were sequenced using pCAS2-RT-F.

Primers sequences for the minigene assay were as follows:

pCAS2-RT-F: 5′-CTGACCCTGCTGACCCTCCT-3′

pCAS2-RT-R: 5′-TTGCTGAGAAGGCGTGGTAGAG-3′

Nested PCR

The splicing variant (*EPHA4:* c.1318+10344A>G) was characterized by nested

PCR. RNA was extracted from the whole blood of the patient using the TRIzon

Reagent (CWBIO, Hangzhou, China) according to the manufacturer's guidelines.

The cDNA was synthesized using HiScript II 1st Strand cDNA Synthesis Kit with a

gDNA wiper (Vazyme, Nanjing, China) according to the manufacturer's guidelines.

Nested PCR was performed as described previously (*Yao and Tavis 2005*). Nested

PCR reactions in 50 μl, included 2 μl cDNA from the reverse transcription reaction

as the template for the first round of PCR or 2 μl of first-round PCR product as the

template for the second PCR, 1.5 μl 10 μM sense primer, 1.5 μl 10 μM anti-sense

primer, 5 μl nucleotide mix (2 mM each dNTP), 5 μl 10× KOD Buffer, 1 μl 1 unit/μl

Kod-Plus-Neo polymerase, 3 μl 25 mM $MgSO_4$ and 31 μl ddH_2O. The PCR program

was (94°C for 2 min, 98°C for 10 s, 55°C for 30 s, 68°C for 60 s) × 20 cycles and

then (94°C for 2 min, 98°C for 10 s, 57°C for 30 s, 68°C for 60 s) × 30 cycles.

Primers sequences:

For the first round:

F: 5′-GGCTCCTGTGTCAACAACTC-3′

R: 5′-GTTGGGATCTTCGTACGTAA-3′

For the second round:

F: 5′-AACTGCCTATGCAACGCTGG-3′

R: 5′-AGCTGCAATGAGAATTACC-3′

Western blots

For the missense variant in *EPHA4* (c.2546G>A, p.Cys849Tyr), wild-type and

mutant EPHA4 proteins were expressed from EPHA4-C1-pEGFP plasmids

introduced into the 293T cells. Cells were cultured in six-well plates and transfected

with DNA (2 mg/well) using Lipofectamine 3000 reagent (Thermo Fisher Scientific,

Waltham, MA, USA). After 48 hours, cells were harvested, and protein extracts were

prepared as described (***Ding et al. 2017***). The mutant proteins and wild-type proteins

were fused to GFP. The expression of the two proteins was compared by western

blots with an GFP antibody using the ECL detection system. The GFP antibody was

purchased from Cell Signaling Technology (Cell Signaling, Danvers, MA, USA).

The CDK5 antibody was purchased from Santa Cruz Biotechnology (Santa Cruz,

CA, USA). A phospho-specific antibody against CDK5 phosphorylated at Tyr

(GeneTex, Irvine, CA 92606 US) was purchased from GeneTex. Western blot

experiments were repeated twice with similar results for the replicates.

Genotyping and imputation of GWAS

In the East Asian Japanese cohort, genotyping was performed by Illumina

Human610 Genotyping BeadChip, Illumina HumanOmniExpressExome   and

HumanOmniExpress as our previous GWASs (***Kou et al. 2019, Ogura et al. 2015,***

Takahashi et al. 2011, Kou et al. 2013). For quality control (QC), subjects with call

rate < 0.98, high-degree relatedness with other subjects and outliers of East Asian

ethnicity were excluded. For variants QC, the exclusion criteria of variants were as

follows: call rate < 0.99, P-value for Hardy-Weinberg equilibrium $< 1.0 \times 10^{-6}$ and

minor allele count < 10. The reference panel for imputation, namely JEWEL7K, was

composed of 1000 Genomes Project phase 3 (v5) (***Auton et al. 2015***) and Japanese

whole-genome sequence data (***Das et al. 2016***) with 3256 high-depth subjects (≥30

read counts) and 4216 low-depth subjects (≤ 15 read counts). Using EAGLE 2.4.1

(https://alkesgroup.broadinstitute.org/Eagle/) to determine the haplotypes, pre-phasing

was conducted. Genotypes were imputed using Minimac4 (version1.0.0) (***Das et al.***

2016). After imputation, we excluded variants with minor allele frequency (MAF) <

0.005 and low imputation quality ($R^2 < 0.3$).

In the East Asian Hong Kong cohort, samples were genotyped with Illumina

Human Omni ZhongHua-8 Beadchips. Illumina Genome Studio v2.0 was used to

convert raw data into PLINK format. The QC steps of samples and variants were

described in a previous study (***Marees et al. 2018***). Genotype phasing and imputing

were executed using SHAPEIT v2.r900 (***Delaneau, Marchini and Zagury 2012***) and

IMPUTE2 (***Marchini et al. 2007***). The imputed data was filtered using the following

parameters: INFO > 0.6, Certainty > 0.8, and MAF > 0.01. Association analysis was

performed using PLINK v1.9 logistic regression with covariates: sex, age and top 20

principal components of the variance-standardized relationship matrix.

East Asian GWAS meta-analysis for IS

For the meta-analysis of the four datasets (three Japanese datasets and one

Hong Kong dataset), an inverse-variance-based method was performed by METAL

(version2011-03-25) (***Willer, Li and Abecasis 2010***). SNPs in three or more of the

four cohorts were used in subsequent analyses.

Gene-based common variant analysis and eQTL analysis

SNPs in *EPHA4* or within 20 kb flanking *EPHA4* with a significant association

with IS were retrieved. SNPs were matched to potential eQTLs according to the

GTEx database (v8, https://gtexportal.org). Gene-based common variant analyses

were performed using Multi-marker Analysis of GenoMic Annotation (MAGMA)

(*de Leeuw et al. 2015*) and FUMA (*Watanabe et al. 2017*) using default settings

with LD information from the 1000 Genomes Project East Asian population (1KGP

EAS) as a reference. SNPs located within two kb upstream and one kb downstream

from *EPHA4* were included in the gene-based analysis.

Zebrafish strains, mutants, and morphants

Zebrafish Tuebingen (TU) strains were maintained at 28°C on a 14-hour/10-hour

light/dark cycle. Embryos were raised at 28.5°C in E3 medium (5 mM NaCl, 0.17

mM KCl, 0.39 mM $CaCl_2$, 0.67 mM $MgSO_4$) following standard protocols. Zebrafish

have two homologs of *EPHA4*, *epha4a* and *epha4b*. The CRISPR/Cas9 system was

used to generate zebrafish *epha4a* and *epha4b* mutants. To increase efficiency, we

injected Cas9 mRNA together with multiple single guide RNAs (sgRNAs) for each

gene (Table S7). The sgRNA sequences for *epha4a* and *epha4b* are listed in Table S7.

Morpholino sequences for *epha4a* and *efnb3b* knockdown analysis are also listed in

Table S7.

Micro CT imaging

Adult zebrafish *epha4a* mutants or wild-type siblings were euthanized with

tricaine methanesulfonate and fixed in 4% paraformaldehyde. Micro CT images were

captured using a PerkinElmer Quantum GX2. Planar images acquired over 360° of

rotation were reconstructed using QuantumGX. Three-dimensional renders of the

skeleton were made with Analyze 12.0 software (AnalyzeDirect).

Behavioral recordings and analysis

For behavior analysis, individual zebrafish larvae were transferred into 24-well

plate with fresh E3 medium at 8 dpf. Then, the plate was placed inside the

Daniovision (Noldus) observation chamber for further behavior analysis.

Video-tracking of swimming activity and further statistical analysis were performed

using the EthoVision™ XT10 software. Behavioral data were shown as total swim

distance (mm), average velocity (mm/s), average relative turn angle (degree), average

absolute turn angle (degree) and average absolute angular velocity (degree/s) at a

duration for 4 min.

For tactile stimulation, zebrafish larvae were placed in a concave slide, and tactile

stimulation was performed with glass capillaries to the head or tail sides. Larval

startle responses were recorded using a high-speed video camera (Mikrotron, EoSens

Mini1) at 1000 fps. Automated analysis of larval movement was performed using the

FLOTE software package (*Jain et al. 2014*).

For optogenetic studies, the Tg(Gal4^{s1020t}; *UAS*:ChR2) transgenic embryos were

injected with control, *epha4a* or *efnb3b* morpholinos at one cell stage. At 5 dpf, the

head of injected larva was mounted in 1% low melting-point agarose (Sigma) and the

body was exposed in glass-bottom dishes (WPI). Leica M165FC fluorescence

microscope was used to irradiate with a blue laser at 488nm wavelength. Larval

responses were recorded using a high-speed video camera (Mikrotron, EoSens Mini1)

at 500 fps. Body curvature analysis of larval movement was performed using the

FLOTE software package.

Whole-mount *in situ* hybridization and immunofluorescence

The primer sequences used to amplify *epha4a*, *epha4b* and *rfng* genes were listed

in Table S7. Probe synthesis and whole-mount in situ hybridization were performed

according to standard protocols. For reticulospinal neurons immunostaining, embryos

were fixed in 2% trichloroacetic acid at 48 hpf for 3-4 h, washed twice in 0.5% Triton

X-100 in PBS and blocked in 0.5% Triton X-100, 10% normal goat serum, 0.1%

bovine serum albumin (Solarbio) in PBS for 1 h. The embryos were stained by

monoclonal anti-neurofilament 160 antibody (Sigma-Aldrich) overnight at 4°C, then

stained by goat anti-mouse Alexa Fluor 488 (Invitrogen) after washing.

Analysis of neuronal calcium signals activity

Control, *epha4a* or *efnb3b* morpholinos were injected into Tg(*elavl3*:GAL4; UAS:GCaMP) transgenic embryos at one cell stage. At 24 hpf, embryos were paralyzed with 0.5 mg/ml α-Bungarotoxin (AlomoneLabs), then mounted in 1% low melting-point agarose (Sigma) in glass-bottom dishes (WPI). Neuronal calcium signal images were collected with IXON-L-888 EMCCD camera equipped on Dragonfly 200 Spinning Disk Confocal Microscope using a 20×/0.55 objective within 1 min at a frame rate of 10 fps.

Images were analyzed with imageJ software. To quantify the change in fluorescence intensity, a region of interest (ROI) was defined, and the fluorescence intensity F_t of different frames was normalized to F_t/F_{min} based on the minimum fluorescence intensity F_{min} in all frames. To compare the left-right alternation pattern of neuronal calcium signals, the ratio of the left-side calcium signals frequency (f_{left}) to the right-side calcium signals frequency (f_{right}) was logarithmically transformed (lg-transformed).

Quantification and statistical analysis

Statistical analyses were performed in SPSS (version 15.0). Unpaired Student's t-test, Welch one-way ANOVA or two-way ANOVA followed by Turkey's multiple comparison test were applied when appropriate. All experiments were replicated at least three times independently. $P<0.05$ was considered statistically significant.

Study approval

Approval for the study was obtained from the ethics committee at the Peking Union Medical College Hospital (JS-098, JS-2364), the medical ethics committee of the Keio University Hospital (No. 20080129), the ethical committee of RIKEN Yokohama Institute (No. H20-17(8)), and the Institutional Review Board of the University Texas Southwestern Medical Center (protocol STU 112010-150). Written informed consent was obtained from each participating individuals and families in

721   the three cohorts. For the control group, the protocols were approved by the ethics

722   committee at Peking Union Medical College Hospital. All zebrafish studies were

723   approved by the Animal Care Committee of the Ocean University of China (Animal

724   protocol number: OUC2012316).

Acknowledgments

We appreciate all the patients, their families and clinical research coordinators, including physicians who participated in this project. The authors also acknowledge the Texas Advanced Computing Center (TACC) at The University of Texas at Austin for providing computing resources that have contributed to the results related to the TSRHC cohort. URL: http://www.tacc.utexas.edu.

Funding

National Natural Science Foundation of China 81822030 (NW)

National Natural Science Foundation of China 82102522 (LW)

National Natural Science Foundation of China 31991194 & 32125015 (CZ)

National Natural Science Foundation of China 82172382 (TJZ)

CAMS Innovation Fund for Medical Sciences 2021-I2M-1-051 (TJZ, NW)

CAMS Innovation Fund for Medical Sciences 2021-I2M-1-052 (ZW)

CAMS Innovation Fund for Medical Sciences 2020-I2M-C&T-B-030 (TJZ)

Beijing Natural Science Foundation 7222133 (SW)

Non-profit Central Research Institute Fund of Chinese Academy of Medical Sciences No. 2019PT320025 (NW)

National High Level Hospital Clinical Research Funding 2022-PUMCH-D-004 (TJZ)

National High Level Hospital Clinical Research Funding 2022-PUMCH-C-033 (NW)

Shandong Natural Science Foundation ZR202102210113 (LW)

Shandong Province Taishan Scholar Project (LW)

Gabriella Miller Kids First Program grant X01 HL132375-01A1 (JR)

Data and materials availability

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

References

Andersson, L. S., M. Larhammar, F. Memic, H. Wootz, D. Schwochow, C.-J. Rubin, K. Patra,

T. Arnason, L. Wellbring & G. Hjälm (2012) Mutations in DMRT3 affect locomotion in

horses and spinal circuit function in mice. *Nature,* 488**,** 642-646.

Auton, A., L. D. Brooks, R. M. Durbin, E. P. Garrison, H. M. Kang, J. O. Korbel, J. L.

Marchini, S. McCarthy, G. A. McVean & G. R. Abecasis (2015) A global reference for

human genetic variation. *Nature,* 526**,** 68-74.

Bagnat, M. & R. S. Gray (2020) Development of a straight vertebrate body axis.

Development, 147.

Borgius, L., H. Nishimaru, V. Caldeira, Y. Kunugise, P. Löw, R. Reig, S. Itohara, T. Iwasato

& O. Kiehn (2014) Spinal glutamatergic neurons defined by EphA4 signaling are essential

components of normal locomotor circuits. *J. Neurosci.,* 34**,** 3841-3853.

Boswell, C. W. & B. Ciruna (2017) Understanding idiopathic scoliosis: a new zebrafish

school of thought. *Trends Genet.,* 33**,** 183-196.

Buchan, J., D. Alvarado, G. Haller, C. Cruchaga, M. Harms, T. Zhang, M. Willing, D.

Grange, A. Braverman, N. Miller, J. Morcuende, N. Tang, T. Lam, B. Ng, J. Cheng, M.

Dobbs & C. Gurnett (2014) Rare variants in FBN1 and FBN2 are associated with severe

adolescent idiopathic scoliosis. *Hum. Mol. Genet.,* 23**,** 5271-82.

Cavone, L., T. McCann, L. K. Drake, E. A. Aguzzi, A.-M. Oprişoreanu, E. Pedersen, S.

Sandi, J. Selvarajah, T. M. Tsarouchas & D. Wehner (2021) A unique macrophage

subpopulation signals directly to progenitor cells to promote regenerative neurogenesis in

the zebrafish spinal cord. *Dev. cell,* 56**,** 1617-1630. e6.

Cayuso, J., Q. Xu, M. Addison & D. G. Wilkinson (2019) Actomyosin regulation by Eph

receptor signaling couples boundary cell formation to border sharpness. *Elife,* 8.

Chen, N., S. Zhao, A. Jolly, L. Wang, H. Pan, J. Yuan, S. Chen, A. Koch, C. Ma & W. Tian

(2021) Perturbations of genes essential for Müllerian duct and Wölffian duct development in

Mayer-Rokitansky-Küster-Hauser syndrome. *Am. J. Hum. Genet.,* 108**,** 337-345.

Cheng, J. C., R. M. Castelein, W. C. Chu, A. J. Danielsson, M. B. Dobbs, T. B. Grivas, C. A.

Gurnett, K. D. Luk, A. Moreau, P. O. Newton, I. A. Stokes, S. L. Weinstein & R. G. Burwell

(2015) Adolescent idiopathic scoliosis. *Nat. Rev. Dis. Primers,* 1**,** 15030.

Consortium, G. (2020) The GTEx Consortium atlas of genetic regulatory effects across

human tissues. *Science,* 369**,** 1318-1330.

Cooke, J. E., H. A. Kemp & C. B. Moens (2005) EphA4 is required for cell adhesion and

rhombomere-boundary formation in the zebrafish. *Curr Biol,* 15**,** 536-42.

Das, S., L. Forer, S. Schönherr, C. Sidore, A. E. Locke, A. Kwong, S. I. Vrieze, E. Y. Chew,

S. Levy & M. McGue (2016) Next-generation genotype imputation service and methods.

Nat. Genet., 48**,** 1284-1287.

de Leeuw, C. A., J. M. Mooij, T. Heskes & D. Posthuma (2015) MAGMA: generalized

gene-set analysis of GWAS data. *PLoS Comput. Biol.,* 11**,** e1004219.

Delaneau, O., J. Marchini & J.-F. Zagury (2012) A linear complexity phasing method for

thousands of genomes. *Nat. Methods,* 9**,** 179-181.

Ding, L., Y. Shen, J. Ni, Y. Ou, Y. Ou & H. Liu (2017) EphA4 promotes cell proliferation

and cell adhesion–mediated drug resistance via the AKT pathway in multiple myeloma.

Tumour Biol., 39**,** 1010428317694298.

Egea, J. & R. Klein (2007) Bidirectional Eph-ephrin signaling during axon guidance. *Trends*

Cell Biol, 17**,** 230-8.

Fan, Y.-H., Y.-Q. Song, D. Chan, Y. Takahashi, S. Ikegawa, M. Matsumoto, I. Kou, K. S.

Cheah, P. Sham & K. Cheung (2012) SNP rs11190870 near LBX1 is associated with

adolescent idiopathic scoliosis in southern Chinese. *J. Hum. Genet.,* 57**,** 244-246.

Fidelin, K., L. Djenoune, C. Stokes, A. Prendergast, J. Gomez, A. Baradel, F. Del Bene & C.

Wyart (2015) State-dependent modulation of locomotion by GABAergic spinal sensory

neurons. *Curr. Biol.,* 25**,** 3035-3047.

Flanagan, J. G. & P. Vanderhaeghen (1998) The ephrins and Eph receptors in neural

development. *Annu Rev Neurosci,* 21**,** 309-45.

Fu, W.-Y., Y. Chen, M. Sahin, X.-S. Zhao, L. Shi, J. B. Bikoff, K.-O. Lai, W.-H. Yung, A. K.

Fu & M. E. Greenberg (2007) Cdk5 regulates EphA4-mediated dendritic spine retraction

through an ephexin1-dependent mechanism. *Nat. Neurosci.,* 10**,** 67-76.

Gao, X., D. Gordon, D. Zhang, R. Browne, C. Helms, J. Gillum, S. Weber, S. Devroy, S.

Swaney, M. Dobbs, J. Morcuende, V. Sheffield, M. Lovett, A. Bowcock, J. Herring & C.

Wise (2007) CHD7 gene polymorphisms are associated with susceptibility to idiopathic

scoliosis. *Am. J. Hum. Genet.,* 80**,** 957-65.

Grimes, D. T., C. W. Boswell, N. F. Morante, R. M. Henkelman, R. D. Burdine & B. Ciruna

(2016) Zebrafish models of idiopathic scoliosis link cerebrospinal fluid flow defects to spine

curvature. *Science,* 352**,** 1341-4.

Guo, L., H. Yamashita, I. Kou, A. Takimoto, M. Meguro-Horike, S. Horike, T. Sakuma, S.

Miura, T. Adachi, T. Yamamoto, S. Ikegawa, Y. Hiraki & C. Shukunami (2016) Functional

Investigation of a Non-coding Variant Associated with Adolescent Idiopathic Scoliosis in

Zebrafish: Elevated Expression of the Ladybird Homeobox Gene Causes Body Axis

Deformation. *PLoS Genet.,* 12**,** e1005802.

Hale, M. E., H. R. Katz, M. Y. Peek & R. T. Fremont (2016) Neural circuits that drive startle

behavior, with a focus on the Mauthner cells and spiral fiber neurons of fishes. *J Neurogenet,*

30**,** 89-100.

Haller, G., D. Alvarado, K. Mccall, P. Yang, C. Cruchaga, M. Harms, A. Goate, M. Willing, J.

Morcuende, E. Baschal, N. Miller, C. Wise, M. Dobbs & C. Gurnett (2016) A polygenic

burden of rare variants across extracellular matrix genes among individuals with adolescent

idiopathic scoliosis. *Hum. Mol. Genet.,* 25**,** 202-9.

Hresko, M. (2013) Clinical practice. Idiopathic scoliosis in adolescents. *N. Engl. J. Med.,*

368**,** 834-41.

Ioannidis, N. M., J. H. Rothstein, V. Pejaver, S. Middha, S. K. McDonnell, S. Baheti, A.

Musolf, Q. Li, E. Holzinger & D. Karyadi (2016) REVEL: an ensemble method for

predicting the pathogenicity of rare missense variants. *Am. J. Hum. Genet.,* 99**,** 877-885.

Iwasato, T., H. Katoh, H. Nishimaru, Y. Ishikawa, H. Inoue, Y. M. Saito, R. Ando, M. Iwama,

R. Takahashi & M. Negishi (2007) Rac-GAP α-chimerin regulates motor-circuit formation

as a key mediator of EphrinB3/EphA4 forward signaling. *Cell,* 130**,** 742-753.

Jain, R. A., H. Bell, A. Lim, C.-B. Chien & M. Granato (2014) Mirror movement-like

defects in startle behavior of zebrafish dcc mutants are caused by aberrant midline guidance

of identified descending hindbrain neurons. *J. Neurosci.,* 34**,** 2898-2909.

Kania, A. & R. Klein (2016) Mechanisms of ephrin–Eph signalling in development,

physiology and disease. *Nat. Rev. Mol. Cell Biol.,* 17**,** 240-256.

Karczewski, K. J., L. C. Francioli, G. Tiao, B. B. Cummings, J. Alfoldi, Q. Wang, R. L.

Collins, K. M. Laricchia, A. Ganna, D. P. Birnbaum, L. D. Gauthier, H. Brand, M.

Solomonson, N. A. Watts, D. Rhodes, M. Singer-Berk, E. M. England, E. G. Seaby, J. A.

Kosmicki, R. K. Walters, K. Tashman, Y. Farjoun, E. Banks, T. Poterba, A. Wang, C. Seed, N.

Whiffin, J. X. Chong, K. E. Samocha, E. Pierce-Hoffman, Z. Zappala, A. H.

O'Donnell-Luria, E. V. Minikel, B. Weisburd, M. Lek, J. S. Ware, C. Vittal, I. M. Armean, L.

Bergelson, K. Cibulskis, K. M. Connolly, M. Covarrubias, S. Donnelly, S. Ferriera, S.

Gabriel, J. Gentry, N. Gupta, T. Jeandet, D. Kaplan, C. Llanwarne, R. Munshi, S. Novod, N.

Petrillo, D. Roazen, V. Ruano-Rubio, A. Saltzman, M. Schleicher, J. Soto, K. Tibbetts, C.

Tolonen, G. Wade, M. E. Talkowski, C. Genome Aggregation Database, B. M. Neale, M. J.

Daly & D. G. MacArthur (2020) The mutational constraint spectrum quantified from

variation in 141,456 humans. *Nature,* 581**,** 434-443.

Kiehn, O. (2006) Locomotor circuits in the mammalian spinal cord. *Annu Rev Neurosci,* 29**,**

279-306.

Kircher, M., D. M. Witten, P. Jain, B. J. O'roak, G. M. Cooper & J. Shendure (2014) A

general framework for estimating the relative pathogenicity of human genetic variants. *Nat.*

Genet., 46**,** 310-315.

Kou, I., N. Otomo, K. Takeda, Y. Momozawa, H. Lu, M. Kubo, Y. Kamatani, Y. Ogura, Y.

Takahashi, M. Nakajima, S. Minami, K. Uno, N. Kawakami, M. Ito, I. Yonezawa, K.

Watanabe, T. Kaito, H. Yanagida, H. Taneichi, K. Harimaya, Y. Taniguchi, H. Shigematsu, T.

Iida, S. Demura, R. Sugawara, N. Fujita, M. Yagi, E. Okada, N. Hosogane, K. Kono, M.

Nakamura, K. Chiba, T. Kotani, T. Sakuma, T. Akazawa, T. Suzuki, K. Nishida, K. Kakutani,

T. Tsuji, H. Sudo, A. Iwata, T. Sato, S. Inami, M. Matsumoto, C. Terao, K. Watanabe & S.

Ikegawa (2019) Genome-wide association study identifies 14 previously unreported

susceptibility loci for adolescent idiopathic scoliosis in Japanese. *Nat. Commun.,* 10**,** 3685.

Kou, I., Y. Takahashi, T. Johnson, A. Takahashi, L. Guo, J. Dai, X. Qiu, S. Sharma, A.

Takimoto, Y. Ogura, H. Jiang, H. Yan, K. Kono, N. Kawakami, K. Uno, M. Ito, S. Minami,

H. Yanagida, H. Taneichi, N. Hosono, T. Tsuji, T. Suzuki, H. Sudo, T. Kotani, I. Yonezawa,

D. Londono, D. Gordon, J. Herring, K. Watanabe, K. Chiba, N. Kamatani, Q. Jiang, Y.

Hiraki, M. Kubo, Y. Toyama, T. Tsunoda, C. Wise, Y. Qiu, C. Shukunami, M. Matsumoto &

S. Ikegawa (2013) Genetic variants in GPR126 are associated with adolescent idiopathic

scoliosis. *Nat. Genet.,* 45**,** 676-9.

Kramers-de Quervain, I. A., R. Muller, A. Stacoff, D. Grob & E. Stussi (2004) Gait analysis

in patients with idiopathic scoliosis. *Eur. Spine J.,* 13**,** 449-56.

Kullander, K., S. J. Butt, J. M. Lebret, L. Lundfald, C. E. Restrepo, A. Rydström, R. Klein &

O. Kiehn (2003) Role of EphA4 and EphrinB3 in local neuronal circuits that control walking.

Science, 299**,** 1889-1892.

Li, C., R. Chen, X. Fan, J. Luo, J. Qian, J. Wang, B. Xie, Y. Shen & S. Chen (2015) EPHA4

haploinsufficiency is responsible for the short stature of a patient with 2q35-q36.2 deletion

and Waardenburg syndrome. *BMC Med. Genet.,* 16**,** 23.

Li, W., Y. Li, L. Zhang, H. Guo, D. Tian, Y. Li, Y. Peng, Y. Zheng, Y. Dai, K. Xia, X. Lan, B.

Wang & Z. Hu (2016) AKAP2 identified as a novel gene mutated in a Chinese family with

adolescent idiopathic scoliosis. *J. Med. Genet.,* 53**,** 488-93.

Luk, K. D., C. F. Lee, K. M. Cheung, J. C. Cheng, B. K. Ng, T. P. Lam, K. H. Mak, P. S. Yip

& D. Y. Fong (2010) Clinical effectiveness of school screening for adolescent idiopathic

scoliosis: a large population-based retrospective cohort study. *Spine,* 35**,** 1607-14.

Marchini, J., B. Howie, S. Myers, G. McVean & P. Donnelly (2007) A new multipoint

method for genome-wide association studies by imputation of genotypes. *Nat. Genet.,* 39**,**

906-913.

Marder, E. & D. Bucher (2001) Central pattern generators and the control of rhythmic

movements. *Curr Biol,* 11**,** R986-96.

Marees, A. T., H. de Kluiver, S. Stringer, F. Vorspan, E. Curis, C. Marie-Claire & E. M.

Derks (2018) A tutorial on conducting genome-wide association studies: Quality control and

statistical analysis. *Int. J. Methods Psychiatr. Res.,* 27**,** e1608.

Miller, N. (2007) Genetics of familial idiopathic scoliosis. *Clin. Orthop. Relat. Res.,* 462**,**

6-10.

Miller, N., C. Justice, B. Marosy, K. Doheny, E. Pugh, J. Zhang, H. Dietz & A. Wilson (2005)

Identification of candidate regions for familial idiopathic scoliosis. *Spine,* 30**,** 1181-7.

Mukhopadhyay, N., M. Bishop, M. Mortillo, P. Chopra, J. Hetmanski, M. Taub, L. Moreno,

L. Valencia-Ramirez, C. Restrepo, G. Wehby, J. Hecht, F. Deleyiannis, A. Butali, S.

Weinberg, T. Beaty, J. Murray, E. Leslie, E. Feingold & M. Marazita (2020) Whole genome

sequencing of orofacial cleft trios from the Gabriella Miller Kids First Pediatric Research

Consortium identifies a new locus on chromosome 21. *Hum. Genet.,* 139**,** 215-226.

Nishida, M., T. Nagura, N. Fujita, N. Hosogane, T. Tsuji, M. Nakamura, M. Matsumoto & K.

Watanabe (2017) Position of the major curve influences asymmetrical trunk kinematics

during gait in adolescent idiopathic scoliosis. *Gait Posture,* 51**,** 142-148.

Ogura, Y., I. Kou, S. Miura, A. Takahashi, L. Xu, K. Takeda, Y. Takahashi, K. Kono, N.

Kawakami & K. Uno (2015) A functional SNP in BNC2 is associated with adolescent

idiopathic scoliosis. *Am. J. Hum. Genet.,* 97**,** 337-342.

Shamah, S., M. Lin, J. Goldberg, S. Estrach, M. Sahin, L. Hu, M. Bazalakova, R. Neve, G.

Corfas, A. Debant & M. Greenberg (2001) EphA receptors regulate growth cone dynamics

through the novel guanine nucleotide exchange factor ephexin. *Cell,* 105**,** 233-44.

Shimode, M., A. Ryouji & N. Kozo (2003) Asymmetry of premotor time in the back muscles

of adolescent idiopathic scoliosis. *Spine,* 28**,** 2535-2539.

Szklarczyk, D., A. L. Gable, D. Lyon, A. Junge, S. Wyder, J. Huerta-Cepas, M. Simonovic,

N. T. Doncheva, J. H. Morris, P. Bork, L. J. Jensen & C. V. Mering (2019) STRING v11:

protein-protein association networks with increased coverage, supporting functional

discovery in genome-wide experimental datasets. *Nucleic Acids Res,* 47**,** D607-D613.

Takahashi, Y., I. Kou, A. Takahashi, T. A. Johnson, K. Kono, N. Kawakami, K. Uno, M. Ito,

S. Minami & H. Yanagida (2011) A genome-wide association study identifies common

variants near LBX1 associated with adolescent idiopathic scoliosis. *Nat. Genet.,* 43**,**

1237-1240.

Talpalar, A. E., J. Bouvier, L. Borgius, G. Fortin, A. Pierani & O. Kiehn (2013) Dual-mode

operation of neuronal networks involved in left-right alternation. *Nature,* 500**,** 85-8.

Tang, N. L., M. B. Dobbs, C. A. Gurnett, Y. Qiu, T. Lam, J. C. Cheng & N. Hadley-Miller

(2021) A Decade in Review after Idiopathic Scoliosis Was First Called a Complex Trait—A

Tribute to the Late Dr. Yves Cotrel for His Support in Studies of Etiology of Scoliosis.

Genes, 12**,** 1033.

Tassabehji, M., A. P. Read, V. E. Newton, M. Patton, P. Gruss, R. Harris & T. Strachan (1993)

Mutations in the PAX3 gene causing Waardenburg syndrome type 1 and type 2. *Nat. Genet.,*

3**,** 26-30.

Valentino, B., L. Maccauro, G. Mango, F. Melito & A. Fabozzo (1985) Electromyography

for the investigation and early diagnosis of scoliosis. *Anat. Clin.,* 7**,** 55-9.

Watanabe, K., E. Taskesen, A. van Bochoven & D. Posthuma (2017) Functional mapping

and annotation of genetic associations with FUMA. *Nat. Commun.,* 8**,** 1826.

Weinstein, S., L. Dolan, J. Wright & M. Dobbs (2013) Effects of bracing in adolescents with

idiopathic scoliosis. *N. Engl. J. Med.,* 369**,** 1512-21.

Weinstein, S. L., L. A. Dolan, K. F. Spratt, K. K. Peterson, M. J. Spoonamore & I. V. Ponseti

(2003) Health and function of patients with untreated idiopathic scoliosis: a 50-year natural

history study. *JAMA,* 289**,** 559-67.

Willer, C. J., Y. Li & G. R. Abecasis (2010) METAL: fast and efficient meta-analysis of

genomewide association scans. *Bioinformatics,* 26**,** 2190-1.

Wu, M. Y., M. Carbo-Tano, O. Mirat, F. X. Lejeune, J. Roussel, F. B. Quan, K. Fidelin & C.

Wyart (2021) Spinal sensory neurons project onto the hindbrain to stabilize posture and

enhance locomotor speed. *Curr. Biol.,* 31**,** 3315-3329 e5.

Wyart, C., F. Del Bene, E. Warp, E. K. Scott, D. Trauner, H. Baier & E. Y. Isacoff (2009)

Optogenetic dissection of a behavioural module in the vertebrate spinal cord. *Nature,* 461**,**

407-10.

Xie, H., M. Li, Y. Kang, J. Zhang & C. Zhao (2022) Zebrafish: an important model for

understanding scoliosis. *Cell. Mol. Life. Sci.,* 79**,** 1-16.

Yao, E. & J. E. Tavis (2005) A general method for nested RT-PCR amplification and

sequencing the complete HCV genotype 1 open reading frame. *Virol. J.,* 2**,** 1-9.

Zhao, S., Y. Zhang, W. Chen, W. Li, S. Wang, L. Wang, Y. Zhao, M. Lin, Y. Ye & J. Lin

(2021) Diagnostic yield and clinical impact of exome sequencing in early-onset scoliosis

(EOS). *J. Med. Genet.,* 58**,** 41-47.

Zhou, W., W. Bi, Z. Zhao, K. K. Dey, K. A. Jagadeesh, K. J. Karczewski, M. J. Daly, B. M.

Neale & S. Lee (2022) SAIGE-GENE+ improves the efficiency and accuracy of set-based

rare variant association tests. *Nat. Genet.,* 54**,** 1466-1469.

Zhu, Z., N. Tang, L. Xu, X. Qin, S. Mao, Y. Song, L. Liu, F. Li, P. Liu, L. Yi, J. Chang, L.

Jiang, B. Ng, B. Shi, W. Zhang, J. Qiao, X. Sun, X. Qiu, Z. Wang, F. Wang, D. Xie, L. Chen,

Z. Chen, M. Jin, X. Han, Z. Hu, Z. Zhang, Z. Liu, F. Zhu, B. Qian, Y. Yu, B. Wang, K. Lee,

W. Lee, T. Lam, Y. Qiu & J. Cheng (2015) Genome-wide association study identifies new

susceptibility loci for adolescent idiopathic scoliosis in Chinese girls. *Nat. Commun.,* 6**,**

8355.

Table 1. Dominantly inherited variants identified in *EPHA4* and *NGEF*.

Patient ID	Chr	Gene	Ethnicity	Inheritance pattern	cDNA change	AA change	Variant type	ExAC PLI	CADD	GnomAD frequency	In-house frequency
SCO2003P0846	2	*EPHA4*	Chinese	De novo	c.1443+1G>C	NA	Splice donor	1	21.4	0	0
SCO2003P2146	2	*EPHA4*	Chinese	De novo	c.2546G>A	p.Cys849Tyr	Missense	1	28.2	0	0
SCO2003P2080	2	*EPHA4*	Chinese	De novo	c.1318+10344A>G	NA	Intronic	1	NA	0	0
SCO2003P3202	2	*EPHA4*	Chinese	De novo	2q35-36.2 4.6 Mb deletion	NA	CNV	NA	NA	0	0
SCO2003P3332	2	*NGEF*	Chinese	De novo	c.1A>G	p.Met1?	Start lost	0.95	12.8	0	0
TSRHC01	2	*NGEF*	Non-Hispanic White	AD	c.857C>T	p.Ala286Val	Missense	0.95	29.6	0.00002	0

Abbreviations: Chr (chromosomal localization); AA (amino acid); AD (autosomal dominant); CNV (copy number variant); ExAC PLI (probability of being loss-of-function intolerant from Exome Aggregation Consortium); CADD (Combined Annotation Dependent Depletion score); gnomAD (Genome Aggregation Database); NA (not applicable).

970
Figure 1. Clinical and genetic information on IS patients and functional effect of
***EPHA4* variants. A:** Locations of three *EPHA4* single nucleotide variants relative to
the protein domains (top panel) and exons 1-18 (bottom panel). **B-E:** Pedigrees and
spinal radiographs of four probands with dominant gene variants. Sanger sequencing
confirmed the variants. The arrows indicate the probands. The term +/+ denotes the
wild-type, and cDNA change/+ denotes the heterozygous variant. **F:** Western blot
analysis of *EPHA4*-c.2546G>A variant showing the protein expression levels of
EPHA4 and *CDK5* and the amount of phosphorylated CDK5 (pCDK5) expressed
from the *EPHA4*-mutant plasmid. WT: wild type. **G:** Schematic representation of the

effect of the *EPHA4*-c.1443+1G>C mutation on the splicing process. This variant

induced a new splicing site (red box). The yellow box indicates the splicing acceptor.

H: Schematic representation of the effect of the *EPHA4*-c.1318+10344A>G mutation

on the splicing process. This variant induced a new splicing site (red box). The yellow

box indicates the splicing acceptor. **I:** The minigene assay result showed that the

c.1443+1G>C variant introduced a new splicing site, resulting in a 36-bp in-frame

deletion in exon 6. **J:** The nested PCR showed that the c.1318+10344A>G variant

induced exon 5 skipping, resulting in a 339-bp in-frame deletion. **K:** NCBI RefSeq

genes included in 2q35-q36.2 from UCSC Genome Browser. *EPHA4* is shown by the

red oval.

990
Figure 2. Scoliosis in zebrafish *epha4a* mutants. A: Diagram of the protein domains,
genomic structures, and sequences of wild-type and corresponding *epha4a* mutants.
Red arrows indicate mutation sites. Blue boxes indicate open reading frames.
Underlined sequences indicate the protospacer adjacent motif (PAM) region, and red
fonts indicate Cas9 binding sites. LBD: ligand binding domain; FN3: fibronectin type
3 domain; TMD: transmembrane domain; PTKc: catalytic domain of the protein
tyrosine kinases; SAM: sterile alpha motif. **B:** Sanger sequencing results confirmed
the deletion of the target region in *epha4a* mutant transcripts. **C:** Bar graph showing
the percentages of adult zebrafish with normal, mild, or severe body axis defects in
wild-type (n=76), *epha4a* heterozygote (n=95), or *epha4a* mutants (n=116). **D:**
Representative images of wild-type and *epha4a* mutants. Micro CT images are shown
on the right. Lateral and dorsal views are shown. Scale bar: 1 cm.

1003
Figure 3. Abnormal left-right swimming pattern in the absence of Epha4a. A:
Dot plots showing the swimming distance of each 8 dpf larva at a duration of 4 min
(N=54 for wild-type and N=60 for *epha4a* mutants). **B:** Dot plots showing the
swimming velocity of wild-type and mutant larvae as indicated. **C:** Diagram showing
the turning angle (θ) of the larvae during swimming. **D:** Scatter plot showing the
relative turning angle of wild-type and mutant larvae. The relative angles were
calculated by the sum of turning angles during fish swimming with left (positive) or
right (negative) turns. The *epha4a* mutants favored turning to one side of their
directions compared with those of wild-type larvae. **E:** Dot plots showing the average
absolute turning angle of wild-type and mutant larvae as indicated. **F:** Dot plots
showing the average absolute angular velocity of wild-type and mutant larvae as
indicated. **G:** Bar graph showing the percentages of 5 dpf zebrafish larvae with
different reactions after tactile stimulation. N=10 for each group; the numbers of
tactile stimulations are indicated in each column. **H:** Representative images of the
total body curvature measurements in zebrafish larvae, with values as the sum of α
and β angles shown in the figure. **I:** Representative time-series images of 5 dpf
wild-type and *epha4a* mutant zebrafish larvae after tactile stimulation to the head.
Each panel represents the points of maximal body curvature for the C-bend and
counterbend after the tactile startle response. **J, K:** A plot of body curvature angles as
measured in panel (H) during swimming in response to tactile head stimulation in 5
dpf wild-type (N=5 larvae, n=10 stimuli) and *epha4a* mutants (N=5 larvae, n=7
stimuli). **L:** The maximum curvature angles during the first C-bend and counterbend
after tactile stimulation in wild-type and *epha4a* mutant larvae.

Figure 4. Uncoordinated left-right activation of spinal cord neurons in the absence of Epha4a. A, A': Fluorescent images showing the dorsal-view of 24 hpf Tg(*elavl3*:GAL4; *UAS*:GCaMP) double transgenic larvae. The corresponding movies are shown in Movie S5 and S6. **B, B':** Line charts showing the quantification of fluorescence changes of the region of interests (ROIs, circled in A, A') in control and *epha4a* morphants. **C:** Scatter plot showing the distribution trend of the ratio of the calcium signal frequency between left and right in control (N=15 larvae, n=38 experiments) and *epha4a* morphants (N=15 larvae, n=41 experiments). **D:** Statistical graph of the ratio of the calcium signal frequency between left and right in control and *epha4a* morphants. Scale bars: 50 μm in panel **(A, A')**.

Figure 5. Disorganized neural patterning in *epha4a* mutants. A: Confocal images showing reticulospinal neuronal axons in 48 hours post-fertilization (hpf) wild-type and *epha4a* mutant larvae visualized with anti-neurofilament antibody RMO44. Asterisks indicate the cross sites of Mauthner axons. The green arrow indicates the cell body of the r5/MiD2 neuron in a wild-type larva. The red arrow points to the ipsilaterally projected axon of r6/MiD3 in the mutant larva, which is normally projected to the other side in wild-type fish. **B:** Statistical chart showing the distance between r4 and r7 of 48 hpf wild-type and *epha4a* mutants. **C:** Scatter plot showing the distance (d) between the center line and the intersection site of Mauthner axons as indicated in panel C′. **D:** The ratio of the distance between the intersection site of Mauthner axons and bilateral axon bundles in 48 hpf wild-type (N=11 larvae) and *epha4a* mutants (N=12 larvae). The ratios were calculated as in panel D′. Scale bars: 20 μm in panel **(A)**.

1051
Figure 6. Aberrant swimming as a result of abnormal extension of CSF-cNs
axons. A: Fluorescent images showing the distribution of ascending axons of
CSF-cNs marked by Tg(pkd2l1:GAL4,UAS:Kaede) in 2 dpf *epha4a* mutant larvae.
Yellow line indicates the midline and the red arrows indicate aberrantly extended
axons in *epha4a*[+/-] and *epha4a*[-/-] larvae. **B:** Bar graph showing the percentages of
abnormal extension of CSF-cNs axons in 2 dpf wild-type and *epha4a* mutant; the
numbers of larvae are indicated in each column. **C, C':** Superimposed frames of tail
oscillations in 5 dpf control and *epha4a* morphants. **D, D':** A plot of body curvature
angles in panel (C) and (C'). **E:** Percentages of tendentious swing in control (N=8
larvae, n=24 experiments) and *epha4a* morphants (N=11 larvae, n=33 experiments).
The percentages were calculated by the ratio of tendentious tail oscillation during the
first eight swings. Scale bars: 50 μm in panel **(A)**.

1064
1065 **Figure 7. IS patients with potential *NGEF* variants. A:** Protein structure of NGEF
1066 protein with the position of potential variants. **B, C:** Pedigrees and spinal radiographs
1067 of two probands with dominant gene variants. Sanger sequencing results are shown on
1068 the right.

1069
Figure 8. The proposed mechanism of IS mediated by *EPHA4* dysfunction. In
healthy individuals, EphrinB3-activated EPHA4 phosphorylates CDK5, leading to the
phosphorylation of Ephexin, a protein encoded by NGEF. Phosphorylated Ephexin
can regulate axon guidance through either activating RHOA or suppressing CDC42
and RAC1 signaling . These processes are critical to maintain the normal function of
the CPG, the local neural network that provides coordinated bilateral muscle control.
Impairment of the *EPHA4* pathway and CPG may cause an imbalance of the motor
drive from the spinal cord during development, thus causing the uncoordinated
left/right swimming behavior in zebrafish larvae and the asymmetry of the bilateral
muscular pull in a young child. Although the appearance is normal in early childhood,
the dysfunction produces a scoliotic curve during the growth spurt.

Supplementary Materials for

Impaired central pattern generators due to abnormal EPHA4 signaling leads to idiopathic scoliosis

Lianlei Wang *et al.*

Corresponding author:

Nan Wu, M.D., E-mail: dr.wunan@pumch.cn

Chengtian Zhao, Ph.D., E-mail: chengtian_zhao@ouc.edu.cn

Terry Jianguo Zhang, M.D., E-mail: zhangjianguo@pumch.cn

Supplementary Fig. 1. Flowchart for identification of causative genes. The processes of candidate gene mapping and gene-based burden analysis were described. The variants in *EPHA4* and *NGEF* identified in each cohort were displayed. Abbreviations: SNP (single nucleotide polymorphism); eQTL (expression quantitative trait locus); IS (idiopathic scoliosis).

A

B

1097   **Supplementary Fig. 2. The splicing analysis by the Alamut software.** The results
1098   of Alamut software by four algorithms (SpliceSiteFinder-like, MaxEntScan,
1099   NNSPLICE, and GeneSplicer).

Supplementary Fig. 3. Phenotypes of *epha4a* mutants. A: Representative images showing the notochord and external phontypes of wild-type and *epha4a* mutants. The red dashed lines indicate the area of the notochord. **B:** Whole-mount in situ hybridization results showing the expression of *rfng* gene in 18 hpf wild type, mutant or morphant larvae as indicated. *rfng* gene marks boundary cells in the hindbrain. The red arrows indicate boundary expression. The numbers of embryos analyzed are shown on top right. Scale bars: 200 μm in panel (A), 100 μm in panel (B).

Supplementary Fig. 4. Zebrafish *epha4b* mutants exhibited body axis defects during development. A: Diagram showing the protein domains, genomic structures, and sequences of the wild-type and corresponding *epha4b* mutants. Red arrows indicate mutation sites. **B:** Representative images of *epha4b* mutants. **C:** Bar graph showing the percentages of adult fish with normal and body axis defects in wild-type (n=39) and *epha4b* mutants (n=43). **D:** Bright-field and GFP fluorescent images showing scoliosis *epha4b* mutants at 28 dpf as indicated by *Tg(Ola.Sp7:NLS-GFP)* transgene, which labels the bone skeleton. Scale bars: 1 cm in panel (B), 2 mm in panel (D).

Supplementary Fig. 5. Expression pattern of *epha4* and *efnb3* in the spinal cord.
A: Whole-mount in situ hybridization results showing the expression of *epha4a* and *epha4b* genes in the spinal cord in 24 dpf zebrafish larvae. **B:** Violin plot showing *epha4a, epha4b, efnb3a,* and *efnb3b* gene expression in different cell types. Scale bars: 200 μm in panel (A).

Supplementary Fig. 6. Left-right coordination defects in *efnb3b* morphants. **A:** Fluorescent images showing the distribution of ascending axons of CSF-cNs marked by Tg*(pkd2l1:GAL4,UAS:Kaede)* in 2 dpf *efnb3b* morphants. The red arrows indicate aberrantly extended axons in *efnb3b* morphants. **B:** Fluorescent images showing the dorsal-view of 24 hpf Tg(*elavl3*:GAL4; *UAS*:GCaMP) transgenic larvae. **C:** The line chart showing the quantification of fluorescence changes of the ROIs in control morphants and *efnb3b* morphants as indicated in panel B. **D:** Superimposed frames of tail oscillations in 5dpf control and *efnb3b* morphants. **E:** A plot of body curvature angles in panel (D). Scale bars: 50 μm in panel (A, B).

1130
Supplementary Fig. 7. Proteins in STRING that interact with EPHA4. Protein
interaction network analysis, as illustrated by STRING v11.5, was used to identify
rare variants in genes related to *EPHA4*. Known and predicted protein-protein
interactions are included in the STRING database. Multiple proteins, including
Ephexin, are associated with EPHA4 physically or functionally. Ephexin, encoded by
NGEF, is a neuronal guanine exchange factor.

Supplementary Table 1. Summary of the 14 studies and their corresponding SNPs included in the candidate genes mapping.

Citation	SNPs
Fan et al., J Hum. Genet. 2012	rs11190870 (p= 9.1×10^{-10})
Zhu et al., Hum. Mol. Genet. 2017	rs7593846 (p= 1.19×10^{-13})
	rs7633294 (p= 1.85×10^{-12})
	rs6570507 (p= 1.14×10^{-11})
	rs6047663 (p= 1.61×10^{-15})
Zhu et al., Nat. Commun. 2015	rs678741 (p= 9.68×10^{-37})
	rs4940576 (p= 2.22×10^{-12})
	rs13398147 (p= 7.59×10^{-13})
	rs241215 (p= 2.95×10^{-9})
Takahashi et al., Nat. Genet. 2011	rs11190870 (p= 1.24×10^{-19})
	rs625039 (p= 8.13×10^{-15})
	rs11598564 (p= 5.98×10^{-14})
Sharma et al., Hum. Mol. Genet. 2011	rs1400180 (p= 6.35×10^{-8})
Kou et al., Nat. Genet. 2013	rs6570507 (p= 1.27×10^{-14})
Xu et al., Spine Deform. 2018	rs169311 (p= 2.10×10^{-8})
Grauers et al., Spine J. 2015	rs11190870 (p= 7.0×10^{-18})
Ogura et al., Am. J of Hum. Genet. 2015	rs3904778 (p= 2.46×10^{-13})
Khanshour et al., Hum. Mol. Genet. 2018	rs4513093 (p= 1.71×10^{-15})
	rs1455114 (p= 2.99×10^{-8})
	rs687621 (p= 7.29×10^{-10})
	rs10756785 (p= 7.00×10^{-10})
Sharma et al., Nat. Commun. 2015	rs6137473 (p= 3.12×10^{-8})
Miyake et al., PLoS One 2013	rs11190870 (p= 2.80×10^{-18})
	rs625039 (p= 1.28×10^{-15})
	rs12946942 (p= 6.43×10^{-12})
	rs11598564 (p= 9.77×10^{-12})
	rs6570507 (p= 3.78×10^{-8})
	rs9496346 (p= 1.00×10^{-8})

Chettier et al., PLoS One 2015	rs11190878 (p= 4.18×10^{-9})
Kou et al., Nat. Commun. 2019	rs9389985 (p= 3.51×10^{-20})
	rs7028900 (p= 2.19×10^{-17})
	rs144131194 (p= 1.35×10^{-11})
	rs6047716 (p= 1.45×10^{-11})
	rs141903557 (p= 9.78×10^{-11})
	rs11205303 (p= 1.62×10^{-10})
	rs12029076 (p= 2.17×10^{-10})
	rs1978060 (p= 3.26×10^{-10})
	rs2467146 (p= 5.96×10^{-10})
	rs11787412 (p= 1.32×10^{-9})
	rs188915802 (p= 1.94×10^{-9})
	rs658839 (p= 3.15×10^{-9})
	rs2194285 (p= 8.69×10^{-9})
	rs160335 (p= 9.10×10^{-9})
	rs482012 (p= 2.30×10^{-8})
	rs11341092 (p= 2.92×10^{-8})
	rs17011903 (p= 3.56×10^{-8})
	rs397948882 (p= 3.66×10^{-8})
	rs12149832 (p= 4.40×10^{-8})

1139

Rank	Gene Symbol	*P* value	Variants in Cases (N = 411)	Variants in Controls (N = 3,800)	OR (95%CI)
1	*EPHA4*	0.045533	3	17	4.09 (0.946-17.6)
2	*ARVCF*	0.067987	6	24	3.05 (0.807-11.6)
3	*TXNRD2*	0.092043	3	11	3.60 (0.729-17.8)
4	*TRIM17*	0.115033	3	11	3.53 (0.795-15.7)
5	*SLC25A1*	0.119166	3	9	3.39 (0.817-14.1)
6	*MICALL2*	0.199131	0	24	0.30 (0.0176-4.94)
7	*MTMR11*	0.249127	5	35	2.05 (0.663-6.36)
8	*MMP2*	0.252208	4	17	3.03 (0.613-15.0)
9	*BCKDHB*	0.293142	1	16	0.643 (0.0469-8.83)
10	*TTK*	0.293162	0	17	0.583 (0.0334-10.2)

1140   **Supplementary Table 2. Summarized results of burden analysis.**

11	*SLC2A6*	0.316088	1	21	0.675 (0.049-9.3)
12	*SOX6*	0.317369	2	25	0.822 (0.0949-7.11)
13	*WNT3A*	0.329116	1	11	0.814 (0.058-11.4)
14	*SDR42E1*	0.33538	0	11	0.631 (0.0359-11.1)
15	*GPR126*	0.33628	0	12	0.699 (0.0393-12.4)
16	*PIGN*	0.338169	1	22	0.663 (0.0869-5.06)
17	*POLL*	0.348494	0	11	0.668 (0.0378-11.8)
18	*PAX1*	0.350574	0	19	0.649 (0.0368-11.4)
19	*INSC*	0.363501	0	13	0.744 (0.0416-13.3)
20	*IBA57*	0.388644	0	8	0.744 (0.0416-13.3)

1141   The p-values (Fisher's Exact Test) of 20 top ranked gene were calculated on the
1142 basis of distribution differences of rare variant between the control group and the
1143 PUMCH IS cohort. Abbreviations: OR (odds ratio); CI (confidence interval).

1144   **Supplementary Table 4. Search strategies for each database.**

Search Number	Key Words	Results
MEDLINE (via Pubmed.gov)		
1#	Idiopathic scoliosis, GWAS/	69
2#	Idiopathic scoliosis, SNP/	56
3#	Idiopathic scoliosis, single nucleotide polymorphism/	149
4#	Idiopathic scoliosis, variant/	129
Web of Science (via Clarivate Analytics)		
1#	Idiopathic scoliosis, GWAS/	42
2#	Idiopathic scoliosis, SNP/	71
3#	Idiopathic scoliosis, single nucleotide polymorphism/	198
4#	Idiopathic scoliosis, variant/	227

1145 **Supplementary Table 5. Inclusion and exclusion criteria of literature review.**

	Inclusion Criteria	**Exclusion Criteria**
Patient Recruitment	Idiopathic scoliosis	Congenital scoliosis
	Adolescent idiopathic scoliosis	Neuromuscular scoliosis
	Early-onset idiopathic scoliosis	Syndromic scoliosis
		Secondary scoliosis
		Degenerative scoliosis
Study Type	Genome Wide Association Study (GWAS)	Linkage study
	Meta-analysis of GWAS	Candidate gene study
		Exome sequencing
		Whole genome sequencing
Study Design	Case-control	-

 Supplementary Table 6. Sequencing information of PUMCH IS cohort.

Characteristics	Cases		Controls	
	Trio	**Singleton**	**Trio**	**Singleton**
Exome sequencing	41	113	2,021	224
Genome sequencing	116	148	483	1,072
Sum	157	261	2,504	1,296
Passing QC	155	256	2,504	1,296

 Abbreviations: QC (quality control); IS (idiopathic scoliosis).

1148 **Supplementary Table 7. Sequences information of sgRNA and primers for**

1149 **zebrafish study.**

sgRNA sequences used to generate zebrafish *epha4a* and *epha4b* mutants	
epha4a sgRNA-1	AAGATTAAAGGTCTCCTTGC
epha4a sgRNA-2	GTATCGCTCTTTGTCATTGT
epha4a sgRNA-3	GAAGCTTTCATCAGCCGCAA
epha4a sgRNA-4	CCGTGTTCAGCTTCATGATG
epha4b sgRNA-1	GCCAAGTTCAACACCG-CCAG
epha4b sgRNA-2	AGTCACCGTATCGGGAAACT
Primers for mutant genotype	
epha4a-forward	GAGCTCAGCGGGTCTACATC
epha4a-reverse	GCGTACCTCCACTAGCGATG
epha4b-forward	GGTGTCTCTGAGGGTCTTTT
epha4b-reverse	CCTGACAGTAGTCCTTATTCTCCTC
Morpholinos used to knock down the expression of *epha4a* and *efnb3b*	
epha4a MO	AACACAAGCGCAGCCATTGGTGTC
efnb3b MO1	GTCTATTTACTCCCATCAAAGCCGT
efnb3b MO2	GAAATCCCGAATTCCGTCTATTTAC
Primers used for in situ hybridization analysis	
epha4a probe forward	AGAATACGCCAATCAGGACG

epha4a probe reverse	CCTTCATGTTCTTCATAGCC
epha4b probe forward	TTTGGTGGAGGTTAGAGGGT
epha4b probe reverse	GTTGTTTAGGGGTTGGGCTC
rfng probe forward	GGTGCTGCTTATTCCCTCCTT
rfng probe reverse	GCGCTAAACCTCGACTGATGC

Supplementary Movie 1: Video showing the swimming of three wild-type fish.

Supplementary Movie 2: Video showing the swimming of three *epha4a* mutants with severe or mild scoliosis.

Supplementary Movie 3: High-speed video showing the startle response in a wild-type larva at 5 dpf triggered by head tactile stimulation. Time units: ms.

Supplementary Movie 4: High-speed video showing the abnormal startle response of an *epha4a* mutant larva at 5 dpf triggered by head tactile stimulation. Time units: ms.

Supplementary Movie 5: Alternated activation of calcium signaling in motor neurons of 24 hpf wild type Tg(*elavl3*:GAL4; *UAS*: GCaMP) transgenic larva.

Supplementary Movie 6: Abnormal activation of calcium signaling in motor neurons of 24 hpf Tg(*elavl3*:GAL4; *UAS*: GCaMP) transgenic larva injected with *epha4a* MO.

Supplementary Movie 7: Tail oscillation after light activation of 5dpf Tg(Gal4^{s1020t}; *UAS*:ChR2) double transgene larva.

Supplementary Movie 8: Tail oscillation after light activation of 5dpf Tg(Gal4^{s1020t}; *UAS*:ChR2) double transgene larva injected with *epha4a* MO.