Decoding auditory working memory content from EEG aftereffects of auditory-cortical TMS

Işıl Uluç[1,2], Mohammad Daneshzand[1,2], Mainak Jas[1,2], Parker Kotlarz[1,2], Kaisu Lankinen[1,2], Jennifer L. Fiedler[1], Fahimeh Mamashli[1,2], Netri Pajankar[1], Tori Turpin[1], Lucia Navarro de Lara[1,2], Padmavathi Sundaram[1,2], Tommi Raij[1], Aapo Nummenmaa[1,2], Jyrki Ahveninen[1,2]

[1]Athinoula A. Martinos Center for Biomedical Imaging, Dept. of Radiology, Massachusetts General Hospital, Charlestown, MA, USA

[2]Harvard Medical School, Boston, MA, USA

Corresponding author:
Işıl Uluç, Ph.D.
CNY 149, 13th St.
A.A. Martinos Center for Biomedical Imaging,
Department of Radiology,
Massachusetts General Hospital,
Charlestown, MA 02129
iuluc@mgh.harvard.edu
Tel: 617 726 6584, Fax: (617) 726-7422

Abstract

Working memory (WM), short term maintenance of information for goal directed behavior, is

essential to human cognition. Identifying the neural mechanisms supporting WM is a focal point

of neuroscientific research. One prominent theory hypothesizes that WM content is carried in

"activity-silent" brain states involving short-term synaptic changes. Information carried in such

brain states could be decodable from content-specific changes in responses to unrelated "impulse

stimuli". Here, we used single-pulse transcranial magnetic stimulation (spTMS) as the impulse

stimulus and then decoded content maintained in WM from EEG using multivariate pattern

analysis (MVPA) with robust non-parametric permutation testing. The decoding accuracy of WM

content significantly enhanced after spTMS was delivered to the posterior superior temporal

cortex during WM maintenance. Our results show that WM maintenance involves brain states,

which are activity silent relative to other intrinsic processes visible in the EEG signal.

Keywords: Working memory, auditory working memory, transcranial magnetic stimulation, TMS-

EEG, MVPA

Introduction

Working memory (WM), the brain system that enables maintenance and processing of recent information, plays an essential role in daily living. The mechanisms and brain areas underlying WM maintenance have thus been prominent topics for neuroscience research. However, research into its neuronal mechanisms has resulted in seemingly contradictory results that have led to a long-standing controversy. The prevailing hypothesis suggests that information is maintained through persistent firing in the prefrontal cortex (PFC). Conversely, an alternative theory posits that persistent activity is not necessary for WM maintenance and rather maintenance can be dynamical in an 'activity-silent' format via functional connectivity and/or synaptic weights. [1-3]. Much of this research has been conducted in the visual modality only, leaving some of the most ecologically relevant aspects of WM in other sensory modalities relatively underexplored. One such aspect is auditory WM, which enables temporary storage and manipulation of sounds and verbal information, such as spoken language or music.

Initially, WM maintenance was linked to persistent activity of prefrontal neurons that respond to the incoming stimulus and remain activated even after the stimuli have vanished [4-6]. However, subsequent human neuroimaging studies suggested that the content of visual WM could only be decoded from signal-change patterns in sensory and posterior brain areas where persistent activity is not present during WM maintenance [7-9] (however, see also [10-13]). Studies in non-human primates (NHP) have not shown persistent WM related neuronal activity during the maintenance period in sensory areas [14, 15]. The identification of content-specific persistent firing patterns at the sensory level has proven challenging in NHP studies of auditory WM as well [16-18]. At the same time, human studies have managed to decode auditory WM content from fMRI signals [12, 19-21] as well as intracranial EEG signals [22] from auditory cortices. The diverse and contrasting findings have inspired the development of a family of alternative theories suggesting that WM is maintained in a more distributed and dynamic fashion than initially believed [23-25].

In explaining possible mechanisms for 'activity-silent' WM maintenance, a theory proposes that WM is maintained through intermittent bursts of neuronal firing and intervals of short-term

synaptic plasticity (STSP) [26], i.e., transient changes in the strength of synaptic connections

between neurons [27]. Item-specific activation of neurons during the encoding process leads to

presynaptic accumulation of calcium, which facilitates postsynaptic connectivity. Due to this

calcium buffer, which operates on a time scale of seconds, even sparse bursts of firing will be

sufficient to maintain the "activity-silent" WM representations [3, 25, 26]. The network

maintaining a synaptic WM trace will respond in a content-specific fashion, even if the non-

specific input is completely unrelated to the maintained representation [3]. Hence, information

maintained in WM via the activity-silent synaptic mechanisms should be decodable with machine

learning techniques that can classify responses elicited to any unrelated stimulus that broadly

activates the same neuronal population [3]. To test this prediction, recent human EEG and MEG

studies presented participants with "impulse stimuli", such as strong visual or auditory feature

patterns unrelated to the maintained content, during WM maintenance [13, 28-31]. The content

maintained in WM, which is otherwise in an activity-silent (or "hidden") state, became more

readily decodable from EEG or MEG responses to such impulse stimuli [13, 28-31]. A limitation in

many of these studies, however, is that it is difficult to deliver such impulse stimuli to a particular

brain area only.

A non-invasive way to probe hidden brain states that underlie human cognition is transcranial

magnetic stimulation (TMS). Unlike observational methods such as MEG or EEG, TMS allows us to

causally interact with focal areas whose role in WM we intend to evaluate [32]. In studies of

human memory processes, TMS has been used to modulate maintenance of visual WM

representations [33] and to enhance neuronal plasticity in visual cortex [34]. A particular benefit

of using single-pulse TMS as opposed to task-irrelevant sensory stimuli for probing memory-

related brain states that its effects are both temporally and anatomically specific [35]. In a recent

study that used TMS to enhance WM decoding from EEG signals [28], perturbing the WM circuits

during the seemingly activity-silent maintenance period yielded noteworthy results. This

intervention not only augmented the decoding of representations that were stored passively in

memory compared to actively maintained content but also contributed to participants recalling

passively maintained items more effectively from WM. However, to our knowledge, this approach

has so far not been tested in WM studies targeting auditory or other earlier sensory cortex areas,

or in designs applying active and sham TMS in the same participants.

Thanks to recent advances in MRI-guided TMS navigation systems and more focal stimulation

coils, TMS pulses can be delivered to the area of interest at an exact latency. This allows one to

test anatomically and temporally specific hypotheses to develop an understanding of how

sensory areas might be contributing to WM. Here, we investigated whether the content of

auditory WM, which is embedded in an activity silent brain state, can be decoded from cortical

effects of single-pulse TMS, delivered to posterior non-primary auditory cortex during the WM

maintenance period. This non-primary auditory cortex target was in in the left posterior superior

temporal cortex (pSTC). In our multivariate pattern analysis derived from whole-scalp EEG, the

decoding accuracy increases above chance level directly after the TMS pulse in an Active TMS

condition, but not after a Control TMS pulse that was too weak to activate the pSTC target area.

Therefore, our study provides strong evidence for the activity silent theory of WM maintenance.

Methods

Participants

A total of 23 healthy right-handed [36] participants (12 women, 11 men; mean age $\pm$ standard

deviation, SD, = 32 $\pm$ 12 years) were enrolled. One participant was excluded due to excessive

movement artifacts (facial movement artifacts in more than 50% of trials) and another due to

their chance-level behavioral performance, resulting in a final cohort of 21 participants (11

women, 10 men; mean age $\pm$ SD=31$\pm$10 years) for the Active TMS session. Significantly better

than chance level performance is essential in WM experiments to ensure that task performance

is not a result from a mere guessing. As for the Control TMS session, two participants opted not

to continue the study and one dataset was rejected due to excessive noise, resulting in a final

sample of 18 participants (9 women, 9 men; mean age $\pm$ SD = 31 $\pm$ 11 years). The same

participants attended Active TMS and Control TMS conditions to eliminate variability in EEG

responses from different participants in different conditions. All participants had normal or

corrected to normal vision and self-reported normal hearing. The participants provided written

informed consent and were informed that they could withdraw at any time. A monetary

compensation was given for each visit. The study design, protocol, and consent form were

approved by the Mass General Brigham Institutional Review Board.

Stimuli and experimental paradigm

Auditory stimuli

We employed non-conceptual, parametrically varied ripple sounds as WM items (**Fig. 1a**). Such

stimuli do not allow verbal memorization strategies. The ripple-velocity pool was individualized

based on each participant's pre-determined ripple-velocity discrimination thresholds. Just

noticeable difference (JND) values were calculated individually for each participant using a 2-

down, 1-up staircase algorithm [12, 13, 22]. Based on these values, we created four auditory

ripple sound stimuli, ensuring that each stimulus was positioned 1.5 JNDs apart from its closest

neighbor in velocity. We used four different ripple sounds as to-be-remembered stimuli. The same

four ripple stimuli were also presented as test stimuli. The order of the memory items was

pseudo-randomized. The participants were naïve to the number of memory items. The auditory

stimuli were presented at a comfortable listening level through Sensimetrics S14 Insert

headphones (Sensimetrics, Malden, MA) that provide high-quality acoustic stimulus delivery

while attenuating TMS click noise, analogous to our previous studies [37].

<u>Experimental paradigm</u>

Figure 1 shows the retro-cue experimental paradigm used during the TMS-EEG recordings. Each

trial started with a "!" presented on the screen. It was followed by two different consecutive ripple

sounds (memory items) that were presented for 750 ms with a 250 ms interstimulus interval.

Ripple sounds were followed by a visual retro-cue indicating whether the participant had to

remember the first sound ("1" on the screen) or the second sound ("2" on the screen). This was

followed by a 4-second maintenance period, where a TMS pulse was delivered in the middle. We

delivered the TMS pulse at the 2 s mark to have the same amount of signal before and after the

TMS pulse during maintenance for balanced comparison of results. Next, a test sound was

presented. The task was to determine whether the test sound was the same or different from the

memorized item. Participants responded with a mouse click with their right hand. An index finger

click indicated that the sounds were the same and a middle finger press indicated that they were

different. Finally, the screen showed whether the participant had responded correctly or

incorrectly. One run of the experiment consisted of 48 trials and one session consisted of 6 runs.

Thus, one session had 288 runs in total.

Figure 1. Task design. **(a)** Examples of the modulation patterns for ripple sounds **(b)** The target brain area for the TMS pulse, adapted from Uluç and colleagues [21]. **(c** The auditory WM retro-cue paradigm. The timeline of events in one trial is depicted.

Structural MRI Data Acquisition

T1-weighted anatomical images were acquired for with a multi-echo MPRAGE pulse sequence (TR=2510 ms; 4 echoes with TEs=1.64 ms, 3.5 ms, 5.36 ms, and 7.22 ms; 176 or 208 (to cover the ears) sagittal slices with 1×1×1 mm^3 voxels, 256×256 mm^2 matrix; flip angle = 7°) [38] in a 3T Siemens Trio MRI scanner (Siemens Medical Systems, Erlangen, Germany) using a 32-channel head coil.

TMS-EEG Data Acquisition

To be used as stimulus amplitude, resting motor threshold (rMT) of each participant was measured by sending a pulse to the left motor cortex thumb area and measuring the response from first dorsal interosseous muscle of the dominant right hand. From peak to peak, the smallest stimulation intensity resulting in 5/10 responses with amplitudes was equal to or greater than 50 uV. After the the rMT visit, participants completed two single-blind sessions. TMS pulses were

delivered either 1) at 100% of individual rMT to the posterior nonprimary auditory area pSTC in

the left hemisphere ("Active TMS") at 45° angle relative to the reference vector [0 0 -1] (A 0-

degree rotation relative to this reference vector means the coil handle is oriented from anterior

to posterior. The rotation angle increases counterclockwise around the superior-inferior axis) or

2) at 100% of rMT at the same location and same angle but with a 20 mm plastic block between

the coil and scalp ("Control TMS"). The pulses were delivered 2 s into the 4 s maintenance period.

EEG, horizontal EOG, and ECG data were sampled at 25 kHz with a 64-channel active EEG system

(ActiChamp, Brain Products GmbH, Gilching, Germany). TMS was delivered with a MagPro X100

w/ MagOption stimulator and a C-B60 figure-of-eight coil (MagVenture, Farum, Denmark). The

plastic block used in the control sessions was built in-house and was the same shape as the TMS

coil. The order of Active and Control TMS sessions was counterbalanced across the participants.

In both Active TMS and Control TMS conditions, the TMS coil clicks were masked with 8 kHz low

pass filtered white noise throughout the experiment. The white noise and the sound stimuli were

presented through Sensimetrics S14 Insert headphones (Sensimetrics, Malden, MA) with Comply

Canal In-Ear Tips (Hearing Components, Inc., North Oakdale, MN) that have a Noise Reduction

Rating (NRR) of above 29 dB. The sound level of the noise mask was measured using Larson Davis

sound level meter LXT2 with a Larson Davis RA0038 coupler (Larson Davis, New York, NY): The

level of the white noise was at 73 dB SPL and auditory items was at 86 dB SPL. Additionally,

subjective report from each participant was taken that they did not hear the TMS clicks or other

background noise.

TMS Neuronavigation

Continuous recording of the head position and orientation relative to the TMS coil was achieved

through a commercial TMS neuronavigation system (LOCALITE GmbH, Bonn, Germany) with an

optical camera and passive trackers (Polaris Spectra, Northern Digital Inc., Waterloo, Ontario).

The participant's registration to their anatomical data were all done in the Localite

neuronavigation software. The reconstructed MRI images were used in the Localite

neuronavigation system (LOCALITE GmbH, Germany) to guide the TMS procedure with MRI.

E-field Calculation

Data from one participant were discarded due to technical problems for Active TMS and Control

TMS sessions. To confirm that we had stimulated the intended cortical target, we computed the

TMS-induced Electric fields (E-field) using the Boundary Element Method accelerated by Fast

Multipole method (BEM-FMM) MATLAB toolbox implementation [39, 40]. The TMS coil

locations/orientations were extracted from the TMS navigation software. The participant-specific

anatomically realistic high-resolution head models were generated from the T1-weighted images

using the SimNIBS toolbox [41]. The model included five distinct layers of scalp, skull, cerebro-

spinal fluid (CSF), grey matter, and white matter, with the assumption of uniform conductivity

within each layer. The E-fields were calculated on a cortical surface halfway between the grey

and white matter surfaces. For group-level visualization, the individual E-field maps were

resampled to the FreeSurfer template brain fsaverage (version 6.2) and averaged across

participants [42].

Basic EEG Preprocessing and Analysis

EEG was preprocessed using MNE Python [43]. We used an established, rigorous preprocessing

procedure [44, 45]. The data were first detrended, and after selection and interpolation of noisy

channels (on average 4, channels), they were epoched to exclude any potential TMS pulse

artifacts. Two consecutive ICAs were calculated for the concatenated epochs [44, 45]. The first

ICA was used to remove remaining TMS related artifacts (3 independent components were

removed), and then the second ICA was performed to exclude physiological artifacts (on average,

5 independent components). Afterwards, we applied a 60-Hz notch filter to remove line noise

and a 150 Hz low pass filter. The data were then downsampled to 1 kHz. We did not use any high

pass filtering as it might introduce artifacts in the signal and tamper with later TMS-evoked

potential components [46]. Finally, all epochs were visually inspected for remaining artifacts and

noisy epochs were rejected. Signals from the occipital Iz electrode were excluded from all

analyses due to excessive noise in most of the datasets.

To display the time course of brain activity, we calculated event-related potentials (ERPs)

separately for each memory item during the memory period as well as a grand average for the

whole trial. For memory item comparison, the cleaned data were separated into four groups

according to cued memory item. Then, the separated data from all runs were concatenated and

averaged across all trials for each memory item within each participant. The data were then

averaged across all participants per memory item. For grand averages, the averages were

calculated across all trials irrespective of memory item. Topographical maps were calculated with

a time window of 0.5 s with equal weights for all trials.

Cross-participant Multivariate Pattern Analysis

Figure 2 shows the MVPA pipeline for the cross-participant classification analysis. We conducted

the decoding analysis by employing the support vector machine (SVM) implementation from

libsvm [47] as provided in the MATLAB/Octave CoSMoMVPA package [48]. To help generalize the

analysis results to a larger population, we performed the classification across the participants

[13]. For the MVPA analysis, to ensure that any TMS artifacts did not bias the results, the time

window starting 50 ms before and ending 50 ms after the TMS pulse was excluded from the

analysis. Using CoSMoMVPA and Fieldtrip toolboxes in each participant, we first balanced the

number of trials for each class and then calculated the class-specific averages for each participant.

Next, for spatial feature selection, we used principal component analysis (PCA) in MATLAB to

transform the data to "virtual channels", extracting the first eight principal components (PC) (**Fig.**

2). To keep the dimensionality constant to allow cross-validation, we selected the number of PCs

based on the grand average of cumulative variance explained across all participants, conditions,

and WM classes. The number of selected PCs refers to the point where the slope of the tangent

decreased below one in a normalized plot with both dimensions scaled between zero and one.

Calculating these spatial PCAs separately for each class in each participant ensured that no

leakage of information occurred between participants/classes. In each task condition, including

periods before and after the pulse in the TMS and Control conditions, this resulted in a two-

dimensional $\left(N_{Subjects} \times 4 \right) \times 8$ feature matrix that was entered into the SVM

(in the TMS conditions $N_{Subjects} = 21$, in the control conditions $N_{Subjects} = 18$).

The classification was conducted as a temporal searchlight analysis [48] with a 300 ms sliding

window at 3 ms steps, and was done separately for the periods before and after the TMS pulse.

The classification was performed using a leave-one out cross-validation procedure: the data sets

were partitioned to training and test sets such that the class-specific sub-averages of one

participant used as the test set and those from the rest of the participants as the training set. The

decoding accuracy was averaged across all iterations. For each condition (active TMS, Control

TMS) and maintenance period (before or after the TMS pulse), the analyses resulted in time series

with decoding accuracies of each searchlight centroid (**Fig. 2**).

Figure 2. MVPA pipeline. Preprocessed whole head EEG data was entered into a PCA for spatial feature

selection. The cut-off for the PC selection (n_{PC}=8) was determined based on the elbow in the grand-average

cumulative variance curve, calculated across all conditions, WM classes, and participants. Then the data

was entered into a searchlight analysis with cross-participant 4-class SVM classification. Leave-one-out

method was used for cross-validation. The analysis resulted with a decoding accuracy (DAc) time series

where DAc is the value assigned to the centroid of searchlight sliding window. For statistical significance,

we used maximum statistics with 500 permutations.

Statistical Significance, Cross-participant MVPA

In our cross-participant MVPA approach, the data of one participant were, iteratively, used as the

test set, to evaluate the model trained in the other participants. This analysis yields one decoding

accuracy value for the entire group at each time point. Instead of conventional one-sample t-

tests, we therefore determined the statistical significance of our cross-participant decoding

results using robust cluster-based permutation testing, which handles multiple comparison

problems using a *maximum-statistic* strategy [13]. Analogous temporal cluster-based maximum-

statistic approaches have been widely used procedures in univariate analyses of ERP and MEG

data [49]. In this procedure, we first generated 500 unique permutations of the true item-content

labels of the classifier. The temporal searchlight analysis was repeated with these permuted labels

to generate a distribution of decoding accuracies for each time point. For each permutation, the

time series of decoding accuracies were converted to *z*-values. This was done by comparing each

decoding-accuracy value to the respective permutation distribution at the same time point.

Continuous clusters with *z*>1.65 (i.e., *p*<0.05) were then identified in each permutation and the

respective cluster sums of *z*-values were calculated. From each permutation, the largest cluster

sum across all conditions was entered to the maximum-statistic null distribution. Analogously to

the conventional procedure [49], each cluster identified from the analysis with true content labels

was then compared to this null distribution, to determine their statistical significance. Clusters

with $p_{Corrected}$<0.05 were considered statistically significant.

Within-participant Multivariate Pattern Analysis

The within participant analysis is conducted based on the same principle as the cross-participant

decoding analysis (Figure 2), by employing the SVM implementation from libsvm [47] and

MATLAB/Octave CoSMoMVPA package [48] . The period 50 ms before and after the TMS pulse

was not entered into the analysis. Using CoSMoMVPA and Fieldtrip toolboxes in each participant,

we first balanced the number of trials for each class and low-pass filtered the signals at 75 Hz. To

enhance the SNR, subaverages of four trials were calculated with each class. Twenty different

random iterations were calculated of these subaveraged samples. In each iteration, spatial

feature selection was performed using a similar, yet individualized, PCA procedure to that used in

the cross-participant MVPA (range 5-10 PCs, group median = 8 PCs; see **Suppl. Fig. 1**).

Similar to the cross-participant MVPA, within-participant classification was performed using a

temporal searchlight analysis [48] with a 300 ms sliding window in 3 ms steps, conducted

separately for the periods before and after the TMS pulse. A k-fold cross-validation procedure was

used to classify the maintained WM content (k=6 in participants with six runs of data; k=5 in two

participants with five runs of data). In each fold, the model was trained in k-1/k of the samples

and tested in the remaining samples. For each searchlight dataset, the decoding accuracies were

averaged across the folds and iterations. For each participant, condition (active TMS, Control

TMS), and maintenance period (before or after the TMS pulse), the analyses resulted in a time

series with decoding accuracies of each searchlight centroid. Similar to previous studies [30], each

participant's decoding accuracy time courses were smoothed over time with a Gaussian kernel

with FWHM of 9.4 ms for significance testing.

Statistical Significance, Within-participant MVPA

The statistical significance was determined using robust cluster-based permutation testing, which

handles multiple comparison problems using a *maximum-statistic* strategy [13]. For each

participant and TMS condition, we first generated 500 unique permutations of the true item-

content labels of the classifier. The temporal searchlight analysis was repeated with these

permuted labels to generate a distribution of decoding accuracies for each time point. To assign

a p-value for each time point, the original group-mean decoding accuracy value, found from

classifiers with true labels, was compared with this permutation distribution. To improve the

precision, we modeled the empirical permutation distribution using a Gaussian fit. Continuous

clusters with $p<0.05$ were then identified in each permutation and the respective cluster sums

decoding accuracies were calculated. From each permutation, the largest cluster sum across all

conditions was entered to the maximum-statistic null distribution. Analogous to the conventional

procedure [49], each cluster identified from the analysis with true content labels was then

compared to this null distribution, to determine their statistical significance. Clusters with

$p_{Corrected}<0.05$ were considered statistically significant.

317 **Results**

318 *Behavioral performance*

319 The participants' behavioral percent corrects were 79.6 ± 6.5% (mean ± SD) in the Active TMS

320 sessions and 78.7± 6.2% in the Control TMS sessions. Although we see a slight increase in

321 behavioral performance in Active TMS session, the increase is not statistically significant. We

322 additionally calculated the percent correct of answers as a function of the ripple-velocity distance

323 from memory item to probe. This analysis revealed a consistent relationship between

324 participants' ability to reject non-matching probes and the difference in ripple velocity between

325 the probe and WM item. In the TMS session, the percent correct of answers for match trials was

326 84.5 ± 6.3%. For non-match trials with JND distance 1, the percent correct was 60.4 ± 13.9%; for

327 JND distance 2, it was 85.7 ± 8.6%, and for non-match trials with JND distance 3, it was 93.9 ±

328 4.9%. In the Control TMS session, match trial percent correct was 84.1 ± 5.7%. Non-match trial

329 JND distance 1 percent correct was 58.0 ± 14.9%; JND distance 2 percent correct was 2 87.1 ±

330 10.2%; and JND distance 3 percent correct 92.2 ± 9.0%.

331

332 *Multivariate Pattern Analysis*

333 *Cross-participant MVPA*

334 We conducted a four-class cross-participant classification analysis to determine whether the

335 single TMS pulses to left pSTC enhanced the decoding of memorized content from the ERPs. The

336 analysis employed a temporal searchlight approach, in which the decoding was performed based

337 on the spatiotemporal pattern of EEG activity within a 300-ms sliding window. The statistical

338 significance was verified through a robust cross-participant cross-validation and cluster-based

339 maximum-statistic permutation procedure. According to these analyses, in the Active TMS

340 condition, the MVPA decoding accuracy for memory content rose significantly above chance level

341 during the first few hundreds of milliseconds after the TMS pulse ($p_{Corrected}$ <0.05, cluster-based

342 maximum-statistic permutation test; cluster sum of normalized accuracy = 137.2; **Fig. 3d**). No

statistically significant decoding results were observed in any other time period in the active TMS

condition or in the Control condition (for an additional analysis of the stability of decoding, see

Suppl. Fig. 2).

Figure 3. The results of E-field calculations and searchlight MVPA decoding of WM content from EEG.

(a) Group median E-field maps for the Active TMS condition. **(b)** Group median E-field maps for the Control

TMS condition. **(c)** Null distribution for 500 permutations, utilized to determine the statistical significance

of decoding accuracies. From each permutation, the maximum cluster sum of normalized decoding

accuracy was identified and added to this null distribution. The vertical dotted line illustrates the critical

value for $p_{corrected} < 0.05$ (cluster sum(z) = 130). **(d)** Decoding accuracies in the cross-participant four-class

MVPA (% of correctly classified trials). The time series reflect the SVM decoding accuracies at the centroid

of each sliding 300-ms searchlight. These decoding accuracies were derived from an iterative leave-one-

participant out cross-validation procedure: In each searchlight time window, the data of each participant

was used once as the test set and those from the rest of the remaining participants as the training set. The

light red line denotes the Active TMS condition and the light blue line the Control TMS condition. The

dotted horizontal line indicates the chance level of decoding-accuracy in a four-class classification (25%).

The time window when the decoding accuracy was significantly higher than chance level in the Active TMS

condition is shown in dark red ($p_{Corrected}$<0.05, non-parametric cluster-based permutation test).

Within-participant MVPA

We additionally conducted a within-participant four-class SVM searchlight to test the effect of the

TMS pulse on the individual brain activity. We employed a similar searchlight approach with a

300-ms sliding window as in the cross-participant analysis. Consistent with the cross-participant

results, in the Active TMS condition, the group average of MVPA decoding accuracy for memory

content rose significantly above chance level during the first few hundreds of milliseconds after

the TMS pulse ($p_{Corrected}$ <0.01, cluster-based maximum-statistic permutation test; cluster sum of

decoding accuracy = 24.5; **Fig. 4**). No statistically significant decoding results were observed in

any other time period in active TMS or control TMS conditions.

Figure 4. The results of within-participant searchlight MVPA decoding of WM content from EEG for Control

and Active TMS conditions. **(a)** Decoding accuracies in the within-participant four-class MVPA. The time-

resolved decoding reflects the accuracies at the centroid of each sliding 300-ms searchlight. The thin red

line denotes the Active TMS condition and the thin blue line the Control TMS condition. The dotted

horizontal line indicates the chance level of decoding-accuracy in a four-class classification (25%). The time

window when the decoding accuracy was significantly higher than chance level in the Active TMS condition

is shown in dark red ($p_{Corrected}$<0.05, non-parametric cluster-based permutation test). **(b)** Null distribution

for 500 permutations, utilized to determine the statistical significance of decoding accuracies. From each

permutation, the maximum cluster sum of decoding accuracy was identified and added to this null

distribution. The vertical dotted line illustrates the critical value for $p_{corrected}$<0.05 (cluster sum of decoding

accuracy = 24.5).

E-field Calculations

Figures 3 a-b depict the median of the E-field calculations for Active TMS and Control TMS

conditions, respectively, thresholded at 60 V/m [50]. The average E-field in the target area

(Talairach -60, -38, 15) was 74.11 V/m for the Active TMS condition and 34.75 V/m for the Control

TMS condition.

Control Analyses

To control whether MVPA results are driven by differences in ERP amplitude between different

conditions, we averaged the response to different cued content during WM period across

participants. No systematic differences were found between signals for different cued content

during the maintenance period before or after the TMS pulse. We also performed a grand average

of the ERP data to observe the ERP time course during the task trial. **Figure 5** depicts the grand

average ERPs calculated for the WM maintenance period as the time of interest and TMS evoked

responses (TEP) for Active TMS and Control TMS sessions. As expected, the ERP results revealed

a N2/P3 response to the visual retro-cue at the beginning of the maintenance period. A pattern

of TMS-elicited ERP deflections was detected 2-2.5 s into the maintenance period. The auditory

component reflects a typical TEP elicited by the TMS pulse (**Fig. 5c**). No comparable effect is

observed in the Control TMS condition (**Fig. 5d**), although the auditory click sound was identical

to the Active condition. This suggests that the click sounds were sufficiently masked by the

constant noise masking used in all sessions. Finally, we did not observe any persistent elevation

in ERP during the WM maintenance period **(Fig. 5)**. To control the data quality, we calculated grand averages for the item presentation and probe and response period **(Suppl. Fig. 3-4)**. The averaged responses for these periods were as expected.

Figure 5. Topographical and butterfly plots of ERP time courses for time of interest (TOI) in the active TMS condition. Different colors in the ERP plots refer to different electrodes. The electrode map above the figures denotes the locations of electrodes. The timeline starts from $t = 0 s$ at the visual retro-cue. **(a)** TOI ERP topographical plots and time courses for WM before TMS pulse. The plot depicts the results for the visual retro-cue that starts the maintenance period. **(b)** TOI ERP topographical plots and time courses after TMS pulse. The timeline starts from $t = 0 s$ at the TMS pulse. **(c)** TMS evoked response for Active TMS session. **(d)** TMS evoked response for Control TMS session.

To control whether the decoding results were driven by a difference in TEP in different memory conditions due to TMS pulse, we also performed a repeated measures ANOVA across ERPs of different memory conditions. We did not find any significant differences in corrected or uncorrected level between averaged ERPs across different conditions (35-100ms: $F_{3,60}=0.65$, $p=0.59$; 100-200ms: $F_{3,60}=0.22$, $p=0.88$; 200-300ms: $F_{3,60}=0.36$, $p=0.78$). We also tested the ERPs across the memory period after the TMS pulse to test whether the auditory click artifact might

bias our main analysis results. We found no significant differences in the trial-averaged EEG

patterns following the TMS pulses across the memory conditions during the whole 2-s memory

period after the TMS pulse ($F_{3,60}=0.28$, $p=0.84$).

Finally, we also performed a control decoding analysis with the same parameters as our main

analysis using the task-irrelevant (i.e., "un-cued") items, which were to be forgotten after the

retro-cue. This control MVPA showed no significant decoding for neither Active TMS nor Control

TMS conditions at any time during the WM retention period.

Discussion

Here, we investigated auditory WM using a "perturbation approach", which combines MRI-navigated single pulse TMS with simultaneous EEG recordings, to unravel content-specific mnemonic states from EEG otherwise obscured by the much larger EEG "background" activity. To decode WM content from EEG signals during the maintenance period, we employed a temporal-searchlight MVPA with robust cross-participant cross-validation and non-parametric permutation testing. As predicted, the decoding accuracy of auditory WM content rose significantly above chance level after a single TMS pulse was delivered to the non-primary auditory areas in the left pSTC. Further, the Control TMS condition (otherwise the same as active single pulse TMS but with a 20-mm plastic block between the coil and the participant's scalp) found no significant decoding accuracy.

One possible explanation for the present finding is offered by the synaptic theory of WM [26]. According to this model, WM information is coded to content-specific patterns of functional connectivity, which result from activity-based STSP in the synaptic terminals of neurons that are strongly activated at the encoding stage [26]. Instead of persistent neuronal activity, this model predicts that only sparse bursts of neuronal oscillations and firing activity are necessary to maintain this otherwise activity-silent mnemonic brain state [3, 51-53]. Notably, although synaptic states are not directly measurable by non-invasive recordings, computational modeling predicts that a circuit that maintains information by content-specific changes of synaptic weights responds differently to other impulse stimuli (until the STSP decays) [26]. These content-specific responses to external impulses might not only provide a way for the maintained content to be read out at the recall stage, but they could also allow one to "ping" the maintained content with externally generated pulses such as single pulse TMS [13, 28-30].

Whereas the original synaptic theory refers to local circuits in prefrontal cortices [26], it has also been proposed that this synaptic model can also provide a way to explain how WM information might be represented in sensory areas , which presumably cannot support persistent neuronal firing in the absence of sensory stimulation [54-56]. However, previous studies probing activity-silent states of WM have been limited to TMS-based perturbation of association areas [28], or to

using auditory or visual "impulse stimuli" [13, 31] that might activate a wide array of brain

networks beyond sensory areas involved in orienting to task-irrelevant stimulus changes. TMS

offers a more direct way to focally perturb cortical neurons [33, 57] therefore increasing the

likelihood that the content-related signals originate from the targeted sensory areas rather than

from higher-order brain regions. The present results suggest that WM content can be decoded at

a high accuracy from EEG responses to TMS pulses directed to pSTC. They could thus offer new

insights into the role of activity-silent WM processes in the sensory cortices.

Alternative explanations for enhanced decodability of WM content, which follows an irrelevant

impulse stimulus (i.e., "pinging effects"), have been recently presented. In a recent reanalysis of

previous studies [28, 30], Barbosa et al. [58] attributed "pinging effects" of visual WM content to

reduced trial-to-trial variability of EEG signals, which was observed after the strong visual impulse

stimuli that had been used to facilitate the decoding of (presumably activity-silent) WM content

(see also [59]). Their reasoning was that such a reduction of variability across trials could have

enhanced the performance of algorithm because of enhanced SNR, rather than a genuine WM

reactivation effect. It is, however, important to note that there were notable differences between

the pinging effects of visual impulses vs. TMS-induced perturbations. In contrast to the effect of

strong visual impulse stimuli (reanalysis the data in [60]), TMS perturbations increased, rather

than decreased, the variability of signals from trial-to-trial (reanalysis of [28]). *These results led*

Barbosa et al. to conclude that TMS-induced enhancement of WM decoding from EEG data could,

nonetheless, reflect an activity-silent mechanism of WM. Notably, consistent with Barbosa et al.,

the present analyses provide no evidence of TMS-induced reduction of trial-to-trial variability of

EEG signals, which could have explained the enhancement of the decoding accuracy of auditory

WM content.

Using measures such as EEG to probe synaptic processes is supported by the notion that EEG

signals primarily result from post-synaptic processes in apical dendrites of cortical pyramidal

neurons [61-63]. These neurons constitute fundamental components of the canonical cortical

circuit that presumably supports WM [25, 64-66]. An inherent limitation of non-invasive

measures, however, is that they do not conclusively rule out other alternative explanations. In

addition to an activity-silent synaptic state, the present enhancement of WM decoding by TMS
pulses could also result from perturbation of an activity-based maintenance process. Cellular-
level studies demonstrate that single pulse TMS activates a broad population of cell bodies in the
cortex [57, 67]. The rapid firing of these neurons after the TMS pulse, which is followed by a
refractory period, disrupts the cortical network, resetting the stimulated region [57, 67]. The
present results could therefore also be arguably consistent with a subthreshold attractor model
[59, 68]. The subthreshold attractor model suggests that WM-related persistent activity tends to
be attracted to a bump state that emerges in varying locations across this network [69, 70]. By
interfering with such activation patterns, TMS pulses might result in content-specific signal
changes that are recordable by EEG. However, a challenge for such a model in the present context
is that the TMS pulse, which tends to overwrite the neuronal activity in the stimulated area, would
disrupt the content-specific population activity in the stimulated network. Neurophysiological
studies provide experimental evidence indicating that distractor events disrupt content-specific
firing activities rather than amplifying them to a discernible level in the mass action of neurons
[71]. Therefore, while it is not entirely incompatible, the subthreshold attractor model does not
adequately describe our results because we found no evidence of impaired WM performance in
the Active TMS vs. Control TMS condition.

Another alternative for a hidden state (whether it is due to synaptic plasticity or to other means)
of WM content in sensory areas is that the WM maintenance is carried through a recurrent neural
network where the PFC shapes and transforms the WM representations according to task
demands [54, 72]. The recurrent model is another possible explanation for how the synaptic
weights could be formed and maintained in the posterior and sensory brain areas. Indeed, it has
been recently shown that the WM content can be effectively maintained by a neuronal behavior
explained by a combination of activity-based and activity-silent models of WM [73].

Another important consideration is that, although the initial E-field exceeded the stimulation
threshold only in our targeted STG site, single-pulse TMS could influence not only local but also
distant neural circuits through axonal and transsynaptic propagation, potentially affecting other
cortical and subcortical areas starting already at the first tens of milliseconds after the TMS pulse.

The action potentials generated by the TMS-induced electric field may propagate along the axons in both anterograde and retrograde directions, facilitating forward and backward information flow within the stimulated pathway [74]. Computational modeling studies of TMS-EEG effects suggest that recurrent network feedback to the target regions begins driving TEP responses already around 100 ms post-stimulation, whereas only the earlier TEP components can be attributed to local reverberatory activity within the stimulated region [75]. Roughly consistent with previous impulse-stimulus and TMS studies of visual WM [28, 30], here the significant increases of decoding accuracy occurred about 200-300 ms after the TMS pulse, peaking slightly earlier in the within-participant than cross-participant analyses. The observed effects may result from feedforward propagation of activity from STG to other areas and/or subsequent feedback influences from other regions back to STG.

Our cross-participant decoding results indicate that the states that were revealed by the TMS pulse were stable across participants[76]. This is argued to be similar to the difference between fixed- and mixed-effects analyses [77]. In the context of MVPA, this distinction allows for the identification of differences in local computations. Significant prediction in a cross-participant model indicates that the stimulus-related information encoded by a time-resolved neuronal activity pattern stays relatively consistent across participants [78].

Some inherent limitations of EEG interpretations during a combined TMS-EEG study include several types of artifacts such as direct muscle/sensory nerve stimulation, somatosensory sensation related to the vibration of the coil, and acoustical clicks. There is a possibility that the improved decoding following a TMS pulse is attributable not only to its neurophysiological effects on the target area but also to non-specific effects associated with unrelated physiological events. Here, we attempted to control these biases with a Control TMS condition. Adding a hard plastic block provides similar tactile and auditory sensation as Active TMS but with subthreshold brain stimulation. To mitigate the effect of acoustical clicks we also used a continuous noise masker stimulus and TMS compatible insert earphones accompanied with earplugs that attenuate the background noise [37]. It is also worth noting that the same TMS stimulator output level was used for Control and Active TMS sessions for each participant, resulting also in exactly the same

sound level of the TMS click sound. Our continuous noise masking should have mainly eliminated

the possibility of TMS-evoked auditory effects (**Fig. 5 a-b**). Further, such effects should be similar

across the control and active TMS conditions as well as between the different memory conditions.

It is thus unlikely that the decoding results would be biased by any auditory artifacts. The

differences in WM decoding between Active and Control sessions thus cannot follow from the

click sound, per se. Finally, TMS was applied at a fixed 2-s latency, which might have created an

anticipation effect However, such an anticipation effect should have been identical in the Active

TMS and Control TMS conditions, making it unlikely that our main results were influenced by such

an effect.

To conclude, using TMS-EEG and cross-participant MVPA, the present study suggests

maintenance of WM content involves an "activity-silent" brain state in auditory brain areas. The

study also demonstrates the power of TMS as a way to probe information content embedded in

EEG signals.

Acknowledgments

This project was supported by NIH grants R01DC016915, R01DC016765, R01DC017991,

1R01DC020891, R01MH128421, R01MH130490-01A1, R01NS126337-03, R01NS126337,

S10OD028668 and P41EB030006.

Open practices statement

The deidentified data and code to reproduce the main findings will be made available on

https://dataverse.harvard.edu/dataverse/isiluluc/.

Declaration of Interest

Aapo Nummenmaa and Lucia Navarro de Lara are named inventors in patents and patent

applications related to TMS.

Tommi Raij and Mohammad Daneshzand are named inventors in patent applications related to

TMS.

References

[1]   D'Esposito M, Postle BR. The Cognitive Neuroscience of Working Memory. Annual Review
of Psychology 2015;66(1):115-42.
[2]   Constantinidis C, Klingberg T. The neuroscience of working memory capacity and training.
Nat Rev Neurosci 2016;17(7):438-49.
[3]   Stokes MG. 'Activity-silent' working memory in prefrontal cortex: a dynamic coding
framework. Trends Cogn Sci 2015;19(7):394-405.
[4]   Goldman-Rakic PS. Cellular basis of working memory. Neuron 1995;14(3):477-85.
[5]   Funahashi S, Bruce CJ, Goldman-Rakic PS. Mnemonic coding of visual space in the
monkey's dorsolateral prefrontal cortex. Journal of Neurophysiology 1989;61(2):331-49.
[6]   Constantinidis C, Williams GV, Goldman-Rakic PS. A role for inhibition in shaping the
temporal flow of information in prefrontal cortex. Nature Neuroscience 2002;5(2):175-80.
[7]   Harrison SA, Tong F. Decoding reveals the contents of visual working memory in early
visual areas. Nature 2009;458(7238):632-5.
[8]   Christophel TB, Haynes JD. Decoding complex flow-field patterns in visual working
memory. NeuroImage 2014;91:43-51.
[9]   Postle BR, Druzgal TJ, D'Esposito M. Seeking the Neural Substrates of Visual Working
Memory Storage. Cortex 2003;39(4):927-46.
[10]   Uluc I, Schmidt TT, Wu YH, Blankenburg F. Content-specific codes of parametric auditory
working memory in humans. Neuroimage 2018;183:254-62.
[11]   Wu YH, Uluc I, Schmidt TT, Tertel K, Kirilina E, Blankenburg F. Overlapping frontoparietal
networks for tactile and visual parametric working memory representations. Neuroimage
2018;166:325-34.
[12]   Ahveninen J, Uluc I, Raij T, Nummenmaa A, Mamashli F. Spectrotemporal content of
human auditory working memory represented in functional connectivity patterns. Commun Biol
2023;6(1):294.
[13]   Mamashli F, Khan S, Hamalainen M, Jas M, Raij T, Stufflebeam SM, et al. Synchronization
patterns reveal neuronal coding of working memory content. Cell Rep 2021;36(8):109566.
[14]   Sreenivasan KK, Curtis CE, D'Esposito M. Revisiting the role of persistent neural activity
during working memory. Trends in Cognitive Sciences 2014;18(2):82-9.
[15]   Lemus L, Hernandez A, Romo R. Neural codes for perceptual discrimination of acoustic
flutter in the primate auditory cortex. Proc Natl Acad Sci 2009;106(23):9471-6.
[16]   Scott BH, Mishkin M, Yin P. Neural correlates of auditory short-term memory in rostral
superior temporal cortex. Curr Biol 2014;24(23):2767-75.
[17]   Bigelow J, Poremba A. Achilles' ear? Inferior human short-term and recognition memory
in the auditory modality. PLoS One 2014;9(2):e89914.
[18]   Ng CW, Plakke B, Poremba A. Neural correlates of auditory recognition memory in the
primate dorsal temporal pole. J Neurophysiol 2014;111(3):455-69.
[19]   Kumar S, Joseph S, Gander PE, Barascud N, Halpern AR, Griffiths TD. A Brain System for
Auditory Working Memory. J Neurosci 2016;36(16):4492-505.

[20] Czoschke S, Fischer C, Bahador T, Bledowski C, Kaiser J. Decoding Concurrent
Representations of Pitch and Location in Auditory Working Memory. The Journal of Neuroscience
2021;41(21):4658-66.
[21] Uluç I, Schmidt TT, Wu YH, Blankenburg F. Content-specific codes of parametric auditory
working memory in humans. NeuroImage 2018;183.
[22] Uluç I, Peled N, Paulk AC, Bush A, Gumenyuk V, Kotlarz P, et al. Decoding auditory working
memory content from intracranial high frequency activity in humans. bioRxiv
2023:2023.08.04.552073.
[23] Christophel TB, Klink PC, Spitzer B, Roelfsema PR, Haynes JD. The Distributed Nature of
Working Memory. Trends Cogn Sci 2017;21(2):111-24.
[24] Rose NS. The Dynamic-Processing Model of Working Memory. Current Directions in
Psychological Science 2020;29(4):378-87.
[25] Miller EK, Lundqvist M, Bastos AM. Working Memory 2.0. Neuron 2018;100(2):463-75.
[26] Mongillo G, Barak O, Tsodyks M. Synaptic theory of working memory. Science
2008;319(5869):1543-6.
[27] Zucker RS, Regehr WG. Short-term synaptic plasticity. Annu Rev Physiol 2002;64:355-405.
[28] Rose NS, LaRocque JJ, Riggall AC, Gosseries O, Starrett MJ, Meyering EE, et al. Reactivation
of latent working memories with transcranial magnetic stimulation. Science
2016;354(6316):1136-9.
[29] Wolff MJ, Ding J, Myers NE, Stokes MG. Revealing hidden states in visual working memory
using electroencephalography. Front Syst Neurosci 2015;9:123.
[30] Wolff MJ, Jochim J, Akyurek EG, Stokes MG. Dynamic hidden states underlying working-
memory-guided behavior. Nat Neurosci 2017.
[31] Wolff MJ, Kandemir G, Stokes MG, Akyurek EG. Unimodal and Bimodal Access to Sensory
Working Memories by Auditory and Visual Impulses. J Neurosci 2020;40(3):671-81.
[32] Muri RM, Nyffeler T. Using transcranial magnetic stimulation to probe decision-making
and memory. Prog Brain Res 2008;171:413-8.
[33] Silvanto J, Cattaneo Z. Transcranial magnetic stimulation reveals the content of visual
short-term memory in the visual cortex. Neuroimage 2010;50(4):1683-9.
[34] Kozyrev V, Staadt R, Eysel UT, Jancke D. TMS-induced neuronal plasticity enables targeted
remodeling of visual cortical maps. Proc Natl Acad Sci U S A 2018;115(25):6476-81.
[35] Romero MC, Davare M, Armendariz M, Janssen P. Neural effects of transcranial magnetic
stimulation at the single-cell level. Nat Commun 2019;10(1):2642.
[36] Oldfield R. The assessment and analysis of handedness: The Edinburgh Inventory.
Neuropsychologia 1971;9:97-113.
[37] Ahveninen J, Huang S, Nummenmaa A, Belliveau JW, Hung AY, Jääskeläinen IP, et al.
Evidence for distinct human auditory cortex regions for sound location versus identity processing.
Nat Commun 2013;4:2585.
[38] van der Kouwe AJ, Benner T, Salat DH, Fischl B. Brain morphometry with multiecho
MPRAGE. Neuroimage 2008;40(2):559-69.
[39] Makarov SN, Noetscher GM, Raij T, Nummenmaa A. A Quasi-Static Boundary Element
Approach With Fast Multipole Acceleration for High-Resolution Bioelectromagnetic Models. IEEE
Trans Biomed Eng 2018;65(12):2675-83.

[40]   Makarov SN, Wartman WA, Daneshzand M, Fujimoto K, Raij T, Nummenmaa A. A software
toolkit for TMS electric-field modeling with boundary element fast multipole method: an efficient
MATLAB implementation. J Neural Eng 2020;17(4):046023.

[41]   Saturnino GB, Puonti O, Nielsen JD, Antonenko D, Madsen KH, Thielscher A. SimNIBS 2.1:
A Comprehensive Pipeline for Individualized Electric Field Modelling for Transcranial Brain
Stimulation. In: Makarov S, Horner M, Noetscher G, editors. Brain and Human Body Modeling:
Computational Human Modeling at EMBC 2018, Cham (CH); 2019, p. 3-25.

[42]   Fischl B. FreeSurfer. Neuroimage 2012;62(2):774-81.

[43]   Gramfort A, Luessi M, Larson E, Engemann DA, Strohmeier D, Brodbeck C, et al. MEG and
EEG data analysis with MNE-Python. Front Neurosci 2013;7:267.

[44]   Al E, Stephani T, Engelhardt M, Haegens S, Villringer A, Nikulin VV. Cardiac activity impacts
cortical motor excitability. PLoS Biol 2023;21(11):e3002393.

[45]   Rogasch NC, Sullivan C, Thomson RH, Rose NS, Bailey NW, Fitzgerald PB, et al. Analysing
concurrent transcranial magnetic stimulation and electroencephalographic data: A review and
introduction to the open-source TESA software. Neuroimage 2017;147:934-51.

[46]   Hernandez-Pavon JC, Veniero D, Bergmann TO, Belardinelli P, Bortoletto M, Casarotto S, et
al. TMS combined with EEG: Recommendations and open issues for data collection and analysis.
Brain Stimul 2023;16(2):567-93.

[47]   Chang C-C, Lin C-J. LIBSVM: A library for support vector machines. ACM Trans Intell Syst
Technol 2011;2(3):1-27.

[48]   Oosterhof NN, Connolly AC, Haxby JV. CoSMoMVPA: Multi-Modal Multivariate Pattern
Analysis of Neuroimaging Data in Matlab/GNU Octave. Front Neuroinform 2016;10:27.

[49]   Maris E, Oostenveld R. Nonparametric statistical testing of EEG- and MEG-data. J Neurosci
Methods 2007;164(1):177-90.

[50]   Numssen O, Kuhnke P, Weise K, Hartwigsen G. Electric-field-based dosing for TMS. Imaging
Neuroscience 2024;2:1-12.

[51]   Buschman TJ, Miller EK. Working Memory Is Complex and Dynamic, Like Your Thoughts. J
Cogn Neurosci 2022;35(1):17-23.

[52]   Lundqvist M, Rose J, Herman P, Brincat SL, Buschman TJ, Miller EK. Gamma and Beta Bursts
Underlie Working Memory. Neuron 2016;90(1):152-64.

[53]   Postle BR. The cognitive neuroscience of visual short-term memory. *Current Opinion in*
Behavioral Sciences. 1. 2015:40-6.

[54]   Mejias JF, Wang XJ. Mechanisms of distributed working memory in a large-scale network
of macaque neocortex. Elife 2022;11.

[55]   Wang XJ. Theory of the Multiregional Neocortex: Large-Scale Neural Dynamics and
Distributed Cognition. Annual Review of Neuroscience 2022;45(1):533-60.

[56]   Comeaux P, Clark K, Noudoost B. A recruitment through coherence theory of working
memory. Progress in Neurobiology 2023;228:102491.

[57]   Murphy SC, Palmer LM, Nyffeler T, Muri RM, Larkum ME. Transcranial magnetic
stimulation (TMS) inhibits cortical dendrites. Elife 2016;5.

[58]   Barbosa J, Lozano-Soldevilla D, Compte A. Pinging the brain with visual impulses reveals
electrically active, not activity-silent, working memories. PLoS Biol 2021;19(10):e3001436.

[59]   Christophel TB, Iamshchinina P, Yan C, Allefeld C, Haynes JD. Cortical specialization for
attended versus unattended working memory. Nat Neurosci 2018;21(4):494-6.

[60] Wolff MJ, Jochim J, Akyurek EG, Stokes MG. Dynamic hidden states underlying working-
memory-guided behavior. Nat Neurosci 2017;20(6):864-71.

[61] Hämäläinen M, Hari R, Ilmoniemi R, Knuutila J, Lounasmaa O. Magnetoencephalography
-theory, instrumentation, and applications to noninvasive studies of the working human brain.
Rev Mod Phys 1993;65:413-97.

[62] Buzsaki G, Anastassiou CA, Koch C. The origin of extracellular fields and currents--EEG,
ECoG, LFP and spikes. Nat Rev Neurosci 2012;13(6):407-20.

[63] Murakami S, Okada Y. Contributions of principal neocortical neurons to
magnetoencephalography and electroencephalography signals. J Physiol 2006;575(Pt 3):925-36.

[64] Kilpatrick ZP, Ermentrout B, Doiron B. Optimizing working memory with heterogeneity of
recurrent cortical excitation. J Neurosci 2013;33(48):18999-9011.

[65] Wang XJ. Synaptic basis of cortical persistent activity: the importance of NMDA receptors
to working memory. J Neurosci 1999;19(21):9587-603.

[66] Vogel P, Hahn J, Duvarci S, Sigurdsson T. Prefrontal pyramidal neurons are critical for all
phases of working memory. Cell Reports 2022;39(2):110659.

[67] Pashut T, Magidov D, Ben-Porat H, Wolfus S, Friedman A, Perel E, et al. Patch-clamp
recordings of rat neurons from acute brain slices of the somatosensory cortex during magnetic
stimulation. Front Cell Neurosci 2014;8:145.

[68] Jaffe RJ, Constantinidis C. Working Memory: From Neural Activity to the Sentient Mind.
Compr Physiol 2021;11(4):2547-87.

[69] Wimmer K, Nykamp DQ, Constantinidis C, Compte A. Bump attractor dynamics in
prefrontal cortex explains behavioral precision in spatial working memory. Nat Neurosci
2014;17(3):431-9.

[70] Itskov V, Hansel D, Tsodyks M. Short-Term Facilitation may Stabilize Parametric Working
Memory Trace. Front Comput Neurosci 2011;5:40.

[71] Miller EK, Li L, Desimone R. Activity of neurons in anterior inferior temporal cortex during
a short-term memory task. J Neurosci 1993;13(4):1460-78.

[72] Panichello MF, Buschman TJ. Shared mechanisms underlie the control of working memory
and attention. Nature 2021;592(7855):601-5.

[73] Barbosa J, Stein H, Martinez RL, Galan-Gadea A, Li S, Dalmau J, et al. Interplay between
persistent activity and activity-silent dynamics in the prefrontal cortex underlies serial biases in
working memory. Nat Neurosci 2020;23(8):1016-24.

[74] Siebner HR, Funke K, Aberra AS, Antal A, Bestmann S, Chen R, et al. Transcranial magnetic
stimulation of the brain: What is stimulated? - A consensus and critical position paper. Clin
Neurophysiol 2022;140:59-97.

[75] Momi D, Wang Z, Griffiths JD. TMS-evoked responses are driven by recurrent large-scale
network dynamics. eLife 2023;12:e83232.

[76] Clithero JA, Smith DV, Carter RM, Huettel SA. Within- and cross-participant classifiers
reveal different neural coding of information. Neuroimage 2011;56(2):699-708.

[77] Friston KJ, Stephan KE, Lund TE, Morcom A, Kiebel S. Mixed-effects and fMRI studies.
Neuroimage 2005;24(1):244-52.

[78] Kay KN, Naselaris T, Prenger RJ, Gallant JL. Identifying natural images from human brain
activity. Nature 2008;452(7185):352-5.