Long-read sequencing and genome assembly of natural history collection samples and challenging specimens

Bernhard Bein [1,2,3#], Ioannis Chrysostomakis [4#], Larissa S. Arantes [5,6#], Tom Brown [5,6#], Charlotte Gerheim [1,2], Tilman Schell [1,2], Clement Schneider [7], Evgeny Leushkin [1,2], Zeyuan Chen [2], Julia Sigwart [1,2], Vanessa Gonzalez [8], Nur Leena W.S. Wong [9], Fabricio R. Santos [10], Mozes P. K. Blom [11], Frieder Mayer [11], Camila J. Mazzoni [5,6], Astrid Böhne [4], Sylke Winkler [12,13], Carola Greve [1,2], Michael Hiller [1,2,3*]

[1] LOEWE Centre for Translational Biodiversity Genomics, Senckenberganlage 25, 60325 Frankfurt, Germany
[2] Senckenberg Research Institute, Senckenberganlage 25, 60325 Frankfurt, Germany
[3] Institute of Cell Biology and Neuroscience, Faculty of Biosciences, Goethe University Frankfurt, Max-von-Laue-Str. 9, 60438 Frankfurt, Germany
[4] Center for Molecular Biodiversity Research, Leibniz Institute for the Analysis of Biodiversity Change, Museum Koenig Bonn, Adenauerallee 127, 53113 Bonn, Germany
[5] Berlin Center for Genomics in Biodiversity Research (BeGenDiv), Königin-Luise-Straße 2-4, 14195 Berlin, Germany
[6] Department of Evolutionary Genetics, Leibniz Institute for Zoo and Wildlife Research, Alfred-Kowalke-Straße 17, 10315 Berlin, Germany
[7] Senckenberg Research Institute, Am Museum 1, 02826 Görlitz, Germany
[8] Global Genome Initiative, National Museum of Natural History, Smithsonian Institution, Washington, DC 20013, USA
[9] International Institute of Aquaculture and Aquatic Sciences, Universiti Putra Malaysia, 71050 Negeri Sembilan, Malaysia
[10] Laboratório de Biodiversidade e Evolução Molecular, Departamento de Genética, Ecologia e Evolução, Universidade Federal de Minas Gerais, Belo Horizonte, Minas Gerais, Brazil
[11] Museum für Naturkunde, Leibniz Institute for Evolution and Biodiversity Science, Invalidenstraße 43, 10115, Berlin, Germany
[12] Max Planck Institute of Molecular Cell Biology and Genetics, Pfotenhauerstr. 108, 01307 Dresden, Germany
[13] DRESDEN concept Genome Center, Technische Universität Dresden, 01062 Dresden, Germany

shared first authors
* michael.hiller@senckenberg.de

Abstract

Museum collections harbor millions of samples, largely unutilized for long-read sequencing. Here, we use ethanol-preserved samples containing kilobase-sized DNA to show that amplification-free protocols can yield contiguous genome assemblies. Additionally, using a modified amplification-based protocol, employing an alternative polymerase to overcome PCR bias, we assembled the 3.1 Gb maned sloth genome, surpassing the previous 500 Mb protocol size limit. Our protocol also improves assemblies of other difficult-to-sequence molluscs and arthropods, including millimeter-sized organisms. By highlighting collections as valuable sample resources and facilitating genome assembly of tiny and challenging organisms, our study advances efforts to obtain reference genomes of all eukaryotes.

Keywords

long-read sequencing, PCR amplification, genome assembly, museum collections

Background

High-quality genomes provide a powerful basis for understanding phylogenetic relationships, discovering fundamental principles of evolutionary processes, applying genomic methods to characterize, monitor and preserve biodiversity, and ultimately revealing the genetic blueprint underlying the fascinating diversity of life on our planet. Therefore, generating high-quality genomes of eukaryotic species has become a central goal in biological sciences [1]. Advances in short-read sequencing technology (with Illumina as the most prominent platform) enabled sequencing the genomes of a few thousand eukaryotes to date [2–5]. However, because eukaryotic genomes are often large and rich in repetitive DNA sequences, genome assembly from short reads ranging from 100 to 300 bp in size results in fragmented and incomplete assemblies [2,3,5], posing many limitations to downstream analyses. To generate highly contiguous genomes, the field has shifted to adopting long-read sequencing platforms from PacBio or Oxford Nanopore Technologies that can sequence DNA fragments with sizes of many kilobase pairs (kb) at once. Such long reads span most genomic repeats and do not suffer from the sequencing biases of short-read platforms in regions with very high or low GC content. Thus, long reads result in highly contiguous and complete genome assemblies, culminating in telomere-to-telomere assemblies [6,7], and consequently enable complete genome annotations and comprehensive analyses [8–16].

A key limitation of long-read sequencing is the availability of high molecular weight DNA, ideally with fragment sizes of 50 kb or more. To obtain samples delivering such DNA, the best practice is to acquire fresh samples (which may require sacrificing an individual), flash-freeze in liquid nitrogen, and preserve samples permanently at -80°C until DNA is extracted (https://www.earthbiogenome.org/sample-collection-processing-standards). Such protocols are not practical or not possible for (i) rare or endangered species, where sacrificing even a single living individual is not permitted, (ii) species which are difficult to sample in the field (e.g. cetaceans), or (iii) situations where liquid nitrogen and freezer capacity is not practicable

83　(e.g. in remote areas). Therefore, sample availability is a key challenge for biodiversity
84　genomics [17].
85
An alternative to get access to valuable or rare species that comprise Earth's biodiversity are
samples that are available in museums and other research collections that house millions of
specimens worldwide, including samples from extinct species [18]. As one example
demonstrating the value of such collections for biodiversity genomics, several hundred bird
genomes have been generated from dry samples stored in museum collections [3]. However,
since DNA of dry samples often exhibits various degrees of degradation, short-read
sequencing was the only feasible technology, resulting in fragmented bird assemblies with
an average contiguity of 43 kb (measured as contig N50 values, which state that 50% of the
assembly consists of contiguous DNA segments – called contigs – of at least that size).
Nevertheless, this and other studies using dry museum samples and short-read sequencing
approaches, including marker-based sequencing and genome skimming, provided valuable
insights into taxonomy, phylogenomics and conservation genomics [19–21].
98
In addition to dry material, collections worldwide also contain many millions of samples
preserved in ethanol. In comparison to the logistical challenges associated with bringing liquid
nitrogen to field trips and transporting flash-frozen samples without breaking the cold chain,
preserving and transporting collected samples in ethanol is a notably simpler task. Since
kilobase-sized DNA can be preserved in such samples [22,23], we explored whether ethanol-
preserved samples are also suitable for long-read sequencing. We reasoned that even if DNA
fragment sizes are substantially shorter than 50 kb, successfully sequencing reads of a few
kilobases in size increases read length by at least an order of magnitude compared to short-
read sequencing approaches, which in turn will improve assembly contiguity. In particular, we
focused on the PacBio high-fidelity (HiFi) read protocol that instead of generating error-prone
reads from "as long as possible" DNA fragments, sequences medium-sized fragments (10-15
kb) but with a high base accuracy of 99.8% [24]. HiFi sequencing enables assemblies that are
both more contiguous and have a higher base accuracy than assemblies obtained with longer
but more error-prone reads [7,16,24,25], making it a promising technology to apply to ethanol-
preserved samples.
114
In this study, we explored the utility of ethanol-preserved samples from collections for HiFi
sequencing. Although we encountered DNA degradation and sample contamination as
expected problems in some samples, we also successfully demonstrate that HiFi reads can
be obtained from ethanol-preserved samples containing kilobase-sized DNA, either using
amplification-free protocols or by using a modified amplification-based protocol that effectively
addresses issues associated with HiFi sequencing and PCR bias. Using this modified protocol,
we generate a high-quality assembly of the 3.1 Gb genome of the maned sloth *Bradypus*
torquatus, demonstrating that the previous genome size limit of 500 Mb can be substantially
extended. Beyond collection samples, we further show that our modified protocol improves
the contiguity of assemblies of species belonging to other phyla such as Mollusca
(Gastropoda, Bivalvia) and Arthropoda (Collembola), where amplification is often required for
long-read sequencing. The efficacy of this protocol facilitates genome assembly of challenging
taxa and suggests that collections can serve as valuable sample sources for long-read
sequencing.

Results

HiFi sequencing of ethanol-preserved samples with an amplification-free protocol

To investigate the effectiveness of PacBio HiFi sequencing from ethanol-preserved collection samples, we focused on vertebrates and used samples of four mammals (three-toed jerboa *Dipus sagitta*, pen-tailed treeshrew *Ptilocercus lowii*, long-eared flying mouse *Idiurus macrotis*, maned sloth *Bradypus torquatus*), two squamates (European blind snake *Xerotyphlops vermicularis*, slow worm *Anguis fragilis)* and two fishes (the catfish species *Cathorops nuchalis* and *Cathorops wayuu*), all lacking a genome assembly (Table 1, Supplementary Table 1). All samples were collected in the field and preserved in technical or 96% ethanol. Apart from the maned sloth and the catfishes, all samples were kept at room temperature. The samples of the maned sloth and catfish were kept most of the time in a freezer at -20ºC; however, in contrast to flash-frozen samples, freezing did not occur immediately after sampling and they were kept at room temperature for extended periods of time, including during transportation.

We used a modified Circulomics Nanobind disk and a phenol/chloroform based protocol for the extraction of genomic DNA (Methods). For *Dipus sagitta*, *Ptilocercus lowii*, and *Xerotyphlops vermicularis*, we did not obtain a sufficient amount of DNA (< 400 ng) and/or DNA fragments were shorter than 0.18 kb (Supplementary Table 1), showing that DNA is too degraded to proceed with library preparation. For four species (*Anguis fragilis*, *Idiurus macrotis*, *Cathorops nuchalis*, and *Cathorops wayuu*), the amount of DNA and the DNA fragment sizes were sufficient to prepare an amplification-free PacBio low input library [26] (Supplementary Table 1). We sequenced all libraries on a PacBio Sequel IIe system, disabling on-board calling of HiFi reads and instead applying the computationally expensive DeepConsensus method [27] to maximize HiFi read yield and length. For *Bradypus torquatus*, we did not obtain enough DNA and therefore proceeded with a PacBio ultra-low input library (see below).

For the two catfish species, *Cathorops nuchalis* and *Cathorops wayuu*, we sequenced two SMRT cells each and obtained HiFi reads with an average length of 8,832 and 8,783 bp, respectively, providing a total of 43.8 and 41.2 Gb, which corresponds to a coverages of ~17X and ~16.5X (Supplementary Table 1). Using HiFiasm with different parameters [28], we were able to obtain a contig assembly for both species with a total length of 2.6 and 2.59 Gb and a contig N50 value of 3.2 and 2.1 Mb (Supplementary Table 2). To assess gene completeness, we used compleasm [29] with the set of 3,640 ray-finned fish (Actinopterygii) near-universally conserved genes (ODB10) [30], which showed that 96.65% of these genes are fully present in the primary assembly of *C. nuchalis* and 95.6% in that of *C. wayuu*. Although additional HiFi data would be needed to improve contiguity and HiC data would be required to scaffold the contigs into chromosome-level scaffolds, our catfish samples exemplify that an adequate genome assembly can be obtained from 10-year-old, ethanol-preserved tissues.

In contrast to the catfish, we obtained very low sequencing yields for *Idiurus macrotis* and *Anguis fragilis*, with only 0.3 Gb and 0.04 Gb of HiFi data (Supplementary Table 1). Quality metrics showed that the polymerase N50 raw read lengths were very short and the local base

rates were low. For example, while the library from *Anguis* met the requirements for PacBio
sequencing with a mean fragment length of 12.2 kb, both the local base rate of 1.64 (expected
~2.8) and the polymerase N50 raw read length of 32.3 kb (expected at least 200 kb) are very
low and insufficient to produce HiFi reads of most DNA fragments in the library. This indicates
that factors such as DNA damage, metabolites bound to the DNA, or contaminants
precipitated with the DNA inhibit the polymerase, highlighting sequencing challenges for
ethanol-preserved samples stored at room temperature.
181

species	year sampled	preservation	type of sample
northern three-toed jerboa (*Dipus sagitta*)	2006 & 1961	technical ethanol, room temperature	muscle, skin
pen-tailed treeshrew (*Ptilocercus lowii*)	1967		muscle
long-eared flying mouse (*Idiurus macrotis*)	2000		skin with hair
maned sloth (*Bradypus torquatus*)	2003	likely pure ethanol, mostly at -20°C (otherwise room temperature)	clogged blood
European blind snake (*Xerotyphlops vermicularis*)	2004 & 2011	technical ethanol, room temperature	skin and muscle
slow worm (*Anguis fragilis*)	2021		muscle from tail cross-section
catfish (*Cathorops nuchalis*)	2014	pure ethanol, transported multiple times at room temperature until final storage at -20°C	fin
catfish (*Cathorops wayuu*)	2014		fin

182 Table 1: Overview of the species and samples.
183
184

HiFi sequencing with the amplification-based ultra-low input protocol

We reasoned that a PCR-based amplification step prior to library preparation could render the
Idiurus macrotis and *Anguis fragilis* samples amenable to sequencing, as this procedure
should yield intact DNA devoid of potential polymerase-inhibiting metabolites. To this end, we
applied the PacBio ultra-low input library protocol [31] to the samples of *Idiurus macrotis* and
Anguis fragilis. Although this protocol was originally designed for small specimens providing
very limited DNA amounts [32] and is recommended only for genome sizes of up to 500 Mb,
the protocol includes a PCR amplification step using two different undisclosed polymerases
targeting DNA with average and high GC contents, respectively. For simplicity, we refer to
these polymerases as "A" and "B" in the following to distinguish them from a third polymerase
"C" that we also investigate as described below. We also generated an ultra-low input library
for the *Bradypus torquatus* sample that did not contain enough DNA for the low input protocol.
197

198   Indeed, for *Idiurus macrotis* and *Anguis fragilis*, sequencing another SMRT cell each produced
199   10 and 19.6 Gb in HiFi reads with an average HiFi read length of 4,854 bp and 7,552 bp. The
200   first SMRT cell for *Bradypus torquatus* yielded 29.9 Gb in HiFi reads with an average HiFi read
201   length of 10,850 bp (Supplementary Table 1).
202
For *Idiurus macrotis* and *Anguis fragilis*, we investigated whether a DNA repair step applied to
the DNA extract before preparing the ultra-low library would increase HiFi read length and
yield (Methods). In contrast to the previous sequencing results, adding the DNA repair step
produced shorter HiFi reads (average read length 4,270 vs. 4,854 bp for *Idiurus macrotis* and
5,609 vs. 7,552 bp for *Anguis fragilis*) and a lower yield (6.4 vs. 10 Gb for *Idiurus macrotis* and
12.6 vs. 19.6 Gb for *Anguis fragilis*), suggesting that the DNA repair process is not
advantageous for these samples (Supplementary Table 1).
210
Next, we investigated whether the sequenced DNA was contaminated with bacteria, fungi or
other microorganisms. While little contamination was found in the *Bradypus torquatus* sample
(~200 kb mostly assigned to plants), the *Anguis fragilis* data had higher levels of contamination
(~200 Mb assigned to various bacterial groups), and the vast majority of the sequencing data
obtained from the *Idiurus macrotis* sample were contamination (~75 Mb assigned to various
groups of bacteria) (Supplementary Figures 1, 2). High levels of contamination (71-90% of
sequenced reads) were also detected for three additional ethanol-preserved samples, where
we directly applied the ultra-low input protocol: Russian desman (*Desmana moschata*)
sampled in 1947, Hazel dormouse (*Muscardinus avellanarius*) sampled in 2016, and a *Anguis*
fragilis sample from 1878 (Supplementary Tables 1, 3). Together, while sample contamination
with bacteria, protists and bacterial viruses or cross-contamination with human DNA is another
challenge related to samples obtained from collections [33–35], our tests also show that
amplifying DNA in the ultra-low input protocol prior to library preparation can enable PacBio
HiFi sequencing of samples where the amplification-free low input library protocol failed.
225
226

PCR bias in the current protocol prevents high-quality assemblies of larger genomes

To investigate the feasibility of using the ultra-low input protocol to obtain a high-quality
assembly of a genome that substantially exceeds the recommended size limit of 500 Mb, we
focused on the maned sloth that has an estimated genome size exceeding 3 Gb and showed
a low level of contamination. To obtain sufficient read coverage for genome assembly, we
generated two additional libraries using the PacBio ultra-low protocol and sequenced four
additional SMRT cells. In total, all five SMRT cells provided 140.2 Gb of HiFi reads, a total
coverage of ~45X, with an average read length of 10.6 kb. However, using this data, we only
obtained an assembly with a contig N50 of 405 kb (Figure 1, brown dashed line), which is
unexpectedly low as similar read coverages typically yield mammalian assemblies with contig
N50 values exceeding several megabases. Using compleasm [29] with the set of 11,366 near-
universally conserved eutheria genes (ODB10) showed that only 85.3% of these genes are
fully present in our assembly. Similarly, using TOGA [36] to determine how many of the 18,430
ancestral placental mammal coding genes have an intact reading frame, revealed that only

68% of the ancestral genes are intact. Together, this indicates not only a low assembly
contiguity but also a high level of incompleteness.
244

245
Figure 1: Contiguity of *B. torquatus* assemblies generated with data from ultra-low input libraries
prepared with polymerases A/B and/or C at different coverages.
Assembly contiguity visualized as N(x) graphs that show contig sizes on the Y-axis, for which x percent
of the assembly consists of contigs of at least that size. The N50 and N90 values are shown as vertical
grey lines and indicate contig sizes for which 50% and 90% of the assembly consists of contigs of at
least that size, respectively. Assemblies involving polymerase C read data are shown as solid lines,
assemblies generated from polymerase A/B data are shown as dashed lines. Colors refer to different
comparisons discussed in the text and summarized in the inset.
254
255
To further investigate the reasons for the poor quality of this assembly, we aligned our
Bradypus torquatus HiFi reads against the high-quality genome of a related sloth species,
Choloepus didactylus [11]. Despite both species being separated for 30 My [37], we observed
that 84.3% of the *Choloepus didactylus* genome was covered with *B. torquatus* HiFi reads at
an average coverage of 38X. Inspecting the mapped reads in a genome browser revealed
larger genomic regions, often spanning many kilobases, that completely lack any mapped
reads (Figure 2). Since several of these regions contain highly-conserved genes, we reasoned
that these read dropouts are probably not caused by high divergence between the sloth
species. Instead, it is likely that despite relying on two polymerases, the PacBio ultra-low
protocol has PCR bias on larger genomes, resulting in genomic regions that lack any reads.

266
Figure 2: PCR bias in reads produced with polymerase A/B.
UCSC genome browser screenshots of the *Choloepus didactylus* assembly, together with the TOGA
gene annotation and mapped HiFi reads of *B. torquatus* produced either with polymerase A/B or
polymerase C. The TOGA gene annotation is shown in blue with boxes representing coding exons,
connecting horizontal lines representing introns, and arrowheads indicating the direction of transcription
(+ or - strand). Mapped HiFi reads are shown below as boxes with orange tickmarks representing

insertions in the *B. torquatus* reads relative to the *C. didactylus* assembly. Reads in blue and red align
to the + and - strand, respectively.
(A) In the *HOXA* gene cluster, several regions, often covering parts or entire *HOX* genes, lack any reads
produced with polymerase A/B (highlighted in blue). In contrast, these regions have a coverage of HiFi
reads produced with polymerase C, which is sufficient for assembly.
(B) While reads produced with polymerase A/B do not cover the *CHRM4* and *MDK* genes, polymerase
C reads cover the entire locus.
280
281

A different polymerase alleviates PCR bias and enables highly-complete assemblies of larger genomes

To alleviate PCR bias, we adapted the ultra-low input protocol and used a different
polymerase, KOD Xtreme™ Hot Start DNA Polymerase (Merck). According to the specification
sheet, this polymerase amplifies DNA fragments up to 24 kb at high fidelity, including
templates with up to 90% GC content, which could help to overcome the underrepresentation
of very low or high GC regions of the PacBio ultra-low input protocol. For simplicity, we refer
to this polymerase as "C" in the following. Using a single library, we sequenced another three
SMRT cells for the maned sloth, providing 91.7 Gb (corresponding to an additional 27X
coverage) of reads with an average length of 10.2 kb.
292
Performing genome assembly using all HiFi reads obtained with polymerase A/B and C,
produced a 3.13 Gb assembly with a contig N50 of 4.88 Mb (Figure 1, black line,
Supplementary Table 4), which is 12 times higher than the previous assembly generated from
reads obtained with polymerase A/B. Gene completeness estimated with compleasm
improved from 85.3% to 96.4% and the percentage of intact ancestral placental mammal
genes inferred with TOGA increased from 68 to 88.6%. Furthermore, mapping the polymerase
C HiFi reads to the *C. didactylus* assembly covered the regions that completely lacked any
read before (Figure 2). Consistent with a higher PCR bias for polymerase A/B, we found that
the normalized read coverage in exonic and repeat regions is biased towards a lower coverage
for the polymerase A/B data compared to polymerase C to data (Supplementary Figure 3).
This confirms that previous read dropouts were not caused by sequence divergence between
both sloth species or selective degradation of certain genomic regions in our sample, but by
PCR bias associated with the polymerases in the PacBio ultra-low input protocol.
306
To directly compare the effect of polymerase A/B vs. C, taking differences in read coverage
from the individual SMRT cells out of the equation, we downsampled our data to equal
coverage and performed a number of tests (Supplementary Table 4). Since DNA fragments
generated by polymerase A and B are pooled during the library preparation, we cannot
investigate the effect of those two polymerases individually. Using an equal, downsampled
coverage of ~11X, we found that the assembly produced from only polymerase C reads
outperformed the assembly produced from only polymerase A/B reads by exhibiting a
substantially higher contiguity (contig N50 1.22 Mb vs. 264 kb) and gene completeness (89.3%
vs. 77.0% completely detected genes) (Figure 1, light blue lines). Remarkably, an assembly
obtained from the complete polymerase C read data is substantially better than an assembly
obtained from the complete polymerase A/B read data (contig N50 3.5 Mb vs. 405 kb, 96.4%

vs. 85.3% completely detected genes) (Figure 1, brown lines), despite the polymerase C data
having a substantially lower coverage (27X vs. 45X for polymerase A/B).
320
We next investigated how the number of libraries produced with polymerase A/B influences
assembly, as additional libraries may increase complexity and reduce bias. However,
sampling an equal coverage of ~20X from either one, two or three libraries results in very
similar assemblies in terms of contiguity and gene completeness (Figure 1, red/orange/yellow
lines; Supplementary Table 4), indicating that inherent bias of polymerase A/B hampers
assembly quality that cannot be overcome by producing several libraries.
327
328   Together, these tests show that *B. torquatus* assemblies generated with polymerase C reads
329   are substantially better. To our knowledge, we provide the first high-quality contig assembly
330   of a 3.1 GB genome that was produced using an adapted ultra-low input protocol combining
331   polymerase A/B and C.
332

Chromosome-level assembly of the maned sloth

To obtain a final scaffolded assembly of *Bradypus torquatus*, we used the Arima HiC protocol,
which is applicable to ethanol-preserved samples [23,38], to generate 97.5 Gb in long-range
read pair data. Using the automated scaffolding software yahs [39] and manual curation, our
contig assembly could be scaffolded into chromosome-level scaffolds (Figure 3A). This final
assembly consists of 2,915 scaffolds and 5,022 contigs. The scaffold N50 and N90 values are
157 Mb and 61.3 Mb, respectively (Figure 3B). The contig N50 and N90 values are 4.75 Mb
and 519 kb, respectively. Using Merqury [40] with the HiFi reads, we estimate a high base
accuracy (QV=46.7), which represents an upper bound as the HiFi reads were also used for
assembly. The assembly has a compleasm gene completeness score of 97.3% based on the
eutheria ODB10 database with n=11,366 genes (Supplementary Table 4) and contains
90.72% of ancestral placental mammal genes.
345

346
Figure 3: Chromosome-scale assembly of *Bradypus torquatus*.

(A) HiC interaction map after automated scaffolding by yahs and manual curation. The HiC map shows
interactions in 3-dimensional space between two regions of the genome. Darker colors indicate a higher
number of interactions. The region of low interaction between scaffold 7 and all other scaffolds indicates
this scaffold is the X chromosome, which was confirmed as this scaffold aligns to the human X
chromosome.
(B) Snail plot showing lengths of all scaffolds, together with the longest scaffold (red), and the N50 (dark
orange) and N90 length (light orange). The outer ring shows the GC content of the genome.

355
356
357

In comparison to existing genome assemblies of xenarthran species, our final assembly clearly
outperforms the short-read based assembly of the sloth *Choloepus hoffmanni* in terms of
contiguity and the number of intact ancestral placental mammal genes (Figure 4). Although
other long-read based xenarthran assemblies, which were most likely generated from flash-
frozen samples obtained from zoos and captive colonies, have even higher contiguities, our
Bradypus torquatus assembly is a valuable addition for xenarthran and, more generally,
mammalian comparative genomics.

366
Figure 4: Comparison of xenarthran genome assemblies.
(A) Visualization of contig sizes of available xenarthran genome assemblies. Each bar represents the
total assembly size. Contigs shorter than 1 Mb are not visualized individually, but shown as the grey
portion of each bar. The final *B. torquatus* assembly and its preliminary assembly generated only from

polymerase A/B reads are in red font. Assembly source or accession is listed in this panel, the
sequencing technology used is listed in the inset in panel B.
(B) Visualization of assembly contiguity as an N(x) graph, showing contig sizes on the Y-axis, for which
x percent of the assembly consists of contigs of at least that size. Assembly order in the legend (inset)
is sorted by contig N50 value.
(C) TOGA classification of 18,430 ancestral placental mammal genes showing the number of genes
that have an intact reading frame (blue bar, number is given in white font), inactivating mutations (e.g.
frameshifts, stop codon, splice site mutations or exon deletions; orange bar), or missing coding
sequence parts often caused by assembly gaps or fragmentation (gray bar). Assemblies are sorted by
the number of intact genes.
381
382

Polymerase C improves assemblies for various species

We next explored whether polymerase C can also help to improve assemblies of other
species, using samples not obtained from collections. To provide a fair comparison, we
randomly downsampled the larger data set to obtain an equal coverage of HiFi reads
generated with polymerases A/B and C. To compare these polymerases for another mammal,
we used the human HG002 sample and generated assemblies for both human haplotypes.
Using an equal coverage of 23.5X, the polymerase A/B read data produced a 2.96 Gb
assembly for haplotype 1 with a contig N50 value of 642 kb, whereas the polymerase C data
generated a 3.03 Gb assembly with a substantially higher contig N50 value of 2.8 Mb, a 4.4
fold increase in contiguity. Consistently, gene completeness assessed with compleasm
(mammalia_odb10) increased substantially from 81.2 to 98.6%. Similar results were obtained
for the haplotype 2 assembly, where the polymerase A/B read data produced a 2.9 Gb
assembly with a contig N50 value of 558.8 kb and a gene completeness of 77.3%, whereas
the polymerase C read data produced a 3 Gb assembly with a contig N50 value of 2 Mb and
a gene completeness of 97.8%.
398
Since PCR amplification may produce chimeric reads [41], we used available non-amplified
human HiFi reads produced from the HG002 sample as a baseline to compare the amount of
chimeric HiFi reads generated by polymerase A/B and C. We mapped reads to the HG002
assembly [6] and computed the number of reads with supplementary alignments, which
indicate chimers. We found that the fraction of chimeric alignments is very low (≤0.81%) across
all three libraries, with polymerase C reads having the lowest fraction (Supplementary Figure
4A). We next included available HG002 read data obtained by Multiple Displacement
Amplification (MDA) in this comparison. Consistent with previous observations [41,42], the
majority of MDA alignments (69.3%) are chimers, which is further supported by the
observation that the primary alignment lengths are much shorter than the MDA reads
(Supplementary Figure 4). We therefore conclude that long range PCR amplification used in
the original and modified ultra-low input protocol does not create more chimeric reads than
non-amplified libraries and orders of magnitude fewer chimeric reads than MDA libraries.
412
413 Next, we explored the application of polymerase C to three non-vertebrate taxa covering two
414 additional phyla, Mollusca (two taxonomic classes: Gastropoda and Bivalvia) and Arthropoda
415 (Collembola), using taxa where genome sequencing efforts often rely on the amplification-

416 based protocols because of low sequencing performance with low input protocol or very small
417 DNA amounts.
418
For the sacoglossan gastropod *Elysia timida* (Mollusca), previous sequencing libraries created
with the low input protocol resulted in very poor sequencing performance. Therefore, we
applied the ultra-low input protocols, and compared two SMRT cells produced with polymerase
A/B, providing 16.6 and 20.8 Gb yield in reads with an N50 length of 6.5 and 5.8 kb, to one
SMRT cell produced with polymerase C, providing 23 Gb yield in reads with an N50 length of
7 kb (Supplementary Table 5). After subsampling to equal read coverage of 26.4X,
polymerase A/B and C read data generated assemblies with similar contig N50 values of 347.1
kb for polymerase A/B and 331 kb for polymerase C (Figure 5, Supplementary Table 5). Using
all polymerase A/B read data with a coverage of 42.5X increased the contig N50 value to
472.6 kb. Importantly, adding the 23 Gb of polymerase C reads, increased the contig N50
value 1.4 fold to 675.8 kb (Figure 5). While the gene completeness (metazoa_odb10) of 97.7
and 97.8% is similar between these assemblies, polymerase C data helped to improve
assembly contiguity for this mollusc.
432
To understand why polymerase C alone does not result in a more contiguous assembly, we
mapped both polymerase A/B and C reads to the *Elysia timida* assembly with highest
contiguity, generated from all read data. This showed that both polymerases A/B and C exhibit
bias; however, bias of one polymerase can be compensated by reads of the other
(Supplementary Figure 5), indicating that these polymerases may have taxon-specific
differences.
439
For the marine bivalve *Scintilla philippinensis* with an estimated genome size of 1.3 Gb, we
compared assemblies produced from 22.3 Gb of reads obtained from ultra-low input libraries
using polymerase A/B or C, which corresponds to a coverage of 17.1X. While the polymerase
A/B reads produced a 1.77 Gb assembly with a contig N50 value of 43.3 kb, the polymerase
C read data produced a 1.88 Gb assembly with a 1.2 fold increased contig N50 value of 52.1
kb. Gene completeness (metazoa_odb10) improved slightly from 89.1% (polymerase A/B) to
89.9% (polymerase C). Combining all polymerase A/B and C read data (coverage of 36.1X)
produced a 1.86 Gb assembly with an even higher contig N50 value of 75.1 kb (Figure 5,
Supplementary Table 5) and a higher gene completeness of 93.5%. Mapping polymerase A/B
and C reads to the assembly generated with all data also revealed regions that were covered
only by reads from one polymerase (Supplementary Figure 6A,B). While polymerase C reads
improved assembly contiguity of both mollusc species, the resulting assemblies have a
comparatively low contiguity, highlighting the challenges of sequencing molluscan DNA.
453
We next tested our adjusted protocol on a species having a very small body size, where
amplification of the limited amount of genomic DNA is required for long-read sequencing and
genome assembly [32]. We used an ethanol-preserved, whole single specimen of the
springtail *Podura aquatica* (Arthropoda: Collembola), which has a body size of only 1.5 mm
and an expected genome size of 200-300 Mb. The polymerase A/B run yielded 13.9 Gb of
HiFi reads with an N50 length of 9.6 kb. The polymerase C run yielded 21.7 Gb of reads but
with a lower N50 read length of 5.7 kb, which is likely explained by sequencing DNA one year
after the initial extraction (the entire specimen was used for the initial DNA extraction).

Strikingly, at an estimated coverage of ~50X, the polymerase A/B read data produced a 278.5
Mb assembly with a contig N50 value of only 919 kb, whereas the polymerase C data
generated a 269.3 Mb assembly with a contig N50 value of 2.77 Mb (Figure 5, Supplementary
Table 5). This represents a 3 fold increase in contiguity, despite the polymerase C reads being
substantially shorter. Gene completeness (arthropoda_odb10) increased slightly from 92.8%
for polymerase A/B assembly to 93.4% for polymerase C assembly. Combining all polymerase
A/B and C read data resulted in a 284.7 Mb assembly with an even higher contig N50 value
of 5.74 Mb and the same gene completeness of 93.4%. Similar to *Elysia* and *Scintilla*, aligning
reads to the most contiguous assembly showed complementary coverage dropouts
(Supplementary Figure 6C,D).

473
Figure 5: Impact of polymerase C on assemblies of mollusc and collembola species.
Assembly contiguity visualized as N(x) graphs that show contig sizes on the Y-axis, for which x percent
of the assembly consists of contigs of at least that size (N50 and N90 values are indicated). Assemblies
are generated with an equal (downsampled) coverage of reads from polymerase A/B (dotted lines) and
C (dashed lines). Assemblies generated with all data are shown as solid lines. Colors refer to different
species.

Together, these tests confirm that polymerase C improves the assembly contiguity and
sometimes gene completeness for a broad range of species, including species that rely on
amplification-based library preparation protocols because their small size does not provide
enough DNA from a single individual or because naturally-occurring metabolites presumably
inhibit the polymerase during sequencing.

Discussion

Our investigation into utilizing collection samples for long-read sequencing confirms that ethanol-preserved samples can contain kilobase-sized DNA, long enough for long-read sequencing [22,23]. For the two catfish species, we found that amplification-free protocols generated sequencing data sufficient to generate assemblies with contig N50 values surpassing 2 Mb. Application of amplification-free protocols is recommended whenever feasible, as they will not suffer from PCR bias. Our other tests indicate that mammal or reptile samples may necessitate amplification-based protocols. It remains to be investigated for which taxonomic groups amplification-free protocols are generally successful. We demonstrate that PCR bias associated with the amplification-based PacBio ultra-low input protocol can be overcome or at least mitigated by employing an alternative polymerase. As a proof of concept, the contiguous 3.1 Gb genome assembly of *B. torquatus* shows that a modified amplification-based protocol can produce high-quality assemblies of gigabase-sized genomes.

Contamination caused by sample decomposition, human handlers, or commensal bacteria is expected for collection samples that have been stored under non-sterile conditions [33]. It is difficult to assess contamination prior to sequencing, and we find different levels of contamination in our samples, ranging from most of sequenced reads stemming from contaminants to almost no contamination. Analyzing a low coverage of sequencing reads for contamination before sequencing a sample to the coverage required for assembly could therefore be a cost-efficient strategy to select those samples that contain sufficiently low contamination levels. Furthermore, the resulting assemblies should be carefully screened for contamination using existing methods [43,44].

Consistent with previous observations [34], we find that sample age alone is not an accurate predictor of input DNA quality and sample suitability for sequencing. For example, while the *B. torquatus* sample was collected in 2003, several younger samples exhibited high degrees of DNA degradation (Supplementary Table 1). Hence, in addition to sample age, other factors such as storage temperature and conditions, storage medium, or tissue type likely influence DNA quality. From our experience, samples consistently stored at -20°C and preserved in 96% ethanol perform well, but a systematic assessment of larger sample numbers is needed to substantiate this.

Our study has a number of implications. First, the modified ultra-low input protocol improves genome assembly of small specimens, where amplification is a requirement to obtain enough DNA for sequencing. For example, the contiguity of the *Podura aquatica* genome increased to an N50 of 5.7 Mb, and thus substantially exceeds the minimum standards of 100 kb set by the Earth Biogenome Project for small species with limited DNA amounts [1]. The modified protocol will likely not only be beneficial for species with diminutive body sizes that represent a very large but mostly uncharacterized part of Earth's biodiversity, but also in cases where only very limited amounts of material from non-lethal samplings (biopsies from human patients or bat wing punches) are available. Second, long-read sequencing remains a challenge for molluscs and other taxonomic groups, where satisfactory sequencing outputs often require amplification-based protocols. Although achieving highly contiguous assemblies with

megabase contig N50 values remains challenging for these species, our investigations
suggest that employing a combination of different polymerases can at least help to improve
assembly contiguity. Third, while the PacBio ultra-low input protocol was previously limited to
genome sizes of up to 500 Mb, the successful application of the modified protocol to *B.
torquatus* with its 3.1 Gb genome extends its applicability to a broad range of species with
larger genome sizes. Together, the improved efficiency of the modified ultra-low input protocol
opens avenues for generating contiguous genomes across various species.
540
Our study raises the question of finding polymerases with minimal bias. While our tests with
B. torquatus and human indicate that polymerase C shows satisfactory performance for
mammals, we found that polymerase C also appears to exhibit bias for samples of molluscs
and collembola, albeit a different bias compared to polymerase A/B (Supplementary Figures
5, 6). Anticipating that DNA amplification will constitute a key step in the genome sequencing
procedure for numerous collection samples, challenging species, and species with diminutive
body sizes, future investigations could focus on identifying the most appropriate polymerase
or combination of polymerases that exhibit minimal bias for specific taxonomic groups.
549
Apart from the ultra-low input protocol, several new approaches have recently been developed
to make small amounts of input DNA accessible for long read sequencing. This includes the
above-mentioned MDA [41,42,45], adapter ligation via tagmentation [46,47], and Picogram
input multimodal sequencing (PiMmS) [48]. We show here that the ultra-low input protocol
produces very few chimeric reads in contrast to MDA. Furthermore, the ultra-low input protocol
can generate average read lengths of ~10 kb, which is similar to read lengths generated by
PiMmS [48], but substantially longer than those generated with tagmentation based
approaches (2.5-5 kb averages) [46,47]. Nevertheless, different methods likely have ideal
application ranges that depend on the input sample, its quality and amount of DNA. Future
research should therefore benchmark which library preparation method is optimal for which
sample type.
561

Conclusions

563 Our work suggests that collections can complement flash-frozen material as a sample source
564 for biodiversity genomics, especially for species that are hard to sample because of rarity,
565 protection status or other reasons. Thus, natural history collections as extensive archives of
566 biodiversity can help to achieve the ambitious goal of generating reference genomes for all life
567 on Earth.

Material and Methods

Sample sources

For *Bradypus torquatus*, we used a sample of ~50 mg of clogged blood, preserved in ethanol. This sample was collected in 2003 under Brazilian license SISGEN number AF86294 and CITES number 138261. For *Idiurus macrotis*, we used ~12 mg of skin with hair preserved in technical ethanol at room temperature. For the *Anguis fragilis*, we used ~51 mg (for the one collected in 2021) and ~3 mg (for the one collected in 1878) of muscle tissue from a tail cross-section. Both samples were preserved in technical ethanol at room temperature. For both *Cathorops* species, we used fin samples stored in ethanol in frozen collections at the Leibniz Institute for the Analysis of Biodiversity Change (LIB) Bonn. Originally, fin clips of specimens acquired from local fishermen were taken in 2014, immediately placed into ethanol, but subsequently transported multiple times at room temperature until final storage at -20°C. The exact time between catch and sampling is unknown but was likely a few hours. For *Desmana moschata*, we used ~9 mg of muscle and skin tissue that was preserved in ethanol at room temperature. For *Muscardinus avellanarius*, we used ~19 mg of foot tissue that was preserved in technical ethanol at room temperature. For *Dipus sagitta*, we used ~12 mg (individual 95545) and ~16 mg (individual 95541) of muscle tissue and ~30 mg (individual 56492) of skin. All three samples were preserved in technical ethanol at room temperature. For *Ptilocercus lowii*, we used ~8 mg of muscle tissue that was preserved in technical ethanol at room temperature. For *Xerotyphlops vermicularis*, we used ~5 mg (individual collected in 2004) and ~3 mg (individual collected in 2011) of skin and muscle tissue preserved in technical ethanol at room temperature. For *Elysia timida*, we used a whole specimen (~1 cm body length) from our living culture, which we immediately homogenized for DNA extraction after euthanization. This sample was collected under license ESNC 205 issued by the Spanish "Dirección General de Biodiversidad, Bosques y Desertificación del Ministerio para la Transición Ecológica y el Reto Demográfico". For *Scintilla philippinensis*, we used ~20 mg of muscle tissue preserved in ethanol, collected in Johor Malaysia under a collaboration agreement between Senckenberg and Universiti Putra Malaysia. For *Podura aquatica*, we used a single whole specimen (~1.5 mm body length) killed and immediately preserved in 96% ethanol. Two libraries were produced either with polymerase A/B or polymerase C (below), and while the polymerase A/B experiment was done within the month following DNA extraction, the polymerase C experiment was conducted one year after DNA extraction, using DNA preserved at -20°C in TE buffer. Supplementary Table 1 lists sample sources, accessions and additional details.

DNA extraction

High molecular weight (HMW) gDNA was extracted from ethanol-preserved tissues of *Bradypus torquatus*, using a modified protocol version of the Circulomics Nanobind Tissue Big DNA kit, including the ethanol removing step described in 'Guide and overview – Nanobind tissue kit'. We retrieved gDNA bound to the Nanobind disk as well as unbound gDNA in the precipitation solution. The gDNA bound to the Nanobind disk was eluted after several washing steps. The unbound gDNA in the precipitation solution was precipitated by centrifugation (18.000 xg for 30 min at 4°C). The resulting pellet was washed twice with 75% ice-cold ethanol, air dried for 20 min at room temperature and resuspended in 1x elution buffer. For both gDNA extractions we performed standard quality control, which involved Qubit quantification,

613 Nanodrop measurement, and pulse-field gel electrophoresis making use of the Femto Pulse
614 system (Agilent Technologies).
615
For *Idiurus macrotis*, *Desmana moschata*, *Muscardinus avellanarius*, *Cathorops nuchalis*,
Cathorops wayuu, *Xerotyphlops vermicularis, Dipus sagitta*, *Ptilocercus lowii* and the two
Anguis fragilis samples, gDNA was extracted according to the protocol of [49]. DNA
concentration and DNA fragment length were assessed using the Qubit dsDNA BR Assay kit
on the Qubit Fluorometer (Thermo Fisher Scientific) and the Genomic DNA Screen Tape on
the Agilent 4150 TapeStation system (Agilent Technologies). For *Elysia timida* and *Scintilla*
philippinensis, gDNA was extracted using a CTAB-based method [50] and a bead-based
protocol [51], respectively, including a pre-wash with sorbitol. The MagAttract HMW DNA Kit
from Qiagen was used to extract gDNA from *Podura aquatica.* For these gDNA extractions,
DNA concentration and DNA fragment length were assessed using Qubit quantification
(Thermo Fisher Scientific), the Agilent 2200 TapeStation system (Agilent Technologies) and
the Femto Pulse system (Agilent Technologies).
628
629 All details on the DNA yield and DNA fragment sizes can be found in Supplementary Table 1.
630

631 ## Low input PacBio HiFi library preparation
632 The low input protocol allows generating PacBio libraries for samples with limited DNA content
633 without amplification [26]. We prepared low input PacBio HiFi libraries according to the
634 instructions of the SMRTbell Express Prep Kit v2.0, except for the libraries of *Cathorops*
635 *nuchalis* and *Cathorops wayuu* which were prepared with the SMRTbell prep kit v3.0.
636

Ultra-low input PacBio HiFi library preparation
PacBio ultra-low input HiFi libraries were prepared with the SMRTbell Express Template Prep
Kit 2.0 according to the 'Procedure & Checklist - Preparing HiFi SMRTbell® Libraries from
Ultra-Low DNA Input' (PN 101-987-800 Version 02). To reduce potential PCR bias of
polymerase A/B, we used in our modified protocol a third PCR reaction, making use of
Polymerase C (KOD Xtreme™ Hot Start DNA Polymerase, Merck PN 71975), which is
optimized for the amplification of long strands and GC-rich DNA templates.
644
645 The amplified DNA from two PCR reactions with polymerase A and B was pooled equimolarly.
646 PCR fragments from polymerase C amplification were kept separately and processed
647 independently from the pooled fragments produced with polymerase A and B. Purified and
648 pooled amplified DNA libraries were size selected to remove smaller fragments
649 (Supplementary Table 1).
650
651 For *Anguis fragilis* and *Idiurus macrotis*, we prepared two additional libraries with DNA extracts
652 to which a DNA repair step was applied using the Sequential Reaction Protocol for PreCR
653 Repair Mix (New England BioLabs) prior to the actual library preparation.
654

PacBio sequencing
A total of 27 SMRT 8M cells were sequenced in CCS mode using the PacBio Sequel II / IIe
instrument. For low input libraries, where possible, libraries were loaded at an on-plate

concentration of 80 pM using adaptive loading and the Sequel II Binding kit 2.2 or 3.2 (Pacific
Biosciences, Menlo Park, CA). Ultra-low input libraries were loaded with up to 80 pM on plate
where possible using the SEQUEL II binding kit 2.2 or 3.2, and the sequencing kit 2.0. Pre-
extension time was 2 hours, run time was 30 hours.
662

663 HiC for scaffolding the *B. torquatus* assembly

Chromatin conformation capture was done using the Arima HiC+ Kit (Material Nr. A410110),
following the user guide for animal tissues (ARIMA-HiC 2.0 kit Document Nr: A160162 v00)
and processing 28 mg of tissue with the standard input approach. The subsequent Illumina
library preparation followed the ARIMA user guide for Library preparation using the Kapa
Hyper Prep kit (ARIMA Document Part Number A160139 v00). The barcoded HiC libraries
were run on an S4 flow cell of a NovaSeq6000 with 200 cycles.
670

671 Comparing polymerase A/B and C read assemblies

Aiming to evaluate the impact of libraries generated with polymerase A/B vs. C on the genome
assembly quality, we combined different datasets with varying coverages, library complexities
(number of libraries) and polymerase combinations (only A/B, only C, and A/B+C). For tests
that did not involve all read data, we randomly subsampled reads. Subsequently, we
assembled the read data into a contig assembly, as described below, and compared the
summary metrics, including contig N50, number of contigs and gene completeness. All results
are listed in Supplementary Tables 4 and 5.
679

680 Contig assembly

HiFi reads were called using a pipeline consisting of PacBio's tools ccs 6.4.0
(https://github.com/PacificBiosciences/ccs) and actc 0.3.1
(https://github.com/PacificBiosciences/actc) as well as samtools 1.15 [52] and
DeepConsensus 0.2.0 or 1.2.0 [27]. All commands were executed as recommended in the
respective guide for DeepConsensus
(https://github.com/google/deepconsensus/blob/v0.2.0/docs/quick_start.md; e.g. ccs --all). To
remove PCR adapters and PCR duplicates, which might originate from the PCR amplification
during the ultra-low library preparation, PacBio's tools lima 2.6.0
(https://github.com/PacificBiosciences/barcoding) with options "--num-threads 67 --split-bam-
named --same" and pbmarkdup 1.0.2-0 with options "--num-threads 67 --log-level INFO --log-
file pbmarkdup.log --cross-library --rmdup"
(https://github.com/PacificBiosciences/pbmarkdup) were applied to samples prepared with the
ultra-low library preparation protocol. For the Catfish samples *Cathorops nuchalis* and *C.*
wayuu that were sequenced using the low-input library preparation protocol, PacBio
sequencing adapters were removed with HiFiAdapterFilt [53]. The resulting reads were
merged and then decontaminated with kraken2 v. 2.1.3 [54] using the kraken2 PlusPFP
database downloaded in March 2023, with a confidence score of 0.51.
698

After HiFi calling, we used hifiasm v0.19.5 [28,55] to assemble HiFi reads obtained from the
Cathorops nuchalis, C. wayuu, Idiurus, Anguis, Elysia, Scintilla and *Podura* samples. For the
two catfish samples *Cathorops nuchalis* and *C. wayuu*, because of suboptimal performance
with default parameters, we tested several hifiasm options before deciding which parameters

produce the best assembly in terms of gene completeness and contiguity (Supplementary
Table 2). To this end, we estimated the genome profile of these two species with FastK
(https://github.com/thegenemyers/FASTK)   and   Genescope.FK
(https://github.com/thegenemyers/GENESCOPE.FK) with k=30 to find the homozygous peak
that was then passed to hifiasm (Supplementary Table 2). In all other cases, we applied default
parameters with strict haplotig purging (-l3 parameter), and for the *Elysia* sample, we
additionally used available Arima HiC data for assembly phasing.
710
Contiguity statistics were calculated with Quast 5.0.2 [56], gfastats v. 1.3.6
(https://github.com/vgl-hub/gfastats)   and   Merqury.FK
(https://github.com/thegenemyers/MERQURY.FK). Gene completeness was evaluated with
BUSCO 5.5.0 [30] as well as compleasm 0.2.5 [29]. We used the eutherian_odb10 dataset for
Bradypus torquatus, and actinopterygii_odb10 for *C. nuchalis* and *C. wayuu*, the
mammalia_odb10 dataset for human, the arthropoda_odb10 dataset for *Podura aquatica,* and
the metazoa_odb10 dataset for *Elysia timida* and *Scintilla philippinensis*.
718
For *B. torquatus*, we initially obtained hifiasm (v0.19.5) assemblies that were of a size
expected from four haplotypes of this genome, consisting of a large number of small contigs
(Supplementary Table 6). Similar results were obtained with HiCanu (v2.2) [57], which is
designed to break contigs at all joins in the assembly graph, meaning any divergences
between the four theoretical haplotypes would result in a new contig (in our case over 200,000
assembled contigs totaling almost 12 Gb of sequence). This indicated that the tissue samples
we obtained for this species originated from two different individuals. While the accuracy of
the PacBio HiFi reads should in principle allow to distinguish all four haplotypes, *B. torquatus*
is expected to have a very low heterozygosity and high in-breeding rate due to small population
size, which results in assembly graphs where many regions collapse all haplotypes due to the
lack of sequence variation.
730
To overcome this problem, we used the assembler Flye (v2.9.2) [58], which allows users to
set the read error rate as an argument. Flye has been previously suggested by the developers
as a method for collapsing sequences from highly diverged haplotypes into a single "pseudo-
haplotype" sequence (https://github.com/fenderglass/Flye/issues/636). Here, we found that a
read error-rate of 3% produced the most contiguous assembly, when combined with a reduced
read-overlap of 5 kb (Supplementary Table 6). The latter deviates from the default value
selected by Flye, which Flye would determine by the N90 of the input reads (in our case the
N90 was 9 kb for the HiFi library, which had a modal read length of ~10 kb). We then removed
retained haplotigs using purge-dups [59].
740

Contamination detection and read coverage analysis

Specimens stored in liquid preservation media are prone to various levels of DNA
contamination from non-target organisms [33], caused by different handling and storage
conditions that are often hard to retrace [60]. To detect levels of contamination from
exogenous DNA in our assemblies, we used NCBI's Foreign Contamination Screen (FCS
0.5.0) [43], which flags both putative adapter sequences (FCS-adaptor) and contigs assigned
to non-target species (FCS-GX). Both FCS tools were executed from the provided singularity

container using singularity 1.2.4. FCS-adaptor was executed through the provided bash script
(run_fcsadaptor.sh) with the option for eukaryotes (--euk). FCS-GX was executed by the
python wrapper (fcs.py screen genome) together with the corresponding NCBI taxonomy ID
and the GX database (as of Dec 5th, 2023). Furthermore, to visualize contamination across
the respective contig-level assemblies before FCS-filtering, we used blobtoolkit v4.1.4 [44],
which assigns all contigs from a given assembly to a taxonomic group based on best blast hits
(Supplementary Figure 2).
755
Additionally, to assess pre-assembly read quality, we mapped reads obtained from samples
of *Bradypus torquatus*, *Anguis fragilis* and *Idiurus macrotis* to available reference genomes of
closely related species *Choloepus didactylus* (GCA_015220235.1), *Elgaria multicarinata*
(GCA_023053635.1) and *Pedetes capensis* (GCA_007922755.1). Similarly, to identify regions
of PCR coverage dropouts, we aligned reads from polymerase A/B or C libraries to the best
(defined as highest contig N50, Supplementary Table 5) assemblies obtained for *Podura*,
Scintilla and *Elysia*, and visually inspected mapped reads (Supplementary Figures 5, 6).
763
To further quantify PCR bias, we calculated the normalized coverage (coverage of each
nucleotide divided by the average coverage) of each polymerase A/B and C *Bradypus*
torquatus library, using either the *Choloepus didactylus* genome or the best assembly of
Bradypus torquatus. We also calculated normalized coverage of a non-amplified human library
(downloaded from https://downloads.pacbcloud.com/public/revio/2022Q4/HG002-rep1/; last
accessed 19 Sep 2024) as well as polymerase C (produced in this study) and A/B amplified
libraries (NCBI, BioProject PRJNA657245, accessions SRR12454519 and SRR12454520)
sequenced from the human cell line HG002, using the human HG002 assembly [6] (v.1.1,
maternal haplotype). We then computed normalized coverage across nucleotides assigned to
exonic and repeat sequences. For *C. didactylus* and *B. torquatus*, exons were annotated by
TOGA v1.0.0 and repeats were annotated with RepeatModeler [61] and RepeatMasker v4.1.4
(https://www.repeatmasker.org/). For human, exons were annotated by RefSeq (v110 from
CHM13, JHU v5.2), https://ccb.jhu.edu/T2T.shtml) and annotated repeats [62] were
downloaded from the UCSC table browser [63] (last accessed 19 Sep 2024). Read mapping
was performed using minimap2 v2.26 [64] with HiFi read mapping parameters (--ax map-hifi),
and absolute coverage per base and across annotations was computed with samtools v1.17
[52], using the 'samtools depth' and 'samtools bedcov' commands, respectively. For the
HG002 gene annotation, we filtered the annotation to only include coding exons to enable a
fair comparison with the TOGA annotations that do not include non-coding transcripts or
UTRs.
784

Scaffolding the final *B. torquatus* genome

To scaffold these *B. torquatus* contigs, we mapped HiC reads to the contig assembly using
bwa-mem (v.0.7.17) [65], before the resulting HiC alignment file was filtered, sorted and
deduplicated with pairtools parse, pairtools sort and pairtools dedup (v0.3.0), respectively. The
processed HiC alignments were then used as input for scaffolder yahs (v1.2a.1.patch) [39]. A
full list of commands is given in Supplementary Note 1. After initial automated scaffolding with
yahs, we ran multiple rounds of manual curation based on the HiC interaction maps. This
involved re-ordering and re-orienting the scaffolded sequences based on sequences close to

each other in the genome, which are expected to have a higher number of HiC interactions than those further apart. Using this method, we were able to obtain chromosome-level scaffolds of the 24 autosomes and the X chromosome. This assembly was then again screened for adapter and foreign sequence contaminants using NCBI's FCS-adaptor and FCS-GX tools [43]. We subsequently removed contaminant sequences by applying the python wrapper (fcs.py clean genome) together with the action report from "screen genome" and setting the minimum sequence length to 1 bp (--min-seq-len 1).

Read chimer analysis

To investigate whether our modified amplification based protocol creates more chimeric reads, we mapped reads (all obtained from the human HG002 sample) against the HG002 reference genome [6] (v.1.1, maternal haplotype, https://github.com/marbl/hg002?tab=readme-ov-file), using minimap2 v2.26 [64] with HiFi read mapping parameters (--ax map-hifi). We used reads amplified with polymerase C and polymerase A/B (NCBI BioProject PRJNA657245, accessions SRR12454519 and SRR12454520), as well as the non-amplified reads (https://downloads.pacbcloud.com/public/revio/2022Q4/HG002-rep1/; last accessed 19 Sep 2024), and reads amplified with MDA (NCBI BioProject PRJNA1005794, accession SRR25653511). To calculate the fraction of alignments classified as primary alignments, secondary alignments, supplementary alignments and unmapped, we counted the flags assigned by minimap using samtools v1.17 [52] with the command 'samtools view'. Raw read lengths and alignment lengths of primary and supplementary alignments were extracted from raw fastq-files and sam-files created by minimap2, respectively.

Competing interests

The authors have no competing interests.

Acknowledgment

We thank Deniz Kaya from PacBio for advice and suggestions on adapting the ultra-low input protocol, and Sarah Kingan, Juniper Lake, Ian McLaughlin, Aaron Wenger and Jonas Korlach from PacBio for the polymerase C runs for the human sample. We also thank the Genome Technology Center (RGTC) at Radboudumc for the use of the Sequencing Core Facility (Nijmegen, The Netherlands), which provided the PacBio SMRT sequencing service on the Sequel IIe platform, the Long Read Team of the DRESDEN Concept Genome Center, part of the MPI-CBG and the technology platform of the CMCB at the TU Dresden, supported by DFG (INST 269/768-1), and the HPC Service of FUB-IT, Freie Universität Berlin, for computing time (doi:10.17169/refubium-26754). We acknowledge Irina Ruf (Senckenberg Frankfurt), Carles Galià Camps (University of Barcelona), Madlen Stange, Claudia Koch, Morris Flecks, Jan Decher and Christian Montermann (Leibniz Institute for the Analysis of Biodiversity Change) for providing samples and Sandra Kukowka for helping in subsampling specimens.

Funding

This work was supported by a grant from the Leibniz Association's Competition Procedure (K419/2021) and the LOEWE-Centre for Translational Biodiversity Genomics (TBG) funded by the Hessen State Ministry of Higher Education, Research and the Arts (LOEWE/1/10/519/03/03.001(0014)/52).

Data and Code Availability

The raw sequencing data and assemblies for *Bradypus torquatus* are available at NCBI under BioProject PRJEB73341 and BioSample SAMEA115348596. An improved version of the *Elysia timida* assembly after incorporating additional polymerase C reads and Hi-C scaffolding [66] is available under Bioproject PRJNA1119176 and Biosample SAMN42332041. The *Scintilla philippinensis* assembly and raw sequencing data are available under Bioproject PRJNA1120792. Genome assemblies and raw sequencing data of both catfish genomes are available under Bioproject PRJNA1162287 (*Cathorops nuchalis*) and PRJNA1162286 (*Cathorops wayuu*). The *Podura aquatica* assembly and sequencing data are available under Bioproject PRJNA1163304. Raw reads and assemblies obtained with polymerase C for HG002 are available on https://downloads.pacbcloud.com/public/revio/2023Q3/KODXtreme/. The TOGA annotation for the *B. torquatus* is available at https://genome.senckenberg.de/download/TOGA/. Ultra-low input based assemblies generated in this study are also available at https://genome.senckenberg.de/download/GenomesCollectionsPolC. No new computer code was generated in this study.

References

1. Lewin HA, Robinson GE, Kress WJ, Baker WJ, Coddington J, Crandall KA, et al. Earth BioGenome Project: Sequencing life for the future of life. Proc Natl Acad Sci U S A. 2018;115:4325–33.

2. Zoonomia Consortium. A comparative genomics multitool for scientific discovery and conservation. Nature. 2020;587:240–5.

3. Feng S, Stiller J, Deng Y, Armstrong J, Fang Q, Reeve AH, et al. Dense sampling of bird diversity increases power of comparative genomics. Nature. 2020;587:252–7.

4. Ronco F, Matschiner M, Böhne A, Boila A, Büscher HH, El Taher A, et al. Drivers and dynamics of a massive adaptive radiation in cichlid fishes. Nature. 2021;589:76–81.

5. Kuderna LFK, Gao H, Janiak MC, Kuhlwilm M, Orkin JD, Bataillon T, et al. A global catalog of whole-genome diversity from 233 primate species. Science. 2023;380:906–13.

6. Nurk S, Koren S, Rhie A, Rautiainen M, Bzikadze AV, Mikheenko A, et al. The complete sequence of a human genome. Science. 2022;376:44–53.

7. Chen J, Wang Z, Tan K, Huang W, Shi J, Li T, et al. A complete telomere-to-telomere assembly of the maize genome. Nat Genet. 2023;55:1221–31.

8. Warren WC, Harris RA, Haukness M, Fiddes IT, Murali SC, Fernandes J, et al. Sequence diversity analyses of an improved rhesus macaque genome enhance its biomedical utility. Science [Internet]. 2020;370. Available from: http://dx.doi.org/10.1126/science.abc6617

9. Kautt AF, Kratochwil CF, Nater A, Machado-Schiaffino G, Olave M, Henning F, et al. Contrasting
signatures of genomic divergence during sympatric speciation. Nature. 2020;588:106–11.

10. Jebb D, Huang Z, Pippel M, Hughes GM, Lavrichenko K, Devanna P, et al. Six reference-quality
genomes reveal evolution of bat adaptations. Nature. 2020;583:578–84.

11. Rhie A, McCarthy SA, Fedrigo O, Damas J, Formenti G, Koren S, et al. Towards complete and
error-free genome assemblies of all vertebrate species. Nature. 2021;592:737–46.

12. Blumer M, Brown T, Freitas MB, Destro AL, Oliveira JA, Morales AE, et al. Gene losses in the
common vampire bat illuminate molecular adaptations to blood feeding. Sci Adv. 2022;8:eabm6494.

13. Vollger MR, Guitart X, Dishuck PC, Mercuri L, Harvey WT, Gershman A, et al. Segmental
duplications and their variation in a complete human genome. Science. 2022;376:eabj6965.

14. Osipova E, Barsacchi R, Brown T, Sadanandan K, Gaede AH, Monte A, et al. Loss of a
gluconeogenic muscle enzyme contributed to adaptive metabolic traits in hummingbirds. Science.
2023;379:185–90.

15. Shao Y, Zhou L, Li F, Zhao L, Zhang B-L, Shao F, et al. Phylogenomic analyses provide insights
into primate evolution. Science. 2023;380:913–24.

16. Hiller M, Morales A, Ahmed A, Hilgers L, Kirilenko B, Kontopoulos D, et al. Reference-quality bat
genomes illuminate adaptations to viral tolerance and disease resistance [Internet]. Research Square.
2023 [cited 2023 Sep 18]. Available from: https://www.researchsquare.com/article/rs-2557682/latest

17. Blom MPK. Opportunities and challenges for high-quality biodiversity tissue archives in the age of
long-read sequencing. Mol Ecol. 2021;30:5935–48.

18. Johnson KR, Owens IFP, Global Collection Group. A global approach for natural history museum
collections. Science. 2023;379:1192–4.

19. Espeland M, Breinholt J, Willmott KR, Warren AD, Vila R, Toussaint EFA, et al. A Comprehensive
and Dated Phylogenomic Analysis of Butterflies. Curr Biol. 2018;28:770–8.e5.

20. Heinicke MP, Nielsen SV, Bauer AM, Kelly R, Geneva AJ, Daza JD, et al. Reappraising the
evolutionary history of the largest known gecko, the presumably extinct Hoplodactylus delcourti, via
high-throughput sequencing of archival DNA. Sci Rep. 2023;13:1–12.

21. Tan HZ, Jansen JJFJ, Allport GA, Garg KM, Chattopadhyay B, Irestedt M, et al. Megafaunal
extinctions, not climate change, may explain Holocene genetic diversity declines in Numenius
shorebirds. Elife [Internet]. 2023;12. Available from: http://dx.doi.org/10.7554/eLife.85422

22. Mulcahy DG, Macdonald KS 3rd, Brady SG, Meyer C, Barker KB, Coddington J. Greater than X
kb: a quantitative assessment of preservation conditions on genomic DNA quality, and a proposed
standard for genome-quality DNA. PeerJ. 2016;4:e2528.

23. Dahn HA, Mountcastle J, Balacco J, Winkler S, Bista I, Schmitt AD, et al. Benchmarking ultra-high
molecular weight DNA preservation methods for long-read and long-range sequencing. Gigascience
[Internet]. 2022;11. Available from: http://dx.doi.org/10.1093/gigascience/giac068

24. Wenger AM, Peluso P, Rowell WJ, Chang P-C, Hall RJ, Concepcion GT, et al. Accurate circular
consensus long-read sequencing improves variant detection and assembly of a human genome. Nat
Biotechnol. 2019;37:1155–62.

25. Sharma P, Al-Dossary O, Alsubaie B, Al-Mssallem I, Nath O, Mitter N, et al. Improvements in the
sequencing and assembly of plant genomes. GigaByte. 2021;2021:gigabyte24.

26. Kingan SB, Heaton H, Cudini J, Lambert CC, Baybayan P, Galvin BD, et al. A high-quality DE
novo genome assembly from a single mosquito using PacBio sequencing. Genes . 2019;10:62.

27. Baid G, Cook DE, Shafin K, Yun T, Llinares-López F, Berthet Q, et al. DeepConsensus improves
the accuracy of sequences with a gap-aware sequence transformer. Nat Biotechnol. 2023;41:232–8.

28. Cheng H, Concepcion GT, Feng X, Zhang H, Li H. Haplotype-resolved de novo assembly using
phased assembly graphs with hifiasm. Nat Methods. 2021;18:170–5.

29. Huang N, Li H. compleasm: a faster and more accurate reimplementation of BUSCO.
Bioinformatics [Internet]. 2023;39. Available from: http://dx.doi.org/10.1093/bioinformatics/btad595

30. Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. BUSCO Update: Novel and
Streamlined Workflows along with Broader and Deeper Phylogenetic Coverage for Scoring of
Eukaryotic, Prokaryotic, and Viral Genomes. Mol Biol Evol. 2021;38:4647–54.

31. PacBio Biosciences. Now available: Ultra-low DNA input workflow for SMRT sequencing
[Internet]. 2020 [cited 2023]. Available from: https://www.pacb.com/blog/introducing-the-ultra-low-
input-protocol-for-smrt-sequencing/

32. Schneider C, Woehle C, Greve C, D'Haese CA, Wolf M, Hiller M, et al. Two high-quality de novo
genomes from single ethanol-preserved specimens of tiny metazoans (Collembola). Gigascience
[Internet]. 2021;10. Available from: http://dx.doi.org/10.1093/gigascience/giab035

33. Raxworthy CJ, Smith BT. Mining museums for historical DNA: advances and challenges in
museomics. Trends Ecol Evol. 2021;36:1049–60.

34. Irestedt M, Thörn F, Müller IA, Jønsson KA, Ericson PGP, Blom MPK. A guide to avian
museomics: Insights gained from resequencing hundreds of avian study skins. Mol Ecol Resour.
2022;22:2672–84.

35. Strunov A, Kirchner S, Schindelar J, Kruckenhauser L, Haring E, Kapun M. Historic museum
samples provide evidence for a recent replacement of Wolbachia types in European Drosophila
melanogaster. Mol Biol Evol [Internet]. 2023;40. Available from:
http://dx.doi.org/10.1093/molbev/msad258

36. Kirilenko BM, Munegowda C, Osipova E, Jebb D, Sharma V, Blumer M, et al. Integrating gene
annotation with orthology inference at scale. Science. 2023;380:eabn3107.

37. Gibb GC, Condamine FL, Kuch M, Enk J, Moraes-Barros N, Superina M, et al. Shotgun
Mitogenomics Provides a Reference Phylogenetic Framework and Timescale for Living Xenarthrans.
Mol Biol Evol. 2016;33:621–42.

38. Osipova E, Ko M-C, Petricek KM, Yung Wa Sin S, Brown T, Winkler S, et al. Convergent and
lineage-specific genomic changes contribute to adaptations in sugar-consuming birds [Internet].
bioRxiv. 2024. Available from: http://dx.doi.org/10.1101/2024.08.30.610474

39. Zhou C, McCarthy SA, Durbin R. YaHS: yet another Hi-C scaffolding tool. Bioinformatics
[Internet]. 2023;39. Available from: http://dx.doi.org/10.1093/bioinformatics/btac808

40. Rhie A, Walenz BP, Koren S, Phillippy AM. Merqury: reference-free quality, completeness, and
phasing assessment for genome assemblies. Genome Biol. 2020;21:245.

41. Hård J, Mold JE, Eisfeldt J, Tellgren-Roth C, Häggqvist S, Bunikis I, et al. Long-read whole-
genome analysis of human single cells. Nat Commun. 2023;14:5164.

42. Lee Y-C, Ke H-M, Liu Y-C, Lee H-H, Wang M-C, Tseng Y-C, et al. Single-worm long-read
sequencing reveals genome diversity in free-living nematodes. Nucleic Acids Res. 2023;51:8035–47.

43. Astashyn A, Tvedte ES, Sweeney D, Sapojnikov V, Bouk N, Joukov V, et al. Rapid and sensitive
detection of genome contamination at scale with FCS-GX. bioRxiv [Internet]. 2023; Available from:
http://dx.doi.org/10.1101/2023.06.02.543519

44. Challis R, Richards E, Rajan J, Cochrane G, Blaxter M. BlobToolKit - interactive quality
assessment of genome assemblies. G3 . 2020;10:1361–74.

45. Roberts NG, Gilmore MJ, Struck TH, Kocot KM. Multiple Displacement Amplification Facilitates
SMRT Sequencing of Microscopic Animals and the Genome of the Gastrotrich Lepidodermella

squamata (Dujardin, 1841) [Internet]. Genomics. bioRxiv; 2024. Available from: https://biorxiv.org/content/10.1101/2024.01.17.576123v1

46. Jia H, Tan S, Cai Y, Guo Y, Shen J, Zhang Y, et al. Low-input PacBio sequencing generates high-quality individual fly genomes and characterizes mutational processes. Nat Commun. 2024;15:5644.

47. Nanda AS, Wu K, Irkliyenko I, Woo B, Ostrowski MS, Clugston AS, et al. Direct transposition of native DNA for sensitive multimodal single-molecule sequencing. Nat Genet. 2024;56:1300–9.

48. Stevens L, Martínez-Ugalde I, King E, Wagah M, Absolon D, Bancroft R, et al. Ancient diversity in host-parasite interaction genes in a model parasitic nematode. Nat Commun. 2023;14:7776.

49. Sambrook J, Russell WD. Protocol: DNA isolation from mammalian tissue. Molecular Cloning: A Laboratory Manual (Sambrook J. 2001;

50. Murray MG, Thompson WF. Rapid isolation of high molecular weight plant DNA. Nucleic Acids Res. 1980;8:4321–5.

51. Mayjonade B, Gouzy J, Donnadieu C, Pouilly N, Marande W, Callot C, et al. Extraction of high-molecular-weight genomic DNA for long-read sequencing of single molecules. Biotechniques. 2016;61:203–5.

52. Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, et al. Twelve years of SAMtools and BCFtools. Gigascience [Internet]. 2021;10. Available from: http://dx.doi.org/10.1093/gigascience/giab008

53. Sim SB, Corpuz RL, Simmonds TJ, Geib SM. HiFiAdapterFilt, a memory efficient read processing pipeline, prevents occurrence of adapter sequence in PacBio HiFi reads and their negative impacts on genome assembly. BMC Genomics. 2022;23:157.

54. Wood DE, Lu J, Langmead B. Improved metagenomic analysis with Kraken 2. Genome Biol. 2019;20:257.

55. Cheng H, Jarvis ED, Fedrigo O, Koepfli K-P, Urban L, Gemmell NJ, et al. Haplotype-resolved assembly of diploid genomes without parental data. Nat Biotechnol. 2022;40:1332–5.

56. Mikheenko A, Prjibelski A, Saveliev V, Antipov D, Gurevich A. Versatile genome assembly evaluation with QUAST-LG. Bioinformatics. 2018;34:i142–50.

57. Nurk S, Walenz BP, Rhie A, Vollger MR, Logsdon GA, Grothe R, et al. HiCanu: accurate assembly of segmental duplications, satellites, and allelic variants from high-fidelity long reads. Genome Res. 2020;30:1291–305.

58. Kolmogorov M, Yuan J, Lin Y, Pevzner PA. Assembly of long, error-prone reads using repeat graphs. Nat Biotechnol. 2019;37:540–6.

59. Guan D, McCarthy SA, Wood J, Howe K, Wang Y, Durbin R. Identifying and removing haplotypic duplication in primary genome assemblies. Bioinformatics. 2020;36:2896–8.

60. Ruiz-Gartzia I, Lizano E, Marques-Bonet T, Kelley JL. Recovering the genomes hidden in museum wet collections. Mol Ecol Resour. 2022;22:2127–9.

61. Flynn JM, Hubley R, Goubert C, Rosen J, Clark AG, Feschotte C, et al. RepeatModeler2 for automated genomic discovery of transposable element families. Proc Natl Acad Sci U S A. 2020;117:9451–7.

62. Hoyt SJ, Storer JM, Hartley GA, Grady PGS, Gershman A, de Lima LG, et al. From telomere to telomere: The transcriptional and epigenetic state of human repeat elements. Science. 2022;376:eabk3112.

63. Karolchik D, Hinrichs AS, Furey TS, Roskin KM, Sugnet CW, Haussler D, et al. The UCSC Table Browser data retrieval tool. Nucleic Acids Res. 2004;32:D493–6.

64. Li H. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics. 2018;34:3094–100.

65. Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM [Internet].
arXiv [q-bio.GN]. 2013. Available from: http://arxiv.org/abs/1303.3997

66. Männer L, Schell T, Spies J, Galià-Camps C, Baranski D, Ben Hamadou A, et al. Chromosome-
level genome assembly of the sacoglossan sea slug Elysia timida (Risso, 1818) [Internet]. Genomics.
bioRxiv; 2024. Available from: https://www.biorxiv.org/content/10.1101/2024.06.04.597355v1

1019

1020

1021