EyeHex toolbox for complete segmentation of ommatidia in fruit fly eyes

Huy Tran[1,2*], Nathalie Dostatni[2], Ariane Ramaekers[2*]

[1] Faculty of Medicine and Health Technology, Tampere University, Tampere, Finland

[2] Laboratory of Nuclear Dynamics, UMR3664, Institut Curie, Sorbonne Université, CNRS, Paris, France

* Corresponding authors

E-mails: huy.tran@tuni.fi , ariane.ramaekers@curie.fr

Abstract

Variation in *Drosophila* compound eye size is studied across research fields, from evolutionary biology to biomedical studies, requiring the collection of large datasets to ensure robust statistical analyses. To address this, we present EyeHex, a tool for automatic segmentation of fruit fly compound eyes from brightfield and scanning electron microscopy (SEM) images. EyeHex features two integrated modules: the first utilizes machine learning to generate probability maps of the eye and ommatidia locations, while the second, a hard-coded module, leverages the hexagonal organization of the compound eye to map individual ommatidia. This iterative segmentation process, which adds one ommatidium at a time based on registered neighbors, ensures robustness to local perturbations. EyeHex also includes an analysis tool that calculates key metrics of the eye, such as ommatidia count and diameter distribution across the eye. With minimal user input for training and application, EyeHex achieves exceptional accuracy (>99.6% compared to manual counts on SEM images) and adapts to different fly strains, species, and image types. EyeHex offers a cost-effective, rapid, and flexible pipeline for extracting detailed statistical data on *Drosophila* compound eye variation, making it a valuable resource for high-throughput studies.

Keywords: *Drosophila*, Compound eye, Image segmentation, Ommatidia number, Ommatidia diameter, Acute zone, Morphological variation.

Background

Insect compound eyes consist of crystal-like lattices of elementary eyes, called facets or ommatidia. Each ommatidium is constituted of a fixed number of cells, including photoreceptors that sense light and transmit visual information to the brain, as well as a set of accessory cells (Nériec and Desplan, 2016; Wolff and Ready, 1993). Compound eye size varies greatly among insect groups and species due to differences in the number or diameter of

individual facets. In fruit flies (*Drosophila* genus), the number of ommatidia per eye ranges from approximately 600 to 1500, while ommatidia diameter varies from approximately 15 to 22 μm (Arif et al., 2013; Gaspar et al., 2019; Keesey et al., 2019; Posnien et al., 2012; Ramaekers et al., 2019; Reis et al., 2020a). Genetic variation is the primary driver of eye size variation among fruit flies, but environmental conditions such as food quality, larval density and temperature also play a role (Currea et al., 2018). On a wider phylogenetic range, the variation in eye size is much more spectacular. For example, the number of ommatidia varies from zero in some ant species for instance, up to approximately 20.000 in dragonflies (Land and Nilsson, 2012).

Variation in compound eye size is investigated from various scientific perspectives, ranging from evolutionary studies to biomedical research. Evolutionary ecologists aim to identify how eye size variation affects vision and visually guided behaviors, potentially resulting in adaptation to novel ecological niches. Research has shown that larger eyes account for improved vision in dim light or acuity (reviewed in (Land, 2005; Land and Nilsson, 2012; Warrant, 2008; Warrant and Nilsson, 1998). In contrast, reduced eye size is common in insects adapted to darkness, such as cave-dwelling beetles, which rely little or not at all on vision (Luo et al., 2019, 2018).

Variation in compound eye size, especially in fruit flies, is also of interest to developmental or evolutionary-developmental biologists who investigate the genetic and developmental origins of morphological variation (reviewed in (Casares and McGregor, 2020)). Furthermore, it serves as a genetic model for human eye conditions, such as aniridia or retinitis pigmentosa (reviewed in (Gaspar et al., 2019)) and a "test-tube" for studying various human pathologies (e.g. Zika virus pathogenicity (Harsh et al., 2020) or molecular determinants of spinal muscular atrophy (Maccallini et al., 2020)).

Until now, the development of larger scale studies has been hindered by the lack of a

simple, accurate, and cost-effective method to fully segment ommatidia. In most reports,

ommatidia were manually counted from Scanning Electron Microscopy (SEM) images with

limited throughput (Arif et al., 2013; Keesey et al., 2019; Posnien et al., 2012; Ramaekers et

al., 2019). Statistical analyses that requires larger datasets were performed using either

estimations of ommatidia numbers (Ramaekers et al., 2019) or measurements of eye surface

area or length instead of ommatidia counts (Arif et al., 2013; Harsh et al., 2020; Norry and

Gomez, 2017). In a recent report, researchers applied a commercial segmentation tool (Amira

v.2019.2, Thermo Fischer Scientific) to high-quality images obtained by X-ray tomography

(Gaspar et al., 2020). Brightfield imaging could be a desirable alternative to the more laborious

and expensive SEM or X-ray tomography technologies. However, due to the prominent

curvature of compound eyes, segmenting ommatidia from 2D images is non-trivial. Two

features present specific challenges. The first feature is the diverse degree of ocular hair (named

bristles) covering ommatidia in 2D eye images. Depending on their location on the eye surface,

the level of coverage varies from minimal (e.g., Fig. 1, black frames) to substantial (e.g., Fig.

1, green, blue and orange frames). The second feature is the dome-shaped transparent corneas,

which act as lenses overlaying the ommatidia. This causes the ommatidia to reflect light

differently depending on their orientation and the angle of the incoming light, hence the varying

levels of reflection observed in brightfield images (Fig. 1A). A recent study on eye size

variation in the *Drosophila virilis* group exploited this property, capturing and segmenting the

reflection of light by each ommatidia, rather than imaging the ommatidia themselves (Reis et

al., 2020a).

Another recent method, named Ommatidia Detection Algorithm (ODA) (Currea et al.,

2023), applies Fourier Transform to detect ommatidia on images of glue eye molds, light

microscopy and SEM images of various species of insects. This method detects the periodic

patterns of ommatidia in the compound eyes with filters in the frequency domain of the images.

However, when applied to 2D images, these two hard-coded methods are limited by the high

variability of the ommatidia imaging angles in distinct regions of the compound eye (Fig. 1),

particularly at the edges where ommatidia are imaged at markedly slanted angles. This may

result in the misdetection of ommatidia or the misidentification of ommatidia's centroid

locations.

In this study, we present EyeHex, a comprehensive toolbox for the automated

segmentation of ommatidia in *Drosophila* compound eyes. EyeHex consists of two modules

(Fig. 2): Module 1 uses a machine learning-based classifier (Trainable Weka Segmentation,

Figure 1. The many facets of the eye**.** Due to the convex structure of the compound eye, the aspect of ommatidia observed on 2D images varies greatly in different eye regions. Images shown here were acquired using (A) a macroscope (this study) and (B) by SEM (Ramaekers et al., 2019). The zoom-in regions are shown on the right side of each panel with the corresponding frame colors. Note the differences in reflection, shape and inter-ommatidial hair aspect.

ImageJ (Arganda-Carreras et al., 2017) to isolate the compound eye from the background and

delineate individual ommatidia. Users can customize training data, enabling broad

compatibility with diverse image types, from brightfield and SEM images, acquired using

various settings. Module 2 is hard-coded and leverages the hexagonal organization of

ommatidia to precisely map their location and quantify their number. This module requires

minimal user input, ensuring cost-effective and efficient characterization of *Drosophila* eye

size variations. The pipeline is coupled with an analytical tool that automatically generates

high-precision metrics, such as ommatidia counts along the eye anterior-posterior axis and

ommatidia diameter distribution across the eye.

Methods

Data acquisition

Drosophila strains and culture:

Fly stocks were cultured on standard cornmeal diet food at 25°C in density-controlled

conditions: batches of 10 females and 10 males were raised at 25°C and transferred into a fresh

vial every day. For each vial, eye measures were performed only on females eclosing during

the first two days of eclosion. Genotypes used in this study were: for _Drosophila melanogaster_:

Canton-S BH (Ramaekers et al., 2019); Hikone-AS (Kyoto DGRC 105668); DGRP-208

(WT25174; BDSC 25174); w;;;+/Df(4)J2 (abbreviated as WTJ2; Ramaekers et al., 2019); for

Drosophila pseudoobscura: Catalina Island, California, Isofemale line (National Drosophila

Species Stock Center, 14011-121.121).

Scanning electron microscopy (SEM) images:

SEM images are reused from (Ramaekers et al., 2019).

Light-microscopy images (Ramaekers et al., 2019 and this study):

Light macroscopy images were either reused from (Ramaekers et al., 2019; WTJ2) or

newly acquired (Hikone-AS). In the latter case, they were acquired from non-fixed, freshly cut

adult heads glued laterally on glass slides using a solvent-free all material glue (UHU® - cat

#64481). 3D image stacks (Z distance = 8 μm) were acquired at 400-600X magnification with

Figure 2. Overview of ommatidia segmentation steps using EyeHex toolbox. The process consists of two phases. The '*Training and Classification phase*' involves the generation of the training data with the assisted segmentation of a subset of ommatidia (module A) for the machine learning module, which converts raw input images into probability maps of the eye region and facet region (module B). In the '*Complete segmentation phase*', ommatidia are automatically and exhaustively segmented by expanding a hexagonal grid to fit the facet probability map within the eye region (module C). The user then validates and manually corrects this ommatidia grid using a provided graphical user interface (module D). The outputs include the labelled images for the ommatidia facet and boundary regions (top right), the ommatidia count and the location of each individual ommatidium (bottom right).

a Keyence digital macroscope VHX 2000 using optical zoom lens VH-Z20R/W. Single focused

2D images were generated using Stack Focuser plugin from ImageJ (Schindelin et al., 2012).

EyeHex workflow

The EyeHex toolbox protocol consists of two successive steps: (1) Training and

Classification, and (2) Segmentation (Fig. 2).

Automatic compound eye and ommatidia classification with machine learning

<u>Training data preparation</u>

This step is required when a new image type (e.g., SEM, brightfield microscope…) is

introduced. A first set of 2D sample images is used as training data for the classification of the

eye area (vs. non-eye area) and the ommatidia surfaces (vs. ommatidia boundaries) (Module

A, Fig. 2). Using the Graphical User Interface (GUI) included in the EyeHex toolbox, users are

invited to manually delineate the compound eye area and segment small groups of ommatidia

at different locations of the eye (e.g., center vs. edges), thus presenting different shapes,

reflections and bristle coverage (Fig. 1A). The machine learning module then uses this training

data to convert each raw input image into probability maps of the eye area and ommatidia

surfaces (module B, Fig. 2). Provided that the imaging settings are unperturbed, a single

training image is sufficient to achieve accurate classification.

<u>Automatic classification</u>

Classification is performed using the Fiji Weka plug-in (Arganda-Carreras et al., 2017)

through a provided macro. Training datasets are used to train two separate pixel-based Fast

Random Forest classifiers, each containing 100 decision trees (Breiman, 2001). The trained

classifier will then convert the focused 2D images into probability maps of the eye area and the

ommatidia surfaces. The compound eye region is defined by thresholding the eye probability

map with the Otsu algorithm (Otsu, 1979).

Automatic segmentation

EyeHex automatic segmentation takes advantage of the remarkably robust hexagonal

organization of the compound eye. During this step, ommatidia will be automatically and

comprehensively segmented by expanding a hexagonal grid to fit the ommatidia probability

map within the eye area (module C, Fig. 2). Users can validate and manually correct the

segmentation process for each image using the provided GUI (module D, Fig. 2). The output

of the segmentation consists of (1) a labeled image carrying the information on the location of

ommatidia surfaces and boundaries, (2) the count of ommatidia number and (3) data on each

ommatidium (2D position and coordinates on the hexagonal grid).

<u>The ommatidia hexagonal grid</u>

Figure 3. The ommatidia hexagonal grid. (A) The grid is seeded from three original ommatidia (purple). Each ommatidium is identified by a unique integer coordinate (x_{hex}, y_{hex}) on the hexagonal grid, alongside its continuous Cartesian coordinate (x_{2D}, y_{2D}) on the ommatidia probability map. Note that the hexagonal axes and the Cartesian axes are not necessarily parallel. (B) New ommatidia (red) are added individually to the grid by mirroring the seed ommatidia (purple): ommatidium 4 is spawned by mirroring ommatidium 1 through ommatidium 2 and 3; ommatidium 5 by mirroring ommatidium 2 through ommatidium 1 and 3 and ommatidium 6 by mirroring ommatidium 3 through ommatidium1 and 2. (C) New ommatidia (red solid circles) are segmented and added to the outer row of existing hexagonal grid set by 3 seed ommatidia (purple), shown here on top of the ommatidia probability map I_{prob} generated during classification phase.

EyeHex segmentation module maps each ommatidium to a unique coordinate on a

hexagonal grid. Thus, each ommatidium is characterized by two sets of coordinates: an integer

coordinate (x_{hex}, y_{hex}) on the hexagonal grid and a continuous Cartesian coordinate (x, y) on

the 2D ommatidia probability map (Fig. 3A). Note that each ommatidium of the hexagonal grid

shares edges with 6 neighbors, instead of 4 as in a normal orthogonal grid. The segmentation

process includes three steps:

Step 1: Setting the origin and orientation of the hexagonal grid:

The origin and orientation of the grid are established based on three original ommatidia

adjacent to one another (purple cells in Fig. 3A; with coordinates (x_{hex}, y_{hex}) equal to (0,0),

(1,0), (0,1)), selected by the user at the center of compound eye.

Step 2: Expansion of the hexagonal grid:

The hexagonal grid is expanded by adding individual ommatidia to the current grid. For

each three adjacent ommatidia (referred as omtd1, omtd2, omtd3 ; purple cells in Fig. 3B), we

have their respective coordinates on the hexagonal grid (x_{hex1}, y_{hex1}), (x_{hex2}, y_{hex2}),

(x_{hex3}, y_{hex3}) and on the Cartesian grid (x_1, y_1), (x_2, y_2), (x_3, y_3). Those coordinates are used

to predict the position of a fourth ommatidium, omtd4, which mirrors omtd1 through omtd2

and omtd3 (Fig. 3B) via Eq. 1-4:

$$x_{hex4} = x_{hex2} + x_{hex3} - x_{hex1}, \tag{1}$$

$$y_{hex4} = y_{hex2} + y_{hex3} - y_{hex1}, \tag{2}$$

$$x_4 = x_2 + x_3 - x_1, \tag{3}$$

$$y_4 = y_2 + y_3 - y_1. \tag{4}$$

177   Based on the ommatidia probability map, the program will find an ommatidium within

178   the vicinity of the mirrored position (x_4, y_4) and add omtd4 to the hexagonal grid at the

179   coordinates (x_{hex4}, y_{hex4}) (see below). Only the closest ommatidium within a distance of

180    $L_{grid}/2$ from the predicted position (x_4, y_4) is added to the grid. Here, L_{grid} is the average

181   length of the hexagonal ommatidial edges:

$$L_{grid} = \sum_{i,j=1..3} \frac{(x_i - x_j)^2 + (y_i - y_j)^2}{6}. \tag{5}$$

During the automatic hexagonal expansion (Fig. 3C), new ommatidia close to the

mirrored position that best fits the ommatidia facet probability map (see below) will always be

added to the perimeter of the grid, unless they are found outside the compound eye region (Fig.

3C).

<u>Fitting new ommatidia to the ommatidia probability map</u>

From the three adjacent ommatidia (omtd1, omtd2, omtd3), we look for the position of

omtd4 (x_4, y_4) based on the facet probability map (denoted as $I_{prob}(x, y)$) generated with the

machine learning module. The general strategy is "find Mickey pattern" with omtd2 and omtd3

being the "ear" and omtd4 being the "face". A template image I_{mouse} of size $m \times m$ (m in

pixel) is constructed with three filled identical circles of radius $0.225 \times m$, the centers of which

form a perfect angle with edge $0.5 \times m$ (Fig. 4A). The pattern is then smoothened with a 2D

gaussian filter with $\sigma = m/20$. We also set an "ignore" region (with NaN pixel value, dashed

blue in Fig. 4A) on I_{mouse} to avoid interference from surrounding ommatidia.

We called (x_2', y_2'), (x_3', y_3') and (x_4', y_4') the coordinates of the centers of the three

circles in the template image I_{mouse}. We find a linear transformation $[M]_B^{B'}$ from the coordinate

vectors B' to the coordinate vectors B, in which:

$$B' = [v'_a \quad v'_b] = \begin{bmatrix} x_4' - x_2' & x_4' - x_3' \\ y_4' - y_2' & y_4' - y_3' \end{bmatrix} \tag{6}$$

$$\text{and } B = [v_a \quad v_b] = \begin{bmatrix} x_4 - x_2 & x_4 - x_3 \\ y_4 - y_2 & y_4 - y_3 \end{bmatrix}. \tag{7}$$

198   For each non-NaN pixel in position (i, j) in I_{mouse}, we find the corresponding

199   coordinate $[M]_B^{B'}(i, j)$ on I_{prob} (Fig. 4B).

200   The objective function for the fit of the new omtd4 position $\Delta(x_4, y_4)$ is given by the

201   squared error between the facet probability map and the transformed mouse image:

$$\Delta(x_4, y_4) = \sum_{i,j=1..m} \left(I_{prob}([M]_B^{B'}(i, j)) - \theta \times I_{mouse}(i, j) \right)^2, \tag{8}$$

202    with θ being a free parameter to adjust for pixel brightness.

203    Due to frequent irregularities on the probability map, we introduce additional

Figure 4. '*Mickey patterns*' on the facet probability map. (A) The coordinate vectors $B' = (v'_a, v'_b)$ (left) for the mouse image I_{mouse} (right). The objective function does not consider the blue dashed regions (NaN value). (B) Example of the new coordinate vectors $B = (v_a, v_b)$ from ommatidium 2, ommatidium 3 and ommatidium 4 position (left) with the transformed I_{mouse} in this coordinate (right).

204 constraints to preserve the integrity of the hexagonal grid and account for the fact that the facet

205 orientation changes gradually along the grid: The objective function $\Delta(x_4, y_4)$ is set to infinity

206 when the new omtd4 position is farther than $0.3 \times L_{grid}$ from the predicted position that

207 mirrors omtd1, and when the omtd4 top view angle is too steep ($<20^o$) or too different from

208 that of omtd1.

209    **Manual correction**

210    Following the automatic hexagonal grid expansion, the user can manually remove or

211 add ommatidia using the provided GUI. After manual correction, newly added ommatidia are

212 mapped and automatically given their own coordinate in the existing hexagonal grid.

213    **Data analyses**

214    <u>Measurement of ommatidia size</u>

215    When multi-focused brightfield images are used, the z-coordinate of each ommatidium

216 can be determined from its position in the original image stack. We use this information to

217 estimate the size of an ommatidium by averaging its Euclidean distances in the three-

218 dimensional space to its adjacent ommatidia. These values are fitted to a polynomial surface to

generate a smooth profile of ommatidia altitude and spacing across the eye, as shown in Figure

5B and 5C.

Ommatidia column identification and column profiles

The ommatidia are arranged in columns that reflect the progressive specification of R8

ommatidia founder cells along the anterior-posterior axis of the eye primordium (eye-antennal

disc) (Ready et al., 1976). To locate these columns, the origin of the ommatidia hexagonal grid

(x_{hex}=0, y_{hex}=0) is shifted to the detected center-most ommatidia. The grid is then reoriented

so that both the x- and y-coordinates of the most anterior ommatidium in this grid take positive

values (Fig. 5A). The sum of the new xy-coordinates of the ommatidia (x_{hex}+y_{hex}) defines the

column index that corresponds to their birth order during R8 specification. This column index

increases from the posterior-dorsal to anterior-ventral regions of the eye (example in Fig. 5D).

From the reoriented grid, we calculate the "column profile" as the number of ommatidia per

column, which provides information about the eye geometry (example in Fig. 5E).

Averaging of morphological profiles

We accumulate measurements of individual ommatidia from different eyes based on

their unique coordinate in the hexagonal grid (x_{hex}, y_{hex}). This allows us to calculate the

average ommatidia spacing in different regions of the grid. For visualization, we project this

average spacing profile onto a reference eye, where the Cartesian coordinate for each

segmented ommatidia is known (example in Fig. 5D).

Outputs

After manual correction of segmentation and analysis of each eye, EyeHex

automatically exports the label image of the identified ommatidia and eye region. These label

images can be added to the training data to improve subsequent classifications. Also exported

is a comma-separated-values (.csv) file containing information about each ommatidium's

index, its Cartesian and hexagonal coordinates, and its birth order during R8 specification.

Results

The EyeHex toolbox

The EyeHex toolbox is a set of MATLAB scripts that allows users to segment all visible

ommatidia from 2D images of *Drosophila* compound eyes in a semi-automatic manner.

Limited user attention through Graphical User Interfaces (GUIs) is required during the training

and manual correction step. EyeHex output includes labeled eye images depicting the

ommatidia masks (Fig. 2E), as well as ommatidia counts and positions in .csv format (Fig. 2F)

for further statistical analyses. The toolbox, its manual and sample images are available for

download on GitHub at https://github.com/huytran216/EyeHex-toolbox.

Segmentation on SEM images

The gold standard for quantifying ommatidia numbers relies on manual counts performed on

2D compound eye images acquired via scanning electron microscopy (SEM). To evaluate the

efficiency of EyeHex segmentation, we compared the number of ommatidia generated by

EyeHex toolbox with manual counts from previously published SEM datasets (Ramaekers et

al., 2019) for three *D. melanogaster* strains : Hikone-AS (Fig. S1), Canton-S BH (Fig. S2), and

DGRP-208 (Fig. S3). For training datasets, we manually segmented approximately 80

ommatidia (~ 10% of the total per eye) in a single SEM image. Subsequent steps included

classification, automatic segmentation, manual corrections, and counting (Table 1; Suppl. File

2). Compared to manual counts, EyeHex ommatidia numbers deviated by 0 to a maximum of

8 ommatidia per eye, achieving an average accuracy exceeding 99.6 % (Table 1, Suppl. File

2). We further tested EyeHex to SEM images of *D. pseudoobscura*, a different *Drosophila*

species with significantly larger compound eyes (*D. pseudoobscura*: ~1200 ommatidia/eye *vs*

D. melanogaster: ~ 800 ommatidia/eye), and compared the EyeHex results to published manual

counts (Ramaekers et al., 2019). For this species, EyeHex also yielded highly accurate counts,

with differences ranging from 0 to 3 ommatidia per eye (99.9 % accuracy; Table 1, Fig. S4,

Suppl. File 2). These results demonstrate our tool's applicability beyond *D. melanogaster* to

other *Drosophila* species.

Segmentation on brightfield images

The sample preparation and image acquisition procedure for SEM images are relatively

time-consuming and expensive. To reduce time and costs, we applied EyeHex on brightfield

images acquired using a macroscope on non-fixed samples. Our previous work estimated

ommatidia numbers in such images by assuming a perfect elliptical fit for the ommatidia

hexagonal grid (see Ramaekers et al., 2019 for the method), but direct manual counts were not

performed. To assess whether EyeHex achieves accuracy comparable to SEM with brightfield

images, we generated training datasets by manually segmenting 150 ommatidia from two

newly acquired Hikone-AS brightfield images. We then applied EyeHex for classification,

segmentation and counting (Fig. 5A-B and Table 1, Suppl. File 2). Notably, the Hikone-AS

strain used here may differ from that in our previous study (Ramaekers et al., 2019), possibly

because of genetic drift or slightly different culture conditions, as experiments were conducted

in different laboratories over 5 years apart. Despite this, EyeHex-derived counts closely

matched previous manual SEM counts (Table 1), indicating that brightfield imaging combined

with EyeHex provides a reliable representation of biological samples. In contrast, our earlier

elliptical estimation method systematically overestimated the number of ommatidia by 20 to

40 ommatidia on average (Ramaekers et al. 2019 and Table 1). We also tested EyeHex on *white*

mutant eyes (strain WTJ2), as this is one of the most commonly used genetic markers in fruit

flies. Those eyes lack the red pigmentation and exhibit reduced contrast. While ommatidia

segmentation accuracy remained comparable to other datasets, more manual correction was

required - up to 14% of the total ommatidia (Table 1, Suppl. File 2). This primarily involved

falsely detected ommatidia at the eye boundary, caused by overestimation of the eye region

during the classification step. However, ommatidia segmentation performed equally well

compared to the other datasets, confirming EyeHex's adaptability to low-contrast samples.

Table 1: Ommatidia counts on SEM and brightfield images.

Strain	Image type	Image collection	n	EyeHex	EyeHex	Manual	Ellipse
				Pre-correction	*Post-correction*		
Hikone-AS	SEM	*Ramaekers et al. 2019*	6	722.7 ± 26.2	741.2 ± 21.8	739.7 ± 21.8*	764.5 ± 26.7
Canton-S[BH]	SEM	*Ramaekers et al. 2019*	9	948.3 ± 31.4	857.8 ± 30.1	859.3 ± 27.4*	888.2 ± 33.1
DGRP-208	SEM	*Ramaekers et al. 2019*	8	859.0 ± 82.1	754.5 ± 44.1	754.6 ± 43.4*	797.8 ± 48.1
D. pseudoobscura Catalina Island	SEM	*Ramaekers et al. 2019*	5	1192.8 ± 14.6	1176.2 ± 22.2	1175.0 ± 22.1*	1216.9 ± 22.0
Hikone-AS	Brightfield	This study	12	787.6 ± 18.2	752.7 ± 15.6	/	778.7 ± 18.3
WTJ2 (white eyes)	Brightfield	*Ramaekers et al. 2019*	8	870.3 ± 73.3	789.9 ± 27.2	/	819.6 ± 28.7*

Ommatidia counts (mean ± standard deviation) for SEM or brightfield images. The images were either reused from a previous study (Ramaekers et al., 2019) or newly collected. We compare ommatidia counts performed using EyeHex segmentation before and after manual correction, together with fully manual counts ('Manual') or elliptic estimations ('Ellipse'). All strains are from D. melanogaster *species, except where otherwise indicated. n: sample size; *: counts from Ramaekers et al., 2019.*

Ommatidia size profile and ommatidia column count

Insect compound eyes often exhibit local specializations – named acute or bright zones

– that enhance acuity or signal-to-noise ratio in specific regions of the visual field (Horridge,

1978) (Land, 1997). The presence and location of these acute/bright zones correlate with

distinct visually guided behaviors. For example, upward-pointing zones are frequently

associated to predatory behavior, whereas male-specific forward-pointing zones are linked to

sexual pursuit. In many insect species, including flies, these zones are characterized by

enlarged ommatidia. In addition to quantifying ommatidia number, EyeHex can automatically

detect these regions of enlarged ommatidia by analyzing ommatidia size variation across the

eye. By analyzing brightfield multifocus image stacks, EyeHex can reconstruct the 3D position

of each ommatidium (Fig. 5C). This, in turn, enables the calculation of the average distance in

3D space between each ommatidium and its 6 adjacent neighbors (see Methods), providing a

proxy for ommatidial diameter across the eye. In all eyes analyzed (e.g., Fig. 5D), ommatidial

density and diameter varied by up to 20 % along the posterior-dorsal to anterior-ventral axis of

the eye, consistent with observations by (Gaspar et al., 2020) and (Buffry et al., 2024) (Fig. 5D

and Fig. S7). Projecting density maps from multiple specimens onto a common reference

revealed a consistent zone of enlarged ommatidia in the anterior-ventral part of the eye (black

crosses in Fig. 5D and Fig. S7). In *Drosophila*, acute zones have been documented in males,

where they occupy an antero-dorsal position. These zones are reportedly absent in females,

though the existence and location of analogous structures in female eyes remain unclear. Since

our study focused on female eyes, further investigation is required to determine whether the

observed antero-ventral ommatidial enlargement represents an acute zone in this sex. Notably,

at the extreme edge of the eye, ommatidia diameters measured by EyeHex were unexpectedly

small (<12 µm). In these regions, some ommatidia adopt an elongated, non-hexagonal shape,

and the diameters reported by EyeHex reflect their reduced width. However, identifying the

xy- and z-coordinates of highly slanted ommatidia can be challenging, making the

determination of inter-ommatidial spacing at the very edge of the eye prone to errors. Thus,

such small diameter may not always fully represent true morphological dimensions.

Figure 5. EyeHex output: Quantitative analysis of compound eye features. A) Focused brightfield image of a Hikone-AS eye, with the overlayed hexagonal grid axes. **B)** Detected ommatidia centers, sorted into columns (separated by color). Ommatidia in the most posterior (rightmost) column are highlighted in bold. **C)** Contour map of ommatidia altitude (in μm, calculated relative to the lowest focal plane). **D)** Contour map of the average spacing between adjacent ommatidia (in μm) projected onto the eye in panel A. Black crosses indicate the locations of greatest inter-ommatidial distance (largest ommatidia) for individual Hikone-AS eyes; data from *n*=12 eyes. **E)** Number of ommatidia in each column from posterior (P) to anterior (A) (line and shading represent the mean and standard deviation, respectively, n=12). **F)** Comparison of mean ommatidia count profiles between left (blue curve, n =8) eyes and right (red curve, n=4) eyes. Scale bars in C and D: 200 μm. Eye orientation is shown in panel A.

In the compound eye, ommatidia are arranged in regularly spaced columns along the

anterior-posterior (A-P) axis of the eye (Fig. 5B). This organization reflects the progressive

determination and spacing of ommatidia founder cells (termed R8 cells) along the A-P axis of

the eye primordium during the progression of a differentiation wave named '*morphogenetic*

furrow' (Ready et al., 1976) (Wolff and Ready, 1993) (for a recent review, see (Warren and

Kumar, 2023)). Thus, quantifying the number of A-P columns in adult eyes provides a read-

out of the R8 determination process and of its robustness. Here, we used EyeHex to extract this

information from a set of brightfield eye images from 12 individuals of the Hikone-AS wild-

type genetic background. Each eye consisted of 33.89 ± 1.05 columns (mean and standard deviation). We also extracted '*column profiles*' by plotting the number of ommatidia per column along the A-P axis of the eye (Fig. 5B), thereby providing a simple representation of the eye geometry. We then aligned the 12 column profiles for comparison. The mean aligned column profile from 12 brightfield eye images is shown in Figure 5E. Interestingly, the standard deviation (shaded area of the plot, Fig. 5E) of the mean is on average ~1.08 ommatidia, with a maximum of 3.73 ommatidia per column. Therefore, the highly inbred Hikone-AS laboratory individuals show very little variation in eye organization and size, indicating that despite its complexity, compound eye development is remarkably robust. Comparison of the mean aligned column profiles between left and right eyes from separate individuals also did not reveal any significant bias (an average difference of 0.45 ommatidia per column) (Fig. 5F).

Discussion

The EyeHex toolbox automates segmentation and mapping in 2D space of ommatidia in *Drosophila* compound eyes. Coupled with a dedicated analysis pipeline, it quantifies metrics of the eye organization, including total ommatidia number, ommatidia count profiles along the A-P axis, and contour maps of ommatidial diameters. Using a machine learning-based approach, EyeHex adapts to diverse image types, with performance improving as datasets expand. Existing tools for automated ommatidial segmentation vary in their methodologies and limitations. Reis *et al.* (Reis et al., 2020b) used brightfield images to segment ommatidia reflections as discrete cells separated by a predefined minimum distance – a strategy prone to false negatives in case of damaged or non-reflective ommatidia. In Gaspar *et al.* (Gaspar et al., 2020), ommatidia segmentation was hard-coded and restricted to 3D X-ray tomography images. The segmentation method developed by Currea *et al.* (Currea et al., 2023) is applicable to various image types and exploits periodic ommatidia patterns via filtering in the frequency domain (Fourier transforms), fitting hexagonal/orthogonal grids to overlapping regions. While

efficient, this semi-global approach may struggle with extreme warping, common at the compound eye boundary. By contrast, EyeHex employs an adaptive, local strategy: ommatidia are identified sequentially and integrated into the existing eye grid based on their three nearest neighbors. This accommodates local distortions in the compound eye surface, including warping at the eye edges, while preserving global hexagonal organization. In addition, because EyeHex can accommodate non-uniform ommatidia spacing, the segmentation process should not be affected by minor disruptions or irregularities in the eye lattice found in some mutants. Similarly, EyeHex is also effective for quantifying eye morphology in non-*melanogaster* *Drosophila* species, which share conserved eye organization, as shown in this study for large-eyed *D. pseudoobscura*. It is also adaptable to images exhibiting lower contrast, such as *white* mutant eyes. However, mutant eyes with severe structural abnormalities may not be accurately segmented by EyeHex. Similarly, highly bulged eyes (obscuring edges in brightfield image stacks) may require alternative imaging techniques such as μCT scans (Gaspar et al., 2020) or indirect imaging of eye molds flattened by incision (Ramirez-Esquivel et al., 2017).

Manual counts of ommatidia from SEM images ommatidia remain the gold standard for ommatidia enumeration. EyeHex achieves > 99.6 % concordance with this gold standard after minimal manual corrections. Applied to macroscope brightfield images of unfixed samples, it replicates SEM-derived statistics, maintaining high accuracy while streamlining sample preparation and processing. Nevertheless, fully automated segmentation methods cannot achieve 100% accuracy – a critical consideration given interindividual variation in ommatidial counts (relative standard deviation < 2% in inbred lines; this study, (Currea et al., 2018; Gaspar et al., 2020). Thus, user-guided inspection and manual correction, integrated into EyeHex toolbox, remains essential.

Compound eyes, though widespread across the animal kingdom, exhibit lower acuity than other eye types, such as the vertebrate camera eyes (Kirschfeld, 1976; Land, 1997; Mallock, 1894).

To achieve human-level resolution, compound eyes should consist of millions of ommatidia,

reaching a radius of 6 meters. Many species compensate via specialized zones ("acute" or

"bright" zones), often characterized by enlarged ommatidia, which enhance acuity for specific

parts of the visual field (Land, 1997). Despite the adaptive relevance of this trait, most

Drosophila studies report only average or local measures of ommatidial diameter (Gaspar et

al., 2020; Keesey et al., 2019; Posnien et al., 2012; Ramaekers et al., 2019). Exceptions include

a recent study applying a 3D version of the ODA segmentation tool (Currea et al., 2023) to

synchrotron microtomography images (Buffry et al., 2024). Similarly, EyeHex derives 3D

altitude data from brightfield multifocus stacks to map ommatidial diameter across the eye

surface, enabling identification of potential acute zones or subtle patterns of variation.

Variation in *Drosophila* compound eye size is a broadly used model in evolutionary

developmental biology (Casares and McGregor, 2020). Despite well-documented eye size

variation, technical limitations have historically limited large-scale studies, precluding the

compound eye's use in addressing fundamental evo-devo questions. For instance, quantifying

subtle morphological variation – a critical metric for assessing developmental precision,

stochasticity, or robustness – demands expansive datasets. This bottleneck has relegated studies

of phenomena like fluctuating asymmetry in *Drosophila* to simpler traits (e.g., wing

morphology and mechanosensory hair patterns, (Debat et al., 2009; Polak and Starmer, 2001),

which are more readily quantified. The compound eye, with its extensively characterized

development, offers an ideal system to explore these questions in a complex tissue, uniquely

positioned to link developmental mechanisms with phenotypic outcome. EyeHex bridges this

methodological gap by enabling providing accurate segmentation and extraction of eye

morphology statistics across imaging platforms, including accessible brightfield macroscopy.

Our pipeline - from imaging unfixed heads to EyeHex analysis – offers a rapid, cost-effective

solution for large-scale quantification of eye morphology.

List of abbreviations

A-P: Anterior-Posterior

μCT: Micro Computed Tomography

SEM: Scanning Electron Microscopy

GUI: Graphical User Interface

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

EyeHex toolbox: code and user manual available to download from https://github.com/huytran216/EyeHex-toolbox.

Examples of segmentation results and analysis for SEM images (Hikone-AS, Canton-S, DGRP-208, *Drosophila pseudoobscura*) and brightfield images (Hikone-AS, WTJ2): Supplementary_file1.pdf

Ommatidia counts: Supplementary_file2.xlsx

Competing interests

The authors declare that they have no competing interests.

Author's contributions

AR collected the data (sample preparation and imaging), HT created EyeHex toolbox and analyzed the data, AR and HT interpreted the data. AR, HT and ND drafted the manuscript.

Acknowledgments

The authors want to thank Virginie Courtier (CNRS, Institut Jacques Monod, Paris) for

providing access to the Keyence macroscope.

Fundings

HT is supported by Research Council of Finland (354987), Tampere University's

Health Data Science Profile. This work was also supported by "Emergence" grant from

Sorbonne Université to AR.

References

Arganda-Carreras, I., Kaynig, V., Rueden, C., Eliceiri, K.W., Schindelin, J., Cardona, A.,
Sebastian Seung, H., 2017. Trainable Weka Segmentation: a machine learning tool for
microscopy pixel classification. Bioinformatics 33, 2424–2426.
https://doi.org/10.1093/bioinformatics/btx180
Arif, S., Hilbrant, M., Hopfen, C., Almudi, I., Nunes, M.D.S., Posnien, N., Kuncheria, L.,
Tanaka, K., Mitteroecker, P., Schlötterer, C., Mcgregor, A.P., 2013. Genetic and
developmental analysis of differences in eye and face morphology between Drosophila
simulans and Drosophila mauritiana. Evolution and Development 15, 257–267.
https://doi.org/10.1111/ede.12027
Breiman, L., 2001. Random Forests. Machine Learning 45, 5–32.
https://doi.org/10.1023/A:1010933404324
Buffry, A.D., Currea, J.P., Franke-Gerth, F.A., Palavalli-Nettimi, R., Bodey, A.J., Rau, C.,
Samadi, N., Gstöhl, S.J., Schlepütz, C.M., McGregor, A.P., Sumner-Rooney, L.,
Theobald, J., Kittelmann, M., 2024. Evolution of compound eye morphology underlies
differences in vision between closely related Drosophila species. BMC Biol 22, 1–15.
https://doi.org/10.1186/s12915-024-01864-7
Casares, F., McGregor, A.P., 2020. The evolution and development of eye size in flies. Wiley
Interdisciplinary Reviews: Developmental Biology. https://doi.org/10.1002/wdev.380
Currea, J.P., Smith, J.L., Theobald, J.C., 2018. Small fruit flies sacrifice temporal acuity to
maintain contrast sensitivity. Vision Research 149, 1–8.
https://doi.org/10.1016/j.visres.2018.05.007
Currea, J.P., Sondhi, Y., Kawahara, A.Y., Theobald, J., 2023. Measuring compound eye optics
with microscope and microCT images. Commun Biol 6, 1–12.
https://doi.org/10.1038/s42003-023-04575-x
Debat, V., Debelle, A., Dworkin, I., 2009. PLASTICITY, CANALIZATION, AND
DEVELOPMENTAL STABILITY OF THE DROSOPHILA WING: JOINT
EFFECTS OF MUTATIONS AND DEVELOPMENTAL TEMPERATURE.
Evolution 63, 2864–2876. https://doi.org/10.1111/j.1558-5646.2009.00774.x
Gaspar, P., Almudi, I., Nunes, M.D.S., McGregor, A.P., 2019. Human eye conditions: insights
from the fly eye. Human Genetics 138, 973–991. https://doi.org/10.1007/s00439-018-
1948-2

Gaspar, P., Arif, S., Sumner-Rooney, L., Kittelmann, M., Bodey, A.J., Stern, D.L., Nunes, M.D.S., McGregor, A.P., 2020. Characterization of the genetic architecture underlying eye size variation within drosophila melanogaster and drosophila simulans. G3: Genes, Genomes, Genetics 10, 1005–1018. https://doi.org/10.1534/g3.119.400877

Harsh, S., Fu, Y., Kenney, E., Han, Z., Eleftherianos, I., 2020. Zika virus non-structural protein NS4A restricts eye growth in *Drosophila* through regulation of JAK/STAT signaling. Disease Models & Mechanisms dmm.040816. https://doi.org/10.1242/dmm.040816

Horridge, G.A., 1978. The separation of visual axes in apposition compound eyes. Philosophical transactions of the Royal Society of London. Series B, Biological sciences 285, 1–59. https://doi.org/10.1098/rstb.1978.0093

Keesey, I.W., Grabe, V., Gruber, L., Koerte, S., Obiero, G.F., Bolton, G., Khallaf, M.A., Kunert, G., Lavista-Llanos, S., Valenzano, D.R., Rybak, J., Barrett, B.A., Knaden, M., Hansson, B.S., 2019. Inverse resource allocation between vision and olfaction across the genus Drosophila. Nature Communications 10, 1–16. https://doi.org/10.1038/s41467-019-09087-z

Kirschfeld, K., 1976. The Resolution of Lens and Compound Eyes, in: Zettler, F., Weiler, R. (Eds.), Neural Principles in Vision, Proceedings in Life Sciences. Springer, Berlin, Heidelberg, pp. 354–370. https://doi.org/10.1007/978-3-642-66432-8_19

Land, M.F., 2005. The optical structures of animal eyes. Current Biology 15, R319–R323. https://doi.org/10.1016/j.cub.2005.04.041

Land, M.F., 1997. VISUAL ACUITY IN INSECTS. Annu. Rev. Entomol. 42, 147–177. https://doi.org/10.1146/annurev.ento.42.1.147

Land, M.F., Nilsson, D.-E., 2012. Animal Eyes, Oxford Animal Biology Series. Oxford University Press, Oxford.

Luo, X.Z., Antunes-Carvalho, C., Wipfler, B., Ribera, I., Beutel, R.G., 2019. The cephalic morphology of the troglobiontic cholevine species Troglocharinus ferreri (Coleoptera, Leiodidae). Journal of Morphology 280, 1207–1221. https://doi.org/10.1002/jmor.21025

Luo, X.-Z., Wipfler, B., Ribera, I., Liang, H.-B., Tian, M.-Y., Ge, S.-Q., Beutel, R.G., 2018. The cephalic morphology of free-living and cave-dwelling species of trechine ground beetles from China (Coleoptera, Carabidae). Org Divers Evol 18, 125–142. https://doi.org/10.1007/s13127-017-0351-5

Maccallini, P., Bavasso, F., Scatolini, L., Bucciarelli, E., Noviello, G., Lisi, V., Palumbo, V., D'Angeli, S., Cacchione, S., Cenci, G., Ciapponi, L., Wakefield, J.G., Gatti, M., Raffa, G.D., 2020. Intimate functional interactions between TGS1 and the Smn complex revealed by an analysis of the Drosophila eye development. PLoS Genet 16, e1008815. https://doi.org/10.1371/journal.pgen.1008815

Mallock, A., 1894. I. Insect sight and the defining power of composite eyes. Proceedings of the Royal Society of London 55, 85–90. https://doi.org/10.1098/rspl.1894.0016

Nériec, N., Desplan, C., 2016. From The Eye To The Brain: Development Of The Drosophila Visual System. Curr Top Dev Biol 116, 247–271. https://doi.org/10.1016/bs.ctdb.2015.11.032

Norry, F.M., Gomez, F.H., 2017. Quantitative trait loci and antagonistic associations for two developmentally related traits in the drosophila head. Journal of Insect Science 17, 19. https://doi.org/10.1093/jisesa/iew115

Otsu, N., 1979. A threshold selection method from gray-level histograms. IEEE Transaction on System, Man and Cybernetics C, 62–66. https://doi.org/10.1109/TSMC.1979.4310076

Polak, M., Starmer, W.T., 2001. THE QUANTITATIVE GENETICS OF FLUCTUATING
ASYMMETRY. evol 55, 498–511. https://doi.org/10.1554/0014-
3820(2001)055[0498:TQGOFA]2.0.CO;2
Posnien, N., Hopfen, C., Hilbrant, M., Ramos-Womack, M., Murat, S., Schönauer, A., Herbert,
S.L., Nunes, M.D.S., Arif, S., Breuker, C.J., Schlötterer, C., Mitteroecker, P.,
McGregor, A.P., 2012. Evolution of eye morphology and Rhodopsin expression in the
Drosophila melanogaster species subgroup. PLoS ONE 7, 1–11.
https://doi.org/10.1371/journal.pone.0037346
Ramaekers, A., Claeys, A., Kapun, M., Mouchel-Vielh, E., Potier, D., Weinberger, S.,
Grillenzoni, N., Dardalhon-Cuménal, D., Yan, J., Wolf, R., Flatt, T., Buchner, E.,
Hassan, B.A., 2019. Altering the Temporal Regulation of One Transcription Factor
Drives Evolutionary Trade-Offs between Head Sensory Organs. Developmental Cell
780–792. https://doi.org/10.1016/j.devcel.2019.07.027
Ramirez-Esquivel, F., Ribi, W.A., Narendra, A., 2017. Techniques for Investigating the
Anatomy of the Ant Visual System. J Vis Exp 56339. https://doi.org/10.3791/56339
Ready, D.F., Hanson, T.E., Benzer, S., 1976. Development of the Drosophila retina, a
neurocrystalline lattice. Developmental Biology 53, 217–240.
https://doi.org/10.1016/0012-1606(76)90225-6
Reis, M., Wiegleb, G., Claude, J., Lata, R., Horchler, B., Ha, N.-T., Reimer, C., Vieira, C.P.,
Vieira, J., Posnien, N., 2020a. Multiple loci linked to inversions are associated with eye
size variation in species of the Drosophila virilis phylad. Scientific Reports 10, 12832.
https://doi.org/10.1038/s41598-020-69719-z
Reis, M., Wiegleb, G., Claude, J., Lata, R., Horchler, B., Ha, N.-T., Reimer, C., Vieira, C.P.,
Vieira, J., Posnien, N., 2020b. Multiple loci linked to inversions are associated with eye
size variation in species of the Drosophila virilis phylad. Sci Rep 10, 12832.
https://doi.org/10.1038/s41598-020-69719-z
Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch,
S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D.J., Hartenstein, V.,
Eliceiri, K., Tomancak, P., Cardona, A., 2012. Fiji: an open-source platform for
biological-image analysis. Nature Methods 9, 676–682.
https://doi.org/10.1038/nmeth.2019
Warrant, E.J., 2008. Seeing in the dark: Vision and visual behaviour in nocturnal bees and
wasps. Journal of Experimental Biology 211, 1737–1746.
https://doi.org/10.1242/jeb.015396
Warrant, E.J., Nilsson, D.E., 1998. Absorption of white light in photoreceptors. Vision Res 38,
195–207. https://doi.org/10.1016/s0042-6989(97)00151-x
Warren, J., Kumar, J.P., 2023. Patterning of the Drosophila retina by the morphogenetic
furrow. Frontiers in Cell and Developmental Biology 11.
Wolff, T., Ready, D.F., 1993. Pattern formation in the Drosophila retina, in: The Development
of Drosophila Melanogaster. Cold Spring Harbor Laboratory Press, pp. 1277–1325.
556