Research Advance

N-acetylation of α-synuclein enhances synaptic vesicle clustering mediated by

α-synuclein and lysophosphatidylcholine

Chuchu Wang[1,2]#, Chunyu Zhao[1,2]#, Xiao Hu[3], Jiali Qiang[1,2], Zhenying Liu[1,2], Jinge Gu[1,2], Shengnan
Zhang[1,2], Dan Li[4,5], Yaoyang Zhang[1], Jacqueline Burré[6], Jiajie Diao[3]*, and Cong Liu[1,7]*,

1. Interdisciplinary Research Center on Biology and Chemistry, Shanghai Institute of Organic Chemistry,

Chinese Academy of Sciences, Shanghai 201210, China.

2. University of Chinese Academy of Sciences, Beijing 100049, China.

3. Department of Cancer Biology, University of Cincinnati College of Medicine, Cincinnati, OH 45267,

USA.

4.Bio-X Institutes, Key Laboratory for the Genetics of Developmental and Neuropsychiatric Disorders

(Ministry of Education), Shanghai Jiao Tong University, Shanghai 200030, China;

5. Zhangjiang Institute for Advanced Study, Shanghai Jiao Tong University, Shanghai 200240, China;

6. Brain and Mind Research Institute & Appel Institute for Alzheimer's Disease Research, Weill Cornell

Medicine, New York, NY 10021, USA.

7. State Key Laboratory of Chemical Biology, Shanghai Institute of Organic Chemistry, Chinese

Academy of Sciences, Shanghai, 200032, China

#These authors contributed equally to this work.

*To whom correspondence should be addressed. E-mails: jiajie.diao@uc.edu (J.D.) and

liulab@sioc.ac.cn (C.L.)

Abstract

Post-translational modifications (PTMs) of α-synuclein (α-syn) such as acetylation and phosphorylation

play important yet distinct roles in regulating α-syn conformation, membrane binding, and amyloid

aggregation. However, how PTMs regulate α-syn function in presynaptic terminals remains unclear.

Previously, we reported that α-syn clusters synaptic vesicles (SV) [1], and neutral phospholipid

lysophosphatidylcholine (LPC) can mediate this clustering [2]. Here, based on our previous findings, we

further demonstrate that N-terminal acetylation, which occurs under physiological condition and is

irreversible in mammalian cells, significantly enhances the functional activity of α-syn in clustering SVs.

Mechanistic studies reveal that this enhancement is caused by the N-acetylation-promoted insertion of

α-syn's N-terminus and increased intermolecular interactions on the LPC-containing membrane. Our

work demonstrates that N-acetylation fine-tunes α-syn–LPC interaction for mediating α-syn's function

in SV clustering.

Key words: N-acetylation, α-synuclein, vesicle clustering

Introduction

α-Syn is a highly abundant presynaptic protein in neuron. It is involved in synaptic vesicle (SV) clustering and the assembly of soluble N-ethylmaleimide sensitive factor receptor (SNARE) complex for mediating SV trafficking and neurotransmitter release [1,3]. Abnormal amyloid aggregation of α-syn and deposition into Lewy Bodies (LBs) is a pathological hallmark of Parkinson's disease [4,5]. Different types of post-translational modifications (PTMs), e.g. acetylation, ubiquitination, and phosphorylation have been identified to modify α-syn under physiological and pathological conditions [5-9]. Phosphorylation of S129 and Y39 is elevated, and this PTM-modified α-syn accumulates in LBs, implying a direct relationship between PTMs and α-syn pathology [7,8]. Rather than disease-related PTMs, additional PTMs e.g. N-acetylation and O-GlcNAcylation of α-syn were found also under normal condition [10,11]. Specifically, N-acetylation can influence the conformation and increase the helicity of the N-terminal area of α-syn, alters its membrane-binding behavior, and is critical for oligomer formation and amyloid aggregation kinetics [12-14]. However, it remains unknown whether N-acetylation directly regulates the physiological function of α-syn in clustering SVs.

Results

N-terminal acetylation enhances SV clustering induced by α-syn.

In this study, we sought to investigate whether N-acetylation could modulate α-syn's function in promoting SV clustering. Firstly, we prepared recombinant N-terminally acetylated α-syn (Ac-α-syn) as well as unmodified α-syn (un-α-syn). N-terminal acetylation leads to a significant chemical shift perturbation of the first nine residues and H50 of α-syn in nuclear magnetic resonance (NMR) spectrum (**Figure 1a, left**). Intriguingly, in-cell NMR experiments show that all of the un-α-syn is N-terminally acetylated once it is transfected into HEK-293T cells by electroporation (**Figure 1a, right**), which is in consistent with a previous report [10]. This result confirms the physiological relevance of N-terminal acetylation of α-syn. Notably, N-terminal acetylation is believed to be irreversible in mammalian cells due to lack of a known N-terminal de-acetyltransferase. Hence, it is crucial to investigate the impact of N-acetylation on the function of α-syn. To further probe the influence of N-acetylation on α-syn-mediated SV clustering, we incubated mouse SVs with various α-synuclein variants (**Figure 1b, upper**). We next measured the extent of SV clustering by utilizing dynamic light scattering (DLS) to monitor alterations in particle diameters, as well as using negatively stained transmission electron microscopy (TEM) (**Figure 1b, middle and lower**). Notably, N-acetylation significantly amplified

α-syn's capacity to cluster SVs, which requires the N-terminal 30 residues (**Figure 1c-d**). These findings

imply that N-acetylation augments SV clustering.

LPC mediates the enhancement of vesicle clustering by N-terminal acetylation of α-syn.

In our previous work, we found that α-syn can bind to both negatively charged dioleoyl-phosphoserine

(DOPS) [1] and neutral lysophosphatidylcholine (LPC) [2]. Building on this, we sought to examine how

N-acetylation affects SV clustering that arises from the interaction between α-syn and either of these

lipids. To assess clustering functionality with specific lipid compositions, we synthesized liposomes

mimicking SVs that contained either LPC or DOPS. We then used a single-vesicle clustering assay to

track the clustering mediated by Ac-α-syn and un-α-syn (**Figure 2a**). Importantly, we observed that

N-acetylation considerably boosted the clustering ability of α-syn with LPC-containing liposomes

(**Figure 2b**). In contrast, N-acetylation did not increase the clustering of liposomes containing DOPS

(**Figure 2c**). Therefore, LPC is responsible for the increased SV clustering activity by N-terminally

acetylated α-syn.

N-terminal acetylation increases the α-syn–LPC interaction.

The interaction between α-syn and lipids is crucial for clustering SVs [1,2,15]. To elucidate the structural

basis of how N-acetylation impacts the α-syn−LPC interaction, we employed NMR measurements and

coarse-grained dissipative particle dynamics (DPD) simulations. First, we carried out solution NMR

titration analyses to examine the α-syn−LPC interaction with residue-specific resolution. Adding

increasing concentrations of pure LPC micelles to unmodified α-syn resulted in a dose-dependent and

continuous attenuation of signals for the first 100 residues of α-syn, indicating these residues can

interact with LPC (**Figure 3a, upper**). Notably, N-acetylation significantly elevated the binding affinity

of the N-terminal segment (residues 1-30), particularly the initial 10 residues, towards LPC, leading to a

rapid binding pattern during titration (**Figure 3a, lower**). Moreover, Ac-α-syn demonstrated a sequential

binding behavior with LPC, where the N-terminal region (residues 1-30) bound initially, succeeded by

the central region (residues 30-100) as the LPC/Ac-α-syn ratio increased (**Figure 3a, lower**). Similar

trends were observed in NMR titrations using DOPC/LPC-mixed liposomes (**Figure 3b**). On the other

hand, N-acetylation only slightly weakened the interaction between the N-terminal region (residues

1-30) and DOPS, particularly at low DOPS/α-syn ratios, resulting in a continuous signal attenuation for

the first 100 residues (**Figure 3c**). These findings reveal that N-acetylation considerably enhances the

N-terminal region's binding affinity, particularly the first 10 residues, to LPC, while modestly reducing
the same region's affinity to DOPS.

To provide insights that are more relevant to physiological conditions, we extended our investigation to
examine how Ac-α-syn interacts with SV membranes. By utilizing a protocol established before [16], we
prepared monodisperse SVs isolated from normal mouse brains and titrated them with Ac-α-syn at an
SV/α-syn ratio of 260:3000, which is a physiologically relevant ratio within presynaptic terminals as
determined in previous work [17]. The HSQC spectrum revealed signal attenuation, notably in the
N-terminal region and specifically within the first 10 residues of α-syn (**Figure 3d**). These findings
closely mirrored those observed when Ac-α-syn was titrated with LPC micelles and DOPC/LPC-mixed
liposomes at low lipid/α-syn ratios of 10:1 and 100:1, respectively (**Figure 3e**). These results further
substantiate the critical role played by the N-terminal region of Ac-α-syn in SV membrane binding.

Ac-α-syn binding on LPC shows high intermolecular interactions
The intermolecular interactions between α-syn monomers [15] or between α-syn and VAMP2 [1] are also
essential for clustering SVs. To further characterize the accessibility of α-syn to LPC and DOPS, we
performed cross-linking experiments coupled with mass spectrometry (XL-MS). We mixed [15]N-labeled
and unlabeled Ac-α-syn at a 1:1 ratio, and selected the cross-linked fragments containing both
[15]N-labeled and unlabeled Ac-α-syn to map the intermolecular cross-linking pattern. We found the
accessibility of α-syn to LPC to be much higher than that to DOPS (**Figure 4a**). The number of
cross-linked peptide pairs, indicative of their close-interaction, detected in LPC-induced multimers was
over twice more than that of DOPS (**Table S1-6**). This result demonstrates high intermolecular
interactions between Ac-α-syn and LPC.

Discussion
In summary, the interaction between α-syn and lipids plays a crucial role in both the protein's
physiological functions in SV trafficking and its pathological aggregation, as observed in
neurodegenerative conditions [18]. Understanding how PTMs like N-acetylation modulate these
interactions and their subsequent effects is a critical area of study. Notably, in addition to in-cell NMR
findings that demonstrate spontaneous acetylation of α-syn, N-acetylated α-syn has been observed in PD
patients' brain tissue, underscoring its physiological relevance [19]. Our research reveals that N-acetylation

enhances the binding of the N-terminal region of α-syn to LPC, thereby facilitating its association with

SVs. LPC is a neutral lipid with an inverted-cone shape, known to create high membrane curvature and

more lipid packing defects [20]. N-acetylation effectively neutralizes the positive charge of α-syn's

N-terminus, fostering its insertion into LPC-rich membranes through hydrophobic interactions.

Conversely, N-acetylation significantly reduces the affinity of α-syn's N-terminal region for negatively

charged PS-containing membranes, where the interaction is mainly electrostatic in nature. Furthermore,

compared to DOPS, the interaction between N-terminally acetylated α-syn and LPC also triggers more

intermolecular interactions that support the functional α-syn for SV clustering (**Figure 4b**).

Most recently, we reported that α-syn also interacts with the neutral phospholipid LPC, which could be

enriched in synaptosomes. LPC can facilitate α-syn to perform its function on clustering SVs [2]. However,

the importance of the α-syn–LPC interaction remains unclear. Since α-syn is N-terminally acetylated in

mammalian cells, the involvement of LPC, not anionic phospholipids, in N-acetylation-enhanced SV

clustering demonstrates the importance of our recently reported α-syn–LPC interaction for mediating

α-syn's function in SV trafficking. Furthermore, the amount of anionic phospholipids on SVs could

changes with aging and disease [21], which may alter bindings modes of α-syn. The α-syn–LPC interaction

enhanced by N-acetylation could serve as an alternative mechanism for maintaining the SV clustering

function of α-syn.

Materials and Methods

Lipids. Lipids used in this study were purchased from Avanti Polar Lipids as follows: 16:0 LPC (1-palmitoyl-2-hydroxy-sn-glycero-3-phosphocholine) (855675), DOPC (1,2-dioleoyl-sn-glycero-3-phosphocholine) (850375), DOPS (1,2-dioleoyl-sn-glycero-3-phospho-L-serine) (840035), POPC (1-palmitoyl-2-oleoyl-glycero-3-phosphocholine) (850457), POPE (1-palmitoyl-2-oleoyl-sn-glycero-3-phosphoethanolamine) (850757), biotin-DPPE (1,2-dipalmitoyl-sn-glycero-3-phosphoethanolamine-N-(cap biotinyl)) (870277), cholesterol (70000).

Protein purification. N-terminal unmodified human α-syn (un-α-syn) and N-terminal 30-residue truncated α-syn (Δ30 α-syn) were cloned into the pET22 vector. For N-terminal acetylated α-syn (Ac-α-syn), the fission yeast NatB complex (N-acetyltransferase) gene was co-expressed with α-syn in bacteria to acetylate its N-terminus [22]. ^{15}N isotope-labeled Ac-α-syn or α-syn were produced in M9 minimal medium with uniformly labeled $^{15}NH_4Cl$ (1 g/L, Cambridge Isotope Laboratories). Not isotope-labeled Ac-α-syn or α-syn were produced in LB medium. Expression was induced by 1 mM isopropyl-1-thio-D-galactopyranoside (IPTG) at 37 °C in *E.coli* strain BL21-DE3, which OD_{600} reached 0.8-1.0. The collected bacteria were lysed in 100 mM Tris-HCl (pH 8.0), 1 mM EDTA, 1 mM phenylmethylsulfonyl fluoride (PMSF). The lysates were boiled for 10 min followed by centrifugation at 16,000 g for 30 min. 20 mg/ml streptomycin was mixed with supernatants. After a 30-min centrifugation at 16,000 g, the pH value of supernatants was adjusted to 3.5 by 2 M HCl, followed by centrifugation at 16,000 g for 30 min. The supernatants were then dialyzed in 25 mM Tris-HCl (pH 8.0) overnight at 4 °C. The dialyzed protein solution was filtered through a 0.22 μm membrane (Merck Millipore, SLGP033RB) and injected onto an ion-exchange Q column (GE Healthcare, 17515601), followed by further purification via size exclusion chromatography (GE Healthcare, Superdex^{TM} 75).

Isolation of synaptic vesicles from mouse brains. Neuronal synaptic vesicles (SV) were isolated from 8-week-old C57BL6 male mice (Lingchang Shanghai) by following published protocols [23]. The experiment was performed at 4 °C or on ice with pre-cooled reagents. Four fresh brains (~0.5 g for each) were homogenized in 25 mL 4 mM HEPES-NaOH (pH 7.4) and 320 mM sucrose buffer (HB) with protease inhibitors by using a PTFE pestle in a 40 mL glass tube (Sigma, P7984). The homogenate was centrifuged at 1,500 g for 10 min. The supernatant (S1) was collected and kept on ice. The pellet was resuspended with 25 mL HB and homogenized, followed by centrifugation at 1,500 g for 10 min. Then the supernatant (S2) combined with S1 were centrifuged at 20,000 g for 20 min. The pellet which contains synaptosomes was resuspended with 2 mL HB and then homogenized in 20 mL H_2O followed by adding 50 μL of 1M HEPES-NaOH (pH 7.4) and protease inhibitors. The homogenate was placed on ice for 30 min, and centrifuged at 20,000 g for 20min. The supernatant was ultra-centrifuged at 70,000 g for 45 min. The pellet was resuspended in buffers indicated in each analysis and homogenized by using a PTFE pestle in a 3 mL glass tube (Sigma, P7734). To further disrupt any remaining SV clusters before experiments, the homogenized SVs were drawn first through a 20-gauge hypodermic needle (Fisher Scientific, 0556121) attached to a 10-ml syringe, and then through a 27-gauge needle (Fisher Scientific, 22557176) and expelled. Mice experiments were conducted according to the protocols approved by Animal Care Committee of the Interdisciplinary Research Center on Biology and Chemistry (IRCBC), Chinese Academy of Sciences (CAS).

α-Syn-mediated SV clustering monitored by dynamic light scattering (DLS). The DLS experiments were performed using a DynaPro Nanostar instrument (Wyatt Technology, Santa Barbara, CA). A 50 μL

aliquot of each sample containing homogenous SV (total protein concentration of 50 μg/mL) and α-syn
variants (3 μM) was used for DLS measurement. Buffer used in this assay was 50 mM HEPES-KOH
(pH 7.4), 150 mM KCl. The following parameters were set for all measurements: correlation function
low cutoff was 1.5 μs; correlation function high cutoff was 6×10^4 μs; peak radius low cutoff was 0.5
nm; peak radius high cutoff was 10^4 nm. The results were reported as the average of 10 consecutive
autocorrelation functions (acquisition time = 5 s), and each sample was repeated measured by three
times.
206
Negatively stained transmission electron microscopy (TEM). An aliquot of 5 μL sample was pipetted
onto a glow discharged carbon-coated copper grid. 5 μL 3% (w/v) uranyl acetate was used for
negatively staining after washing the grid twice with 5 μL distilled H_2O. The grid was air-dried after
removing extra uranyl acetate by filter paper. TEM images were obtained by a electron microscope
(Thermo Fisher FEI Tecnai G2 TEM) at 120 kV equipped with a LaB_6 gun and a FEI BM Eagle CCD
camera.
213
Liposome preparation. DOPS dissolved in chloroform was evaporated using a dry nitrogen stream.
The dried lipid film was hydrated in buffers used for cross-linking (indicated below), and then sonicated
in a water bath at 65 °C for 10 min. To mimic the size of synaptic vesicles, the hydrated lipids were
extruded 41 times at 65 °C through a poly-carbonate film with a pore size of 50 nm (Whatman
Nucleopore Track-Etch) by using an extruder apparatus (Avanti Polar Lipids, 610000). LPC powder was
dissolved in buffers used for cross-linking for further analysis.
220
Single-vesicle clustering assay. The protein-free liposomes (~100 nm) of POPC, POPE, cholesterol,
LPC or DOPS, and biotin-DPPE or DiD (Invitrogen, D307) were prepared in the single-vesicle
clustering buffer (25 mM HEPES-NaOH (pH 7.4), 100 mM NaCl) to a total lipid concentration of ~10
mM. The group of unlabeled liposomes contain POPC, POPE, cholesterol, LPC or DOPS and
biotin-DPPE with a molar ratio of (49.9~59.9):20:20:(0~10):0.1. The group of labeled liposomes contain
POPC, POPE, cholesterol, LPC or DOPS and DiD with a molar ratio of (49~59):20:20:(0~10):1. LPC or
DOPS on both vesicles was varied from 0 to 10%, in expense of POPC.
228
For the clustering experiments, the unlabeled liposomes (100 μM) were injected into the sample
chamber and incubated for 30 min, followed by buffer exchange of 300 μL single-vesicle buffer. 10 μM
DiD labeled liposomes were incubated with 20 nM α-syn or Ac-α-syn, and loaded into the sample
chamber with 30 min incubation at room temperature. Unbound DiD liposomes and proteins were
removed by buffer exchange of 300 μL single-vesicle buffer. Sample slides with multiple channels were
monitored in a wide-field TIR fluorescence microscope. Data were acquired with a CCD camera, and
analyzed with the smCamera program. Six images were taken at random locations within each channel
on the quartz slide. The number of liposome interactions (clustering) was determined by counting the
number of fluorescent spots from emission of DiD upon excitation at 633 nm. Details were described
previously [1,24].
239
Delivering ^{15}N-un-α-syn into mammalian cells by using electroporation method. Electroporation
was performed with Neon transfection system (Invitrogen, MPK5000). Lyophilized ^{15}N-un-α-syn was
dissolved in Buffer R (Invitrogen, MPK10025) to final concentration of 300 μM. HEK-293T cells
(ATCC, CRL-3216) were collected and washed with PBS for 3 times to get rid of medium. Cells were
resuspendedin ^{15}N-un-α-syn solution at 8×10^7 cells per mL. Eight aliquots of 100 μL cell-protein

mixture were pipetted and electroporated. Pulse program was: 1,400 V pulse voltage, 20 ms pulse width,
1 pulse number. Electroporated cells were added into four 10 cm-diameter dishes with pre-warmed
culture medium. Cells were allowed to recover for 5 h, and were then harvested by trypsinization and
washed with PBS four time. Resuspended cells were washed in 160 μL pH-stable L-15 medium (Gibco,
11415064) and 40 μL D$_2$O, and allowed to settle in a 5 mm NMR tube by gentle sedimentation with a
hand-driven centrifuge. Thesupernatant was discarded. Finally, ~500 μL sedimented cell slurry was
prepared for NMR measurement.
252
In-cell and in vitro solution NMR spectroscopy. In-cell NMR experiments were carried out at 25 °C
on a Bruker 900 MHz spectrometer equipped with a cryogenic probe by following a published paper [16].
Bruker standard SOFAST-HMQC pulse sequence [25,26] was used and the ^{1}H shape pulse efficient was
optimized for collecting the 2D NMR spectrum of the in-cell samples with 80 scans. The delay time
(D1) was set to 0.29 s, and 1024 and 128 complex points were used for ^{1}H and ^{15}N, respectively. The
experiment duration time is 61 min 16 s. Cell viability before and after the SOFAST-HMQC NMR
experiment was assessed by trypan blue staining. Cell viability remained above 90% after the NMR
experiments.
261
In vitro NMR experiments were carried out at 25 °C on an 800 MHz Agilent spectrometer equipped
with a cryogenic probe (Agilent Technologies). The buffer used in all NMR titration assay was 50 mM
sodium phosphate buffer (pH 6.5) containing 50 mM NaCl and 10% D$_2$O (v/v). All of the LPC or
liposomes were prepared with a total molar concentration of 50 mM, and gradually added to the 500 μL
50 μM N-acetylated or unmodified α-syn in a serious of NMR titration assay. In NMR SV-titration
experiment, each spectrum was acquired from separately prepared samples with increasing amounts of
SV in 50 μM N-acetylated α-syn. The molar concentration of SVs was calculated according to
parameters from a published paper [27].
270
271 All the NMR data were processed using NMRPipe and analyzed by SPARKY [28]. Backbone resonance
272 assignment of N-acetylated and unmodified α-syn was accomplished according to a published paper [10]
273 and a Biological Magnetic Resonance Bank (BMRB) entry 6968 [29], respectively.
274
α-Syn multimer cross-linking experiment
LPC micelles, DOPS liposomes and α-syn variants were mixed in cross-linking-buffer (10 mM
HEPES-NaOH, pH 7.4). The stock of cross-linker DSS (Thermo Scientific, 21658) was dissolved in
DMSO to 40 mM. 50 μM protein and 2 mM DSS were incubated at indicated concentrations of lipid in
the cross-linking-buffer at 25 °C for 1 h. Crosslinking was terminated by addition of 100 mM Tris-HCl
(pH 7.4) at 25 °C for 10 min. For identification of the inter-molecular cross-linked peptide pairs from
α-syn multimers by MS analysis, half amount of α-syn was ^{15}N-labeled α-syn.
282
Protein cross-linking mass spectrometry (XL-MS) analysis
The cross-linked samples were precipitated using acetone at −20 °C for 30 min. Precipitates were dried
in air and resuspended in 8 M urea in 100 mM Tris-HCl (pH 8.5). Then, 5 mM TCEP was adding for 20
min incubation, followed by adding 10 mM iodoacetamide for 15 min for alkylation reaction in the dark.
Then, the samples were diluted to 2 M urea in 100 mM Tris-HCl (pH 8.5), and digested with trypsin
(molar ratio of protein to trypsin was 50:1) at 37 °C for 16 h in the presence of 1 mM CaCl$_2$ and 20 mM
methylamine. Digestion was terminated by adding formic acid to 5% final concentration (v/v). After
desalting by C18 desalting tips, the digested samples were air-dried and stored at -80 °C until analysis.

291
The digested peptides were analyzed by online nanoflow liquid chromatography tandem mass
spectrometry (LC−MS/MS). Briefly, nano LC−MS/MS-experiments were performed on an EASY-nLC
1000 system (Thermo Scientific) connected to an Orbitrap Q Exactive HF (Thermo Scientific) through a
nanoelectrospray ion source. The peptides were separated on a nano-column (C18, 100 μm × 15 cm, 1.9
μm, 120 Å) and further analyzed using an Orbitrap Q Exactive HF mass spectrometer. A 60-min
gradient was used in the nano LC at 300 nL/min flow rate: 0-4 min from 4% buffer B (80% ACN, 0.1%
FA), 96% buffer A (0.1% FA in water) to 8% buffer B, 4-45 min from 8% buffer B to 22% buffer B,
45-53min from 22% buffer B to 30% buffer B, then increasing to 95% buffer B in the next 41min, and
stay 95% buffer B for 3min. One full-scan mass spectrum (350-1500 m/z) at a resolution of 60,000
followed by HCD fragmentation and detection of the fragment ions (scan range from 350 to 1,500 m/z)
in Orbitrap at a 27% normalized collision energy was repeated continuously.

303
The cross-linked peptides pairs were identified by pLink [30] firstly. The pLink search parameters used in
this study includes: (1) 20 ppm for a window of precursor mass tolerance. (2) 4 ppm for a window of
fragment mass tolerance. (3) Lys for cross-linking sites of the cross-linker DSS. (4) 138.0680796 Da for
xlink mass shift. (5) 156.0786442 Da for monolink mass shift. (6) 57.02146 Da for fixed
carbamidomethyl-Cys modification mass shift. (7) 15.99491 Da for variable Met modification mass shift.
(8) 4-100 amino acids per chain for peptide length. (9) 400-10,000 Da per chain for peptide mass. (9) 2
missed cleavage sites per chain for trypsin digestion of protein. The searching results were filtered with
a false rate of less than 5% and E-values of less than 0.0001.

312
The pLink readout-data were filtered with a score less than 10^{-6} and mass error for crosslinking
assignments less than 4 ppm. To identifying the inter-protein cross-linked peptide pairs, the spectrum
($\Delta RT < 0.5$ s) of the two cross-linked peptide segments need to be manually checked with four peaks of
^{14}N-peptide and ^{14}N-peptide, ^{14}N-peptide and ^{15}N-peptide, ^{15}N-peptide and ^{14}N-peptide and ^{15}N-peptide
and ^{15}N-peptide.

Acknowledgments
C.L. and D.L. thank the support from the National Natural Science Foundation (NSF) of China
(92353302 and 32170683 to D.L.; 82188101 and 32171236 to C.L.), the Science and Technology
Commission of Shanghai Municipality (STCSM) (Grant No. 20XD1425000 and 22JC1410400 to C.L.),
the Shanghai Pilot Program for Basic Research – Chinese Academy of Science, Shanghai Branch (Grant
No. CYJ-SHFY-2022-005 to C.L.), the CAS Project for Young Scientists in Basic Research (Grant
No.YSBR-095 to C.L.). This work was supported by the National Institutes of Health (R01NS102181,
R01NS113960, 1RF1NS126342 and R21NS127939 to J.B.), and the Michael J Fox Foundation (16661
to J.D. and 16164 to J.B.).

Author contributions: J.D., and C.L. designed and directed the study. C.W., C.Z., Z.L., J.G., and S.Z.
prepared proteins, performed cellular and cross-linking experiments, and carried out EM, NMR, and MS
measurements. X.H. performed single-vesicle experiments. D.L., Y.Z., J.B., J.D., and C.L. analyzed the
data, discussed the results, and wrote the manuscript.

333
334 **Competing interests:** The authors declare no competing financial interests.

Figure 1 N-terminal acetylation enhances SV clustering induced by α-syn. (**a**) α-syn is N-terminally acetylated in cells. Upper: Schematic representation of the delivery of [15]N-labeled N-terminally unmodified α-synuclein (un-α-syn) into mammalian cells through electroporation. Lower: Comparisons of 2D [1]H-[15]N HSQC spectra of N-acetylated α-syn (Ac-α-syn) (black) with in-buffer un-α-syn (blue) and in-cell un-α-syn (red). Distinct assignments for NMR cross-peaks corresponding to amino acids in Ac-α-syn and un-α-syn are enclosed and labeled. (**b**) Scheme of experimental design for the functional study of Ac-α-syn on SV clustering. SVs isolated from mouse brain were added to Ac-α-syn (Upper) for measuring size distribution by DLS (middle), and the sample was further visualized by using negatively stained TEM (lower). (**c**) The influences of N-acetylation of α-syn and (**d**) α-syn without the N-terminal 30 aa on SV clustering measured by DLS. The X-axis represents number percent of the single SV and clustered SVs counted by DLS. Error bars are standard deviations from 3 biological replicates. **, p-value < 0.01; ***, p-value < 0.001; analysis by Student's t test. (**e**) Representative negatively stained TEM images of the single SV and clustered SVs in SV samples with no α-syn (grey) and Ac-α-syn (red), respectively.

Figure 2 LPC mediates the enhancement of vesicle clustering by N-terminal acetylation of α-syn.
(**a**) Scheme of single-vesicle clustering assay for the functional study of Ac-α-syn on vesicle clustering. Vesicles were prepared with different amounts of LPC, and labeled with DiD (red) and DiI (green), respectively. A saturated layer DiD-vesicles were immobilized on the imaging surface. Free DiI-vesicles were injected into the system with Ac-α-syn. Green laser illumination imaged the DiI-vesicles that clustered with DiD-vesicles. The enhancement of LPC (**b**) and DOPS (**c**) on single vesicle clustering probability by Ac-α-syn and un-α-syn, respectively. Error bars are standard deviations from 6 random imaging locations in the same sample channel. *** indicates p-value < 0.001, analysis by Student's t test.

Figure 3 N-terminal acetylation increases the α-syn–LPC interaction. Comparisons of
residue-resolved NMR signal intensity ratios (I/I_0) of un-α-syn (upper) and Ac-α-syn (lower) during
titration with LPC micelles (**a**), LPC-containing liposomes (DOPC:LPC = 4:1, mol:mol) (**b**), and DOPS
liposomes (**c**) at indicated protein/lipid molar ratios. Dash lines highlights the residue positions of 30 and
95. (**d**) SVs isolated from mouse brains were employed for NMR titration with ^{15}N-Ac-α-syn,
approximating the physiological ratio (α-syn:SV = 4000:300, mol:mol). Residue-resolved NMR signal
intensity ratios (I/I_0) of Ac-α-syn titrated by SVs to that in solution. The molar ratios of SV to Ac-α-syn
are indicated. LPC titration in the Ac-α-syn/LPC ratio of 1:10 (blue curve) is overlaid on the SV titration.
(**e**) 2D ^{1}H-^{15}N HSQC spectra of NMR for un-α-syn with LPC micelles and Ac-α-syn with LPC micelles,
LPC-containing liposomes, and mouse SVs. The NMR cross-peak of the first 10 residues are highlighted
and magnified, as depicted on the right side of each spectrum set (Note: the first and second residues of
un-α-syn cannot be assigned).

Figure 4 Ac-α-syn binding on LPC shows high intermolecular interactions. (**a**) The cross-linking
patterns of Ac-α-syn in the presence of LPC and DOPS mapped by MS at the protein/lipid molar ratio of
1:50. Lines present the inter-molecular cross-linked residues between two individual ^{15}N-labeled and
unlabeled Ac-α-syn. The grayscale of the lines corresponds to the frequency of the cross-linked pairs
identified in three individual experiments. Source data are provided in **Tables S1-S6**. (**b**) Ac-α-syn binds
strongly to LPC through the N-terminal region (red arrow), and leaves more unbound NAC and
C-terminus for intermolecular interaction (green arrow). In contrast, N-ternimal acetalytion reduces
α-syn's binding to DOPS, and due to the negtively charged headgroup of PS, Ac-α-syn binding on
DOPS extends to NAC region, which limits intermolecular interactions.

References:

1. Diao, J., Burre, J., Vivona, S., Cipriano, D.J., Sharma, M., Kyoung, M., Sudhof, T.C., and Brunger, A.T. (2013). Native alpha-synuclein induces clustering of synaptic-vesicle mimics via binding to phospholipids and synaptobrevin-2/VAMP2. Elife *2*, e00592. 10.7554/eLife.00592.

2. Lai, Y., Zhao, C., Tian, Z., Wang, C., Fan, J., Hu, X., Tu, J., Li, T., Leitz, J., Pfuetzner, R.A., et al. (2023). Neutral lysophosphatidylcholine mediates α-synuclein-induced synaptic vesicle clustering. Proc Natl Acad Sci U S A *120*, e2310174120. 10.1073/pnas.2310174120.

3. Burre, J., Sharma, M., Tsetsenis, T., Buchman, V., Etherton, M.R., and Sudhof, T.C. (2010). alpha-Synuclein Promotes SNARE-Complex Assembly in Vivo and in Vitro. Science *329*, 1663-1667. Doi 10.1126/Science.1195227.

4. Goedert, M., Spillantini, M.G., Del Tredici, K., and Braak, H. (2013). 100 years of Lewy pathology. Nat Rev Neurol *9*, 13-24. 10.1038/nrneurol.2012.242.

5. Uversky, V.N., and Eliezer, D. (2009). Biophysics of Parkinson's disease: structure and aggregation of alpha-synuclein. Curr Protein Pept Sci *10*, 483-499. 10.2174/138920309789351921.

6. Fauvet, B., Fares, M.B., Samuel, F., Dikiy, I., Tandon, A., Eliezer, D., and Lashuel, H.A. (2012). Characterization of semisynthetic and naturally Nα-acetylated α-synuclein in vitro and in intact cells: implications for aggregation and cellular properties of α-synuclein. J Biol Chem *287*, 28243-28262. 10.1074/jbc.M112.383711.

7. Oueslati, A. (2016). Implication of Alpha-Synuclein Phosphorylation at S129 in Synucleinopathies: What Have We Learned in the Last Decade? J Parkinsons Dis *6*, 39-51. 10.3233/jpd-160779.

8. Mahul-Mellier, A.L., Fauvet, B., Gysbers, A., Dikiy, I., Oueslati, A., Georgeon, S., Lamontanara, A.J., Bisquertt, A., Eliezer, D., Masliah, E., et al. (2014). c-Abl phosphorylates α-synuclein and regulates its degradation: implication for α-synuclein clearance and contribution to the pathogenesis of Parkinson's disease. Human molecular genetics *23*, 2858-2879. 10.1093/hmg/ddt674.

9. Zhang, H., Zhu, L., Sun, L., Zhi, Y., Ding, J., Yuan, Y.S., Shen, F.F., Li, X., Ji, P., Wang, Z., et al. (2019). Phosphorylated α-synuclein deposits in sural nerve deriving from Schwann cells: A biomarker for Parkinson's disease. Parkinsonism Relat Disord *60*, 57-63. 10.1016/j.parkreldis.2018.10.003.

10. Theillet, F.X., Binolfi, A., Bekei, B., Martorana, A., Rose, H.M., Stuiver, M., Verzini, S., Lorenz, D., van Rossum, M., Goldfarb, D., and Selenko, P. (2016). Structural disorder of monomeric alpha-synuclein persists in mammalian cells. Nature *530*, 45-50. 10.1038/nature16531.

11. Marotta, N.P., Lin, Y.H., Lewis, Y.E., Ambroso, M.R., Zaro, B.W., Roth, M.T., Arnold, D.B., Langen, R., and Pratt, M.R. (2015). O-GlcNAc modification blocks the aggregation and toxicity of the protein α-synuclein associated with Parkinson's disease. Nat Chem *7*, 913-920. 10.1038/nchem.2361.

12. Bu, B., Tong, X., Li, D., Hu, Y., He, W., Zhao, C., Hu, R., Li, X., Shao, Y., Liu, C., et al. (2017). N-Terminal Acetylation Preserves alpha-Synuclein from Oligomerization by Blocking Intermolecular Hydrogen Bonds. ACS Chem Neurosci *8*, 2145-2151. 10.1021/acschemneuro.7b00250.

13. Runfola, M., De Simone, A., Vendruscolo, M., Dobson, C.M., and Fusco, G. (2020). The N-terminal Acetylation of α-Synuclein Changes the Affinity for Lipid Membranes but not the Structural Properties of the Bound State. Sci Rep *10*, 204. 10.1038/s41598-019-57023-4.

14. Trexler, A.J., and Rhoades, E. (2012). N-Terminal acetylation is critical for forming α-helical
oligomer of α-synuclein. Protein science : a publication of the Protein Society *21*, 601-605.
10.1002/pro.2056.
15. Fusco, G., Pape, T., Stephens, A.D., Mahou, P., Costa, A.R., Kaminski, C.F., Kaminski Schierle,
G.S., Vendruscolo, M., Veglia, G., Dobson, C.M., and De Simone, A. (2016). Structural basis of
synaptic vesicle assembly promoted by alpha-synuclein. Nat Commun *7*, 12563.
10.1038/ncomms12563.
16. Wang, C., Tu, J., Zhang, S., Cai, B., Liu, Z., Hou, S., Zhong, Q., Hu, X., Liu, W., Li, G., et al.
(2020). Different regions of synaptic vesicle membrane regulate VAMP2 conformation for the
SNARE assembly. Nat Commun *11*, 1531. 10.1038/s41467-020-15270-4.
17. Wilhelm, B.G., Mandad, S., Truckenbrodt, S., Kröhnert, K., Schäfer, C., Rammner, B., Koo, S.J.,
Claßen, G.A., Krauss, M., Haucke, V., et al. (2014). Composition of isolated synaptic boutons
reveals the amounts of vesicle trafficking proteins. Science *344*, 1023-1028.
10.1126/science.1252884.
18. Burré, J., Sharma, M., and Südhof, T.C. (2018). Cell Biology and Pathophysiology of
α-Synuclein. Cold Spring Harb Perspect Med *8*. 10.1101/cshperspect.a024091.
19. Anderson, J.P., Walker, D.E., Goldstein, J.M., de Laat, R., Banducci, K., Caccavello, R.J.,
Barbour, R., Huang, J., Kling, K., Lee, M., et al. (2006). Phosphorylation of Ser-129 is the
dominant pathological modification of alpha-synuclein in familial and sporadic Lewy body
disease. J Biol Chem *281*, 29739-29752. 10.1074/jbc.M600933200.
20. Lauwers, E., Goodchild, R., and Verstreken, P. (2016). Membrane Lipids in Presynaptic
Function and Disease. Neuron *90*, 11-25. 10.1016/j.neuron.2016.02.033.
21. Emre, C., Do, K.V., Jun, B., Hjorth, E., Alcalde, S.G., Kautzmann, M.I., Gordon, W.C., Nilsson,
P., Bazan, N.G., and Schultzberg, M. (2021). Age-related changes in brain phospholipids and
bioactive lipids in the APP knock-in mouse model of Alzheimer's disease. Acta Neuropathol
Commun *9*, 116. 10.1186/s40478-021-01216-4.
22. Johnson, M., Coulton, A.T., Geeves, M.A., and Mulvihill, D.P. (2010). Targeted amino-terminal
acetylation of recombinant proteins in E. coli. Plos One *5*, e15801.
10.1371/journal.pone.0015801.
23. Ahmed, S., Holt, M., Riedel, D., and Jahn, R. (2013). Small-scale isolation of synaptic vesicles
from mammalian brain. Nat Protoc *8*, 998-1009. 10.1038/nprot.2013.053.
24. Diao, J., Ishitsuka, Y., Lee, H., Joo, C., Su, Z.L., Syed, S., Shin, Y.K., Yoon, T.Y., and Ha, T.
(2012). A single vesicle-vesicle fusion assay for in vitro studies of SNAREs and accessory
proteins. Nature Protocols *7*, 921-934. Doi 10.1038/Nprot.2012.020.
25. Schanda, P., and Brutscher, B. (2005). Very fast two-dimensional NMR spectroscopy for
real-time investigation of dynamic events in proteins on the time scale of seconds. J Am Chem
Soc *127*, 8014-8015. 10.1021/ja051306e.
26. Schanda, P., Kupce, E., and Brutscher, B. (2005). SOFAST-HMQC experiments for recording
two-dimensional heteronuclear correlation spectra of proteins within a few seconds. J Biomol
NMR *33*, 199-211. 10.1007/s10858-005-4425-x.
27. Takamori, S., Holt, M., Stenius, K., Lemke, E.A., Gronborg, M., Riedel, D., Urlaub, H., Schenck,
S., Brugger, B., Ringler, P., et al. (2006). Molecular anatomy of a trafficking organelle. Cell *127*,
831-846. Doi 10.1016/J.Cell.2006.10.030.
28. Delaglio, F., Grzesiek, S., Vuister, G.W., Zhu, G., Pfeifer, J., and Bax, A. (1995). NMRPipe: a
multidimensional spectral processing system based on UNIX pipes. J Biomol NMR *6*, 277-293.
10.1007/bf00197809.

29.   Bermel, W., Bertini, I., Felli, I.C., Lee, Y.M., Luchinat, C., and Pierattelli, R. (2006). Protonless
475   NMR experiments for sequence-specific assignment of backbone nuclei in unfolded proteins. J
476   Am Chem Soc *128*, 3918-3919. 10.1021/ja0582206.
477   30.   Fan, S.B., Meng, J.M., Lu, S., Zhang, K., Yang, H., Chi, H., Sun, R.X., Dong, M.Q., and He,
478   S.M. (2015). Using pLink to Analyze Cross-Linked Peptides. Curr Protoc Bioinformatics *49*,
479   8.21.21-28.21.19. 10.1002/0471250953.bi0821s49.
480