Opto-controlled C9orf72 poly-PR forms anisotropic condensates causative of TDP-43 pathology in the nucleus

Rachel E Hodgson*, Jessica Rayment, Wan-Ping Huang, Anna Sanchez Avila, Tatyana A Shelkovnikova*

Sheffield Institute for Translational Neuroscience and Neuroscience Institute, University of Sheffield, Sheffield, UK

Abstract: Proteinaceous inclusions formed by *C9orf72* derived dipeptide-repeat (DPR) proteins are a histopathological hallmark in ~50% of familial amyotrophic lateral sclerosis/frontotemporal dementia (ALS/FTD) cases. However DPR aggregation/inclusion formation could not be efficiently recapitulated in cell models for four out of five DPRs. In this study, using optogenetics, we achieved chemical-free poly-PR condensation/aggregation in cultured cells, with spatial and temporal control. Strikingly, nuclear poly-PR condensates had anisotropic, hollow-centre appearance, resembling anisosomes formed by aberrant TDP-43 species, and their growth was limited by RNA. These condensates induced abnormal TDP-43 granulation in the nucleus without the activation of stress response. Cytoplasmic poly-PR aggregates that formed under prolonged light stimulation were more persistent than its nuclear condensates, selectively sequestered TDP-43 in a demixed state and surrounded spontaneous stress granules. Our data suggest that poly-PR anisotropic condensation in the nucleus, causative of nuclear TDP-43 dysfunction, may constitute an early pathological event in C9-ALS/FTD. Anisosome-type condensates may represent a more common cellular pathology in neurodegeneration than previously thought.

corresponding authors: r.hodgson1@sheffield.ac.uk, t.shelkovnikova@sheffield.ac.uk

Key words: C9orf72, ALS, FTD, DPR, poly-PR, optogenetics, condensate, TDP-43

Highlights:

- Optogenetics can be used to model *C9orf72* DPR condensation in cultured cells.
- Opto-PR forms hollow nuclear condensates, and RNA limits their growth by fusion.
- Opto-PR condensation leads to stress-independent TDP-43 pathology in the nucleus.
- Cytoplasmic poly-PR assemblies are persistent and selectively sequester TDP-43.

Graphical abstract:

Introduction

A G_4C_2 hexanucleotide repeat expansion (HRE) in the first intron of the *C9orf72* gene is the
most common genetic alteration associated with amyotrophic lateral sclerosis (ALS) and
frontotemporal dementia (FTD) (DeJesus-Hernandez et al, 2011; Renton et al, 2011). Healthy
individuals commonly carry 2 repeats, while C9-ALS/FTD is associated with ≥30 repeat lengths
(Van Mossevelde et al, 2017). Production of dipeptide repeat (DPR) proteins from the *C9orf72-*
HRE transcripts is one of the proposed mechanisms of repeat toxicity (Ash et al, 2013; Mori et
al, 2013; Zu et al, 2013). Both sense and antisense C9orf72-HRE transcripts are translated in
all reading frames via non-canonical, repeat-associated non-AUG (RAN) translation to produce
five DPRs: poly-GA, -GR (sense), -PR, -PA (antisense) and -GP (both sense and antisense). All
five DPRs have been detected in the patient CNS, primarily as cytoplasmic and less often
nuclear inclusions in neurons and glia (Gendron *et al*, 2013; Mackenzie *et al*, 2015; Saberi *et al*,
2018; Sakae *et al*, 2018; Schludi *et al*, 2015). Whilst poly-GA and -GP are the most abundant,
arginine containing DPRs (R-DPRs) poly-GR and poly-PR are the most toxic species in cells
and *in vivo* (Kwon *et al*, 2014; Mizielinska *et al*, 2014; Moens *et al*, 2019). For example, nuclear
poly-PR aggregation was sufficient to induce ALS-like phenotypes in non-human primates (Xu
et al, 2023), and expression of this DPR was extremely toxic in mice (LaClair *et al*, 2020; Zhang
et al, 2019).

R-DPR inclusion pathology seen in patients and mouse models (Chew *et al*, 2019; Choi *et al*,
2019; Cook *et al*, 2020) is not easily reproducible in cultured cells, where R-DPRs typically
display diffuse distribution (outside the nucleolus), even when overexpressed and independent
of the repeat length; this is in contrast to poly-GA that readily forms large cytoplasmic
aggregates (Frottin *et al*, 2021; Hartmann *et al*, 2018; Liu *et al*, 2022; Lopez-Gonzalez *et al*,
2016; Vanneste *et al*, 2019). High arginine content renders R-DPRs highly hydrophilic and
hence soluble. Although poly-PR and poly-GR have similar biochemical properties, molecular
dynamics simulations revealed that poly-PR is capable of (limited) self-association – forming
either dimers or small amorphous oligomers, whereas poly-GR is not likely to form any stable
oligomeric species (Zheng *et al*, 2021). Consistently, *in vitro,* R-DPRs undergo phase
separation only in the presence of additional agents – RNA or proteins (Balendra *et al*, 2023;
Boeynaems *et al*, 2017; Lee *et al*, 2016; Zhang *et al.*, 2019). This suggests that certain factors
in the cellular environment drive R-DPR loss of solubility in ALS/FTD. Sensitive immunoassays
revealed that soluble DPRs are less abundant in the more affected brain regions, as compared
to the relatively spared regions (Quaegebeur *et al*, 2020).

Intermediate products of R-DPR aggregation may contribute to cellular dysfunction, however
due to difficulty of modelling this molecular event, their role could not be investigated in the
cellular setting. We hypothesised that the use of optogenetics (Park *et al*, 2017; Shin *et al*,
2017) will allow circumventing the intrinsic solubility of R-DPRs, by triggering and maintaining
an oligomerised R-DPR state. Indeed, using Cry2olig tagging, we were able to reliably induce
condensation of poly-PR ("opto-PR") in cultured cells, with spatial and temporal control. Using
this model, we show that: i) Poly-PR can form nuclear condensates with a specific ordered
arrangement reminiscent of TDP-43 "anisosomes" (Yu *et al.,* 2021), and these assemblies
trigger abnormal nuclear granulation of TDP-43. ii) Poly-PR cytoplasmic aggregates can form
by clustering around stress granules and can nucleate cytoplasmic TDP-43 aggregates, whilst
maintaining a TDP-43/DPR demixed state. These findings unveil a converging molecular
mechanism for aberrant C9orf72-DPR and TDP-43 species – formation of ordered nuclear
condensates. Furthermore, they link putative early disease-stage C9-ALS/FTD species to TDP-

43 pathology – the main correlate of neurodegeneration. Finally, they provide a possible
mechanism of large DPR inclusion nucleation/seeding in the cytoplasm.
83
Results

Optogenetic modeling of C9orf72 poly-PR condensation in cells

We utilised Cry2olig, a Cry2 variant with high oligomerisation capacity (Taslimi *et al*, 2014), in
an attempt to induce and maintain R-DPR self-association. 'Opto-DPR' constructs were
generated for codon-optimised expression of poly-PR, poly-GR and additionally poly-GP (36
repeats), tagged with Cry2olig and mCherry on the N-terminus (Fig.1A). Opto-DPRs
demonstrated a subcellular distribution pattern in HeLa cells typical for the 30-1000 repeat
range (Bennion Callister *et al*, 2016; Kanekura *et al*, 2018; Liu *et al.*, 2022; Vanneste *et al.*,
2019), where poly-GR and -GP were predominantly cytoplasmic and poly-PR was
predominantly nuclear, with high enrichment in the nucleolus; none of the DPRs showed signs
of aggregation (Fig.1B). Upon single-pulse light stimulation on a custom blue-light array,
Cry2olig vector control, opto-PR and opto-GP but not opto-GR readily formed visible
clusters/foci (Fig.1B), consistent with the low poly-GR self-association capacity reported in
simulation studies (Zheng *et al.*, 2021). Opto-PR foci were small (<500 nm), dot-like and
exclusively nuclear, whereas opto-GP formed large amorphous cytoplasmic inclusions (Fig.1B).
We focused on opto-PR thereafter, using opto-GR as a control.

Nuclear opto-PR foci/condensates were negative for the two nucleolar markers tested, fibrillarin
(FBL) and UPF1, confirming that they are not merely fragments of the nucleolus (Fig.S1A).
Continuous 3-h blue-light stimulation led to an increased opto-PR condensate size, as
compared to single pulse (Fig.1C). By super-resolution microscopy (SRM), the condensates
induced by a single light pulse were found to represent a mixed population of ~100 nm dot-like
and ~250 nm spherical, hollow-centre assemblies resembling anisotropic vesicle-like structures
formed by arginine-rich peptide/RNA mixtures *in vitro* (Alshareedah *et al*, 2020). Furthermore,
the larger condensates formed after 3-h continuous light exposure were found to represent
multiples of these spheroids (Fig.1D). Although Cry2olig protein on its own also formed clusters
throughout the cell in response to blue light (Fig.1B) (Taslimi *et al.*, 2014), these structures
appeared disordered/filamentous and were clearly different from opto-PR condensates (Fig.1D).
Opto-PR condensates could be detected by a PR-repeat specific antibody used in human tissue
studies (Fig.1E). Therefore, poly-PR confers a specific architecture to light-inducible
condensates.

We next set up an imaging approach for simultaneous induction and tracking of opto-PR
condensation on a confocal high-content imaging system. Even extremely short blue-light
exposures and low 488 nm laser power (50 ms/5%) were sufficient to induce opto-PR foci; a
combination of 500 ms exposure/80% laser power was used in subsequent experiments as
consistently and robustly inducing condensates (Fig.S1B). Visible opto-PR condensates
appeared within 2 min post-pulse and could be maintained by both short- and long-interval
repetitive opto-stimulation (every 2 min or every 15 min, respectively, Fig.1F). Condensates
were reversible, typically resolving within ~14 min after the last light pulse (Fig.1F). Opto-PR
condensate nucleation was concentration-dependent, with more structures forming in higher-
expressing cells (R=0.65) (Fig.1G; Fig.S1C). FRAP analysis revealed limited dynamics of opto-
PR within these assemblies, with low recovery after photobleaching of the entire structure,
despite a significant amount of diffuse opto-PR in the nucleoplasm (Fig.1H). A fraction of cells
(22.0±8.7%) developed persistent opto-PR condensates after continuous light stimulation,
which were still detectable after 3 h of recovery in the dark (Fig.1I).

Poly-PR was previously shown to interact with nucleophosmin (NPM1) and to co-partition with
this protein into phase-separated droplets *in vitro* (Lee et al., 2016; White et al, 2019).
Consistent with this, opto-PR condensates stained positive for NPM1, where NPM1 formed a
"shell" around the condensates, suggesting its secondary recruitment (Fig.1J). Interestingly,
opto-stimulation induced opto-PR signal segregation in the nucleolus (Fig.S1D). In this, we
observed intra-nucleolar demixing of opto-PR from NPM1, where the proteins fully co-localised
in the granular component (GC) under dark conditions but formed two distinct phases after
opto-stimulation (Fig.S1E,F). In contrast, opto-PR and FBL (the latter residing in the nucleolar
dense fibrillar component, DFC) showed no co-localisation both under dark and light conditions
(Fig.S1E,F).

R-DPRs were shown to promiscuously interact with membraneless organelle (MLO)
components – RNA and proteins with low-complexity domains, leading to wide-spread MLO
dysfunction (Kwon *et al.*, 2014; Lee *et al.*, 2016; Lin *et al*, 2016; Liu *et al.*, 2022). We
investigated the effect of opto-PR and its condensation on MLOs, focusing on those in the
nucleus due to the predominantly nuclear localisation of this DPR. Systematic analysis of four
MLOs – nuclear bodies (Gems, Cajal bodies, paraspeckles and speckles) revealed only minor
changes in the number and size in the presence of diffuse or condensed opto-PR (Fig.S2).
Cytoplasmic stress granules (SGs) were not induced by opto-PR with or without blue-light
stimulation (3 h continuous), consistent with its mainly nuclear localisation (data not shown).

Thus, microscopically visible DPR self-assembly/condensation in cultured cells can be achieved
using Cry2olig tagging, allowing the formation of DPR-specific, ordered assemblies. Opto-PR
condensates are characterised by concentration-dependent growth and low dynamic properties
associated with persistence, and sequester NPM1.

RNA limits the growth of anisotropic poly-PR condensates in cells

We next asked whether our opto-model can be used to characterise potential modifiers of poly-
PR condensation. RNA was previously found to promote R-DPR phase separation *in vitro*
(Balendra *et al.*, 2023; Boeynaems *et al.*, 2017; Gittings *et al*, 2020). More recently, using RNA-
protein crosslinking, R-DPRs have been shown to bind RNA in cells, in particular ribosomal
RNA (rRNA), with a preference for GA-rich sequences (Balendra *et al.*, 2023; Ortega *et al*,
2023). Using electrophoretic mobility shift assay (EMSA) with an RNA oligonucleotide
representing a naturally occurring RNA sequence containing 5xGA repeats (Clip34nt)
(Bhardwaj *et al*, 2013), we indeed observed that synthetic poly-PR and -GR peptides (10-mers)
form complexes with RNA (Fig.2A).

Having confirmed RNA binding, we next studied the impact of RNA depletion on opto-PR
condensates. Opto-PR expressing cells were treated with two transcriptional blockers, a global
inhibitor (actinomycin D) and an RNA polymerase II-specific inhibitor (5,6-Dichloro-1-β-D-
ribofuranosylbenzimidazole, DRB), followed by induction and tracking of opto-PR condensates
in individual cells for up to 4 h (long-interval repetitive stimulation, in the presence of the
inhibitor). Both inhibitors caused dramatic nucleolar shrinking confirming their activity in cells
(Fig.2B) (Shav-Tal *et al*, 2005). Opto-PR condensates formed in RNA-depleted conditions
appeared noticeably larger compared to control (Fig.2B). Similar result was obtained with a 3-h
continuous opto-stimulation on blue-light array, and quantification confirmed a significant
increase in the opto-PR condensate size (Fig.2C). These larger structures remained negative
for the nucleolar marker FBL (Fig.2C). DRB is a reversible inhibitor, which allowed us to analyse
possible changes in the stability of RNA-depleted opto-PR condensates. Cells were stimulated
for 3 h in the presence or absence of DRB followed by 2 h of recovery in the dark. Opto-PR
condensates formed in the presence of DRB appeared significantly more persisting, still

detectable following the recovery in 63±6.4% cells, compared to 30±7.8% cells in control
condition (Fig.2D).

Increase in the opto-PR condensate size upon actinomycin D treatment was accompanied by a
decrease in their number (Fig.2C), suggesting clustering or fusion. SRM analysis revealed that
opto-PR condensates in actinomycin D treated cells were no longer clusters of individual ~250
nm spherical units seen in the untreated or DRB condition but instead represented larger (>500
nm) hollow-centre spheres (Fig.2E). RNA on the surface of condensates has been shown to
limit their fusion, including in cells (Cochard *et al*, 2022). Actinomycin D but not DRB depletes
ribosomal RNA (rRNA), and we observed dense rRNA signal on the opto-PR condensate
surface which disappeared after actinomycin D treatment (Fig.2F). Therefore, the opto-PR
condensates in actinomycin D treated cells may form by fusion of the smaller units due rRNA
depletion from the surface, followed by relaxation into a larger spherical assembly.

Another putative modifier of R-DPR self-association/condensation is arginine dimethylation
(DMA), which increases the fluidity of R-DPR droplets *in vitro* and was found enriched in R-DPR
inclusions in C9-ALS/FTD patient tissue (Gittings *et al.*, 2020). We employed two small
molecule methyltransferase inhibitors, MS023 inhibiting five type-I methyltransferases that
synthesise asymmetric DMA (aDMA) and EPZ015666, a specific inhibitor of PRMT5
responsible for most symmetric DMA (sDMA) (Chan-Penebre *et al*, 2015; Eram *et al*, 2016).
Opto-PR expressing cells were treated with MS023 and EPZ015666 for 24 h and then exposed
to blue-light for 3 h continuously, followed by opto-PR condensate quantification. We observed
a mild decrease in the condensate number after aDMA depletion without changes in their size
or structure (Fig.S3). Thus, removing DMA marks may attenuate opto-PR condensate
nucleation, although this effect is small.

Therefore, our cellular opto-PR model can be utilised to analyse modifiers of poly-PR self-
assembly in the cellular context, as exemplified by the modulatory effect of RNA that we have
uncovered.

203   *Poly-PR condensation induces nuclear TDP-43 pathology*

R-DPR interactomes are enriched in RNA-binding proteins (RBPs) (Kwon *et al.*, 2014; Lee *et*
al., 2016). We therefore examined the effect of nuclear opto-PR condensation on ALS/FTD
relevant RBPs. TDP-43, FUS, NONO and SFPQ, all tagged with GFP, were co-expressed with
opto-PR and their subcellular distribution was examined with and without opto-stimulation.
Opto-PR presence *per se* did not affect RBP distribution (Fig.3A; Fig.S4A). However, light-
stimulated opto-PR expressing cells displayed a striking nuclear condensation phenotype for
TDP-43 but not other RBPs analysed (Fig.3A; Fig.S4A). Although we did observe TDP-43
condensation in a fraction of light-stimulated Cry2olig-expressing, it was significantly smaller
than in opto-PR cultures; nor was it observed in light-stimulated opto-GR expressing cells
(Fig.3B). Opto-PR assemblies were frequently found in the physical contact with TDP-43
condensates (44±0.5% of all opto-PR foci) (Fig.3C), suggestive of a direct nucleating effect of
oligomerising opto-PR. Fractionation confirmed reduced solubility of TDP-43 GFP after
induction of opto-PR condensation but not in light-stimulated opto-GR expressing cells (Fig.3D).
This analysis also confirmed reduced solubility of opto-PR but not opto-GR in light-stimulated
cells (Fig.3D).

TDP-43 GFP nuclear condensates induced by opto-PR were strikingly similar in their
morphology to the condensates formed during the recovery from arsenite stress (Wang *et al.*,
2020; Cohen *et al.,* 2015; Huang, Ellis *et al.,* manuscript in revision). Like these stress-induced
foci, TDP-43 condensates induced by opto-PR were devoid of polyA+ RNA (Fig.3E). We
therefore asked whether TDP-43 condensation elicited by opto-PR was due to an upregulated

stress response. Classic cellular stress markers, phosho-eIF2α, GADD34 and ATF4, were not
altered in opto-PR expressing cells after 3-h light stimulation (Fig.S4B,C). This is in contrast to
the dramatic upregulation of these markers during the recovery from sodium arsenite stress
used as a positive control (Fig.S4C). In addition to demonstrating the stress-unrelated nature of
TDP-43 condensates induced by opto-PR self-assembly, this experiment also confirmed a lack
of phototoxicity in our model.

TDP-43 acetylation was shown to impair its RNA binding and enhance aggregation, with
acetylated TDP-43 inclusions detected in sALS (Cohen *et al*, 2015). We investigated the effect
of opto-PR condensation on an acetylation-mimic TDP-43 mutant, K145Q (Cohen *et al.*, 2015).
In agreement with the published data, TDP-43 K145Q was prone to spontaneously forming
nuclear foci/granules in naïve cells (Fig.S4D). Opto-PR condensation potentiated the pro-
aggregating effect of acetylation, further increasing nuclear granulation of TDP-43 K145Q
(Fig.3F,G).

We next examined the effect of R-DPRs on TDP-43 higher-order assembly *in vitro,* using the
condensate immunodetection/imaging assay we recently developed (Hodgson, Huang *et al.,*
2024, doi: 10.2139/ssrn.4721338) (Fig.3H). Both poly-PR and poly-GR peptides were included
in these studies. "Supernatant" fraction of recombinant TDP-43 (after removal of preformed
aggregates) containing small protein clusters and soluble protein was incubated with equimolar
amounts of synthetic R-DPR peptides, followed by sedimentation and fixation of TDP-43
clusters on coverslips for immunostaining/imaging; in parallel, samples were fractionated by
centrifugation for western blot analysis (Fig.3H). Addition of both R-DPRs significantly
enhanced TDP-43 clustering – manifested as an increased cluster size and decreased cluster
number, with poly-GR being more potent (Fig. 3I). Fractionation confirmed increased
partitioning of TDP-43 to the pellet fraction in the presence of R-DPRs (Fig.3J). In contrast,
addition of a peptide with a "generic" sequence moderately enriched in proline and containing
no arginine residues (V5: GKPIPNPLLGLDST) did not induce TDP-43 clustering (Fig.S4E),
ruling out a non-specific molecular crowding effect of R-DPRs.

Collectively, these results suggest that poly-PR condensation can directly cause changes to the
nuclear distribution of TDP-43 without activation of stress signaling.

Cytoplasmic poly-PR assemblies are persistent and selectively sequester TDP-43

Having characterised the nuclear phenotypes, we asked whether our opto-PR model is
amenable to reproducing the cytoplasmic pathology typical for DPRs. Continuous 24-h long
stimulation on the blue-light array resulted in significant redistribution of opto-PR to the
cytoplasm, with cytoplasmic foci formation in 32% of cells (>300 transfected cells analysed;
Fig.4A,B). This was accompanied by a reduction in the incidence of nuclear opto-PR
condensates (from 94% cells after 3 h to 9% after 24 h of stimulation) (Fig.4A,B). Nuclear and
cytoplasmic opto-PR foci induced by 24-h opto-stimulation were persisting, with no significant
decline observed after 8 h of recovery in the dark (Fig.4A,B). This is in contrast to the nuclear
opto-PR condensates forming after a 3-h stimulation that were largely cleared after 3 h of
recovery in the dark (Fig.4A; Fig.1I). Furthermore, we found that in a small proportion of cells
that developed spontaneous SGs, opto-PR assemblies surrounded SGs (Fig.4C). Opto-PR
redistribution to the cytoplasm was not due to the nuclear membrane damage/nuclear pore
complex disruption, since nuclear retention of several RBPs was not affected in these cells
(Fig.4D). Therefore, impaired nuclear import of opto-PR under these conditions could be due to
its submicroscopic oligomerisation in the cytoplasm. Strikingly, endogenous TDP-43 but not
other ALS-related RBPs (FUS, NONO, SFPQ) were found to be enriched in the cytoplasmic
opto-PR foci (Fig.4D). In contrast, Cry2olig-only cytoplasmic structures induced by 24-h light

stimulation were negative for TDP-43 (Fig.4E). SRM revealed that cytoplasmic opto-PR
assemblies were ~250 nm structures that, unlike nuclear condensates, were not hollow
(Fig.4F). It also revealed that in these assemblies, opto-PR and TDP-43 remained demixed,
with the TDP-43 signal primarily on the surface (Fig.4F). A small fraction of cytoplasmic opto-
PR foci were positive for p62 (Fig.4G) but none of them stained positive for ubiquitin (data not
shown).

We next examined whether opto-PR accumulated/aggregated in the cytoplasm after prolonged
opto-stimulation affects stress-induced SGs (Boeynaems *et al*., 2017; Marmor-Kollet *et al*,
2020; Wen *et al*, 2014). Opto-PR and Cry2olig-only expressing cells subjected to long opto-
stimulation or kept in the dark were analysed during the recovery from NaAsO$_2$ (3 h time-point),
using G3BP1 as a marker. SG disassembly was delayed in light-stimulated opto-PR expressing
cells compared to unstimulated opto-PR cells or Cry2-only expressing (stimulated or
unstimulated) cells (Fig.4H). This finding further validates our cellular model as capable of
reproducing the key molecular effects of R-DPRs on the cellular RNA/RNP granule metabolism.

Therefore, our opto-model is amenable to the induction of cytoplasmic poly-PR accumulation
and aggregation, with its cytoplasmic assemblies being significantly more persistent than those
in the nucleus. Our data also point to a role for SGs in the growth of cytoplasmic poly-PR
aggregates as well as a role for poly-PR assemblies in "nucleating" TDP-43 aggregation in the
cytoplasm.

Discussion

A vast body of knowledge on C9-DPR related disease mechanisms has accumulated in the
past decade yet it remains unclear whether, and if so how, DPR self-assembly, resulting in a
C9-ALS/FTD hallmark pathology, contributes to the disease. Aggregation intermediates, and
especially smaller, highly reactive oligomeric species, have been validated as toxic/pathogenic
in the case of other neurodegeneration-linked proteins, such as tau and alpha-synuclein
(Bengoa-Vergniory *et al*, 2017; Choi & Gandhi, 2018; Jucker & Walker, 2018). It is possible that
equivalent DPR aggregation products play a role in C9-ALS/FTD. Poly-PR is the least abundant
DPR in human tissue (Mackenzie et al., 2015; Davidson et al., 2016) despite theoretically it
should be expressed at the levels similar to other DPRs (except poly-GP produced from both
strands). Although this can be due to a variation in the expression mechanisms and antibody
detection, it is also possible that neurons that accumulate poly-PR are lost early in disease due
to its high toxicity, including its aggregation products. Although the inclusions of all five DPRs in
the patient CNS are morphologically similar (Gendron *et al*., 2013), DPRs other than GA fail to
form microscopically visible assemblies in cell models (Frottin *et al*., 2021; Liu *et al*., 2022; Zhou
et al, 2017). In order to circumvent the high solubility of R-DPRs in cells, we harnessed the
Cry2olig opto-module (Taslimi *et al*., 2014) successfully used previously to promote self-
association of neurodegeneration-linked proteins (Berard *et al*, 2022; Jiang *et al*, 2021; Mann *et
al*, 2019). In line with the molecular dynamics predictions (Zheng *et al*., 2021), poly-GR's self-
association capacity was too low even when facilitated by Cry2olig, not yielding visible
condensation in cells. However we succeeded in achieving the condensation of the
oligomerisation-competent poly-PR.

Two key phenotypes – alterations to the nucleolus and SGs – validate our opto-PR model in the
context of the existing literature (Kwon *et al*., 2014; Marmor-Kollet *et al*., 2020; White *et al*.,
2019). In our model, SG disassembly was impaired by cytoplasmic opto-PR (but not Cry2olig-
only). This can be caused by altered composition, and hence dynamics, of SGs formed in the
opto-PR rich milieu. We also observed sequestration of NPM1, a confirmed R-DPR interactor,
into nuclear opto-PR condensates. Beyond these two phenotypes however, we failed to detect

the ubiquitous MLO disruption by poly-PR reported in some studies, and its condensation also
had limited effect on MLOs. This may be due to high variability of these phenotypes depending
on the cell type, DPR levels and repeat length. It would be important to establish consensus
phenotypes that can be used for the benchmarking of novel cellular DPR models, and the
nucleolar and SG pathology validated in multiple studies are the prime candidate readouts.

We found that poly-PR confers a specific ordered arrangement to the opto-induced nuclear
condensates – sphere with hollow centre. Arginine-rich peptides form such anisotropic
structures in the presence of RNA *in vitro*, both under RNA and peptide excess conditions
(Alshareedah *et al.*, 2020). Adopting this model, we speculate that upon interaction with cellular
RNA, the small opto-PR oligomers nucleated with the aid of Cry2olig, form nanocondensates
with a neutral "head" and a charged "tail" that subsequently coalesce into RNA-coated micelles
(~100 nm granules). These transition into a vesicle-like conformation "layered" with RNA on
both the internal and external surface (~250 nm granules). Upon (r)RNA depletion, these
condensates undergo fusion into a larger (>500 nm) hollow-centre structure (Fig.S5A).
Anisotropic condensate is a non-equilibrium state that may require ATP to be established and
maintained (Bergmann *et al*, 2023), and in our system, this state is fuelled by the light-induced
Cry2olig oligomerisation. Poly-PR was shown to form nuclear structures in C9-ALS/FTD,
varying from compact inclusions to less dense "territories", in multiple studies (Cooper-Knock et
al., 2015; Mori et al., 2013; Davidson et al., 2016; Wen et al., 2014). Nuclear poly-PR
condensates not overlapping with nucleolar markers also form in transgenic mice (Zhang *et al.*,
2019). It would be interesting to establish whether the structures seen in mice adopt a hollow-
centre structure during their biogenesis. Anisotropic nuclear assemblies ("anisosomes") are
formed by acetylated TDP-43 in cultured cells and *in vivo* (Yu *et al*, 2021). RBPs complexed
with polyadenylated RNA form similar structures in cell models of spinal muscular atrophy
(Narcis *et al*, 2018). Furthermore, nuclear condensates of RNA-binding deficient TDP-43 and
ALS-linked CREST protein (Kukharsky *et al*, 2015) also possess this typical morphology
(Fig.S5B). Remarkably, DDX3X mutations causative of neurodevelopmental disorders also form
cytoplasmic hollow condensates, and those composed of an aggressive RNA-binding deficient
mutant display low recovery in FRAP (Owens *et al*, 2023). It is possible that specific changes to
the cellular metabolism in neurological disease, e.g. altered protein and RNA stoichiometries,
favour this assembly type. RNA facilitates LLPS of poly-PR *in vitro* (Balendra et al., 2023)
however cellular RNA restricts the growth of the non-dynamic opto-PR condensates in cells.
Together with the reports on the solubilising effect of RNA on RBPs (Grese *et al*, 2021;
Maharana *et al*, 2018; Mann *et al.*, 2019; Shelkovnikova *et al*, 2014a) and on the wide-spread
RNA degradation in ALS (Tank *et al*, 2018), our findings suggest than declining RNA levels may
be a common factor underlying protein aggregation across ALS subtypes including C9-ALS.

Transient nuclear TDP-43 condensation is a hallmark of stress response (Cohen *et al.*, 2015;
Wang *et al.*, 2020). We recently found that this molecular event leads to TDP-43 loss of function
and prolonged STMN2 depletion and is dysregulated by TDP-43 mutations (Huang, Ellis *et al.,*
2024, manuscript in revision). These phenotypes may become persistent with chronic or
repetitive stress, precipitating the disease. Here we show that poly-PR self-assembly causes
TDP-43 condensation in the absence of stress signalling, indicating the convergence between
C9-ALS and other ALS subtypes in this putative disease mechanism. Consistently, recent use
of a specific RNA aptamer revealed abnormal nuclear TDP-43 granulation in motor neurons in
ALS tissue (Spence et al., 2024).

Cytoplasmic poly-PR assemblies in our opto-model are structurally dissimilar to and more
persistent than the nuclear condensates, probably reflecting a different environment in the two
cellular compartments. This higher stability is consistent with a higher frequency of poly-PR
cytoplasmic inclusions in patients (Zu *et al.*, 2013). DPR inclusions are rarely found to co-

deposit with TDP-43 inclusions in the patient tissue, with clear anatomical region specificity (Al-
Sarraj et al, 2011; Davidson et al., 2016). Some histopathological studies found evidence of
DPR aggregation preceding the cytoplasmic TDP-43 pathology (Baborie *et al*, 2015; Vatsavayai
et al, 2016). TDP-43 joins the cytoplasmic opto-PR foci where the two proteins remained in two
phases. It is tempting to speculate that transient cytoplasmic poly-PR assemblies "seed" TDP-
43 pathology. SGs may play an equivalent role for cytoplasmic poly-PR inclusions,
concentrating the initial poly-PR assemblies in a confined space and promoting their
coalescence into a larger aggregate. Indeed, a recent study showed that cytoplasmic TDP-43
aggregates can be nucleated within SGs, as a separate phase, and left behind after SG
dissolution (Yan *et al*, 2024). Inclusions C9-ALS/FTD are likely composed of several DPRs, and
their co-expression leads to different phenotypes *in vivo* (West *et al.,* 2020). Opto-GP readily
undergoes cytoplasmic condensation in our model, enabling future opto-DPR co-aggregation
studies.

Limitations of the study. We used a relatively short repeat length (which nevertheless is in the
patient range), whereas the Cry2olig-mCherry is a large tag. However, multiple studies have
demonstrated that DPR repeat lengths of 30-1000 have identical subcellular localisation with
similar toxicity (Bennion Callister *et al.*, 2016; Kwon *et al.*, 2014; Miyagi *et al*, 2023; White *et al.*,
2019). We cannot exclude that longer repeats will undergo condensation more readily, which
should be tested in future studies. In addition, for this proof-of-principle study, non-neuronal
cells were used for the ease of expression and imaging. To address potential cell-specificity of
the phenotypes revealed in this study, the opto-PR model should be transferred into a (human)
neuronal system.

Method details

Plasmids

The CRY2olig-mCherry backbone vector was a gift from Chandra Tucker (Addgene plasmid #
60032) (Taslimi *et al.*, 2014). DPR plasmids were a gift from Kurt De Vos (Bauer *et al*, 2022).
Codon-optimised DPR sequences were cloned into the CRY2olig-mCherry vector using
standard techniques. These plasmids are available via Addgene. Plasmids for the expression of
GFP-tagged TDP-43, FUS, NONO, SFPQ and CREST (N-terminal tag, in pEGFP-C1 vector)
were generated previously (Kukharsky *et al.*, 2015; Shelkovnikova *et al* 2014b). TDP-43
K145Q and F147/149L in pEGFP-C1 vector were generated using PCR mutagenesis.

Cell culture, transfection and treatments

HeLa cells were obtained from ATCC via Sigma, cultured in Dulbecco's Modified Eagle
Medium/Nutrient Mixture F-12 (DMEM/F-12) supplemented with 10% foetal bovine serum (FBS)
and penicillin-streptomycin. For time-lapse imaging, cells were plated on PhenoPlate-96 (black,
optically clear bottom, PerkinElmer) at a density of 2×10^4. For all other experiments, cells were
seeded in 24 well plates, with or without coverslips dependent on the application, at a density of
5×10^4 cells, unless otherwise stated. Transfection was performed 24 h prior to blue light
stimulation, using either Lipofectamine 2000 (ThermoFisher) or jetPRIME (Jena Bioscience)
according to the manufacturer's instructions. For transcriptional inhibition cells were treated with
2.5 µg/ml actinomycin D or 5,6-dichloro-1-beta-D-ribofuranosylbenzimidazole (DRB) (both
Sigma). For inhibition of arginine methylation, cells were treated with 10 µM MS-023 or
EPZ015666 (both ApexBio). To induce cellular stress, cells were treated with 250 µM $NaAsO_2$
(Sigma). Incubation times for treatments within individual experiments are indicated in
respective Figure legends.

Opto-stimulation

Cells expressing opto-constructs were stimulated with a 488 nm laser on Opera Phenix HCS
(500 ms, 80% laser power) for live cell time-lapse imaging (under full environmental control
conditions), or on a custom-built LED blue-light array housed in a humidified incubator
maintained at 37℃ with 5 % CO_2. Cells were protected from light between experiments and
during fixation.

Immunocytochemistry and RNA-FISH

Immunocytochemistry was performed as described elsewhere (Shelkovnikova *et al*, 2018) using
the following commercially available antibodies: FBL (rabbit polyclonal, Proteintech, 16021-1-
AP); NPM1 (mouse monoclonal, Proteintech, 60096-1-Ig); UPF1 (rabbit polyclonal, Proteintech,
23379-1-AP); PNN (rabbit polyclonal, Proteintech, 18266-1-AP); coilin p80 (mouse monoclonal,
BD Biosciences, 612074); SMN (mouse monoclonal, BD Biosciences, 610647); G3BP1 (rabbit
polyclonal, Proteintech, 13057-2-AP); repeat-PR (rabbit polyclonal, Proteintech, 23979-1-AP);
TDP-43 (rabbit polyclonal C-terminal, Sigma); FUS (mouse monoclonal, Santa Cruz, [4H11], sc-
47711); NONO (rabbit polyclonal, Proteintech, 11058-1-AP); SFPQ (rabbit monoclonal, Abcam,
ab177149); ribosomal RNA (mouse monoclonal Y10b, NB100-662); p62/SQSTM1 (mouse
monoclonal, MAB8028R, R&D Biosystems); RNA-FISH using a Cy5-labelled $dT_{(30)}$ DNA
oligonucleotide probe (Sigma) was performed as described earlier (Shelkovnikova *et al*, 2017).

Microscopy

Conventional fluorescence microscopy was performed using 100x objective on an Olympus
BX57 upright microscope equipped with an ORCA-Flash 4.0 camera (Hamamatsu) and
cellSens Dimension software (Olympus). Super-resolution microscopy was performed using a
63x oil immersion objective on a ZEISS 980 laser scanning confocal microscope (LSM) with
Airyscan 2 detector and ZEN Blue software. Time-lapse microscopy was performed using a 40x
objective on Opera Phenix HCS, and Harmony 4.9 software was used for image processing and
analysis (all PerkinElmer). Image processing and profile drawing was done using Image J or
ZEN Blue software. Condensate/aggregate quantification was done manually or on Image J in a
blinded manner.

Fluorescent recovery after photobleaching (FRAP)

Cells seeded at a density of 2.8 x 10^5 in glass-bottomed 35mm dishes (Ibidi), were transfected
and 24 h post-transfection, subjected to 3-hr continuous stimulation prior to FRAP analysis.
Imaging was performed using a 63x oil immersion objective on a ZEISS LSM 800 confocal
microscope, equipped with a humidified incubation chamber maintained at 37℃ with 5 % CO_2.
FRAP acquisition was performed on condensates formed after 3 h of continuous stimulation. A
circular region of interest (ROI) around each condensate was bleached using a 568 nm laser at
100 % laser power. Images were captured pre-bleach, immediately following bleach and at x
200 ms intervals during recovery. The mean fluorescence intensity within the ROI was
determined for each image using ZEN blue software. Intensity values were corrected for
bleaching during imaging and normalised to pre-bleach intensity. Average values were plotted
and FRAP curves fitted using a one-phase association equation in GraphPad Prism 9 software.

Electrophoretic mobility shift assay (EMSA)

Cy5-labeled RNA oligonucleotide 5'- GAGAGAGCGCGUGCAGAGACUUGGUGGUGCAUAA-3'
("Clip34nt") (Bhardwaj *et al.*, 2013) was custom-synthesised by Eurofins. Poly-GR_{x10} and poly-
PR_{x10} were custom-made by Biosynth. Lyophilised peptides were resuspended in water at 50
mM and stored at -80℃ before use. RNA was incubated at 250nM with 250nM-10µM peptides
in the EMSA buffer (50 mM Tris-HCl, pH 7.5, 100 mM KCl, 2 mM $MgCl_2$, 100 mM β-
mercaptoethanol, 0.1 mg/ml BSA) for 15 min at RT with gentle shaking. Samples were analysed

on 6% native acrylamide gel in TBE buffer, followed by imaging on Licor Odissey FC (700nm
channel).

Cell lysis, fractionation, and western blotting

For fractionation, cells were lysed in 1 % Titon-X-100 in PBS for 5 min at RT. Lysates were
incubated on ice for 30 minutes with vortexing at 5 min intervals and centrifuged at 13,000rpm
for 20 min at 4℃. Supernatant and pellet were mixed with 2x Laemmli buffer (2xLB), vortexed
and heated at 95℃ for 10 min. For western blotting, proteins were resolved on a 10% Mini-
PROTEAN® TGX™ hand-cast protein gel and transferred to a PVDF membrane. Gels were
stained with Gelcode™ (ThermoFisher) post-transfer for total protein. Membranes were blocked
for 1 h in 4 % milk/TBST and then incubated with the following primary antibodies (1:1000
dilution) at 4℃ overnight: mCherry (rabbit polyclonal, Proteintech, 26765-1-AP); GFP (mouse
monoclonal, 66002-1-Ig, Proteintech); TDP-43 (rabbit polyclonal, C-terminal, Sigma); eIF2α
(rabbit polyclonal, Cell Signaling, 9722), eIF2αP (rabbit polyclonal, Abcam, antibody [E90],
ab32157); repeat-PR (rabbit polyclonal, Proteintech, 23979-1-AP). Following primary antibody
incubation, blots were washed with TBST and incubated with an appropriate HRP secondary
antibody (GE Healthcare) for 1 h at room temperature. Signal was detected using Clarity Max
Western ECL Substrate (Bio-Rad) and imaged and quantified using Licor Odissey FC/Image
Studio software.

RNA expression analysis

481 Total RNA was extracted using GenElute total mammalian RNA kit (Sigma) in accordance with
482 the manufacturer's instructions. First-strand cDNA synthesis was performed using 500 ng of
483 RNA with random primers (ThermoFisher) and MMLV reverse transcriptase (Promega). qRT-
484 PCR was performed using qPCRBIO SyGreen Lo-ROX (PCRbio), and GAPDH was used for
485 normalisation. Primer sequences are provided in a previous study (Shelkovnikova *et al.*, 2017).

In vitro analysis of TDP-43 condensation

In vitro TDP-43 clustering analysis with immunodetection and imaging were performed as
described in Hodgson, Huang et al., 2024 (doi: 10.2139/ssrn.4721338). Briefly, 1 μM total
recombinant TDP-43 (R&D Biosystems, AP-190-100) or supernatant (after centrifuging at
13,300 rpm for 1 min) was mixed with 1 μM poly-PR/GR peptides (as above, BioSynth) or a
generic peptide (V5: GKPIPNPLLGLDST, Proteintech) in the assay buffer, and incubated for 10
min. Samples were sedimented and fixed with glutaraldehyde on coverslips, blocked with 1%
BSA in PBS for 1 h at RT and incubated with an anti-TDP-43 antibody (1:5000, mouse
monoclonal, R&D Biosystems, MAB77781) in the blocking buffer for 2 h. After washes, TDP-43
protein clusters were visualised using anti-mouse AlexaFluor488 antibody (1:2000,
ThermoFisher), incubated for 1 h at RT. Coverslips were mounted on a glass slide using Immu-
mount (ThermoFisher). Images were taken using Olympus BX57 upright microscope and
ORCA-Flash 4.0 camera and processed using cellSens Dimension software (Olympus).
Quantification of assemblies was done using the 'Analyze particles' tool of Image J. For
confirmatory western blot analysis, recombinant TDP-43 samples incubated with or without the
peptides as above (for 10 min, and an additional set – for 4 h) were centrifuged at 1,000xg for
10 min. Pellet and supernatant were analysed by western blot using a C-terminal TDP-43
antibody (Sigma).

Statistical analysis

Analysis was done using respective tests on GraphPad Prism 9 software. N corresponds to the
number of biological replicates. Error bars represent S.D. unless indicated otherwise.

Author contributions

Rachel E. Hodgson: Conceptualization; Data curation; Formal analysis; Investigation;
Methodology; Writing - original draft; Writing - review and editing. **Jessica Rayment:** Formal
analysis; Investigation; Methodology. **Wan-Ping Huang**: Formal analysis; Investigation;
Methodology; Writing - review and editing. **Anna Sanchez Avila**: Data curation; Investigation.
Tatyana A. Shelkovnikova: Conceptualization; Supervision; Funding acquisition; Project
administration; Writing - review and editing.

Disclosure and competing interests statement

516 The authors declare no competing interests.

Acknowledgements

518 This work was supported by the UKRI Future Leaders Fellowship (MR/W004615/1), MRC
519 (MR/W028522/1) and BBSRC (BB/V014110/1) standard grants, and MND Association
520 fellowship/grant (Shelkovnikova/Oct17/968-799). We also acknowledge the MRC grant
521 MR/X012077/1 for Airyscan 2 LMF.

References

Al-Sarraj S, King A, Troakes C, Smith B, Maekawa S, Bodi I, Rogelj B, Al-Chalabi A, Hortobagyi T, Shaw CE (2011) p62 positive, TDP-43 negative, neuronal cytoplasmic and intranuclear inclusions in the cerebellum and hippocampus define the pathology of C9orf72-linked FTLD and MND/ALS. *Acta neuropathologica* 122: 691-702

Alshareedah I, Moosa MM, Raju M, Potoyan DA, Banerjee PR (2020) Phase transition of RNA-protein complexes into ordered hollow condensates. *Proc Natl Acad Sci U S A* 117: 15650-15658

Ash PE, Bieniek KF, Gendron TF, Caulfield T, Lin WL, Dejesus-Hernandez M, van Blitterswijk MM, Jansen-West K, Paul JW, 3rd, Rademakers R *et al* (2013) Unconventional translation of C9ORF72 GGGGCC expansion generates insoluble polypeptides specific to c9FTD/ALS. *Neuron* 77: 639-646

Baborie A, Griffiths TD, Jaros E, Perry R, McKeith IG, Burn DJ, Masuda-Suzukake M, Hasegawa M, Rollinson S, Pickering-Brown S *et al* (2015) Accumulation of dipeptide repeat proteins predates that of TDP-43 in frontotemporal lobar degeneration associated with hexanucleotide repeat expansions in C9ORF72 gene. *Neuropathol Appl Neurobiol* 41: 601-612

Balendra R, Ruiz de Los Mozos I, Odeh HM, Glaria I, Milioto C, Wilson KM, Ule AM, Hallegger M, Masino L, Martin S *et al* (2023) Transcriptome-wide RNA binding analysis of C9orf72 poly(PR) dipeptides. *Life Sci Alliance* 6

Bauer CS, Webster CP, Shaw AC, Kok JR, Castelli LM, Lin YH, Smith EF, Illanes-Alvarez F, Higginbottom A, Shaw PJ *et al* (2022) Loss of TMEM106B exacerbates C9ALS/FTD DPR pathology by disrupting autophagosome maturation. *Front Cell Neurosci* 16: 1061559

Bengoa-Vergniory N, Roberts RF, Wade-Martins R, Alegre-Abarrategui J (2017) Alpha-synuclein oligomers: a new hope. *Acta neuropathologica* 134: 819-838

Bennion Callister J, Ryan S, Sim J, Rollinson S, Pickering-Brown SM (2016) Modelling C9orf72 dipeptide repeat proteins of a physiologically relevant size. *Human molecular genetics* 25: 5069-5082

Berard M, Sheta R, Malvaut S, Rodriguez-Aller R, Teixeira M, Idi W, Turmel R, Alpaugh M, Dubois M, Dahmene M *et al* (2022) A light-inducible protein clustering system for in vivo analysis of alpha-synuclein aggregation in Parkinson disease. *PLoS Biol* 20: e3001578

Bergmann AM, Bauermann J, Bartolucci G, Donau C, Stasi M, Holtmannspotter AL, Julicher F, Weber CA, Boekhoven J (2023) Liquid spherical shells are a non-equilibrium steady state of active droplets. *Nat Commun* 14: 6552

Bhardwaj A, Myers MP, Buratti E, Baralle FE (2013) Characterizing TDP-43 interaction with its RNA targets. *Nucleic Acids Res* 41: 5062-5074

Boeynaems S, Bogaert E, Kovacs D, Konijnenberg A, Timmerman E, Volkov A, Guharoy M, De Decker M, Jaspers T, Ryan VH *et al* (2017) Phase Separation of C9orf72 Dipeptide Repeats Perturbs Stress Granule Dynamics. *Molecular cell* 65: 1044-1055 e1045

Chan-Penebre E, Kuplast KG, Majer CR, Boriack-Sjodin PA, Wigle TJ, Johnston LD, Rioux N, Munchhof MJ, Jin L, Jacques SL *et al* (2015) A selective inhibitor of PRMT5 with in vivo and in vitro potency in MCL models. *Nat Chem Biol* 11: 432-437

Chew J, Cook C, Gendron TF, Jansen-West K, Del Rosso G, Daughrity LM, Castanedes-Casey M, Kurti A, Stankowski JN, Disney MD *et al* (2019) Aberrant deposition of stress granule-resident proteins linked to C9orf72-associated TDP-43 proteinopathy. *Mol Neurodegener* 14: 9

Choi ML, Gandhi S (2018) Crucial role of protein oligomerization in the pathogenesis of Alzheimer's and Parkinson's diseases. *Febs J* 285: 3631-3644

Choi SY, Lopez-Gonzalez R, Krishnan G, Phillips HL, Li AN, Seeley WW, Yao WD, Almeida S, Gao FB (2019) C9ORF72-ALS/FTD-associated poly(GR) binds Atp5a1 and compromises mitochondrial function in vivo. *Nat Neurosci* 22: 851-862

Cochard A, Garcia-Jove Navarro M, Piroska L, Kashida S, Kress M, Weil D, Gueroui Z (2022) RNA at the surface of phase-separated condensates impacts their size and number. *Biophys J* 121: 1675-1690

Cohen TJ, Hwang AW, Restrepo CR, Yuan CX, Trojanowski JQ, Lee VM (2015) An acetylation switch controls TDP-43 function and aggregation propensity. *Nat Commun* 6: 5845

Cook CN, Wu Y, Odeh HM, Gendron TF, Jansen-West K, Del Rosso G, Yue M, Jiang P, Gomes E, Tong J *et al* (2020) C9orf72 poly(GR) aggregation induces TDP-43 proteinopathy. *Sci Transl Med* 12

Cooper-Knock J, Higginbottom A, Stopford MJ, Highley JR, Ince PG, Wharton SB, Pickering-Brown S, Kirby J, Hautbergue GM, Shaw PJ (2015) Antisense RNA foci in the motor neurons of C9ORF72-ALS patients are associated with TDP-43 proteinopathy. *Acta neuropathologica* 130: 63-75

Davidson Y, Robinson AC, Liu X, Wu D, Troakes C, Rollinson S, Masuda-Suzukake M, Suzuki G, Nonaka T, Shi J et al (2016) Neurodegeneration in frontotemporal lobar degeneration and motor neurone disease associated with expansions in C9orf72 is linked to TDP-43 pathology and not associated with aggregated forms of dipeptide repeat proteins. *Neuropathol Appl Neurobiol* 42: 242-254

DeJesus-Hernandez M, Mackenzie IR, Boeve BF, Boxer AL, Baker M, Rutherford NJ, Nicholson AM, Finch NA, Flynn H, Adamson J *et al* (2011) Expanded GGGGCC hexanucleotide repeat in noncoding region of C9ORF72 causes chromosome 9p-linked FTD and ALS. *Neuron* 72: 245-256

Eram MS, Shen Y, Szewczyk M, Wu H, Senisterra G, Li F, Butler KV, Kaniskan HU, Speed BA, Dela Sena C *et al* (2016) A Potent, Selective, and Cell-Active Inhibitor of Human Type I Protein Arginine Methyltransferases. *ACS Chem Biol* 11: 772-781

Frottin F, Perez-Berlanga M, Hartl FU, Hipp MS (2021) Multiple pathways of toxicity induced by C9orf72 dipeptide repeat aggregates and G(4)C(2) RNA in a cellular model. *Elife* 10

Gendron TF, Bieniek KF, Zhang YJ, Jansen-West K, Ash PE, Caulfield T, Daughrity L, Dunmore JH, Castanedes-Casey M, Chew J *et al* (2013) Antisense transcripts of the expanded C9ORF72 hexanucleotide repeat form nuclear RNA foci and undergo repeat-associated non-ATG translation in c9FTD/ALS. *Acta neuropathologica* 126: 829-844

Gittings LM, Boeynaems S, Lightwood D, Clargo A, Topia S, Nakayama L, Troakes C, Mann DMA, Gitler AD, Lashley T *et al* (2020) Symmetric dimethylation of poly-GR correlates with disease duration in C9orf72 FTLD and ALS and reduces poly-GR phase separation and toxicity. *Acta neuropathologica* 139: 407-410

Grese ZR, Bastos AC, Mamede LD, French RL, Miller TM, Ayala YM (2021) Specific RNA interactions promote TDP-43 multivalent phase separation and maintain liquid properties. *EMBO Rep* 22: e53632

Hartmann H, Hornburg D, Czuppa M, Bader J, Michaelsen M, Farny D, Arzberger T, Mann M, Meissner F, Edbauer D (2018) Proteomics and C9orf72 neuropathology identify ribosomes as poly-GR/PR interactors driving toxicity. *Life Sci Alliance* 1: e201800070

Hodgson R, Huang WP, Kumar V, Haiyan A, Chalakova Z, Rayment J, Ellis BCS, Stender EGP, van Vugt, JJFA, Project MinE ALS Sequencing Consortium, Locker N, Pitout I, Fletcher S, Cooper-Knock J, Shelkovnikova T (2024) TDP-43 is a Master Regulator of Paraspeckle Condensation. Available at SSRN: https://ssrn.com/abstract=4721338 or http://dx.doi.org/10.2139/ssrn.4721338

Huang WP, Ellis BCS, Hodgson R, Sanchez Avila A, Kumar V, Rayment J, Moll T, Shelkovnikova TA. Stress-induced TDP-43 nuclear condensation leads to splicing loss-of-function and STMN2 depletion. *Manuscript in revision*

Jiang L, Lin W, Zhang C, Ash PEA, Verma M, Kwan J, van Vliet E, Yang Z, Cruz AL, Boudeau S *et al* (2021) Interaction of tau with HNRNPA2B1 and N(6)-methyladenosine RNA mediates the progression of tauopathy. *Molecular cell* 81: 4209-4227 e4212

Jucker M, Walker LC (2018) Propagation and spread of pathogenic protein assemblies in neurodegenerative diseases. *Nat Neurosci* 21: 1341-1349

Kanekura K, Harada Y, Fujimoto M, Yagi T, Hayamizu Y, Nagaoka K, Kuroda M (2018) Characterization of membrane penetration and cytotoxicity of C9orf72-encoding arginine-rich dipeptides. *Sci Rep* 8: 12740

Kukharsky MS, Quintiero A, Matsumoto T, Matsukawa K, An H, Hashimoto T, Iwatsubo T,
Buchman VL, Shelkovnikova TA (2015) Calcium-responsive transactivator (CREST) protein
shares a set of structural and functional traits with other proteins associated with amyotrophic
lateral sclerosis. *Mol Neurodegener* 10: 20

Kwon I, Xiang S, Kato M, Wu L, Theodoropoulos P, Wang T, Kim J, Yun J, Xie Y,
McKnight SL (2014) Poly-dipeptides encoded by the C9orf72 repeats bind nucleoli, impede
RNA biogenesis, and kill cells. *Science* 345: 1139-1145

LaClair KD, Zhou Q, Michaelsen M, Wefers B, Brill MS, Janjic A, Rathkolb B, Farny D,
Cygan M, de Angelis MH *et al* (2020) Congenic expression of poly-GA but not poly-PR in mice
triggers selective neuron loss and interferon responses found in C9orf72 ALS. *Acta
neuropathologica* 140: 121-142

Lee KH, Zhang P, Kim HJ, Mitrea DM, Sarkar M, Freibaum BD, Cika J, Coughlin M,
Messing J, Molliex A *et al* (2016) C9orf72 Dipeptide Repeats Impair the Assembly, Dynamics,
and Function of Membrane-Less Organelles. *Cell* 167: 774-788 e717

Lin Y, Mori E, Kato M, Xiang S, Wu L, Kwon I, McKnight SL (2016) Toxic PR Poly-
Dipeptides Encoded by the C9orf72 Repeat Expansion Target LC Domain Polymers. *Cell* 167:
789-802 e712

Liu F, Morderer D, Wren MC, Vettleson-Trutza SA, Wang Y, Rabichow BE, Salemi MR,
Phinney BS, Oskarsson B, Dickson DW *et al* (2022) Proximity proteomics of C9orf72 dipeptide
repeat proteins identifies molecular chaperones as modifiers of poly-GA aggregation. *Acta
Neuropathol Commun* 10: 22

Lopez-Gonzalez R, Lu Y, Gendron TF, Karydas A, Tran H, Yang D, Petrucelli L, Miller BL,
Almeida S, Gao FB (2016) Poly(GR) in C9ORF72-Related ALS/FTD Compromises
Mitochondrial Function and Increases Oxidative Stress and DNA Damage in iPSC-Derived
Motor Neurons. *Neuron* 92: 383-391

Mackenzie IR, Frick P, Grasser FA, Gendron TF, Petrucelli L, Cashman NR, Edbauer D,
Kremmer E, Prudlo J, Troost D *et al* (2015) Quantitative analysis and clinico-pathological
correlations of different dipeptide repeat protein pathologies in C9ORF72 mutation carriers.
Acta neuropathologica 130: 845-861

Maharana S, Wang J, Papadopoulos DK, Richter D, Pozniakovsky A, Poser I, Bickle M,
Rizk S, Guillen-Boixet J, Franzmann TM *et al* (2018) RNA buffers the phase separation
behavior of prion-like RNA binding proteins. *Science* 360: 918-921

Mann JR, Gleixner AM, Mauna JC, Gomes E, DeChellis-Marks MR, Needham PG,
Copley KE, Hurtle B, Portz B, Pyles NJ *et al* (2019) RNA Binding Antagonizes Neurotoxic
Phase Transitions of TDP-43. *Neuron* 102: 321-338 e328

Marmor-Kollet H, Siany A, Kedersha N, Knafo N, Rivkin N, Danino YM, Moens TG,
Olender T, Sheban D, Cohen N *et al* (2020) Spatiotemporal Proteomic Analysis of Stress
Granule Disassembly Using APEX Reveals Regulation by SUMOylation and Links to ALS
Pathogenesis. *Molecular cell* 80: 876-891 e876

Miyagi T, Ueda K, Sugimoto M, Yagi T, Ito D, Yamazaki R, Narumi S, Hayamizu Y, Uji IH,
Kuroda M *et al* (2023) Differential toxicity and localization of arginine-rich C9ORF72 dipeptide
repeat proteins depend on de-clustering of positive charges. *iScience* 26: 106957

Mizielinska S, Gronke S, Niccoli T, Ridler CE, Clayton EL, Devoy A, Moens T, Norona FE,
Woollacott IOC, Pietrzyk J *et al* (2014) C9orf72 repeat expansions cause neurodegeneration in
Drosophila through arginine-rich proteins. *Science* 345: 1192-1194

Moens TG, Niccoli T, Wilson KM, Atilano ML, Birsa N, Gittings LM, Holbling BV, Dyson
MC, Thoeng A, Neeves J *et al* (2019) C9orf72 arginine-rich dipeptide proteins interact with
ribosomal proteins in vivo to induce a toxic translational arrest that is rescued by eIF1A. *Acta
neuropathologica* 137: 487-500

Mori K, Weng SM, Arzberger T, May S, Rentzsch K, Kremmer E, Schmid B, Kretzschmar
HA, Cruts M, Van Broeckhoven C *et al* (2013) The C9orf72 GGGGCC repeat is translated into
aggregating dipeptide-repeat proteins in FTLD/ALS. *Science* 339: 1335-1338

Narcis JO, Tapia O, Tarabal O, Piedrafita L, Caldero J, Berciano MT, Lafarga M (2018) Accumulation of poly(A) RNA in nuclear granules enriched in Sam68 in motor neurons from the SMNDelta7 mouse model of SMA. *Sci Rep* 8: 9646

Ortega JA, Sasselli IR, Boccitto M, Fleming AC, Fortuna TR, Li Y, Sato K, Clemons TD, McKenna ED, Nguyen TP *et al* (2023) CLIP-Seq analysis enables the design of protective ribosomal RNA bait oligonucleotides against C9ORF72 ALS/FTD poly-GR pathophysiology. *Sci Adv* 9: eadf7997

Owens MC, Shen H, Yanas A, Mendoza-Figueroa MS, Lavorando E, Wei X, Shweta H, Tang HY, Goldman YE, Liu KF (2023) Mutant forms of DDX3X with diminished catalysis form hollow condensates that exhibit sex-specific regulation. *bioRxiv*

Park H, Kim NY, Lee S, Kim N, Kim J, Heo WD (2017) Optogenetic protein clustering through fluorescent protein tagging and extension of CRY2. *Nat Commun* 8: 30

Quaegebeur A, Glaria I, Lashley T, Isaacs AM (2020) Soluble and insoluble dipeptide repeat protein measurements in C9orf72-frontotemporal dementia brains show regional differential solubility and correlation of poly-GR with clinical severity. *Acta Neuropathol Commun* 8: 184

Renton AE, Majounie E, Waite A, Simon-Sanchez J, Rollinson S, Gibbs JR, Schymick JC, Laaksovirta H, van Swieten JC, Myllykangas L *et al* (2011) A hexanucleotide repeat expansion in C9ORF72 is the cause of chromosome 9p21-linked ALS-FTD. *Neuron* 72: 257-268

Saberi S, Stauffer JE, Jiang J, Garcia SD, Taylor AE, Schulte D, Ohkubo T, Schloffman CL, Maldonado M, Baughn M *et al* (2018) Sense-encoded poly-GR dipeptide repeat proteins correlate to neurodegeneration and uniquely co-localize with TDP-43 in dendrites of repeat-expanded C9orf72 amyotrophic lateral sclerosis. *Acta neuropathologica* 135: 459-474

Sakae N, Bieniek KF, Zhang YJ, Ross K, Gendron TF, Murray ME, Rademakers R, Petrucelli L, Dickson DW (2018) Poly-GR dipeptide repeat polymers correlate with neurodegeneration and Clinicopathological subtypes in C9ORF72-related brain disease. *Acta Neuropathol Commun* 6: 63

Schludi MH, May S, Grasser FA, Rentzsch K, Kremmer E, Kupper C, Klopstock T, German Consortium for Frontotemporal Lobar D, Bavarian Brain Banking A, Arzberger T *et al* (2015) Distribution of dipeptide repeat proteins in cellular models and C9orf72 mutation cases suggests link to transcriptional silencing. *Acta neuropathologica* 130: 537-555

Shav-Tal Y, Blechman J, Darzacq X, Montagna C, Dye BT, Patton JG, Singer RH, Zipori D (2005) Dynamic sorting of nuclear components into distinct nucleolar caps during transcriptional inhibition. *Mol Biol Cell* 16: 2395-2413

Shelkovnikova TA, Dimasi P, Kukharsky MS, An H, Quintiero A, Schirmer C, Buee L, Galas MC, Buchman VL (2017) Chronically stressed or stress-preconditioned neurons fail to maintain stress granule assembly. *Cell Death Dis* 8: e2788

Shelkovnikova TA, Kukharsky MS, An H, Dimasi P, Alexeeva S, Shabir O, Heath PR, Buchman VL (2018) Protective paraspeckle hyper-assembly downstream of TDP-43 loss of function in amyotrophic lateral sclerosis. *Mol Neurodegener* 13: 30

Shelkovnikova TA, Robinson HK, Southcombe JA, Ninkina N, Buchman VL (2014a) Multistep process of FUS aggregation in the cell cytoplasm involves RNA-dependent and RNA-independent mechanisms. *Human molecular genetics.* 23: 5211-26

Shelkovnikova TA, Robinson HK, Troakes C, Ninkina N, Buchman VL (2014b) Compromised paraspeckle formation as a pathogenic factor in FUSopathies. *Human molecular genetics* 23: 2298-2312

Shin Y, Berry J, Pannucci N, Haataja MP, Toettcher JE, Brangwynne CP (2017) Spatiotemporal Control of Intracellular Phase Transitions Using Light-Activated optoDroplets. *Cell* 168: 159-171 e114

Spence H, Waldron FM, Saleeb RS, Brown AL, Rifai OM, Gilodi M, Read F, Roberts K, Milne G, Wilkinson D, *et al* (2023) RNA aptamer reveals nuclear TDP-43 pathology is an early aggregation event that coincides with STMN-2 cryptic splicing and precedes clinical manifestation in ALS. *Acta Neuropathologica* 147:50.

Tank EM, Figueroa-Romero C, Hinder LM, Bedi K, Archbold HC, Li X, Weskamp K,
Safren N, Paez-Colasante X, Pacut C *et al* (2018) Abnormal RNA stability in amyotrophic lateral
sclerosis. *Nat Commun* 9: 2845

Taslimi A, Vrana JD, Chen D, Borinskaya S, Mayer BJ, Kennedy MJ, Tucker CL (2014)
An optimized optogenetic clustering tool for probing protein interaction and function. *Nat*
Commun 5: 4925

Van Mossevelde S, van der Zee J, Cruts M, Van Broeckhoven C (2017) Relationship
between C9orf72 repeat size and clinical phenotype. *Curr Opin Genet Dev* 44: 117-124

Vanneste J, Vercruysse T, Boeynaems S, Sicart A, Van Damme P, Daelemans D, Van
Den Bosch L (2019) C9orf72-generated poly-GR and poly-PR do not directly interfere with
nucleocytoplasmic transport. *Sci Rep* 9: 15728

Vatsavayai SC, Yoon SJ, Gardner RC, Gendron TF, Vargas JN, Trujillo A, Pribadi M,
Phillips JJ, Gaus SE, Hixson JD *et al* (2016) Timing and significance of pathological features in
C9orf72 expansion-associated frontotemporal dementia. *Brain* 139: 3202-3216

Wang C, Duan Y, Duan G, Wang Q, Zhang K, Deng X, Qian B, Gu J, Ma Z, Zhang S *et al*
(2020) Stress Induces Dynamic, Cytotoxicity-Antagonizing TDP-43 Nuclear Bodies via
Paraspeckle LncRNA NEAT1-Mediated Liquid-Liquid Phase Separation. *Molecular cell* 79: 443-
458 e447

Wen X, Tan W, Westergard T, Krishnamurthy K, Markandaiah SS, Shi Y, Lin S, Shneider
NA, Monaghan J, Pandey UB *et al* (2014) Antisense proline-arginine RAN dipeptides linked to
C9ORF72-ALS/FTD form toxic nuclear aggregates that initiate in vitro and in vivo neuronal
death. *Neuron* 84: 1213-1225

West RJH, Sharpe JL, Voelzmann A, Munro AL, Hahn I, Baines RA, Pickering-Brown S
(2020) Co-expression of C9orf72 related dipeptide-repeats over 1000 repeat units reveals age-
and combination-specific phenotypic profiles in Drosophila. *Acta Neuropathol Commun 8: 158*

White MR, Mitrea DM, Zhang P, Stanley CB, Cassidy DE, Nourse A, Phillips AH, Tolbert
M, Taylor JP, Kriwacki RW (2019) C9orf72 Poly(PR) Dipeptide Repeats Disturb Biomolecular
Phase Separation and Disrupt Nucleolar Function. *Molecular cell* 74: 713-728 e716

Xu L, Wang D, Zhao L, Yang Z, Liu X, Li X, Yuan T, Wang Y, Huang T, Bian N *et al*
(2023) C9orf72 poly(PR) aggregation in nucleus induces ALS/FTD-related neurodegeneration
in cynomolgus monkeys. *Neurobiol Dis* 184: 106197

Yan X, Kuster D, Mohanty P, Nijssen J, Pombo-Garcia K, Rizuan A, Franzmann TM,
Sergeeva A, Passos PM, George L *et al* (2024) Intra-condensate demixing of TDP-43 inside
stress granules generates pathological aggregates. *bioRxiv*

Yu H, Lu S, Gasior K, Singh D, Vazquez-Sanchez S, Tapia O, Toprani D, Beccari MS,
Yates JR, 3rd, Da Cruz S *et al* (2021) HSP70 chaperones RNA-free TDP-43 into anisotropic
intranuclear liquid spherical shells. *Science* 371

Zhang YJ, Guo L, Gonzales PK, Gendron TF, Wu Y, Jansen-West K, O'Raw AD, Pickles
SR, Prudencio M, Carlomagno Y *et al* (2019) Heterochromatin anomalies and double-stranded
RNA accumulation underlie C9orf72 poly(PR) toxicity. *Science* 363

Zheng S, Sahimi A, Shing KS, Sahimi M (2021) Molecular Dynamics Study of Structure,
Folding, and Aggregation of Poly-PR and Poly-GR Proteins. *Biophys J* 120: 64-72

Zhou Q, Lehmer C, Michaelsen M, Mori K, Alterauge D, Baumjohann D, Schludi MH,
Greiling J, Farny D, Flatley A *et al* (2017) Antibodies inhibit transmission and aggregation of
C9orf72 poly-GA dipeptide repeat proteins. *EMBO Mol Med* 9: 687-702

Zu T, Liu Y, Banez-Coronel M, Reid T, Pletnikova O, Lewis J, Miller TM, Harms MB,
Falchook AE, Subramony SH *et al* (2013) RAN proteins and RNA foci from antisense
transcripts in C9ORF72 ALS and frontotemporal dementia. *Proc Natl Acad Sci U S A* 110:
E4968-4977

Figure legends

Figure 1. An optogenetic cellular system for controllable *C9orf72* DPR condensation.

(A) Opto-DPR approach.

(B) Opto-DPR condensation can be induced by a single-pulse blue light stimulation. HeLa cells expressing the respective opto-DPR or empty Cry2olig-mCherry vector were analysed 24 h post-transfection. Blue-light array (single-pulse) was used. Scale bar 10 μm.

(C) Continuous opto-stimulation induces larger opto-PR condensates as compared to single light pulse. HeLa cells expressing opto-PR or empty vector were subjected to either a single-pulse or a 3-h continuous blue-light stimulation. 30 cells per condition were analysed from a representative experiment. **p<0.01, ****p<0.0001, Kruskal-Wallis test with Dunn's post-hoc test. Scale bar 10 μm.

(D) Super-resolution microscopy (SRM) demonstrates structural differences between Cry2olig-only and opto-PR assemblies and between the condensates formed after single-pulse and continuous (3-h long) blue-light stimulation. Representative images and graphical representation are shown.

(E) Poly-PR, detected using a PR-repeat antibody, is enriched within opto-PR condensate rims. Representative image is shown.

(F) Opto-PR condensate induction and tracking using confocal longitudinal imaging. Opto-PR expressing cells were stimulated with a 488 nm laser (at 80% for 500 ms), coupled with mCherry imaging. Cells were stimulated/imaged either every 2 min ("short-interval") or every 15 min for up to 4 h ("long-interval"). Alternatively, cells with preformed condensates were imaged every 2 min without stimulation ("dissolution"). Representative images are shown. Scale bar, 10 μm.

(G) Opto-PR condensate nucleation is concentration-dependent. mCherry fluorescence intensity was measured in the nucleoplasm of individual cells, outside the nucleolus, and the number of condensates was quantified in the same cells at the peak of their assembly (7-min interval repetitive stimulation for 49 min). 75 cells were analysed. Also see Fig.S1C.

(H) FRAP analysis after full opto-PR condensate photobleaching reveals low protein mobility between the condensate and nucleoplasm. Representative image and FRAP curve for 25 cells from a representative experiment are shown. Error bars represent SEM. Scale bar, 10 μm.

(I) Nuclear opto-PR condensates can become persistent. Opto-PR condensate were induced by 3-h continuous stimulation on blue-light array and the proportion of condensate-positive cells was quantified immediately post-stimulation or after 3 h of recovery in the dark. N=3 (300 cells analysed in total).

(J) Opto-PR condensates are positive for nucleophosmin (NPM1). Opto-PR condensates induced by a 3-h continuous opto-stimulation were analysed by SRM. Representative images and profile plots are shown. Scale bar, 2 μm.

Figure 2. RNA limits opto-PR condensation in the nucleus.

(A) Electrophoretic mobility shift assay (EMSA) with a natural GA-rich RNA sequence reveals R-DPR binding to RNA. Cy5-labelled synthetic nucleotide "Clip34nt" (34-mer) and synthetic poly-PR and poly-GR peptides (10 repeats) were used. Representative gel is shown.

(B) RNA depletion promotes opto-PR condensate growth. Opto-PR expressing cells were pre-treated with actinomycin D or DRB for 1 h, followed by long-interval repetitive blue-light stimulation (every 15 min) coupled with time-lapse imaging for up to 4 h (in the presence of the inhibitor). Representative images are shown. Scale bar, 10 μm.

(C) Opto-PR condensates formed under conditions of actinomycin D-induced RNA depletion
are larger in size and less numerous than in RNA-sufficient cells. Cells were opto-stimulated for
3 h continuously. Representative images and quantification are shown. Note that the larger
condensates remain FBL-negative. 30 and 60 cells per condition were included in analysis for
condensate size and number, respectively, from a representative experiment. **p<0.01,
***p<0.001, Student's *t* test. Scale bar, 5 µm.

(D) Opto-PR condensates formed in RNA-depleted conditions are more persistent. Opto-PR
expressing cells were light-stimulated for 3 h continuously, with or without DRB addition,
followed by DRB removal and recovery for 2 h in the dark. Representative images and
quantification are shown. N=3 (150 cells analysed in total). **p<0.01, Mann-Whitey *U* test. Scale
bar, 10 µm.

(E) Opto-PR condensates formed in actinomycin D-treated cells are structurally different, as
revealed by SRM. Representative images of condensates of a comparable size from control,
DRB- or actinomycin D-treated cells induced by 3-h continuous opto-stimulation are shown,
alongside with graphical representation.

(F) Ribosomal (r)RNA depletion from the nucleus and opto-PR condensates in actinomycin D
treated cells. Representative images are shown. Scale bar, 5 µm.

Figure 3. Nuclear opto-PR condensation induces TDP-43 pathology.

(A,B) Opto-PR self-assembly induces nuclear condensation of co-expressed TDP-43 GFP.
Cells were opto-stimulated for 3 h continuously. Representative images (A) and quantification
(B) are shown. Scale bar, 10 µm. N=3 (≥100 cells analysed per biological repeat). **p<0.01,
***p<0.001, mixed-effects analysis with Sidak's multiple comparisons test.

(C) Spatial proximity of opto-PR and TDP-43 GFP nuclear condensates as revealed by SRM.
Representative image is shown.

(D) Reduced TDP-43 GFP solubility upon induction of opto-PR condensation. Cells expressing
opto-PR were subjected to 3-h continuous blue-light stimulation before lysis and fractionation.
Western blots and quantification of the relative TDP-43 GFP amount in the supernatant fraction
are shown for a representative experiment. S, supernatant; P, pellet.

(E) TDP-43 condensates induced by opto-PR self-assembly are devoid of polyA+ RNA. Cells
after 3-h continuous opto-stimulation were analysed by poly(dT) RNA-FISH. Representative
image is shown. Scale bar, 5 µm.

(F,G) Opto-PR self-assembly promotes nuclear condensation of a TDP-43 acetylation mimic
mutant (K145Q). Representative images (F) and quantification (G) are shown. 50 cells were
analysed from a representative experiment. *p<0.05, **p<0.01, Student's *t* test. Scale bar, 5
µm.

(H-J) R-DPRs promote TDP-43 clustering *in vitro.* Recombinant TDP-43 (supernatant fraction
depleted of large aggregates; 1 µM) was incubated with equimolar amounts of poly-PR$_{(x10)}$ or
poly-GR$_{(x10)}$ peptides and analysed by immunostaining and fractionation/western blot.
Methodology (H), immunostaining/imaging (I) and fractionation/western blotting (J) data are
shown. In I, 10 min incubation was used and five fields of view from a representative
experiment were analysed; ***p<0.001, ****p<0.0001, one-way ANOVA with Dunnett's post-hoc
test. In J, representative western blot and band intensity quantification for the supernatant are
shown. S, supernatant; P, pellet. Scale bar, 20 µm.

Figure 4. Prolonged opto-stimulation elicits cytoplasmic opto-PR/TDP-43 pathology.

(A,B) Prolonged, 24-h blue-light stimulation leads to cytoplasmic opto-PR redistribution and aggregation, with persistent assembly formation. Proportion of cells with opto-PR assemblies (nuclear and cytoplasmic) was quantified after 24-h blue-light array stimulation with and without 8-h recovery in the dark. Representative images (A) and quantification (B) are shown. Images for a 3-h stimulation and recovery are included for comparison. N=3 (300 cells analysed per condition). n.s., non-significant. Scale bar, 10 µm.

(C) Cytoplasmic opto-DPR foci surround spontaneous stress granules (SGs, visualised using G3BP1 as a marker). Representative image is shown. Cells were opto-stimulated for 24 h continuously. Scale bar, 5 µm.

(D) Endogenous TDP-43 but, not other RBPs, joins opto-PR assemblies induced by prolonged opto-stimulation. Also note normal nuclear localisation of all RBPs in cells with cytoplasmic opto-PR. Cells were opto-stimulated for 24 h continuously. Representative images are shown. Scale bars, 10 µm.

(E) Cry2only cytoplasmic assemblies are negative for TDP-43. Cells were opto-stimulated for 24 h continuously. Representative images are shown. Scale bars, 10 µm.

(F) TDP-43 remains demixed within opto-PR within cytoplasmic assemblies, as revealed by SRM. Cells were opto-stimulated for 24 h continuously. Representative image is shown.

(G) Cytoplasmic opto-PR assemblies are occasionally positive for p62. Representative image is shown. Insets 1 and 2 shows examples of p62-positive and -negative assemblies, respectively. Scale bars, 10 µm.

(H) SG dissolution is affected in cells with cytoplasmically localised opto-PR. SGs Representative were induced $NaAsO_2$ in cells expressing either opto-PR or Cry2olig-only, subjected to 24-h long opto-stimulation or kept in the dark. Efficiency of SG dissolution was analysed 3 h into the recovery (post-$NaAsO_2$ removal). Representative images and quantification are shown. N=3 (120 cells analysed per condition). *$p<0.05$, Two-way ANOVA with Sidak's multiple comparisons test. Scale bars, 10 µm.

900   **Figure 1**

901
902
903
904

905 **Figure 2.**

Figure 4.