C9orf72 poly-PR condensation induces nuclear TDP-43 pathology
and is inhibited by RNA in an optogenetic cell model

Hodgson RE*, Rayment J, Huang W-P, Sanchez Avila A, Shelkovnikova TA*

Sheffield Institute for Translational Neuroscience and Neuroscience Institute, University of
Sheffield, Sheffield, UK

Abstract: Proteinaceous inclusions formed by *C9orf72* derived dipeptide-repeat (DPR)
proteins are a histopathological hallmark in ~50% of familial amyotrophic lateral
sclerosis/frontotemporal dementia (ALS/FTD) cases. However DPR aggregation/inclusion
formation could not be efficiently recapitulated in cell models for four out of five DPRs. We
utilised the Cry2olig optogenetic module to achieve chemical-free and controllable poly-PR
condensation/aggregation in cultured cells. This approach revealed that poly-PR forms
anisotropic condensates in the nucleus and that RNA limits their fusion-dependent growth.
Poly-PR self-assembly induced nuclear TDP-43 condensation without activation of cellular
stress response. Poly-PR cytoplasmic redistribution and aggregation could be also achieved
with prolonged light stimulation. Cytoplasmic poly-PR assemblies were more persistent than
its nuclear condensates, selectively sequestered TDP-43 and surrounded spontaneous
stress granules. Our data suggest that poly-PR condensation in the nucleus and cytoplasm,
causative of TDP-43 dysfunction, may constitute an early pathological event in C9-ALS/FTD.
The opto-DPR platform described here is a useful tool for modelling cytopathologies elicited
by DPR aggregation for mechanistic research and drug discovery.

corresponding authors: r.hodgson1@sheffield.ac.uk, t.shelkovnikova@sheffield.ac.uk

Key words: ALS, FTD, C9orf72, DPR, poly-PR, optogenetics, condensate, TDP-43,
membraneless organelle (MLO)

Highlights:

- Optogenetics can be used to model *C9orf72* DPR condensation in cultured cells.
- Opto-PR forms hollow nuclear condensates, and RNA limits their growth by fusion.
- Opto-PR self-assembly leads to stress-independent nuclear TDP-43 condensation.
- Prolonged opto-stimulation induces persistent cytoplasmic opto-PR/TDP-43
assemblies.

Graphical abstract:

31
32

Introduction

A G_4C_2 hexanucleotide repeat expansion (HRE) in the first intron of the *C9orf72* gene is the most common genetic alteration associated with amyotrophic lateral sclerosis (ALS) and frontotemporal dementia (FTD) (DeJesus-Hernandez et al, 2011; Renton et al, 2011). Healthy individuals commonly carry 2 repeats, while C9-ALS/FTD is associated with ≥30 repeat lengths (Van Mossevelde et al, 2017). Production of dipeptide repeat (DPR) proteins from the *C9orf72*-HRE transcripts is one of the proposed mechanisms of repeat toxicity (Ash et al, 2013; Mori et al, 2013; Zu et al, 2013). Both sense and antisense C9orf72-HRE transcripts are translated in all reading frames via non-canonical, repeat-associated non-AUG (RAN) translation to produce five DPRs: poly-GA, -GR (sense), -PR, -PA (antisense) and -GP (both sense and antisense). All five DPRs have been detected in the patient CNS, primarily as cytoplasmic and less often nuclear inclusions in neurons and glia (Gendron *et al*, 2013; Mackenzie *et al*, 2015; Saberi *et al*, 2018; Sakae *et al*, 2018; Schludi *et al*, 2015). Whilst poly-GA and -GP are the most abundant, arginine containing DPRs (R-DPRs) poly-GR and poly-PR are the most toxic species in cell and *in vivo* models (Kwon *et al*, 2014; Mizielinska *et al*, 2014; Moens *et al*, 2019). For example, nuclear poly-PR aggregation was sufficient to induce ALS-like phenotypes in a non-human primate model (Xu *et al*, 2023), and expression of this DPR was extremely toxic in mice (LaClair *et al*, 2020; Zhang *et al*, 2019).

R-DPR inclusion pathology seen in patients and mouse models (Chew *et al*, 2019; Choi *et al*, 2019; Cook *et al*, 2020) is not easily reproducible in cultured cells, where R-DPRs typically display diffuse distribution (outside the nucleolus), even when overexpressed and independent of the repeat length; this is in contrast to poly-GA that readily forms large cytoplasmic aggregates (Frottin *et al*, 2021; Hartmann *et al*, 2018; Liu *et al*, 2022; Lopez-Gonzalez *et al*, 2016; Vanneste *et al*, 2019). High arginine content renders R-DPRs highly hydrophilic and hence soluble. Although poly-PR and poly-GR have similar biochemical properties, molecular dynamics simulations revealed that poly-PR is capable of (limited) self-association – forming either dimers or small amorphous oligomers, whereas poly-GR is not likely to form any stable oligomeric species (Zheng *et al*, 2021). Consistently, *in vitro,* R-DPRs undergo phase separation only in the presence of additional agents – RNA or proteins (Balendra *et al*, 2023; Boeynaems *et al*, 2017; Lee *et al*, 2016; Zhang *et al.*, 2019). This suggests that certain factors in the cellular environment drive R-DPR loss of solubility in ALS/FTD. Sensitive immunoassays revealed that soluble DPRs are less abundant in the brain regions more affected by neurodegeneration as compared to the relatively spared regions (Quaegebeur *et al*, 2020).

Intermediate products of R-DPR aggregation may contribute to cellular dysfunction, however due to difficulty of modelling this molecular event, their role could not be investigated in the cellular setting. We hypothesised that the use of optogenetics (Park *et al*, 2017; Shin *et al*, 2017) will allow circumventing the intrinsic solubility of R-DPRs, by triggering and maintaining an oligomerised R-DPR state. Indeed, using Cry2olig tagging, we were able to reliably induce condensation of poly-PR ("opto-PR") in cultured cells, with spatial and temporal control. This cellular model was validated by demonstrating NPM1 partitioning into nuclear opto-PR condensates and enhancement of stress granule assembly by cytoplasmic opto-PR. Using this model, we revealed that: i) RNA modulates poly-PR condensation in cells; ii) poly-PR self-assembly causes a TDP-43 nuclear condensation phenotype; iii) poly-PR assemblies are more stable in the cytoplasm than nucleus; iv) TDP-43 can co-deposit with cytoplasmic poly-PR assemblies in a demixed state. Thus, opto-induction of DPR

80 pathology is an accessible and straightforward approach to model patient-relevant
81 phenotypes in cultured cells for mechanistic studies and drug discovery.
82

83 Results

84 *Optogenetic modeling of C9orf72 poly-PR condensation in cells*

We utilised Cry2olig, a Cry2 variant with high oligomerisation capacity (Taslimi *et al*, 2014),
in an attempt to induce and maintain R-DPR self-association. 'Opto-DPR' constructs were
generated for codon-optimised expression of poly-PR, poly-GR and additionally poly-GP (36
repeats), tagged with Cry2olig and mCherry on the N-terminus (Fig.1A). Opto-DPRs
demonstrated a subcellular distribution pattern in HeLa cells typical for the 30-1000 repeat
range (Bennion Callister *et al*, 2016; Kanekura *et al*, 2018; Liu *et al*., 2022; Vanneste *et al*.,
2019), where poly-GR and -GP were predominantly cytoplasmic and poly-PR was
predominantly nuclear, with high enrichment in the nucleolus; none of the DPRs showed
signs of aggregation (Fig.1B). Upon single-pulse light stimulation on a custom blue-light
array, Cry2olig vector control, opto-PR and opto-GP but not opto-GR readily formed visible
clusters/foci (Fig. 1B), consistent with the low poly-GR self-association capacity reported in
simulation studies (Zheng *et al*., 2021). Opto-PR foci were small (<500 nm), dot-like and
exclusively nuclear, whereas opto-GP formed large amorphous cytoplasmic inclusions (Fig.
1B). We focused on opto-PR thereafter, using opto-GR as a control.

Nuclear opto-PR foci/condensates were negative for the two nucleolar markers tested,
fibrillarin (FBL) and UPF1, confirming that they are not merely fragments of the nucleolus
(Fig.S1A). Continuous 3-h blue-light stimulation led to an increased opto-PR condensate
size, as compared to single pulse (Fig.1C). By super-resolution microscopy (SRM), the
condensates induced by a single light pulse were found to represent a mixed population of
~100 nm dot-like and ~250 nm spherical, hollow-centre assemblies resembling anisotropic
vesicle-like structures formed by arginine-rich peptide/RNA mixtures *in vitro* (Alshareedah *et*
al, 2020). Furthermore, the larger condensates formed after 3-h continuous light exposure
were found to represent multiples of these spheroids (Fig.1D). Although Cry2olig protein on
its own also formed clusters throughout the cell in response to blue light (Fig. 1B) (Taslimi *et*
al., 2014), these structures appeared disordered/filamentous and were clearly different from
opto-PR condensates (Fig.1D). Staining with a PR-repeat specific antibody detected poly-PR
primarily within the rims of the hollow-centre spheroids (Fig.1E). Therefore, poly-PR confers
a specific architecture to light-inducible condensates.

We next set up an imaging approach for simultaneous induction and tracking of opto-PR
condensation on a confocal high-content imaging system. Even extremely short blue-light
exposures and low 488 nm laser power (50 ms/5%) were sufficient to induce opto-PR foci; a
combination of 500 ms exposure/80% laser power was used in subsequent experiments as
consistently and robustly inducing condensates (Fig.S1B). Visible opto-PR condensates
appeared within 2 min post-pulse and could be maintained by both short- and long-interval
repetitive opto-stimulation (every 2 min or every 15 min, respectively, Fig.1F). Condensates
were reversible, typically resolving within ~14 min after the last light pulse (Fig. 1F). Opto-PR
condensate nucleation was concentration-dependent, with more structures forming in higher-
expressing cells (R=0.65) (Fig.1G; Fig.S1C). FRAP analysis revealed limited dynamics of
opto-PR within these assemblies, with low recovery after photobleaching of the entire
structure, despite a significant amount of diffuse opto-PR in the nucleoplasm (Fig.1H). A

fraction of cells (22.0±8.7%) developed persistent opto-PR condensates after continuous
light stimulation, which were still detectable after 3 h of recovery in the dark (Fig.1I).

Poly-PR was previously shown to interact with nucleophosmin (NPM1) and to co-partition
with this protein into phase-separated droplets *in vitro* (Lee et al., 2016; White et al, 2019).
Consistent with this, opto-PR condensates stained positive for NPM1, where NPM1 was
found primarily on the condensate surface, suggesting its secondary recruitment (Fig.1J).
Interestingly, opto-stimulation induced opto-PR signal segregation in the nucleolus
(Fig.S1D). In this, we observed intra-nucleolar demixing of opto-PR from NPM1, where the
proteins fully co-localised in the granular component (GC) under dark conditions but formed
two distinct phases after opto-stimulation (Fig.S1E,F). In contrast, opto-PR and FBL (the
latter residing in the nucleolar dense fibrillar component, DFC) showed no co-localisation
both under dark and light conditions (Fig.S1E,F).

R-DPRs were shown to promiscuously interact with membraneless organelle (MLO)
components – RNA and proteins with low-complexity domains, leading to wide-spread MLO
dysfunction (Kwon *et al.*, 2014; Lee *et al.*, 2016; Lin *et al*, 2016; Liu *et al.*, 2022). We
investigated the effect of opto-PR and its condensation on MLOs, focusing on those in the
nucleus due to the predominantly nuclear localisation of this DPR. Systematic analysis of
four MLOs – nuclear bodies (Gems, Cajal bodies, paraspeckles and speckles) revealed only
minor changes in the number and size in the presence of diffuse or condensed opto-PR
(Fig.S2). Cytoplasmic stress granules (SGs) were not induced by opto-PR with or without
blue-light stimulation (3 h continuous), consistent with its mainly nuclear localisation (data
not shown).

Thus, microscopically visible DPR self-assembly/condensation in cultured cells can be
achieved using Cry2olig tagging, allowing the formation of DPR-specific, ordered
assemblies. Opto-PR condensates are characterised by concentration-dependent growth
and low dynamic properties associated with persistence, and sequester NPM1.

RNA limits poly-PR condensate growth in cells

We next asked whether our opto-model can be used to characterise potential modifiers of
poly-PR condensation. RNA was previously found to promote R-DPR phase separation *in*
vitro (Balendra *et al.*, 2023; Boeynaems *et al.*, 2017; Gittings *et al*, 2020). More recently,
using RNA-protein crosslinking, R-DPRs have been shown to bind RNA in cells, in particular
ribosomal RNA (rRNA), with a preference for GA-rich sequences (Balendra *et al.*, 2023;
Ortega *et al*, 2023). Using electrophoretic mobility shift assay (EMSA) with an RNA
oligonucleotide representing a naturally occurring RNA sequence containing 5xGA repeats
(Clip34nt) (Bhardwaj *et al*, 2013), we indeed observed that synthetic poly-PR and -GR
peptides (10-mers) form complexes with RNA (Fig.2A).

Having confirmed RNA binding, we next studied the impact of RNA depletion on opto-PR
condensates. Opto-PR expressing cells were treated with two transcriptional blockers, a
global inhibitor (actinomycin D) and an RNA polymerase II-specific inhibitor (5,6-Dichloro-1-
β-D-ribofuranosylbenzimidazole, DRB), followed by induction and tracking of opto-PR
condensates in individual cells for up to 4 h (long-interval repetitive stimulation, in the
presence of the inhibitor). Both inhibitors caused dramatic nucleolar shrinking confirming
their activity in cells (Fig.2B) (Shav-Tal *et al*, 2005). Opto-PR condensates formed in RNA-
depleted conditions appeared noticeably larger compared to control (Fig.2B). Similar result
was obtained with a 3-h continuous opto-stimulation on blue-light array, and quantification

confirmed a significant increase in the opto-PR condensate size (Fig.2C). These larger
structures remained negative for the nucleolar marker FBL (Fig.2C). DRB is a reversible
inhibitor, which allowed us to analyse possible changes in the stability of RNA-depleted opto-
PR condensates. Cells were stimulated for 3 h in the presence or absence of DRB followed
by 2 h of recovery in the dark. Opto-PR condensates formed in the presence of DRB
appeared significantly more persisting, still detectable following the recovery in 63±6.4%
cells, compared to 30±7.8% cells in control condition (Fig.2D).

Increase in the opto-PR condensate size upon actinomycin D treatment was accompanied
by a decrease in their number (Fig.2C), suggesting clustering or fusion. SRM analysis
revealed that opto-PR condensates in actinomycin D treated cells were no longer clusters of
individual ~250 nm spherical units seen in the untreated or DRB condition but instead
represented larger (>500 nm) hollow-centre spheres (Fig.2E). RNA on the surface of
condensates has been shown to limit their fusion, including in cells (Cochard *et al*, 2022).
Actinomycin D but not DRB depletes ribosomal RNA (rRNA), and we observed dense rRNA
signal on the opto-PR condensate surface which disappeared after actinomycin D treatment
(Fig.2F). Therefore, the opto-PR condensates in actinomycin D treated cells may form by
fusion of the smaller units due rRNA depletion from the surface, followed by relaxation into a
larger spherical assembly.

Another putative modifier of R-DPR self-association/condensation is arginine dimethylation
(DMA), which increases the fluidity of R-DPR droplets *in vitro* and was found enriched in R-
DPR inclusions in C9-ALS/FTD patient tissue (Gittings *et al.*, 2020). We employed two small
molecule methyltransferase inhibitors, MS023 inhibiting five type-I methyltransferases that
synthesise asymmetric DMA (aDMA) and EPZ015666, a specific inhibitor of PRMT5
responsible for most symmetric DMA (sDMA) (Chan-Penebre *et al*, 2015; Eram *et al*, 2016).
Opto-PR expressing cells were treated with MS023 and EPZ015666 for 24 h and then
exposed to blue-light for 3 h continuously, followed by opto-PR condensate quantification.
We observed a mild decrease in the condensate number after aDMA depletion without
changes in their size or structure (Fig.S3). Thus, removing DMA marks may attenuate opto-
PR condensate nucleation, although this effect is small.

Therefore, our cellular opto-PR model can be utilised to analyse modifiers of poly-PR self-
assembly in the cellular context, as exemplified by the modulatory effect of RNA that we
have uncovered.

Poly-PR condensation induces nuclear TDP-43 pathology

R-DPR interactomes are enriched in RNA-binding proteins (RBPs) (Kwon *et al.*, 2014; Lee *et*
al., 2016). We therefore examined the effect of nuclear opto-PR condensation on ALS/FTD
relevant RBPs. TDP-43, FUS, NONO and SFPQ, all tagged with GFP, were co-expressed
with opto-PR and their subcellular distribution was examined with and without opto-
stimulation. Opto-PR presence *per se* did not affect RBP distribution (Fig.3A; Fig.S4A,B).
However, light-stimulated opto-PR expressing cells displayed a striking nuclear
condensation phenotype for TDP-43 but not other RBPs analysed (Fig.3A; Fig.S4A).
Although we did observe TDP-43 condensation in a fraction of light-stimulated Cry2olig-
expressing, it was significantly less than in opto-PR cultures; nor was it observed in light-
stimulated opto-GR expressing cells (Fig.3B). Opto-PR assemblies were frequently found in
the physical contact with TDP-43 condensates (44±0.5% of all opto-PR foci), suggestive of a
direct nucleating effect of oligomerising opto-PR (Fig.3C). Fractionation demonstrated
reduced solubility of TDP-43 GFP, alongside with endogenous TDP-43, after induction of

opto-PR condensation (Fig.3D). In contrast, opto-GR solubility was unaltered (Fig.3D),
consistent with lack of visible self-assembly (Fig.1B). TDP-43 GFP nuclear condensates
induced by opto-PR were strikingly similar to the condensates formed during the recovery
from arsenite stress (Wang *et al*, 2020) (Huang, Ellis, et al., manuscript in revision). Like
these stress-induced foci, TDP-43 condensates induced by opto-PR were devoid of polyA+
RNA (Fig.3E). We therefore asked whether TDP-43 condensation elicited by opto-PR was
due to an upregulated stress response. Classic cellular stress markers, phosho-eIF2α,
GADD34 and ATF4, were not altered in opto-PR expressing cells after 3-h light stimulation
(Fig.S4B,C). This is in contrast to their dramatic upregulation during the recovery from
sodium arsenite stress used as a positive control (Fig.S4C). In addition to demonstrating the
stress-unrelated nature of TDP-43 condensates induced by opto-PR, this also confirms a
lack of phototoxicity in our model.

TDP-43 acetylation was shown to impair its RNA binding and enhance aggregation, with
acetylated TDP-43 inclusions detected in sALS (Cohen *et al*, 2015). We investigated the
effect of opto-PR self-assembly on the acetylation-mimic TDP-43 mutant, K145Q (Cohen *et*
al., 2015). In agreement with the published data, TDP-43 K145Q was prone to
spontaneously forming nuclear foci/granules in naïve cells (Fig.S4D). Opto-PR condensation
potentiated the pro-aggregating effect of acetylation, further increasing nuclear granulation of
TDP-43 K145Q (Fig.3F,G).

We next examined the effect of R-DPRs on TDP-43 higher-order assembly *in vitro,* using the
condensate immunodetection/imaging assay we recently developed (Hodgson, Huang et al.,
2024, doi: 10.2139/ssrn.4721338) (Fig.3H). Both poly-PR and poly-GR peptides were
included in these studies. "Supernatant" fraction of recombinant TDP-43 (after removal of
preformed aggregates) containing small protein clusters and soluble protein was incubated
with equimolar amounts of synthetic R-DPR peptides, followed by sedimentation and fixation
of TDP-43 clusters on coverslips for immunostaining/imaging; in parallel, samples were
fractionated by centrifugation for western blot analysis (Fig.3H). Addition of both R-DPRs
significantly enhanced TDP-43 clustering – manifested as an increased cluster size and
decreased cluster number, with poly-GR being more potent (Fig. 3I). Fractionation confirmed
increased partitioning of TDP-43 to the pellet fraction in the presence of R-DPRs (Fig.3J).

Collectively, these results suggest that poly-PR condensation can directly cause changes to
the nuclear distribution of TDP-43 and its disease-linked variant, without activation of stress
signaling.

Cytoplasmic poly-PR assemblies are persistent and selectively sequester TDP-43

Having characterised the nuclear phenotypes, we asked whether our opto-PR model is
amenable to reproducing the cytoplasmic pathology typical for DPRs. Continuous 24-h long
stimulation on the blue-light array resulted in significant redistribution of opto-PR to the
cytoplasm, with cytoplasmic foci formation in 32% of cells (>300 transfected cells analysed;
Fig. 4A,B). This was accompanied by a reduction in the incidence of nuclear opto-PR
condensates (from 94% after 3 h to 9% after 24 h of stimulation) (Fig. 4A,B). Nuclear and
cytoplasmic opto-PR foci induced by 24-h opto-stimulation were persisting, with no
significant decline observed after 8 h of recovery in the dark (Fig. 4A,B). This is in contrast to
the nuclear opto-PR condensates forming after a 3-h stimulation that were largely cleared
after 3 h of recovery in the dark (Fig.4A; Fig.1I). Furthermore, we found that in a small
proportion of cells that developed spontaneous SGs, these were surrounded by opto-PR

assemblies (Fig.4C). Opto-PR redistribution to the cytoplasm was not due to the nuclear
membrane damage/nuclear pore complex disruption, since nuclear retention of several
RBPs was not affected in these cells (Fig.4D). One possible explanation is impaired nuclear
import of opto-PR due to submicroscopic oligomerisation in the cytoplasm. Strikingly,
endogenous TDP-43 but not other ALS-related RBPs (FUS, NONO, SFPQ) were found to be
enriched in the cytoplasmic opto-PR foci (Fig.4D). In contrast, Cry2olig-only cytoplasmic
structures induced by light were negative for TDP-43 (Fig.4E). SRM revealed that
cytoplasmic opto-PR assemblies were ~250 nm structures that, unlike nuclear condensates,
were not hollow (Fig.4F). Furthermore, in these assemblies, opto-PR and TDP-43 remained
demixed, with the TDP-43 signal primarily on the surface (Fig.4F).

We next examined whether opto-PR accumulated/aggregated in the cytoplasm after
prolonged opto-stimulation affects stress-induced SGs (Boeynaems *et al.*, 2017; Marmor-
Kollet *et al*, 2020; Wen *et al*, 2014). Opto-PR expressing cells subjected to long opto-
stimulation or kept in the dark were analysed during $NaAsO_2$ exposure (15 and 60 min) and
during the recovery from $NaAsO_2$ (3 h), using G3BP1 as a marker. We detected accelerated
SG assembly (15 min time-point) and delayed SG disassembly (3 h recovery time-point) in
light-stimulated compared to unstimulated cells (Fig.4G). This finding further validates our
opto-model as capable of reproducing the key molecular effects of R-DPRs on the cellular
RNA/RNP granule metabolism.

Therefore, our opto-model is amenable to the induction of cytoplasmic poly-PR accumulation
and aggregation, with its cytoplasmic assemblies being significantly more persistent than
those in the nucleus. Our data also suggest a potential role for SGs in the growth of
cytoplasmic poly-PR aggregates as well as a role for poly-PR assemblies in "nucleating"
TDP-43 aggregation in the cytoplasm.

Discussion

A vast body of knowledge on C9-DPR related disease mechanisms has accumulated in the
past decade yet it remains unclear whether, and if so how, DPR aggregation – a C9-
ALS/FTD hallmark pathology – contributes to the disease. Aggregation intermediates, and
especially smaller, highly reactive oligomeric species, have been validated as
toxic/pathogenic in the case of other neurodegeneration-linked proteins, such as tau and
alpha-synuclein (Bengoa-Vergniory *et al*, 2017; Choi & Gandhi, 2018; Jucker & Walker,
2018). It is possible that equivalent DPR aggregation products play a role in C9-ALS/FTD.
Although the inclusions of all five DPRs in the patient CNS are morphologically similar
(Gendron *et al.*, 2013), in cellular models, DPRs other than GA fail to form microscopically
visible assemblies (Frottin *et al.*, 2021; Liu *et al.*, 2022; Zhou *et al*, 2017). In order to
circumvent the high solubility of R-DPRs in cells, we harnessed the Cry2olig opto-module
(Taslimi *et al.*, 2014) successfully used previously to promote self-association of
neurodegeneration-linked proteins (Berard *et al*, 2022; Jiang *et al*, 2021; Mann *et al*, 2019).
In line with the molecular dynamics predictions (Zheng *et al.*, 2021), poly-GR's self-
association capacity was too low even when facilitated by Cry2olig, not yielding visible
condensation in cells. However we succeeded in achieving the condensation of the
oligomerisation-competent poly-PR.

Two key phenotypes – alterations to the nucleolus and SGs – validate our opto-PR model in
the context of the existing literature (Kwon *et al.*, 2014; Marmor-Kollet *et al.*, 2020; White *et*
al., 2019). In our model, SG assembly is enhanced upon opto-PR cytoplasmic

accumulation/aggregation. This can be due to increased nucleation of SGs by oligomerised
opto-PR and/or due to their altered composition. We also observed sequestration of NPM1,
a key R-DPR interactor, into opto-PR condensates. Beyond these two phenotypes however,
we failed to detect the ubiquitous MLO disruption by poly-PR reported in some studies, and
its condensation also had limited effect on MLOs. This may be due to high variability of these
phenotypes depending on the cell type, DPR levels and repeat length. This calls for
establishing consensus phenotypes that can be used for the benchmarking of novel cellular
DPR models, with the nucleolar and SG pathology being the prime candidates.

We found that poly-PR confers a specific ordered arrangement to the opto-induced
condensates – sphere with hollow centre. Arginine-rich peptides form such anisotropic
structures in the presence of RNA *in vitro*, both under RNA and peptide excess conditions
(Alshareedah *et al.*, 2020). Adopting this model, we speculate that upon interaction with
cellular RNA, the small opto-PR oligomers, nucleated with the aid of Cry2olig, form
nanocondensates with a neutral "head" and a charged "tail" that subsequently coalesce into
RNA-coated micelles (~100 nm granules). These transition into a vesicle-like conformation
"layered" with RNA on both the internal and external surface (~250 nm granules). Upon
(r)RNA depletion, these condensates undergo fusion into a larger (>500 nm) hollow-centre
structure (Fig.S5A). Anisotropic condensate is a non-equilibrium state that may require ATP
to be established and maintained (Bergmann *et al*, 2023), and in our system, this state is
fuelled by the light-induced Cry2olig oligomerisation. Nuclear poly-PR condensates not
overlapping with nucleolar markers form in transgenic mice (Zhang *et al.*, 2019), and it would
be interesting to establish whether they adopt a hollow-centre structure at any point during
their biogenesis. Anisotropic nuclear assemblies are detected in an increasing number of
neuro-diseases, for example, they are formed by acetylated TDP-43 (Yu *et al*, 2021), RNA-
binding deficient TDP-43 (Wang *et al.*, 2020) (Fig.S5B) and RBPs/polyA+ RNA in cell
models of spinal muscular atrophy (Narcis *et al*, 2018). Furthermore, we found that nuclear
foci of ALS-linked CREST protein and its mutants (Kukharsky *et al*, 2015) also possess this
typical morphology (Fig.S5B). Remarkably, DDX3X mutations causative of
neurodevelopmental disorders form cytoplasmic hollow condensates, and those composed
of an aggressive RNA-binding deficient mutant display low recovery in FRAP (Owens *et al*,
2023). One can speculate that neurological disease states are characterised by specific
changes to the neuronal metabolism, e.g. altered protein and RNA stoichiometries, that
favour biogenesis of this assembly type.

RNA facilitates LLPS of poly-PR *in vitro* (Balendra et al., 2023) however cellular RNA
restricts the growth of the non-dynamic (based on FRAP analysis) opto-PR condensates in
cells – similar to TDP-43 (Grese *et al*, 2021; Mann *et al.*, 2019). Together with the reports on
the role of RNA in buffering pathological RBP aggregation (Maharana *et al*, 2018; Mann *et al.*, 2019; Shelkovnikova *et al*, 2014a) and on the wide-spread RNA degradation in ALS
al., 2019; Shelkovnikova *et al*, 2014a) and on the wide-spread RNA degradation in ALS
(Tank *et al*, 2018), our findings suggest than declining RNA levels may be a common factor
underlying protein aggregation across different ALS subtypes, including C9-ALS.

Transient nuclear TDP-43 condensation is a hallmark of stress response (Cohen *et al.*,
2015; Wang *et al.*, 2020). We recently found that this response leads to TDP-43 loss of
function and prolonged STMN2 depletion and is dysregulated by TDP-43 mutations (Huang,
Ellis et al., 2024, manuscript in revision). These phenotypes may become persistent with
chronic or repetitive stress, precipitating the disease. Here we show that poly-PR self-
assembly causes TDP-43 condensation in the absence of stress signalling, indicating the
convergence between C9-ALS and other ALS subtypes in this putative disease mechanism.

Consistently, recent use of a specific RNA aptamer revealed nuclear TDP-43 granulation in
motor neurons in ALS tissue (Spence et al., 2023, doi: 10.1101/2023.10.24.563701).

Cytoplasmic poly-PR assemblies in our opto-model are structurally dissimilar to and more
persistent than the nuclear condensates, probably reflecting a different environment in the
two cellular compartments. This higher stability is consistent with a higher frequency of poly-
PR cytoplasmic inclusions in patients (Zu *et al.*, 2013). DPR inclusions are rarely found to
co-deposit with TDP-43 inclusions in the patient tissue, with clear anatomical region
specificity (Al-Sarraj et al, 2011). However, some histopathological studies find evidence of
DPR aggregation preceding the cytoplasmic TDP-43 pathology (Baborie *et al*, 2015;
Vatsavayai *et al*, 2016). TDP-43 was found to join the cytoplasmic opto-PR foci where the
two proteins remained demixed. It is tempting to speculate that transient cytoplasmic poly-
PR assemblies "seed" TDP-43 pathology. SGs may play an equivalent role for cytoplasmic
poly-PR inclusions, concentrating the initial poly-PR assemblies in a confined space and
promoting their coalescence into a larger aggregate. Indeed, a recent study showed that
cytoplasmic TDP-43 aggregates can be nucleated within SGs, as a separate phase, and left
behind after SG dissolution (Yan *et al*, 2024).

Our model has several limitations. We used a relatively short repeat length (which
nevertheless is in the patient range), whereas the Cry2olig-mCherry is a large tag. However,
multiple studies have demonstrated that DPR repeat lengths of 30-1000 have identical
subcellular localisation with similar toxicity (Bennion Callister *et al.*, 2016; Kwon *et al.*, 2014;
Miyagi *et al*, 2023; White *et al.*, 2019). We cannot exclude that longer repeats will undergo
condensation more readily, which should be tested in future studies. In addition, for this
proof-of-principle study, non-neuronal cells were used for the ease of imaging. To address
potential cell-specificity of the phenotypes revealed in this study, the opto-PR model should
be transferred into a (human) neuronal system.

Materials and methods
Plasmids
The CRY2olig-mCherry backbone vector was a gift from Chandra Tucker (Addgene plasmid
60032) (Taslimi *et al.*, 2014). DPR plasmids were a gift from Kurt De Vos (Bauer *et al*,
2022). Codon-optimised DPR sequences were cloned into the CRY2olig-mCherry vector
using standard techniques. These plasmids are available via Addgene. Plasmids for the
expression of GFP-tagged TDP-43, FUS, NONO, SFPQ and CREST (N-terminal tag, in
pEGFP-C1 vector) were generated previously (Kukharsky *et al.*, 2015; Shelkovnikova *et al*
2014b). TDP-43 K145Q and F147/149L in pEGFP-C1 vector were generated using PCR
mutagenesis.

Cell culture, transfection and treatments
HeLa cells were obtained from ATCC via Sigma, cultured in Dulbecco's Modified Eagle
Medium/Nutrient Mixture F-12 (DMEM/F-12) supplemented with 10% foetal bovine serum
(FBS) and penicillin-streptomycin. For time-lapse imaging, cells were plated on PhenoPlate-
96 (black, optically clear bottom, PerkinElmer) at a density of 2×10^4. For all other
experiments, cells were seeded in 24 well plates, with or without coverslips dependent on
the application, at a density of 5×10^4 cells, unless otherwise stated. Transfection was
performed 24 h prior to blue light stimulation, using either Lipofectamine 2000
(ThermoFisher) or jetPRIME (Jena Bioscience) according to the manufacturer's instructions.
For transcriptional inhibition cells were treated with 2.5 µg/ml actinomycin D or 5,6-dichloro-

1-beta-D-ribofuranosylbenzimidazole (DRB) (both Sigma). For inhibition of arginine
methylation, cells were treated with 10 µM MS-023 or EPZ015666 (both ApexBio). To induce
cellular stress, cells were treated with 250 µM NaAsO$_2$ (Sigma). Incubation times for
treatments within individual experiments are indicated in respective Figure legends.
407
Opto-stimulation
Cells expressing opto-constructs were stimulated with a 488 nm laser on Opera Phenix HCS
(500 ms, 80% laser power) for live cell time-lapse imaging (under full environmental control
conditions), or on a custom-built LED blue-light array housed in a humidified incubator
maintained at 37℃ with 5 % CO$_2$. Cells were protected from light between experiments and
during fixation.
414
Microscopy
Conventional fluorescence microscopy was performed using 100x objective on an Olympus
BX57 upright microscope equipped with an ORCA-Flash 4.0 camera (Hamamatsu) and
cellSens Dimension software (Olympus). Super-resolution microscopy was performed using
a 63x oil immersion objective on a ZEISS 980 laser scanning confocal microscope (LSM)
with Airyscan 2 detector and ZEN Blue software. Time-lapse microscopy was performed
using a 40x objective on Opera Phenix HCS, and Harmony 4.9 software was used for image
processing and analysis (all PerkinElmer). Image processing and profile drawing was done
using Image J or ZEN Blue software. Condensate/aggregate quantification was done
manually or on Image J in a blinded manner.
425
Fluorescent recovery after photobleaching (FRAP)
Cells seeded at a density of 2.8 x 10^5 in glass-bottomed 35mm dishes (Ibidi), were
transfected and 24 h post-transfection, subjected to 3-hr continuous stimulation prior to
FRAP analysis. Imaging was performed using a 63x oil immersion objective on a ZEISS
LSM 800 confocal microscope, equipped with a humidified incubation chamber maintained
at 37℃ with 5 % CO$_2$. FRAP acquisition was performed on condensates formed after 3 h of
continuous stimulation. A circular region of interest (ROI) around each condensate was
bleached using a 568 nm laser at 100 % laser power. Images were captured pre-bleach,
immediately following bleach and at x 200 ms intervals during recovery. The mean
fluorescence intensity within the ROI was determined for each image using ZEN blue
software. Intensity values were corrected for bleaching during imaging and normalised to
pre-bleach intensity. Average values were plotted and FRAP curves fitted using a one-phase
association equation in GraphPad Prism 9 software.
439
Immunocytochemistry and RNA-FISH
Immunocytochemistry was performed as described elsewhere (Shelkovnikova *et al*, 2018)
using the following commercially available antibodies: FBL (rabbit polyclonal, Proteintech,
16021-1-AP); NPM1 (mouse monoclonal, Proteintech, 60096-1-Ig); UPF1 (rabbit polyclonal,
Proteintech, 23379-1-AP); PNN (rabbit polyclonal, Proteintech, 18266-1-AP); coilin p80
(mouse monoclonal, BD Biosciences, 612074); SMN (mouse monoclonal, BD Biosciences,
610647); G3BP1 (rabbit polyclonal, Proteintech, 13057-2-AP); repeat-PR (rabbit polyclonal,
Proteintech, 23979-1-AP); TDP-43 (rabbit polyclonal C-terminal, Sigma); FUS (mouse
monoclonal, Santa Cruz, [4H11], sc-47711); NONO (rabbit polyclonal, Proteintech, 11058-1-
AP); SFPQ (rabbit monoclonal, Abcam, ab177149). RNA-FISH using a Cy5-labelled dT$_{(30)}$

DNA oligonucleotide probe (Sigma) was performed as described earlier (Shelkovnikova *et*
al, 2017).
.
Electrophoretic mobility shift assay (EMSA)
Cy5-labeled RNA oligonucleotide 5'-
GAGAGAGCGCGUGCAGAGACUUGGUGGUGCAUAA-3' ("Clip34nt") (Bhardwaj *et al*.,
2013) was custom-synthesised by Eurofins. Poly-GR$_{x10}$ and poly-PR$_{x10}$ were custom-made
by Biosynth. Lyophilised peptides were resuspended in water at 50 mM and stored at -80℃
before use. RNA was incubated at 250nM with 250nM-10µM peptides in the EMSA buffer
(50 mM Tris-HCl, pH 7.5, 100 mM KCl, 2 mM MgCl$_2$, 100 mM β-mercaptoethanol, 0.1 mg/ml
BSA) for 15 min at RT with gentle shaking. Samples were analysed on 6% native acrylamide
gel in TBE buffer, followed by imaging on Licor Odissey FC (700nm channel).
462
Cell lysis, fractionation, and western blotting
For fractionation, cells were lysed in 1 % Titon-X-100 in PBS for 5 min at RT. Lysates were
incubated on ice for 30 minutes with vortexing at 5 min intervals and centrifuged at
13,000rpm for 20 min at 4℃. Supernatant and pellet were mixed with 2x Laemmli buffer
(2xLB), vortexed and heated at 95℃ for 10 min. For western blotting, proteins were resolved
on a 10% Mini-PROTEAN® TGX™ hand-cast protein gel and transferred to a PVDF
membrane. Gels were stained with Gelcode™ (ThermoFisher) post-transfer for total protein.
Membranes were blocked for 1 h in 4 % milk/TBST and then incubated with the following
primary antibodies (1:1000 dilution) at 4℃ overnight: mCherry (rabbit polyclonal, Proteintech,
26765-1-AP); GFP (mouse monoclonal, 66002-1-Ig, Proteintech); TDP-43 (rabbit polyclonal,
C-terminal, Sigma); eIF2α (rabbit polyclonal, Cell Signaling, 9722), eIF2αP (rabbit polyclonal,
Abcam, antibody [E90], ab32157); repeat-PR (rabbit polyclonal, Proteintech, 23979-1-AP).
Following primary antibody incubation, blots were washed with TBST and incubated with an
appropriate HRP secondary antibody (GE Healthcare) for 1 h at room temperature. Signal
was detected using Clarity Max Western ECL Substrate (Bio-Rad) and imaged and
quantified using Licor Odissey FC/Image Studio software.
479
RNA expression analysis
Total RNA was extracted using GenElute total mammalian RNA kit (Sigma) in accordance
with the manufacturer's instructions. First-strand cDNA synthesis was performed using 500
ng of RNA with random primers (ThermoFisher) and MMLV reverse transcriptase
(Promega). qRT-PCR was performed using qPCRBIO SyGreen Lo-ROX (PCRbio), and
GAPDH was used for normalisation. Primer sequences are provided in a previous study
(Shelkovnikova *et al*., 2017).
487
In vitro analysis of TDP-43 clustering
In vitro TDP-43 clustering analysis with immunodetection and imaging were performed as
described in Hodgson, Huang et al., 2024 (doi: 10.2139/ssrn.4721338). Briefly, 1 µM total
recombinant TDP-43 (R&D Biosystems, AP-190-100) or supernatant (after centrifuging at
13,300 rpm for 1 min) was mixed with 1 µM poly-PR/GR peptides described above in the
assay buffer, and incubated for 10 min. Samples were sedimented and fixed with
glutaraldehyde on coverslips, blocked with 1% BSA in PBS for 1 h at RT and incubated with
an anti-TDP-43 antibody (1:5000, mouse monoclonal, R&D Biosystems, MAB77781) in the
blocking buffer for 2 h. After washes, TDP-43 protein clusters were visualised using anti-
mouse AlexaFluor488 antibody (1:2000, ThermoFisher), incubated for 1 h at RT. Coverslips

were mounted on a glass slide using Immu-mount (ThermoFisher). Images were taken using
Olympus BX57 upright microscope and ORCA-Flash 4.0 camera and processed using
cellSens Dimension software (Olympus). Quantification of assemblies was done using the
'Analyze particles' tool of Image J. For confirmatory western blot analysis, recombinant TDP-
43 samples incubated with or without the peptides as above (for 10 min, and an additional
set – for 4 h) were centrifuged at 1,000xg for 10 min. Pellet and supernatant were analysed
by western blot using a C-terminal TDP-43 antibody (Sigma).
Statistical analysis
Analysis was done using respective tests on GraphPad Prism 9 software. N corresponds to
the number of biological replicates. Error bars represent S.E.M.
508

Author contributions

Rachel E. Hodgson: Conceptualization; Data curation; Formal analysis; Investigation;
Methodology; Writing - original draft; Writing - review and editing. **Jessica Rayment:** Formal
analysis; Investigation; Methodology. **Wan-Ping Huang**: Formal analysis; Investigation;
Methodology; Writing - review and editing. **Anna Sanchez Avila**: Data curation;
Investigation. **Tatyana A. Shelkovnikova:** Conceptualization; Supervision; Funding
acquisition; Project administration; Writing - review and editing.
516

Disclosure and competing interests statement

518 The authors declare no competing interests.
519

Acknowledgements

521 This work was supported by the UKRI Future Leaders Fellowship (MR/W004615/1), MRC
522 (MR/W028522/1) and BBSRC (BB/V014110/1) standard grants, and MND Association
523 fellowship/grant (Shelkovnikova/Oct17/968-799). We also acknowledge the MRC grant
524 MR/X012077/1 for Airyscan 2 LMF.

References

Al-Sarraj S, King A, Troakes C, Smith B, Maekawa S, Bodi I, Rogelj B, Al-Chalabi A,
Hortobagyi T, Shaw CE (2011) p62 positive, TDP-43 negative, neuronal cytoplasmic and
intranuclear inclusions in the cerebellum and hippocampus define the pathology of C9orf72-
linked FTLD and MND/ALS. *Acta neuropathologica* 122: 691-702

Alshareedah I, Moosa MM, Raju M, Potoyan DA, Banerjee PR (2020) Phase transition
of RNA-protein complexes into ordered hollow condensates. *Proc Natl Acad Sci U S A* 117:
15650-15658

Ash PE, Bieniek KF, Gendron TF, Caulfield T, Lin WL, Dejesus-Hernandez M, van
Blitterswijk MM, Jansen-West K, Paul JW, 3rd, Rademakers R *et al* (2013) Unconventional
translation of C9ORF72 GGGGCC expansion generates insoluble polypeptides specific to
c9FTD/ALS. *Neuron* 77: 639-646

Baborie A, Griffiths TD, Jaros E, Perry R, McKeith IG, Burn DJ, Masuda-Suzukake M,
Hasegawa M, Rollinson S, Pickering-Brown S *et al* (2015) Accumulation of dipeptide repeat
proteins predates that of TDP-43 in frontotemporal lobar degeneration associated with
hexanucleotide repeat expansions in C9ORF72 gene. *Neuropathol Appl Neurobiol* 41: 601-
612

Balendra R, Ruiz de Los Mozos I, Odeh HM, Glaria I, Milioto C, Wilson KM, Ule AM,
Hallegger M, Masino L, Martin S *et al* (2023) Transcriptome-wide RNA binding analysis of
C9orf72 poly(PR) dipeptides. *Life Sci Alliance* 6

Bauer CS, Webster CP, Shaw AC, Kok JR, Castelli LM, Lin YH, Smith EF, Illanes-Alvarez F, Higginbottom A, Shaw PJ *et al* (2022) Loss of TMEM106B exacerbates C9ALS/FTD DPR pathology by disrupting autophagosome maturation. *Front Cell Neurosci* 16: 1061559

Bengoa-Vergniory N, Roberts RF, Wade-Martins R, Alegre-Abarrategui J (2017) Alpha-synuclein oligomers: a new hope. *Acta neuropathologica* 134: 819-838

Bennion Callister J, Ryan S, Sim J, Rollinson S, Pickering-Brown SM (2016) Modelling C9orf72 dipeptide repeat proteins of a physiologically relevant size. *Human molecular genetics* 25: 5069-5082

Berard M, Sheta R, Malvaut S, Rodriguez-Aller R, Teixeira M, Idi W, Turmel R, Alpaugh M, Dubois M, Dahmene M *et al* (2022) A light-inducible protein clustering system for in vivo analysis of alpha-synuclein aggregation in Parkinson disease. *PLoS Biol* 20: e3001578

Bergmann AM, Bauermann J, Bartolucci G, Donau C, Stasi M, Holtmannspotter AL, Julicher F, Weber CA, Boekhoven J (2023) Liquid spherical shells are a non-equilibrium steady state of active droplets. *Nat Commun* 14: 6552

Bhardwaj A, Myers MP, Buratti E, Baralle FE (2013) Characterizing TDP-43 interaction with its RNA targets. *Nucleic Acids Res* 41: 5062-5074

Boeynaems S, Bogaert E, Kovacs D, Konijnenberg A, Timmerman E, Volkov A, Guharoy M, De Decker M, Jaspers T, Ryan VH *et al* (2017) Phase Separation of C9orf72 Dipeptide Repeats Perturbs Stress Granule Dynamics. *Molecular cell* 65: 1044-1055 e1045

Chan-Penebre E, Kuplast KG, Majer CR, Boriack-Sjodin PA, Wigle TJ, Johnston LD, Rioux N, Munchhof MJ, Jin L, Jacques SL *et al* (2015) A selective inhibitor of PRMT5 with in vivo and in vitro potency in MCL models. *Nat Chem Biol* 11: 432-437

Chew J, Cook C, Gendron TF, Jansen-West K, Del Rosso G, Daughrity LM, Castanedes-Casey M, Kurti A, Stankowski JN, Disney MD *et al* (2019) Aberrant deposition of stress granule-resident proteins linked to C9orf72-associated TDP-43 proteinopathy. *Mol Neurodegener* 14: 9

Choi ML, Gandhi S (2018) Crucial role of protein oligomerization in the pathogenesis of Alzheimer's and Parkinson's diseases. *Febs J* 285: 3631-3644

Choi SY, Lopez-Gonzalez R, Krishnan G, Phillips HL, Li AN, Seeley WW, Yao WD, Almeida S, Gao FB (2019) C9ORF72-ALS/FTD-associated poly(GR) binds Atp5a1 and compromises mitochondrial function in vivo. *Nat Neurosci* 22: 851-862

Cochard A, Garcia-Jove Navarro M, Piroska L, Kashida S, Kress M, Weil D, Gueroui Z (2022) RNA at the surface of phase-separated condensates impacts their size and number. *Biophys J* 121: 1675-1690

Cohen TJ, Hwang AW, Restrepo CR, Yuan CX, Trojanowski JQ, Lee VM (2015) An acetylation switch controls TDP-43 function and aggregation propensity. *Nat Commun* 6: 5845

Cook CN, Wu Y, Odeh HM, Gendron TF, Jansen-West K, Del Rosso G, Yue M, Jiang P, Gomes E, Tong J *et al* (2020) C9orf72 poly(GR) aggregation induces TDP-43 proteinopathy. *Sci Transl Med* 12

DeJesus-Hernandez M, Mackenzie IR, Boeve BF, Boxer AL, Baker M, Rutherford NJ, Nicholson AM, Finch NA, Flynn H, Adamson J *et al* (2011) Expanded GGGGCC hexanucleotide repeat in noncoding region of C9ORF72 causes chromosome 9p-linked FTD and ALS. *Neuron* 72: 245-256

Eram MS, Shen Y, Szewczyk M, Wu H, Senisterra G, Li F, Butler KV, Kaniskan HU, Speed BA, Dela Sena C *et al* (2016) A Potent, Selective, and Cell-Active Inhibitor of Human Type I Protein Arginine Methyltransferases. *ACS Chem Biol* 11: 772-781

Frottin F, Perez-Berlanga M, Hartl FU, Hipp MS (2021) Multiple pathways of toxicity
induced by C9orf72 dipeptide repeat aggregates and G(4)C(2) RNA in a cellular model. *Elife*
10

Gendron TF, Bieniek KF, Zhang YJ, Jansen-West K, Ash PE, Caulfield T, Daughrity L,
Dunmore JH, Castanedes-Casey M, Chew J *et al* (2013) Antisense transcripts of the
expanded C9ORF72 hexanucleotide repeat form nuclear RNA foci and undergo repeat-
associated non-ATG translation in c9FTD/ALS. *Acta neuropathologica* 126: 829-844

Gittings LM, Boeynaems S, Lightwood D, Clargo A, Topia S, Nakayama L, Troakes C,
Mann DMA, Gitler AD, Lashley T *et al* (2020) Symmetric dimethylation of poly-GR correlates
with disease duration in C9orf72 FTLD and ALS and reduces poly-GR phase separation and
toxicity. *Acta neuropathologica* 139: 407-410

Grese ZR, Bastos AC, Mamede LD, French RL, Miller TM, Ayala YM (2021) Specific
RNA interactions promote TDP-43 multivalent phase separation and maintain liquid
properties. *EMBO Rep* 22: e53632

Hartmann H, Hornburg D, Czuppa M, Bader J, Michaelsen M, Farny D, Arzberger T,
Mann M, Meissner F, Edbauer D (2018) Proteomics and C9orf72 neuropathology identify
ribosomes as poly-GR/PR interactors driving toxicity. *Life Sci Alliance* 1: e201800070

Hodgson R, Huang WP, Kumar V, Haiyan A, Chalakova Z, Rayment J, Ellis BCS,
Stender EGP, van Vugt, JJFA, Project MinE ALS Sequencing Consortium, Locker N, Pitout
I, Fletcher S, Cooper-Knock J, Shelkovnikova T (2024) TDP-43 is a Master Regulator of
Paraspeckle Condensation. Available at SSRN: https://ssrn.com/abstract=4721338 or
http://dx.doi.org/10.2139/ssrn.4721338

Huang WP, Ellis BCS, Hodgson R, Sanchez Avila A, Kumar V, Rayment J, Moll T,
Shelkovnikova TA. Stress-induced TDP-43 nuclear condensation leads to splicing loss-of-
function and STMN2 depletion. *Manuscript in revision*

Jiang L, Lin W, Zhang C, Ash PEA, Verma M, Kwan J, van Vliet E, Yang Z, Cruz AL,
Boudeau S *et al* (2021) Interaction of tau with HNRNPA2B1 and N(6)-methyladenosine RNA
mediates the progression of tauopathy. *Molecular cell* 81: 4209-4227 e4212

Jucker M, Walker LC (2018) Propagation and spread of pathogenic protein assemblies
in neurodegenerative diseases. *Nat Neurosci* 21: 1341-1349

Kanekura K, Harada Y, Fujimoto M, Yagi T, Hayamizu Y, Nagaoka K, Kuroda M
(2018) Characterization of membrane penetration and cytotoxicity of C9orf72-encoding
arginine-rich dipeptides. *Sci Rep* 8: 12740

Kukharsky MS, Quintiero A, Matsumoto T, Matsukawa K, An H, Hashimoto T,
Iwatsubo T, Buchman VL, Shelkovnikova TA (2015) Calcium-responsive transactivator
(CREST) protein shares a set of structural and functional traits with other proteins
associated with amyotrophic lateral sclerosis. *Mol Neurodegener* 10: 20

Kwon I, Xiang S, Kato M, Wu L, Theodoropoulos P, Wang T, Kim J, Yun J, Xie Y,
McKnight SL (2014) Poly-dipeptides encoded by the C9orf72 repeats bind nucleoli, impede
RNA biogenesis, and kill cells. *Science* 345: 1139-1145

LaClair KD, Zhou Q, Michaelsen M, Wefers B, Brill MS, Janjic A, Rathkolb B, Farny D,
Cygan M, de Angelis MH *et al* (2020) Congenic expression of poly-GA but not poly-PR in
mice triggers selective neuron loss and interferon responses found in C9orf72 ALS. *Acta*
neuropathologica 140: 121-142

Lee KH, Zhang P, Kim HJ, Mitrea DM, Sarkar M, Freibaum BD, Cika J, Coughlin M,
Messing J, Molliex A *et al* (2016) C9orf72 Dipeptide Repeats Impair the Assembly,
Dynamics, and Function of Membrane-Less Organelles. *Cell* 167: 774-788 e717

Lin Y, Mori E, Kato M, Xiang S, Wu L, Kwon I, McKnight SL (2016) Toxic PR Poly-
Dipeptides Encoded by the C9orf72 Repeat Expansion Target LC Domain Polymers. *Cell*
167: 789-802 e712

Liu F, Morderer D, Wren MC, Vettleson-Trutza SA, Wang Y, Rabichow BE, Salemi
MR, Phinney BS, Oskarsson B, Dickson DW *et al* (2022) Proximity proteomics of C9orf72
dipeptide repeat proteins identifies molecular chaperones as modifiers of poly-GA
aggregation. *Acta Neuropathol Commun* 10: 22
Lopez-Gonzalez R, Lu Y, Gendron TF, Karydas A, Tran H, Yang D, Petrucelli L, Miller
BL, Almeida S, Gao FB (2016) Poly(GR) in C9ORF72-Related ALS/FTD Compromises
Mitochondrial Function and Increases Oxidative Stress and DNA Damage in iPSC-Derived
Motor Neurons. *Neuron* 92: 383-391
Mackenzie IR, Frick P, Grasser FA, Gendron TF, Petrucelli L, Cashman NR, Edbauer
D, Kremmer E, Prudlo J, Troost D *et al* (2015) Quantitative analysis and clinico-pathological
correlations of different dipeptide repeat protein pathologies in C9ORF72 mutation carriers.
Acta neuropathologica 130: 845-861
Maharana S, Wang J, Papadopoulos DK, Richter D, Pozniakovsky A, Poser I, Bickle
M, Rizk S, Guillen-Boixet J, Franzmann TM *et al* (2018) RNA buffers the phase separation
behavior of prion-like RNA binding proteins. *Science* 360: 918-921
Mann JR, Gleixner AM, Mauna JC, Gomes E, DeChellis-Marks MR, Needham PG,
Copley KE, Hurtle B, Portz B, Pyles NJ *et al* (2019) RNA Binding Antagonizes Neurotoxic
Phase Transitions of TDP-43. *Neuron* 102: 321-338 e328
Marmor-Kollet H, Siany A, Kedersha N, Knafo N, Rivkin N, Danino YM, Moens TG,
Olender T, Sheban D, Cohen N *et al* (2020) Spatiotemporal Proteomic Analysis of Stress
Granule Disassembly Using APEX Reveals Regulation by SUMOylation and Links to ALS
Pathogenesis. *Molecular cell* 80: 876-891 e876
Miyagi T, Ueda K, Sugimoto M, Yagi T, Ito D, Yamazaki R, Narumi S, Hayamizu Y, Uji
IH, Kuroda M *et al* (2023) Differential toxicity and localization of arginine-rich C9ORF72
dipeptide repeat proteins depend on de-clustering of positive charges. *iScience* 26: 106957
Mizielinska S, Gronke S, Niccoli T, Ridler CE, Clayton EL, Devoy A, Moens T, Norona
FE, Woollacott IOC, Pietrzyk J *et al* (2014) C9orf72 repeat expansions cause
neurodegeneration in Drosophila through arginine-rich proteins. *Science* 345: 1192-1194
Moens TG, Niccoli T, Wilson KM, Atilano ML, Birsa N, Gittings LM, Holbling BV, Dyson
MC, Thoeng A, Neeves J *et al* (2019) C9orf72 arginine-rich dipeptide proteins interact with
ribosomal proteins in vivo to induce a toxic translational arrest that is rescued by eIF1A. *Acta
neuropathologica* 137: 487-500
Mori K, Weng SM, Arzberger T, May S, Rentzsch K, Kremmer E, Schmid B,
Kretzschmar HA, Cruts M, Van Broeckhoven C *et al* (2013) The C9orf72 GGGGCC repeat is
translated into aggregating dipeptide-repeat proteins in FTLD/ALS. *Science* 339: 1335-1338
Narcis JO, Tapia O, Tarabal O, Piedrafita L, Caldero J, Berciano MT, Lafarga M (2018)
Accumulation of poly(A) RNA in nuclear granules enriched in Sam68 in motor neurons from
the SMNDelta7 mouse model of SMA. *Sci Rep* 8: 9646
Ortega JA, Sasselli IR, Boccitto M, Fleming AC, Fortuna TR, Li Y, Sato K, Clemons
TD, McKenna ED, Nguyen TP *et al* (2023) CLIP-Seq analysis enables the design of
protective ribosomal RNA bait oligonucleotides against C9ORF72 ALS/FTD poly-GR
pathophysiology. *Sci Adv* 9: eadf7997
Owens MC, Shen H, Yanas A, Mendoza-Figueroa MS, Lavorando E, Wei X, Shweta
H, Tang HY, Goldman YE, Liu KF (2023) Mutant forms of DDX3X with diminished catalysis
form hollow condensates that exhibit sex-specific regulation. *bioRxiv*
Park H, Kim NY, Lee S, Kim N, Kim J, Heo WD (2017) Optogenetic protein clustering
through fluorescent protein tagging and extension of CRY2. *Nat Commun* 8: 30
Quaegebeur A, Glaria I, Lashley T, Isaacs AM (2020) Soluble and insoluble dipeptide
repeat protein measurements in C9orf72-frontotemporal dementia brains show regional
differential solubility and correlation of poly-GR with clinical severity. *Acta Neuropathol
Commun* 8: 184

Renton AE, Majounie E, Waite A, Simon-Sanchez J, Rollinson S, Gibbs JR, Schymick
JC, Laaksovirta H, van Swieten JC, Myllykangas L *et al* (2011) A hexanucleotide repeat
expansion in C9ORF72 is the cause of chromosome 9p21-linked ALS-FTD. *Neuron* 72: 257-
268

Saberi S, Stauffer JE, Jiang J, Garcia SD, Taylor AE, Schulte D, Ohkubo T,
Schloffman CL, Maldonado M, Baughn M *et al* (2018) Sense-encoded poly-GR dipeptide
repeat proteins correlate to neurodegeneration and uniquely co-localize with TDP-43 in
dendrites of repeat-expanded C9orf72 amyotrophic lateral sclerosis. *Acta neuropathologica*
135: 459-474

Sakae N, Bieniek KF, Zhang YJ, Ross K, Gendron TF, Murray ME, Rademakers R,
Petrucelli L, Dickson DW (2018) Poly-GR dipeptide repeat polymers correlate with
neurodegeneration and Clinicopathological subtypes in C9ORF72-related brain disease.
Acta Neuropathol Commun 6: 63

Schludi MH, May S, Grasser FA, Rentzsch K, Kremmer E, Kupper C, Klopstock T,
German Consortium for Frontotemporal Lobar D, Bavarian Brain Banking A, Arzberger T *et*
al (2015) Distribution of dipeptide repeat proteins in cellular models and C9orf72 mutation
cases suggests link to transcriptional silencing. *Acta neuropathologica* 130: 537-555

Shav-Tal Y, Blechman J, Darzacq X, Montagna C, Dye BT, Patton JG, Singer RH,
Zipori D (2005) Dynamic sorting of nuclear components into distinct nucleolar caps during
transcriptional inhibition. *Mol Biol Cell* 16: 2395-2413

Shelkovnikova TA, Dimasi P, Kukharsky MS, An H, Quintiero A, Schirmer C, Buee L,
Galas MC, Buchman VL (2017) Chronically stressed or stress-preconditioned neurons fail to
maintain stress granule assembly. *Cell Death Dis* 8: e2788

Shelkovnikova TA, Kukharsky MS, An H, Dimasi P, Alexeeva S, Shabir O, Heath PR,
Buchman VL (2018) Protective paraspeckle hyper-assembly downstream of TDP-43 loss of
function in amyotrophic lateral sclerosis. *Mol Neurodegener* 13: 30

Shelkovnikova TA, Robinson HK, Southcombe JA, Ninkina N, Buchman VL (2014a)
Multistep process of FUS aggregation in the cell cytoplasm involves RNA-dependent and
RNA-independent mechanisms. *Human molecular genetics*

Shelkovnikova TA, Robinson HK, Troakes C, Ninkina N, Buchman VL (2014b)
Compromised paraspeckle formation as a pathogenic factor in FUSopathies. *Human*
molecular genetics 23: 2298-2312

Shin Y, Berry J, Pannucci N, Haataja MP, Toettcher JE, Brangwynne CP (2017)
Spatiotemporal Control of Intracellular Phase Transitions Using Light-Activated optoDroplets.
Cell 168: 159-171 e114

Spence H, Waldron FM, Saleeb RS, Brown AL, Rifai OM, Gilodi M, Read F, Roberts K,
Milne G, Wilkinson D, *et al* (2023) RNA aptamer reveals nuclear TDP-43 pathology is an
early aggregation event that coincides with STMN-2 cryptic splicing and precedes clinical
manifestation in ALS. bioRxiv. doi: https://doi.org/10.1101/2023.10.24.563701

Tank EM, Figueroa-Romero C, Hinder LM, Bedi K, Archbold HC, Li X, Weskamp K,
Safren N, Paez-Colasante X, Pacut C *et al* (2018) Abnormal RNA stability in amyotrophic
lateral sclerosis. *Nat Commun* 9: 2845

Taslimi A, Vrana JD, Chen D, Borinskaya S, Mayer BJ, Kennedy MJ, Tucker CL
(2014) An optimized optogenetic clustering tool for probing protein interaction and function.
Nat Commun 5: 4925

Van Mossevelde S, van der Zee J, Cruts M, Van Broeckhoven C (2017) Relationship
between C9orf72 repeat size and clinical phenotype. *Curr Opin Genet Dev* 44: 117-124

Vanneste J, Vercruysse T, Boeynaems S, Sicart A, Van Damme P, Daelemans D, Van
Den Bosch L (2019) C9orf72-generated poly-GR and poly-PR do not directly interfere with
nucleocytoplasmic transport. *Sci Rep* 9: 15728

Vatsavayai SC, Yoon SJ, Gardner RC, Gendron TF, Vargas JN, Trujillo A, Pribadi M,
Phillips JJ, Gaus SE, Hixson JD *et al* (2016) Timing and significance of pathological features
in C9orf72 expansion-associated frontotemporal dementia. *Brain : a journal of neurology*
139: 3202-3216

Wang C, Duan Y, Duan G, Wang Q, Zhang K, Deng X, Qian B, Gu J, Ma Z, Zhang S
et al (2020) Stress Induces Dynamic, Cytotoxicity-Antagonizing TDP-43 Nuclear Bodies via
Paraspeckle LncRNA NEAT1-Mediated Liquid-Liquid Phase Separation. *Molecular cell* 79:
443-458 e447

Wen X, Tan W, Westergard T, Krishnamurthy K, Markandaiah SS, Shi Y, Lin S,
Shneider NA, Monaghan J, Pandey UB *et al* (2014) Antisense proline-arginine RAN
dipeptides linked to C9ORF72-ALS/FTD form toxic nuclear aggregates that initiate in vitro
and in vivo neuronal death. *Neuron* 84: 1213-1225

White MR, Mitrea DM, Zhang P, Stanley CB, Cassidy DE, Nourse A, Phillips AH,
Tolbert M, Taylor JP, Kriwacki RW (2019) C9orf72 Poly(PR) Dipeptide Repeats Disturb
Biomolecular Phase Separation and Disrupt Nucleolar Function. *Molecular cell* 74: 713-728
e716

Xu L, Wang D, Zhao L, Yang Z, Liu X, Li X, Yuan T, Wang Y, Huang T, Bian N *et al*
(2023) C9orf72 poly(PR) aggregation in nucleus induces ALS/FTD-related
neurodegeneration in cynomolgus monkeys. *Neurobiol Dis* 184: 106197

Yan X, Kuster D, Mohanty P, Nijssen J, Pombo-Garcia K, Rizuan A, Franzmann TM,
Sergeeva A, Passos PM, George L *et al* (2024) Intra-condensate demixing of TDP-43 inside
stress granules generates pathological aggregates. *bioRxiv*

Yu H, Lu S, Gasior K, Singh D, Vazquez-Sanchez S, Tapia O, Toprani D, Beccari MS,
Yates JR, 3rd, Da Cruz S *et al* (2021) HSP70 chaperones RNA-free TDP-43 into anisotropic
intranuclear liquid spherical shells. *Science* 371

Zhang YJ, Guo L, Gonzales PK, Gendron TF, Wu Y, Jansen-West K, O'Raw AD,
Pickles SR, Prudencio M, Carlomagno Y *et al* (2019) Heterochromatin anomalies and
double-stranded RNA accumulation underlie C9orf72 poly(PR) toxicity. *Science* 363

Zheng S, Sahimi A, Shing KS, Sahimi M (2021) Molecular Dynamics Study of
Structure, Folding, and Aggregation of Poly-PR and Poly-GR Proteins. *Biophys J* 120: 64-72

Zhou Q, Lehmer C, Michaelsen M, Mori K, Alterauge D, Baumjohann D, Schludi MH,
Greiling J, Farny D, Flatley A *et al* (2017) Antibodies inhibit transmission and aggregation of
C9orf72 poly-GA dipeptide repeat proteins. *EMBO Mol Med* 9: 687-702

Zu T, Liu Y, Banez-Coronel M, Reid T, Pletnikova O, Lewis J, Miller TM, Harms MB,
Falchook AE, Subramony SH *et al* (2013) RAN proteins and RNA foci from antisense
transcripts in C9ORF72 ALS and frontotemporal dementia. *Proc Natl Acad Sci U S A* 110:
E4968-4977

 Figure legends

Figure 1. An optogenetic cellular system for controllable *C9orf72* DPR condensation.

(A) Opto-DPR approach.

(B) Opto-DPR condensation can be induced by a single-pulse blue light stimulation. HeLa
cells expressing the respective opto-DPR or empty Cry2olig-mCherry vector were analysed
24 h post-transfection. Blue-light array (single-pulse) was used. Scale bar 10 µm.

(C) Continuous opto-stimulation induces larger opto-PR condensates as compared to single
light pulse. HeLa cells expressing opto-PR or empty vector were subjected to either a single-
pulse or a 3-h continuous blue-light stimulation. 30 cells per condition were analysed from a
representative experiment. **p<0.01, ****p<0.0001, Kruskal-Wallis test with Dunn's post-hoc
test. Scale bar 10 µm.

(D) Super-resolution microscopy (SRM) demonstrates structural differences between
Cry2olig-only and opto-PR assemblies and between the condensates formed after single-
pulse and continuous (3-h long) blue-light stimulation. Representative images and graphical
representation are shown.

(E) Poly-PR, detected using a PR-repeat antibody, is enriched within opto-PR condensate
rims. Representative image is shown.

(F) Opto-PR condensate induction and tracking using confocal longitudinal imaging. Opto-
PR expressing cells were stimulated with a 488 nm laser (at 80% for 500 ms), coupled with
mCherry imaging. Cells were stimulated/imaged either every 2 min ("short-interval") or every
15 min for up to 4 h ("long-interval"). Alternatively, cells with preformed condensates were
imaged every 2 min without stimulation ("dissolution"). Representative images are shown.
Scale bar, 10 µm.

(G) Opto-PR condensate nucleation is concentration-dependent. mCherry fluorescence
intensity was measured in the nucleoplasm of individual cells, outside the nucleolus, and the
number of condensates was quantified in the same cells at the peak of their assembly (7-min
interval repetitive stimulation for 49 min). 75 cells were analysed. Also see Fig.S1C.

(H) FRAP analysis after full opto-PR condensate photobleaching reveals low protein mobility
between the condensate and nucleoplasm. Representative image and FRAP curve for 25
cells from a representative experiment are shown. Scale bar, 10 µm.

(I) Nuclear opto-PR condensates can become persistent. Opto-PR condensate were induced
by 3-h continuous stimulation on blue-light array and the proportion of condensate-positive
cells was quantified immediately post-stimulation or after 3 h of recovery in the dark. N=3
(300 cells in total).

(J) Opto-PR condensates are positive for nucleophosmin (NPM1). Opto-PR condensates
induced by a 3-h continuous opto-stimulation were analysed by SRM. Representative
images and profile plots are shown. Scale bar, 2 µm.

Figure 2. RNA limits opto-PR condensation in the nucleus.

(A) Electrophoretic mobility shift assay (EMSA) with a natural GA-rich RNA sequence
reveals R-DPR binding to RNA. Cy5-labelled synthetic nucleotide "Clip34nt" (34-mer) and
synthetic poly-PR and poly-GR peptides (10 repeats) were used. Representative gel is
shown.

(B) RNA depletion promotes opto-PR condensate growth. Opto-PR expressing cells were
pre-treated with actinomycin D or DRB for 1 h, followed by long-interval repetitive blue-light

stimulation (every 15 min) coupled with time-lapse imaging for up to 4 h (in the presence of
the inhibitor). Representative images are shown. Scale bar, 10 µm.

(C) Opto-PR condensates formed under conditions of actinomycin D-induced RNA depletion
are larger in size and less numerous than in RNA-sufficient cells. Cells were opto-stimulated
for 3 h continuously. Representative images and quantification are shown. Note that the
larger aggregates remain FBL-negative. 30 and 60 cells per condition were included in
analysis for condensate size and number, respectively, from a representative experiment.
**p<0.01, ***p<0.001, Student's *t* test. Scale bar, 5 µm.

(D) Opto-PR condensates formed in RNA-depleted conditions are more persistent. Opto-PR
expressing cells were light-stimulated for 3 h continuously, with or without DRB addition,
followed by DRB removal and recovery for 2 h in the dark. Representative images and
quantification are shown. N=3 (50 cells in total). **p<0.01, Mann-Whitey *U* test. Scale bar, 10
µm.

(E) Opto-PR condensates formed in actinomycin D-treated cells are structurally different, as
revealed by SRM. Representative images of condensates of a comparable size from control,
DRB- or actinomycin D-treated cells induced by 3-h continuous opto-stimulation are shown,
alongside with graphical representation.

(F) Ribosomal (r)RNA depletion from the nucleus and opto-PR condensates in actinomycin
D treated cells. Representative images are shown. Scale bar, 5 µm.

Figure 3. Nuclear opto-PR condensation induces TDP-43 pathology.

(A,B) Opto-PR self-assembly induces nuclear condensation of co-expressed TDP-43 GFP.
Cells were opto-stimulated for 3 h continuously. Representative images (A) and
quantification (B) are shown. Scale bar, 10 µm. N=4 (≥100 cells per biological repeat).
**p<0.01, ***p<0.001, mixed-effects analysis with Sidak's multiple comparisons test.

(C) Spatial proximity of opto-PR and TDP-43 GFP nuclear condensates as revealed by
SRM. Representative image is shown.

(D) Reduced TDP-43 GFP solubility upon induction of opto-PR condensation. Cells
expressing opto-PR were subjected to 3-h continuous blue-light stimulation before lysis and
fractionation. Western blots and quantification of the relative TDP-43 GFP amount in the
supernatant fraction are shown for a representative experiment. S, supernatant; P, pellet.

(E) TDP-43 condensates induced by opto-PR self-assembly are devoid of polyA+ RNA.
Cells after 3-h continuous opto-stimulation were analysed by poly(dT) RNA-FISH.
Representative image is shown. Scale bar, 5 µm.

(F,G) Opto-PR aggregation induces nuclear condensation of a TDP-43 acetylation mimic
mutant (K145Q). Representative images (F) and quantification (G) are shown. 50 cells were
analysed from a representative experiment. *p<0.05, **p<0.01, Student's *t* test. Scale bar, 5
µm.

(H-J) R-DPRs promote TDP-43 clustering *in vitro.* Recombinant TDP-43 (supernatant
fraction depleted of large aggregates; 1 µM) was incubated with equimolar amounts of poly-
PR$_{(x10)}$ or poly-GR$_{(x10)}$ peptides for 10 min or 4 h and analysed by immunostaining and
fractionation/western blot. Methodology (H), immunostaining/imaging (I) and
fractionation/western blotting (J) data are shown. In I, five fields of view from a
representative experiment were analysed, ***p<0.001, ****p<0.0001, one-way ANOVA with

Dunnett's post-hoc test. In J, representative western blot and band intensity quantification for
the supernatant are shown. S, supernatant; P, pellet. Scale bar, 20 µm.
875

Figure 4. Prolonged opto-stimulation elicits cytoplasmic opto-PR/TDP-43 pathology.

(A,B) Prolonged, 24-h blue-light stimulation leads to cytoplasmic opto-PR redistribution and
aggregation, with persistent assembly formation. Proportion of cells with opto-PR assemblies
(nuclear and cytoplasmic) was quantified after 24-h blue-light array stimulation with and
without 8-h recovery in the dark. Representative images (A) and quantification (B) are
shown. Images for a 3-h stimulation and recovery are included for comparison. N=3 (300
cells per condition). n.s., non-significant. Scale bar, 10 µm.

(C) Cytoplasmic opto-DPR foci surround spontaneous stress granules (SGs, visualised
using G3BP1 as a marker). Representative image is shown. Cells were opto-stimulated for
24 h continuously. Scale bar, 5 µm.

(D) Endogenous TDP-43 but, not other RBPs, joins opto-PR assemblies induced by
prolonged opto-stimulation. Also note normal nuclear localisation of all RBPs in cells with
cytoplasmic opto-PR. Cells were opto-stimulated for 24 h continuously. Representative
images are shown. Scale bars, 10 µm.

(E) Cry2only cytoplasmic assemblies are negative for TDP-43. Cells were opto-stimulated
for 24 h continuously. Representative images are shown. Scale bars, 10 µm.

(F) TDP-43 remains demixed within opto-PR within cytoplasmic assemblies, as revealed by
SRM. Cells were opto-stimulated for 24 h continuously. Representative image is shown.

(G) SG assembly and dissolution are affected in cells with cytoplasmically localised opto-PR.
SG were studied in cells subjected to 24-h long opto-stimulation or kept in the dark, prior to
NaAsO2 stress. Cells were analysed either after 15 min of NaAsO2 stress ("assembly") or 3
h into the recovery from NaAsO2 ("dissolution"). Representative images and quantification
are shown. N=3 (120 cells per condition). *p<0.05, **p<0.01, Mann-Whitney *U* test. Scale
bars, 10 µm.

A

Poly-GR(x10)   poly-PR(x10)

μM: 0 0.25 0.5 2.5 5.0 7.5 10.0   0 0.25 0.5 2.5 5.0 7.5 10.0

Cy5-labelled synthetic RNA olig.:
*GAGAGA*GCGCGUGC*AGAGA*CUUGGUGGUGCAUAA

B

0 hr   1 hr   2 hr   3 hr   4 hr

control

act. D

DRB

Repetitive stimulation; 15 min intervals

C

3 hr   Zoom-in

control   PR   FBL

act. D   PR   FBL

Opto-PR / FBL / DAPI

Condensate area (a.u.)

Control   act. D   **

Condensates / cell

Control   act.D   ***

D

control   DRB

3 hr light / 2 hr dark

Opto-PR

% cells with Opto-PR condensates

Control   DRB   **

E

control   DRB   act. D

Opto-PR

500 nm

RNA depletion

F

3hr

control

Opto-PR / rRNA / DAPI

Profile plot

Opto-PR / rRNA

500 nm

act. D

A

Opto-PR | TDP-43 GFP | Overlay + DAPI

Dark

Opto-PR | TDP-43 GFP | Overlay + DAPI

3 hr

B

☐ Unstimulated
☐ 3hr continuous

% cells with nuclear TDP-43-GFP condensation

** *** ****
**** ns

Cry2olig   OptoPR   OptoGR

C

TDP-43 GFP

Overlay

Opto-PR / TDP-43 GFP

500 nm

D

Trans-fection — TDP-43 GFP
Cry2olig | Opto-PR | Opto-GR
Blue light − + − + − +
P S P S | P S P S | P S P S

mCherry
PR-rep
GFP   32% 26% 37% 20% 43% 40%
TDP-43
Total protein

% TDP-43 in pellet fraction

E

Opto-PR | TDP-43 GFP | PolyA

3 hr

F

Opto-PR | TDP-43^{K145Q} GFP | Overlay + DAPI

Dark

Opto-PR | TDP-43^{K145Q} GFP | Overlay + DAPI

3 hr

G

TDP-43^{K145Q} GFP condensate area (a.u.)

*

Control   3 hr

No. TDP-43^{K145Q} GFP condensates / cell

**

Control   3 hr

H

13,000rpm → *Remove pre-formed aggregates*

TDP-43

Small clusters + soluble

poly-PR$_{x10}$ or poly-GR$_{x10}$ + Incubate

Sediment clusters → Fix → IF detection

1000xg → Pellet (P) + Supernatant (S)

I

TDP-43
− | poly-PR$_{(x10)}$ | poly-GR$_{(x10)}$

No. TDP-43 clusters / field

**** ****

Ctrl + PR$_{(x10)}$ + GR$_{(x10)}$

Size of TDP-43 clusters (a.u.)

*** ****

Ctrl + PR$_{(x10)}$ + GR$_{(x10)}$

J

10 min   4 hr
total | Ctrl +PR +GR | Ctrl +PR +GR
S P S P S P | S P S P S P

TDP-43   100% 27% 54% 66% 7% 3%

% TDP-43 in supernatant relative to control

A

3 hr | 3 hr / 3 hr dark

24 hr | 24 hr / 8 hr dark

Opto-PR / DAPI

B

- ▨ Nuclear
- ▨ Cytoplamic
- ▨ Nuclear and cytoplasmic
- ☐ Diffuse

Proportion of cells

ns

24 hr | 24 hr + R

C

Overlay + DAPI | Opto-PR | G3BP1

D

Overlay + DAPI | Opto-PR | TDP-43

Overlay + DAPI | Opto-PR | NONO

Overlay + DAPI | Opto-PR | SFPQ

Overlay + DAPI | Opto-PR | FUS

E

Overlay + DAPI | Cry2olig | TDP-43

F

Individual opto-PR/TDP-43 aggregate/SRM

Overlay | Opto-PR | TDP-43

500 nm

G

SG assembly: *15 min NaASO₂*

Dark | 24 hr

Opto-PR / G3BP1 / DAPI

Proportion of cells

**

Dark | 24 hr

SG dissolution: *1 hr NaASO₂ + 3 hr Recovery*

Dark | 24 hr

Proportion of cells

*

Dark | 24 hr