Understanding the complex interplay between tau, amyloid and the network in the spatiotemporal progression of Alzheimer's Disease

Statistical and computational modeling demonstrates how amyloid and tau interact with each other and the connectome to produce Alzheimer progression observable on human neuroimaging.

Authors: Ashish Raj [1,2] [*] and Justin Torok[1]

Affiliations:

[1]Department of Radiology, University of California at San Francisco, USA

  1 Irving St, AC-116, Box 028, Parnassus Campus, San Francisco, CA 94122

[2]Bakar Computational Health Sciences Institute, UCSF

*Correspondence to: ashish.raj@ucsf.edu

Short title: Co-evolving amyloid and tau in AD

Number of Figures: 6

Number of Tables: 3

Approx Word Count: 6000 (excl Abstract, Refs and legends)

ABSTRACT

It is well known that Aβ and tau proteins are deposited stereotypically in brain regions to cause Alzheimer's disease. The interaction of amyloid and tau in neurodegenerative diseases is a central feature and key to understanding AD pathophysiology. However their mechanisms are controversial, and many aspects do not fit current theories that rely on cell-autonomous factors. While cell culture and animal studies point to various interaction mechanisms between amyloid and tau, their causal direction and mode (local, remote or network-mediated) remain unknown in human subjects. Further, cross-protein interaction is yet to be reconciled with canonical observations that the two species do not co-localize significantly either in space or in time, and do not target the same neuronal populations. To answer these questions quantitatively, in this study we employed a mathematical reaction-diffusion model encoding the biophysical mechanisms underlying self-assembly, trans-neuronal network propagation and enzymtic cross-species coupling of amyloid and tau. We first established that the spatiotemporal evolution of theoretical tau and Aβ correctly predicts empirical patterns of regional Aβ, tau and atrophy. Remarkably, the introduction of a 1-way Aβ→tau interaction was critical to the models' success. In comparison, both the non-interacting and the 2-way interaction models were significantly worse. We also found that network-mediated spread is essential; alternative modes of spread involving proximity or fiber length fare much worse. This mathematical exposition of the "pas de deux" of co-evolving proteins provides crucial quantitative and whole-brain support to the concept of amyloid-facilitated-

tauopathy rather than the classic amyloid-cascade or pure-tau hypotheses, and helps explain certain known but poorly understood aspects of AD.

INTRODUCTION

Alzheimer's disease (AD) involves widespread and progressive deposition of amyloid beta (Aβ) protein in cortical plaques and tau tangles[1,2]. Aβ usually first appears in frontal regions and subsequently spreads to allocortical, diencephalic, brainstem, striatal and basal forebrain regions [2,3]. In contrast to Aβ, tau tangles appear first in locus coeruleus, then entorhinal cortex, followed by an orderly spread into hippocampus, amygdala, temporal lobe, basal forebrain, and isocortical association areas[1]. The dominant "amyloid cascade hypothesis" [7] , posits that amyloid is the upstream factor, whose early abnormal accumulation in the brain causes a cascade of downstream events that recruit misfolded tau. However, this hypothesis has encountered several difficulties, including the spectacular failure of many large clinical trials of amyloid-targeting therapies, and the fact that the temporal and regional distribution of Aβ is quite dissociated from that of tau [4–6]. as well as of downstream atrophy and cognitive deficits. This has led to the search of other mechanisms, especially the role of tau, which is increasingly considered more central to D pathophysiology. However, it is unlikely that tau alone can explain AD pathogenesis, due to overwhelming mechanistic, genetic and demographic evidence of amyloid involvement.

Among potential alternative concepts put forward to reconcile these difficulties, a prominent one is the emerging concept of network-based transmission of both amyloid and tau. Evidence is accumulating in favor of self-assembly and trans-neuronal propagation of amyloid and tau, indeed, of all neurodegenerative pathologies [19–22]. Unlike conventional assumptions about spatial spread of pathology in the brain, this emerging concept implies spread along axonal projections, and the resulting networked spread has repeatedly been substantiated by neuroimaging [9,10,23,24]. Network-mediated spread of amyloid and tau offers an attractive means of reconciling above difficulties: the two species may interact locally, but the effects of these interactions do not remain local, and may propagate across neural circuits to distant regions. Given that the two species have unique and different "epicenters" or seeding loci, this would lead to the apparent observation of their deposition in separate non-overlapping regions. Thus, the cross-species interaction, in the background of network-mediated propagation, may present a plausible framework for understanding AD progression.

Many experimental and mechanistic studies are available that point to a complex interaction between amyloid and tau. It is well known that Aβ facilitates the aggregation of tau and influences the course and severity of downstream atrophy [16–18]. Mechanistic studies point to various modes of interaction [17,30–32].

While these studies provide important mechanistic evidence in model organisms, they may be sensitive to idiosyncratic methodological choices, leading to a diversity of mechanistic possibilities that may not generalize and may not be germane to human disease. Key aspects of brain-wide propagation of tau and amyloid, and the exact mechanistic form their mutual interactions, therefore remain unsupported empirically in human AD. Since experimental testing of mechanistic hypotheses is difficult in humans, here we took a somewhat unorthodox approach, by relying on a *computational* rather than *experimental* interrogation of the mechanistic interaction between tau and amyloid. We first formulated a mathematical model that recapitulates the spatiotemporal evolution of human AD, based on recent advances in the mathematical modeling of reaction-diffusion or network processes that have emerged as a powerful means of evaluating the brain-wide consequences of biophysical mechanisms underlying self-assembly and propagation of neurodegenerative pathologies [8–15]. On top of this base network model, we then built various *interaction models* that allow Aβ and tau species to enzymatically interact in local neural populations. We compared the performance of each interaction model via thorough statistical adjudication, by accumulating model evidence on large neuroimaging (tau and amyloid PET, MRI) datasets of AD spectrum subjects. We devised a fitting procedure to obtain model parameters that best match individual subjects' regional disease patterns. In this manner we were not only able to determine the most well-supported modes of cross-species interactions, but also their applicable kinetic rates. Although spatial and network spread models of single pathological species have been previously reported[8–15], and data-driven models of multiple biomarkers are also available[15,28,29], this study is unique in modeling and evaluating directly on empirical data the network-mediated transmission and interaction of tau and amyloid *jointly*.

We show that both network propagation and amyloid-tau interaction are necessary to recapitulate human AD data. Our data conclusively support that network-mediated spread, with 1-way interaction, whereby amyloid facilitates local tau aggregation, is the most parsimonious and accurate model, yielding correlations above 0.7 against empirical tau topography. Other interactions, including bidirectional ones and those involving enhanced pathology spread instead of aggregation, were not well supported on statistical tests. In totality, this "toxic *pas de deux*" of co-evolving

tau and amyloid pathologies provides critical numerical support to mechanistic hypotheses not possible to be tested directly in humans. Our computational testbed may become an important future tool for the generation and testing of novel mechanistic hypotheses with the potential to obviate expensive and time-consuming mechanistic studies in model organisms.

Figure 1. **Study design, work flow and goals.** Using the structural network connectome (**1**), we deploy the extended Network Diffusion Model (eNDM) that seeks to recapitulate trans-neuronal spread of amyloid and tau pathology (**2**). Startign with the base model of independent eNDMs for both amyloid and tau, we successively introduce various models of interaction between the two (**3**) – these models are listed in **Table 1**. From the ADNI database we extract regional distributions of tau and amyloid SUVr from raw PET images (**4**), whose group average patterns are used to statistically evaluate against each computational model (**5**). Using model evidence criteria we adjudicate between all possible interaction models, and select the best performing one, yielding a numerical assessment of the most likely mode of interaction between amyloid and tau. This adjudicated model is then fit to individual patients' regional pathology data using Bayesian inference of model parameters (**6**). The fitted model is capable of not only predicting the patient's pathology pattern but also their specific seeding sites – a potential marker of translational interest. The model outcomes may in future translational applications be used to predict future pathology, neurodegeneration and cognitive state of the patient (**7**).

MATERIALS AND METHODS

Mathematical Model of Network Transmission

Notation. Brain's anatomic connectivity network is defined on the graph $\mathcal{G} = \{\mathcal{V}, \mathcal{E}\}$ whose nodes $v_i \in \mathcal{V}, i \in [1, N_{roi}]$ represent grey matter structures, and edges $e_{i,j} \in \mathcal{E}$ represent fiber connectivity. Structures v_i comes from parcellation of brain MRI with N_{roi} gray matter regions, and connection strength of the edge between any pair of

regions i,j, $c_{i,j}$ is measured by fiber tractography [61], giving the network connectivity matrix $C = \{c_{i,j}, (i,j) \in \mathcal{E}\}$. Each node has an in-degree and an out-degree given by row and column sums of the matrix C: $\mathbf{d}_{row,i} = \sum_j c_{i,j}$, $\mathbf{d}_{col,j} = \sum_i c_{i,j}$. Define H as the graph Laplacian matrix $H = I - diag(\mathbf{d}_{row} \cdot \mathbf{d}_{col})^{-1/2} C$. This Laplacian definition normalizes each node's connectivity by the geometric mean of its in/out-degree, to accommodate brain parcellations of varying size and connectivity, and emphasizes nodes with high imbalance between in-coming and out-going connections. We define two time-varying vectors containing levels of $A\beta$ and tau at each node: $\mathbf{x}_{A\beta}(t)$ and $\mathbf{x}_\tau(t)$ respectively.

Non-linear reaction + network diffusion with local growth and cross-tau-Aβ interaction

Here we propose a complete and realistic model of pathology dynamics that incorporates the production, growth and spread of both Aβ and tau, as well as the interaction between them.

1) **The No-interaction model**:

Evolution of tau:
$$\frac{d\mathbf{x}_\tau(t)}{dt} = -\beta H \mathbf{x}_\tau(t) + \alpha f_\tau(t)\mathbf{e}_{ERC} \qquad \textbf{(1a)}$$

Evolution of Aβ:
$$\frac{d\mathbf{x}_{A\beta}(t)}{dt} = -\beta H \mathbf{x}_{A\beta}(t) + \alpha f_{A\beta}(t)(\mathbf{m} \cdot \mathbf{x}_{APP}) \qquad \textbf{(1b)}$$

The first term on the right side in Eqs (**1**) above represents network diffusion, following our prior proposal [23], whereby pathology spread occurs by diffusion restricted along network connections. This involves the connectome's Laplacian matrix H and the diffusivity rate constant β. Diffusive spread implies its rate depends both concentration gradients and the amount of connectivity between regions. This model captures trans-neuronal propagation as a connectivity- rather than distance-based process, enacted via active axonal transport followed by membrane- or exocytotic-processes into extracellular space. Fiber length does not enter this model since there is no evidence that axonal transport efficiency is dependent on fiber length. **Evolution of tau**: The second term of Eq (**1a**) introduces a focal seeding event, at the entorhinal cortex (EC), i.e. where all misfolded pathology is produced; for which there is historically accepted evidence that pathology begins in EC. No other region is capable of producing misfolded pathology, but is able to further transmit it. **Evolution of Aβ**: Aβ spread is modeled identically to tau, with the same global diffusivity constant β. Unlike tau, Aβ appears to have a diffuse production mediated by metabolism – this is

encoded by the second term in Eq (**1b**). Since onset of production is gradual, followed by eventual decline and plateauing, we govern the production terms by long-duration gamma-shaped driving functions $f_{A\beta}$ and f_{τ}; see **SI: Note 3** and **Figure S2**. The production of misfolded amyloid is assumed to be proportional to both its baseline metabolism **m** measured via FDG-PET of healthy subjects, as well as the region's pool of available Amyloid Precursor Protein (APP), $\mathbf{x}_{APP}$, from which $A\beta$ is cleaved.

Modeling the network-mediated interaction between tau and amyloid

Several possibilities of cross-interaction exist; here we specify those having the highest empirical or mechanistic support.

2) **1-way interaction model: Amyloid affects tau aggregation**. A new cross-species interaction term was added via coupling coefficient γ to simulate a potential role for amyloid to govern the rate of growth of misfolded tau [16–18]:

Amyloid-facilitated tau aggregation: $\frac{d\mathbf{x}_{\boldsymbol{\tau}}(t)}{dt} = -\beta H\mathbf{x}_{\boldsymbol{\tau}}(t) + \alpha f_{\tau}(t)\mathbf{e}_{EC} + \gamma(\mathbf{x}_{\mathbf{A\beta}}(t) \cdot \mathbf{x}_{\boldsymbol{\tau}}(t) \cdot (1 - \mathbf{x}_{\boldsymbol{\tau}}(t)/K))$ (**2a**)

Note the last term has a logistic growth term of the form $x(1 - \frac{x}{K})$. This term is required to ensure that tau does not grow exponentially indefinitely, and recognizes that realistic tau growth must be limited by overall capacity of healthy tau from which toxic tau can be cleaved. The capacity level is controlled by the parameter K. The equation for amyloid evolution (**2b**) remains the same as (**1b**).

3) **1-way interaction model: Amyloid affects tau diffusion into the network**. In this model amyloid does not influence tau aggregation but increases the rate of tau diffusivity into thE network in areas where amyloid id high and less diffusivity in areas where amyloid is low. This model simply modulates the scalar diffusivity constant in Eq (**1a**) by allowing it to be regionally-varying:

Amyloid-facilitated tau diffusion: $\frac{d\mathbf{x}_{\boldsymbol{\tau}}(t)}{dt} = -\beta \cdot diag\{\mathbf{x}_{\mathbf{A\beta}}(t)\} \cdot H\mathbf{x}_{\boldsymbol{\tau}}(t) + \alpha f_{\tau}(t)\mathbf{e}_{EC}$ (**3a**)

In this model tau in every region experiences a different net diffusivity, depending on the local level of amyloid. The equation for amyloid evolution (**3b**) remains the same as (**1b**).

4) **Tau to amyloid interaction model**. The reverse effect of tau on Aβ production does not have firm pathophysiological support [16]. However, for completeness we also explored the alternative interaction via the coefficient η. The equation for tau evolution (**4a**) remains the same as (**1a**).

Tau-facilitated Aβ: $\quad \frac{d\mathbf{x_{Aβ}}(t)}{dt} = -\beta H \mathbf{x_{Aβ}}(t) + \alpha f_{Aβ}(t)(\mathbf{m} \cdot \mathbf{x}_{APP}) + \eta(\mathbf{x_{τ}}(t) \cdot \mathbf{x_{Aβ}}(t) \cdot (1 - \mathbf{x_{Aβ}}(t)/K))$ $\qquad$ (**4b**)

5) **Two-way interaction model**. Finally we included in our analyses a bidirectional model whereby both species influence each other. Hence Eq 5a for tau evolution is the same as Eq (**2a**) and Eq (**5b**) for amyloid evolution is the same as Eq (**4b**).

6) **Proximity-based spread**. We simulated proximity-based spread using the same formulation as Eqns (**1a** and **1b**), but replacing the connectome-derived Laplacian with adjacency matrix computed from inter-regional Euclidean distance: $C_{dist} = \exp\left(-\frac{D}{\sigma}\right)$, where $D = \{d_{ij}\}$ is the matrix of distance between regions i,j, and σ is the standard deviation over all elements of D. Laplacian H_{dist} was defined analogously to the connectome Laplacian H, and replaced that term in Eq (**1**). We also evaluated a Gaussian relationship instead of exponential, but the results were generally worse and were not further evaluated. Denote the full model by Eq (**6a**, **6b**).

7) **Fiber distance-driven spread**. We simulated spread along fiber projections using the same formulation as Eq (**1**), but replacing connectivity with fiber distance, as per $C_{fiberdist} = \exp\left(-\frac{L}{\sigma}\right)$, where $L = \{l_{ij}\}$. Here l_{ij} is the average length of streamlines between regions i,j. Laplacian $H_{fiberdist}$ was defined analogously to the connectome Laplacian H. Denote the full model by Eq (**7a**, **7b**).

Implementation. The above theoretical models were numerically solved using MATAB's **ode45()** solver, jointly for both amyloid and tau, even in those cases when some parameter choices would lead to decoupled equations. Initial condition was set such that $x_{\tau,Aβ}(t = 0) = 0$; however note that the seeding and production of the two species are non-zero, via the term $f(\cdot)$.

Experimental Design

Subjects and data. Data used in this study were obtained from the ADNI [33] database (http://adni.loni.usc.edu); consisting of 531 ADNI-3 subjects who had at least one exam of all three: MRI, AV1451-PET and AV45-PET, available by 1/1/2021. Demographic information is in **Table S1**. These data were processed to obtain regional statistics of pathology and atrophy, the latter used in this analysis as a surrogate measure for tau tangles/neurodegeneration. Anatomic connectomes were computed from healthy diffusion MRI and tractography algorithms. The primary dataset was evaluated on the 86-region Desikan atlas and the Supplementary dataset on the 90-region AAL atlas, using similar processing pipelines; see **SI: Note 1**. To remove AV1451-PET scans' considerable non-specific binding and the effect of iron in thalamus and striatum, their values were removed from subsequent analysis. The proposed network model was applied to canonical healthy connectomes, and model patterns compared against the ADNI regional data.

Statistical Analysis and model testing

Both group statistics (t-statistics for each of EMCI, LMCI and AD groups) and individual fitting was performed. Empirical regional AV1451-PET tau was supplemented with MRI-derived regional atrophy to leverage larger sample size; atrophy is a useful surrogate for tau, with strong association [44]. The statistical test of choice is Pearson correlation strength, R, and its two-tailed p. Detailed model fitting to individual subjects is described in **SI: Note 6**.

We corrected each Pearson p-value with an inflation factor of 50 – the number of unique comparisons. We also performed extensive permutation tests as noted in earlier with 500 random permutations. A comparison of the reported R with "null" distributions of randomized R gives a pertinent and additional measure of significance. We further obtained Fisher's transform and its p-values to assess significant improvements between models. AIC scores were also computed to assess whether different models with varying complexity give comparable performance.

Data and code Availability

All patient data used in this study were obtained from the public ADNI study. Users interested in obtaining these data can do so by directly visiting the website http://adni.loni.usc.edu. Nonetheless, for the purpose of convenience and to facilitate experimenting with the presented approach, group regional SUVr, atrophy and metabolism, etc are being made available within the code repository, and can be found at https://github.com/Raj-Lab-UCSF/Aggregation-Network-Diffusion . There are no restrictions on the use of this data. All computer code and group statistics used in producing the current results will be shared publicly and without limitations at our laboratory's GitHub site:

RESULTS

I. Cross-sectional relationships between tau, amyloid and atrophy

The public ADNI3 study [33] was processed with home-grown software pipelines to obtain reginal values of the three imaging biomarkers: atrophy. Tau and amyloid; see demographic data in **Table S1**. First we ascertained what the ADNI data at large are telling us about how the three imaging biomarkers are related to each other, and to the network. In **Figure 2** we depict these relationships across the whole ADNI database of how induvial subjects triplets (tau, atrophy, amyloid) are related. **Figure 2A** show the three biomarkers' associations –in the scatter plots in top row each dot represents a single subject's single brain region. At this aggregate level clearly there is a moderate yet significant relationship between atrophy and tau, but not between atrophy and amyloid. There is a strong relationship between tau and amyloid – cementing the prior expectations that amyloid and tau are colocalized. The bottom row shows histograms of the correlation strengths of the three disease groups (EMCI, LMCI, AD) separately. It demonstrates a prominent stage-dependent effect, whereby atrophy is more tightly related to tau in later than earlier stages. The situation is reversed for the tau-amyloid associations: earlier stages have a stronger association than later stages. These data point to the well-known finding that amyloid plays a role early in AD pathophysiology, and at later stages it has a plateauing behavior and is no longer predictive.

Stage-dependent relationship to network connectivity

Figure 2B shows analogous results, but with the x-axis depicting the network connectivity strength of each region to the canonical seed region of bilateral ERC. This data attempts to test at the population level the hypothesis that biomarkers are predicted by connectivity to the pathology origination site. At the aggregate level we find moderate but significant evidence for an association between EC-connectivity and all three biomarkers; however the association with amyloid is in fact negative. These results are also stage-dependent; with earlier stages giving a stronger association with connectivity for atrophy and tau, and the reverse for amyloid.

Longitudinal relationships between tau, amyloid and network connectivity

The longitudinal behavior of tau was assessed using the first difference between baseline and year-1 visit, the most generalizable longitudinal measure for subjects with varying number of visits. **Figure 2C** shows these relationships at the aggregate level (top) and for each stage (bottom). A moderate but significant association with change of tau was found for baseline tau, but not baseline amyloid. We then generalized the network hypothesis to assess whether baseline pattern of tau weighted by each region's network connectivity would predict the change of tau. This hypothesis involves the possibility of remote effect, whereby tau at a region may after the change of tau at a distant but connected region. We used a simple model of spread of tau from baseline pattern along the network, given by the well-known network diffusion model (Raj 2012). As shown in the figure, this hypothesis was found to have significant support for baseline tau, but not for amyloid. As with earlier results, there is a distinct stage dependency, such that earlier stages show a stronger association for baseline tau but not for amyloid.

Taken together, these results demonstrate that there is a significant cross-sectional and longitudinal relationship between tau and amyloid distributions in the Alzheimer brain, and this relationship is stage-dependent.

Figure 2. **Cross-sectional and longitudinal relationships between tau, amyloid, atrophy and the network.** **2A**: Scatter plots (top) and histograms of correlation strengths (bottom) are shown. There is a moderate yet significant relationship between atrophy and tau and amyloid and tau, but not between atrophy and amyloid. There is a distinct stage-dependent effect whereby later stages give stronger associations with respect to tau but the opposite for amyloid. **2B**: Using bilateral ERC as the canonical pathology seeding site, it was found that its network connectivity to a region is moderately associated with atrophy and tau but negatively associated with amyloid. Accompanying

histograms suggest a stage dependency, with earlier stages giving a stronger association with connectivity for atrophy and tau, and the reverse for amyloid. **2C**: change of tau from baseline to year 1 is associated moderately with baseline tau but not baseline amyloid. To assess the hypothesis that baseline pathology may affect change of tau in a distant but connected region we employed the simple association proposed by the well-known network diffusion model: $\frac{d\tau}{dt} \propto -L\tau$, where L is the network Laplacian matrix (see model and cite ??). This remote-association was significant for baseline tau but not for amyloid, and again there was a stage-dependent behavior for tau.

II. Evidence-based development and adjudication of competing hypotheses of protein aggregation, network transmission and interaction

The above data snapshots provide empirical support to the hypothesis of a cross-species interaction between Aβ and tau; however it does not point to the exact mechanistic form these interactions might take, nor does it give a model of brain-wide protein propagation. Whether these data are consistent with connectome-mediated spread, or with other modes f spread, e.g. based on proximity or along fiber distances, is also unclear. In this section we will first develop a base model of network transmission of tau and Aβ. From the base model we will create various extensions that will implement different potential interactions, as listed in the Model section of Methods. **Table 1** summarizes these 7 competing hypotheses, and the computational model used to assess and adjudicate each hypothesis quantitatively.

Table 1: **Alternative/competing hypotheses of tau-amyloid interaction to be tested in the background of transmission mediated by network connectivity**. Brief description of each hypothesis is given, along with the mathematical differential equation corresponding to it. References to supporting literature for each hypothesis is provided. Two additional models of spread are given, pertaining to spread via proximity and spread via fiber distance – for these models we use the base case of no interaction between tau and amyloid.

Hypothesis	Description	Model Equation	Supporting Reference
1) No interaction	Tau and amyloid beta propagate independently on the connectivity network (Base model)	1a, 1b	
2) $A\beta \rightarrow tau$ aggregation	Tau and amyloid propagate on the network, but local presence of amyloid enzymatically facilitates the aggregation of tau	2a, 2b	
3) $A\beta \rightarrow tau$ diffusion	Tau and amyloid propagate on the network, but local presence of amyloid increases the spread/transmission of tau	3a, 3b	
4) Tau → $A\beta$ aggregation	Local presence of tau enzymatically increases the rate of aggregation of amyloid	4a, 4b	

5) Tau ←→ $A\beta$	Tau and amyloid propagate on the network, and the local presence of each protein enzymatically facilitates the aggregation of the other protein	5a, 5b	
6) Proximity-based transmission	Tau and amyloid beta propagate within the brain based on proximity (Euclidean distance between regions) rather than via the connectome. They are independent of each other (i.e. under the base model 1 of no-interaction)	6a, 6b	
7) Fiber Distance-based transmission	Tau and amyloid beta propagate within the brain based on Fiber distance between regions rather than via the connectome. Fiber distance is the aergae length of white matter tracts that cnnect any two regions. They are independent of each other.	7a, 7b	

Illustration of base model of joint evolution of Aβ and tau

The numerical solution $(\mathbf{x}_{A\beta}(t), \mathbf{x}_\tau(t))$ of the joint Aβ-tau model (Eqs **1a,b**) was evaluated on the canonical healthy connectome under the Desikan-Killiany parcellation (86 regions). In order to assess the model behavior broadly, in the following set of results we used a canonical model specification, given by the coarsely optimized (default) parameter values. Please refer to **SI: Note 6** and **Table S2** for details. **Figure 3** left column shows the evolution of amyloid regional distribution, as it propagates into the structural network, eventually establishing itself in frontal and medial areas of the brain. Model amyloid evolution appears to recapitulate the classic amyloid progression, e.g. Thal's five pathologic stages [2] and amyloid PET patterns [3]; proceeding from medial frontal and precuneus, areas with high baseline metabolism, into the wider network, only slowly entering temporal cortices. The middle column shows the evolution of tau on the same network, starting from seeding event in the bilateral ERC.

Network transmission of Aβ and tau, with cross-species interaction

Each of the 7 interaction models, encompassing the most common mechanistic hypotheses in the field, were implemented as described in Model section. For illustration purposes here we describe the most promising of those, given by the 1-way interaction from amyloid to tau; specifically: **Aβ-facilitated tau aggregation** [16–18] encompassed in (Eq **2**). The resulting tau progression, illustrated in **Figure 3A** right column, remained confined to the medial temporal lobe until sufficient levels of Aβ had spread to and accumulated there (at $t \approx 10$). There onward, in contrast to the "pure tau" evolution, it took on a more aggressive trajectory, spreading first to nearby limbic, then basal forebrain, then parietal, lateral occipital and other neocortical areas, in close concordance with Braak's six tau stages [1]. Thus, the facilitation of tau by amyloid does not lead to colocalization of the two until late stages, helping explain why regional Aβ patterns do not coincide with tau and atrophy patterns. Conversely, the absence of the interaction term

led modeled tau to remain confined to the temporal lobe (**Figure 3A**, middle), mirroring primary age-related tauopathy (PART), a new classification for mild neurofibrillary degeneration in the medial temporal lobe, but no Aβ plaques [34]. For comparison, the AAL-connectome-based model evolution is shown in **Figure S4**, with very similar behavior, indicating that the choice of atlas or processing pipeline did not drive results.

A

B

C

Figure 3. **The evolution of model dynamics of amyloid and EC-seeded tau**. **A**: **Spatiotemporal evolution** of the model evaluated by numerical integration of Eqs (1a, 2a), on the 86x86 healthy Desikan connectome using default model parameters. The first column shows the evolution of amyloid, the second column of amyloid-facilitated tau and the third column of tau only, putatively a model of primary age-related tauopathy (PART), which remains confined to the temporal cortex and is relatively lower in magnitude compared to AD. In comparison, the amyloid-facilitated tauopathy model (middle column) shows much more profuse tau deposition in temporal areas with the arrival of amyloid into temporal cortices, putatively recapitulating AD tau. Each sphere represents a brain region, its diameter is proportional to pathology burden at the region, and is color coded by lobe (blue = frontal, purple = parietal, red = occipital, green = temporal, cyan = cingulate, black = subcortical). **B**: **Predicting Braak stages using computational model.** Using the time-of-arrival calculation of each group's fitted eNDM model, we developed a 6-stage "computational Braak" staging system. The predicted Braak stages strongly agree with the original Braak stages, applied to the DK atlas parcellation, achieving $R = 0.76, p < 10^{-6}$ for the best-adjudicated model (right). In comparison, the non-interacting model (left), while being significant at $R = 0.64$, is substantially worse, suggesting that amyloid→ tau interaction is a necessary factor in Braak staging. **C**: **Gestalt of spatiotemporal progression of AD-related amyloid and tau**. Following diffuse production of amyloid **(red)** in proportion to metabolism, and focal production of tau **(blue)** at the EC, aggregation into plaques and tangles occur at networked sites following graph topology. Tau pathology is further aggravated by the arrival of amyloid in temporal areas. Finally the classic AD spatial patterns are established – frontal-dominant for amyloid and temporal dominant for tau.

Supplemental Figure S3 shows the global accumulation of theoretical pathology over model time, evaluated at default parameters. All proteins increase over time, but amyloid-facilitated tau diverges dramatically from the non-facilitated tau at around t=15, mirroring **Figure 3**. **Figure S3** makes it clear that while the "pure" tau model also captures empirical data, it does so far slower and achieves far less correlation strength than the facilitated version.

Predicting Braak stages using computational model. Using the time-of-arrival calculation of each group's fitted eNDM model, we developed a 6-stage "computational Braak" staging system. The predicted Braak stages strongly

agree with the original Braak stages, applied to the DK atlas parcellation, achieving $R = 0.76, p < 10^{-6}$ for the best-adjudicated model (right). In comparison, the non-interacting model (left), while being significant at $R = 0.64$, is substantially worse, suggesting that amyloid$\rightarrow$ tau interaction is a necessary factor in Braak staging.

To summarize these impressions, we find that **amyloid$\rightarrow$tau** interaction is an essential component of the spatiotemporal propagation of the two species, without which the computational model does not fully recapitulate empirical amyloid and tau spatial patterns in patients. An overall picture that emerges can be depicted (**Figure 3C**) whereby diffuse production of amyloid, potentially in proportion to metabolism, and focal production of tau at the EC, propagate into the network, even as tau pathology is further aggravated by the arrival of amyloid in temporal areas.

Group level empirical validation and model fitting

We next performed model fitting on empirical group data of the proposed model initiated at the canonical EC-seeding of tau. For model fitting we developed a robust maximum a posteriori (MAP) inference procedure described in detail in **SI: Note 6**. Cross-sectional group ADNI data were correlated against the fitted model's evolution ($\mathbf{x_{A\beta}}(t)$, $\mathbf{x_{\tau}}(t)$) at every time t, and Pearson's R was recorded. The resulting "R-t curves", shown in **Figure 4** displayed a characteristic peak as more amyloid and tau pathology diffused into the network it increasingly recapitulated cross sectional data. Subsequently model diverged from empirical pattern, decreasing R. Since our model posits that both amyloid and tau evolution happens on the same time axis, we report the model instant that maximized the posterior, rather than peak R for either tau or amyloid separately. The R value resulting from this "shared time", called t_{max}, is recorded for both amyloid and tau (denoted by square markers) and considered as model evidence. Optimal fitted parameters shown in **Table S2** indicate substantial differences between groups. **Table 2** tabulates the resulting R statistics. All reported R values that are moderately to highly significant as denoted by * (p < 0.05) and ** (p < 0.005), **corrected for multiple comparisons.** Statistics for all model forms are contained in **Table 2**; however for illustration purposes below we highlight the best, most parsimonious interaction model (1-way amyloid $\rightarrow$ tau interaction).

Amyloid. Correlations between model and empirical AV45 SUVr are shown in **Figure 4** and **Table 2**. The peak R was: $R = 0.71$ (EMCI), $R = 0.49$ (LMCI) and $R = 0.70$ (AD). All correlations were highly significant. R-t curves shown suggest that the correspondence reaches a peak and then plateaus. Note that the model time t has arbitrary units that

may not be directly comparable to empirical duration in years. **Tau.** The network propagation of modeled tau starting from its seeding in EC was computed and the correspondence between model-predicted regional tau deposition and group-average empirical AV1451-PET data, was assessed (**Figure 4, Table 2**). Unlike the amyloid case, the R-t curves of tau and atrophy show a slow and steady rise and peak without a plateau effect. The R between the model at $t = t_{max}$, and empirical tau PET data, was: $R = 0.51$ (EMCI), $R = 0.75$ (LMCI) and $R = 0.73$ (AD). The latter two correlations were extremely highly significant. Striatal AV1451 uptake is considered non-specific binding and was suppressed in these comparisons. Correlation with EMCI group is poor, likely due to lower PET uptake and inter-subject heterogeneity. **Atrophy**. Since tau is highly co-localized with regional atrophy, we also show a comparison with MRI-derived group atrophy. However, we did not fit the model again to atrophy, instead borrowing the tau-fitted models for each cohort, under the assumption that tau is the primary, and atrophy is a surrogate for tau. The resulting R between the model at $t = t_{max}$ and empirical atrophy was: $R = 0.37$ (EMCI), $R = 0.60$ (LMCI) and $R = 0.66$ (AD). EMCI atrophy gives the lowest correlation, likely due to the low effect size measurable on MRI in this cohort.

Time of peak: The t_{max} of maximum posterior followed the expected order: t_{max} for AD > LMCI>EMCI. It is challenging to fit an accurate time axis to empirical data, since it does not have a measure of pathology duration – which may never be known. This order also supports the ability of the fitted time of arrival as an accurate Braak staging of the disease (see above). Interestingly, the peak for tau occurs more than a decade (in model "years") after the plateau seen for amyloid – perhaps recapitulating well-known clinical and pathological examinations that suggest a decade-long delay between the two processes.

Figure 4. **Validating evolution of model tau seeded at ERC against empirical data**. **Left column**: The top panel shows the behavior of model evidence R against model time t, suggesting that the model reaches a peak correspondence with empirical data relatively early, then plateaus. The model time t_{max} at which the model was evaluated against empirical data is denoted by square markers. The "glass brain" renderings show empirical amyloid-PET SUVr patterns from all 3 diagnostic groups. The R statistic between the model and empirical data are shown alongside. **Middle column:** The top panel shows that the R between the model and empirical tau data grows steadily, hits a peak, then declines. Notice that the peak for tau occurs more than a decade (in model "years") after the plateau seen for amyloid. R is strongest for LMCI and AD and weakest for EMCI. The cross-sectional empirical tau-PET SUVr patterns from all 3 diagnostic groups are shown. The **right-most column** shows results for MRI-derived atrophy. The best-fitting model amyloid and tau patterns appear strikingly similar to the empirical PET data, which a quick visual comparison of **Figures 3** and **4** will confirm. The match between the model and empirical data are strong for LMCI and AD, and weakest for EMCI. Note that t_{max} for each case is identical between amyloid, tau and atrophy – this point may not always coincide with the actual peak R for a given case.

Quantitative adjudication of different models of amyloid-tau interaction

We evaluated five theoretical interaction models: **1)** No-interaction model Eqn (**1**); **2)** 1-way interaction model (Eq **2**), whereby tau affects amyloid but not vice versa; **3)** 1-way interaction whereby amyloid affects tau aggregation but not

vice versa (Eqs **3**); **4)** 1-way interaction whereby amyloid affects tau diffusion into the network, but not vice versa (Eqs **4**); and **5)** 2-way interaction model (Eq **5**). Each model was evaluated in identical fashion after full MAP inference and identification of a single unique operating time t_{max} that spans both amyloid and tau data. Statistical comparison of these fitted models is shown in **Table 2**. Both the fitted Pearson's R is reported, along with model comparison using Fisher's R-to-z transform and the p-value of significant differences between models; p<0.05 are highlighted.

These are our findings: The best overall model is the 1-way interaction **$A\beta\rightarrow$ tau aggregation**. In contrast, the 1-way interaction $A\beta\rightarrow$ tau diffusion does not significantly improve upon the base (No-interaction) model, suggesting that amyloid primarily facilitates the aggregation of tau, not its enhanced spread into the network. The 2-way model is insignificantly different from the 1-way model. Both interaction models are significantly better than the no-interaction model. Interestingly, 1-way tau$\rightarrow A\beta$ model is indistinguishable from No-interaction model, suggesting that tau does not exert a clinically relevant effect on amyloid.

To confirm these impressions and to rule out improved performance due to higher degrees of freedom of additional coupling coefficients, we obtained the Aikeke information criterion (AIC) for each model. Model order was defined as the number of non-zero parameters. The lowest AIC is always achieved by 1-way $A\beta\rightarrow$ tau interaction model, supporting our earlier conclusion. AIC for $A\beta\rightarrow$ tau interaction is always higher than for tau$\rightarrow A\beta$ or 2-way interaction, confirming that additional tau$\rightarrow$amyloid interaction does not further improve model fit. The five models are ordered descendingly in AIC:

(Best) $A\beta\rightarrow$tau aggregation (local) **>** $A\beta\rightarrow$tau aggregation (remote) **>** tau$\leftarrow\rightarrow A\beta$ **>** $A\beta\rightarrow$tau diffusion **>** no-interaction **>** tau$\rightarrow A\beta$ aggregation **(Worst).**

The amyloid component is equally strong for all models and its parameters are largely unaltered – this might reflect the well known plateau effect of amyloid, although further investigation will be warranted.

Table 2: **Peak Pearson's R of correlation between model and empirical regional statistics**. Highly significant correlations: ** (p < 0.001), moderately significant: * (p < 0.01), after correction for multiple comparisons – i.e. inflation by n=50. Six theoretical interaction models' evaluations are presented: the No-interaction model, without the interaction term ($\gamma = 0$); the 1-way interaction model, whereby tau affects amyloid but not vice versa; another 1-way model whereby amyloid affects tau but not vice versa; and the 2-way interaction model where both tau and amyloid affect each other via γ. Three different modes of causal interaction from amyloid to tau are given: where amyloid influences the diffusion of tau; where it influences only the aggregation of tau; and where amyloid influences tau aggregation remotely, via long-range fiber connections. Aikeke information criterion (AIC) is reported for each model in brackets. There is no difference between models in amyloid fits. For tau and atrophy, the lowest overall AIC (**boldface**), and highest R (**boldface**), are achieved by the 1-way interaction from amyloid to tau aggregation; consequently it is highlighted by green box around it. The remote-interaction and 2-way interaction models are insufficiently different from this 1-way model. Comparisons between these models via Fisher's t-test are contained in **Table S3**.

Empirical data	No- interaction Model	1-way interaction, tau→$A\beta$ aggregation	1-way interaction, $A\beta$→tau diffusion	**1-way interaction, $A\beta$→tau aggregation**	1-way (remote) interaction $C \cdot A\beta$→tau aggregation	2-way interaction $A\beta$←→tau
EMCI Aβ	0.71 ** (163)	0.71 ** (163)	0.71 ** (163)	0.71 ** (163)	0.71 ** (163)	0.71 ** (166)
LMCI Aβ	0.49 * (223)	0.49 * (223)	0.49 * (223)	0.49 * (223)	0.49 * (223)	0.45 * (229)
AD Aβ	0.70 ** (333)	0.70 ** (333)	0.70 ** (333)	0.70 ** (333)	0.70 ** (333)	0.70 ** (335)
EMCI tau	0.33 (200)	0.38 (202)	0.35 (201)	**0.48** * (**190**)	**0.48** * (192)	0.36 (203)
LMCI tau	0.60 ** (232)	0.60 ** (234)	0.49 * (247)	**0.75** ** (**206**)	0.73 ** (211)	0.70 ** (219)
AD tau	0.57 ** (322)	0.51 * (332)	0.52 * (331)	**0.73** ** (**296**)	0.71 ** (301)	0.65 ** (315)
EMCI atrophy	0.34 (**171**)	0.32 (176)	0.33 (175)	0.35 (174)	0.33 (178)	**0.37** (175)
LMCI atrophy	0.60** (**158**)	0.60** (162)	0.56* (171)	**0.62** ** (**158**)	0.61 ** (159)	0.48 * (184)
AD atrophy	0.52* (316)	0.38 (331)	0.39 (330)	**0.66** ** (**299**)	0.62 ** (304)	0.58 ** (312)

Permutation testing to demonstrate disease specificity

Various permutation tests were deployed to determine whether the presented model only recapitulates empirical regional distributions when it is applied in the correct region order and to the correct human connectome, contained in **SI: Notes 7** and **Figures S5** and **S6**. Under 500 random permutations of atrophy and tau (**Figure S5-A**), and under 500 random permutations of the connectome itself (**Figure S5-B**) we found that the above-reported R values of the true data are highly significant compared to these "null" distributions ($p < 10^{-3}$ for all groups). This was true whether for Pearson correlation or Spearman (**Figure S6**) as the performance metric.

Comparison of different modes of spread

We implemented two alternative modes of spread: a) inter-regional pathology spread depends only on the shortest Euclidean distance; and b) transmission between regions is inversely proportional to the average length of fiber projections between regions. See **Methods** and quantitative comparison in **Table 3**. Again, the model in each case was refitted using the MAP estimator, hence each fitted model may represent the best-case scenario for that hypothesis. P-value of Fisher's R-to-z transform of significant differences between models was evaluated, accounting for the correlation between dependent models. We found that : **a)** For both tau and amyloid evolution, connectome-mediated spread model gave the closest correspondence to empirical data; and **b)** Euclidean spatial spread was slightly but not significantly superior to fiber distance-based spread. For these comparisons we chose the best interaction model identified in **Table 1** ($A\beta \rightarrow tau$ aggregation) for all three networks. However, we repeated these comparisons using other interaction terms as well, and did not find significant differences in performance (Fisher's p-value = N.S.).

Table 3: **Pearson's R of correlation between alternative spread models and empirical regional statistics**. Fisher's R-to-z transform was computed between the connectome and the alternative modes of spread, and its p-value is shown alongside. The connectome-mediated model significantly outperforms the two distance-based models (p values denoted in red font).

Empirical data	$A\beta \rightarrow tau$ interaction, Connectivity	Fiber Distance	Fisher R-to-z, Fiber vs C	Euclidean Distance	Fisher R-to-z, Euclidean vs C
EMCI Aβ	0.71	0.56	**p < 0.01**	0.57	**p < 0.01**
LMCI Aβ	0.49	0.43	p = 0.10	0.42	p = 0.09
AD Aβ	0.70	0.56	**p < 0.01**	0.57	**p < 0.01**

EMCI tau	0.48	0.25	**p < 0.01**	0.24	**p < 0.01**
LMCI tau	0.75	0.31	**p < 0.01**	0.35	**p < 0.01**
AD tau	0.73	0.34	**p < 0.01**	0.36	**p < 0.01**
EMCI atrophy	0.35	0.00	**p < 0.01**	0.18	**p = 0.03**
LMCI atrophy	0.62	0.29	**p < 0.01**	0.41	**p = 0.025**
AD atrophy	0.66	0.17	**p < 0.01**	0.22	**p < 0.01**

Translational aspects

Having established the network model's validity on group data, and having determined which modes of the tau-amyloid-network interactions are relevant and empirically supported, we devote the last section to showcase two key results related to translational aspects in patients: etiologic heterogeneity and the capacity to predict an individual subject's spatiotemporal trajectory of AD. These results will serve as the foundation of future diagnostic and prognostic applications.

A) Uncovering etiologic heterogeneity: Repeated seeding to assess alternative seeding sites

Entorhinal cortex (EC) was chosen above as the canonical tau seeding site. To establish other regions' other regions' seeding plausibility we repeatedly simulated the selected network interaction model seeded from every possible region bilaterally. For each seed region, peak Pearson's R between model and ADNI tau PET data is shown in **Figure 5A**. EC is amongst the best overall cortical seeding site, while hippocampus (HP) is the best subcortical site. Other prominent seeding sites include Parahippocampal gyrus (PHP) and Fusiform gyrus (Fus), which are adjoining EC and appear frequently similar in tau uptake to EC on PET imaging. Thus the quantification of these structures as likely seeding locations affirms our model's relevance and plausibility. The evolution of pathology from some of these non-EC sites are illustrated in **Figures S8** and **S9**, for HP-seeding and IT-seeding, respectively. They remain substantially similar to **Figure 3**; however a notable difference from EC seeding is that HP seeding leads to higher involvement of medial temporal and subcortical structures. Amongst the above 5 regions, EC is unique in giving consistently one of the highest seeding likelihood despite having less levels of empirical tau deposition than others.

A

Figure 5. Translational applications. A: Bar charts of repeated seeding of each region in turn. For each seed region, peak Pearson's R over model time, corresponding to the correlation between the best interaction model (amyloid-facilitated tau aggregation) and empirical tau PET data, is shown, for each of the 3 diagnostic groups: EMCI

(top), LMCI (middle) and AD (bottom). Entorhinal cortex (EC) is amongst the best cortical seeding sites and Hippocampus (HP) the best subcortical site. Other likely seeding sites include Fusiform (Fus) and Parahippocampal gyrus (PHP). **B**: **Fitting model to individual subjects.** Histogram of Rmax achieved by fitting the model parameters to individual subjects' baseline regional tau data, after imposing a common seeding site located at the ERC (Left) or after using the best seeding site for each subject (Middle). Amyloid histograms are shown on Right panel. Here the best model combination from **Table 2** was used: 1-way interaction ($A\beta \rightarrow tau$) on the structural connectivity graph, and the evaluation was performed on tau-PET scans in ADNI3. Average Rmax over all subjects within each diagnostic group is shown in red dashed line.

B) Individual subject fitting and prediction

For translational applications we must necessarily move away from above group comparisons to model inference on individual subjects, while accommodating both etiologic heterogeneity as well as potentially subject-specific model parametrization. In order to achieve this we deployed our robust the maximum a posteriori (MAP) inference procedure (see **SI: Note 6**) on individual subjects' multimodal regional biomarker data from ADNI-3 (see **Table S1** for demographic details), under our best-adjudicated model (i.e. network-mediated spread of tau and the 1-way interaction best $A\beta \rightarrow tau$ aggregation). **Figure 5B** gives the histograms of Pearson correlation R_{max} between observed and model-predicted regional tau distribution in each subject. For comparison we show the results of both canonical (entorhinal) seeding and each individual's best seeding site. EC seeding was significantly worse (paired t-test after Fisher R-to-z: $p < 10^{-7}$, corrected), implying that a common seeding site may not be appropriate for all subjects, and revealing a potential etiological factor behind observed heterogeneity in AD-spectrum subjects. With individual-specific seeding and model fitting, we were able to achieve excellent prediction of the subjects' tau distributions, with Rmax ranging widely up to a maximum of 0.9 and mean of 0.65 for LMCI and AD, and slightly lower for EMCI subjects. The mean Rmax across individuals in **Figure 5B** roughly tracks, but slightly lower than, the Rmax achieved on the group-averaged tau data of **Figure 4** and **Table 2**; which is expected due to heterogeneity as well as measurement noise in individual subjects. It is also noteworthy that canonical EC seeding fails in approximately half of the subjects in eMCI and LMCI cohorts ($e.g. R_{max} < 0.3$) but only a small minority of diagnosed AD patients: indicating that higher stages have lower etiologic heterogeneity.

DISCUSSION

The protein-protein interaction of amyloid and tau in neurodegenerative diseases is a central feature and key to understanding AD pathophysiology [17,30–32], now commonly called the A-T-N model [38]. However, this model has been difficult to reconcile with the observations of dissociated spatial distribution of tau and amyloid. Critical aspects of brain-wide propagation of tau and amyloid, and the exact mechanistic form their mutual interactions, remain unsupported empirically in human disease. Hence the cause-effect mechanisms by which amyloid and tau regulate each other and cause downstream neurodegeneration and symptomatology, remain poorly understood. Prior mechanistic studies in model organisms reveal a plethora of potential interactions, yet these possibilities are difficult to test directly in humans. This is problematic since animal models do not fully recapitulate human disease. Further, data from experimental systems are sensitive to idiosyncratic methodological choices, leading to a diversity of mechanistic possibilities that may not generalize and may not be germane to human disease. Yet, the interrogation of these aspects is critical to achieve disease understanding and future therapeutic options.

In this study we took a computational rather than experimental approach to address these issues directly in human AD. Our objectives were twofold: <u>First</u>, to test whether a mathematical encoding of network spread due to trans-neuronal proteopathic transmission [19,23,39], combined with local interaction between amyloid and tau pathologies, is capable of recapitulating observed pathology progression in AD; <u>Second</u>, to infer quantitatively the factors of cross-species interactions, their causal direction, and their associated kinetic rate parameters in human AD brain.

Our major findings are: <u>First</u>, the local production of Aβ driven by glucose metabolism followed by subsequent network spread correctly recapitulates the spatial distribution of empirical Aβ, especially in the frontal lobe, the region of highest glucose metabolism. <u>Second</u>, starting from EC, model tau recapitulates empirical tau in temporal areas, followed by network ramification in wider cortices. <u>Third</u>, introduction of Aβ→tau interaction in the model recapitulated the concept of amyloid-facilitated tauopathy, and was critical for success in recapitulating empirical data. <u>Fourth</u>, 2-way interaction is equal or worse than 1-way mode in recapitulating empirical patterns, and the reverse interaction (tau→Aβ) gives the poorest model evidence. <u>Fifth</u>, connectome-mediated spread outperforms other modes of spread, whether by proximity or by fiber length. <u>Sixth</u>, we verified these group-average results on individual subjects, both at baseline and longitudinally, and found essentially equivalent results. These results support

an amyloid-facilitation rather than the classic amyloid-cascade hypothesis, such that both amyloid and tau are capable of independently ramifying within the network, but with an interaction between the two. As discussed below, these results help explain certain known but poorly understood aspects of AD.

Network transmission drives divergent spatiotemporal progression of Aβ and tau

We find that Aβ production driven by glucose metabolism and local APP pool, followed by a certain amount of network dissemination, recapitulates regional pattern of empirical amyloid deposition (see early peak in the R-t curves of **Figure 4**). Neural activity is known to regulate the production and secretion of Aβ[3]. In young transgenic mice, neural activity is related to Aβ release and later deposition of plaques[25]. Neural activity modulates the release of cleavage products of the amyloid precursor protein [40], and Aβ secretion could be affected by neural activity [41] through synaptic exocytosis[26]. In humans, Aβ release parallels fluctuations in synaptic activity in human sleep/wake cycles [42]. AV45 uptake is higher in hubs, multimodal cortices[43] and default mode network, all characterized by higher baseline metabolism[27]. These data provide strong molecular and imaging basis of the amyloid production model (Eq 1a).

For the simulations that were explicitly seeded at EC, their ability to correctly identify mesial temporal involvement is not surprising. However, model pathology increased (**Figure 3**) and fit improved with model time (see R-t curves), suggesting that network spread into other areas was essential to improve the model's performance. **Repeated seeding** shown in **Figure 5A** confirms that EC and adjacent temporal areas are excellent tau seeding locations. Given the preponderance of prior evidence, this serves to further enhance the plausibility of the proposed model.

Once Aβ and tau pathologies are anchored at their respective early sites, subsequent network transmission via the non-linear ODE model of Eqns (1,2) correctly recapitulated the spatiotemporal progression of both Aβ and tau. Thal's five pathologic stages [2] and amyloid PET patterns [3,35] were correctly recapitulated by the model's Aβ evolution, proceeding from medial frontal and precuneus, areas with high baseline metabolism, into the wider network, only slowly entering temporal cortices. These results correlated significantly with amyloid PET scans of disease cohorts. In contrast to Aβ, model tau proceeded outward from EC, to lateroinferior temporal cortices, thence to parietal, lateral occipital and medial frontal areas, in close concordance with Braak's six tau stages [36]. The R-t curve of the tau model against AV1451-PET data (**Figure 4**) yielded highly significant $R = 0.75$ (LMCI), $R = 0.73$ (AD). Against group amyloid, the model yielded very high significance: $R = 0.71$ (EMCI), $R = 0.70$ (AD) Further, the model correlated with MRI-derived regional atrophy of EMCI, LMCI and AD cohorts with high significance. MRI atrophy is an excellent

surrogate for underlying tau, as known from post-mortem studies [44,45]. The presented correlations are significantly stronger than 500 random permutations (**Figure S5**), suggesting that the model is specific to the human connectome and brain topography.

That the spatiotemporal patterning of $A\beta$, tau and atrophy are distinct is well-known but not fully understood[4] and appears at odds with the prevailing amyloid cascade hypothesis [3,46]. Soluble Aβ is sometimes invoked to help explain the discrepancy, but this has been questioned[3]. The divergent evolution of theoretical amyloid and tau on the same network suggests that network spread combined with metabolic drivers might be the governing mechanism behind these long-observed discrepancies. Importantly, this spatial divergence holds even in the presence of amyloid-facilitation.

Aβ-facilitated tau ramification

Large-scale connectivity approaches have yet to address the role of protein-specific mechanisms in disease evolution [47]. Generic networked spread, e.g. our previous study [23], cannot capture the divergence between amyloid and tau, which here was made possible by the introduction of new interaction terms. Our third major result is that Aβ-tau interaction played an important role in forcing the extra-temporal spread of tau. The presence of Aβ is known to influence the course and severity of tau and atrophy [16–18]. Ittner and Götz [17] suggested three modes of interaction: (1) Aβ drives tau pathology; (2) synergistic toxic effects of Aβ and tau; and (3) tau may mediate Aβ toxicity. Aβ and pathological tau co-localize in synapses [48,49], and tau is essential for Aβ-induced neurotoxicity [50]. Aβ seeded exogenously in tau transgenic mice elicited aggressive tau pathology in retrogradely connected regions [18]. Amyloid pathology accelerated tau deposition in double transgenic mice, but the reverse effect was not observed [16]. A "seminal cell biological event" in AD pathogenesis was suggested, whereby acute, tau-dependent loss of microtubule integrity is caused by exposure of neurons to readily diffusible Aβ [32]. However some studies have not found an explicit amyloid-mediated increase in tau in humans. It was suggested that tau propagation across connected regions is unlikely to be a gain of function mediated by the presence of amyloid-β [51]. Nonetheless, there are other mechanistic routes to enact this interaction, e.g. Aβ-related microglial activation promotes local tau-hyperphosphorylation [52], increasing the likelihood of tau spread across connected neurons [53]. Microglia PET in AD confirmed early microglial activation in inferior temporal sites of tau accumulation [54]. Hence it is possible that instead of a direct interaction between amyloid

and tau, there may be an indirect mediation via microglial activation, tau hyperphosphorylation, increased tau load and subsequent enhanced network spread of tau. An excellent review of tau-amyloid synergy is available [55].

In our mathematical exposition of these aspects, modeled tau seeded at EC remained confined to the temporal lobe until sufficient levels of Aβ had spread to and accumulated in temporal cortices (**Figure 3**). Thereupon amyloid-facilitated model tau started diverging from pure tau, taking on a more aggressive trajectory, spreading first to nearby limbic, then basal forebrain, then other neocortical areas. Thus, the facilitation of tau by amyloid does not lead to colocalization of the two until late stages, further helping to explain why regional Aβ patterns do not coincide with tau and atrophy patterns [56,57]. These results concord with neuroimaging, e.g. Sepulcre et al found several convergence zones where amyloid and tau might undergo potential interactions, especially in inferior–lateral temporal areas and entorhinal cortex [58], which they link to the high density of tau in dystrophic neurites in the inferior–lateral and posterior temporal areas [59]. Using a local-to-distributed approach they find specifically that tau accumulation in these temporal areas relates to "massive Aβ elsewhere in the brain" (see, e.g. Fig 6 therein), suggesting these areas as critical regions for linking both pathologies at the large-scale level. Franzmeier 2019 show that tau uptake in amyloid-negative healthy elderly is restricted to (inferior lateral) temporal cortex, but is observed in extra-temporal areas in amyloid-positive controls, MCI and AD, in increasing order[60] (Figure 2 therein). Remarkably, our theoretical modeling also gives the same regions (inferior temporal and EC – see **Figure 3**) as key areas of convergence between amyloid and tau, where the "arrival" of the former is accompanied by prominent deposition of the latter.

Conversely, the absence of the interaction term led modeled tau to remain confined to the temporal lobe (**Figures 3**), mirroring primary age-related tauopathy (PART), a new classification for mild neurofibrillary degeneration in the medial temporal lobe, but no Aβ plaques [34]. Thus, medial temporal NFTs may be involved in two divergent processes: AD and PART. Purely tau-specific abnormalities, e.g. the MAPT gene H1 haplotype, would predispose the subject to PART, whereas additional Aβ abnormalities, e.g. the dysregulation of presenilin, APP or APOE ε4 allele, would cause AD predisposition [34]. Our simulations provide remarkable support to this concept. The non-facilitated tau model was less successful than amyloid-facilitated tau model (**Table 2**, **Figure 3**). Overall, the pure-tau model evolved far slower (in model time) to reach peak correlation with empirical data compared to facilitated-tau model. This result might impart one of the first algorithmic evidence of the coupling between the two protein species as a relevant factor in AD spatiotemporal progression.

Two-way interaction model. The interaction between tau and amyloid can take many forms; a benefit of mathematical modeling is that such interactions can be "observed" *in silico*. We evaluated all four modes of interaction between amyloid and tau. Our 4th major result is that the 2-way model is insignificantly different from the 1-way model for both amyloid and tau, while they are identical for amyloid (**Table 2**). The tau→amyloid interaction performed relatively worse on tau data, suggesting that tau driving amyloid is not a clinically relevant factor in pathology ramification, and its addition gives poorer AIC scores. That amyloid results are identical for both modes is somewhat surprising. We think this was due to amyloid evolution peaking before a significant tau-influence was observed. Hence, by the time tau can begin to influence amyloid, the pear R against empirical amyloid distribution has already been reached.

Non-connectome-mediated spread. Can modes of spread other than trans-neuronal anatomic spread also explain empirical data equally well? Our 5th major result is that alternative proximity- and fiber distance-based spread models did not give good correspondence with empirical tau, and were essentially identical for amyloid (**Table 3**). The amyloid result mirrors our previous exploration in mouse data[37], where too connectivity was not better than spatial spread. The tau result potentially lends a new mechanistic insight about the protein-protein interaction in AD.

A parsimonious explanation of spatiotemporal progression of coupled pathologies in AD

The overall picture that emerges was distilled in **Figure 3C**. Both amyloid and tau are subject to identical networked spread processes. Amyloid production is driven by APP and metabolism, while tau is seeded in mesial temporal cortex, e.g. the EC; but otherwise both pathologies can be diffusely propagated, with no specific role for selective vulnerability due to cell type or cytoarchitecture. From the first temporal tau aggregates, network transmission continues and when sufficient amyloid has arrived at temporal cortices, tau pathology is aggravated. Finally, classic, spatially divergent amyloid and tau patterns are established. This model can explain many empirical discrepancies noted in the literature, yet has the benefit of parsimony, universality and consistency.

Broader Implications. Scientifically, this study bolsters the hypothesis of trans-neuronal transmission by demonstrating its role in humans, an aspect that cannot be studied directly. Since trans-neuronal spread is a common feature of neurodegeneration, the proposed model may be equally applicable to co-morbid pathologies seen in other disorders like Parkinson's, Lewy Body, frontotemporal and other dementias. Clinically, our model could provide a unique opportunity for computational tracking and prediction of individual patients, especially regarding their future

progression. Although multiple AD imaging biomarkers are now routinely acquired, including MRI, FDG-, AV45- and AV1451-PET, the relationships between them are poorly understood, hampering clinical explorations. Since the model can successfully capture the relationship between various evolving biomarkers, it will enable multi-modality integration and tracking of patients' imaging data; e.g. **Figure 5**, where individual subjects at baseline and longitudinally were successfully fitted to the joint amyloid-tau model. We showed that subject-specific seeding sites gave much higher accuracy than uniform common seeding of EC – revealing important inter-subject heterogeneity of etiology and ramification.

However, presented model will need to be strengthened with machine learning and further model inference. Several effective data-driven multi-modality ML methods are available, e.g. the multifactorial model [15] or event-based models [28,29]. In contrast the current study focuses on modeling a specific process of aggregation and evolution of co-morbid pathologies. There is a role for both approaches. Successful tracking and prognostication can improve the odds of early detection, inform clinical care, provide patients visibility into the future, and serve as outcome measures in clinical trials of disease-modifying drugs.

Limitations and future work. Neuroimaging software pipelines have several limitations in image resolution, noise and artifacts [23]. DTI suffers from susceptibility artifacts and poor resolution. PET has poor resolution compared to MRI, and AV45 and AV1451 tracers show significant non-specific binding. Tractography can under-estimate crossing fibers and long tracts. Small subcortical structures can present challenges in inferring connectivity. This study was not designed to achieve staging by fitting a quantitative time-axis in the model – which would ideally utilize longitudinal data and a measure of pathology duration.

ACKNOWLEDGEMENTS

Authors wish to acknowledge assistance in gathering ADNI data by Dr Duygu Tosun, Daren Ma and Areez Malik.

FUNDING

This research was supported by the following grants from the National Institutes of Health: R01NS092802, R01EB022717, RF1AG062196, R56AG064873

COMPETING INTERESTS

None of the authors have a competing financial interest in relation to the work described herein.

AUTHOR CONTRIBUTIONS

AR conceptualized this study and developed the mathematical model, performed all correlations and statistical tests, and wrote the manuscript. JT assisted with data collection, coding and paper writing.

SUPPLEMENTAL INFORMATION

This section includes:

Supplemental Figures (Figures S1 to S9)
Supplemental Table S1, S2
Supplemental Experimental Procedures (Note 1 – Note 9)

Note 1. **Image processing and analysis of ADNI data**

Image acquisition

High resolution T1-weighted MR images were available from the ADNI website from multi-site 3.0 T GE scanners using scanner-specific T1-weighted sagittal 3D MPRAGE sequences (8-channel coil, TR = 400 ms, TE = min full, flip-angle = 11°, slice thickness = 1.2 mm, resolution = 256 × 256 mm and FOV = 26 cm. AV45- and FDG-PET data are also available for download from on multiple instruments of varying resolution and following different platform-specific acquisition protocols. PET data in ADNI underwent standardized image pre-processing correction steps to increase data uniformity across the multicenter acquisitions. More detailed information on the different imaging protocols employed across ADNI sites and standardized image pre-processing steps for MRI and PET acquisitions can be found on the ADNI web site (http://adni.loni.usc.edu/methods/). Demographic information is summarized in **Table S1**.

Image processing

The two datasets were processed independently with different processing streams, the former at Weill Cornell using image processing pipelines established at the Raj laboratory on the AAL atlas parcellation with 90 cerebral regions, and the latter using pipelines developed at the UCSF site on the 86-region Desikan atlas parcellation. Although both pipelines are derived from widely used tools (SPM, FSL, Freesurfer, Camino, etc) and have many common elements, their independent processing enables testing whether the proposed models are specific to our pipelines and atlas, or whether they are more generally applicable. Detailed processing steps are available elsewhere [9,65]. Briefly, T1 images were normalized into MNI space and segmented using SPM8's unified coregistration and segmentation scheme. GM was further parcellated into the 116 region of interest (ROI) AAL atlas [66] using the IBASPM package [67] within. The post-processed AV45-PET and AV1451-PET images were downloaded from ADNI; these images were already normalized to the same common space, scaled by cerebellar values and resliced to uniform voxel resolution. PET images from all normal controls were used to create an average image, which was then normalized to MNI space using SPM8's linear and non-linear transformation. This transformation was then applied to all individual's PET images, which were then resliced to the same resolution as the 90-region atlas at the Cornell site and to the 86-region atlas at the UCSF site. Mean AV1451- and AV45-PET signals were then calculated over each of the GM ROIs.

The raw imaging data were then converted into regional group statistics, following established approaches in neuroimaging. Groupwise, unpaired t-tests were computed between the MCI/AD groups and the normal controls for the AV45-PET and AV1451-PET signal averages in each GM region. T-statistics were calculated to maintain the convention that higher positive value signifies more pathology. Next, since subsequent analysis warrants positive and range-bound measures of regional effects, we scaled the t-statistic to have a mean of 1, and replaced all negative entries (that would represent artifactual hypertrophy) by 0.

Connectomes. Connectomes corresponding to the 90-region AAL parcellation were extracted from 69 healthy subjects, whose data was acquired under a previous study at Cornell, using a processing pipeline that is well established in the Raj laboratory; details can be found in [65]. In short, T1 images were segmented and atlased using SPM8/IBASPM, orientation functions constructed using spherical deconvolution and probabilistic fiber tracking performed in subject's DTI space at 2mm voxel resolution. The average connectivity across 69 subjects was computed for each region pair, to obtain a canonical healthy connectome C. A different canonical healthy connectome corresponding to the 86-region Desikan atlas was generated from the same 69-subject Cornell data, using the same Cornell pipeline.

Table S1. Sample size of the ADNI-3 diagnostic cohorts used in this study. AD: Alzheimer's Disease patients, EMCI: early mild cognitive impairment, LMCI: late MCI, HC: healthy controls. These subjects represent the most complete set of subjects included in the ADNI-3 study as of 1/1/2020 who had all 3 imaging data (AV45-PET, AV1451-PET and MRI). The sample size listed here are for those subjects who had more than one tau PET scan.

	Diagnosis	N	Age range
ADNI 3	AD	68	73.8 (8.12)
	EMCI	137	72.1 (5.91)
	LMCI	75	71.9 (7.61)
	HC	251	74.10 (6.69)

Note 2. Spatial Distribution of APP and MAPT

The rate of Aβ accumulation is modeled in our approach as the product of two starting distributions: a) the distribution of APP, the precursor protein from which Aβ is cleaved; and b) the distribution of regional metabolism, as measured from FDG-PET scans of healthy age-matched individuals in the ADNI cohort. APP regional patterns are not available from in vivo scans, hence a rough spatial pattern vector of APP was computed from publicly available human gene expression data and from post-mortem histology data. We relied on two studies to obtain the pattern of APP in the human brain. First, we used the Allen Brain Institute's human brain gene expression data, from which the APP expression values were resolved on all available sample locations in 6 brains. This data, shown in **Figure S1** (left), strongly suggests that APP distribution in the brain is highly heterogeneous, with above-average expression in brainstem, subcortical and frontal cortices, but below average expression in posterior and cerebellar structures. This closely matches a previous histology study of post-mortem brains, which also found that APP, along with all the other Aβ-producing factors (PS1, BACE1, APP-CTFβ) are usually under-expressed in the posterior and parietal neocortex, as well as in cerebellum [63]. For example, the expression of APOE, which plays a strong role in the processing of APP, is similarly distributed, with below average levels in posterior cortices and above-average levels in striatal and subcortical structures (**Figure S1**, right). The study by Shinohara et al [63] suggested that the numerical values of APP distribution in posterior cortices is approximately 50% compared to other cortices, especially in orbitofrontal cortex. Since APP levels from either our in-house or the study by Shinohara et al were not quantitatively available from the individual brain regions used in this study, we simply use a 2-level APP distribution, where all brain regions are assigned the APP level of 1, except the occipital regions, whose APP level is assigned 0.5. Note that the cerebellum is not part of this study, hence its APP level was not used. Some striatal areas, in particular putamen and caudate nucleus, show higher APP levels than the rest of the brain, as shown in [63]. Therefore these two structures were assigned the relative score of 1.1.

Tau accumulation is not assigned a specific regional driver, however, since misfolded tau derived from physiologic tau, its concentration in turn is expected to be governed by the expression of microtubule associated protein tau (MAPT). MAPT expression is now quantitatively known in various parts of the brain, and in general is much lower in occipital and striatal regions [64]. Hence we assign a rough map of MAPT with the value of 1 everywhere except occipital, putamen and caudate, where it is assigned 0.5.

Healthy baseline metabolic load   Healthy APP gene expression   Healthy MAPT gene expression

Figure S1. Spatial distribution in the human brain of factors driving the production of amyloid in the proposed model of Eq 1. These factors are: baseline glucose metabolic load and APP gene expression, and together they govern the spatial distribution of amyloid production, as per Eq 1. Recall that tau is deemed produced only at the seeding site, which is ERC throughout the manuscript unless otherwise noted. To provide some context to the tau production aspect, on the right panel we show the spatial distribution of the MAPT gene. MAPT is regarded as the pool from which misfolded tau is cleaved. Cleary, MAPT does not show appreciable spatial concentration in the human brain, occurring diffusely throughout, with the exception of visual cortices. The metabolic load is the t-score of healthy FGD-PET SUVr in the cognitively normal subjects in ADNI. Spheres are located at the site of the gene expression sample, collected over 6 healthy young brain hemispheres by the Allen Institute. Sphere diameter is proportional to the z-score with respect to overall mean and standard deviation (+/- 2 std are shown here) and color-coded by lobe as in the main manuscript. Amongst cerebral regions, posterior and occipital regions have the least overall levels of APP and MAPT, while brainstem and subcortical structures have the highest levels. Frontal regions are generally above average in both.

Note 3. Driving function to model production of pathology at local sites

The coupled differential equation given by Eq (4a, 4b) has a production term for both tau and $A\beta$. This is modeled in this study as smooth but time-limited driving functions, in recognition of the fact that abrupt, discontinuous production functions are biophysically implausible, and that the production terms cannot continue *ad infinitum*, and must eventually decline in response to amyloid saturation, cell death and other factors. The function chosen for this purpose is one shaped like the Gamma curve given by $f_\tau(t) = \frac{1}{T_\tau} e^{-t/T_\tau}$, shown in **Figure S2**. The function depends on a single parameter T_τ that controls the duration of production of tau; analogously for amyloid. Trial and error resulted in the best choice of this parameter as $T_\tau = T_{A\beta} = 5\ years$. This function reproduces the expectation that at the beginning the rate of production is low, and steadily increases with increasing pathology. The rate of production then plateaus in response to countervailing forces in the brain. Finally, as neurodegeneration takes hold and amyloid and tau begin to be increasingly sequestered into plaques and tangles, the rate of production of soluble pathology begins to decline. It is stressed that the shape used here, Gamma function, is only one of many possibilities. Its choice was motivated by its smoothness, time boundedness, unimodal peak, and its ubiquity in mathematical modeling. This is clearly a design choice but not an essential one; any reasonable bounded driving function of sufficiently long duration will suffice.

Figure S2. **Gamma-shaped production function for both tau and amyloid production**. Related to Experimental Procedures, Supplemental Experimental Procedures: **Note 3**. The y-axis represents the rate of production of each entity as a function of time, with t=0 at disease onset. This function gives low production in the beginning, hitting a peak in the middle of disease course, and thence declining.

Note 4. Global pathology burden predicted by the model

Figure S3 shows the global accumulation of theoretical pathology over model time, evaluated at default parameters. All proteins increase over time, but amyloid-facilitated tau diverges dramatically from the non-facilitated tau at around t=15, mirroring main **Figure 3**. Clearly, while the "pure" tau model also captures empirical data, it does so far slower and achieves far less correlation strength than the facilitated version.

Figure S3. **Global burden of modeled pathology and the effect of amyloid facilitation. Top:** The evolution of global pathology burden as evaluated by numerical integration of Eqs (1) under default model parameters. **Bottom**: R-t curves showing the correlation between the non-facilitated tau model and empirical tau and atrophy data. Note the significantly longer time scale compared to **Figure 4**, and the lower peak R.

Note 5. Influence of Atlas

The two atlases used in this study, Desikan and AAL, give comparable results when the model is applied to each. **Figure S4** (left) shows the second eigenmode of the Laplacian from each connectome, with highly similar results. In both cases the highest values are in temporal cortices, which in this study was used to explain the early formation of tau fibrils in these regions. Similarly, the two atlases gave comparable distribution of empirical atrophy measured from MRI of the ADNI cohort, as shown in **Figure S4** (right). Note that neither the datasets, nor the atlases, nor the image processing pipeline used in constructing the two regional distributions are identical; this was described in **Supplemental Experimental Procedures: Note 1**. Thus, we conclude that the choice of connectome, atlas

parcellation, processing pipeline or the specific stratification of the ADNI cohort are responsible for achieving the results shown in the main text.

A

B

Figure S4. The evolution of model dynamics of amyloid and tau, evaluated by numerical integration of Eqs (1a,b), on the 90x90 healthy AAL connectome, under default model parameters. The first column shows the evolution of amyloid, the second column of amyloid-facilitated tau and the third column of tau only. Note the AAL atlas does not separate out entorhinal cortex, which is included within the parahippocampal gyrus. Spheres are color-coded by lobe, but color assignment is different from **Figure 3**.

Note 6. Parameter fitting for individual cases

The full models in Eqns (1,2) have 6 unknown global parameters: $\beta, \alpha, \gamma, \eta, \ T_{A\beta}, T_\tau, K$. Fortunately, the model is not overly sensitive to parameter choice, since all these parameters are related in one way or another to the time-scale of the neurodegenerative process, thereby primarily governing the scale of the time axis, not the overall behavior. We set the time-specific parameters $T_{A\beta}, T_\tau$ at the outset heuristically over 20 trials to yield a temporal response that roughly matched the duration of Alzheimer pathology dissemination, i.e. 30 years [62]. This yielded $T_{A\beta} = T_\tau = 10 \ years$. The parameter K, which represents the capacity of overall pool of tau and amyloid, and thereby governs the logistic growth terms in the equations, was set at 5 for all subjects and simulations. The exact choice appears to not be important; similar model behavior was observed for other choices, like 10 or 20.

Other model parameters were optimized in two steps. First, starting with initial guesses of all parameters at 1.00, we employed brute-force grid search, over a range with 10 equally-spaced points per parameter. This yielded the parameter values shown in **Table 3**. We call this coarsely-optimized model specification the "default parameters", which was used for assessing the model qualitatively and to understand its behavior over different seeding sites. Please note that the temporal units used here are shown as "years" for illustration only; since the model time as arbitrary units, so do the above rate constants.

Next, using the default parameters as starting guesses, we implemented a maximum a posteriori (MAP) estimator for the model, which jointly maximized a likelihood function as well as priors on each parameter. For this purpose we chose relatively shallow mutually-independent priors given by Gaussians around the initial guess of each parameter. The hyperparameter that weighs the prior term was obtained via trail and error, and set at 0.01; the method is not sensitive to the hyperparameter choice, since it is usually very small in order to not overly rely on the priors. We selected a likelihood function that is given by the strength of correlation to empirical data rather than more typical residual norm:

$$\Pr\big(x_\tau(t), x_{A\beta}(t)|\theta\big) \propto \exp\big(-R(x_\tau(t,\theta), x_\tau^{ADNI})\big) \cdot \exp\big(-R\big(x_{A\beta}(t,\theta), x_{A\beta}^{ADNI}\big)\big)$$

$$t_{max}|\theta = \arg\max_t \prod_i \Pr\big(x_\tau(t+\Delta t_i), x_{A\beta}(t+\Delta t_i)|\theta\big)$$

where $R(\cdot,\cdot)$ is the correlation operator, θ is the collection of model parameters, and the model's evolution is governed by these parameters via the ODE. Note that the fits employ all available longitudinal visits t_i, where by convention we define time gaps by $\Delta t_i = t_i - t_1$. The resulting MAP problem then becomes a non-linear cost minimization problem, which was solved using MATLAB's fmincon() subroutine. Since fmincon() calls the ODE solver during each iteration, the process can become time consuming. Fortunately we found that using K=50 iterations was quite adequate; there was no discernible difference by using much higher number of iterations. Overall execution time for a single subject fitting ranged from 5 to 20 seconds.

Table S2: **Model parameters and defaults used in optimization.** Parameters set after coarse grid search ("coarsely optimized" or default parameters) and after further refinement using the Maximum a Posteriori estimation procedure ("Optimal values"), using the default parameters as initial guess. All parameters have units of $year^{-1}$, since they are rate parameters.

Empirical data used for fitting	Coarsely optimized (default) parameters					Fitted parameters				
	β,	β_a,	α,	α_a,	γ	β,	β_a,	α,	α_a,	γ
EMCI	0.4,	1.00,	2.00,	2.00,	0.0	0.22,	0.86,	1.00,	1.33,	1.33
LMCI	1.00,	1.00,	2.00,	2.00,	0.20	0.28,	1.38,	0.99,	1.43,	1.44
AD	1.00,	1.00,	2.00,	2.00,	0.20	0.28,	0.97,	1.02,	1.22,	1.20

The other spread models (other than connectome-based spread) were optimized in the same fashion. Their optimal values did not change much or in interesting ways, likely due to the normalization of the adjacency matrices implicit in the definition of the Laplacian.

Model fitting to individual subjects. We applied the same MAP estimator to individual subjects, whether they had a single baseline scan or multiple longitudinal scans. In the latter case, the likelihood function above was evaluated jointly at all available longitudinal samples, keeping the time gap between them intact. Both amyloid and tau evolution occurs jointly in time in our model, hence in main **Figure 4** we report the correlation achieved by both tau and amyloid at the same time instant t_{max}, when the posterior probability peaks. It is possible to impose a delay between tau and amyloid onsets within our model, but we opted not to do so due to a lack of experimental data on that aspect.

Note 7. Permutation testing to demonstrate disease specificity

To determine whether the presented model displays AD-specificity and connectome-specificity, two permutation tests were devised:

A) 500 random permutations of the group AD atrophy pattern were obtained and repeatedly tested against the network model (**Figure S5-A)**. If the model is not specific to AD, a significant proportion of these tests should be

positive. In fact, randomly permuted R cluster near zero; hence reported R values are highly significant compared to this "null" distribution ($p < 10^{-3}$ for all groups). To simplify presentation and avoid excessive computational burden, in these simulations only linear and cross-product terms were retained in the models; this choice does not alter the overall system behavior while making computations dramatically faster.

B) The above null models tend to destroy the correlation structure in regional tau and atrophy. To avoid this and to test whether the true anatomic network structure is necessary to replicate our results, we performed 500 random permutations of the connectome itself, and correlated the (refitted) model seeded at EC against empirical (unchanging) regional pattern of tau-PET. Each connectome realization randomly scrambled the upper triangular portion of the true human connectome, and generated a symmetric matrix from that. Thus, these connectomes preserve actual connectivity values, but randomize network topology. If the model is not specific to the connectome, a significant proportion of these tests should be positive. The "null" R distributions shown in **Figure S5-B** are significantly smaller compared to the true reported R ($p < 10^{-3}$ for all groups). The results for atrophy and amyloid are largely the same and not reported here. Note there is a hard lower limit in each case, which reflect the correlation between EC seed and regional tau prior to any network spread, regardless of the connectome used.

Taken together, these results confirm that the proposed model only recapitulates empirical regional distributions when it is applied in the correct region order and to the correct human connectome.

Figure S5. **Permutation testing**. Pearson's R of correlation strength between proposed model in Eq (1a,b) solved via numerical integration, and empirical regional patterns of each diagnosis group (EMCI, LMCI, AD) in the ADNI cohort. **A**: Modeled vs empirical tau (top row), atrophy (middle) and Aβ data (bottom) are shown alongside the R-t curves corresponding to 500 random permutations of the regional patterns of the AD cohort. The random

permutations lead to peak correlations with the model that are centered near 0, suggesting that the actual topography of the regional values is critical to the results we have reported. Histograms of peak R from random permutations are shown in right column, alongside R reported for empirical data, indicating that the latter are highly significant under permutation testing. **B**: Histograms of peak R against reginal tau from random permutations of the connectome, alongside R reported for true connectome, indicating that the latter are highly significant under permutation testing.

Note 8. Spearman correlations, compared to Pearson in Figure S5

The previous figure reported the agreement between the model and empirical regional distribution of biomarkers using Pearson correlation. It is now demonstrated that the results are similar when using a non-parametric correlation statistic, i.e. Spearman rank correlation. Spearman does not require that the data being compared are drawn from Gaussian distributions, hence it may be appropriate in the current context. **Figure S6** shows that the results using Spearman are generally similar to the main Pearson results in the main text but with lower significance. Pearson was preferred in this study because it is not as non-linear as Spearman, which is overly sensitive to low effect sizes, which generally pertain to brain regions that do not participate in AD pathology. These noisy values are largely irrelevant in the current context, yet Spearman rank is overly sensitive to these values. However, the fact that both Spearman and Pearson are in rough agreement gives additional confidence that the data reported in the main study are robust.

Figure S6. **Spearman correlation results of default model against empirical regional statistics**. Compared to Pearson correlation curves in main text, these results are very similar, if noisier due to non-linearities inherent in Spearman rank calculations. Both atrophy and amyloid patterns of the theoretical model are specific to AD phenotypes, and do not correlate well with either random permutations or MCI data for amyloid-negative subjects (dotted blue curves).

Note 9: Repeated seeding experiments to assess alternative seeding sites

Other than the canonical Entorhinal cortex (EC) seeding site, we also assessed other regions' seeding plausibility. For this, we repeatedly simulated the joint model seeded from each region bilaterally. For each seed region, peak Pearson's R between model and ADNI tau PET data were shown in **Figure 5**. EC is amongst the best overall cortical seeding site, while hippocampus (HP) is the best subcortical site. Other prominent seeding sites include Inferotemporal cortex (IT), Parahippocampal gyrus (PHP) and Fusiform gyrus (Fus). The evolution of pathology from these non-EC sites are shown below in **Figures S8** and **S9**, for HP-seeding and IT-seeding, respectively. They remain substantially similar to main **Figure 4**; however a notable difference from EC seeding is that HP seeding leads to higher involvement of medial temporal and subcortical structures. Amongst the above 5 regions, EC is unique in giving consistently one of the highest seeding likelihood despite having less levels of empirical tau deposition than others.

Figure S8. **Evolution of default model's tau after seeding the Hippocampus, compared against empirical data.** The first column shows the modeled evolution of amyloid-facilitated tau, while the second column shows cross-sectional empirical tau-PET SUVr patterns from all 3 diagnostic groups. The top panel shows the behavior of peak R against model time t. The match between the model and empirical tau are strongest for E/LMCI and weakest for AD. The right-most column shows results for MRI-derived atrophy. The match between the model and empirical data are strongest for AD and weakest for EMCI. Peak R is highly significant in all cases. These results are similar to the base case presented in the main manuscript, **Figure 4**, which had EC seeding of tau. The key difference is that the baseline correlation (left-most end of the R-t curves shown in top row) start off higher than does EC – this is likely due to the stronger baseline levels of both tau and atrophy in hippocampus compared to EC. Another notable difference from EC seeding is that HP seeding lead to higher involvement of medial temporal and subcortical structures (denoted in black).

Figure S9. **Evolution of default model tau after seeding the Inferotemporal Cortex, compared against empirical data.** The first column shows the modeled evolution of amyloid-facilitated tau, while the second column shows cross-sectional empirical tau-PET SUVr patterns from all 3 diagnostic groups. The top panel shows the behavior of peak R against model time t. The match between the model and empirical tau are strongest for LMCI and AD, and weakest for EMCI. The right-most column shows results for MRI-derived atrophy. The match between the model and empirical data are strongest for AD and weakest for E/LMCI. Peak R is highly significant in all cases. These results are similar to the base case presented in the main manuscript, **Figure 4**, which had EC seeding of tau. The key difference is that the baseline correlation of tau (left-most end of the R-t curves shown in top row) start off higher than does EC – this is likely due to the stronger baseline levels of tau (but not atrophy) in IT compared to EC.

Table S3: **Model comparison using Fisher's R-to-z transformed t-test results pertaining to main data in Table 2**. Six theoretical models are compared for statistically significant difference in performance: the No-interaction model, without the interaction term ($\gamma = 0$); the 1-way interaction model, whereby tau affects amyloid but not vice versa; another 1-way model whereby amyloid affects tau but not vice versa; and the 2-way interaction model where both tau and amyloid affect each other via γ. Three different modes of causal interaction from amyloid to tau are given: where amyloid influences the diffusion of tau; where it influences only the aggregation of tau; and where amyloid influences aggregation of tau remotely, i.e. via long-range connections, rather than locally. Fisher's R-to-z transform was computed, and the *1-sided* p-value of significant differences between each model-pair was evaluated; hence only improvements from base model are assessed. P-values that are smaller than the significance threshold ($p < 0.05$) are highlighted in bold red font, and indicate comparisons where one model is significantly different than another. For conciseness, Fisher comparisons are shown only for model-pairs "closest" to each other. Overall, the most significantly improved model from the base model is the 1-way interaction from amyloid to tau aggregation; consequently it is highlighted by green box around it. The remote interaction and the 2-way models are not sufficiently different from the 1-way model.

Model A:	1-way interaction tau→$A\beta$ aggregation	1-way interaction $A\beta$→ tau diffusion	1-way interaction $A\beta$→ tau aggregation	1-way (remote) interaction $C \cdot A\beta$→tau aggregation	2-way interaction $A\beta\leftrightarrow$tau aggregation
Model B:	No-interaction	No-interaction	No-interaction	1-way interaction $A\beta$→ tau aggregation	1-way interaction $A\beta$→ tau aggregation
EMCI Aβ	p = 1	p = 1	p = 1	p = 1	p = 1
LMCI Aβ	p = 1	p = 1	p = 1	p = 1	p = 1
AD Aβ	p = 1	p = 1	p = 1	p = 1	p = 1
EMCI tau	p = 1	p = 1	**p < 0.01**	p = 1	p = 1
LMCI tau	p = 1	p = 1	**p < 0.01**	p = 1	p = 1
AD tau	p = 1	p = 1	**p < 0.01**	p = 1	p = 1
EMCI atr	p = 1	p = 1	p = 1	p = 1	p = 1
LMCI atr	p = 1	p = 1	p = 1	p = 1	p = 1
AD atr	**p < 0.01**	p = 1	**p < 0.01**	p = 1	p = 1

References

1. Braak, H. & Braak, E. Neuropathological stageing of Alzheimer-related changes. *Acta Neuropathol.* **82**, 239–259 (1991).

2. Thal, D. R., Rüb, U., Orantes, M. & Braak, H. Phases of A beta-deposition in the human brain and its relevance for the development of AD. *Neurology* **58**, 1791–800 (2002).

3. Jagust, W. J. & Mormino, E. C. Lifespan brain activity, β-amyloid, and Alzheimer's disease. *Trends Cogn. Sci.* **15**, 520–6 (2011).

4. La Joie, R. *et al.* Region-specific hierarchy between atrophy, hypometabolism, and β-amyloid (Aβ) load in Alzheimer's disease dementia. *J. Neurosci.* **32**, 16265–73 (2012).

5. Rabinovici, G. D. *et al.* Increased metabolic vulnerability in early-onset Alzheimer's disease is not related to amyloid burden. *Brain* **133**, 512–28 (2010).

6. Jack Jr, C. R. *et al.* Hypothetical model of dynamic biomarkers of the Alzheimer's pathological cascade. *Lancet Neurol.* **9**, 119–128 (2010).

7. Hardy, J. & Selkoe, D. J. The amyloid hypothesis of Alzheimer's disease: progress and problems on the road to therapeutics. *Science* **297**, 353–6 (2002).

8.   Raj, A., Kuceyeski, A. & Weiner, M. A Network Diffusion Model of Disease Progression in Dementia. *Neuron* **73**, 1204–1215 (2012).

9.   Raj, A. *et al.* Network Diffusion Model of Progression Predicts Longitudinal Patterns of Atrophy and Metabolism in Alzheimer's Disease. *Cell Rep.* **in print**, 359–369 (2015).

10.   Pievani, M., de Haan, W., Wu, T., Seeley, W. W. & Frisoni, G. B. Functional network disruption in the degenerative dementias. *Lancet. Neurol.* **10**, 829–43 (2011).

11.   Zhou, J., Gennatas, E. D., Kramer, J. H., Miller, B. L. & Seeley, W. W. Predicting regional neurodegeneration from the healthy brain functional connectome. *Neuron* **73**, 1216–27 (2012).

12.   Weickenmeier, J., Jucker, M., Goriely, A. & Kuhl, E. A physics-based model explains the prion-like features of neurodegeneration in Alzheimer's disease, Parkinson's disease, and amyotrophic lateral sclerosis. *J. Mech. Phys. Solids* **124**, 264–281 (2019).

13.   Fornari, S., Schäfer, A., Jucker, M., Goriely, A. & Kuhl, E. Prion-like spreading of Alzheimer's disease within the brain's connectome. *J. R. Soc. Interface* **16**, (2019).

14.   Iturria-Medina, Y., Sotero, R. C., Toussaint, P. J. & Evans, A. C. Epidemic Spreading Model to Characterize Misfolded Proteins Propagation in Aging and Associated Neurodegenerative Disorders. *PLoS Comput. Biol.* **10**, e1003956 (2014).

15.   Iturria-Medina, Y., Carbonell, F. M., Sotero, R. C., Chouinard-Decorte, F. & Evans, A. C. Multifactorial causal model of brain (dis)organization and therapeutic intervention: Application to Alzheimer's disease. *Neuroimage* **152**, 60–77 (2017).

16.   Hurtado, D. E. *et al.* A{beta} accelerates the spatiotemporal progression of tau pathology and augments tau amyloidosis in an Alzheimer mouse model. *Am. J. Pathol.* **177**, 1977–88 (2010).

17.   Ittner, L. M. & Götz, J. Amyloid-β and tau--a toxic pas de deux in Alzheimer's disease. *Nat. Rev. Neurosci.* **12**, 65–72 (2011).

18.   Götz, J., Chen, F., van Dorpe, J. & Nitsch, R. M. Formation of neurofibrillary tangles in P301l tau transgenic mice induced by Abeta 42 fibrils. *Science* **293**, 1491–5 (2001).

19.   Jucker, M. & Walker, L. C. Self-propagation of pathogenic protein aggregates in neurodegenerative diseases. *Nature* **501**, 45–51 (2013).

20.   Frost, B. & Diamond, M. I. Prion-like mechanisms in neurodegenerative diseases. *Nat. Rev. Neurosci.* **11**, 155–

9 (2010).

21. Clavaguera, F. *et al.* Transmission and spreading of tauopathy in transgenic mouse brain. *Nat. Cell Biol.* **11**, 909–13 (2009).

22. Iba, M. *et al.* Tau pathology spread in PS19 tau transgenic mice following locus coeruleus (LC) injections of synthetic tau fibrils is determined by the LC's afferent and efferent connections. *Acta Neuropathol.* **130**, 349–62 (2015).

23. Raj, A., Kuceyeski, A. & Weiner, M. A network diffusion model of disease progression in dementia. *Neuron* **73**, 1204–15 (2012).

24. Zhou, J., Gennatas, E. D., Kramer, J. H., Miller, B. L. & Seeley, W. W. Predicting regional neurodegeneration from the healthy brain functional connectome. *Neuron* **73**, 1216–27 (2012).

25. Bero, A. W. *et al.* Neuronal activity regulates the regional vulnerability to amyloid-β deposition. *Nat. Neurosci.* **14**, 750–756 (2011).

26. Cirrito, J. R. *et al.* Synaptic activity regulates interstitial fluid amyloid-beta levels in vivo. *Neuron* **48**, 913–22 (2005).

27. Buckner, R. L. *et al.* Molecular, structural, and functional characterization of Alzheimer's disease: evidence for a relationship between default activity, amyloid, and memory. *J. Neurosci.* **25**, 7709–17 (2005).

28. Young, A. L. *et al.* A data-driven model of biomarker changes in sporadic Alzheimer's disease. *Brain* **137**, 2564–2577 (2014).

29. Oxtoby, N. P. *et al.* Data-Driven Sequence of Changes to Anatomical Brain Connectivity in Sporadic Alzheimer's Disease. *Front. Neurol.* **8**, 580 (2017).

30. Walker, L. C., Lynn, D. G. & Chernoff, Y. O. A standard model of Alzheimer's disease? *Prion* vol. 12 261–265 (2018).

31. Kara, E., Marks, J. D. & Aguzzi, A. Toxic Protein Spread in Neurodegeneration: Reality versus Fantasy. *Trends in Molecular Medicine* vol. 24 1007–1020 (2018).

32. King, M. E. *et al.* Tau-dependent microtubule disassembly initiated by prefibrillar β-amyloid. *J. Cell Biol.* (2006) doi:10.1083/jcb.200605187.

33. Weiner, M. W. *et al.* The Alzheimer's Disease Neuroimaging Initiative: a review of papers published since its inception. *Alzheimers. Dement.* **8**, S1-68 (2012).

34. Crary, J. F. *et al.* Primary age-related tauopathy (PART): a common pathology associated with human aging. *Acta Neuropathol.* **128**, 755–66 (2014).

35. Landau, S. M. *et al.* Amyloid deposition, hypometabolism, and longitudinal cognitive decline. *Ann. Neurol.* **72**, 578–86 (2012).

36. Braak, H. & Braak, E. Evolution of the neuropathology of Alzheimer's disease. *Acta Neurol. Scand. Suppl.* **165**, 3–12 (1996).

37. Mezias, C. & Raj, A. Analysis of Amyloid-β Pathology Spread in Mouse Models Suggests Spread Is Driven by Spatial Proximity, Not Connectivity. *Front. Neurol.* **8**, 653 (2017).

38. Jack, C. R. *et al.* A/T/N: An unbiased descriptive classification scheme for Alzheimer disease biomarkers. *Neurology* (2016) doi:10.1212/WNL.0000000000002923.

39. Seeley, W. W., Crawford, R. K., Zhou, J., Miller, B. L. & Greicius, M. D. Neurodegenerative diseases target large-scale human brain networks. *Neuron* **62**, 42–52 (2009).

40. Nitsch, R. M., Farber, S. A., Growdon, J. H. & Wurtman, R. J. Release of amyloid beta-protein precursor derivatives by electrical depolarization of rat hippocampal slices. *Proc. Natl. Acad. Sci. U. S. A.* **90**, 5191–3 (1993).

41. Kamenetz, F. *et al.* APP processing and synaptic function. *Neuron* **37**, 925–37 (2003).

42. Gilestro, G. F., Tononi, G. & Cirelli, C. Widespread Changes in Synaptic Markers as a Function of Sleep and Wakefulness in Drosophila. *Science (80-.).* **324**, 109–112 (2009).

43. Buckner, R. L. *et al.* Cortical hubs revealed by intrinsic functional connectivity: mapping, assessment of stability, and relation to Alzheimer's disease. *J. Neurosci.* **29**, 1860–73 (2009).

44. Whitwell, J. L. *et al.* MRI correlates of neurofibrillary tangle pathology at autopsy: a voxel-based morphometry study. *Neurology* **71**, 743–9 (2008).

45. Vemuri, P. *et al.* Antemortem MRI based STructural Abnormality iNDex (STAND)-scores correlate with postmortem Braak neurofibrillary tangle stage. *Neuroimage* **42**, 559–67 (2008).

46. Fjell, A. M. & Walhovd, K. B. Neuroimaging results impose new views on Alzheimer's disease--the role of amyloid revised. *Mol. Neurobiol.* **45**, 153–72 (2012).

47. Warren, J. D., Rohrer, J. D. & Hardy, J. Disintegrating brain networks: from syndromes to molecular nexopathies. *Neuron* **73**, 1060–2 (2012).

48. Fein, J. A. *et al.* Co-localization of amyloid beta and tau pathology in Alzheimer's disease synaptosomes. *Am. J. Pathol.* **172**, 1683–92 (2008).

49. Takahashi, R. H., Capetillo-Zarate, E., Lin, M. T., Milner, T. A. & Gouras, G. K. Co-occurrence of Alzheimer's disease ß-amyloid and τ pathologies at synapses. *Neurobiol. Aging* **31**, 1145–52 (2010).

50. Amadoro, G. *et al.* Endogenous Aβ causes cell death via early tau hyperphosphorylation. *Neurobiol. Aging* **32**, 969–90 (2011).

51. Franzmeier, N. *et al.* Functional brain architecture is associated with the rate of tau accumulation in Alzheimer's disease. *Nat. Commun.* **11**, 1–17 (2020).

52. Heurtaux, T. *et al.* Microglial activation depends on beta-amyloid conformation: role of the formylpeptide receptor 2. *J. Neurochem.* **114**, 576–586 (2010).

53. Maphis, N. *et al.* Reactive microglia drive tau pathology and contribute to the spreading of pathological tau in the brain. *Brain* **138**, 1738–1755 (2015).

54. Hamelin, L. *et al.* Early and protective microglial activation in Alzheimer's disease: a prospective study using 18 F-DPA-714 PET imaging. *Brain* **139**, 1252–1264 (2016).

55. Busche, M. A. *et al.* Tau impairs neural circuits, dominating amyloid-β effects, in Alzheimer models in vivo. *Nat. Neurosci.* **22**, 57–64 (2019).

56. Vemuri, P. *et al.* MRI and CSF biomarkers in normal, MCI, and AD subjects: predicting future clinical change. *Neurology* **73**, 294–301 (2009).

57. Jack, C. R. & Holtzman, D. M. Biomarker Modeling of Alzheimer's Disease. *Neuron* **80**, 1347–1358 (2013).

58. Sepulcre, J. *et al.* In vivo tau, amyloid, and gray matter profiles in the aging brain. *J. Neurosci.* **36**, 7364–7374 (2016).

59. Marquié, M. *et al.* Validating novel tau positron emission tomography tracer [F-18]-AV-1451 (T807) on postmortem brain tissue. *Ann. Neurol.* **78**, 787–800 (2015).

60. Franzmeier, N. *et al.* Functional connectivity associated with tau levels in ageing, Alzheimer's, and small vessel disease. *Brain* **142**, 1093–1107 (2019).

61. Behrens, T. E. J., Berg, H. J., Jbabdi, S., Rushworth, M. F. S. & Woolrich, M. W. Probabilistic diffusion tractography with multiple fibre orientations: What can we gain? *Neuroimage* **34**, 144–55 (2007).

62. Villemagne, V. L. *et al.* Amyloid β deposition, neurodegeneration, and cognitive decline in sporadic

Alzheimer's disease: a prospective cohort study. *Lancet Neurol.* **12**, 357–67 (2013).

63.   Shinohara, M. *et al.* Regional distribution of synaptic markers and APP correlate with distinct clinicopathological features in sporadic and familial Alzheimer's disease. *Brain* **137**, 1533–49 (2014).

64.   Trabzuni, D. *et al.* MAPT expression and splicing is differentially regulated by brain region: relation to genotype and implication for tauopathies. *Hum. Mol. Genet.* **21**, 4094–103 (2012).

65.   Kuceyeski, A., Maruta, J., Relkin, N. & Raj, A. The Network Modification (NeMo) Tool: elucidating the effect of white matter integrity changes on cortical and subcortical structural connectivity. *Brain Connect.* **3**, 451–63 (2013).

66.   Tzourio-Mazoyer, N. *et al.* Automated anatomical labeling of activations in SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject brain. *Neuroimage* **15**, 273–89 (2002).

67.   Alemán-Gómez, Y., Melie-García, L. & Valdés-Hernandez, P. IBASPM: Toolbox for automatic parcellation of brain structures. in *Presented at the 12th Annual Meeting of the Organization for Human Brain Mapping* (2005).