1

Prior exposure to pathogens augments host heterogeneity in susceptibility and has key

epidemiological consequences

Dana M. Hawley*,[1], Anna A. Pérez-Umphrey[1], James S. Adelman[2], Arietta E. Fleming-Davies[3],

Jesse Garrett-Larsen[1], Steven J. Geary[4], Lauren M. Childs[$,5], Kate E. Langwig[$,*,1]

[1]Department of Biological Sciences, Virginia Tech, Blacksburg, VA, USA

[2]Department of Biological Sciences, University of Memphis, Memphis, TN, USA

[3]Department of Biology, University of San Diego, San Diego, CA, USA

[4]Department of Pathobiology & Veterinary Science, University of Connecticut, Storrs, CT, USA

[5]Department of Mathematics and Virginia Tech Center for Mathematics of Biosystems, Virginia

Tech, Blacksburg, VA, USA

*Corresponding authors

Email: hawleyd@vt.edu, klangwig@gmail.com

[$]These authors contributed equally to the work

Abstract. Pathogen epidemics are key threats to human and wildlife health. Across systems,

host protection from pathogens following initial exposure is often incomplete, resulting in

recurrent epidemics through partially-immune hosts. Variation in population-level protection has

important consequences for epidemic dynamics, but how acquired protection influences inter-

individual heterogeneity in susceptibility and its epidemiological consequences remains

understudied. We experimentally investigated whether prior exposure (none, low-dose, or high-

dose) to a bacterial pathogen alters host heterogeneity in susceptibility among songbirds. Hosts

with no prior pathogen exposure had little variation in protection, but heterogeneity in

susceptibility was significantly augmented by prior pathogen exposure, with the highest

variability detected in hosts given high-dose prior exposure. An epidemiological model

parameterized with experimental data found that heterogeneity in susceptibility from prior

exposure more than halved epidemic sizes compared with a homogeneous population with

identical mean protection. However, because infection-induced mortality was also greatly

reduced in hosts with prior pathogen exposure, reductions in epidemic size were smaller than

expected in hosts with prior exposure. These results highlight the importance of variable

protection from prior exposure and/or vaccination in driving population-level heterogeneity and

epidemiological dynamics.

Introduction

Pathogen epidemics are increasing in frequency in humans and other animals (1,2), underscoring the need to characterize key sources of heterogeneity that influence epidemiological dynamics for pathogens. Hosts in a population often show considerable heterogeneity in contact rates (e.g. 3,4), infectiousness (e.g. 5), and even pathogen susceptibility (e.g. 6), defined here as the probability of infection given exposure. The individual-level traits that predict host susceptibility to a given pathogen, including sex (7), genotypic variants (8), and immune history (9), have been widely described for some systems. Nonetheless, the coarse discretization of these traits into mathematical models often inaccurately describes the extent of heterogeneous infection risk present among hosts in a population (10–13). A key approach to quantifying such heterogeneous risk at the population level is to estimate the shape of the distribution of host susceptibility (14–16), either using epidemiological data (e.g. 17,18) or experimental studies of non-human animals (e.g. 19,20). Overall, such population-level approaches to quantifying host susceptibility distributions are particularly important because the shape of such distributions, and particularly the extent of inter-individual heterogeneity present in a host population, can have key downstream consequences for epidemic dynamics (e.g. 14,16,21,22), Thus, it is critical to understand how factors such as acquired protection from prior pathogen exposure alter the degree of population-level heterogeneity in susceptibility, and its epidemiological consequences.

Host reinfection is a common but understudied feature of many host-pathogen systems (23–28), including SARS-Cov-2 (29), pneumococcal disease (30), influenza (31), malaria (32), and other diseases of agricultural and wildlife significance. Because the protection acquired from prior pathogen infection is often incomplete and/or wanes over time, reinfections occur even in systems where hosts have lower mean susceptibility during secondary exposures, relative to individuals exposed for the first time (33–36). While there is growing appreciation for the pervasiveness of heterogeneity in acquired host protection in response to vaccination or

infection across systems (31,35,37,38), prior work has largely focused on how host protection

from vaccination or prior infection influences mean population traits, rather than variability

among individuals in a given population (10). As such, it remains unknown how host protection

acquired from prior pathogen infection alters the shape of the susceptibility distribution present

in a given host population.

Prior pathogen exposure can modify inter-individual heterogeneity in susceptibility via

several mechanisms. First, prior exposure to pathogens could generate incomplete (or "leaky")

protection similarly for all exposed hosts (35), such that all exposed hosts acquire identical

levels of protection against reinfection. Alternatively, prior pathogen exposure could result in

relatively complete acquired protection for some proportion of hosts, while others remain

relatively unprotected (termed "all-or-nothing" immunity (35)). Epidemiological models often

select either wholly leaky protection of hosts that allow every host an identical rate of reinfection,

or all-or-nothing protection such that some hosts remain in the recovered class while a subset is

routed back to the susceptible class, representing two extremes of how heterogeneity in

susceptibility is incorporated into disease models (10,20). Whether and how prior exposure to a

pathogen alters the shape of the susceptibility distribution for a given host population likely

depends on the mode of action of acquired protection ["leaky", "all-or-nothing" (10,20), or some

intermediate (38)], as well as the extent of prior pathogen exposure that hosts experience (39).

For example, host exposure to low pathogen doses, which can occur during the natural

transmission process for many pathogens (e.g. 40–42), might be more likely than higher

exposure doses to generate incomplete host protection (43,44) and thus to induce population-

level heterogeneity in susceptibility.

Regardless of the underlying mechanisms, effects of prior exposure on population-level

heterogeneity in host susceptibility are particularly important to characterize because this

heterogeneity can have key epidemiological consequences, including lower predicted outbreak

sizes (20) and higher rates of reinfection (22). This is in part due to the process of cohort

selection, through which the most susceptible individuals are infected first in any given epidemic

time step, leaving lower mean susceptibility among remaining uninfected hosts in a population

(45). Experimental quantification of population-level heterogeneity in susceptibility draws on

quantitative microbial risk assessment approaches (46) to estimate susceptibility distributions

using a series of increasing pathogen challenge doses. Because only the most susceptible

hosts become infected at low challenge doses, whereas even the least susceptible hosts

become infected at the highest challenge doses, dose-response models can be used to

estimate the degree of host heterogeneity in susceptibility in a given population (20).

Importantly, by using controlled pathogen exposure doses, experimental studies allow direct

quantification of host heterogeneity in susceptibility without the confounds of variation in

exposure, a key additional source of heterogeneity in free-living populations (47). Thus, dose-

response studies of non-human animals are particularly useful for isolating effects of factors

such as vaccination (20) or maternal parasite exposure (19) on the degree of heterogeneity

present in a host population as a whole. Here we apply such experimental approaches to ask

how prior pathogen exposure alters the degree of inter-individual heterogeneity in susceptibility.

Robustly quantifying changes in population susceptibility distributions that occur with prior

pathogen exposure is challenging due to large sample size requirements of experimental dose

response studies. Nonetheless, studies of host susceptibility distributions are critical for

adequate characterization of host-pathogen dynamics, including the potential downstream

effects of inter-individual heterogeneity on pathogen strain coexistence and virulence evolution

(21,48,49).

Here we investigate effects of host prior exposure on inter-individual heterogeneity in

susceptibility using a tractable wildlife-pathogen system: a common North American songbird

species (house finch; *Haemorhous mexicanus*) and its bacterial pathogen, *Mycoplasma*

gallisepticum (MG), which causes severe conjunctivitis. The potential effects of prior pathogen

exposure on population-level heterogeneity in susceptibility are particularly relevant in this

system because MG exposure doses are variable across free-living hosts: susceptible birds

contact MG that was deposited onto bird feeder surfaces in variable amounts by infected

individuals (50,51). Further, although the conjunctivitis caused by MG indirectly results in

significant mortality in finches, free-living hosts recover at high rates from initial infection (52),

leaving a pool of recovered hosts in natural populations. Experimental studies show that

recovered hosts have significant acquired protection relative to pathogen-naive hosts, but can

still be successfully reinfected at high rates, even with homologous pathogen strains (39,53).

Finally, our work to date suggests that experimental variation in the degree of prior exposure to

MG (in both dose and number of exposures) results in heterogeneous protection from

reinfection among finches, as measured by conjunctival pathogen loads following reinfection

challenge (39,43,44). Whether prior exposure to MG specifically alters inter-individual

heterogeneity in susceptibility, a population-level trait that requires dose-response approaches

to robustly quantify, has not yet been examined.

To test how variation in prior pathogen exposure alters population-level distributions of

susceptibility, we first experimentally varied the degree of prior pathogen exposure categorically

(none, low-dose, or high-dose) in wild caught-finches that we confirmed had no exposure to MG

prior to capture. After recovery from prior exposure treatments, we measured host susceptibility

(0|1) to secondary dose challenge (Figure 1), fitting dose-response models for each treatment

group to determine whether host prior exposure treatment altered the degree of population-level

heterogeneity in susceptibility. Specifically, for each treatment group, we considered population-

level distributions of susceptibility that represent either identical infection probabilities

(homogeneous model) or models that represent inter-host variation in rates of infection

(heterogeneous model, where numerical susceptibility values were estimated according to a

fitted gamma distribution (16,19,20,54)). Importantly, our models allowed us to disentangle

effects of prior exposure on mean susceptibility from changes in heterogeneity in susceptibility.

Lastly, to determine effects of exposure-induced population heterogeneity on resulting epidemic

dynamics, we parameterized an SIR model for this system with susceptibility distributions and

expected mortality rates in the wild estimated from our experimental data.

Figure 1. Experimental design for quantifying how prior exposure to *Mycoplasma gallisepticum* alters host heterogeneity in susceptibility in house finches (n=150 for final analysis). Five weeks after one of three prior exposure (PE) treatments (none, low, or high dose; left panel), all birds received a secondary dose challenge (right panel) to assess heterogeneity in susceptibility. Our primary data set was whether birds became infected (0|1) in response to a given secondary dose (right). *However, to improve model fits, we also used bird responses to low-dose PE treatment (left, asterisk), which fell intermediate to our highest secondary challenge doses (300 and 7000 Color Changing Units [CCU]/mL), to quantify the proportion of birds with no prior exposure (at the time of low-dose PE) that become infected at a 750 CCU/mL dose. Made in Biorender.

Results

Prior exposure to pathogens augments population-level heterogeneity in susceptibility.

Birds with no prior pathogen exposure at the time of secondary dose challenge had low

variability in host susceptibility (coef of variation [CoV] from gamma distribution = 0.899) and

there was no support that heterogeneous dose-response models were an improvement over

models assuming homogeneous host susceptibility (defined here as identical infection

probabilities across hosts in a population) (Fig 2a; deviance homogeneous = 1.248; deviance

heterogeneous = 3.307). Birds with no prior pathogen exposure also had high mean

susceptibility to infection (mean susceptibility [fitted gamma distribution]=1.181). In contrast,

dose-response curves were shallower and fewer birds were infected upon secondary exposure

in both the low (mean susceptibility = 0.446 [fitted gamma], CoV [fitted gamma] = 1.630; Fig 2b),

and high-dose (mean susceptibility = 0.192 [fitted gamma], CoV [fitted gamma] = 2.511; Fig 2c)

prior exposure groups. Both prior exposure treatments were better described by models

assuming heterogeneous versus homogeneous susceptibility (Fig 2; likelihood ratio tests

heterogeneous versus homogeneous (46), low dose: P = 0.020, high dose: P = 0.033).

169
Figure 2. Dose response curves for house finch susceptibility to secondary challenge with
Mycoplasma gallisepticum across prior exposure treatments. Points (+/- 1SE) show the fraction
of individual birds (n=10-12 birds for most points; individual responses are 0|1) infected at each
secondary exposure dose, shown as Color Changing Units (CCU)/uL. Lines indicate model fits,
with blue indicating gamma (heterogeneous) model fits, and red dashed lines indicating
homogeneous model fits. Panel labels show prior exposure treatment (birds in the no prior
exposure treatment were pathogen-naive at the time of secondary dose challenge). In hosts
with prior pathogen exposure (low and high-dose prior exposure groups), the gamma model
(which accounts for inter-individual heterogeneity) was better supported via likelihood ratio tests.
180

Fig 3. Host susceptibility distributions for house finches from variable prior exposure treatments:
no prior exposure (A,B); low-dose (C); or high-dose (D). Colored lines show estimated
susceptibility distributions from either homogeneously (A) or gamma-distributed (B-D) models
(note distinct axes for the two models). In (A), host infection probability per 1000 bacterial
particles (*p*) is shown as the single best fit parameter *p* (dotted vertical lines represent 1
standard error) for the homogeneous model, which was the best fit model for the no prior
exposure group (see *Results*). In (B-D), the best fit parameters (shape and scale) for gamma
distributions (teal lines) are listed for each group, and vertical gray lines indicate mean
susceptibility (*x*) for that treatment. Lighter shading represents 95% confidence regions for
gamma distributions, obtained by bootstrapping chi-squared residuals to create 1,000
pseudoreplicates of infection data and then refitting the model to pseudoreplicates, as per
(20,46). The gamma model was the best fit for only the low-dose and high-dose groups.
Gamma estimates are also shown for the no prior exposure group (B) because this allowed
more equivalent comparisons for certain SIR simulations (see *Methods*).

To estimate effects of prior exposure on host mortality rates in the wild, we used clinical scores

of eye disease in response to secondary dose challenge to estimate mean mortality rates for

each prior exposure treatment group (as per (34); see *Methods*). As expected, predicted host

mortality rates decreased with the degree of prior host exposure (Table S2), declining from

0.05894/day in birds with no prior exposure, to 0.02806/day in birds with prior low-dose

exposure, and to a negligible value ($2.483*10^{-17}$) in birds with prior high-dose exposure.

Exposure-induced heterogeneity in susceptibility reduces epidemic size. Using these

empirically-derived susceptibility distributions and predicted mortality rates (Fig 3, Table S2) to

parameterize an SIR model of this system, we find that the total epidemic size is largest

(97.13% of the population) for a host population with no prior pathogen exposure, while for low

and high-dose prior exposure groups, the total epidemic size is significantly reduced (33.67%

and 16.13% of the population, respectively; blue bars, Fig 4A). Because prior pathogen

exposure also protects hosts from mortality, there is no mortality during the epidemic in the

high-dose prior exposure population, while there is 16.28% mortality in the low-dose prior

exposure population, and more than 64.37% mortality in the naive host population (darker

colored portion of bars, Fig 4A; dashed lines, Fig 4C).

We also examined whether the observed reductions in epidemic size with host prior exposure

are driven primarily by changes in the susceptibility distribution, versus reductions in mean host

susceptibility (blue versus red bars within each prior exposure treatment group, Fig 4A, B). For

empirically determined best fit parameters, epidemic size for a paired simulation assuming

homogeneous rather than heterogeneous susceptibility (while holding mean susceptibility

constant) was larger by 49% and 56%, under low dose and high dose prior exposure,

respectively (low dose, fitted mortality: 49.12 epidemic size difference [hom-het]; high dose,

fitted mortality: 56.55 epidemic size difference [hom-het]). We simulated the heterogeneous and

homogeneous models using the parameter estimates obtained from bootstrapping chi-squared

residuals (see Fig 3). Assuming homogeneous versus heterogeneous susceptibility (while

holding mean susceptibility constant for a given prior exposure treatment) results in consistently

larger outbreaks (Fig S1; difference in epidemic size = homogeneous [red]-heterogeneous

[blue] outbreak size for a given mean susceptibility). Indeed, across all simulations, outbreaks

are never smaller in models assuming homogeneous versus heterogeneous susceptibility (all P

< 0.0001; Fig S1; epidemic size differences [95% CIs of hom-het epidemic size] for each

simulation: low dose, fitted mortality [2.078,49.12]; high dose, fitted mortality [0.8994,61.29]; low

dose, fixed mortality [0.1187,40.77]; high dose, fixed mortality [0.010,36.35]). Together, these

results indicate effects of host heterogeneity *per se* on outbreak size, which act above and

beyond the effects of lower mean susceptibility from prior pathogen exposure.

Reductions in mean susceptibility with host prior exposure also contribute to smaller outbreak

sizes, but the extent depends on the assumed level of host mortality. When controlling for

mortality differences (i.e., all populations are assumed to have mortality equivalent to the naïve,

no-prior exposure population), the outbreak size of both groups with prior pathogen exposure

(low or high) is reduced (Fig 4B,D). In the case of simulations assuming homogeneous

susceptibility (red bars, Fig 4B) while accounting for differences in mean protection with prior

exposure, this reduction is *entirely* due to lower mean host susceptibility resulting from prior

pathogen exposure. However, when heterogeneity in susceptibility and reduced mean

susceptibility are both accounted for, further reductions in outbreak size and mortality are found

for the low-dose prior exposure group (blue bars and lines, Fig 4B,D). Further reductions from

heterogeneity *per se* are not possible for the high-dose prior exposure group, for which our

model produced an outbreak size of nearly 0 from changes in mean susceptibility alone.

In the more biologically realistic setting (Fig 4A,C) with changes in susceptibility and host

mortality in response to prior exposure both accounted for, we find a more complicated pattern.

If we assume a homogeneous population (red bars) with the mean susceptibility found

empirically for low and high-dose prior exposure groups (and mortality rates set as empirically

determined values), the epidemic sizes drastically increase (82.79% and 72.68% of the

population for the low and high prior exposure, respectively) relative to models accounting for

host heterogeneity in susceptibility (blue bars and lines; Fig 4A,C). The large outbreak in the

high-dose prior exposure population (despite very low mean susceptibility in this group) is due

to the absence of disease-induced mortality in this population, resulting in a higher basic

reproductive number because all individuals survive the entire infectious period.

Fig 4. Cumulative epidemic size (proportion of individuals who ever entered the infectious class) (A,B); dynamics of cumulative infections and deaths (top: C, D; deaths: dashed lines) over time, and numbers of infectious hosts over time (bottom: C, D) in our SIR model, as a function of prior exposure (abbreviated as "exp.") treatment (x-axis for A,B; columns for C,D). Total starting population was 100; darker shading within color represents individuals (within cumulative totals) that did not survive the epidemic. Blue bars and lines use fitted heterogeneous susceptibility (gamma distribution) for all prior exposure groups (Fig 3, Table S2). Red bars and lines assume homogeneous susceptibility for all groups, with mean susceptibility for each group equal to that of the fitted heterogeneous distribution for that treatment group. Left panels (A,C) use fitted mortality rates, whereby estimated mortality rates declined with prior exposure, while right panels (B,D) assume the fixed mortality rate of all groups is equal to that of the no prior exposure birds.

Discussion

Discussion

Here we use a naturally-occurring host-pathogen system characterized by high rates of

reinfection (43) to test how acquired protection from variable degrees of prior pathogen

exposure influences population-level distributions of host susceptibility. Using experimental

infection data for birds across three distinct prior exposure treatments, we tested for population-

level heterogeneity by statistically comparing dose response models that assume either

identical susceptibilities among hosts in a population (homogeneous model) or inter-host

variation in susceptibility (heterogeneous model). We find that prior exposure to either low or

high doses of pathogen significantly augments the extent of population-level heterogeneity in

susceptibility relative to a pathogen-naive host population, with population-level heterogeneity

increasing with the degree of prior pathogen exposure, from none, low, to high-dose. In addition,

our SIR model demonstrates that the observed changes in the host susceptibility distribution

with prior pathogen exposure have key effects on epidemic dynamics. After accounting for

estimated effects of prior exposure on host mortality, changes in population-level heterogeneity

in susceptibility had stronger effects on outbreak dynamics than reductions in mean host

susceptibility, resulting in dramatically lower outbreak sizes.

The past several decades have brought growing recognition that many host-pathogen

systems are characterized by reinfection potential, whether shortly following initial infection and

recovery, or over longer timescales after any host acquired protection from prior infection has

waned (23–26,29,55–58). Despite the ubiquity of highly specific immune memory that allows

hosts to respond rapidly to and effectively resist reinfection with the same pathogen, the

acquired protection generated by prior pathogen exposure is often incomplete, even in

vertebrate taxa with highly specific antigen repertoires and associated immune memory (59).

Thus, some degree of reinfection is still possible, despite lower mean susceptibility in hosts with

acquired protection (33,48). Given the importance of reinfection in many systems, it is critical to

characterize variability in key epidemiological traits for host populations made up of individuals

with some degree of prior pathogen exposure. For example, two longitudinal studies of repeated

malarial episodes in children documented substantial heterogeneity in the length of time until

Plasmodium falciparum reinfection (60) among children and the degree of acquired protection

generated against clinical malaria (60,61). However, this heterogeneity was attributed primarily

to variation among children in the extent of repeated exposure to the parasite. Because such

heterogeneity in exposure to pathogens is common in natural populations (47), it can be

challenging to quantify effects of acquired protection on heterogeneity in host susceptibility *per*

se without the use of controlled experiments. Nonetheless, it is critical to characterize effects of

acquired protection on host susceptibility distributions in particular, because population-level

heterogeneity in susceptibility can influence the likelihood and severity of outbreaks

(3,10,15,62), as well as the evolution of virulence for pathogens (21,48,49,63).

One intuitive prediction is that host acquired protection from prior exposure to pathogens

should homogenize variation in susceptibility in a population by reducing susceptibility to

reinfection similarly for all individuals. Instead, we found that the protection acquired from prior

exposure to a naturally-occurring bacterial pathogen, whether at low or high doses, significantly

augmented population-level heterogeneity in susceptibility in house finches. Importantly, while

our dose-response estimates for birds with no prior exposure to MG are well supported by

models assuming homogeneous susceptibility, this does not indicate that there is no biological

variation in susceptibility present within this treatment group. Instead, this result suggests that

variation in susceptibility among house finches without prior pathogen exposure is of low

magnitude, which is further supported by the low coefficient of variation of the estimated gamma

distribution for this group. Prior studies in this system find notable variation among both

individuals and populations in other types of host responses to experimental MG infection,

including infection severity (pathogen loads), disease severity (eye scores), and tolerance (per-

pathogen disease severity) (e.g. 64–67), even for birds with no prior exposure to MG infection.

Whether host susceptibility *per se* tends to be less variable than other types of host responses,

both in this system and others, is an interesting topic for future inquiry, particularly given the key

epidemiological consequences of heterogeneity in host susceptibility (14–16).

Few studies have explicitly tested how population-level distributions of host susceptibility

change in the presence of prior pathogen exposure or host protection. However, two

experimental studies that examined how other forms of acquired protection (transgenerational

pathogen exposure, and vaccination) influence the shape of host susceptibility distributions

found surprisingly similar results to what we report here. First, Ben-Ami et al. (19) showed that

maternal exposure to a high dose of a bacterial pathogen in *Daphnia magna* widened the

susceptibility distribution of the offspring without affecting its mean. Second, Langwig et al. (20)

found that vaccination of rainbow trout against a virus augmented population-level heterogeneity

in susceptibility to that virus relative to unvaccinated individuals, while also reducing mean

susceptibility. Interestingly, the results from this vaccination study qualitatively match what we

found for house finches, even though the protection that hosts acquire from active infection can

be stronger and/or more variable in strength or duration than that acquired by vaccination (e.g.

68,69). Further, our detected pattern of increasing heterogeneity in susceptibility with higher

degrees of prior pathogen exposure, from no, low-dose, to high-dose prior exposure treatments,

suggests that stronger stimulation of host acquired immune responses by high pathogen doses

(in this case, 30000 CCU/mL) can, at least in this system, lead to even more variability across

individuals in a population. Intriguingly, we observed highly variable responses to prior exposure

despite using an MG strain (VA1994) isolated from free-living house finches over 25 years ago.

Because significant host evolution has occurred since then (64,67), one might expect less

variable host responses to the strain used here, compared to a more recent MG strain, if host

resistance traits have been favored or even fixed in the host population (70). Nonetheless, prior

exposure to an older, basal house finch MG strain still induced significantly higher levels of host

heterogeneity in susceptibility, even at low prior exposure doses. Because strain traits such as

virulence may influence the degree of host heterogeneity in a population (48), future work

should examine whether prior exposure to MG strains more virulent and derived than the strain

used here generates even higher degrees of heterogeneity in susceptibility in a host population.

We used an SIR model parameterized empirically to predict how the population-level

susceptibility distributions induced by host prior exposure would influence epidemic dynamics in

this system. Consistent with prior work using SIR (20,22) and dynamic network approaches

(71), we show that population-level heterogeneity in susceptibility (here as a result of prior

pathogen exposure) suppresses overall outbreak size relative to models that assume

homogeneous host susceptibility. Importantly, under either set of mortality assumptions used in

our model, outbreak size is always reduced when prior exposure generates heterogeneity in

susceptibility, compared to the homogeneous case that still accounts for reductions in mean

host susceptibility with prior exposure. These results indicate direct effects of exposure-induced

heterogeneity *per se* on outbreak size, effects which act above and beyond effects of mean

susceptibility from prior pathogen exposure on outbreak size. Moreover, under the more realistic

assumption of host protection from mortality after prior pathogen exposure, the reductions on

overall epidemic size in our models appear to be driven predominantly by increased population-

level heterogeneity in susceptibility. This observation is consistent with herd immunity

derivations in a similar SEIR model, which finds strong dependence on the coefficient of

variation (72).

An earlier SIR model of vaccine-induced heterogeneity similarly found that augmented

population-level heterogeneity in susceptibility reduced outbreak size (20), but this model did

not incorporate potential changes in host mortality rate. Reductions in mortality rate in hosts with

prior pathogen exposure, which we estimated were significant in the house finch system, are

likely common in hosts with acquired protection, because such protection often reduces host

disease and mortality more strongly than infection risk (e.g. 73,74). In our model results, the

degree to which prior exposure-induced reductions in mean susceptibility drive outbreak size

depends on assumptions of mortality rates in our model. In our model with empirically-derived

mortality rates, infected birds in the prior exposure groups are more likely to survive infection

(lighter color sections of bars; Fig 4A). Intriguingly, these higher survival rates nearly outweigh

the protective effects of reduced mean susceptibility in previously-exposed hosts, resulting in

relatively little reduction in epidemic size in both prior exposure groups (assuming

homogeneous susceptibility for both groups; red bars, Fig 4A), despite significantly lower mean

host susceptibility in birds with prior exposure. In contrast, when mortality rates are held

constant regardless of prior exposure treatment, exposure-induced reductions in mean host

susceptibility alone result in substantially smaller outbreak sizes (red bars; Fig 4B). These

results indicate that effects of prior exposure on mortality rate and susceptibility distributions are

both critical to account for in epidemiological models, as they may interact to drive outbreak

size.

One limitation of our SIR model is that it assumes that epidemics begin in host

populations entirely composed of individuals of one prior exposure type. While such models are

likely realistic representations of host populations at certain times of the year (e.g., epidemics in

late summer that occur in juvenile flocks composed of fully susceptible finches (75,76)), future

models should also consider scenarios whereby populations are composed of some individuals

with no prior exposure, and others with varying degrees of prior pathogen exposure. Developing

a simulated population to such a state would require knowing how the level of susceptibility of

individual birds is altered by prior exposure, for which no data exists. Another limitation is the

choice of an SIR rather than an SIRS model that allows for loss of immunity over time, as is the

case in the house finch-MG model system, in which protection wanes over approximately one

year (53). To disentangle effects of population-level heterogeneity of susceptibility induced by

prior exposure, we focus this study on a single, short timescale epidemic while ignoring

demographic effects and re-infection, similar to previous work (38,77,78). Allowing for

reinfection in our model would require an understanding of how susceptibility and infectivity are

linked, because such correlations have an impact on the progression of epidemics (77,78).

However, such correlations are particularly challenging to quantify in natural host-pathogen

systems. Overall, our work represents an important first step in incorporating one key source

(prior exposure) of population-level heterogeneity in susceptibility into mathematical models.

Although the importance of individual variation in infection-derived immunity for epidemiological

dynamics is increasingly recognized (31,35,79), to our knowledge, our study is the first to

explicitly quantify how acquired protection from prior pathogen infection influences population-

level distributions of host susceptibility.

In summary, our results highlight the key importance of variable protection from prior

pathogen exposure in driving host heterogeneity in susceptibility, a population-level trait with

key downstream consequences for epidemiological and evolutionary dynamics (15,22,48).

Because the host protection acquired from a prior infection (or vaccination) is often incomplete

and/or wanes over time in diverse vertebrate systems (e.g. 29,58), infection-induced

heterogeneity in susceptibility is likely a common phenomenon for many animal populations,

including humans. While work to date has largely focused on characterizing innate, individual-

level sources of host heterogeneity in susceptibility [sex (80), genotype (81), etc.], our results

show that population-level heterogeneity in susceptibility can be readily "induced" by prior

pathogen exposure, with key downstream consequences.

Methods

Bird Capture and Housing

We captured 157 mixed-sex, hatch-year house finches (aged by plumage (82)) in

Montgomery County, Virginia using a combination of cage traps and mist nets in June-Aug

2021. Capture and collection were approved by Virginia Department of Game and Inland

Fisheries (066646) and USFWS (MB158404), and all handling and care procedures were

approved by the Virginia Tech (VT) IACUC. Birds were housed in pairs for a two-week minimum

quarantine period in an indoor animal facility, and captured every 3-4 days to assess potential

clinical signs of mycoplasmal conjunctivitis. Any birds with signs of conjunctivitis (see below)

were immediately isolated, along with their cage-mates, and were not used in experiments.

For all birds that never showed clinical signs of conjunctivitis during quarantine, blood

samples were collected 14-18 days post-capture and run for ELISA via a commercial IDEXX kit

(99-06729) using methods previously described (83). Only birds seronegative via ELISA on

days 14-18 post-capture were assigned to MG-exposed treatment groups (Table S1; all birds

except n=3 "prior sham-0 challenge" birds; see Supplement). All birds were housed individually

and then resampled one week before prior exposure treatments for final confirmation of

seronegativity just prior to experiment initiation.

Experimental Design and Timeline

We created variation in prior exposure using three categorical treatments (Figure 1,

Table S1): no prior exposure to MG (inoculation with the same volume of sterile Frey's media as

a "sham" treatment), low-dose prior exposure (dose of 750 Color Changing Units [CCU]/mL of

MG), or high-dose prior exposure (dose of 30,000 CCU/mL of MG). Birds were infected with an

MG strain (VA1994 isolate of MG; [7994-1 (6P) 9/17/2018]) that was isolated shortly following

the pathogen's emergence in house finches; despite significant host and pathogen evolution

since then (64,84,85), this 1994 strain still results in pathogen load and eye scores trajectories

that are qualitatively similar to later-isolated strains (VA2013 (64)). Birds were inoculated with 70

uL of their assigned treatment concentration directly into the conjunctivae via micropipette

droplets, and then allowed to recover (with recovery quantified at day 41 post-infection; see

below).

Forty-two days after prior exposure treatment, birds were given a secondary dose

challenge with the same MG strain (again, 70uL directly inoculated into the conjunctivae). Using

a fully factorial design (Table S1), we inoculated birds from each prior exposure treatment with

one of five MG exposure doses to assess heterogeneity in susceptibility (Figure 1). We selected

prior exposure treatments and challenge doses (also termed "secondary doses") most likely to

generate and detect differences in susceptibility based on our published work on conjunctival

loads (44), and we minimized other inherent sources of variation in susceptibility in our wild-

caught birds by using same-age birds, excluding birds with detectable prior exposure at capture,

and randomizing assignments of sex within treatment groups, such that sex ratios were similar

across treatment combinations. Because of the potential for birds to remain infected from prior

exposure treatments until the time of secondary challenge (42 days post-prior exposure), we

included a set of controls (n=10 total birds) that received prior exposure doses of pathogen

(either low or high-dose) and then received a control inoculation of sterile media at day 42.

These 10 birds were all qPCR negative for MG infection by day 41 and during all three

subsequent sample periods post-secondary challenge (see Supplement).

Because our main interest was in how prior exposure treatment alters host susceptibility

(1|0) to secondary dose challenge, we generally do not analyze or present data collected prior

to secondary challenge (day 0), with one exception. To improve our dose response model fits

for the "no prior exposure" treatment group, we included susceptibility data generated in

response to the initial, low-dose prior exposure treatment. We use only the low-dose prior

exposure treatment for this purpose, and not the high-dose prior exposure treatment (30000

CCU / mL) because the low-dose treatment uniquely provided an exposure dose (750 CCU /

mL) that fell intermediate to those used in our secondary challenge doses (which were

otherwise 30, 100, 300, and 7000 CCU/mL), informing a key part of the dose response curve.

Thus, each of the 50 finches given a low-dose prior exposure treatment and then one of five

secondary challenge doses (Table S1) contributed two data points to the analyses: (i) their

susceptibility (1|0) to the "no-prior exposure" dose of 750 CCU/mL, with data collected on days -

42 to -28 prior to secondary challenge; and (ii) their susceptibility (1|0) to one of five secondary

doses, with data collected on days 4-14 post-secondary challenge. All other birds contributed

only one data point to the data-set: susceptibility (1|0) to one of five secondary doses, with data

collected on days 4-14 post-secondary challenge.

Final sample sizes

We began with a balanced design (n=12 birds for all non-0 MG challenge doses in Table

S1; n=157 total birds), but one individual died of unknown causes prior to secondary dose

challenge. A second individual was excluded from all analyses after study completion because,

when ELISA assays on frozen blood samples were completed, that individual was discovered to

have been MG-seropositive just before prior exposure treatments were given. Finally, five

additional individuals were later determined to be qPCR positive one day prior to secondary

challenge (see *Susceptibility and disease sampling*), and thus were eliminated from all analyses

to ensure that our susceptibility data represent true responses to secondary challenge rather

than residual effects of prior exposure treatment. Final sample sizes for all analyses were n=150

individuals (Table S1).

Susceptibility and disease sampling

We quantified whether birds were susceptible to infection or not (Y/N, 1|0) at a given MG

exposure dose via qPCR of conjunctival swab samples collected post-exposure. We also

visually scored clinical signs of conjunctivitis (all scoring was done blind to a bird's treatment) to

use as a proxy for the likelihood of mortality during infection (see below). At each sampling

point, we first scored clinical signs of conjunctivitis on a scale of 0 to 3 for left and right

conjunctiva separately, as per (86), with 0.5 score intervals used when clinical signs were

intermediate between two integer scores. We summed the scores (left plus right conjunctiva)

within a given sample day for a total maximum score of 6 for a given bird per day. After scoring,

each bird's conjunctiva was swabbed for 5s with a sterile cotton swab dipped in tryptose

phosphate broth (TPB). Swabs were swirled in 300uL of sterile TPB and then wrung out into the

sample collection tube. Samples from both eyes were pooled within sampling date for a given

individual and frozen at -20°C until processing. Genomic DNA was extracted from samples

using Qiagen DNeasy 96 Blood and Tissue kits (Qiagen, Valencia, CA) as previously described

(84). Extracted genomic DNA from each sample was used to measure overall numbers of MG

present in the conjunctiva using a qPCR assay targeting the *mgc2* gene of MG using the

primers from (87) and qPCR methods outlined in (84).

Conjunctival swab samples and clinical scores were taken on days 4, 7, 14, and 21 post-

secondary-challenge (and on days 7 and 14 post-prior exposure treatments). However, we only

use conjunctival swab data from the first three sample points (days 4, 7, or 14 post-challenge)

as most relevant for quantifying susceptibility. To account for potential low-level contamination

in our qPCR assay (see Supplement) and the possibility that MG loads from prior exposure

treatment might still be present in some birds at the time of secondary dose challenge, we

considered a bird as susceptible to a given dose if they had a conjunctival MG pathogen load >

50 copies at any of the sample points from day 4 to 14 post-challenge. This cut-off fell above the

highest MG load detected among birds that were given a sham secondary challenge treatment

(but had been previously exposed), and thus should not encompass low MG loads that

represent residual loads from prior exposure treatments. We also sampled all birds in the study

on day 41 post-prior exposure, which was one day prior to secondary challenge treatment (day

42). As noted above, we eliminated 5/155 birds deemed infected on day 41 from all analyses

(see Supplement).

Dose response models and parameterization

We used dose-response models to describe the susceptibility of naive and previously

exposed birds challenged with MG (20,46). We fit the models to data by minimizing the

deviance between the data and the model and compared models using likelihood ratio tests that

accounted for differences in the number of parameters between models (46). We assumed

heterogeneous susceptibility followed a gamma distribution (Equation 1) (10). We compared

models of homogeneous susceptibility (best supported for the no prior exposure group), with

models of gamma distributed susceptibility (best supported for the two prior exposure groups).

With these models we determined the proportion of individuals infected, I_{hom} or I_{het}, via

Equation 1

517
$$I_{hom} = 1 - e^{-pd}$$

518
$$I_{het} = 1 - \int_0^\infty e^{-xd} \frac{x^{k-1} e^{-x/\sigma}}{\sigma^k \Gamma(k)} dx$$

Here, d (dose) is the exposure concentration in CCU/μl. In I_{hom}, the subscript *hom* denotes a

model assuming homogeneous susceptibility with p representing the host infection probability

per 1000 bacterial particles.The model with subscript *het* assumes individuals have

susceptibility (x) that is distributed according to a gamma distribution with shape parameter k

and scale parameter σ. Here, $\Gamma(k)$ is the gamma function evaluated at the shape parameter k.

We calculated coefficients of variation for the gamma models as the standard deviation divided

by the mean.

Fitting mortality data

We use clinical scores of conjunctivitis collected post-secondary challenge (with scores

per bird per day ranging from 0 to 6; see above) to estimate mean mortality rates in the wild for

each prior exposure treatment group. We assume that these ordinal scores monotonically

predict host mortality because several lines of evidence indicate that increasingly severe

conjunctivitis and resulting visual and behavioral impairment result in higher mortality risk for

house finches in the wild: 1) the presence of conjunctivitis is linked with higher mortality in wild

birds (52); 2) higher conjunctivitis severity predicts faster capture time in mock-predation events

(i.e., a human capturing the bird by hand) in captivity (88); and 3) indirect mortality from

predators appears to be the primary source of mortality for infected birds in the wild, because

birds in captivity (a predator-free environment) do not ever succumb to infection, even with

highly virulent strains (84). Thus, MG infection *per se* does not directly cause mortality in house

finches, but mortality in the wild (via predation or the inability to find food) is directly linked with

visual impairment and the associated disease that we measure.

Mortality rate was fit using nonlinear least squares regression (nls function in R) as the

nonlinear inverse logit function, 1/(1+exp(-z)) scaled by 6, where z is the eye lesion score

(integers between 0 to 6). This fit was then scaled by vmax (0.25; maximum mortality rate

observed in the field from (52)) such that the maximum mortality rate is vmax rather than 1. A

similar approach was used in previous work (34).

Epidemic model

We use a Susceptible-Infectious-Recovered (SIR) model to examine the effect of

heterogeneity in susceptibility on outbreak size and impact. For populations with determined

heterogeneity in susceptibility, we consider a continuous distribution of susceptibility, with

susceptibility, *x*, distributed according to the parameters (shape k and scale σ) of the gamma

distribution obtained from the dose response models. We model this with the differential

equation system

Equation 2

558
$$\frac{dS(x)}{dt} = -x\beta IS(x)$$

559
$$\frac{dI}{dt} = \beta I \int xS(x)dx - \gamma I - \mu I$$

$$\frac{dR}{dt} = \gamma I$$

Here, $S(x)$ is the number of susceptible individuals with susceptibility x, and x is a value

between 0 and infinity (note that for simulation purposes, we truncate the integral at 10, which

covers more than 95% of the full susceptibility distribution with our various choices of

parameters). I is the number of infectious individuals, and R is the number of recovered

individuals. Parameters for infectivity (β) and recovery (γ) were taken from a prior empirically-

parameterized model in this system (89) (Table S2), while disease-induced mortality (μ) was fit

as described above (See *Fitting mortality data*) for each prior exposure group (none, low, high).

Infectivity and recovery rates do not depend on prior exposure, while mortality rate does (in the

"fitted mortality" model). A density-dependent rather than frequency-dependent transmission

term was used due to the nature of transmission in this system, which increases with density

(76,90).

For simulation, we discretize the susceptibility distribution of the heterogeneous model into 300

evenly spaced susceptibility classes, represented by the midpoint susceptibility, x, for each

class. We assume a population size of 100 and seed infections with a single infected individual.

The initial population in each susceptibility class is determined empirically from the fitted

distributions by prior exposure treatment, discretized into the 300 evenly spaced susceptibility

classes which are truncated above susceptibility of 10 (See *Dose response models and*

parameterization for details on determination of the fitted distributions). We quantify the total

size of the epidemic as the cumulative fraction of infected individuals, and simulate for 500

days, after which nearly all epidemics have completely resolved. Because we simulate a short-

time scale single epidemic, we ignore demographic effects and assume recovered individuals

cannot be reinfected, despite documented waning of immunity to MG over the course of a year

(53).

To disentangle the effects of heterogeneity in susceptibility and changes in mean susceptibility,

we simulate outbreaks using the homogeneous model with identical mean susceptibility to the

discretized heterogeneous distribution. To generate confidence intervals on outbreak sizes, we

used the parameter estimates obtained from bootstrapping the chi-squared residuals of the

dose-response estimates. We then estimated the probability of obtaining an epidemic size

difference as extreme as observed (hom-het) using the best fit parameters, but with the

heterogeneous case resulting in larger epidemics (het-hom).

Data Availability Statement

The code and datasets generated for this study are available in the GitHub repository, at

https://github.com/klangwig/mg_dose_response_public.

Acknowledgments

We thank Edan Tulman for input on experimental design and MG isolate preparation, Chava

Weitzman, John Brule, and many other Hawley lab members for help with capture and

sampling, and Bambi Kelly for assistance with animal care. Research reported in this publication

was supported by the National Institute Of General Medical Sciences of the National Institutes

of Health under Award Number R01GM144972. The content is solely the responsibility of the

authors and does not necessarily represent the official views of the National Institutes of Health.

Author Contributions

D.M.H.: Conceptualization, Methodology, Resources, Investigation, Writing- Original Draft,

Supervision, Project administration, Funding acquisition

A.P.U.: Investigation, Data Curation, Writing - Review & Editing

J.S.A: Conceptualization, Methodology, Writing-Review and Editing, Funding acquisition

AEFD: Conceptualization, Methodology, Software, Formal Analysis, Writing - Review & Editing,

Funding acquisition

J.N.GL: Investigation, Writing-Review and Editing

S.J.G.: Conceptualization, Methodology, Writing-Review and Editing

L.M.C.: Conceptualization, Methodology, Software, Validation, Formal Analysis, Writing-Review

and Editing, Visualization, Funding acquisition

K.E.L.: Conceptualization, Methodology, Software, Validation, Formal Analysis, Writing-Review

and Editing, Visualization, Funding acquisition

References Cited

1. Smith KF, Goldberg M, Rosenthal S, Carlson L, Chen J, Chen C, et al. Global rise in human infectious disease outbreaks. J R Soc Interface. 2014 Dec 6;11(101):20140950.

2. Schmeller DS, Courchamp F, Killeen G. Biodiversity loss, emerging pathogens and human health risks. Biodivers Conserv. 2020 Aug 13;29(11–12):3095–102.

3. Lloyd-Smith JO, Schreiber SJ, Kopp PE, Getz WM. Superspreading and the effect of individual variation on disease emergence. Nature. 2005 Nov 17;438(7066):355–9.

4. Britton T, Ball F, Trapman P. A mathematical model reveals the influence of population heterogeneity on herd immunity to SARS-CoV-2. Science. 2020 Aug 14;369(6505):846–9.

5. Ke R, Martinez PP, Smith RL, Gibson LL, Mirza A, Conte M, et al. Daily longitudinal sampling of SARS-CoV-2 infection reveals substantial heterogeneity in infectiousness. Nat Microbiol. 2022 May;7(5):640–52.

6. Anderson TL, Nande A, Merenstein C, Raynor B, Oommen A, Kelly BJ, et al. Quantifying individual-level heterogeneity in infectiousness and susceptibility through household studies. Epidemics. 2023 Sep;44:100710.

7. Channappanavar R, Fett C, Mack M, Ten Eyck PP, Meyerholz DK, Perlman S. Sex-Based Differences in Susceptibility to Severe Acute Respiratory Syndrome Coronavirus Infection. J Immunol. 2017 May 15;198(10):4046–53.

8. Fricke-Galindo I, Falfán-Valencia R. Genetics Insight for COVID-19 Susceptibility and Severity: A Review. Front Immunol. 2021 Apr 1;12:622176.

9. Cobey S, Hensley SE. Immune history and influenza virus susceptibility. Curr Opin Virol. 2017 Feb;22:105–11.

10. Gomes MGM, Lipsitch M, Wargo AR, Kurath G, Rebelo C, Medley GF, et al. A missing dimension in measures of vaccination impacts. PLoS Pathog. 2014 Mar;10(3):e1003849.

11. Gomes MGM, Blagborough AM, Langwig KE, Ringwald B. Remodelling selection to optimise disease forecasts and policies. J Phys A: Math Theor. 2024 Feb 23;57(10):103001.

12. Dushoff J, Levin S. The effects of population heterogeneity on disease invasion. Math Biosci. 1995 Jul-Aug;128(1–2):25–40.

13. Novozhilov AS. On the spread of epidemics in a closed heterogeneous population. Math Biosci. 2008 Oct;215(2):177–85.

14. Hickson RI, Roberts MG. How population heterogeneity in susceptibility and infectivity influences epidemic dynamics. J Theor Biol. 2014 Jun 7;350:70–80.

15. Gomes MGM, Aguas R, Lopes JS, Nunes MC, Rebelo C, Rodrigues P, et al. How host heterogeneity governs tuberculosis reinfection? Proc Biol Sci. 2012 Jun 22;279(1737):2473–8.

16. Dwyer G, Elkinton JS, Buonaccorsi JP. Host heterogeneity in susceptibility and disease

dynamics: tests of a mathematical model. Am Nat. 1997 Dec;150(6):685–707.

17. Tuschhoff BM, Kennedy DA. Detecting and quantifying heterogeneity in susceptibility using
contact tracing data [Internet]. bioRxiv. 2023 [cited 2024 Jun 25]. p. 2023.10.04.560944.
Available from: https://www.biorxiv.org/content/10.1101/2023.10.04.560944

18. Gomes MGM, Ferreira MU, Corder RM, King JG, Souto-Maior C, Penha-Gonçalves C, et
al. Individual variation in susceptibility or exposure to SARS-CoV-2 lowers the herd
immunity threshold. J Theor Biol. 2022 May 7;540:111063.

19. Ben-Ami F, Ebert D, Regoes RR. Pathogen dose infectivity curves as a method to analyze
the distribution of host susceptibility: a quantitative assessment of maternal effects after
food stress and pathogen exposure. Am Nat. 2010 Jan;175(1):106–15.

20. Langwig KE, Wargo AR, Jones DR, Viss JR, Rutan BJ, Egan NA, et al. Vaccine Effects on
Heterogeneity in Susceptibility and Implications for Population Health Management. MBio
[Internet]. 2017 Nov 21;8(6). Available from: http://dx.doi.org/10.1128/mBio.00796-17

21. Pugliese A. The role of host population heterogeneity in the evolution of virulence. J Biol
Dyn. 2011 Mar;5(2):104–19.

22. Rodrigues P, Margheri A, Rebelo C, Gomes MGM. Heterogeneity in susceptibility to
infection can explain high reinfection rates. J Theor Biol. 2009 Jul 21;259(2):280–90.

23. Yamamoto T, Nagasawa I, Nojima M, Yoshida K, Kuwabara Y. Sexual transmission and
reinfection of group B streptococci between spouses. J Obstet Gynaecol Res. 1999
Jun;25(3):215–9.

24. Nardell E, McInnis B, Thomas B, Weidhaas S. Exogenous reinfection with tuberculosis in a
shelter for the homeless. N Engl J Med. 1986 Dec 18;315(25):1570–5.

25. Islam N, Krajden M, Shoveller J, Gustafson P, Gilbert M, Buxton JA, et al. Incidence, risk
factors, and prevention of hepatitis C reinfection: a population-based cohort study. Lancet
Gastroenterol Hepatol. 2017 Mar;2(3):200–10.

26. Versteegh FGA, Schellekens JFP, Nagelkerke AF, Roord JJ. Laboratory-confirmed
reinfections with Bordetella pertussis. Acta Paediatr. 2007 Jan 2;91(1):95–7.

27. Bazan JA, Tzeng Y-L, Stephens DS, Carter AM, Brown MA, Snyder B, et al. Repeat
Episodes of Symptomatic Urethritis Due to a Uropathogenic Meningococcal Clade. Sex
Transm Dis. 2020 Jan;47(1):e1–4.

28. Barenkamp SJ, Shurin PA, Marchant CD, Karasic RB, Pelton SI, Howie VM, et al. Do
children with recurrent Haemophilus influenzae otitis media become infected with a new
organism or reacquire the original strain? J Pediatr. 1984 Oct;105(4):533–7.

29. Pilz S, Theiler-Schwetz V, Trummer C, Krause R, Ioannidis JPA. SARS-CoV-2 reinfections:
Overview of efficacy and duration of natural and hybrid immunity. Environ Res. 2022
Jun;209:112911.

30. Sanz JC, Rodríguez-Avial I, Ríos E, Ramos B, Cercenado E. Recurrent pneumococcal
invasive disease in the region of Madrid during a five-year period. Infection. 2014
Jun;42(3):475–83.

31. Ranjeva S, Subramanian R, Fang VJ, Leung GM, Ip DKM, Perera RAPM, et al. Age-
specific differences in the dynamics of protective immunity to influenza. Nat Commun. 2019
Apr 10;10(1):1660.

32. Ranawaka MB, Munesinghe YD, de Silva DM, Carter R, Mendis KN. Boosting of
transmission-blocking immunity during natural Plasmodium vivax infections in humans
depends upon frequent reinfection. Infect Immun. 1988 Jul;56(7):1820–4.

33. Forshey BM, Reiner RC, Olkowski S, Morrison AC, Espinoza A, Long KC, et al. Incomplete
Protection against Dengue Virus Type 2 Re-infection in Peru. PLoS Negl Trop Dis. 2016
Feb;10(2):e0004398.

34. Fleming-Davies AE, Williams PD, Dhondt AA, Dobson AP, Hochachka WM, Leon AE, et al.
Incomplete host immunity favors the evolution of virulence in an emergent pathogen.
Science. 2018 Mar 2;359(6379):1030–3.

35. Le A, King AA, Magpantay FMG, Mesbahi A, Rohani P. The impact of infection-derived
immunity on disease dynamics. J Math Biol. 2021 Nov 12;83(6–7):61.

36. Altarawneh HN, Chemaitelly H, Hasan MR, Ayoub HH, Qassim S, AlMukdad S, et al.
Protection against the Omicron Variant from Previous SARS-CoV-2 Infection. N Engl J
Med. 2022 Mar 31;386(13):1288–90.

37. Gomes MGM, White LJ, Medley GF. Infection, reinfection, and vaccination under
suboptimal immune protection: epidemiological perspectives. J Theor Biol. 2004 Jun
21;228(4):539–49.

38. Zachreson C, Tobin R, Szanyi J, Walker C, Cromer D, Shearer FM, et al. Individual
variation in vaccine immune response can produce bimodal distributions of protection.
Vaccine. 2023 Oct 26;41(45):6630–6.

39. Leon AE, Hawley DM. Host Responses to Pathogen Priming in a Natural Songbird Host.
Ecohealth. 2017 Dec;14(4):793–804.

40. Aiello CM, Nussear KE, Esque TC, Emblidge PG, Sah P, Bansal S, et al. Host contact and
shedding patterns clarify variation in pathogen exposure and transmission in threatened
tortoise Gopherus agassizii: implications for disease modelling and management. J Anim
Ecol. 2016;85(3):829–42.

41. Konrad M, Vyleta ML, Theis FJ, Stock M, Tragust S, Klatt M, et al. Social transfer of
pathogenic fungus promotes active immunisation in ant colonies. PLoS Biol. 2012 Apr
3;10(4):e1001300.

42. Stroeymeyt N, Grasse AV, Crespi A, Mersch DP, Cremer S, Keller L. Social network
plasticity decreases disease transmission in a eusocial insect. Science. 2018 Nov
23;362(6417):941–5.

43. Weitzman CL, Ceja G, Leon AE, Hawley DM. Protection Generated by Prior Exposure to
Pathogens Depends on both Priming and Challenge Dose [Internet]. Vol. 90, Infection and
Immunity. 2022. Available from: http://dx.doi.org/10.1128/iai.00537-21

44. Leon AE, Fleming-Davies AE, Hawley DM. Host exposure history modulates the within-host

advantage of virulence in a songbird-bacterium system. Sci Rep. 2019 Dec 30;9(1):20348.

45. Aalen OO, Valberg M, Grotmol T, Tretli S. Understanding variation in disease risk: the
elusive concept of frailty. Int J Epidemiol. 2015 Aug;44(4):1408–21.

46. Haas CN, Rose JB, Gerba CP. Quantitative Microbial Risk Assessment. John Wiley &
Sons; 1999. 449 p.

47. Sweeny AR, Albery GF. Exposure and susceptibility: The Twin Pillars of infection. Funct
Ecol. 2022 Jul;36(7):1713–26.

48. Fleming-Davies AE, Dukic V, Andreasen V, Dwyer G. Effects of host heterogeneity on
pathogen diversity and evolution. Ecol Lett. 2015 Nov;18(11):1252–61.

49. Miller IF, Metcalf CJ. Vaccine-driven virulence evolution: consequences of unbalanced
reductions in mortality and transmission and implications for pertussis vaccines. J R Soc
Interface. 2019 Dec;16(161):20190642.

50. Adelman JS, Carter AW, Hopkins WA, Hawley DM. Deposition of pathogenic Mycoplasma
gallisepticum onto bird feeders: host pathology is more important than temperature-driven
increases in food intake. Biol Lett. 2013 Oct 23;9(5):20130594.

51. Dhondt AA, Dhondt KV, Hawley DM, Jennelle CS. Experimental evidence for transmission
of Mycoplasma gallisepticum in house finches by fomites. Avian Pathol. 2007
Jun;36(3):205–8.

52. Faustino CR, Jennelle CS, Connolly V, Davis AK, Swarthout EC, Dhondt AA, et al.
Mycoplasma gallisepticum infection dynamics in a house finch population: seasonal
variation in survival, encounter and transmission rate. J Anim Ecol. 2004 Jul;73(4):651–69.

53. Sydenstricker KV, Dhondt AA, Ley DH, Kollias GV. Re-exposure of captive house finches
that recovered from Mycoplasma gallisepticum infection. J Wildl Dis. 2005 Apr;41(2):326–
33.

54. Ben-Ami F, Regoes RR, Ebert D. A quantitative test of the relationship between parasite
dose and infection probability across different host-parasite combinations. Proc Biol Sci.
2008 Apr 7;275(1636):853–9.

55. Koopman JS, Henry CJ, Park JH, Eisenberg MC, Ionides EL, Eisenberg JN. Dynamics
affecting the risk of silent circulation when oral polio vaccination is stopped. Epidemics.
2017 Sep;20:21–36.

56. van den Hoogen BG, Herfst S, de Graaf M, Sprong L, van Lavieren R, van Amerongen G,
et al. Experimental infection of macaques with human metapneumovirus induces transient
protective immunity. J Gen Virol. 2007 Apr;88(Pt 4):1251–9.

57. Singleton R, Etchart N, Hou S, Hyland L. Inability to evoke a long-lasting protective immune
response to respiratory syncytial virus infection in mice correlates with ineffective nasal
antibody responses. J Virol. 2003 Nov;77(21):11303–11.

58. Edridge AWD, Kaczorowska J, Hoste ACR, Bakker M, Klein M, Loens K, et al. Seasonal
coronavirus protective immunity is short-lasting. Nat Med. 2020 Nov;26(11):1691–3.

59. Boehm T. Evolution of vertebrate immunity. Curr Biol. 2012 Sep 11;22(17):R722-32.

60. Rodriguez-Barraquer I, Arinaitwe E, Jagannathan P, Boyle MJ, Tappero J, Muhindo M, et
al. Quantifying Heterogeneous Malaria Exposure and Clinical Protection in a Cohort of
Ugandan Children. J Infect Dis. 2016 Oct 1;214(7):1072–80.

61. Addy JWG, Bediako Y, Ndungu FM, Valetta JJ, Reid AJ, Mwacharo J, et al. 10-year
longitudinal study of malaria in children: Insights into acquisition and maintenance of
naturally acquired immunity. Wellcome Open Res. 2021;6:79.

62. Ferrari MJ, Bansal S, Meyers LA, Bjørnstad ON. Network frailty and the geometry of herd
immunity. Proc Biol Sci. 2006 Nov 7;273(1602):2743–8.

63. Leggett HC, Buckling A, Long GH, Boots M. Generalism and the evolution of parasite
virulence. Trends Ecol Evol. 2013 Oct;28(10):592–6.

64. Henschen AE, Vinkler M, Langager MM, Rowley AA, Dalloul RA, Hawley DM, et al. Rapid
adaptation to a novel pathogen through disease tolerance in a wild songbird. PLoS Pathog.
2023 Jun;19(6):e1011408.

65. Ruden RM, Adelman JS. Disease tolerance alters host competence in a wild songbird. Biol
Lett. 2021 Oct;17(10):20210362.

66. Adelman JS, Kirkpatrick L, Grodio JL, Hawley DM. House Finch Populations Differ in Early
Inflammatory Signaling and Pathogen Tolerance at the Peak ofMycoplasma
gallisepticumInfection [Internet]. Vol. 181, The American Naturalist. 2013. p. 674–89.
Available from: http://dx.doi.org/10.1086/670024

67. Bonneaud C, Tardy L, Giraudeau M, Hill GE, McGraw KJ, Wilson AJ. Evolution of both host
resistance and tolerance to an emerging bacterial pathogen. Evol Lett. 2019 Oct;3(5):544–
54.

68. Wendelboe AM, Van Rie A, Salmaso S, Englund JA. Duration of immunity against pertussis
after natural infection or vaccination. Pediatr Infect Dis J. 2005 May;24(5 Suppl):S58-61.

69. Hobson D. Resistance to reinfection in experimental mouse typhoid. J Hyg . 1957
Sep;55(3):334–43.

70. Duffy MA, Sivars-Becker L. Rapid evolution and ecological host-parasite dynamics. Ecol
Lett. 2007 Jan;10(1):44–53.

71. White LA, Forester JD, Craft ME. Covariation between the physiological and behavioral
components of pathogen transmission: host heterogeneity determines epidemic outcomes.
Oikos. 2018 Apr;127(4):538–52.

72. Montalbán A, Corder RM, Gomes MGM. Herd immunity under individual variation and
reinfection. J Math Biol. 2022 Jun 30;85(1):2.

73. Kapczynski DR, King DJ. Protection of chickens against overt clinical disease and
determination of viral shedding following vaccination with commercially available Newcastle
disease virus vaccines upon challenge with highly virulent virus from the California 2002
exotic Newcastle disease outbreak. Vaccine. 2005 May 16;23(26):3424–33.

74. Khoury DS, Cromer D, Reynaldi A, Schlub TE, Wheatley AK, Juno JA, et al. Neutralizing
antibody levels are highly predictive of immune protection from symptomatic SARS-CoV-2
infection. Nat Med. 2021 Jul;27(7):1205–11.

75. Altizer S, Davis AK, Cook KC, Cherry JJ. Age, sex, and season affect the risk of
mycoplasmal conjunctivitis in a southeastern house finch population. Can J Zool. 2004 May
1;82(5):755–63.

76. Hosseini PR, Dhondt AA, Dobson A. Seasonality and wildlife disease: how seasonal birth,
aggregation and variation in immunity affect the dynamics of Mycoplasma gallisepticum in
house finches. Proc Biol Sci. 2004 Dec 22;271(1557):2569–77.

77. Tkachenko AV, Maslov S, Elbanna A, Wong GN, Weiner ZJ, Goldenfeld N. Time-
dependent heterogeneity leads to transient suppression of the COVID-19 epidemic, not
herd immunity. Proc Natl Acad Sci U S A [Internet]. 2021 Apr 27;118(17). Available from:
http://dx.doi.org/10.1073/pnas.2015972118

78. Rose C, Medford AJ, Goldsmith CF, Vegge T, Weitz JS, Peterson AA. Heterogeneity in
susceptibility dictates the order of epidemic models. J Theor Biol. 2021 Nov 7;528:110839.

79. Carlsson R-M, Childs LM, Feng Z, Glasser JW, Heffernan JM, Li J, et al. Modeling the
waning and boosting of immunity from infection or vaccination. J Theor Biol. 2020 Jul
21;497:110265.

80. Scully EP, Haverfield J, Ursin RL, Tannenbaum C, Klein SL. Considering how biological
sex impacts immune responses and COVID-19 outcomes. Nat Rev Immunol. 2020
Jul;20(7):442–7.

81. Charbonnel N, Pagès M, Sironen T, Henttonen H, Vapalahti O, Mustonen J, et al.
Immunogenetic factors affecting susceptibility of humans and rodents to hantaviruses and
the clinical course of hantaviral disease in humans. Viruses. 2014 May 26;6(5):2214–41.

82. Al-Bowardi NC, Riddle KE, Bragin EA. Pyle, P. 1997. Identification guide to North. Available
from: https://sora.unm.edu/sites/default/files/journals/nabb/v027n01/p0014-p0018.pdf

83. Hawley DM, Grodio J, Frasca S, Kirkpatrick L, Ley DH. Experimental infection of domestic
canaries (Serinus canaria domestica) with Mycoplasma gallisepticum: a new model system
for a wildlife disease. Avian Pathol. 2011 Jun;40(3):321–7.

84. Hawley DM, Osnas EE, Dobson AP, Hochachka WM, Ley DH, Dhondt AA. Parallel patterns
of increased virulence in a recently emerged wildlife pathogen. PLoS Biol. 2013 May
28;11(5):e1001570.

85. Bonneaud C, Giraudeau M, Tardy L, Staley M, Hill GE, McGraw KJ. Rapid Antagonistic
Coevolution in an Emerging Pathogen and Its Vertebrate Host. Curr Biol. 2018 Sep
24;28(18):2978-2983.e5.

86. Sydenstricker KV, Dhondt AA, Hawley DM, Jennelle CS, Kollias HW, Kollias GV.
Characterization of experimental Mycoplasma gallisepticum infection in captive house finch
flocks. Avian Dis. 2006 Mar;50(1):39–44.

87. Grodio JL, Dhondt KV, O'Connell PH, Schat KA. Detection and quantification of

Mycoplasma gallisepticum genome load in conjunctival samples of experimentally infected
house finches (Carpodacus mexicanus) using real-time polymerase chain reaction. Avian
Pathol. 2008 Aug;37(4):385–91.

88. Adelman JS, Mayer C, Hawley DM. Infection reduces anti-predator behaviors in house
finches. J Avian Biol. 2017 Apr;48(4):519–28.

89. Williams PD, Dobson AP, Dhondt KV, Hawley DM, Dhondt AA. Evidence of trade-offs
shaping virulence evolution in an emerging wildlife pathogen. J Evol Biol. 2014
Jun;27(6):1271–8.

90. Hochachka WM, Dhondt AA. Density-dependent decline of host abundance resulting from a
new infectious disease. Proc Natl Acad Sci U S A. 2000 May 9;97(10):5303–6.