Prior exposure to pathogens augments host heterogeneity in susceptibility and has key

epidemiological consequences

Dana M. Hawley*[,1], Anna M. Pérez-Umphrey[1], James S. Adelman[2], Arietta E. Fleming-Davies[3],

Jesse Garrett-Larsen[1], Steven J. Geary[4], Lauren M. Childs[$,5], Kate E. Langwig[$,*,1]

[1]Department of Biological Sciences, Virginia Tech, Blacksburg, VA, USA

[2]Department of Biological Sciences, University of Memphis, Memphis, TN, USA

[3]Department of Biology, University of San Diego, San Diego, CA, USA

[4]Department of Pathobiology & Veterinary Science, University of Connecticut, Storrs, CT, USA

[5]Department of Mathematics, Virginia Tech, Blacksburg, VA, USA

*Corresponding authors

Email: hawleyd@vt.edu, klangwig@vt.edu

[$]These authors contributed equally to the work

Abstract. Pathogen epidemics are key threats to human and wildlife health. Across systems, host protection from pathogens following initial exposure is often incomplete, resulting in recurrent epidemics through partially-immune hosts. Variation in population-level protection has important consequences for epidemic dynamics, but whether acquired protection influences host heterogeneity in susceptibility and its epidemiological consequences remains unexplored. We experimentally investigated whether prior exposure (none, low-dose, or high-dose) to a bacterial pathogen alters host heterogeneity in susceptibility among songbirds. Hosts with no prior pathogen exposure had little variation in protection, but heterogeneity in susceptibility was significantly augmented by prior pathogen exposure, with the highest variability detected in hosts given high-dose prior exposure. An epidemiological model parameterized with experimental data found that heterogeneity in susceptibility from prior exposure more than halved epidemic sizes compared with a homogeneous population with identical mean protection. However, because infection-induced mortality was also greatly reduced in hosts with prior pathogen exposure, reductions in epidemic size were smaller than expected in hosts with prior exposure. These results highlight the importance of variable protection from prior exposure and/or vaccination in driving host heterogeneity and epidemiological dynamics.

Author Summary. Individuals in a population can be highly variable in terms of whether or not they get sick during a pathogen outbreak. This individual variability in susceptibility has important consequences for how widely a disease can spread in a population. Therefore, it is key to understand what drives such variability in susceptibility among individuals. One possibility is that variable levels of standing immunity in a population, whether from vaccination or previous infection, lead to variability in susceptibility among individuals. We tested whether acquired immunity creates more variability in susceptibility among individuals in a host population, using a songbird disease system as a model. We found that birds that had acquired immunity to a

51   bacterial pathogen were far more variable in their susceptibility. We also show that this

52   population-level variability in itself can strongly suppress disease outbreaks.

Introduction

Pathogen epidemics are increasing in frequency in humans and other animals (1,2),
underscoring the need to characterize key sources of heterogeneity that influence both
epidemic and evolutionary dynamics for pathogens. There is growing appreciation among
infectious disease epidemiologists and ecologists for both the extent and downstream
importance of individual variation in traits such as contact rates (e.g. 3,4) and host
infectiousness (e.g. 5). In contrast, the heterogeneity present among individual hosts in
pathogen susceptibility (defined here as the probability of infection given exposure), and the
factors that influence the degree of such population-level heterogeneity in susceptibility, has
received relatively less attention (but see (6,7)). Given the documented importance of host
heterogeneity in susceptibility for both epidemiological and evolutionary dynamics (e.g. 7–11), it
is critical to understand how factors such as acquired protection from prior pathogen exposure
influence the degree of among-individual variation present in a given population.

Host reinfection is a common but understudied feature of many host-pathogen systems
(12–17), including SARS-Cov-2 (18), pneumococcal disease (19), malaria (20), and other
diseases of agricultural and wildlife significance. Because the protection acquired from prior
pathogen infection is often incomplete and/or wanes over time, reinfections occur even in
systems where hosts have lower mean susceptibility during secondary exposures, relative to
individuals exposed for the first time (21–24). While there is growing appreciation for the
pervasiveness of heterogeneity in acquired host protection in response to vaccination or
infection across systems (23,25,26), prior work has largely focused on how host protection from
vaccination or prior infection influences mean population traits, rather than variability among
individuals in a given population (27). As such, it remains unknown how host protection acquired
from prior pathogen exposure alters the degree of inter-individual heterogeneity in susceptibility.

Prior pathogen exposure can modify population-level heterogeneity in susceptibility via
several mechanisms. First, prior exposure to pathogens could generate incomplete (or "leaky")

protection similarly for all exposed hosts (23), such that all hosts receive identical protection

from reinfection. Alternatively, prior pathogen exposure could result in relatively complete

acquired protection for some proportion of hosts, while others remain relatively unprotected

(termed "all-or-nothing" immunity (23)). Epidemiological models often select either wholly leaky

protection of hosts that allow every host an identical rate of reinfection or all-or-nothing

protection such that some hosts remain in the recovered class while a subset is routed back to

the susceptible class, representing two extremes of how heterogeneity is incorporated into

disease models (6,27). Whether and how prior exposure to pathogens alters population-level

heterogeneity in susceptibility in a given system likely depends on the extent of prior pathogen

exposure that hosts experience (28) and the mode of action of such protection ["leaky", "all-or-

nothing" (6,27), or some intermediate (26)]. For example, host exposure to low pathogen doses,

which can occur during the natural transmission process for many pathogens (e.g. 29–31),

might be more likely than higher exposure doses to generate incomplete host protection (32,33)

and thus to induce population-level heterogeneity in susceptibility.

Regardless of the underlying mechanisms, effects of prior exposure on population-level

heterogeneity in host susceptibility are particularly important to characterize because this

heterogeneity can have key epidemiological consequences, including lower predicted outbreak

sizes (6) and higher rates of reinfection (10). This is in part due to the process of cohort

selection, through which the most susceptible individuals are infected first in any given epidemic

time step, leaving lower mean susceptibility among remaining uninfected hosts in a population

(34). Experimental quantification of population-level heterogeneity in susceptibility draws on

quantitative microbial risk assessment approaches (35) to estimate susceptibility distributions

using a series of increasing pathogen challenge doses. Because only the most susceptible

hosts become infected at low challenge doses, whereas even the least susceptible hosts

become infected at the highest challenge doses, dose-response models can be used to

estimate the degree of host heterogeneity in susceptibility in a given population (6). Robustly

quantifying changes in host heterogeneity that occur with prior pathogen exposure is

challenging due to large sample size requirements, but nonetheless critical for adequate

characterization of host-pathogen dynamics, including the potential downstream effects of inter-

individual heterogeneity on pathogen strain coexistence and virulence evolution (7,9,36).

Here we investigate effects of host prior exposure on inter-individual heterogeneity in

susceptibility using a tractable wildlife-pathogen system: a common North American songbird

species (house finch; *Haemorhous mexicanus*) and its widespread bacterial pathogen,

Mycoplasma gallisepticum (MG), which causes severe conjunctivitis. The potential effects of

prior pathogen exposure on host heterogeneity are particularly relevant in this system because

MG exposure doses are variable across free-living hosts: susceptible birds contact MG that was

deposited onto bird feeders in variable amounts by infected individuals (37,38). Further,

although the conjunctivitis caused by MG indirectly results in significant mortality in finches,

free-living hosts recover at high rates from initial infection (39), leaving a pool of recovered hosts

in natural populations. Experimental studies show that recovered hosts have significant

acquired protection relative to pathogen-naive hosts, but can still be successfully reinfected at

high rates, even with homologous pathogen strains (28,40). Finally, our work to date suggests

that experimental variation in the degree of prior exposure to MG (in both dose and number of

exposures) results in heterogeneous protection from reinfection among finches, as measured by

conjunctival pathogen loads following reinfection challenge (28,32,33). Whether prior exposure

to MG specifically alters inter-individual heterogeneity in susceptibility, a population-level trait

that requires dose-response approaches to robustly quantify, has not yet been examined.

To test how variation in prior pathogen exposure alters population-level heterogeneity in

susceptibility, we first experimentally varied the degree of prior pathogen exposure categorically

(none, low-dose, or high-dose) in wild caught-finches that we documented had no exposure to

MG prior to capture. After recovery from prior exposure treatments, we measured host

susceptibility (1|0) to secondary dose challenge (Figure 1), fitting dose-response models for

each treatment group to determine whether host prior exposure treatment altered population-

level heterogeneity in susceptibility. Specifically, for each treatment group, we considered

population-level distributions of susceptibility that represent either identical infection probabilities

(homogeneous model) or models that represent inter-host variation in rates of infection

(heterogeneous model, where numerical susceptibility values were estimated according to a

fitted gamma distribution (6,11,41,42)). Importantly, our models allowed us to disentangle

effects of prior exposure on mean susceptibility from changes in heterogeneity in susceptibility.

Lastly, to determine effects of exposure-induced heterogeneity on resulting epidemic dynamics,

we parameterized an SIR model for this system with susceptibility distributions and expected

mortality rates in the wild estimated from our experimental data.

Figure 1. Experimental design for quantifying how prior exposure to *Mycoplasma gallisepticum* alters host heterogeneity in susceptibility in house finches (n=150 for final analysis). Five weeks after one of three prior exposure (PE) treatments (none, low, or high dose; left panel), all birds received a secondary dose challenge (right panel) to assess heterogeneity in susceptibility. Our primary data set was whether birds became infected (0|1) in response to a given secondary dose (right). *However, to improve model fits, we also used bird responses to low-dose PE treatment (left, asterisk), which fell intermediate to our highest secondary challenge doses (300 and 7000 Color Changing Units [CCU]/mL), to quantify the proportion of birds with no prior exposure (at the time of low-dose PE) that become infected at a 750 CCU/mL dose. Made in Biorender.

Results

Prior exposure to pathogens augments population-level heterogeneity in susceptibility.

Birds with no prior pathogen exposure at the time of secondary dose challenge had low

variability in host susceptibility (coef of variation [CoV] from gamma distribution = 0.899) and

there was no support that heterogeneous dose-response models were an improvement over

models assuming homogeneous host susceptibility (defined here as identical infection

probabilities across hosts in a population) (Fig 2a; deviance homogeneous = 1.248; deviance

heterogeneous = 3.307). Birds with no prior pathogen exposure also had high mean

susceptibility to infection (mean susceptibility [fitted gamma distribution]=1.181). In contrast,

dose-response curves were shallower and fewer birds were infected upon secondary exposure

in both the low (mean susceptibility = 0.446 [fitted gamma], CoV [fitted gamma] = 1.630; Fig 2b),

and high-dose (mean susceptibility = 0.192 [fitted gamma], CoV [fitted gamma] = 2.511; Fig 2c)

prior exposure groups. Both prior exposure treatments were better described by models

assuming heterogeneous versus homogeneous susceptibility (Fig 2; likelihood ratio tests

heterogeneous versus homogeneous (35), low dose: P = 0.020, high dose: P = 0.033).

159
Figure 2. Dose response curves for house finch susceptibility to secondary challenge with
Mycoplasma gallisepticum across prior exposure treatments. Points (+/- 1SE) show the fraction
of individual birds (n=10-12 birds for most points; individual responses are 1|0) infected at each
secondary exposure dose, shown as Color Changing Units (CCU)/uL. Lines indicate model fits,
with blue indicating gamma (heterogeneous) model fits, and red dashed lines indicating
homogeneous model fits. Panel labels show prior exposure treatment (birds in the no prior

exposure treatment were pathogen-naive at the time of secondary dose challenge). In hosts
with prior pathogen exposure (low and high-dose prior exposure groups), the gamma model
(which accounts for inter-individual heterogeneity) was better supported via likelihood ratio tests.

Fig 3. Host susceptibility distributions for house finches from variable prior exposure treatments:
no prior exposure (A,B); low-dose (C); or high-dose (D). Colored lines show estimated
susceptibility distributions from either homogeneously (A) or gamma-distributed (B-D) models
(note distinct axes for the two models). In (A), susceptibility (p) is shown as the single best fit
parameter p (dotted vertical lines represent 1 standard error) for the homogeneous model,
which was the best fit model for the no-prior exposure group (see *Results*). In (B-D), the best fit
parameters (shape and scale) for gamma distributions (teal line) are listed for each group, and
vertical gray lines indicate mean susceptibility (x) for that treatment. Lighter shading represents
95% confidence regions for gamma distributions, obtained by bootstrapping chi-squared
residuals to create 1,000 pseudoreplicates of infection data and then refitting the model to
pseudoreplicates, as per (6,35). The gamma model was the best fit for only the low-dose and
high-dose groups. Gamma estimates are also shown for the no prior exposure group (B)
because this allowed more equivalent comparisons for certain SIR simulations (see *Methods*).

To estimate effects of prior exposure on host mortality rates in the wild, we used clinical scores

of eye disease in response to secondary dose challenge to estimate mean mortality rates for

each prior exposure treatment group (as per (22); see Methods). As expected, predicted host

mortality rates decreased with the degree of prior host exposure (Table S2), declining from

0.05894/day in birds with no prior exposure, to 0.02806/day in birds with prior low-dose

exposure, and to a negligible value ($2.483*10^{-17}$) in birds with prior high-dose exposure.

Exposure-induced heterogeneity in susceptibility reduces epidemic size. Using these

empirically-derived susceptibility distributions and predicted mortality rates (Fig 3, Table S2) to

parameterize an SIR model of this system, we find that the total epidemic size is largest

(90.51% of the population) for a host population with no prior pathogen exposure, while for low

and high-dose prior exposure groups, the total epidemic size is significantly reduced (33.67%

and 16.13%) of the population, respectively; blue bars, Fig 4A). Because prior pathogen

exposure also protects hosts from mortality, there is no mortality during the epidemic in the

high-dose prior exposure population, while there is 16.28% mortality in the low-dose prior

exposure population, and more than 59.98% mortality in the naive host population (darker

colored portion of bars, Fig 4A; dashed lines, Fig 4C).

We also examined whether the observed reductions in epidemic size with host prior exposure

are driven primarily by changes in the susceptibility distribution, versus reductions in mean host

susceptibility (blue versus red bars within each prior exposure treatment group, Fig 4A, B). For

empirically determined best fit parameters, epidemic size for a paired simulation assuming

homogeneous rather than heterogeneous susceptibility was larger by 59% and 56%, under low

dose and high dose prior exposure, respectively (low dose, fitted mortality: 49.12 epidemic size

difference [hom-het]; high dose, fitted mortality: 56.55 epidemic size difference [hom-het]). We

simulated the heterogeneous and homogeneous models using the parameter estimates

obtained from bootstrapping chi-squared residuals (see Fig 3). Assuming homogeneous versus

heterogeneous susceptibility (while holding mean susceptibility constant for a given prior

exposure treatment) results in consistently larger outbreaks (Fig S1; difference in epidemic size

= homogeneous [red]-heterogeneous [blue] outbreak size for a given mean susceptibility).

Indeed, across all simulations, outbreaks are never smaller in models assuming homogeneous

versus heterogeneous susceptibility (all P < 0.0001; Fig S1; epidemic size differences [95% CIs

of hom-het epidemic size] for each simulation: low dose, fitted mortality [4.009,49.15]; high

dose, fitted mortality [0.8994,61.29]; low dose, fixed mortality [0.1198,40.77]; high dose, fixed

mortality [0.010,36.18]). Together, these results indicate effects of host heterogeneity *per se* on

outbreak size, which act above and beyond the effects of lower mean susceptibility from prior

pathogen exposure.

Reductions in mean susceptibility with host prior exposure also contribute to smaller outbreak

sizes, but the extent depends on the assumed level of host mortality. When controlling for

mortality differences (i.e., all populations are assumed to have mortality equivalent to the naïve,

no-prior exposure population), the outbreak size of both groups with prior pathogen exposure

(low or high) is reduced (Fig 4B,D). In the case of simulations assuming homogeneous

susceptibility (red bars, Fig 4B) while accounting for differences in mean protection with prior

exposure, this reduction is *entirely* due to lower mean host susceptibility resulting from prior

pathogen exposure. However, when heterogeneity in susceptibility and reduced mean

susceptibility are both accounted for, further reductions in outbreak size and mortality are found

for the low-dose prior exposure group (blue bars and lines, Fig 4B,D). Further reductions from

heterogeneity *per se* are not possible for the high-dose prior exposure group, for which our

model produced an outbreak size of nearly 0 from changes in mean susceptibility alone.

In the more biologically realistic setting (Fig 4A,C) with changes in susceptibility and host

mortality in response to prior exposure both accounted for, we find a more complicated pattern.

If we assume a homogeneous population (red bars) with the mean susceptibility found

empirically for low and high-dose prior exposure groups (and mortality rates set as empirically

determined values), the epidemic sizes drastically increase (82.79% and 72.68% of the

population for the low and high prior exposure, respectively) relative to models accounting for

host heterogeneity in susceptibility (blue bars and lines; Fig 4A,C). The large outbreak in the

high-dose prior exposure population (despite very low mean susceptibility in this group) is due

to the absence of disease-induced mortality in this population, resulting in a higher basic

reproductive number because all individuals survive the entire infectious period.

Fig 4. Cumulative epidemic size (proportion of individuals who ever entered the infectious class) (A,B); dynamics of cumulative infections and deaths (top: C, D; deaths: dashed lines) over time, and numbers of infectious hosts over time (bottom: C, D) in our SIR model, as a function of prior exposure treatment (x-axis for A,B; columns for C,D). Total starting population was 100; darker shading within color represents individuals (within cumulative totals) that did not survive the epidemic. Blue bars and lines use fitted heterogeneous susceptibility (gamma distribution) for all prior exposure groups (Fig 3, Table S2). Red bars and lines assume homogeneous susceptibility for all groups, with mean susceptibility for each group equal to that of the fitted heterogeneous distribution for that treatment group. Left panels (A,C) use fitted mortality rates, whereby estimated mortality rates declined with prior exposure, while right panels (B,D) assume the fixed mortality rate of all groups is equal to that of the no prior exposure birds.

Discussion

Here we use a naturally-occurring host-pathogen system characterized by high rates of reinfection (32) to test how acquired protection from variable degrees of prior pathogen exposure influences host heterogeneity in susceptibility. Using experimental infection data for birds across three distinct prior exposure treatments, we tested for population-level heterogeneity by statistically comparing dose response models that assume either identical susceptibilities among hosts in a population (homogeneous model) or inter-host variation in susceptibility (heterogeneous model). We find that prior exposure to either low or high doses of pathogen significantly augments population-level heterogeneity in susceptibility relative to a pathogen-naive host population, with the extent of population-level heterogeneity increasing with the degree of prior pathogen exposure, from none, low, to high-dose. In addition, our SIR model demonstrates that the observed changes in host susceptibility with prior exposure have key effects on epidemic dynamics. After accounting for estimated effects of prior exposure on host mortality, changes in population-level heterogeneity had stronger effects on outbreak dynamics than reductions in mean host susceptibility, resulting in dramatically lower outbreak sizes.

The past several decades have brought growing recognition that many host-pathogen systems are characterized by reinfection potential, whether shortly following initial infection and recovery, or over longer timescales after any host acquired protection from prior infection has waned (12–15,18,43–46). Despite the ubiquity of highly specific immune memory that allows hosts to respond rapidly to and effectively resist reinfection with the same pathogen, the acquired protection generated by prior pathogen exposure is often incomplete, even in vertebrate taxa with highly specific antigen repertoires and associated immune memory (47). Thus, some degree of reinfection is still possible, despite lower mean susceptibility in hosts with acquired protection (7,21). Given the importance of reinfection in many systems, it is critical to characterize variability in key epidemiological traits such as susceptibility for host populations

made up of individuals with some degree of acquired protection, because this population-level

heterogeneity can strongly influence the likelihood and severity of outbreaks (3,27,48,49), as

well as the evolution of virulence for pathogens (7,9,36,50).

One intuitive prediction is that host acquired protection from prior exposure to pathogens

should homogenize variation in susceptibility in a population by reducing susceptibility similarly

for all individuals. Instead, we found that the protection acquired from prior exposure to a

naturally-occurring bacterial pathogen, whether at low or high doses, significantly augmented

host heterogeneity in susceptibility in house finches. Importantly, while our dose-response

estimates for birds with no prior exposure to MG are well supported by models assuming

homogeneous susceptibility, this does not indicate that there is no biological variation in

susceptibility present within this treatment group. Instead, this result suggests that variation in

susceptibility among house finches without prior exposure is of sufficiently low magnitude, which

is further supported by the low coefficient of variation of the estimated gamma distribution for

this group. Prior studies in this system find variation among both individuals and populations in

other types of host responses to experimental MG infection, including infection severity

(pathogen loads), disease severity (eye scores), and tolerance (per-pathogen disease severity)

(e.g. 51–54), even for birds with no prior exposure to MG infection. Whether host susceptibility

per se tends to be less variable than other types of host responses, both in this system and

others, is an interesting topic for future inquiry, particularly given the key epidemiological

consequences of host susceptibility *per se*.

Interestingly, the only other study to examine effects of acquired protection (from

vaccination rather than active infection) on host heterogeneity in susceptibility (6) found similar

results: vaccination of rainbow trout against a virus augmented host heterogeneity in

susceptibility relative to unvaccinated individuals. Although the protection that hosts acquire

from active infection can be stronger and/or more variable in strength or duration than that

acquired by vaccination (e.g. 55,56), our susceptibility distributions suggest that host protection

generated by prior exposure to live pathogen, whether low or high doses, significantly augments

host heterogeneity in susceptibility, akin to patterns found for vaccination of trout. Further, the

detected pattern of increasing heterogeneity in susceptibility with higher degrees of prior

pathogen exposure, from no, low-dose, to high-dose prior exposure treatments, suggests that

stronger stimulation of host acquired immune responses by high pathogen doses (in this case,

30000 CCU/mL) can, at least in this system, lead to even more variability across individuals in a

population. Intriguingly, we observed highly variable responses to prior exposure despite using

an MG strain (VA1994) isolated from free-living house finches over 25 years ago. Because

significant host evolution has occurred since then (51,54), one might expect less variable host

responses to the strain used here, compared to a more recent MG strain, if host resistance traits

have been favored or even fixed in the host population. Nonetheless, prior exposure to an older,

basal house finch MG strain still induced significantly higher levels of host heterogeneity in

susceptibility, even at low prior exposure doses. Because strain traits such as virulence may

influence the degree of host heterogeneity in a population (7), future work should examine

whether prior exposure to MG strains more virulent and derived than the strain used here

generates even higher degrees of heterogeneity in susceptibility in a host population.

We used an SIR model parameterized empirically to predict how variation in host prior

exposure would influence epidemic dynamics in this system. Consistent with prior work using

SIR (6,10) and dynamic network approaches (57), we show that population-level heterogeneity

in susceptibility (here as a result of prior pathogen exposure) suppresses overall outbreak size

relative to models that assume homogeneous host susceptibility. Importantly, under either set of

mortality assumptions used in our model, outbreak size is always reduced when prior exposure

generates heterogeneity in susceptibility, compared to the homogeneous case that still accounts

for reductions in mean host susceptibility with prior exposure. These results indicate direct

effects of exposure-induced heterogeneity *per se* on outbreak size, effects which act above and

beyond effects of mean susceptibility from prior pathogen exposure on outbreak size. Moreover,

under the more realistic assumption of host protection from mortality after prior pathogen

exposure, the reductions on overall epidemic size in our models appear to be driven

predominantly by increased host heterogeneity in susceptibility.

An earlier SIR model of vaccine-induced heterogeneity also found that augmented host

heterogeneity in susceptibility reduced outbreak size (6), but this model did not incorporate

potential changes in host mortality rate. Reductions in mortality rate in hosts with prior pathogen

exposure, which we estimated were significant in the house finch system (Table S2), are likely

common in hosts with acquired protection, as such protection often reduces host disease and

mortality more strongly than infection risk (e.g. 58,59). In our model results, the degree to which

prior exposure-induced reductions in mean susceptibility drive outbreak size depends on

assumptions of mortality rates in our model. In our model with empirically-derived mortality

rates, infected birds in the prior exposure groups are more likely to survive infection (lighter

color sections of bars; Fig 4A). Intriguingly, these higher survival rates nearly outweigh the

protective effects of reduced mean susceptibility in previously-exposed hosts, resulting in

relatively little reduction in epidemic size in both prior exposure groups (assuming

homogeneous susceptibility for both groups; red bars, Fig 4A), despite significantly lower mean

host susceptibility in birds with prior exposure. In contrast, when mortality rates are held

constant regardless of prior exposure treatment, exposure-induced reductions in mean host

susceptibility alone result in substantially smaller outbreak sizes (red bars; Fig 4B). These

results indicate that effects of prior exposure on mortality rate and susceptibility distributions are

both critical to account for in epidemiological models, as they may interact to drive outbreak

size.

One limitation of our SIR model is that it assumes that epidemics begin in host

populations entirely composed of individuals of one prior exposure type. While such models are

likely realistic representations of host populations at certain times of the year (e.g., epidemics in

late summer that occur in juvenile flocks composed of fully susceptible finches (60,61)), future

models should also consider scenarios whereby populations are composed of some individuals

with no prior exposure, and others with varying degrees of prior pathogen exposure. Developing

a simulated population to such a state would require knowing how the level of susceptibility of

individual birds is altered by prior exposure, for which no data exists. Another limitation is the

choice of an SIR rather than an SIRS model that allows for loss of immunity over time, as is the

case in the house finch-MG model system, in which protection wanes over approximately one

year (40). To disentangle effects of population-level heterogeneity of susceptibility induced by

prior exposure, we focus this study on a single, short timescale epidemic while ignoring

demographic effects and re-infection, similar to previous work (26,62,63). Allowing for

reinfection in our model would require an understanding of how susceptibility and infectivity are

linked, because such correlations have an impact on the progression of epidemics (62,63).

However, such correlations are particularly challenging to quantify in natural host-pathogen

systems. Overall, our work represents an important first step in incorporating one key source

(prior exposure) of individual variation in susceptibility into mathematical models. Although the

importance of individual variation in infection-derived immunity for epidemiological dynamics is

increasingly recognized (23,64), to our knowledge, our study is the first to explicitly consider

how host prior exposure to pathogens influences population-level heterogeneity in susceptibility.

In summary, our results highlight the key importance of variable protection from prior

exposure and/or vaccination in driving host heterogeneity in susceptibility, a population-level

trait with key downstream consequences for epidemiological and evolutionary dynamics

(7,10,48). Because the host protection acquired from a prior infection (or vaccination) is

incomplete and/or wanes over time in diverse vertebrate systems (e.g. 18,46), exposure-

induced heterogeneity in susceptibility may be a common phenomenon for many animal hosts,

including humans. While work to date has largely focused on innate sources of host

heterogeneity in susceptibility [sex (65), genotype (66), etc.], our results show that population-

level heterogeneity in susceptibility can be readily "induced" by prior pathogen exposure, with

key downstream consequences.

Methods

Bird Capture and Housing

We captured 157 mixed-sex, hatch-year house finches (aged by plumage (67)) in

Montgomery County, Virginia using a combination of cage traps and mist nets in June-Aug

2021. Capture and collection were approved by Virginia Department of Game and Inland

Fisheries (066646) and USFWS (MB158404), and all handling and care procedures were

approved by the Virginia Tech (VT) IACUC. Birds were housed in pairs for a two-week minimum

quarantine period in an indoor animal facility, and captured every 3-4 days to assess potential

clinical signs of mycoplasmal conjunctivitis. Any birds with signs of conjunctivitis (see below)

were immediately isolated, along with their cage-mates, and were not used in experiments.

For all birds that never showed clinical signs of conjunctivitis during quarantine, blood

samples were collected 14-18 days post-capture and run for ELISA via a commercial IDEXX kit

(99-06729) using methods previously described (68). Only birds seronegative via ELISA on

days 14-18 post-capture were assigned to MG-exposed treatment groups (Table S1; all birds

except n=3 "prior sham-0 challenge" birds; see Supplement). All birds were housed individually

and then resampled one week before prior exposure treatments for final confirmation of

seronegativity just prior to experiment initiation.

Experimental Design and Timeline

We created variation in prior exposure using three categorical treatments (Figure 1,

Table S1): no prior exposure to MG (inoculation with the same volume of sterile Frey's media as

a "sham" treatment), low-dose prior exposure (dose of 750 Color Changing Units [CCU]/mL of

MG), or high-dose prior exposure (dose of 30,000 CCU/mL of MG). Birds were inoculated with 70

uL of their assigned treatment concentration directly into the conjunctivae via micropipette

droplets, and then allowed to recover (with recovery quantified at day 41 post-infection; see

below). Birds were infected with an MG strain (VA1994 isolate of MG; [7994-1 (6P) 9/17/2018])

that was isolated shortly following the pathogen's emergence in house finches; despite

significant host and pathogen evolution since then (51,69,70), this 1994 strain still results in

pathogen load and eye scores trajectories that are qualitatively similar to later-isolated strains

(VA2013 (51)).

Forty-two days after prior exposure treatment, birds were given a secondary dose

challenge with the same MG strain (again, 70uL directly inoculated into the conjunctivae). Using

a fully factorial design (Table S1), we inoculated birds from each prior exposure treatment with

one of five MG exposure doses to assess heterogeneity in susceptibility (Figure 1). We selected

prior exposure treatments and challenge doses (also termed "secondary doses") most likely to

generate and detect differences in susceptibility based on our published work on conjunctival

loads (33), and we minimized other inherent sources of variation in susceptibility in our wild-

caught birds by using same-age birds, excluding birds with detectable prior exposure at capture,

and randomizing assignments of sex within treatment groups, such that sex ratios were similar

across treatment combinations. Because of the potential for birds to remain infected from prior

exposure treatments until the time of secondary challenge (42 days post-prior exposure), we

included a set of controls (n=10 total birds) that received prior exposure doses of pathogen

(either low or high-dose) and then received a control inoculation of sterile media at day 42.

These 10 birds were all qPCR negative for MG infection by day 41 and during all three

subsequent sample periods post-secondary challenge (see Supplement).

Because our main interest was in how prior exposure treatment alters host susceptibility

(1|0) to secondary dose challenge, we generally do not analyze or present data collected prior

to secondary challenge (day 0), with one exception. To improve our dose response model fits

for the "no prior exposure" treatment group, we included susceptibility data generated in

response to the initial, low-dose prior exposure treatment. We use only the low-dose prior

exposure treatment for this purpose, and not the high-dose prior exposure treatment (30000

CCU / mL) because the low-dose treatment uniquely provided an exposure dose (750 CCU /

mL) that fell intermediate to those used in our secondary challenge doses (which were

otherwise 30, 100, 300, and 7000 CCU/mL), informing a key part of the dose response curve.

Thus, each of the 50 finches given a low-dose prior exposure treatment and then one of five

secondary challenge doses (Table 1) contributed two data points to the analyses: (i) their

susceptibility (1|0) to the "no-prior exposure" dose of 750 CCU/mL, with data collected on days -

42 to -28 prior to secondary challenge; and (ii) their susceptibility (1|0) to one of five secondary

doses, with data collected on days 4-14 post-secondary challenge. All other birds contributed

only one data point to the data-set: susceptibility (1|0) to one of five secondary doses, with data

collected on days 4-14 post-secondary challenge.

Final sample sizes

We began with a balanced design (n=12 birds for all non-0 MG challenge doses in Table

S1; n=157 total birds), but one individual died of unknown causes prior to secondary dose

challenge. A second individual was excluded from all analyses after study completion because,

when ELISA assays on frozen blood samples were completed, that individual was discovered to

have been MG-seropositive just before prior exposure treatments were given. Finally, five

additional individuals were later determined to be qPCR positive one day prior to secondary

challenge (see *Susceptibility and disease sampling*), and thus were eliminated from all analyses

to ensure that our susceptibility data represent true responses to secondary challenge rather

than residual effects of prior exposure treatment. Final sample sizes for all analyses were n=150

individuals (Table S1).

Susceptibility and disease sampling

We quantified whether birds were susceptible to infection or not (Y/N, 1|0) at a given MG exposure dose via qPCR of conjunctival swab samples collected post-exposure. We also visually scored clinical signs of conjunctivitis (all scoring was done blind to a bird's treatment) to use as a proxy for the likelihood of mortality during infection (see below). At each sampling point, we first scored clinical signs of conjunctivitis on a scale of 0 to 3 for left and right conjunctiva separately, as per (71), with 0.5 score intervals used when clinical signs were intermediate between two integer scores. We summed the scores (left plus right conjunctiva) within a given sample day for a total maximum score of 6 for a given bird per day. After scoring, each bird's conjunctiva was swabbed for 5s with a sterile cotton swab dipped in tryptose phosphate broth (TPB). Swabs were swirled in 300uL of sterile TPB and then wrung out into the sample collection tube. Samples from both eyes were pooled within sampling date for a given individual and frozen at -20°C until processing. Genomic DNA was extracted from samples using Qiagen DNeasy 96 Blood and Tissue kits (Qiagen, Valencia, CA) as previously described(69). Extracted genomic DNA from each sample was used to measure overall numbers of MG present in the conjunctiva using a qPCR assay targeting the *mgc2* gene of MG using the primers from (72) and qPCR methods outlined in (69).

Conjunctival swab samples and clinical scores were taken on days 4, 7, 14, and 21 post-secondary-challenge (and on days 7 and 14 post-prior exposure treatments). However, we only use conjunctival swab data from the first three sample points (days 4, 7, or 14 post-challenge) as most relevant for quantifying susceptibility. To account for potential low-level contamination in our qPCR assay (see Supplement) and the possibility that MG loads from prior exposure treatment might still be present in some birds at the time of secondary dose challenge, we considered a bird as susceptible to a given dose if they had a conjunctival MG pathogen load > 50 copies at any of the sample points from day 4 to 14 post-challenge. This cut-off fell above the highest MG load detected among birds that were given a sham secondary challenge treatment

(but had been previously exposed), and thus should not encompass low MG loads that

represent residual loads from prior exposure treatments. We also sampled all birds in the study

on day 41 post-prior exposure, which was one day prior to secondary challenge treatment (day

42). As noted above, we eliminated 5/155 birds deemed infected on day 41 from all analyses

(see Supplement).

Dose response models and parameterization

We used dose-response models to describe the susceptibility of naive and previously

exposed birds challenged with MG(6,35). We fit the models to data by minimizing the deviance

between the data and the model and compared models using likelihood ratio tests (35). We

assumed heterogeneous susceptibility following a gamma distribution (Equation 1) (27). We

compared models of homogeneous susceptibility (best supported for the no prior exposure

group), with models of gamma distributed susceptibility (best supported for the two prior

exposure groups).

Equation 1

$$I_{hom} = 1 - e^{-pd}$$

$$I_{het} = 1 - \int_0^1 e^{-xpd} f(x) dx$$

Here, d (dose) is the exposure concentration in CCU/μl, and I_{hom} or I_{het} is the proportion of

individuals infected. The subscript hom denotes a model assuming homogeneous susceptibility

where het assumes individuals have susceptibility (x) that is distributed according to a gamma

distribution. p is the per bacterial particle concentration rate of host infection. We calculated

coefficients of variation for the gamma models as the standard deviation divided by the mean.

Fitting mortality data

We use clinical scores of conjunctivitis collected post-secondary challenge (with scores

per bird per day ranging from 0 to 6; see above) to estimate mean mortality rates in the wild for

each prior exposure treatment group. We assume that these ordinal scores monotonically

predict host mortality because several lines of evidence indicate that increasingly severe

conjunctivitis and resulting visual and behavioral impairment result in higher mortality risk for

house finches in the wild: 1) the presence of conjunctivitis is linked with higher mortality in wild

birds (39); 2) higher conjunctivitis severity predicts faster capture time in mock-predation events

(i.e., a human capturing the bird by hand) in captivity (73); and 3) indirect mortality from

predators appears to be the primary source of mortality for infected birds in the wild, because

birds in captivity (a predator-free environment) do not ever succumb to infection, even with

highly virulent strains (69). Thus, MG infection *per se* does not directly cause mortality in house

finches, but mortality in the wild (via predation or the inability to find food) is directly linked with

visual impairment and the associated disease that we measure.

Mortality rate was fit using nonlinear least squares regression (nls function in R) as the

nonlinear inverse logit function, $1/(1+\exp(-x))$ scaled by 6, where x is the eye lesion score

(integers between 0 to 6). This fit was then scaled by vmax (0.25; maximum mortality rate

observed in the field from (39)) such that the maximum mortality rate is vmax rather than 1. A

similar approach was used in previous work (22).

Epidemic model

We use a Susceptible-Infectious-Recovered (SIR) model to examine the effect of

heterogeneity in susceptibility on outbreak size and impact. For populations with determined

heterogeneity in susceptibility, we consider a continuous distribution of susceptibility, with

susceptibility, x, distributed according to the parameters of the gamma distribution obtained

from the dose response models.

Equation 2

$$\frac{dS(x)}{dt} = -x\beta IS(x)$$

$$\frac{dI}{dt} = \beta I \int xS(x)dx - \gamma I - \mu I$$

$$\frac{dR}{dt} = \gamma I$$

Here, $S(x)$ is the number of susceptible individuals with susceptibility x, and x is a value

between 0 and 10. I is the number of infectious individuals, and R is the number of recovered

individuals. Parameters for infectivity (β) and recovery (γ) were taken from a prior empirically-

parameterized model in this system (74) (Table S2), while disease-induced mortality (μ) was fit

as described above (See *Fitting mortality data*) for each prior exposure group (none, low, high).

Infectivity and recovery rates do not depend on prior exposure, while mortality rate does (in the

"fitted mortality" model). A density-dependent rather than frequency-dependent transmission

term was used due to the nature of transmission in this system, which increases with density

(61,75).

For simulation, we discretize the susceptibility distribution of the heterogeneous model into 300

evenly spaced susceptibility classes, represented by the midpoint susceptibility, x, for each

class. We assume a population size of 100 and seed infections with a single infected individual,

and the initial population in each susceptibility class is determined empirically from the fitted

distributions by prior exposure treatment (See *Dose response models and parameterization*).

We quantify the total size of the epidemic as the cumulative fraction of infected individuals, and

simulate for 500 days, after which nearly all epidemics have completely resolved. Because we

simulate a short-time scale single epidemic, we ignore demographic effects and assume

recovered individuals cannot be reinfected, despite documented waning of immunity to MG over

the course of a year (40).

To disentangle the effects of heterogeneity in susceptibility and changes in mean susceptibility,

we simulate outbreaks using the homogeneous model with identical mean susceptibility to the

discretized heterogeneous distribution. To generate confidence intervals on outbreak sizes, we

used the parameter estimates obtained from bootstrapping the chi-squared residuals of the

dose-response estimates. We then estimated the probability of obtaining an epidemic size

difference as extreme as observed (hom-het) using the best fit parameters, but with the

heterogeneous case resulting in larger epidemics (het-hom).

Data Availability Statement

The datasets generated for this study are available in the GitHub repository, at

https://github.com/klangwig/mg_dose_response_public. Data and file code will be archived on

Dryad upon manuscript acceptance.

Acknowledgments

We thank Edan Tulman for input on experimental design and MG isolate preparation, Chava

Weitzman, John Brule, and many other Hawley lab members for help with capture and

sampling, and Bambi Kelly for assistance with animal care. Research reported in this publication

was supported by the National Institute Of General Medical Sciences of the National Institutes

of Health under Award Number R01GM144972. The content is solely the responsibility of the

authors and does not necessarily represent the official views of the National Institutes of Health.

Author Contributions

D.M.H.: Conceptualization, Methodology, Resources, Investigation, Writing- Original Draft,

Supervision, Project administration, Funding acquisition

A.P.U.: Investigation, Data Curation, Writing - Review & Editing

J.S.A: Conceptualization, Methodology, Writing-Review and Editing, Funding acquisition

AEFD: Conceptualization, Methodology, Software, Formal Analysis, Writing - Review & Editing,

Funding acquisition

J.N.GL: Investigation, Writing-Review and Editing

S.J.G.: Conceptualization, Methodology, Writing-Review and Editing

L.M.C.: Conceptualization, Methodology, Software, Validation, Formal Analysis, Writing-Review

and Editing, Visualization, Funding acquisition

K.E.L.: Conceptualization, Methodology, Software, Validation, Formal Analysis, Writing-Review

and Editing, Visualization, Funding acquisition

594
595
596

References Cited

1. Smith KF, Goldberg M, Rosenthal S, Carlson L, Chen J, Chen C, et al. Global rise in human infectious disease outbreaks. J R Soc Interface. 2014 Dec 6;11(101):20140950.

2. Schmeller DS, Courchamp F, Killeen G. Biodiversity loss, emerging pathogens and human health risks. Biodivers Conserv. 2020 Aug 13;29(11–12):3095–102.

3. Lloyd-Smith JO, Schreiber SJ, Kopp PE, Getz WM. Superspreading and the effect of individual variation on disease emergence. Nature. 2005 Nov 17;438(7066):355–9.

4. Britton T, Ball F, Trapman P. A mathematical model reveals the influence of population heterogeneity on herd immunity to SARS-CoV-2. Science. 2020 Aug 14;369(6505):846–9.

5. Ke R, Martinez PP, Smith RL, Gibson LL, Mirza A, Conte M, et al. Daily longitudinal sampling of SARS-CoV-2 infection reveals substantial heterogeneity in infectiousness. Nat Microbiol. 2022 May;7(5):640–52.

6. Langwig KE, Wargo AR, Jones DR, Viss JR, Rutan BJ, Egan NA, et al. Vaccine Effects on Heterogeneity in Susceptibility and Implications for Population Health Management. MBio [Internet]. 2017 Nov 21;8(6). Available from: http://dx.doi.org/10.1128/mBio.00796-17

7. Fleming-Davies AE, Dukic V, Andreasen V, Dwyer G. Effects of host heterogeneity on pathogen diversity and evolution. Ecol Lett. 2015 Nov;18(11):1252–61.

8. Hickson RI, Roberts MG. How population heterogeneity in susceptibility and infectivity influences epidemic dynamics. J Theor Biol. 2014 Jun 7;350:70–80.

9. Pugliese A. The role of host population heterogeneity in the evolution of virulence. J Biol Dyn. 2011 Mar;5(2):104–19.

10. Rodrigues P, Margheri A, Rebelo C, Gomes MGM. Heterogeneity in susceptibility to infection can explain high reinfection rates. J Theor Biol. 2009 Jul 21;259(2):280–90.

11. Dwyer G, Elkinton JS, Buonaccorsi JP. Host heterogeneity in susceptibility and disease dynamics: tests of a mathematical model. Am Nat. 1997 Dec;150(6):685–707.

12. Yamamoto T, Nagasawa I, Nojima M, Yoshida K, Kuwabara Y. Sexual transmission and reinfection of group B streptococci between spouses. J Obstet Gynaecol Res. 1999 Jun;25(3):215–9.

13. Nardell E, McInnis B, Thomas B, Weidhaas S. Exogenous reinfection with tuberculosis in a shelter for the homeless. N Engl J Med. 1986 Dec 18;315(25):1570–5.

14. Islam N, Krajden M, Shoveller J, Gustafson P, Gilbert M, Buxton JA, et al. Incidence, risk factors, and prevention of hepatitis C reinfection: a population-based cohort study. Lancet Gastroenterol Hepatol. 2017 Mar;2(3):200–10.

15. Versteegh FGA, Schellekens JFP, Nagelkerke AF, Roord JJ. Laboratory-confirmed reinfections with Bordetella pertussis. Acta Paediatr. 2007 Jan 2;91(1):95–7.

16. Bazan JA, Tzeng Y-L, Stephens DS, Carter AM, Brown MA, Snyder B, et al. Repeat Episodes of Symptomatic Urethritis Due to a Uropathogenic Meningococcal Clade. Sex

Transm Dis. 2020 Jan;47(1):e1–4.

17. Barenkamp SJ, Shurin PA, Marchant CD, Karasic RB, Pelton SI, Howie VM, et al. Do
children with recurrent Haemophilus influenzae otitis media become infected with a new
organism or reacquire the original strain? J Pediatr. 1984 Oct;105(4):533–7.

18. Pilz S, Theiler-Schwetz V, Trummer C, Krause R, Ioannidis JPA. SARS-CoV-2 reinfections:
Overview of efficacy and duration of natural and hybrid immunity. Environ Res. 2022
Jun;209:112911.

19. Sanz JC, Rodríguez-Avial I, Ríos E, Ramos B, Cercenado E. Recurrent pneumococcal
invasive disease in the region of Madrid during a five-year period. Infection. 2014
Jun;42(3):475–83.

20. Ranawaka MB, Munesinghe YD, de Silva DM, Carter R, Mendis KN. Boosting of
transmission-blocking immunity during natural Plasmodium vivax infections in humans
depends upon frequent reinfection. Infect Immun. 1988 Jul;56(7):1820–4.

21. Forshey BM, Reiner RC, Olkowski S, Morrison AC, Espinoza A, Long KC, et al. Incomplete
Protection against Dengue Virus Type 2 Re-infection in Peru. PLoS Negl Trop Dis. 2016
Feb;10(2):e0004398.

22. Fleming-Davies AE, Williams PD, Dhondt AA, Dobson AP, Hochachka WM, Leon AE, et al.
Incomplete host immunity favors the evolution of virulence in an emergent pathogen.
Science. 2018 Mar 2;359(6379):1030–3.

23. Le A, King AA, Magpantay FMG, Mesbahi A, Rohani P. The impact of infection-derived
immunity on disease dynamics. J Math Biol. 2021 Nov 12;83(6–7):61.

24. Altarawneh HN, Chemaitelly H, Hasan MR, Ayoub HH, Qassim S, AlMukdad S, et al.
Protection against the Omicron Variant from Previous SARS-CoV-2 Infection. N Engl J
Med. 2022 Mar 31;386(13):1288–90.

25. Gomes MGM, White LJ, Medley GF. Infection, reinfection, and vaccination under
suboptimal immune protection: epidemiological perspectives. J Theor Biol. 2004 Jun
21;228(4):539–49.

26. Zachreson C, Tobin R, Szanyi J, Walker C, Cromer D, Shearer FM, et al. Individual
variation in vaccine immune response can produce bimodal distributions of protection.
Vaccine. 2023 Oct 26;41(45):6630–6.

27. Gomes MGM, Lipsitch M, Wargo AR, Kurath G, Rebelo C, Medley GF, et al. A missing
dimension in measures of vaccination impacts. PLoS Pathog. 2014 Mar;10(3):e1003849.

28. Leon AE, Hawley DM. Host Responses to Pathogen Priming in a Natural Songbird Host.
Ecohealth. 2017 Dec;14(4):793–804.

29. Aiello CM, Nussear KE, Esque TC, Emblidge PG, Sah P, Bansal S, et al. Host contact and
shedding patterns clarify variation in pathogen exposure and transmission in threatened
tortoise Gopherus agassizii: implications for disease modelling and management. J Anim
Ecol. 2016;85(3):829–42.

30. Konrad M, Vyleta ML, Theis FJ, Stock M, Tragust S, Klatt M, et al. Social transfer of

pathogenic fungus promotes active immunisation in ant colonies. PLoS Biol. 2012 Apr
3;10(4):e1001300.

31. Stroeymeyt N, Grasse AV, Crespi A, Mersch DP, Cremer S, Keller L. Social network
plasticity decreases disease transmission in a eusocial insect. Science. 2018 Nov
23;362(6417):941–5.

32. Weitzman CL, Ceja G, Leon AE, Hawley DM. Protection Generated by Prior Exposure to
Pathogens Depends on both Priming and Challenge Dose [Internet]. Vol. 90, Infection and
Immunity. 2022. Available from: http://dx.doi.org/10.1128/iai.00537-21

33. Leon AE, Fleming-Davies AE, Hawley DM. Host exposure history modulates the within-host
advantage of virulence in a songbird-bacterium system. Sci Rep. 2019 Dec 30;9(1):20348.

34. Aalen OO, Valberg M, Grotmol T, Tretli S. Understanding variation in disease risk: the
elusive concept of frailty. Int J Epidemiol. 2015 Aug;44(4):1408–21.

35. Haas CN, Rose JB, Gerba CP. Quantitative Microbial Risk Assessment. John Wiley &
Sons; 1999. 449 p.

36. Miller IF, Metcalf CJ. Vaccine-driven virulence evolution: consequences of unbalanced
reductions in mortality and transmission and implications for pertussis vaccines. J R Soc
Interface. 2019 Dec;16(161):20190642.

37. Adelman JS, Carter AW, Hopkins WA, Hawley DM. Deposition of pathogenic Mycoplasma
gallisepticum onto bird feeders: host pathology is more important than temperature-driven
increases in food intake. Biol Lett. 2013 Oct 23;9(5):20130594.

38. Dhondt AA, Dhondt KV, Hawley DM, Jennelle CS. Experimental evidence for transmission
of Mycoplasma gallisepticum in house finches by fomites. Avian Pathol. 2007
Jun;36(3):205–8.

39. Faustino CR, Jennelle CS, Connolly V, Davis AK, Swarthout EC, Dhondt AA, et al.
Mycoplasma gallisepticum infection dynamics in a house finch population: seasonal
variation in survival, encounter and transmission rate. J Anim Ecol. 2004 Jul;73(4):651–69.

40. Sydenstricker KV, Dhondt AA, Ley DH, Kollias GV. Re-exposure of captive house finches
that recovered from Mycoplasma gallisepticum infection. J Wildl Dis. 2005 Apr;41(2):326–
33.

41. Ben-Ami F, Ebert D, Regoes RR. Pathogen dose infectivity curves as a method to analyze
the distribution of host susceptibility: a quantitative assessment of maternal effects after
food stress and pathogen exposure. Am Nat. 2010 Jan;175(1):106–15.

42. Ben-Ami F, Regoes RR, Ebert D. A quantitative test of the relationship between parasite
dose and infection probability across different host-parasite combinations. Proc Biol Sci.
2008 Apr 7;275(1636):853–9.

43. Koopman JS, Henry CJ, Park JH, Eisenberg MC, Ionides EL, Eisenberg JN. Dynamics
affecting the risk of silent circulation when oral polio vaccination is stopped. Epidemics.
2017 Sep;20:21–36.

44. van den Hoogen BG, Herfst S, de Graaf M, Sprong L, van Lavieren R, van Amerongen G,

et al. Experimental infection of macaques with human metapneumovirus induces transient
protective immunity. J Gen Virol. 2007 Apr;88(Pt 4):1251–9.

45. Singleton R, Etchart N, Hou S, Hyland L. Inability to evoke a long-lasting protective immune
response to respiratory syncytial virus infection in mice correlates with ineffective nasal
antibody responses. J Virol. 2003 Nov;77(21):11303–11.

46. Edridge AWD, Kaczorowska J, Hoste ACR, Bakker M, Klein M, Loens K, et al. Seasonal
coronavirus protective immunity is short-lasting. Nat Med. 2020 Nov;26(11):1691–3.

47. Boehm T. Evolution of vertebrate immunity. Curr Biol. 2012 Sep 11;22(17):R722-32.

48. Gomes MGM, Aguas R, Lopes JS, Nunes MC, Rebelo C, Rodrigues P, et al. How host
heterogeneity governs tuberculosis reinfection? Proc Biol Sci. 2012 Jun
22;279(1737):2473–8.

49. Ferrari MJ, Bansal S, Meyers LA, Bjørnstad ON. Network frailty and the geometry of herd
immunity. Proc Biol Sci. 2006 Nov 7;273(1602):2743–8.

50. Leggett HC, Buckling A, Long GH, Boots M. Generalism and the evolution of parasite
virulence. Trends Ecol Evol. 2013 Oct;28(10):592–6.

51. Henschen AE, Vinkler M, Langager MM, Rowley AA, Dalloul RA, Hawley DM, et al. Rapid
adaptation to a novel pathogen through disease tolerance in a wild songbird. PLoS Pathog.
2023 Jun;19(6):e1011408.

52. Ruden RM, Adelman JS. Disease tolerance alters host competence in a wild songbird. Biol
Lett. 2021 Oct;17(10):20210362.

53. Adelman JS, Kirkpatrick L, Grodio JL, Hawley DM. House Finch Populations Differ in Early
Inflammatory Signaling and Pathogen Tolerance at the Peak ofMycoplasma
gallisepticumInfection [Internet]. Vol. 181, The American Naturalist. 2013. p. 674–89.
Available from: http://dx.doi.org/10.1086/670024

54. Bonneaud C, Tardy L, Giraudeau M, Hill GE, McGraw KJ, Wilson AJ. Evolution of both host
resistance and tolerance to an emerging bacterial pathogen. Evol Lett. 2019 Oct;3(5):544–
54.

55. Wendelboe AM, Van Rie A, Salmaso S, Englund JA. Duration of immunity against pertussis
after natural infection or vaccination. Pediatr Infect Dis J. 2005 May;24(5 Suppl):S58-61.

56. Hobson D. Resistance to reinfection in experimental mouse typhoid. J Hyg . 1957
Sep;55(3):334–43.

57. White LA, Forester JD, Craft ME. Covariation between the physiological and behavioral
components of pathogen transmission: host heterogeneity determines epidemic outcomes.
Oikos. 2018 Apr;127(4):538–52.

58. Kapczynski DR, King DJ. Protection of chickens against overt clinical disease and
determination of viral shedding following vaccination with commercially available Newcastle
disease virus vaccines upon challenge with highly virulent virus from the California 2002
exotic Newcastle disease outbreak. Vaccine. 2005 May 16;23(26):3424–33.

59. Khoury DS, Cromer D, Reynaldi A, Schlub TE, Wheatley AK, Juno JA, et al. Neutralizing
antibody levels are highly predictive of immune protection from symptomatic SARS-CoV-2
infection. Nat Med. 2021 Jul;27(7):1205–11.

60. Altizer S, Davis AK, Cook KC, Cherry JJ. Age, sex, and season affect the risk of
mycoplasmal conjunctivitis in a southeastern house finch population. Can J Zool. 2004 May
1;82(5):755–63.

61. Hosseini PR, Dhondt AA, Dobson A. Seasonality and wildlife disease: how seasonal birth,
aggregation and variation in immunity affect the dynamics of Mycoplasma gallisepticum in
house finches. Proc Biol Sci. 2004 Dec 22;271(1557):2569–77.

62. Tkachenko AV, Maslov S, Elbanna A, Wong GN, Weiner ZJ, Goldenfeld N. Time-
dependent heterogeneity leads to transient suppression of the COVID-19 epidemic, not
herd immunity. Proc Natl Acad Sci U S A [Internet]. 2021 Apr 27;118(17). Available from:
http://dx.doi.org/10.1073/pnas.2015972118

63. Rose C, Medford AJ, Goldsmith CF, Vegge T, Weitz JS, Peterson AA. Heterogeneity in
susceptibility dictates the order of epidemic models. J Theor Biol. 2021 Nov 7;528:110839.

64. Carlsson R-M, Childs LM, Feng Z, Glasser JW, Heffernan JM, Li J, et al. Modeling the
waning and boosting of immunity from infection or vaccination. J Theor Biol. 2020 Jul
21;497:110265.

65. Scully EP, Haverfield J, Ursin RL, Tannenbaum C, Klein SL. Considering how biological
sex impacts immune responses and COVID-19 outcomes. Nat Rev Immunol. 2020
Jul;20(7):442–7.

66. Charbonnel N, Pagès M, Sironen T, Henttonen H, Vapalahti O, Mustonen J, et al.
Immunogenetic factors affecting susceptibility of humans and rodents to hantaviruses and
the clinical course of hantaviral disease in humans. Viruses. 2014 May 26;6(5):2214–41.

67. Al-Bowardi NC, Riddle KE, Bragin EA. Pyle, P. 1997. Identification guide to North. Available
from: https://sora.unm.edu/sites/default/files/journals/nabb/v027n01/p0014-p0018.pdf

68. Hawley DM, Grodio J, Frasca S, Kirkpatrick L, Ley DH. Experimental infection of domestic
canaries (Serinus canaria domestica) with Mycoplasma gallisepticum: a new model system
for a wildlife disease. Avian Pathol. 2011 Jun;40(3):321–7.

69. Hawley DM, Osnas EE, Dobson AP, Hochachka WM, Ley DH, Dhondt AA. Parallel patterns
of increased virulence in a recently emerged wildlife pathogen. PLoS Biol. 2013 May
28;11(5):e1001570.

70. Bonneaud C, Giraudeau M, Tardy L, Staley M, Hill GE, McGraw KJ. Rapid Antagonistic
Coevolution in an Emerging Pathogen and Its Vertebrate Host. Curr Biol. 2018 Sep
24;28(18):2978-2983.e5.

71. Sydenstricker KV, Dhondt AA, Hawley DM, Jennelle CS, Kollias HW, Kollias GV.
Characterization of experimental Mycoplasma gallisepticum infection in captive house finch
flocks. Avian Dis. 2006 Mar;50(1):39–44.

72. Grodio JL, Dhondt KV, O'Connell PH, Schat KA. Detection and quantification of

Mycoplasma gallisepticum genome load in conjunctival samples of experimentally infected
house finches (Carpodacus mexicanus) using real-time polymerase chain reaction. Avian
Pathol. 2008 Aug;37(4):385–91.

73. Adelman JS, Mayer C, Hawley DM. Infection reduces anti-predator behaviors in house
finches. J Avian Biol. 2017 Apr;48(4):519–28.

74. Williams PD, Dobson AP, Dhondt KV, Hawley DM, Dhondt AA. Evidence of trade-offs
shaping virulence evolution in an emerging wildlife pathogen. J Evol Biol. 2014
Jun;27(6):1271–8.

75. Hochachka WM, Dhondt AA. Density-dependent decline of host abundance resulting from a
new infectious disease. Proc Natl Acad Sci U S A. 2000 May 9;97(10):5303–6.

800
801

 Figure Captions

Figure 1. Experimental design for quantifying how prior exposure to *Mycoplasma gallisepticum*

alters host heterogeneity in susceptibility in house finches (n=150 for final analysis). Five weeks

after one of three prior exposure (PE) treatments (none, low, or high dose; left panel), all birds

received a secondary dose challenge (right panel) to assess heterogeneity in susceptibility. Our

primary data set was whether birds became infected (0|1) in response to a given secondary

dose (right). *However, to improve model fits, we also used bird responses to low-dose PE

treatment (left, asterisk), which fell intermediate to our highest secondary challenge doses (300

and 7000 Color Changing Units [CCU]/mL), to quantify the proportion of birds with no prior

exposure (at the time of low-dose PE) that become infected at a 750 CCU/mL dose. Made in

Biorender.

Figure 2. Dose response curves for house finch susceptibility to secondary challenge with

Mycoplasma gallisepticum across prior exposure treatments. Points (+/- 1SE) show the fraction

of individual birds (n=10-12 birds for most points; individual responses are 1|0) infected at each

secondary exposure dose, shown as Color Changing Units (CCU)/uL. Lines indicate model fits,

with blue indicating gamma (heterogeneous) model fits, and red dashed lines indicating

homogeneous model fits. Panel labels show prior exposure treatment (birds in the no prior

exposure treatment were pathogen-naive at the time of secondary dose challenge). In hosts

with prior pathogen exposure (low and high-dose prior exposure groups), the gamma model

(which accounts for inter-individual heterogeneity) was better supported via likelihood ratio tests.

Figure 3. Host susceptibility distributions for house finches from variable prior exposure

treatments: no prior exposure (A,B); low-dose (C); or high-dose (D). Colored lines show

estimated susceptibility distributions from either homogeneously (A) or gamma-distributed (B-D)

models (note distinct axes for the two models). In (A), susceptibility (p) is shown as the single

best fit parameter p (dotted vertical lines represent 1 standard error) for the homogeneous

model, which was the best fit model for the no-prior exposure group (see *Results*). In (B-D), the

best fit parameters (shape and scale) for gamma distributions (teal line) are listed for each

group, and vertical gray lines indicate mean susceptibility (x) for that treatment. Lighter shading

represents 95% confidence regions for gamma distributions, obtained by bootstrapping chi-

squared residuals to create 1,000 pseudoreplicates of infection data and then refitting the model

to pseudoreplicates, as per (6,35). The gamma model was the best fit for only the low-dose and

high-dose groups. Gamma estimates are also shown for the no prior exposure group (B)

because this allowed more equivalent comparisons for certain SIR simulations (see *Methods*).

Figure 4. Cumulative epidemic size (proportion of individuals who ever entered the infectious

class) (A,B); dynamics of cumulative infections and deaths (top: C, D; deaths: dashed lines)

over time, and numbers of infectious hosts over time (bottom: C, D) in our SIR model, as a

function of prior exposure treatment (x-axis for A,B; columns for C,D). Total starting population

was 100; darker shading within color represents individuals (within cumulative totals) that did

not survive the epidemic. Blue bars and lines use fitted heterogeneous susceptibility (gamma

distribution) for all prior exposure groups (Fig 3, Table S2). Red bars and lines assume

homogeneous susceptibility with mean susceptibility equal to that of the fitted heterogeneous

distribution. Left panels (A,C) use fitted mortality rates, whereby estimated mortality rates

declined with prior exposure, while right panels (B,D) assume the fixed mortality rate of all

groups is equal to that of the no prior exposure birds.

22