Light as an "ecological trap". A synanthropic incipient speciation taken up by the gecko

Domenico Fulgione[1*], Valeria Maselli[1], Bice Avallone[1], Eleonora Rivieccio[2], Alessandro Mondanaro[3],

Giorgio Giurato[4] and Maria Buglione[1]

1 - Department of Biology, University of Naples Federico II, Naples, Italy, 80126

2 - Department of Humanities Studies, University of Naples Federico II, Naples, Italy, 80138

3 - Department of Earth Sciences, University of Florence, 50121

4 - Department of Medicine, Surgery and Dentistry Medicine and Surgery Campus Laboratory of

Molecular Medicine and Genomics, University of Salerno, Salerno, Italy, 84081

*Correspondence to Domenico Fulgione

Department of Biology, University of Naples Federico II, via Cinthia 26, 80126 Naples, Italy

Phone: 0039 081769130

Email: fulgione@unina.it

Running title: Anthropogenic speciation by light

Abstract

Two distinct and sympatric populations of Moorish gecko (*Tarentola mauritanica*) can be easily

distinguished: the diurnal-dark gecko, which mainly lives on trees, and the nocturnal-pale gecko,

recognizable on anthropic buildings' walls, especially nearby artificial lights, in order to increase the

chances to get preys.

Here we hypothesize that this evolutionary splitting was determined by the choice of the

anthropogenic niche occurring in a population of Moorish gecko back in the time when *Homo*

sapiens used fire in its settlements, luring insects to positive phototaxis. The frequentation of the

new trophic niche created by human through light has increasingly strengthened the distinctive

traits in the nocturnal population of geckos, driving sympatric speciation by selection for background

matching.

We provide empirical evidence supporting this hypothesis, suggesting that the nocturnal-pale

geckos split from the diurnal-dark populations due to the innate plastic variation determined by their

ability to adjust their skin color and later reinforcement of depigmentation.

Our study inaugurates an interesting line of investigation that could shed light on the impact that

humankind has on the evolutionary trajectories of non-domesticated species, endowed with

adaptive plasticity, functional to the exploration and colonization of the anthropic niche.

Keywords

Homo sapiens, Fire Era, Moorish gecko, synanthropism

Introduction

The history of *Homo sapiens* is studded with many events which promoted relationships with wild animals in different ways (Ålund et al. 2023; Hulme-Beaman et al. 2016), changing their evolutionary path or, somehow, impacting their adaptation (Van Bocxlaer et al. 2010; Puckett et al. 2020).

Whereas these "domestication" events are generally recognized as the product of planned actions, aimed to annex useful plant and animal species into the anthropic niche, some organisms established synanthropic relationships beyond the human intention. These latter occurrences are noticeable in many evolutionary processes as the self-domestication developed in mice and rats (Pocock et al. 2004), sparrows (Ravinet et al. 2018), cockroaches (Booth et al. 2011), foxes (Parsons et al. 2020) , and elephants (Raviv et al. 2023).

Humans' connotation as "niche builders" (Kendal et al. 2011), referring to the unintentional exploitation of other species, is not limited to the management of food, but it's also conceivable as an "ecological trap" (Schlaepfer et al. 2002; Hale and Swearer 2016).

Animals may find opportunities to make dens, shelters, and microclimatic buffering (Kamp et al. 2009) within human environments. Artificial light is present mostly in urban agglomeration, but it irremediably spreads to natural ecosystems, this could lead to adaptive changes that influence temporal niches and trophic interactions in different groups of animals (Jägerbrand & Spoelstra, 2023). Several studies documented the effect of artificial light at night on animals (Wang et al. 2021; Baker and Richardson 2006; Beier 2006; Russo et al. 2017), including Moorish gecko, *Tarentola mauritanica* (Martín et al. 2018), mainly focusing on physiological and behavioral conditioning.

Herein, we report data corroborating the hypothesis of an evolutionary splitting determined by synanthropism, occurring in the Moorish gecko, dating back to the time when *Homo sapiens* used fire in its settlements.

The Moorish gecko is a small saurian widespread in North Africa and Europe. This species has the widest range in the Mediterranean Basin compared to all geckos' species (Rieppel 1981). However, because of the close relationship with humans, the gecko has been accidentally introduced in the Balearics (Spain), Madeira island (Portugal), some Balkan islands, Crete, and South America (Carranza et al. 2000; IUCN 2017) (Fig. 1). European populations of Moorish gecko diverged during the early Pliocene, around 4.14 Mya. In fact, the colonization of the Iberian Peninsula by Moorish gecko from North Africa could have been during the Messinian Salinity Crisis, or after the opening of the Strait of Gibraltar during the early Pliocene (Rato et al. 2012).

Figure 1 – **Moorish gecko distribution range.** Native (red) and introduction (yellow) range distribution of Moorish gecko (*Tarentola mauritanica*). The map was by © Rune Midtgaard, "RepFocus - A survey of Reptiles of the World" (https://repfocus.dk). The blue spot is referred to the study site.

The Moorish gecko is active during diurnal and nocturnal hours, occupying a variety of ecological niches, such as tree trunks, houses, and stone walls (Fulgione et al. 2019; Rato et al. 2023; Valverde

2004). This species represents an interesting example of plastic variation, highlighted by the ability

to accommodate its skin color according to the substrate (Vroonen et al. 2012) (Fig. 2).

Figure 2 – Skin color change. Variation in skin colouration of Moorish gecko placed on light (a) and

dark (b) substrate. The color change occurred in 1.5 hours.

Two distinct and sympatric populations can be easily distinguished (Fulgione et al. 2019): the -dark

diurnal-gecko which mainly lives on trees, and the pale-nocturnal gecko, recognizable on anthropic

buildings' walls, especially nearby artificial lights, in order to increase the chances to get preys

(Fulgione et al. 2019; Arnold 2002; Rato and Carretero 2015) (Fig. 3).

Figure 3 – Geckos' morphotypes. Dark-diurnal (A) and pale-nocturnal (B) Moorish geckos (*Tarentola*

mauritanica). They occupy two distinct ecological niches: logs during the day and the walls of houses

at night, respectively.

We hypothesized that the pale gecko, active during the night, may have experienced a prolonged

evolution alongside humans, rooted in the deep past, when *H. sapiens* populations occupied the

first caves and used fire. Since then, the nocturnal gecko has accompanied our species and the

evolution of housing structures.

We provide empirical evidence supporting this hypothesis, suggesting that the nocturnal-pale

geckos split from the diurnal-dark populations due to the innate plastic variation determined by their

ability to adjust their skin color (Vroonen et al. 2012; Fulgione et al. 2019, 2014) and later

reinforcement of depigmentation. This attitude could have been useful when humans added a new

niche dimension to the natural environment, lighting up fires on purpose to get illumination and

heating.

Humans have been using fire for hundreds of thousands of years: the intentional use of fire traces back to the Middle Pleistocene of the Levant (Zohar et al. 2022; Stepka et al. 2022) but understandably spread as a common practice just afterward (Shimelmitz et al. 2014).

Whereas earliest fire usage may suggest cooking and anti-predatory activities, the *daylight extension*, typical of the human social niche, may have appeared later, and could have been linked to the development of language during the latest Pleistocene (Gowlett 2016).

The *daylight extension* hypothesis is consistent with the observation that humans have developed an activity peak during late evening hours, an unusual condition comparing to the mammalian standard (Bowman et al. 2011).

This phenomenon may have unintentionally provided the Moorish gecko with abundant insect preys, especially among insects species characterized by positive phototaxis (i.e. moths, grasshoppers, and flies) (Park and Lee 2017).

This extraordinary new food bonanza could be exploited only using crypticism, a specialization that prevents the geckos from being preys in their turn of nocturnal predators, which is led by their skin-color variation ability (Vroonen et al. 2012). Moreover, geckos with adhesive toepads can easily scale vertical or even inverted surfaces (Gamble et al. 2012), and these extraordinary clinging abilities may have facilitated the use of smooth walls inside the caves.

To test this hypothesis, we have studied two sympatric populations of geckos, a dark-diurnal and a pale-nocturnal one. First, we investigated the two populations' phylogenetic history to date their split age. Then, we produced habitat suitability models for the gecko, and we tested them for a positive association with the human fossil sites dating back to the split age. Molecular dating of the phyletic separation between diurnal and nocturnal geckos, together with the spatial correspondence between geckos and human populations, could prove the beginning of the phenomenon. Moreover, a palaeoecological experiment was performed to demonstrate the effective attraction of Arthropods

on wall lighted with fire. The subsequent reinforcement, through the depigmentation effect observed in pale-nocturnal geckos, was shown using histology of melanosomes and quantified the markers of melanization.

Materials and Methods

Sampling and study area

The study was carried out in South Italy (40°15'N, 14°54'E, Salerno province) (Figure. 1). The area was dominated by Mediterranean vegetation, characterized by olive tree cultivation, rural buildings, and stone walls.

By using nylon loops, we sampled 16 individuals: 6 dark-diurnal geckos collected on the trunks, and 10 pale-nocturnal geckos captured on the wall.

For each gecko, a tip of the tail was collected after its autotomy and preserved in a sterile tube with 70% Et-OH, stored at -20°C until the processing in the laboratory. After tissue collection, the animals were released at the point of capture, except for 6 individuals (3 dark-diurnal geckos and 3 pale-nocturnal geckos) that were sacrificed for sampling tissues useful for histological analyses (see after). The animals were sampled with the permission of the local county authority (Cilento, Vallo di Diano e Alburni National Park, prot. 2013/0010678). All experimental protocols were approved by the Ethical Committee for Animal Experiments University of Naples Federico II (protocol 2013/0032826).

Mitochondrial genomic characterization

DNA extraction and sequencing

Total mitochondrial DNA (mtDNA) was extracted from 25 mg of tissue using miniprep PureLink HiPure Plasmid DNA Purification Kits (Invitrogen), according to manufacturer's recommendations, except for initial mechanical disruption of samples: the procedure followed is 1h in a Phosphate

Buffer Saline (PBS). Blank extractions were performed at each extraction to monitor potential cross-

contaminations. Finally, we checked the concentration and quality of DNA using the Qubit

Fluorometer (Thermofisher Scientific) and the Agilent 4200 TapeStation system (Agilent

Technologies), respectively.

High throughput sequencing of the mitochondrial genomes was performed on the Illumina 2500

platform using the Nextera XT DNA Sample Preparation protocol at the Genomix4Life Srl

(http://www.genomix4life.com/ it/).

Processing of raw sequencing data was performed as reported in (Fulgione et al. 2019). In brief, the

FastQC software v0.11.4 (http://www.bioinformatics. babraham.ac.uk) was used to quality control

check of raw sequence data. The Trimmomatic software v0.35 (Bolger et al. 2014), was

implemented, in paired end mode, to trim and crop the low-quality raw reads (Q < 28), and to

remove the primers and Illumina adapters. Paired reads were processed simultaneously, while

orphan reads were discarded from the subsequent analyses.

The assembly was performed in Geneious version R9.16 (Kearse et al. 2012) in mapping assembler

mode. For each sample, the forward and reverse filtered reads were used to create contiguous

sequences (contigs) in *de novo* assembly procedure. To check the quality of assembly, the contig

sequences have been mapped against a *T. mauritanica* reference sequence (NCBI accession number

EU443255).

Phylogenetic characterization and time tree

To evaluate the haplotype of sequenced mitochondrial genomes, we aligned them in Geneious

version R9.16 with sequences of *T. mauritanica* (NCBI accession number from JQ300539 to

JQ301443), representative of different lineages (Rato et al. 2012), using *Gekko gecko* (accession

number AY282753) as the outgroup.

To infer the split time of pale-nocturnal and dark-diurnal geckos' populations, we elaborated a time-

tree in MEGA11 (Tamura et al. 2021). We used 10 pale-nocturnal, 6 dark-diurnal geckos sequences,

and *T. mauritanica* sequence belonging to Clade I (accession number JQ425050) as outgroup. We

also added a *T. mauritanica* Clade II sequence (accession number JQ425041) to allow a better

resolution of time-lines. Indeed, Clade I and II are temporally the closest (Rato et al. 2012) .

Before that, to evaluate the haplotype for JQ425041 and JQ425050, we aligned these with

sequences of *T.mauritanica* from JQ300539 to JQ301443 (Rato et al. 2012), representative of

different lineages of *T. mauritanica*.

Firstly, an alignment with Muscle was performed considering the 18 mtDNA sequences. The

following parameters were set: gapopn = -400.00, gapextend = 0.00, max iters = 16, 2 iterations,

cluster methods = UPGMA.

Maximum-Likelihood uses several substitution models to correct for multiple changes at the same

site during the evolutionary history of the sequences. To set the best model, we perform a test

considering the lowest Bayesian Information Criteria (BIC). According this criteria the Tamura-Nei

method was selected (BIC:52037,98) for the construction of the phylogenetic tree.

A timetree was inferred by applying the RelTime method (Tamura et al. 2012, 2018) on 18TN93Tree

in MEGA11 (Tamura et al. 2021) using a dataset of 16,618 positions.

Branch lengths were calculated using the Maximum Likelihood (ML) method and the General Time

Reversible substitution model (Nei and Kumar 2000).

The timetree was computed using 1 calibration constraint. Times were not estimated for outgroup

nodes, because the Reltime method uses evolutionary rates from the ingroup to calculate

divergence times and doesn't assume that evolutionary rates in the ingroup clade apply to the

outgroup.

The estimated log likelihood value of the tree is -25,781.91. A discrete Gamma distribution was used

to model evolutionary rate differences among sites (5 categories; +G; parameter = 0.6794).

To confirm the results obtained with MEGA11, the divergence time estimation was performed also

with BEAST v1.14 (Suchard et al. 2018), setting the following parameters: Substitution Model= TN93,

Gamma Category Count= 5, Strick Clock Model and setting priors to Coalescent Constant Population.

Species Distribution models

For both species, we defined the Mediterranean region as the study area, according to the current

and historical (Rato et al. 2012) geographical distribution of *T. mauritanica*.

Then, we generated by random sampling 10000 background points within the background area.

To account for potential sampling biases, pseudoabsences were geographically placed according to

the density of the occurrence data, so pseudoabsences are more abundant where presences are

denser (Phillips and Dudík 2008; Syfert et al. 2013; Roy-Dufresne et al. 2019).

For *H. sapiens* only, the record was divided into 1000-years consecutive time bins according to the

time resolution of bioclimatic predictors.

Subsequently, we partitioned the pseudoabsences proportionally to the number of presences per

time bin.

After extracting climatic values in each occurrence and pseudoabsence datapoints, we accounted

for multicollinearity among predictors by considering the Variance Inflation Factor (VIF).

The latter was assessed by using the function "vifcor" embedded in the "usdm" R package and

selecting a threshold of 0.75 (Zuur et al. 2010; Naimi et al. 2014).

After VIF application, the selected predictors were BIO4 (temperature seasonality), BIO8 (mean

temperature of the wettest quarter), BIO 9 (mean temperature of driest quarter), BIO 13

(Precipitation of Wettest Month), BIO 14 (precipitation of driest month), BIO 15 (Precipitation

Seasonality), BIO 19 (precipitation of coldest quarter).

To estimate the geographic distribution of both species in the past, we adopted the SDM ensemble

approach. Specifically, we trained SDMs through an ensemble forecasting approach, as implemented

in the R package "biomod2" (Thuiller et al. 2009) .

We considered four different algorithms: Maximum Entropy Models (MaxEnt), Generalized Boosted

Models (GBM), Random Forests (RF), and Generalized Linear Models (GLM). As model tuning, we

adopted the default setting described in "biomod" R package. To evaluate the predictive accuracy of

SDMs, we randomly split each occurrence dataset into a 70% sample, used for the calibration of the

model, and the remaining 30%, used to evaluate model performance. Then, we calculated the area

under the receiver operating characteristic curve (AUC (Hanley and McNeil 1982)) and the true skill

statistic (TSS (Allouche et al. 2006)). This entire procedure was iterated 10 times, changing the

randomly selected training/testing datapoints at each iteration. Model averaging was performed by

weighting the individual model projection by their AUC values and averaging the results (Marmion

et al. 2009) after excluding the model with AUC <0.7.

Lastly, SDM predictions were projected in the past to obtain spatial explicit habitat suitability of *H.*

sapiens and *T. mauritanica* in the Mediterranean area during the 60-20 kyr interval.

Human-gecko sympatry

We collected modern occurrence data for *T. mauritanica* from the "Global Biodiversity Information

Facility" (GBIF, www.gbif.org/) database by selecting "human observation" and "machine

observation" as the basis of the record.

The data were further filtered by removing duplicate records and occurrence records devoid of

geographical coordinates.

For the human dataset, we implemented the *H. sapiens* record published in (Raia et al. 2020) and

(Timmermann et al. 2022), by focusing on the fossil presence observations dated in between 60 and

20 kya.

Radiocarbon data were calibrated by using the "Bchron" R package (Haslett and Parnell 2008)

through the "intcal20" curve (Reimer et al. 2020).

As environmental predictors, we used the monthly bioclimatic variables generated through the 2Ma

CESM1.2 simulation (Timmermann et al. 2022) downscaled at 0.5° × 0.5° grid resolution. The native

set of predictors was subsequently converted in the bioclimatic variables according to WorldClim

using the "dismo" R package (Hijmans et al. 2023). Lastly, variables were projected on the Mollweide

coordinate reference system.

To prevent model over-fitting, duplicate occurrences in raster grid were removed. After this step, we

gathered on 520 modern *T. mauritanica* and 740 past *H. sapiens* occurrences.

***Homo sapiens – Tarentola mauritanica* spatial correlation**

To verify whether there was a significant spatial correlation between *H. sapiens* and *T. mauritanica*,

we adopted the Boyce Index (Hirzel et al. 2006). It requires presences datapoints only and measures

how much model predictions differ from a random distribution of the observed presences across

the prediction gradients (Boyce et al. 2002). Boyce Index varies between -1 and +1, whereby positive

values indicate a model which present predictions are consistent with the distribution of presences

in the evaluation dataset, values close to zero mean that the model is not different from a random

model, negative values indicate counter predictions, i.e., predicting poor quality areas where

presences are more frequent (Hirzel et al. 2006).

In our case, we used Boyce Index to compare the prediction of *T. mauritanica* in the past associated

to human presence and the prediction of *T. mauritanica* across the entire background area (i.e.,

Mediterranean area).

Moreover, we trained a Linear Mixed-Effects Model to test for the presence of ecological correlation

between humans and the Moorish gecko in the past. To this aim, we merged the habitat suitability

values of the two species associated with both human presence and pseudoabsence datapoints

across time. Lined Models were fitted by selecting the habitat suitability values as independent and

response variables and by choosing the "time" as a random effect.

Paleoecology simulation

Two walls of 4x4 square meters each were set up on a site within the study area. A wall has a fire

placed at the base and the other was without.

Over the course of 5 hours at night, all the Arthropoda resting on the two walls were captured and

identified through the examination of diagnostic morphological traits under a stereomicroscope

Leica EZ4 W and specialized reference manual (Bellmann 2019). We recorded the Orders and the

corresponding number of individuals and then determine diversity and richness by Past3 software

(Hammer et al. 2001).

Reflectance

To objectively determine the skin coloration of sampled geckos, we measured the skin reflectance

by spectrophotometry (250–1000 nm, AvaSpec-2048-USB2-UA-50; Avantes, Apeldorn, Netherlands)

considered a range between 300 and 700 nm (Vroonen et al. 2012). A white reference tile (WS2;

Avantes) was used as a reference for calibration.

The spectrophotometer probe (with a hole end of 0.2 mm) was held perpendicular to the body

surface of animals and the reflectance (R%) was recorded in three positions on the back. Then, the

average of the integrals subtended the reflectance curves was assumed to be representative of the

whole back of each individual (Vroonen et al. 2012; Fulgione et al. 2019, 2015). The difference

between diurnal/dark and nocturnal/pale groups was assessed by a one-way ANOVA test. And

statistical significance was defined as $p < 0.05$.

Skin histology

Three dark and three light geckos were anaesthetized with 250 mg/kg 1% MS-222 (Sigma Chemical

Co. St. Luis, MO) injected into the intracoelomic cavity and were sacrificed by decapitation. Skin

samples deriving from 3 dark and 3 light geckos' backs, were processed for light microscopy as in

(Avallone et al. 2018). Briefly, after fixation in 2% paraformaldehyde and 2.5% glutaraldehyde (4h at

4 °C) and post-fixation in 2% osmium tetroxide solution (1 h, at 4 °C), they were dehydrated in

ascending series of ethyl alcohol and embedded in Epon 812 resin (Fluka). Semi-thin sections of 1.5

μm were cut with a Super Nova Leica Ultratome and stained with 1% toluidine blue in 1% sodium

tetraborate buffer. Sections were examined with a Zeiss Axiocam camera applied to a Zeiss Axioskop

microscope (Zeiss, Jena, Germany).

Melanocyte stimulating Hormone (α-MSH) and Melanin assay

Plasma and skin were collected from 3 nocturnal/pale and 3 diurnal/dark geckos.

Blood was sampled with a heparinized syringe from interdiscal vertebra windows of the tail. The

plasma was isolated by centrifugation at 2000 g for 10 min at room temperature and samples were

stored at −80 °C until processing in laboratory.

α-MSH levels were determined by the ELISA assay as described in Monti and co-authors (Monti et

al. 2013). The Student's t test was used to test statistical significance difference (p < 0.05) between

the two groups.

To quantify melanin, we used protocol proposed by Kalie and co-authors (Kalie et al. 2013), with the

following excepts: tissues collection in 50 mM Tris-HCl pH 7.4, 300 mM NaCl, 0.5% NP-40 and

protease inhibitors (Roche), and lysing of cells by freeze-thawing (3 cycles of dried ice at -37 °C),

Dounce homogenization (200 strokes) and sonication (2 min, 10 sec on and 10sec off).

Results

Divergence timing

The phylogeny of the considered geckos' populations, using the complete mitochondrial genome,

highlights their monophyly by placing the common ancestor at about 500 thousand years ago (0.28-

0.71 Mya) (Fig. 4). Subsequently, the topology shows a dichotomous division in two clades, dark-

diurnal and pale-nocturnal, also monophyletic. The common ancestor of the pale-nocturnal geckos

seems to be traced back to 40 thousand years ago (range 0.02 – 0.06 Mya).

Figure 4 – **Phylogenetic reconstruction**. Phylogeny and divergence time estimation obtained by molecular-clock analysis of 10 pale-nocturnal (N1-N10) and 6 dark-diurnal geckos (D1-D6). Divergence times of the two gecko's populations were calculated using the complete mitochondrial genome. The red bar on the nodes indicates the 95% credibility intervals of the estimated posterior distributions of the divergence times. Mya: million years ago.

Species Distribution Models

The SDMs developed for humans reached high predictive performances, expressed by the AUC value equal to 0.818 (sd=0.014) and TSS equal to 0.502 (sd=0.028).

The SDMs developed for *T. mauritanica* reached very high performances, expressed by the AUC value

equal to 0.96 (sd=0.002) and the TSS equal to 0.803 (sd=0.008).

We estimated the magnitude of the spatial correlation between humans and the Moorish gecko in

the past using the Boyce Index. Specifically, we obtained a Spearman correlation value equal to

0.422.

Linear Mixed-Effects Model shows a positive and significant relationship between *H. sapiens* and *T.*

mauritanica habitat suitability values (p<0.001) (Fig. 5). Furthermore, the model showed a relevant

age-driven random effect structure (p=0.016).

Figure 5 - **Spatial distribution model of *Tarentola mauritanica* and *Homo sapiens* 40 kylo years age**.

Linear Mixed-Effects Model shows a positive and significant relationship between *H. sapiens* (green

dots) and *T. mauritanica* habitat suitability (red area).

Paleoecology simulation

The paleoecology simulation experiment in the field made possible to determine the effectiveness of a light source (fire) placed at the base of the wall in attracting Arthropods. The number and diversity of taxa sampled to the wall lighted by fire was significantly higher than the wall without input of fire (Fig. 6) (for diversity: Chi squared = 6.250 with 1 degrees of freedom; the two-tailed P = 0.0124. For abundance: Chi squared = 38.111 with 1 degrees of freedom; the two-tailed $P < 0.0001$).

Figure 6 – Paleoecology simulation. Arthropods richness and number of individuals on the walls with (white histogram) and without (grey histogram) a light source. The colored pies represent the proportion of taxa for each Order.

Histological skin analysis

Although the two populations (diurnal and nocturnal) retain the ability to vary the level of darkness/lightness (Fulgione et al. 2019), nocturnal geckos are characterized by a marked depigmentation. The comparison between histological skin samples taken from the backs of pale-

nocturnal and dark-diurnal geckos showed that dark phenotype geckos' skin has numerous, predominantly large, melanophores. In addition, a thick and continuous layer of pigment granules is observed in the connective tissue of the underlying dermis. By contrast, in the skin of pale-nocturnal specimens, the melanophores appear remarkably reduced in size and less abundant, and the layer of pigment surrounding them is exiguous (Fig. 7).

Figure 7 - **Histological skin analysis.** Skin sections stained with toluidine blue. A) Skin from the backs of dark-diurnal gecko, and single melanophores (inset). There is a high concentration of melanophores (thick arrow) localized in the dermal region and a thick layer of melanin granule (arrow). B) Picture A digitally stained to emphasize melanophores and melanin pigment. C) Skin from the backs of a pale-nocturnal gecko and a single melanophore (inset). D) Picture C digitally stained to emphasize melanophores and a layer of melanin pigment.

α-MSH and Melanin assay

The α-MSH hormone, produced by the pituitary gland, stimulates the melanosomes to produce

melanin. The blood levels of α-MSH are significantly lower in nocturnal-pale geckos (Fig. 8A), so they

will consequently produce a little amount of melanin in the dermis (Fig. 8B), which leads to a

decrease of the skin's reflectance (Fig. 8C).

Figure 8 - α-MSH and Melanin assay. A) α-MSH hormone blood concentration, B) melanin assay in

skin, and C) reflectance of the skin in dark-diurnal (D) and pale-nocturnal (N) geckos (one-way

ANOVA test $p<0.05$).

Discussion

There was a moment in the history of life on Earth when humans began to be decisive for the life of

the entire Planet. Three major revolutions shaped human history: Cognitive (70 kya), Agricultural

(10 kya), and Scientific (500 ya) revolutions (Harari 2018), which marked a deep watershed between

H. sapiens before and after this moment. The relationship *H. sapiens* established with animals and

plants has reverberated in the ecology and evolution of organisms, up to the present day.

Investigating the events that marked this phase, identifying either the animals and the plants

involved, their propensity to change, and the impact of the change, brings new evidence to

comprehend the evolutionary processes, the history of life on Earth and ourselves.

The human impact on natural populations can affect their phenotype, playing an important role in

evolution. In wild populations, the propensity to establish a relationship with humans was allowed

by a plastic capacity, whether domesticated, synanthropic or commensal. Indeed, plasticity confers

the ability of niche exploration a prominent potential, especially in order to adapt to the anthropic

niche, due to the peculiar speed of its creation and change.

Our results highlight a remarkable variation, both in morphological and functional terms, of the pale-

nocturnal gecko's population. The latter derived from the dark-diurnal population, and their

common ancestor has been dated to around ca 40,000 years ago. In our hypothesis, starting from

that moment, pale-nocturnal geckos would have been differentiated by extending the boundaries

of their plasticity. Indeed, their depigmentation went beyond the lightning value that the dark-

diurnal gecko could normally be able to achieve. The pale phenotype has been reached through the

reduction of α-MSH hormone as well as the number and size of melanosomes in the dermis

(reinforcement).

This lightening would have responded to the main selective pressure effective in the dark

environment: the highly frequent predation of a nocturnal animal on a light substrate.

However, most likely they would have triggered metabolic implications: a heterothermic lizard can

live in the absence of the sun only if it significantly elevates its metabolism, eating more, as pale-

nocturnal geckos should have accomplished during evolution. However, this point is worthy of future

investigation.

Our study inaugurates a new and interesting line of investigation that could shed light on the impact

that humans have on the evolutionary trajectories of non-domesticated species, endowed with

adaptive plasticity, functional to the exploration and colonization of the anthropic niche.

The case of the Moorish gecko is emblematic. It clearly describes a species with plastic capacities:

its adaptation results from the ability to become cryptic on different substrates. This attitude, in

presence of artificially lit environments, would have been translated into the ability to invade the

new artificial niche, with the subsequent reinforcement of the plastic ability. This leap would have

created the pale-nocturnal population of geckos with a common ancestor when *H. sapiens* deal with

managed fire.

The timing of the human control of fire is a controversial issue (Clark and Harris 1985; Gowlett et al.

1981). The European evidence suggests that Neanderthals and early modern humans used the

controlled and constant fire starting from the Middle Pleistocene onward in Europe, and from the

Late Pleistocene in Europe and South Africa, for many other purposes than just warming and cooking

(Henry et al. 2011; Roebroeks and Villa 2011).

The first ideal case of fire documentation, based on a piece of pitch keeping a human fingerprint, is

dated at ca 48,000 BP on the limits of techniques and compatible with a geological age of

approximately 80,000 years. Although during the Middle Pleistocene, recognizable hearths

demonstrate a social and economic focus on many sites attributable to the early hominins.

Interestingly, in that Era there was a clear sympatry between geckos and humans.

Over time, genetic changes have occurred in the nocturnal population, which shifted the coloration

from an induced plastic response, by the reduction of melanin aggregation in the dermis, to a lighter

coloration, and improved crypticism due to a constitutive reduction of this pigment. Probably, these

genetic changes were subsequently increased in frequency in the pale-nocturnal population through

positive selection, whereas obviously the pale-nocturnal geckos would be hardly targeted by

predators (Fulgione et al. 2019); consequently, they would be more likely to survive and reproduce,

thus passing on the genetic changes underlying the pale phenotype to their offspring.

Plasticity facilitated the initial colonization and survival of the population in a new environment and

allowed time for a subsequent genetic adaptation to express a "fine-tune" response to this

environment. Plastic capabilities of the phenotype can respond to selection, anticipating the

genotype's adaptation process and gaining the colonization of new niches. This new evolutionary

perspective, however fascinating, requires accomplishing several empirical tests in species with

different plastic capacities.

The gecko brings something new to our attention. Indeed, the challenge the gecko populations have

faced is totally unexpected. This is not a new niche, however foreseen by a natural context, the

nocturnal niche with artificial light is a niche's new dimension created by humans. A sort of

environmental change that follows from those possible foreseen by natural systems.

The light of the fire that pierces the night is not a consequence of the evolution of natural systems,

but it's like "pulled-out of a hat rabbit", when nothing suggests this.

Conflict of Interest

The authors declare no conflicts of interest.

Author Contributions: Conceptualization: D.F. and M.B.; Sample collection: D.F., E.R., V.M., and M.B;

Laboratory experiments: V.M., E.R., B.A. and M.B; Analyses of data: D.F., E.R., B.A., A.M., G.G., and

M.B; Supervision: D.F.; Writing original draft: D.F., E.R. and M.B.; Review & editing: D.F., V.M., E.R.,

B.A., A.M., G.G. and M.B. All the authors read and approved the manuscript.

Data Availability

The sequencing data are deposited in GenBank under accession numbers MK275668 - MK275687

Acknowledgments

We thank the Professor Greger Larson for suggestion and revision of the manuscript.

References

Allouche O, Tsoar A, Kadmon R. 2006. Assessing the accuracy of species distribution models: prevalence, kappa and the true skill statistic (TSS): Assessing the accuracy of distribution models. *Journal of Applied Ecology* **43**: 1223–1232.

Ålund M, Cenzer M, Bierne N, Boughman JW, Cerca J, Comerford MS, Culicchi A, Langerhans B, McFarlane SE, Möst MH, et al. 2023. Anthropogenic Change and the Process of Speciation. *Cold Spring Harb Perspect Biol* **15**: a041455.

Arnold N. 2002. *Collins Field Guide to the Reptiles and Amphibians of Britain and Europe*. Collins; New edition. Harper Collins, London.

Avallone B, Tizzano M, Cerciello R, Buglione M, Fulgione D. 2018. Gross anatomy and ultrastructure of Moorish Gecko, Tarentola mauritanica skin. *Tissue and Cell* **51**: 62–67.

Baker BJ, Richardson JML. 2006. The effect of artificial light on male breeding-season behaviour in green frogs, *Rana clamitans melanota*. *Can J Zool* **84**: 1528–1532.

Beier P. 2006. Effects of artificial night lighting on terrestrial mammals. In *Ecological consequences of artificial night lighting*, pp. 19–42, Island Press https://books.google.it/books?hl=it&lr=&id=dEEGtAtR1NcC&oi=fnd&pg=PA19&dq=Beier,+P.+(2006).+Effects+of+artificial+night+lighting+on+terrestrial+mammals.+Ecological+consequences+of+artificial+night+lighting,+19-

42.&ots=86T7d698lL&sig=y9ydAuJYwSQkEuR2212wlDsBVVs&redir_esc=y#v=onepage&q&f

=false.

Bellmann H. 2019. *Guida agli insetti d'Europa. Ediz. illustrata*. Ricca Editore.

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data.

Bioinformatics **30**: 2114–2120.

Booth W, Santangelo RG, Vargo EL, Mukha DV, Schal C. 2011. Population Genetic Structure in German

Cockroaches (Blattella Germanica): Differentiated Islands in an Agricultural Landscape.

Journal of Heredity **102**: 175–183.

Bowman DMJS, Balch J, Artaxo P, Bond WJ, Cochrane MA, D'Antonio CM, DeFries R, Johnston FH,

Keeley JE, Krawchuk MA, et al. 2011. The human dimension of fire regimes on Earth: The

human dimension of fire regimes on Earth. *Journal of Biogeography* **38**: 2223–2236.

Boyce MS, Vernier PR, Nielsen SE, Schmiegelow FKA. 2002. Evaluating resource selection functions.

Ecological Modelling **157**: 281–300.

Carranza S, Arnold EN, Mateo JA, López-Jurado LF. 2000. Long-distance colonization and radiation in

gekkonid lizards, *Tarentola* (Reptilia: Gekkonidae), revealed by mitochondrial DNA

sequences. *Proc R Soc Lond B* **267**: 637–649.

Clark JD, Harris JWK. 1985. Fire and its roles in early hominid lifeways. *Afr Archaeol Rev* **3**: 3–27.

Fulgione D, Buglione M, Rippa D, Trapanese M, Petrelli S, Monti DM, Aria M, Del Giudice R, Maselli

V. 2019. Selection for background matching drives sympatric speciation in Wall Gecko. *Sci

Rep* **9**: 1288.

Fulgione D, Lega C, Trapanese M, Buglione M. 2015. Genetic factors implied in melanin-based

coloration of the Italian wall lizard. *J Zool* **296**: 278–285.

Fulgione D, Trapanese M, Maselli V, Rippa D, Itri F, Avallone B, Van Damme R, Monti DM, Raia P. 2014.

Seeing through the skin: dermal light sensitivity provides cryptism in moorish gecko. *Journal*

of Zoology **294**: 122–128.

Gamble T, Greenbaum E, Jackman TR, Russell AP, Bauer AM. 2012. Repeated Origin and Loss of

Adhesive Toepads in Geckos ed. J. Castresana. *PLoS ONE* **7**: e39429.

Gowlett JAJ. 2016. The discovery of fire by humans: a long and convoluted process. *Phil Trans R Soc*

B **371**: 20150164.

Gowlett JAJ, Harris JWK, Walton D, Wood BA. 1981. Early archaeological sites, hominid remains and

traces of fire from Chesowanja, Kenya. *Nature* **294**: 125–129.

Hale R, Swearer SE. 2016. Ecological traps: current evidence and future directions. *Proc R Soc B* **283**:

20152647.

Hammer Ø, Harper D, Ryan P. 2001. Past: Paleontological Statistics Software Package for Education

and Data Analysis. *Palaeontologia Electronica* **4**: 9.

Hanley JA, McNeil BJ. 1982. The meaning and use of the area under a receiver operating

characteristic (ROC) curve. *Radiology* **143**: 29–36.

Harari YN. 2018. *Sapiens: a brief history of humankind*. First Harper Perennial edition. Harper

Perennial, New York.

Haslett J, Parnell A. 2008. A Simple Monotone Process with Application to Radiocarbon-Dated Depth

Chronologies. *Journal of the Royal Statistical Society Series C: Applied Statistics* **57**: 399–418.

Henry AG, Brooks AS, Piperno DR. 2011. Microfossils in calculus demonstrate consumption of plants

and cooked foods in Neanderthal diets (Shanidar III, Iraq; Spy I and II, Belgium). *Proc Natl*

Acad Sci USA **108**: 486–491.

Hijmans R, Phillips S, Leathwick J, Elith J. 2023. dismo: methods for species distribution modeling,

that is, predicting the environmental similarity of any site to that of the locations of known

occurrences of a species. https://rspatial.org/raster/sdm/.

Hirzel AH, Le Lay G, Helfer V, Randin C, Guisan A. 2006. Evaluating the ability of habitat suitability

models to predict species presences. *Ecological Modelling* **199**: 142–152.

Hulme-Beaman A, Dobney K, Cucchi T, Searle JB. 2016. An Ecological and Evolutionary Framework

for Commensalism in Anthropogenic Environments. *Trends in Ecology & Evolution* **31**: 633–

645.

IUCN. 2017. Tarentola mauritanica: Vogrin, M., Corti, C., Pérez Mellado, V., Baha El Din, S. &

Martínez-Solano, I.: The IUCN Red List of Threatened Species 2017: e.T61578A63716927.

http://www.iucnredlist.org/details/61578/0 (Accessed November 29, 2023).

Kalie E, Razi M, Tooze SA. 2013. ULK1 Regulates Melanin Levels in MNT-1 Cells Independently of

mTORC1 ed. W.-G. Zhu. *PLoS ONE* **8**: e75313.

Kamp J, Sheldon RD, Koshkin MA, Donald PF, Biedermann R. 2009. Post-Soviet steppe management

causes pronounced synanthropy in the globally threatened Sociable Lapwing *Vanellus*

gregarius. *Ibis* **151**: 452–463.

Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz

S, Duran C, et al. 2012. Geneious Basic: An integrated and extendable desktop software

platform for the organization and analysis of sequence data. *Bioinformatics* **28**: 1647–1649.

Kendal J, Tehrani JJ, Odling-Smee J. 2011. Human niche construction in interdisciplinary focus. *Phil*

Trans R Soc B **366**: 785–792.

Marmion M, Parviainen M, Luoto M, Heikkinen RK, Thuiller W. 2009. Evaluation of consensus

methods in predictive species distribution modelling. *Diversity and Distributions* **15**: 59–69.

Martín B, Pérez H, Ferrer M. 2018. Effects of natural and artificial light on the nocturnal behaviour

of the wall gecko. *Animal Biodiversity and Conservation* **41.2**: 209–215.

Monti DM, Raia P, Vroonen J, Maselli V, Van Damme R, Fulgione D. 2013. Physiological change in an

insular lizard population confirms the reversed island syndrome: Physiology in Insular Lizards.

Biol J Linn Soc Lond **108**: 144–150.

Naimi B, Hamm NAS, Groen TA, Skidmore AK, Toxopeus AG. 2014. Where is positional uncertainty a

problem for species distribution modelling? *Ecography* **37**: 191–203.

Nei M, Kumar S. 2000. *Molecular Evolution and Phylogenetics*. Oxford University Press Inc.

Park J-H, Lee H-S. 2017. Phototactic behavioral response of agricultural insects and stored-product

insects to light-emitting diodes (LEDs). *Appl Biol Chem* **60**: 137–144.

Parsons KJ, Rigg A, Conith AJ, Kitchener AC, Harris S, Zhu H. 2020. Skull morphology diverges between

urban and rural populations of red foxes mirroring patterns of domestication and

macroevolution. *Proc R Soc B* **287**: 20200763.

Phillips SJ, Dudík M. 2008. Modeling of species distributions with Maxent: new extensions and a

comprehensive evaluation. *Ecography* **31**: 161–175.

Pocock MJO, Searle JB, White PCL. 2004. Adaptations of animals to commensal habitats: population

dynamics of house mice Mus musculus domesticus on farms: Population dynamics of

commensal house mice. *Journal of Animal Ecology* **73**: 878–888.

Puckett EE, Orton D, Munshi-South J. 2020. Commensal Rats and Humans: Integrating Rodent

Phylogeography and Zooarchaeology to Highlight Connections between Human Societies.

BioEssays **42**: 1900160.

Raia P, Mondanaro A, Melchionna M, Di Febbraro M, Diniz-Filho JAF, Rangel TF, Holden PB,

Carotenuto F, Edwards NR, Lima-Ribeiro MS, et al. 2020. Past Extinctions of Homo Species

Coincided with Increased Vulnerability to Climatic Change. *One Earth* **3**: 480–490.

Rato C, Carranza S, Harris DJ. 2012. Evolutionary history of the genus Tarentola (Gekkota:

Phyllodactylidae) from the Mediterranean Basin, estimated using multilocus sequence data.

BMC Evol Biol **12**: 14.

Rato C, Carretero MA. 2015. Ecophysiology Tracks Phylogeny and Meets Ecological Models in an

Iberian Gecko. *Physiological and Biochemical Zoology* **88**: 564–575.

Rato C, Silva-Rocha I, Sillero N. 2023. What does the future hold for a thermophilic and widely

introduced gecko, Tarentola mauritanica (Squamata: Phyllodactylidae)? *Biol Invasions*.

https://link.springer.com/10.1007/s10530-023-03229-7 (Accessed February 27, 2024).

Ravinet M, Elgvin TO, Trier C, Aliabadian M, Gavrilov A, Sætre G-P. 2018. Signatures of human-

commensalism in the house sparrow genome. *Proc R Soc B* **285**: 20181246.

Raviv L, Jacobson SL, Plotnik JM, Bowman J, Lynch V, Benítez-Burraco A. 2023. Elephants as an animal

model for self-domestication. *Proc Natl Acad Sci USA* **120**: e2208607120.

Reimer PJ, Austin WEN, Bard E, Bayliss A, Blackwell PG, Bronk Ramsey C, Butzin M, Cheng H, Edwards

RL, Friedrich M, et al. 2020. The IntCal20 Northern Hemisphere Radiocarbon Age Calibration

Curve (0–55 cal kBP). *Radiocarbon* **62**: 725–757.

Rieppel O. 1981. Tarentola mauritanica (Linnaeus, 1758) - Wiesbade. In *Handbuch der Reptilien und*

Amphibien Europas, Vol. 1 of, pp. 119–133, Akademische Verlagsgesellschaft, Wiesbaden.

Roebroeks W, Villa P. 2011. On the earliest evidence for habitual use of fire in Europe. *Proc Natl Acad*

Sci USA **108**: 5209–5214.

Roy-Dufresne E, Saltré F, Cooke BD, Mellin C, Mutze G, Cox T, Fordham DA. 2019. Modeling the

distribution of a wide-ranging invasive species using the sampling efforts of expert and citizen

scientists. *Ecology and Evolution* **9**: 11053–11063.

Russo D, Cistrone L, Libralato N, Korine C, Jones G, Ancillotto L. 2017. Adverse effects of artificial

illumination on bat drinking activity. *Animal Conservation* **20**: 492–501.

Schlaepfer MA, Runge MC, Sherman PW. 2002. Ecological and evolutionary traps. *Trends in Ecology*

& Evolution **17**: 474–480.

Shimelmitz R, Kuhn SL, Jelinek AJ, Ronen A, Clark AE, Weinstein-Evron M. 2014. 'Fire at will': The

emergence of habitual fire use 350,000 years ago. *Journal of Human Evolution* **77**: 196–203.

Stepka Z, Azuri I, Horwitz LK, Chazan M, Natalio F. 2022. Hidden signatures of early fire at Evron

Quarry (1.0 to 0.8 Mya). *Proc Natl Acad Sci USA* **119**: e2123439119.

Suchard MA, Lemey P, Baele G, Ayres DL, Drummond AJ, Rambaut A. 2018. Bayesian phylogenetic

and phylodynamic data integration using BEAST 1.10. *Virus Evolution* **4**.

https://academic.oup.com/ve/article/doi/10.1093/ve/vey016/5035211 (Accessed

November 29, 2023).

Syfert MM, Smith MJ, Coomes DA. 2013. Correction: The Effects of Sampling Bias and Model

Complexity on the Predictive Performance of MaxEnt Species Distribution Models ed. D.L.

Roberts. *PLoS ONE* **8**. https://dx.plos.org/10.1371/annotation/35be5dff-7709-4029-8cfa-

f1357e5001f5 (Accessed October 10, 2023).

Tamura K, Battistuzzi FU, Billing-Ross P, Murillo O, Filipski A, Kumar S. 2012. Estimating divergence

times in large molecular phylogenies. *Proc Natl Acad Sci USA* **109**: 19333–19338.

Tamura K, Stecher G, Kumar S. 2021. MEGA11: Molecular Evolutionary Genetics Analysis Version 11

ed. F.U. Battistuzzi. *Molecular Biology and Evolution* **38**: 3022–3027.

Tamura K, Tao Q, Kumar S. 2018. Theoretical Foundation of the RelTime Method for Estimating

Divergence Times from Variable Evolutionary Rates ed. C. Russo. *Molecular Biology and*

Evolution **35**: 1770–1782.

Thuiller W, Lafourcade B, Engler R, Araújo MB. 2009. BIOMOD - a platform for ensemble forecasting

of species distributions. *Ecography* **32**: 369–373.

Timmermann A, Yun K-S, Raia P, Ruan J, Mondanaro A, Zeller E, Zollikofer C, Ponce De León M,

Lemmon D, Willeit M, et al. 2022. Climate effects on archaic human habitats and species

successions. *Nature* **604**: 495–501.

Valverde JA. 2004. *Estructura de una comunidad mediterránea de vertebrados terrestres*. Consejería

de Medio Ambiente, Junta de Andalucía, Sevilla.

Van Bocxlaer I, Loader SP, Roelants K, Biju SD, Menegon M, Bossuyt F. 2010. Gradual Adaptation

Toward a Range-Expansion Phenotype Initiated the Global Radiation of Toads. *Science* **327**:

679–682.

Vroonen J, Vervust B, Fulgione D, Maselli V, Van Damme R. 2012. Physiological colour change in the

Moorish gecko, Tarentola mauritanica (Squamata: Gekkonidae): effects of background, light,

and temperature: PHYSIOLOGICAL COLOUR CHANGE IN THE MOORISH GECKO. *Biological*

Journal of the Linnean Society **107**: 182–191.

Wang J-S, Tuanmu M-N, Hung C-M. 2021. Effects of artificial light at night on the nest-site selection,

reproductive success and behavior of a synanthropic bird. *Environmental Pollution* **288**:

117805.

Zohar I, Alperson-Afil N, Goren-Inbar N, Prévost M, Tütken T, Sisma-Ventura G, Hershkovitz I, Najorka

J. 2022. Evidence for the cooking of fish 780,000 years ago at Gesher Benot Ya'aqov, Israel.

Nat Ecol Evol **6**: 2016–2028.

Zuur AF, Ieno EN, Elphick CS. 2010. A protocol for data exploration to avoid common statistical

problems: *Data exploration*. *Methods in Ecology and Evolution* **1**: 3–14.

Graphical Abstract

This image was painted by Domenico Fulgione