Biological Control of Green Mold in Simulated Post-harvest Chain of Citrus Fruit: Efficacy of

Candida oleophila **Strain O and Molecular Insight into Elicitation of Host Immune System**

Ermes Ivan ROVETTO[a], Federico LA SPADA[a*], Soumia EL BOUMLASY[a,b], Sebastiano CONTI

TAGUALI[a,c], Mario RIOLO[a], Antonella PANE[a], and Santa Olga CACCIOLA[a],

[a]Department of Agriculture, Food and Environment, University of Catania, 95123 Catania, Italy;

[b]Laboratory of Materials-Catalysis, Chemistry Department, Faculty of Science, University

Abdelmalek Essaadi, Tetouan B.P. 2117, Morocco.

[c]Department of Agricultural Science, Mediterranean University of Reggio Calabria, 89122 Reggio

Calabria, Italy

[*]Corresponding author: Federico LA SPADA, Department of Agriculture, Food and Environment,

University of Catania, 95123 Catania, Italy. E-mail address: federico.laspada@unict.it

Abstract

Managing post-harvest decays in citrus fruit without relying on conventional pesticides presents a significant

challenge in modern Plant Pathology. This study aimed to evaluate the efficacy of the biological control

agent *Candida oleophila* strain O in controlling green mold caused by *Penicillium digitatum* throughout

various stages of the post-harvest supply chain. Using a series of *in vivo* experiments, different scenarios of

P. digitatum infections in clementine tangerine, orange, and lemon fruit were examined, with treatments

applied before, during or after infection. The study simulated typical conditions of the citrus supply chain,

including picking, processing in packinghouses, and transportation, as well as cold storage and shelf-life

phases. Results indicated that *C. oleophila* exhibited significant efficacy in reducing green mold symptoms,

even at shelf-life temperatures, making it a practical alternative to conventional fungicides. Additionally, the

study provided insights into the molecular mechanisms underlying the defensive response of citrus fruit to *C.*

oleophila treatment, with up-regulation of defense-related genes observed across different fruit types.

Overall, this study underscores the potential of *C. oleophila* as a sustainable and effective solution for

managing post-harvest decays in citrus fruit within the complexities of the supply chain.

Keywords: *Candida oleophila*, citrus green mold, *Penicillium digitatum*, citrus fruit supply chain, biological

control, plant defense transcriptomic response.

1. Introduction

The primary and globally prevalent postharvest challenge impacting citrus fruit is the green mold caused by *Penicillium digitatum* (Abo-Elnaga, 2013; Kanan and Al-Najar, 2008; Plaza et al., 2004; Rovetto et al., 2023, 2024), a destructive disease accounting approximately 90% of total post-harvest losses (Costa et al. 2019; Ismail and Zhang, 2004). This disease causes considerable damages on major commercial citrus crops, specifically oranges (Elsherbiny et al., 2021), lemons (Ben-Yehoshua et al., 2008), tangerines (Suwannarach et al., 2015), and grapefruit (Shi et al., 2018). The citrus green mold is caused by a wound-mediated infection of the fruit peel (Rovetto et al., 2024). At early stages, the infection appears as a softened area of the peel surrounding the infection point (La Spada et al. 2021); this symptom, known as 'Clear Rot' (Ismail and Zhang, 2004), creates conducive conditions for the development of white mycelium and triggers the rapid decay of fruit. Within a few days, the infection progresses to complete molding of the fruit. Simultaneously, the pathogen develops a substantial quantity of green airborne conidia on the fruit surface, serving as a source of inoculum for new infections (Costa et al., 2019; Ismail and Zhang, 2004; Rovetto et al., 2024; Vu et al., 2018).

Due to the high virulence of *P. digitatum* and the potential for related infections to occur at any point, from the field to the consumers' table, the post-harvest supply chain of citrus fruit—encompassing phases of harvesting, warehouse processing (including treatments and packaging), transportation, and delivery to the final customer—deserves particular attention and the implementation of specific practices of control (Fallanaj et al., 2016). Therefore, care in both harvesting and handlings fruit as well as sanitization measures in packing-houses are effective in preventing the disease to minimize peel punctures, wounds, bruises, compression damages, and other mechanical (El-Otmani et al., 2011; Zacarias et al., 2020). Respecting specific temperatures after the warehouse processing, such as 4-8°C during shipment, is an additional practice for limiting pathogen's development in the short term (Khumalo et al., 2021). While these practices could be easily applicable in short-duration supply chains, such as local or national fruit markets, their applicability faces technical limitations in continental and transcontinental markets, where long-duration shipments of fruit at a continuous cold temperature are not cost effective and, therefore, cannot be guaranteed. In these latter contexts, the major strategy for limiting losses by citrus green mold relies

prevalently on post-harvest treatments of fruit with synthetic fungicides (Ismail and Zhang, 2004). In this

respect, the widely employed fungicide imazalil (IMZ), well known for its strong effectiveness against

Penicillium spp., provides both advantages and drawbacks (Bus et al., 1991, 1992; Davé et al., 1989; Holmes

and Eckert, 1999; Ismail and Zhang, 2004; Moraes Bazioli et al., 2019; Sánchez-Torres and Tuset, 2011;

Smilanick et al., 2005). Indeed, while IMZ application is essential for maintaining fruit quality, its

employment has several drawbacks, including the environmental impact, the presence of toxic residues on

the fruit, that in the EU countries is regulated by stringent legislative measures, and the risk of developing

resistant pathogen isolates, consequently diminishing its efficacy (Arslan, 2015; Davé et al., 1989; EFSA,

2010; Gikas et al., 2022; Holmes and Eckert, 1999; Kanetis et al., 2007; Stracquadanio et al., 2021; Strange

and Scott, 2005). Consequently, this has fostered the development of eco-friendly disease management

strategies for controlling citrus green mold, that can be an alternative to synthetic fungicides or may

contribute to reduce their use (Fenta et al., 2019; Hammami et al., 2022; Janisiewicz and Korsten, 2022; La

Spada et al., 2021; Li et al., 2022; Riolo et al., 2023; Stracquadanio et al., 2020; Sui et al., 2016; Talibi et al.,

2014). These include, among others, synthetic resistance elicitors, that activate natural plant defense

mechanisms, and biological control agents (BCAs) (Battha, 2022; Boller, 1995; Côté et al., 1998; Cluzet et

al., 2004; Droby et al., 2009; Du Jardin, 2015). Among resistance elicitors, salts of phosphonic acid

(phosphonates) have gained attention for their applications in agriculture as either fertilizers or fungicides

(Gómez-Merino and Trejo-Téllez, 2015; Manghi et al., 2021). Presently, phosphonates are banned from

organic agriculture by the EU and are restrictively classified as plant protection products (European

Parliament, 2019). Although phosphonates have been exploited mainly for controlling plant diseases caused

by Oomycetes, potassium phosphonate was reported to be also effective against green mold of citrus fruit

(Amiri and Bompeix, 2011; Cerioni et al., 2013; Strano et al., 2015; Souza Morais Reis et al., 2024). Among

potential BCAs of post-harvest rots of citrus fruit, extensive research has been conducted on various species

of yeasts, including *Candida* spp, *Cryptococcus* spp., *Metschnikowia* spp., *Pichia* spp. and *Rhodotorula* spp.

(Droby et al., 1993; Hammami et al., 2022; Kowalska et al., 2022; Liu et al., 2013; Oztekin et al., 2023;

Zhang et al., 2020). *Candida* species, particularly *Candida oleophila*, are at the forefront of commercial

biopesticides, due to their positive effects on plants both as biological control agent and elicitor of plant

defense response (Kowalska et. al., 2022). Currently, the strategic application of elicitors of plant defense

response is emerging as a promising avenue for fortifying disease resistance in agronomic crops (Boller,

1995; Cluzet et al., 2004; Côté et al., 1998; La Spada et al., 2021). In this respect, it is well known that plants

respond to infections through distinct biochemical changes, which are mediated by the transcription of

various defense genes both locally and distantly from the infection site (Boava et al., 2011). These responses

encompass the hypersensitive response (HR), modifications in the cell wall, production of reactive oxygen

species (ROS), accumulation of phytoalexins, and synthesis of pathogenesis-related (PR) proteins (Boava et

al., 2011; La Spada et al., 2021; Van Loon et al., 2006). Some PR proteins, such as β-1,3-glucanases, can

degrade fungal cell wall constituents, specifically β-1,3-glucan and chitin, exhibiting antifungal properties

(Balasubramanian et. al., 2012); for this reason, their biosynthesis and accumulation are considered a

fundamental defense mechanism (Odjakova and Hadjiivanova, 2001). Other PR proteins, such as

peroxidases play, an additional key role in the plant defense response; produced during lignin synthesis, they

contribute to reinforcing the cell wall, thereby enhancing resistance against numerous pathogens (Miedes et

al., 2014). The antioxidant activity of peroxidase has been documented for its positive correlation with the

enhancement of resistance of citrus fruit to *Penicillium* (Ballester et al., 2006). In addition to PR-proteins, the

phenylalanine ammonia-lyase (PAL), the first enzyme in the phenylpropanoid pathway, increases the plant

defense response to various types of stress (Ke and Saltveit, 1989; Youssef et al., 2014; Wang et al., 2023). In

citrus fruit, the activity of phenylalanine ammonia-lyase (PAL) leads to the production of flavonoids (Arcas

et al., 2000; Del Río et al., 1998) and p-coumaric acid derivatives (Afek et al., 1999; Kim et al., 1991),

potentially bolstering the fruit's protection against pathogens. Several studies have delved into evaluating the

antagonistic activity of *C. oleophila* for controlling citrus green mold (Dukare et al., 2019; Laahlali et al.,

2004; Lassois et al., 2008). Additionally, a study of the 2002 by Droby et al. evaluated the ability of *C.*

oleophila in affecting the plant biosynthesis of some defensive metabolites in peels of grapefruit under *P.*

digitatum infection. Despite this attention, there remains a notable gap in understanding the effectiveness of

C. oleophila in controlling citrus green mold throughout the various stages of the citrus fruit supply chain.

Moreover, there is limited knowledge concerning *C. oleophila*'s ability to modulate the triggering of specific

genetic pathways related to the plant-defense response, and whether this ability varies depending on the type

of citrus fruit.

To advance beyond the current state of the art, this study simulated different scenarios of *P. digitatum*

infections in various types of citrus fruit (clementine tangerines, oranges, and lemons) to assess the potential

of *Candida oleophila* in reducing the incidence of citrus green mold throughout the different stages of the

citrus fruit supply chain. Additionally, the study evaluated *C. oleophila*'s ability to modulate the plant-

defense response by analyzing the transcription of genes encoding β-1,3-glucanases, phenylalanine

ammonia-lyase, and peroxidase in clementine tangerine, orange, and lemon fruit.

2. Materials and Methods

2.1 Fungal plant pathogen

The IMZ-sensitive (MIC < 1 μml^{-1} a.i.) *P. digitatum* strain P1PP0 used in this study was from the collection

of the Laboratory of Molecular Plant Pathology of the Department of Agriculture, Food and Environment of

the University of Catania (Catania, Italy) and was characterized in a previous study (El boumlasy et al.,

2021). Before tests, the *P. digitatum* strain P1PP0 was maintained on Potato Dextrose Agar at 25°C.

2.2. Biocide agent and experimental substances

The substances used in this study were: a 10^8 CFU·mL^{-1} suspension in sterile distilled water (sdw) of the

yeast *Candida oleophila* Montrocher strain O (ID substance I01), belonging to the collection of the

Molecular Plant Pathology Laboratory of the Department of Agriculture, Food and Environment (University

of Catania, Catania, Italy); an eco-friendly sanitizing agent containing hydrogen peroxide and acetic acid (ID

substance P01); a 57% (w/v) potassium phosphonate formulation (ID substance RI01); and an emulsifiable

50% w/v concentrate formulation of imazalil fungicide (ID substance F01). All chemicals used in this study

were kindly provided by DECCO Italia srl (Belpasso, CT, Italy), and their application dosages are reported in

Table 1.

Table 1. List of substances used in this study.

ID of treatment	Type of substance	Substance$_a$/ Mixture	Application dosage
P01	Sanitizing agent - eco-friendly, non-toxic substance	hydrogen peroxide and acetic acid	(0.1 mL/L)
RI01	Solution of fertilisers based on phosphoric anhydride and potassium oxide	potassium phosphonate formulation	0.25%[b]
F01	Emulsifiable concentrate formulation of imazalil	imazalil fungicide	1,000 ppm (a.i.*[a]) - Std**
M01	Bioproduct (living microrganism)	Suspension in sterile distilled water of *Candida oleophila* strain O	10^8 CFU·mL^{-1}

a.i.*, active ingredient.
Std**, suggested dosage.

2.3. Citrus fruit

Fruit used in this study were mature and non-wounded clementine tangerines (*Citrus clementina* Hort.

ex Tan., cv. Comune), oranges (*Citrus x sinensis* (L.) Osbeck, cv. Ovale) and lemons (*Citrus x limon* (L.)

Burm. f., cv. Femminello) collected from an organically cultivated orchard. Before each experiment, fruit

underwent preliminary disinfection through a sequence of processes, including washing in NaClO 1%,

followed by rinsing in sterilized distilled water (sdw), and subsequent air drying at room temperature.

2.4. Evaluating the effectiveness of *Candida oleophila* strain O in controlling citrus green mold

The effectiveness of *Candida oleophila* strain O (M01) in controlling citrus green mold was assessed in

orange, lemon, and clementine tangerine fruit. Tests were conducted to simulate as close as possible various

stages of the citrus postharvest chain, spanning from warehousing (including treatment, packaging, and

storage) to retailing. For each type of fruit, these stages were replicated through two consecutive phases

following treatment: a. 7-day storage under refrigerated conditions (at 6°C) and b. an additional 7-day

storage under shelf-life conditions (at 18°C). These storage conditions were included within three realistic

scenarios of fruit infection by *P. digitatum*: i. before treatment; ii. during treatment; iii. 12 hours post-

treatment. Further details of the conducted tests are provided in subsequent sections.

2.4.1. Fruit inoculated with *P. digitatum* before treatment

The peel of each citrus fruit was wounded by a plastic tip (diameter, 2 mm) at four equidistant points in an

equatorial position along the rind surface. Then, each wound was inoculated with 5 μl of a conidial

suspension (10^6 conidia/ml) of *P. digitatum* strain P1PP0; fruit were then incubated at 20°C for 4 hours. After

the incubation period, inoculated fruit were pre-treated by dipping in P01 (0.1 mL/L) for 30 seconds and

rinsing in sdw for additional 30 seconds, or by a simple dipping in sdw for 30 seconds (control). Immediately

after the pre-treatment, at each wound, fruit were treated with a 65 μl spray of RI01, F01, M01 or sdw

(controls). Then, fruit were organized in five batches corresponding to the following treatments: i. fruit pre-

treated only with sdw (control 1), ii. fruit just pre-treated only with P01 (control 2), iii. fruit pre-treated with

P01 and then treated with RI01, iv. fruit pre-treated with P01 and then treated with F01, v. fruit pre-treated

with P01 and then treated with M01. Each treatment comprised three technical replicates made up of five

citrus fruit each**.**

Fruit from each treatment were air dried, packaged in plastic bags, then incubated at 6°C for 7 days. After the

incubation period, fruit were further incubated at 18°C for other 7 days. For each treatment, the disease

incidence, evaluated as occurrence of disease symptoms (%), was recorded at 7, 10 and 14 days post the last

treatment session.

The above described experimental scheme was replicated for orange, lemon and clementine tangerine fruit.

2.4.2. Fruit inoculated with *P. digitatum* during treatment

The peel of each citrus fruit was wounded by a plastic tip (diameter, 2 mm) at four points along the

equatorial surface. Then, fruit were pre-treated by dipping in P01 (0.1 mL/L) for 30 seconds and rinsing in

sdw for additional 30 seconds, or by a simple dipping in sdw for 30 seconds (control). After the pre-

treatment, each wound was sprayed with 65 µl of RI01, F01, M01 or sdw (controls); then, each wound was

inoculated with 5 µl of a conidial suspension (10^6 conidia/ml) of *P. digitatum* strain P1PP0. Fruit were air

dried and organized in five batches corresponding to the following treatments: i. fruit pre-treated only with

sdw (control 1), ii. fruit pre-treated only with P01 (control 2), iii. fruit pre-treated with P01 and then treated

with RI01, iv. fruit pre-treated with P01 and then treated with F01, v. fruit pre-treated with P01 and then

treated with M01. Each treatment comprised three technical replicates each with five citrus fruit**.** After

drying, fruit from each treatment were packaged in plastic bags and incubated at 6°C for 7 days. After the

incubation period, fruit from any treatment were further incubated at 18°C for other 7 days. For each

treatment, the disease incidence, evaluated as occurrence of disease symptoms (%), was recorded at 7, 10

and 14 days post the last treatment session.

The above-described experimental scheme was replicated for orange, lemon and clementine tangerine fruit.

2.4.3. Fruit inoculated with *P. digitatum* post-treatment

The peel of each citrus fruit was wounded by a plastic tip (diameter, 2 mm) at four points along the

equatorial surface. Then, fruit were pre-treated by dipping in P01 (0.1 mL/L) for 30 seconds and rinsing in

sdw for additional 30 seconds, or by a simple dipping in sdw for 30 seconds (control). After the pre-

treatment, each wound was sprayed with 65 µl of RI01, F01, M01 or sdw (controls); then, each wound of

each fruit was inoculated with 5 µl of a conidial suspension (10^6 conidia/ml) of *P. digitatum* strain P1PP0.

Fruit were air dried and organized in five batches corresponding to the following treatments: i. fruit pre-

treated only with sdw (control 1), ii. fruit pre-treated only with P01 (control 2), iii. fruit pre-treated with P01

and then treated with RI01, iv. fruit pre-treated with P01 and then treated with F01, v. fruit pre-treated with

P01 and then treated with M01. Each treatment had three technical replicates made up of five citrus fruit

each**.** After drying, fruit from each treatment were packaged in plastic bags and incubated at 6°C for 12

hours. After the incubation period, each wound of any fruit was inoculated with 5 µl of a conidial suspension

(10^6 conidia/ml) of *P. digitatum* strain P1PP0. Fruit from all the treatments were packaged again and

incubated at 6°C for additional 7 days. After the incubation period, fruit were further incubated at 18°C for

other 7 days. For each treatment, the disease incidence, evaluated as occurrence of disease symptoms (%),

was recorded at 7, 10 and 14 days post the last treatment session.

The above-described experimental scheme was replicated for orange, lemon and clementine tangerine fruit.

2.5. Tissue sampling for RNA isolation from fruit

For the evaluation of the differential expression of genes involved in the induction of resistance in citrus fruit

(clementine tangerines, oranges and lemons) by the different treatments, a specific test was performed.

Disinfected fruit were individually wounded with a 2 mm plastic tip at eight points on the rind surface in an

equatorial position. The experimental design comprised six treatments as it follows: i. unwounded fruit; fruit

treated with: ii. sdw; iii. P01; iv. RI01; v. F01; or vi. M01. For each treatment, fruit were collected at three

different time points: 0, 24, and 48 h. For each time point, fruit batches comprised three fruit. Fruit of each

treatment were placed in plastic containers and incubated at 18°C for 48 h. At each time point, fragments of

peel (5 mm ×15 mm × 2 mm) were excised from each fruit of any treatment from the treated area. The

excised tissues were rapidly frozen in liquid nitrogen and stored at −80°C until they were used for gene

expression analyses. The experiment was repeated twice with similar results, so only results of the first

experiment are reported.

2.5.1 RNA extraction from fruit peels and synthesis of complementary DNA

Total RNA was extracted from 100 mg of frozen citrus fruit peels (clementine tangerines, oranges and

lemons), which were reduced into a fine powder using liquid nitrogen. The RNeasy Plant Mini Kit (Qiagen,

Venlo, Netherlands) was used for RNA extraction following the manufacturer's protocol. Subsequently, the

extracted RNA underwent treatment with the TURBO DNA-free™ Kit (Invitrogen, Carlsbad, CA, United

States). The concentration of RNA was adjusted to 200 ng/µl, and its integrity was assessed through

denaturing RNA electrophoresis in TAE agarose, as described by Masek et al. (2005). Reverse transcription

was carried out using the High-Capacity cDNA Reverse Transcription Kit (Applied Biosystems™, Foster

City, CA, United States) following the manufacturer's instructions.

2.5.2. Selection of genes and specific primers

A set of distinct primer pairs targeting various genes associated with induced resistance in citrus fruit were

selected from previous studies (La Spada et. al., 2021; Youssef et al., 2014). Details about selected genes and

related primers are reported in Table 2.

Table 2. Gene-specific primers used in quantitative reverse transcription real-time polymerase chain reaction (qRT-PCR).

Target gene (gene symbol)	Genebank accession n.	Gene description	Primers description	Melting temperature (°C)	Amplicone size (bp)
LOC102631140 (Beta tube)	XM_006473602	*Citrus sinensis* beta-tubulin (housekeeping gene)	**Csis_Betatub_For** 5'- TCTGATCTCTGCAACTATGAGTG -3' **Csis_Betatub_Rev** 5'-ATTTGCTGGGTCAGCTCTGG -3'	57.74 60.32	198
LOC18032133 (beta-1,3-glucanase)	XM_024191934	*Citrus sinensis* beta-1,3-glucanase	**Gluc 2R** 5'-ATGGCGTCAAAAAGACTTCG-3' **Gluc 2F** 5'-ATTCGCTTCCTCAACGAAAA-3'	57.03 55.98	164
LOC102620464 (PAL)	XM_006481431	*Citrus sinensis* phenylalanine ammonia-lyase-like	**PAL 1F** 5'-GCTCATGTTTGCCCAATTTT-3' **PAL 2R** 5'-AGAAATTGGAGCTCGGAACA-3'	55.68 57.14	153
gpx1 (POX)	AJ582678.1	*Citrus sinensis* gpx1 gene for phospholipid hydroperoxide glutathione peroxidase	**POX 1R** 5'-TCTGCAAGGGGGTAACAAAC-3' **POX 2F** 5'-GCAAAGGTGGACTTTTTGGA-3'	58.01 56.73	240

2.5.3. Quantitative Real-Time PCR (qRT-PCR) analysis of gene expression

qRT-PCR amplifications were carried out by using the iCycler iQ™ Real-Time PCR Detection System

(QuantGeene 9600, Hangzhou Bioer Technology Co., Ltd., BIOER, 1192 Bin An Rd, Binjiang District,

Hangzhou, 310053, People's Republic of China). Each reaction was carried out in a final volume of 20 μl,

comprising 10 ng of cDNA, 1 μl of 10 μM for each primer, and 10 μl of PowerUp™ SYBR™ Green Master

Mix (2X, Applied Biosystems™, Foster City, CA, United States). For each biological sample collected, qRT-

PCR experiments were performed in triplicate. Thermocycling conditions were 2 min at 50°C (UDG

activation), 2 min at 95°C (Dual-Lock™ DNA polymerase), followed by 40 cycles of two steps: 95°C for 15

s (denaturation) and 59°C (annealing/extension) for 1 min. The quantification of gene expression relative to

the unwounded control sample was carried out by using the $2^{-\Delta\Delta Ct}$ method (Livak and Schmittgen, 2001),

where $\Delta\Delta Ct$ = (Ct of target gene − Ct of reference gene)$_{sample}$ − (Ct of target gene − Ct of reference

gene)$_{calibrator}$ and Ct is the threshold cycle of each transcript, defined as the point at which the amount of

amplified target reaches a fixed threshold above the background fluorescence.

2.6. Statistical analyses

247 Analysis of data was performed by one-way ANOVA followed by Tukey's HSD (Honestly Significant

248 Difference) post hoc test by using the R software. To normalize the distributions, percentage data were

249 transformed into square root values, but untransformed percentages are reported in the respective graphs.

250 Differences at $p \leq 0.05$ were considered significant. The mean ± SD was reported in graphs.

3. Results

3.1. Effectiveness of *Candida oleophila* strain O in controlling citrus green mold

Following sections describe results on *in vivo* tests carried out to evaluate the effectiveness of the *C.*

oleophila strain O in controlling citrus green mold on artificially *P. digitatum*-infected orange, lemon, and

clementine tangerine fruit. In detail, three realistic scenarios of fruit infection were simulated: *P. digitatum*

inoculation i. before treatment, ii. during treatment and iii. 12 hours post-treatment. Additionally, in order to

simulate various stages of the citrus postharvest chain, per each scenario of inoculation the following stages

of storing conditions were assessed: a. 7-day storage under refrigerated conditions (at 6°C), and b. an

additional 7-day storage under shelf-life conditions (at 18°C).

3.1.1. Effectiveness of *Candida oleophila* strain O on clementine tangerines

Results of tests on clementine tangerines are summarized in Fig.1. Seven days after the last treatment

session, none of the clementine tangerine fruit from any treatment or infection scenario showed any

symptoms of disease (Figure 1 - a1, b1, and c1). Ten days after the last treatment session, in any scenario of

infection, clementine tangerines from treatment with M01 and F01 showed the lowest occurrences of disease

symptoms (%), with statistical differences with water treated control; however, in any scenario of infection,

the treatment with M01 determined an occurrence of disease symptoms (%) comparable to that of fruit from

treatment with both RI01 and F01 (Figure 1 – a2, b2, and c2). A similar trend of occurrence of disease

symptoms (%) was also observed fourteen days post the last treatment (Figure 1 – a3, b3, and c3).

Figure 1. Evaluation of occurrence of disease symptoms (%) in clementine tangerine [*Citrus clementina* Hort. ex Tan.,cv. Comune]
fruit in three scenarios of infection by *Penicillium digitatum* strain P1PP0: (a) before treatment; (b) during treatment and (c) 12 hours
post-treatment. For each treatment, the occurrence of disease symptoms (%) was recorded at 7 (1), 10 (2) and 14 (3) days post the last
treatment session. For each time point of measurement, values sharing the same letters are not statistically different according to the
Tukey's HSD (Honestly Significant Difference) test (p ≤ 0.05).
275

3.1.2. Effectiveness of *Candida oleophila* strain O on oranges

Results of tests on oranges are summarized in Fig.2. As observed for clementine tangerines, seven days after

the last treatment no orange fruit from any treatment or infection scenario showed any symptoms of disease

(Figure 2 - a1, b1, and c1).

Ten days after the last treatment, across the three scenarios of infection, treatment with M01 determined

markedly low occurrence of disease symptoms, with mean values comparable to those of treatments with the

fungicide (F01) (Figure 2 – a2, b2, and c2); however, except for the scenario with infection during treatment

(Figure 2 - b2), in which M01 determined a statistical reduction of disease severity comparable just to the

fungicide (F01), when pathogen's infection occurred both before and after treatment, M01 determined an

occurrence of disease symptoms statistically similar to that of treatment with RI01 (Figure 2 a2 and c2).

After fourteen days, across the three scenarios of infection, treatment with M01 determined an occurrence of

disease symptoms (%) significantly lower than untreated control and treatments with P01 and R01 (Figure 2

– a3, b3 and c3); additionally, in the scenario with infection before the treatment (Figure 2 – a3), the

application of M01 determined a value of occurrence of disease symptoms as significant as that determined

by treatment with F01.

Figure 2. Evaluation of occurrence of disease symptoms (%) in orange [*Citrus x sinensis* (L.) Osbeck cv. Ovale] fruit in three
scenarios of infection by *Penicillium digitatum* strain P1PP0: (a) before treatment; (b) during treatment and (c) 12 hours post-
treatment. For each treatment, the occurrence of disease symptoms (%) was recorded at 7 (1), 10 (2) and 14 (3) days post the last
treatment session. For each time point of measurement, values sharing the same letters are not statistically different according to the
Tukey's HSD (Honestly Significant Difference) test ($p \leq 0.05$).

3.1.3. Effectiveness of *Candida oleophila* strain O on lemons

Results of tests on lemons are summarized in Fig.3. As observed previously for clementine tangerine and

orange fruit seven days after the last treatment, no lemon fruit from any treatment or infection scenario

showed any symptoms of disease (Figure 3 - a1, b1, and c1). Ten days after the last treatment, fruit treated

with M01 exhibited a significant low occurrence of disease symptoms, with mean values comparable to

those of fruit treated with the fungicide (F01) in scenarios of both infection before and during treatment

(Figure 3 – a2 and b2); additionally, treatment with M01 resulted in the lowest occurrence of disease

symptoms in the post-treatment infection scenario (Figure 3 – c2). After fourteen days of incubation, when

fruit were inoculated before treatment (Figure 3 – a3), the effectiveness of M01 was quite low, with a value

of occurrence of disease symptom comparable to that determined by RI01. In contrast, M01 demonstrated

strong effectiveness in scenarios with infection both during and after treatment. Specifically, treatment with

M01 resulted in the lowest occurrence of disease symptoms in the scenario of infection during treatment

(Figure 3 – b3). Additionally, a significant low occurrence of disease symptoms by treatment with M01 was

recorded in the scenario with infection after treatment, with a mean value comparable to that of treatment

with fungicide (F01) (Figure 3 – c3).

Figure 3. Evaluation of occurrence of disease symptoms (%) in lemon [*Citrus x limon* (L.) Burm. f., cv. Femminello] fruits in three scenarios of infection by *Penicillium digitatum* strain P1PP0: (a) before treatment; (b) during treatment and (c) 12 hours post-treatment. For each treatment, the occurrence of disease symptoms (%) was recorded at 7 (1), 10 (2) and 14 (3) days post the last treatment session. For each time point of measurement, values sharing the same letters are not statistically different according to the Tukey's HSD (Honestly Significant Difference) test (p ≤ 0.05).

3.2. Regulation of induced resistance-related genes in *C. oleophila*-treated citrus fruit

Following paragraph describes the expression levels of the plant defense-related genes, β-1,3-glucanase-,

phenylalanine ammonia-lyase (PAL)-, and peroxidase (PEROX)-encoding genes measured in peels of

clementine tangerine, orange, and lemon fruit at 0-, 24- and 48-hours post after the treatment with M01 and

the other three products (Table 1).

Regarding clementine tangerine fruit, upregulation in the levels of β-1,3-glucanase-, PAL-, and PEROX-

encoding genes was observed at all time points in peels treated with M01 (Figure 4). Specifically, the

treatment with M01 induced the strongest upregulations of the β-1,3-glucanase-encoding gene at 0- and 48-

hours post-treatment, while the upregulation recorded at 24 hours was similar to that induced by treatment

with F01 (Figure 4a). Concerning the PAL-encoding gene, treatment with M01 showed an increasing trend

of upregulation within the first 24 hours; at 48 hours, treatment with M01 still resulted in the strongest

upregulation, albeit with a mean value lower than that observed at 24 hours post-treatment (Figure 4b). An

increasing trend of expression level was also determined by treatment with M01 towards the PEROX-

encoding gene, with the strongest upregulation recorded at 48 hours post-treatment (Figure 4c).

Figure 4. Differences in the expression levels of β-1,3-glucanases-(a), PAL-(b), and PEROX-(c) encoding genes in clementine tangerine fruits (*Citrus clementina* Hort. ex Tan., cv. Comune), wounded and treated with sterile distilled water (controls) or test substances P01 (a sanitization product containing hydrogen peroxide and acetic acid) at 0.1 mL/L , RI01 (a potassium phosphonate formulation) at the dosage of 0.25%, F01 (emulsifiable 50% w/v concentrate formula of imazalil fungicide) at 1.000 ppm (a.i.) and M01 (*Candida oleophila* strain O) as such; at 0, 24, and 48 h after treatment. Values sharing the same letters are not statistically different according to the Tukey's HSD (Honestly Significant Difference) test (p ≤ 0.05).

331 With reference to orange fruit, no up-regulation for any gene was induced by any treatment at 0 hours after

332 treatment (Figure 5a, b and c). At 24- and 48- hours post-treatment, an increasing trend of up-regulation was

333 observed for both β-1,3-glucanase- and PAL-encoding genes in fruit treated with F01 and M01, with the

334 highest values in this latter treatment (Figure 5a and b). Finally, the PEROX-encoding gene was up-regulated

335 just in the treatment with M01 and exclusively at 48 hours post treatment (Figure 5c).

Figure 5. Differences in the expression levels of β-1,3-glucanases-(a), PAL-(b), and PEROX-(c) encoding genes in orange fruit
(*Citrus x sinensis*) cv. Ovale, wounded and treated with sterile distilled water (controls) or test substances P01 (a sanitization product
containing hydrogen peroxide and acetic acid) at 0.1 mL/L, RI01 (a potassium phosphonate formulation) at the dosage of 0.25%, F01
(emulsifiable 50% w/v concentrate formula of imazalil fungicide) at 1.000 ppm (a.i.) and M01 (*Candida oleophila* strain O) as such;
at 0, 24, and 48 h after treatment. Values sharing the same letters are not statistically different according to the Tukey's HSD
(Honestly Significant Difference) test (p ≤ 0.05).

344 Regarding lemon fruit, no up-regulation for any gene was induced by treatments until 48 hours (Figure 6a, b

345 and c). At this time point, both the β-1,3-glucanase- and PEROX-encoding genes were up-regulated only in

346 the treatment with M01 (Figure 6a and c). Finally, except for the control, up-regulations of PAL-encoding

347 gene were recorded in all treatments, with the highest value in fruit treated with M01 (Figure 6b).

Figure 6. Differences in the expression levels of β-1,3-glucanases-(a), PAL-(b), and PEROX-(c) encoding genes in lemon fruits (*Citrus x limon* L. *Burm. f.,* cv. Femminello), wounded and treated with sterile distilled water (controls) or test substances P01 (a sanitization product containing hydrogen peroxide and acetic acid) at 0.1 mL/L , RI01 (a potassium phosphonate formulation) at the dosage of 0.25%, F01 (emulsifiable 50% w/v concentrate formula of imazalil fungicide) at 1.000 ppm (a.i.) and M01 (*Candida oleophila* strain O) as such; at 0, 24, and 48 h after treatment. Values sharing the same letters are not statistically different according to the Tukey's HSD (Honestly Significant Difference) test (p ≤ 0.05).

4. Discussion

Achieving optimal plant disease control without relying on conventional pesticides is currently one of the

most stimulating challenges for the modern Plant Pathology (Bhatta, 2022; Dukare et al., 2022; Lahlali et al.,

2022; Nega, 2014; Pal and Gardener, 2006). This ambitious goal holds significant relevance in managing

post-harvest decays of fruit from the citrus supply chain. The high incidence of product losses due to molds

caused by *Penicillium* species makes it imperative to adopt carefully pondered and evaluated strategies of

control. Therefore, with the awareness that means of control safe for human health as effective as pesticides

are unlikely to be identified, scientific community has the duty to undertake the study of the efficacy of new

plant disease control means through evaluative protocols that realistically simulate the phases of the complex

framework of post-harvest of citrus fruit (Zhang et al., 2020). Hence, the first step of this study was to

examine different scenarios of *P. digitatum* infections in clementine tangerine, orange, and lemon fruit to

evaluate the potential of *C. oleophila* strain O, a specimen well-known for its high and reliable antagonistic

activity against post-harvest wound pathogens from various crops (Ballet et al., 2016; Bastiaanse et al., 2010;

Dukare et al., 2019; Jijakli and Lepoivre, 1993; Lahlali and Jijakli, 2009; Lahlali et al., 2004, 2011;

Lassois et al., 2008), in reducing the occurrence of green mold throughout the main stages of the post-harvest

supply chain. Previous studies already investigated some aspects related to the efficacy of *C. oleophila* in

controlling diseases of citrus fruit incited by *Penicillium* spp. Brown et al. (2000) evidenced limits of *C.*

oleophila strain O in controlling *Penicillum* molds of citrus fruit, demonstrating that the success in

controlling fruit decay depends on how quickly and well the yeast colonizes injuries to the fruit surface,

including minor injuries involving only oil vesicles. Laahlali et al. (2004) highlighted the good efficacy of *C.*

oleophila strain O in reducing mold severity in 'clementine' and 'valencia-late' orange fruit inoculated with *P.*

digitatum and *P. italicum* after 7 days of incubation at 24°C. A recent study of Hammami et al. (2022) tested

the efficiency of *C. oleophila* strain L12 in reducing severity of molds by *Penicillium* spp. in orange fruit

subjected to different incubation conditions, including both a long time cold storage (30 days at 4°C) and a

particular storage condition consisting of 7 days at room temperature and 30 days at 4 °C. Although these

studies have provided valuable information about the potential of *C. oleopihla* in controlling post-harvest

decays of citrus fruit, the adopted *in vivo* protocols were based on conditions, such us timing and temperature

of incubation, that are far from the reality of the post-harvest supply chain of citrus fruit. Indeed, in the

majority of post-harvest chains of citrus fruit, the operational process typically involves bringing fruit from

the field into the packinghouses (within four hours from the picking), where blemished or damaged fruit are

discarded. Then, marketable fruit undergo a series of steps including washing with detergent and/or

fungicides, rinsing with fresh water, waxing (optional), drying, sorting, stamping, sizing, and packing.

Finally, fruit are either temporarily held (maximum 12 hours) in a cold store (temperature in the range 4-8°C,

depending on the kind of citrus fruit) or directly transported to the market (Malik et al., 2004). The

transportation is one of the most critical steps of the post-harvest supply chain of citrus fruit (Cao et al.,

2019; Chace et al., 1970; Defraeye et al., 2015). It often exposes fruit to mechanical damages due to

improper loading/off-loading procedures, reduced sanitation conditions, and temperatures far from an ideal

continuous cold storage environment effective in preserving fruit from the development of *Penicillium* molds

(Zacarias et al., 2020). These aspects, together with the rapid progression of these fruit molds and the

necessity to minimize post-harvest applications of fungicides, have led citrus producers to realize that the

best strategy for reducing product losses is to shorten as much as possible the overall timing of the supply

chain. Indeed, nowadays the transportation of fruit typically takes about 2-4 hours for shipments destined for

local/regional markets, and approximately 2-7 days for shipments headed to national/continental markets.

Once received by the retailer, fruit are displayed on the store shelves at a temperature of shelf-life (16-18°C)

and usually sold by no more than seven days from the delivery. Therefore, the duration of a post-harvest

supply chain for citrus fruit in a market at a national/continental scale ranges from approximately 10 to 14

days from fruit picking.

In this study, the assessment of tests in different kinds of citrus fruit (clementine tangerines, oranges and

lemons) inoculated with *P. digitatum* before, during, or after treatment with *C. oleophila*, provided findings

that can be directly related to the efficiency of this biological control agent in typical scenarios of infections

of the post-harvest chain of citrus fruit, such as picking, processing in the packinghouse, and loading of fruit

for transportation. Additionally, the incubation of inoculated fruit for a period of 7 days at 6°C and for further

7 days at 18°C, simulated the cold storage shipment and the shelf-life phases typical of a market at

national/continental scale. Therefore, this study offered the unprecedented opportunity to rationally evaluate

the potential of *C. oleophila* for controlling green mold by *P. digitatum* in a real supply chain. In accordance

with the physiology of *P. digitatum*, whose cardinal temperatures for growth fall within the range of 6-37°C,

no citrus fruit from any treatment or infection scenario exhibited any symptoms of green mold after 7 days of

incubation at 6°C. As expected, first symptoms of green mold started to occur at the day 10 day of incubation

post-treatment, namely at the third day at shelf-life temperature (18°C); at this time point, the treatment with

C. oleophila reduced strongly the occurrence of symptoms of green mold in all kinds of tested fruit

(clementine tangerines, oranges and lemons) and scenario of infection, with values significantly lowers than

untreated controls and often comparable to those recorded in fruit treated with the fungicide IMZ. This

marked effectiveness of *C. oleophila* persisted until the 14th day of post-treatment incubation, corresponding

to the seventh day at shelf-life temperature (18°C), with values of occurrence of disease symptoms still

comparable to those attained with IMZ treatments. The significant reduction in the occurrence of green mold

symptoms after treatment with *C. oleophila*, particularly evident at shelf-life temperatures, underscores its

efficacy in mitigating post-harvest fungal infections. This sustained efficacy is particularly noteworthy given

the challenges posed by prolonged storage and transportation periods within the post-harvest supply chain.

Additionally, the comparability of results to those achieved with conventional fungicide treatments, such as

IMZ, highlight the practical potential of *C. oleophila* as an alternative or complementary approach for the

management of green molds by *P. digitatum* in the post-harvest chain of citrus fruit at national/continental

scale.

This study also provided a significant advancement in understanding the potential of *C. oleophila* to trigger

the natural defensive response of various citrus fruit (clementine tangerines, oranges, and lemons). To this

purpose, the modulation of the transcriptional levels of genes encoding β-1,3-glucanases, phenylalanine

ammonia-lyase, and peroxidase in peels of clementine tangerine, orange and lemon fruit was examined at

specific time-points after treatment with *C. oleophila* or comparative substances (potassium phosphonate;

IMZ). Several studies supported the involvement of metabolism of β-1,3-glucanases, phenylalanine

ammonia-lyase and peroxidase enzymes in the activation of defense response of citrus fruit. The β-1,3-

glucanases, an enzyme contributing significantly to the degradation of the fungal cell wall (La Spada et al.

2020, 2021; Youssef et al. 2014), is considered a real molecular marker for postharvest physiological

disorders of citrus fruit (Sanchez-Ballesta et al., 2008). The high concentration of phenylalanine ammonia-

lyase in citrus peels has been positively associated with the triggering of resistance mechanisms against

infections by *P. digitatum*; these mechanisms include the synthesis of antifungal metabolites, such as the

phytoalexins scoparone and scopoletin (Ballester et al., 2010; Droby et al., 2002; Kim et al., 1991; La Spada

et al., 2021; Youssef et al., 2014). The antioxidant enzyme peroxidase participates in the formation of auxin

metabolism, lignin, and suberin, cross-linking cell wall components, synthesizing phytoalexins, and

metabolizing reactive oxygen species (ROS) (Almagro et al., 2009). In this study, *C. oleophila* was the

treatment that elicited the strongest up-regulations in all kinds of test fruit. Specifically, clementine

tangerines resulted the most responsive fruit, with marked up-regulations of the three genes in the treatment

with *C. oleophila* at each time-point of measurement. Orange fruit were less responsive than clementine

tangerines; for these fruit, treatment with *C. oleophila* stimulated the up-regulation of β-1,3-glucanases- and

phenylalanine ammonia-lyase-encoding genes starting from 24 hours post-treatment, with an increasing

trend until 48 hours; In contrast, peroxidase up-regulation in orange fruit occurred only at 48 hours post-

treatment. Finally, the defensive response of lemon fruit was significantly delayed, with up-regulations of the

three genes recorded only after 48 hours. Overall, results from gene expression evidenced that *C. oleophila*

was the most efficient treatment in the elicitation of defense response of fruit peels. This high efficiency of

C. oleophila in activating the studied metabolic pathways is consistent with previous literature. Indeed,

Droby et al. (2002) evidenced that the treatment of grapefruit with the strain O of *C. oleophila* promoted the

biosynthesis of β-1,3-glucanases and phenylalanine ammonia-lyase enzymes, and that their accumulation in

fruit peels showed an increasing trend over time. Another study found that *C. oleophila* applications are also

able to promote the gradual accumulation over time of peroxidase enzymes (Zheng et al., 2021).

In conclusion, this study addressed the pressing challenge of achieving optimal plant disease control in citrus

fruit without relying on conventional pesticides, emphasizing the significance of managing post-harvest

decays in the citrus supply chain. The study focused on evaluating the efficacy of the biological control agent

C. oleophila strain O in controlling green mold caused by *P. digitatum* throughout various stages of the post-

harvest supply chain. Results indicated that *C. oleophila* exhibited significant efficacy in reducing the

occurrence of green mold symptoms, particularly under conditions simulating the post-harvest supply chain

of citrus fruit. This efficacy was evident even at shelf-life temperatures, highlighting the potential of *C.*

oleophila as a practical alternative or complementary approach to conventional fungicides. Moreover, the

study contributed valuable insights into the molecular aspects of the defensive response of citrus fruit to *C.*

oleophila treatment. The up-regulation of genes encoding β-1,3-glucanases, phenylalanine ammonia-lyase,

and peroxidase in fruit peels demonstrated the activation of defense mechanisms. Clementine tangerines

displayed the most pronounced response, while oranges and lemons exhibited delayed but significant up-

regulation of these genes. Overall, the findings supported the potential of *C. oleophila* as biological control

agent not only as pathogen antagonist, but also as elicitor of key defense pathways in citrus fruit. The results

highlighted the usefulness of this biological control agent and its comparability with conventional fungicides,

emphasizing its importance in the context of a post-harvest citrus industry on a national/continental scale.

Funding
474
This study was supported by the project "Smart and innovative packaging, postharvest rot management, and
shipping of organic citrus fruit (BiOrangePack)" under Partnership for Research and Innovation in the
Mediterranean Area (PRIMA) – H2020 (E69C20000130001), the he "Italie–Tunisie Cooperation Program
2014–2020" project "PROMETEO «Un village transfrontalierpour protéger les cultures arboricoles
méditerranéennes enpartageant les connaissances» cod. C-5–2.1–36, CUP 453E25F2100118000 and the
European Union (NextGeneration EU), through the MUR-PNRR project SAMOTHRACE (ECS00000022).
Ermes Ivan Rovetto was supported by a Ph.D. fellowship funded by University of Catania (cycle XXXVI).

483
484
CRediT authorship contribution statement
Ermes Ivan ROVETTO: Writing – original draft, Visualization, Software, Investigation, Formal analysis, Data curation. **Federico LA SPADA**: Writing – review & editing, Writing – original draft, Visualization, Supervision, Methodology, Investigation, Data curation, Conceptualization. **Soumia EL BOUMLASY**: Writing – review & editing, Investigation, Data curation. **Sebastiano CONTI TAGUALI**: Writing – review & editing, Investigation, Data curation. **Mario RIOLO**: Writing – review & editing, Investigation, Data curation. **Antonella PANE**: Writing – review & editing, Validation, Supervision, Resources, Project administration, Funding acquisition. **Santa Olga CACCIOLA**: Writing – review & editing, Validation, Supervision, Resources, Project administration, Funding acquisition, Conceptualization.
494

References

Abo-Elnaga, H. I. G. 2013. Control of green mould of citrus by using *Trichoderma harzianum*, humic acid or garlic. *Archives of Phytopathology and Plant Protection* 46:1118-1126. https://doi.org/10.1080/03235408.2012.760226

Afek, U., Orenstein, J., Carmeli, S., Rodov, V., and Joseph, M. B. 1999. Umbelliferone, a phytoalexin associated with resistance of immature Marsh grapefruit to *Penicillium digitatum*. *Phytochemistry* 50, 1129–1132. https://doi.org/10.1016/S0031-9422(98)00671-2

Almagro, L., Gómez Ros, L. V., Belchi-Navarro, S., Bru, R., Ros Barceló, A., and Pedreño, M. A. 2009. Class III peroxidases in plant defence reactions. *Journal of Experimental Botany* 60(2), 377-390. https://doi.org/10.1093/jxb/ern277

Amiri, A., and Bompeix, G. 2011. Control of *Penicillium expansum* with potassium phosphite and heat treatment. *Crop Protection* 30(2), 222-227. https://doi.org/10.1016/j.cropro.2010.10.010

Arcas, M. C., Botía, J. M., Ortuño, A. M., and Del Río, J. A. 2000. UV irradiation alters the levels of flavonoids involved in the defence mechanism of *Citrus aurantium* fruit against *Penicillium digitatum*. *European Journal of Plant Pathology* 106, 617-622. https://doi.org/10.1023/A%3A1008704102446

Arslan, U. 2015. Evaluation of antifungal activity of mono and dipotassium phosphates against phytopathogenic fungi. *Fresenius Environmental Bulletin* 24(3):810–816 https://doi.org/10.1080/17429145.2017.1397205

Balasubramanian, V., Vashisht, D., Cletus, J., and Sakthivel, N. 2012. Plant β-1, 3-glucanases: their biological functions and transgenic expression against phytopathogenic fungi. *Biotechnology letters* 34, 1983-1990. https://doi.org/10.1007/s10529-012-1012-6

Ballester, A. R., Lafuente, M. T., and González-Candelas, L. 2006. Spatial study of antioxidant enzymes, peroxidase and phenylalanine ammonia-lyase in the citrus fruit–*Penicillium digitatum* interaction. *Postharvest Biology and Technology* 39(2), 115-124. https://doi.org/10.1016/j.postharvbio.2005.10.002

Ballester, A. R., Izquierdo, A., Lafuente, M. T., and González-Candelas, L. 2010. Biochemical and molecular characterization of induced resistance against *Penicillium digitatum* in citrus fruit. *Postharvest Biology and Technology* 56, 31–38. https://doi.org/10.1016/j.postharvbio.2009.10.002

Ballet, N., Souche, J.L. and Vandekerckove, P. 2016. Efficacy of *Candida oleophila*, strain O, in preventing postharvest diseases of fruit. *Acta Horticulturae* 1144, 105-112 https://doi.org/10.17660/ActaHortic.2016.1144.15

Bastiaanse, H., De Bellaire, L. D. L., Lassois, L., Misson, C., and Jijakli, M. H. 2010. Integrated control of crown rot of banana with *Candida oleophila* strain O, calcium chloride and modified atmosphere packaging. *Biological Control* 53(1), 100-107. https://doi/10.1016/j.biocontrol.2009.10.012

Ben-Yehoshua, S., Rodov, V., Nafussi, B., Feng, X., Yen, J., Koltai, T., and Nelkenbaum, U. 2008. Involvement of limonene hydroperoxides formed after oil gland injury in the induction of defense response against *Penicillium digitatum* in lemon fruit. *Journal of Agricultural and Food Chemistry* 56(6), 1889-1895. https://doi.org/10.1021/jf072825z

Bhatta, U. K. 2022. Alternative management approaches of citrus diseases caused by *Penicillium digitatum* (green mold) and *Penicillium italicum* (blue mold). *Frontiers in Plant Science* 12, 833328. https://doi.org/doi: 10.3389/fpls.2021.833328.

Boava, L. P., Cristofani-Yaly, M., Stuart, R. M., and Machado, M. A. 2011. Expression of defense-related genes in response to mechanical wounding and *Phytophthora parasitica* infection in *Poncirus trifoliata* and *Citrus sunki*. *Physiological and Molecular Plant Pathology* 76(2), 119-125. http://dx.doi.org/10.1016/j.pmpp.2011.07.004

Boller, T. 1995. Chemoperception of microbial signals in plant cells. *Annual Review of Plant Physiology and Plant Molecular Biology* 46, 189–214. http://dx.doi.org/10.1146/annurev.pp.46.060195.001201

Brown, G. E., Davis, C., and Chambers, M. 2000. Control of citrus green mold with Aspire is impacted by the type of injury. *Postharvest biology and technology* 18(1), 57-65. https://doi.org/10.1016/S0925-5214(99)00060-5

Bus, V. G. 1992. ED50 levels of *Penicillium digitatum* and *P. italicum* with reduced sensitivity to thiabendazole, benomyl and imazalil. *Postharvest Biology and Technology* 1(4), 305-315. https://doi.org/10.1016/0925-5214(92)90033-L

Bus, V. G., Bongers, A. J., and Risse, L. A. 1991. Occurrence of *Penicillium digitatum* and *P. italicum* resistant to benomyl, thiabendazole and imazalil on citrus fruit from different geographic origins. *Plant Disease* 75(11), 1098-1100. https://doi.org/10.1094/PD-75-1098

Cao, J., Wang, C., Xu, S., Chen, Y., Wang, Y., Li, X., and Sun, C. 2019. The effects of transportation temperature on the decay rate and quality of postharvest Ponkan (*Citrus reticulata* Blanco) fruit in different storage periods. *Scientia Horticulturae* 247, 42-48. https://doi.org/10.1016/j.scienta.2018.12.009

Cerioni, L., Sepulveda, M., Rubio-Ames, Z., Volentini, S.,I., Rodríguez-Montelongo, L. , Smilanick, J. L., Ramallo, J., and Rapisarda, V. A. 2013. Control of lemon postharvest diseases by low-toxicity salts combined with hydrogen peroxide and heat. *Postharvest Biology and Technology* 83, 17-21. https://doi.org/10.1016/j.postharvbio.2013.03.002.

Chace, W. G., Smoot, J. J., and Cubbedge, R. H. 1970. Storage and transportation of Florida citrus fruit. *Storage and transportation of Florida citrus fruit.*

Cluzet, S., Torregrosa, C., Jacquet, C., Lafitte, C., Fournier, J., Mercier, L., ... and Dumas, B. 2004. Gene expression profiling and protection of *Medicago truncatula* against a fungal infection in response to an elicitor from green algae *Ulva* spp. *Plant, Cell & Environment* 27(7), 917-928. https://doi.org/10.1111/j.1365-3040.2004.01197.x

Costa, J. H., Bazioli, J. M., de Moraes Pontes, J. G., and Fill, T. P. 2019. *Penicillium digitatum* infection mechanisms in citrus: What do we know so far? *Fungal Biology* 123(8):584-593. https://doi.org/10.1016/J.FUNBIO.2019.05.004

Côté, F., Ham, K. S., Hahn, M. G., and Bergmann, C. W. 1998. Oligosaccharide elicitors in host-pathogen interactions: generation, perception, and signal transduction. *Subcellular Biochemistry* 29, 385–432. doi: 10.1007/978-1-4899-1707-2_13. https://doi.org/10.1007/978-1-4899-1707-2_13

Davé, B., Sales, M., and Walia, M. 1989. Resistance of different strains of *Penicillium digitatum* to imazalil treatment in California citrus packinghouses. *Proceedings of the Florida State Horticultural Society* 102:178–179

Defraeye, T., Cronje, P., Verboven, P., Opara, U. L., and Nicolai, B. 2015. Exploring ambient loading of citrus fruit into reefer containers for cooling during marine transport using computational fluid dynamics. *Postharvest Biology and Technology* 108, 91-101. https://doi.org/10.1016/j.postharvbio.2015.06.004

Del Río, J. A., Arcas, M. C., Benavente-García, O., and Ortuño, A. 1998. Citrus polymethoxylated flavones can confer resistance against *Phytophthora citrophthora*, *Penicillium digitatum*, and *Geotrichum* species. *Journal of Agricultural and Food Chemistry* 46, 4423–4428. https://doi.org/10.1021/jf980229m

Droby, S., Hofstein, R., Wilson, C.L., Wisniewski, M., Fridlender, B., Cohen, L., Weiss, B., Daus, A., Timar, D., and Chalutz, E. 1993. Pilot testing of *Pichia guilliermondii*: A biocontrol agent of postharvest diseases of citrus fruit. *Biological Control* 3(1), 47-52. https://doi.org/10.1006/bcon.1993.1008

Droby, S., Vinokur, V., Weiss, B., Cohen, L., Daus, A., Goldschmidt, E. E,, and Porat, R. 2002 Induction of resistance to *Penicillium digitatum* in grapefruit by the yeast biocontrol agent *Candida oleophila*. *Phytopathology* 92(4):393-399. https://doi.org/10.1094/PHYTO.2002.92.4.393

Droby, S., Wisniewski, M., Mascarisin, D., and Wilson, C. 2009. Twenty years of postharvest biocontrol research: is it time for a new paradigm?. *Postharvest Biology and Technology* 52, 137-145. https://doe.org/10.1016/j.postharvbio.2008.11.009

Dukare, A. S., Paul, S., Nambi, V. E., Gupta, R. K., Singh, R., Sharma, K., and Vishwakarma, R. K. 2019. Exploitation of microbial antagonists for the control of postharvest diseases of fruit: a review. *Critical Reviews in Food Science and Nutrition* 59(9), 1498-1513. https://doi.org/10.1080/10408398.2017.1417235

Dukare, A. S., Singh, R. K., Jangra, R. K., and Bhushan, B. 2022. Non-fungicides-based promising technologies for managing post-production penicillium induced spoilage in horticultural commodities: a comprehensive review. *Food Reviews International* 38(3), 227-267. https://doi.org/10.1080/87559129.2020.1727497

Du Jardin, P. 2015. Plant biostimulants: Definition, concept, main categories and regulation. *Scientia Horticulturae* 196:3-14. https://doi.org/10.1016/J.SCIENTA.2015.09.021

EFSA 2010. Conclusion on the peer review of the pesticide risk assessment of the active substance imazalil. *EFSA Journal* 8(4):1526.

El Boumlasy, S., La Spada, F., Tuccitto, N., Marletta, G., Luz Mínguez, C., Meca, G., Rovetto, E. I., Pane, A., Debdoubi, A., and Cacciola, S. O. 2021. Inhibitory activity of shrimp waste extracts on fungal and oomycete plant pathogens. *Plants* 10(11):2452. https://doi.org/10.3390/PLANTS10112452

El-Otmani, M., Ait-Oubahou, A., and Zacarías, L. 2011. Citrus spp.: Orange, mandarin, tangerine, clementine, grapefruit, pomelo, lemon and lime. In Postharvest biology and technology of tropical and subtropical fruit. *Woodhead Publishing* 437-516. https://doi.org/10.1533/9780857092762.437

Elsherbiny, E. A., Taher, M. A., Abd El Aziz, M. H., and Mohamed, S. Y. 2021. Action mechanisms and biocontrol of *Purpureocillium lilacinum* against green mould caused by *Penicillium digitatum* in orange fruit. *Journal of Applied Microbiology* 131(3), 1378-1390. https://doi.org/10.1111/jam.15016

European Parliament (2019) Directive 2019/1009/EC of the European Parliament and the Council of 5 June 2019, laying down rules on the making available on the market of EU fertilising products and amending Regulations

(EC) No 1069/2009 and (EC) No 1107/2009 and repealing Regulation (EC) No 2003/2003.
http://data.europa.eu/eli/reg/2019/1009/oj

Fallanaj, F., Ippolito, A., Ligorio, A., Garganese, F., Zavanella, C., and Sanzani, S. M. 2016. Electrolyzed sodium
bicarbonate inhibits *Penicillium digitatum* and induces defence responses against green mould in citrus fruit.
Postharvest Biology and Technology 115, 18-29. http://dx.doi.org/10.1016/j.postharvbio.2015.12.009

Fenta, L., Mekonnen, H., and Gashaw, T. 2019. Biocontrol potential of *Trichoderma* and yeast against post harvest fruit
fungal diseases : A review. *World News of Natural Sciences* 27:153-173.
http://www.doi.org/10.54574/RJPP.15.01

Gikas, G. D., Parlakidis, P., Mavropoulos, T., and Vryzas, Z. 2022. Particularities of fungicides and factors affecting
their fate and removal efficacy: A review. *Sustainability* 14(7):4056. https://doi.org/10.3390/su14074056

Gómez-Merino, F. C., and Trejo-Téllez, L. I. 2015. Biostimulant activity of phosphite in horticulture. *Scientia*
Horticulturae 196, 82–90. doi: 10.1016/j.scienta.2015.09.035

Holmes, G. J., and Eckert, J. W. 1999. Sensitivity of *Penicillium digitatum* and *P. italicum* to postharvest citrus
fungicides in California. *Phytopathology* 89(9):716-721. https://doi.org/10.1094/PHYTO.1999.89.9.716

Ismail, M., and Zhang, J. 2004. Post-harvest citrus diseases and their control. *Outlooks on Pest Management* 15:29–35.
https://doi.org/10.1564/15feb12

Janisiewicz, W.J., and Korsten,L. 2002. Biological control of post-harvest diseases of fruit. *Annual Review of*
Phytopathology 40, 411–41. doi: https://doi.org/10.1146/annurev.phyto.40.120401.130158

Kanan, G. J., and Al-Najar, R. A. 2008. In vitro antifungal activities of various plant crude extracts and fractions against
citrus post-harvest disease agent *Penicillium digitatum. Jordan Journal of Biological Sciences* 1:89-99.
https://doi.org/10.2478/v10045-009-0054-9

Kanetis, L., Förster, H., and Adaskaveg, J. E. 2007. Comparative efficacy of the new postharvest fungicides
azoxystrobin, fludioxonil, and pyrimethanil for managing citrus Green Mold. *Plant Disease* 91(11):1502–1511.
https://doi.org/10.1094/PDIS-91-11-1502

Ke, D., and Saltveit, M.E., 1989. Wounded-induced ethylene production, phenolic metabolism and susceptibility to
russet spotting in iceberg lettuce. *Physiologia Plantarum* 76, 412–418. https://doi.org/10.1111/j.1399-
3054.1989.tb06212.x

Kim, J. J., Ben-Yehoshua, S., Shapiro, B., Henis, Y., and Carmeli, S. 1991. Accumulation of scoparone in heat-treated
lemon fruit inoculated with *Penicillium digitatum* Sacc. *Plant Physiology* 97(3), 880-885.
https://doi.org/10.1104%2Fpp.97.3.880

Kowalska, J., Krzymińska, J., and Tyburski, J. 2022. Yeasts as a potential biological agent in plant disease protection
and yield improvement—A short review. *Agriculture* 12(9), 1404. https://doi.org/10.3390/agriculture12091404

Khumalo, G., Goedhals-Gerber, L. L., Cronje, P., and Berry, T. 2021. The non-conformance of in-transit citrus container
shipments to cold protocol markets: A systematic literature review. *Food Control* 125, 107947.
http://dx.doi.org/10.1016/j.foodcont.2021.107947

La Spada, F., Aloi, F., Coniglione, M., Pane, A., and Cacciola, S. O. 2021. Natural biostimulants elicit plant immune
system in an integrated management strategy of the postharvest Green Mold of orange fruit incited by *Penicillium*
digitatum. Frontiers in Plant Science 12:1-16. https://doi.org/10.3389/fpls.2021.684722

La Spada, F., Stracquadanio, C., Riolo, M., Pane, A., and Cacciola, S. O. 2020. Trichoderma counteracts the challenge
of *Phytophthora nicotianae* infections on tomato by modulating plant defense mechanisms and the expression of
crinkler, necrosis-inducing *Phytophthora* Protein 1, and cellulose-binding elicitor lectin pathogenic effectors.
Frontiers in Plant Science 11:1–16. https://doi.org/10.3389/fpls.2020.583539

Lahlali, R., Ezrari, S., Radouane, N., Kenfaoui, J., Esmaeel, Q., El Hamss, H., ... and Barka, E. A. 2022. Biological
control of plant pathogens: A global perspective. *Microorganisms* 10(3), 596.
https://doi.org/10.3390/microorganisms10030596

Lahlali, R., and Jijakli, M. H. 2009. Enhancement of the biocontrol agent *Candida oleophila* (strain O) survival and
control efficiency under extreme conditions of water activity and relative humidity. *Biological Control* 51(3), 403-
408. https://doi.org/10.1016/j.biocontrol.2009.07.014

Lahlali, R., Raffaele, B., and Jijakli, M. H. 2011. UV protectants for *Candida oleophila* (strain O), a biocontrol agent of
postharvest fruit diseases. *Plant Pathology* 60(2), 288-295.https://doi.org/10.1111/j.1365-3059.2010.02368.x

Lahlali, R., Serrhini, M. N., and Jijakli, H. 2004. Efficacy assessment of *Candida oleophila* (strain O) and *Pichia*
anomala (strain K) against major postharvest diseases of citrus fruit in Morocco. *Communications in agricultural*
and applied biological sciences - Ghent University 69(4). https://api.semanticscholar.org/CorpusID:20323063

Lassois, L., de Bellaire, L. M., and Jijakli, H. 2008. Biological control of crown rot of bananas with *Pichia anomala*
strain K and *Candida oleophila* strain O. *Biological Control* 45 (3):410–418.
https://doi.org/10.1016/j.biocontrol.2008.01.013

Li, Y., Xia, M., He P., Yang, Q., Wu, Y., He, P., Ahmed, A., Li, X., Wang, Y., Munir, S., and He, Y. 2022. Developing
Penicillium digitatum management strategies on post-Harvest citrus fruit with metabolic components and
colonization of *Bacillus subtilis* L1-21. *Journal of Fungi (Basel)* 8(1), 80. https://doi.org/ 10.3390/jof8010080.

Liu, J., Sui, Y., Wisniewski, M., Droby, S., and Liu, Y. 2013. Review: Utilization of antagonistic yeasts to manage
postharvest fungal diseases of fruit. *International Journal of Food Microbiology* 167, 153–160.
https://doi.org/10.1016/j.ijfoodmicro.2013.09.004

Livak, K. J., and Schmittgen, T. D. 2001. Analysis of relative gene expression data using real-time quantitative PCR and
the 2-ΔΔCT method. *Methods* 25, 402–408. https://doi.org/10.1006/meth.2001.1262

Malik, A. U., Singh, Z., Saleem, B. A., and Khan, M. N. 2004. Postharvest handling of fresh citrus fruit: An overview.
In Proceedings of International Conference on Citriculture (pp. 223-230). *Pakistan: The Horticultural Foundation*
https://api.semanticscholar.org/CorpusID:28394380

Manghi, M. C., Masiol, M., Calzavara, R., Graziano, P.L., Peruzzi, E., and Pavoni, B. 2021. The use of phosphonates in
agriculture. Chemical, biological properties and legislative issues. *Chemosphere* 283,131187.
https://doi.org/10.1016/j.chemosphere.2021.131187.

Masek, T., Vopalensky, V., Suchomelova, P., and Pospisek, M. 2005. Denaturing RNA electrophoresis in TAE agarose
gels. *Analytical Biochemistry* 336, 46–50. doi: 10.1016/J.AB.2004.09.010

Miedes, E., Vanholme, R., Boerjan, W., and Molina, A. 2014. The role of the secondary cell wall in plant resistance to
pathogens. *Frontiers in plant science* 5, 358. https://doi.org/10.3389/fpls.2014.00358

Moraes Bazioli, J. M., Belinato, J. R., Costa, J. H., Akiyama, D. Y., de Moraes Pontes J. G., Kupper, K. C., Augusto, F.,
de Carvalho, J. E., and Fill, T.P. 2019. Biological control of citrus postharvest phytopathogens. *Toxins* 11(8):460.
https://doi.org/10.3390/toxins11080460

Nega, A. 2014. Review on concepts in biological control of plant pathogens. J*ournal of Biology, Agriculture and
Healthcare* 4(27), 33-54. https://api.semanticscholar.org/CorpusID:54896133

Odjakova, M., and Hadjiivanova, C. 2001. The complexity of pathogen defense in plants. *Bulgarian Journal of Plant
Physiology* 27(1-2), 101-109. https://www.researchgate.net/publication/253807226

Oztekin, S., Dikmetas, D. N., Devecioglu, D., Acar, E. G., and Karbancioglu-Guler, F. 2023. Recent insights into the
use of antagonistic yeasts for sustainable biomanagement of postharvest pathogenic and mycotoxigenic fungi in
fruit with their prevention strategies against mycotoxins. *Journal of Agricultural and Food Chemistry* 71(26),
9923-9950. https://doi.org/10.1021/acs.jafc.3c00315.

Pal, K.K. and Gardener, B.M. 2006. Biological Control of Plant Pathogens. *The Plant Health Instructor* 2, 1117-1142.
https://doi.org/10.1094/PHI-A-2006-1117-02

Plaza, P., Sanbruno, A., Usall, J., Lamarca, N., Torres, R., Pons, J., and Viñas, I. 2004. Integration of curing treatments
with degreening to control the main postharvest diseases of clementine mandarins. *Postharvest Biology and
Technology* 34(1):29–37. https://doi.org/10.1016/j.postharvbio.2004.03.012

Riolo, M., Luz, C., Santilli, E., Meca, G., and Cacciola, S.O. 2023. Antifungal activity of selected lactic acid bacteria
from olive drupes. *Food Bioscience* 52,102422. https://doi.org/10.1016/j.fbio.2023.102422

Rovetto, E. I., La Spada, F., Aloi, F., Riolo, M., Pane, A., Garbelotto, M., and Cacciola, S. O. 2024. Green solutions and
new technologies for sustainable management of fungus and oomycete diseases in the citrus fruit supply chain.
Journal of Plant Pathology 1-27. https://doi.org/10.1007/s42161-023-01543-6

Rovetto, E. I., Luz, C., La Spada, F., Meca, G., Riolo, M., and Cacciola, S. O. 2023. Diversity of Mycotoxins and Other
Secondary Metabolites Recovered from Blood Oranges Infected by *Colletotrichum*, *Alternaria*, and *Penicillium*
Species. *Toxins* 15(7), 407. https://doi.org/10.3390/toxins15070407

Sanchez-Ballesta, M. T., Zacarias, L., Granell, A., and Lafuente, M. T. 2008. β-1, 3-Glucanase gene expression as a
molecular marker for postharvest physiological disorders in citrus fruit and its hormonal regulation. *Postharvest
Biology and Technology* 48(1), 146-149. https://doi.org/10.1016/j.postharvbio.2007.09.007

Sánchez-Torres, P., and Tuset, J. J. 2011. Molecular insights into fungicide resistance in sensitive and resistant
Penicillium digitatum strains infecting citrus. *Postharvest Biology and Technology* 59(2), 159-165.
https://doi.org/10.1016/j.postharvbio.2010.08.017

Shi, Z., Wang, F., Lu, Y., and Deng, J. 2018. Combination of chitosan and salicylic acid to control postharvest green
mold caused by *Penicillium digitatum* in grapefruit fruit. *Scientia Horticulturae 233*, 54-60.
https://doi.org/10.1016/j.scienta.2018.01.039

Smilanick, J. L., Mansour, M. F., Margosan, D. A., Gabler, F. M., and Goodwine, W. R. 2005. Influence of pH and
NaHCO3 on effectiveness of imazalil to inhibit germination of *Penicillium digitatum* and to control postharvest
green mold on citrus fruit. *Plant Disease* 89(6), 640-648. https://doi.org/10.1094/PD-89-0640

Souza Morais Reis, M., Carvalho Rezende, D., Moreira Mafra, N., Carvalho Machado, L., Lessi dod Reis, L., Silva
Gonçalves, A., and Malaquias da Silva, L.G. 2024. Potassium phospite in the management of green mold in
citrus. *Revista Observatorio de La Economia LatinoAmericana* 22(1), 422-442.
https://doi.org/10.55905/oelv22n1-024

Stracquadanio, C., Luz, C., La Spada, F., Meca, G., and Cacciola, S. O. 2021. Inhibition of mycotoxigenic fungi in
different vegetable matrices by extracts of *Trichoderma* species. *Journal of Fungi (Basel)* 7(6):445.
https://doi.org/10.3390/jof7060445

Stracquadanio, C., Quiles, J. M., Meca, G., and Cacciola, S. O. 2020. Antifungal activity of bioactive metabolites
produced by *Trichoderma asperellum* and *Trichoderma atroviride* in liquid medium. *Journal of Fungi (Basel)*
6(4): 263. https://doi.org/10.3390/jof6040263

Strange, R. N., and Scott, P. R. 2005. Plant Disease: A threat to global food security. *Annual Review of Phytopathology*
43(1):83-116. https://doi.org/10.1146/annurev.phyto.43.113004.133839

Strano, M. C., Di Silvestro, S.; Coniglione, M., and Magnano di San Lio, R. 2015. Decay control of cold stored *Citrus clementina* Hort. ex Tan. fruit by pre- and post-harvest application of potassium phosphite. *Advances in Horticultural Science* 29(2-3), 97-102. https://doi.org/10.13128/ahs-22686

Sui, Y., Wisniewski, M., Droby, S., Norelli, J., and Liu,J. 2016. Recent advances and current status of the use of heat treatments in postharvest disease management systems: Is it time to turn up the heat?. *Trends in Food Science and Technology* 51, 34-40. https://doi.org/10.1016/j.tifs.2016.03.004.

Suwannarach, N., Bussaban, B., Nuangmek, W., Pithakpol, W., Jirawattanakul, B., Matsui, K., and Lumyong, S. 2016. Evaluation of *Muscodor suthepensis* strain CMU☐Cib462 as a postharvest biofumigant for tangerine fruit rot caused by *Penicillium digitatum*. *Journal of the Science of Food and Agriculture* 96(1), 339-345. https://doi.org/10.1002/jsfa.7099

Talibi, I., Boubaker, H., Boudyach, E. H., and Ait Ben Aoumar, A. 2014 Alternative methods for the control of postharvest citrus diseases. *Journal of Applied Microbiology* 117(1):1–17. https://doi.org/10.1111/jam.12495

Van Loon, L. C., Rep, M., and Pieterse, C. M. 2006. Significance of inducible defense-related proteins in infected plants. *Annual Review of Phytopathology* 44, 135-162. https://doi.org/10.1146/annurev.phyto.44.070505.143425

Vu, T. X., Ngo, T. T., Mai, L. T. D., Bui, T. T., Le, D. H., Bui, H. T. V., Nguyen, H. Q., Ngo, B. X., and Tran, V. T. 2018. A highly efficient *Agrobacterium tumefaciens*-mediated transformation system for the postharvest pathogen *Penicillium digitatum* using DsRed and GFP to visualize citrus host colonization. *Journal of Microbiological Methods* 144:134–144. https://doi.org/10.1016/J.MIMET.2017.11.019

Wang, L., Wu, H., Qin, G., and Meng, X. 2014. Chitosan disrupts *Penicillium expansum* and controls postharvest blue mold of jujube fruit. *Food Control* 41(1):56–62. https://doi.org/10.1016/j.foodcont.2013.12.028

Wang, W., Li, T., Chen, J., Zhang, X., Wei, L., Yao, S., and Zeng, K. 2023. A self-regulated transcription factor CsWRKY33 enhances resistance of citrus fruit to *Penicillium digitatum*. *Postharvest Biology and Technology* 198, 112267. https://doi.org/10.1016/j.postharvbio.2023.112267

Youssef, K., Sanzani, S. M., Ligorio, A., Ippolito, A., and Terry, L. A. 2014. Sodium carbonate and bicarbonate treatments induce resistance to postharvest green mould on citrus fruit. *Postharvest Biology and Technology* 87, 61-69. https://doi.org/10.1016/j.postharvbio.2013.08.006

Zacarias, L., Cronje, P. J., and Palou, L. 2020. Postharvest technology of citrus fruit. In The genus citrus. *Woodhead Publishing* 421-446. http://dx.doi.org/10.1016/B978-0-12-812163-4.00021-8

Zhang, X., Li, B., Zhang, Z., Chen, Y., and Tian, S. 2020. Antagonistic Yeasts: A promising alternative to chemical fungicides for controlling postharvest decay of fruit. *Journal of Fungi (Basel)* 6:158. https://doi.org/10.3390/JOF6030158

Zheng, X., Jiang, H., Silvy, E. M., Zhao, S., Chai, X., Wang, B., ... and Prusky, D. 2021. *Candida oleophila* proliferated and accelerated accumulation of suberin poly phenolic and lignin at wound sites of potato tubers. *Foods* 10(6), 1286. https://doi.org/10.3390/foods10061286