Title:

Development of a new genotype–phenotype linked antibody screening system

Authors:

Takashi Watanabe[1*], Hikaru Hata[2], Yoshiki Mochizuki[1], Fumie Yokoyama[1], Tomoko

Hasegawa[1], Naveen Kumar[3], Tomohiro Kurosaki[2,4], Osamu Ohara[1,5] and Hidehiro

Fukuyama[2,6,7]

Affiliations:

[1]Laboratory for Integrative Genomics, RIKEN Center for Integrative Medical Sciences,

Yokohama, 230-0045, Japan

[2]Laboratory for Lymphocyte Differentiation, RIKEN Center for Integrative Medical Sciences,

Yokohama, 230-0045, Japan

[3]Laboratory for Integrated Bioinformatics, RIKEN Center for Integrative Medical Sciences,

Yokohama, 230-0045, Japan

[4]Laboratory of Lymphocyte Differentiation, WPI Immunology Frontier Research Center,

Osaka University, Osaka, 565-0871, Japan

[5]Department of Applied Genomics, Kazusa DNA Research Institute, Chiba 292-0818, Japan

17   [6]Near-InfraRed Photo-Immunotherapy Research Institute, Kansai Medical University,

18   Hirakata, Osaka,573-1010, Japan

19   [7]INSERM EST, Strasbourg Cedex 2, 67200, France

20   *To whom correspondence should be addressed. Tel: +81 45 503 9696; Fax: +81 45 503

21   9694; Email: twatanabe@riken.jp

Impact Statement:

23   A Golden Gate-based dual-expression vector enables rapid screening, facilitating efficient

24   isolation of high-affinity cross-reactive antibodies for therapeutic or diagnostic use, offering

25   a crucial advance for pandemic preparedness.

Keywords:

27   Membrane-bound Ig expression, NGS-based Ig-seq, Golden Gate Cloning, influenza

28   broadly-reactive

Abstract:

Antibodies are powerful tools for the therapy and diagnosis of various diseases. In addition to conventional hybridoma-based screening, recombinant antibody-based screening has become a common choice; however, its application is hampered by two factors: 1) screening starts only after Ig gene cloning and recombinant antibody production, and 2) the antibody is composed of paired chains, heavy and light, commonly expressed from two independent expression vectors. Here, we introduce a method for the rapid screening of recombinant monoclonal antibodies by establishing a Golden Gate-based dual-expression vector and *in vivo* expression of membrane-bound antibodies. Using this system, we demonstrated the efficient isolation of influenza cross-reactive antibodies with high affinity from mouse germinal center B cells over 4 days. This system is particularly useful for isolating therapeutic or diagnostic antibodies (e.g., during foreseen pandemics).

Introduction:

Technical improvements in the rapid isolation of monoclonal antibodies (mAbs) are critical for their diagnostic and therapeutic development. This need became apparent to the general public during the COVID-19 pandemic. Antibodies isolated from convalescent COVID-19 patients are valuable resources for the development of mAb therapeutics and rapid diagnostic tools. Human antibodies can be administered to other patients without the need to humanize in an emergency and with a slight modification of Fc receptor binding (1) to increase the half-life or reduce the risk of antibody-dependent enhancement. Although production costs remain high, the efficiency of generating recombinant antibodies has dramatically improved with the use of mammalian cell lines, and the application of therapeutic antibodies is likely to expand in the near future (2).

The gold standard method for isolating mAbs is the hybridoma technology developed by Kohler and Milstein in 1975 (3), in which antibody-producing B cells are fused with immortal B cells, called myelomas, to produce long-lasting antibody-producing B cells. In addition to hybridoma technology, new methods have been developed to increase mAb screening efficiency. These include direct immortalization of B cells by gene reprogramming using the Epstein–Barr virus (4) or retrovirus-mediated gene transfer (5), cloning of variable region-encoding genes by single-cell PCR (6,7), single-cell culture

screening (8), and *in vitro* screening of recombinant antibody libraries (9–13). Some of

these methods successfully yield high-affinity antibodies in various formats, including

single variable domain on a heavy chain or single-chain fragment variable antibodies.

Although hybridoma technology is highly reliable for the isolation of valuable antibodies

from animal models, such as mice, rats, and hamsters, it is time-consuming and requires

technical skill. The development of a new fusion partner cell line, SPYMEG, has enabled the

production of human hybridomas and opened a new direction for the isolation of human

mAbs (14). Although many of these new technologies have increased the throughput of

mAb isolation, the procedure still requires significant resources and time.

To dramatically enhance the efficiency of mAb isolation, we used next-generation

sequencing (NGS) technology, which has changed the method for sequencing

immunoglobulin (Ig) variable-region genes. For instance, tens of thousands of Ig genes

specific to certain antigens can be identified by combining droplet-based single-cell

isolation with DNA barcode antigen technology, followed by NGS (15). Although Ig genes

can be sequenced at a high throughput, there is no method for screening antibodies in a

high-throughput format that is compatible with NGS technology.

In this study, we developed a new functional screening method that is compatible

with NGS to rapidly identify antigen-specific clones. We first generated an Ig dual-

expression vector using Golden Gate Cloning (16), which enabled the linkage of heavy-

chain variable and light-chain variable DNA fragments obtained from a single-sorted B cell,

followed by the expression of membrane-bound Ig. This single-step procedure enabled the

enrichment of antigen-specific, high-affinity Igs by flow cytometry, which was significantly

faster than conventional cloning-based methods that require sequential steps. To

demonstrate the efficiency of our new method, we screened for potent broadly reactive

antibodies (17–19) against the influenza virus in an experimental mouse model using our

technology. Broad reactivity allows for influenza vaccines that work across seasonal

variations in influenza strains, which can only be accomplished by screening a large

number of candidate antibodies. We first raised cross-reactive B cells against various

hemagglutinin (HA) antigens from the influenza virus by sequential immunization with

heterotypic HA antigens from group 1 influenza. Using this technology, we obtained

several mAbs that bind to other group 1 HA antigens and even to group 2 HA antigens of

the influenza virus. Our technology can also be applied to human antibody screening and

represents a new line of mAb screening that accelerates the isolation of therapeutic and

diagnostic mAbs.

Results:

Rapid membrane-bound dual Ig expression screening system

Cloning-based Ig screening involves several steps: cloning two Ig gene fragments

encoding heavy and light chains independently from a single cell, co-expressing these Ig

chains, and purifying individual recombinant antibodies. These steps are labor intensive.

We established a new antibody screening system for the isolation of valuable mAbs

(details in the method). Eventually, the antigen-binding Ig transformants were collected

by sorting in a bulk fashion, and the unique CDR3 region and clones of interest were

identified using the Ig-seq database (Fig. 1A).

We generated a dual-expression vector to express both Ig heavy and kappa/lambda

in a single-expression vector (Fig. 1B). These plasmids were individually transfected

transiently to the floating human FreeStyle 293 cell line. Igs were expressed on the cell

surface in 2 days; their expression could be confirmed and normalized by *Venus*

expression, since the Ig heavy chain is fused with *Venus* at the cytoplasmic domain tail

(Fig. 1C). The advantage of this system is the rapid enrichment of clones of interest

through bulk screening. In the current single-cell-based cloning/screening, plasmid DNA

extracted from the collected antigen-binding bulk transformants was sequenced for the

heavy chain CDR3 region. The CDR3 region is unique and can be used for the identification

of a clone (20,21). This membrane-bound Ig expression system links antigen-binding

features and genetic information of the Ig repertoire.

Building the Ig-seq database and analysis

As a model experiment, we obtained broadly reactive antibodies against influenza viruses

using multiple HA probes. We prepared two HA proteins as probes: A/Puerto Rico/8/1934

(H1N1), designated as PR8, and A/Okuda/1957 (H2N2), designated as H2. A total of 380

IgG1$^+$ germinal center B cells of either PR8$^+$ (204 cells), H2$^+$ (99 cells), or PR8$^+$H2$^+$ (77

cells) were collected in a single-cell fashion. We obtained sequences of 284 independent

clones. The success rate of cloning the paired Ig fragments was 75.9%. An overview of

heavy chain V-D-J and light chain V-J usage and repertoire clonality is shown in

Supplementary Fig. 1A and B. Mutation rates and CDR3 lengths of the heavy chains of the

three cell populations (PR8$^+$, H2$^+$, and PR8$^+$H2$^+$) were comparable (Supplementary Fig. 1C

and D). These results indicate that broadly reactive antibodies do not require unique

genetic traces to obtain breadth in our experimental setup.

Isolation of H2- and H1-reactive B cells

On day 2 after transfection, 287 transformants (Ig-expressing transformant libraries) were

stained with HA probes (Fig. 2A). As shown in Fig. 2B, we selected three prominent

populations, H1$^+$H2$^+$ (cross), H2$^+$ (H2), and H1$^+$ (H1), and collected them in bulk. In

contrast to bacterial transfection, mammalian cell lines, such as FreeStyle 293 cells, can

contain multiple plasmids in a cell and ectopically express multiple proteins. Therefore, we

decided to mix the transformants instead of plasmids. We sorted 2,981 cells from these

three populations and performed sequencing. Bulk Ig-seq data are referred to as Ig-seq

data, as described above. All the transformants were examined individually. A total of 90

clones from the three populations were verified for their sequences and HA binding by flow

cytometry analysis, as shown in Fig. 2. Among these, 110 "H1$^+$," 67 "H2$^+$," and 13 "cross"

clones were successfully isolated. Then, 138 transformants were verified for their binding

to HAs by flow cytometry. Of these, 81 "H1$^+$," 48 "H2$^+$," and nine "cross" clones were

successfully isolated. Finally, 8 of the 13 "cross" clones obtained in bulk were included in

this group; the binding of one clone (D11p4) was so weak that it was not included in the

gates sorted in bulk. Thus, we established a high-throughput method for sorting "cross"

clones.

Characterization of "cross" clones for broad reactivity against the influenza virus

By generating a secretory form of mAbs for these nine cross clones, including C10p2, we

characterized their affinity and broadness using six different HA antigens from strains

A/Okuda/1957 (H2N2) named for H2, A/Puerto Rico/8/1934 (H1N1) for PR8,

A/California/2009 (X-179A) [H1N1] Pdm09 for Cal, A/Texas/50/2012 (X-223) [H3N2] for

H3, A/Egypt/N03072/2010(H5N1) for H5, and A/Brisbane/59/2007 [H1N1] (19) for Stem.

We determined the affinities (K_d) of the antibodies for the six HA antigens using surface

plasmon resonance analysis (Table 1 and Supplementary Fig. 2), which ranged from 500

to 100 nM. With the highest affinity among them, the C10p2 antibody bound to Cal at K_d

$\simeq 5.66 \times 10^{-10}$ (M). Surprisingly, the antibodies showed a broader spectrum of HA binding

among the influenza strains. Seven of the nine mAbs bound to HA from the highly

pathogenic avian influenza strain H5N1. Furthermore, the six antibodies that bound to H3

were categorized into group 2 (22). In addition, we observed that two of the broadly

reactive antibodies recognized the stem region of the HA of A/Brisbane/59/2007. A

possible scenario for other stem cell-negative antibodies is that they may either bind to

the stem region of other strains or to the head region of HA. We tested whether some of

these antibodies shared features with classic broadly neutralizing antibodies. C179, the

first broadly neutralizing mouse antibody isolated in 1993 (17), reacts with the stem

region of HAs from group 1 influenza virus strains (22). C179 binds to HAs from group 1

influenza viruses, including H1N1, H2N2, and H5N1. A competition assay for the binding of

H2 and PR8 revealed that A6p4 competes with C179 (Fig. 3). In addition, our two-

dimensional phylogeny map indicated that A6p4 was unique to all clones (Supplementary

Fig. 3). Note that we used NSP2 as a negative control antibody, which binds specifically to

the HA from A/California/2009 (X-179A) [H1N1] Pdm09 (23). These results demonstrate

that we were able to isolate a broadly reactive mAb with an affinity comparable to that of

a classic broadly reactive mAb (C179) using our technology.

Discussion:

The discovery of therapeutic antibodies is accumulating, and their demand is increasing

(24). Efficiently isolating one biologically significant antibody from the 10^{13} antibody

repertoire is challenging. The hybridoma method has long been used for this purpose and

remains reliable. However, the fusion rate of B and myeloma cells is < 0.001% in this

method; therefore, the number of hybridomas obtained from one individual B cell is

limited to several hundred clones, limiting its use in the production of mAbs. As an

alternative choice, cloning-based antibody screening has gained popularity, especially for

human antibody discovery, because of the lack of a human myeloma cell line similar to

that used for murine cell fusion partners. The SPYMEG cell line can generate antigen-

specific antibody-producing cells by fusing the murine SP2/0 myeloma cell line and human

megakaryoblastic leukemia cell line MEG-01 with human PBMCs (14, 25). In addition to

gene expression profiling, NGS facilitates antibody repertoire analysis. However, despite

breakthroughs in sequencing technology, the process of linking antibody function and its

genes is still long and labor intensive. To overcome this issue, we developed a new

antibody screening system to directly link antigen–antibody binding (as a function) with a

gene encoding the antibody by expressing its membrane-bound form. Among conventional

cloning-based mAb isolation methods, this system demonstrates several advantages. 1)

Membrane Ig expression can link the antigen-binding feature of membrane-expressed Ig,

which can be linked to Ig DNA sequence information using our plasmid construct. Similar

to the panning procedure used in phage display (26), antigen-binding cells carry the

plasmid encoding the Ig genes through surface Ig so that the selection process in bulk

format enriches the relevant plasmids for use in further experiments. 2) The dual Ig

expression vector links heavy- and light-chain genes, which reduces the plasmid

preparation time and stock by half. 3) Highly reliable Golden Gate Cloning technology

using type IIs restriction enzymes can readily generate plasmid clones. This reduced the

time required to generate an Ig plasmid library. 4) The population profile, defined by the

fluorescence intensity during flow cytometry, directly reflected the affinity of a clone

(Supplementary Fig. 4). Collectively, our technology has streamlined the isolation of mAbs

for therapy and diagnosis.

Conclusion:

Our findings indicate that the developed antibody presentation system facilitates antibody

functional analysis and is well suited for the discovery of antibodies important for

infectious diseases when combined with conventional NGS-based antibody repertoire

analysis. Compared with droplet-based experimental systems, well-based systems are

limited in the number of cells they can process. Furthermore, experiments involving

infectious bacteria and viruses have imposed limitations on human experimentation. To

solve these problems, the automation of experiments will become important in the future.

By combining our screening system with robotic automation of experiments, it will be

possible to obtain useful mAbs for various diseases quickly and in large quantities, which

has broad implications for the development of vaccines against various diseases.

211   Materials and Methods:

212   Protein purification of influenza HAs (H1 and H2) as antigens

A mixture of 25.7 µg of HA, comprising H1 strain A/Puerto Rico/8/1934 (PR8), H1 strain

A/California/7/2009 (X-179A), H2 strain A/Okuda/1957 (H2), H3 strain A/Texas/50/2012

(X-223), or H5 strain A/Egypt/N03072/2010], 1.3 µg of NA (Strain-matched), and 3 µg of

BirA-expressing plasmids were transfected into Expi293 cells using the Expifectamine 293

transfection kit (Thermo Fisher Scientific, Waltham, MA, USA) according to the

manufacturer's instructions. The cells were then cultured in Expi293 Expression Medium

(Thermo Fisher Scientific) supplemented with D-biotin at a final concentration of 100 µM

for *in vivo* BirA biotinylation in a humidified incubator containing 8% CO_2 at 37°C and 125

rpm. If biotinylation was not necessary, transfection was performed without BirA, and cell

culture was carried out without D-biotin. Culture supernatants were harvested at 3 days

post-transfection and filtered through a 0.45 µm filter (Merck Millipore, Burlington, MA,

USA). All proteins were purified with a Talon metal affinity resin (Takara Bio USA,

Mountain View, CA, USA) and dialyzed against pyrogen-free PBS. The oligomeric state and

purity were determined at 25°C by size-exclusion chromatography on a Superdex 200

10/300GL column (GE Healthcare Technologies, Chicago, IL).

HA (H1 and H2) protein immunization

Two individual BALB/c mice were intraperitoneally immunized sequentially, 2 weeks apart,

with 15 µg of H1 (PR8) HA followed by 15 µg of H2 HA protein as an antigen and

supplemented with AddaVax adjuvant (InvivoGen, San Diego, CA, USA). BALB/c mice

were purchased from CLEA Japan (CLEA Japan, Inc., Tokyo, Japan) All animal experiments

were performed using protocols approved by the Institutional Animal Care and Use

Committee (IACUC) of the RIKEN Yokohama Branch.

Fluorescence-activated cell sorting

CD43-negative B cells were collected using AutoMACS (Miltenyi Biotec, Inc., Bergisch

Gladbach, North Rhine-Westphalia, Germany) and stained with IgG1 (Bv510), non-

biotinylated His-tagged purified recombinant H1 (PR8) protein, and biotinylated purified

recombinant H2 protein on ice for 30 min. After washing three times, the cell suspension

was incubated with Alexa Fluor 488-labeled anti-His antibodies on ice for 30 min. After

washing three times, the cell suspension was incubated with Brilliant Violet 421-

conjugated streptavidin on ice for 3 min. Next, the B cells were subjected to single-cell

sorting using a BD FACSAria III (BD Biosciences, Franklin Lakes, NJ, USA) and collected into

96-well plates pre-loaded with lysis buffer. The single cells in the lysis buffer were

immediately snap-frozen on powdered dry ice and stored at −80°C.

Amplification of a paired B cell repertoire amplicon from a single cell

Single cells were dropped into pre-loaded plates containing 4 μL/well of lysis buffer (2

U/μL RNase inhibitor, 0.5× PBS, 10 mM DTT) and 1 μL of 10 μM Oligo(dT)18 Primer. Cell

lysates were incubated at 70°C for 90 s, followed by 35°C for 15 s, and then placed on ice

for 2 min. For first-strand cDNA synthesis, 5 μL of RT mix (2× PCR buffer, 5 mM DTT, 2

mM dNTP Mix, 10 U RNase inhibitor, 100 U SuperScript III Reverse Transcriptase) was

added to the cell lysate and incubated consecutively at 35°C for 5 min, 45°C for 20 min,

70°C for 10 min, and 4°C for chilling. For primer digestion, 5 μL of the primer digestion

mix (1× ExoI Buffer, 7.5 U Exonuclease I) was prepared and incubated at 37°C for 30

min, 80°C for 20 min, and 4°C for chilling. For the poly-A tailing reaction, 5 μL of the poly-

A tailing reaction mix (4× PCR buffer, 5 mM dATP, 0.1 U RNase H, 60 U Terminal

transferase) was prepared and incubated at 37°C for 50 s, 65°C for 10 min, and 4°C for

chilling. For the 2nd strand cDNA synthesis, 25 μL of the amplification mix (1× KAPA HiFi

259 HotStart ReadyMix, 400 nM SMART_dT primer, 10 μL of poly-A tailing product, 1.5 μL of

260 nuclease-free H_2O) was prepared and incubated at 95°C for 3 min, 40°C for 1 min, 65°C

261 for 10 min, and 72°C for 5 min and chilled. For the cDNA amplification using an Ig-specific

262 outer primer set, 20 μL of the outer amplification mix (1× KOD FX Neo buffer, 400 μM

263 dNTPs, 1× UPM, 300 nM outer primer, 3.5 μL of the SMART product, 0.5 U KOD FX Neo

264 polymerase) was prepared and incubated at 94°C for 2 min, followed by 25 cycles of 98°C

265 for 10 s, 60°C for 30 s, and 68°C for 45 s and then chilled. For cDNA amplification with an

266 Ig-specific inner primer set, 20 μL of the inner amplification mix (1× KOD FX Neo buffer,

267 400 μM dNTPs, 500 nM 5' matrix primer mix, 500 nM 3' matrix primer mix, 3.5 μL of the

268 outer cDNA amplified product, 0.5 U KOD FX Neo polymerase) was prepared and

269 incubated at 94°C for 2 min, followed by 25 cycles of 98°C for 10 s, 60°C for 30 s, and

270 68°C for 45 s and then chilled. Finally, 3 μL of the amplified inner cDNA product was used

271 for further plate indexing by amplification. Then, 20 μL of the plate-indexing amplification

272 mix (1× KOD FX Neo buffer, 400 μM dNTPs; 500 nM i5-Fwd or i7-Fwd, 500 nM i7-Rev or

273 i5-Rev; 2 μL of the inner cDNA amplified product; 0.5 U KOD FX Neo polymerase) was

274 prepared and incubated at 94°C for 2 min, followed by 25 cycles at 98°C for 10 s, 60°C

275 for 30 s, and 68°C for 45 s and then chilled. All primers used are listed in Supplementary

276 Table 1 and Supplementary Table 2.

 Paired-end Illumina sequencing strategy and analysis

The libraries were sequenced on a MiSeq instrument (Illumina, San Diego, CA, USA)

following the manufacturer's protocol. As the sequencing length was limited to 2 × 300

bp, we used the following strategy. Amplicons in both orientations were generated using

an Illumina adapter sequence. Then, asymmetric 400+100 nt paired-end sequencing was

performed to obtain high-quality 400 + 400 nt paired-end sequences of the original cDNA

molecule. This approach provided a sufficient sequencing length to capture extra-long Ig

variants. We followed Turchaninova's method (27). Only reads that met the following four

conditions were used as the resulting Ig sequences: 1) reads with a length of 260 nt or

more; 2) cumulative clustered reads representing the top 5% of the entire reads; 3) 5'-

and 3'-reads overlapping a length of 20 bases or more; and 4) a total length of 460 ± 50

bp for heavy chain or 390 ± 40 bp for light chain. All resulting sequence data were

analyzed using IMGT/High V-QUEST (http://www.imgt.org/IMGT_vquest/vquest) for

annotation (28). Using the sequences and annotations, Ig database construction and

visualization of the Ig repertoire were performed using the in-house BONSCI software.

Construction of a mouse IgG/IgK expression plasmid

The BsaI restriction sites were inserted into the paired B cell repertoire amplicon from a

single cell using PCR. The destination vector contained the IL6 signal peptide, EF1a

promoter, ccdB gene, and mouse IgK Fc region. The donor vector contained the mouse

IgG1 Fc region, the *Venus* gene, and the EF1a promoter. The paired B cell repertoire

amplicons, destination vector, and donor vector containing BsaI restriction sites were

assembled using our assembly method. For this step, 10 µL of the assembly mix (1× T4

DNA ligase buffer, 1× BSA, 1 U BsaI restriction enzyme, 40 U T4 DNA ligase, 100 ng

heavy chain amplicon, 100 ng light chain amplicon, 100 ng destination vector, and 100 ng

donor vector) was prepared and incubated for 25 cycles at 37°C for 3 min, 16°C for 4 min,

50°C for 5 min, and 80°C for 5 min, and then chilled. The antibody sequence that entered

the construct was fused to the *Venus* sequence and expressed in membrane form.

Antibody display on FreeStyle 293 cultured cells

A mixture of 1 µg of antibody-expressing plasmid was transfected into FreeStyle 293 cells

using the 293fectin Transfection Reagent (Thermo Fisher Scientific) according to the

manufacturer's instructions and cultured in FreeStyle 293 Expression Medium (Thermo

Fisher Scientific) in a humidified incubator with 8% CO_2 at 37°C and 125 rpm. Antibodies

were displayed on the surface of cultured cells, which enabled the determination of

antigen specificity. Furthermore, the antibody-display cells were tested for binding activity

with Alexa647-labeled H1 and Alexa568-labeled H2 using a BD FACSAria III (BD

Biosciences).

Antibody production

To obtain a large amount of secretory antibodies, PCR fragments from the antibody-

expressing plasmid were cloned into the pcDNA3.4-mIgG1 or pcDNA3.4-kappa vectors. A

mixture of 15 μg of pcDNA3.4-V-gene-mIgG1 vector and 15 μg of pcDNA3.4-V-gene-

kappa vector were transfected into Expi293 cells using the ExpiFectamine 293 Transfection

Kit (Thermo Fisher Scientific), according to the manufacturer's instructions, and cultured in

Expi293 Expression Medium (Thermo Fisher Scientific) in a humidified 8% CO_2 incubator at

37°C and 125 rpm. Then, 5 days post-transfection, the culture supernatants were

harvested, and all proteins were purified with PureSpeed IMAC resin (Mettler Toledo,

Columbus, OH, USA), according to the manufacturer's instructions.

Surface plasmon resonance

Kinetic analyses were performed at 25°C using a BIAcore 3000 machine (GE Healthcare

Technologies). A2p1, B10p2, C10p2, E11p2, G6p2, A6p4, D4p4, D11p4, or G12p4

antibodies were immobilized on a CM5 sensor chip (GE Healthcare Technologies) with an

amine-coupling kit according to the manufacturer's instructions. HA probes were serially

diluted at five different concentrations and injected at a flow rate of 30 μL/min for 3 min

with a dissociation time of 7 min in HBS−EP buffer (10 mM HEPES, pH 7.4, 150 mM NaCl,

3.4 mM EDTA, and 0.005% Surfactant P20). The chip was regenerated using 10 mM

glycine (pH 2.5) as an HA probe.

Data access:

All raw and processed sequencing data generated in this study have been submitted to the

NCBI Gene Expression Omnibus (GEO; https://www.ncbi.nlm.nih.gov/geo/) under

accession number GSE140720.

Acknowledgements:

We thank Mr. Atsuo Kobayashi, Ms. Chieko Okamura, Dr. Seok-Won Kim, Mr. Maxime

Hebrard, Dr. Todd D Taylor, and the RIKEN IMS Genome Platform for their technical

support, and Dr. Christopher Nicholas for critical reading and English editing of the

manuscript. We would like to thank Editage (www.editage.jp) for English language editing.

Funding:

(S) [26221306 to T.K. and H.F.]. The funding sources were not involved in the study

design, data collection and interpretation, or paper submission decisions.

Conflict of interest:

None declared.

Author contributions:

T.W. and H.F. designed the study, the main conceptual ideas, and the proof outline. H.H.,

Y.M., F.Y., T.H., and N.K. collected the data. O.O. and T.K. supervised the project. T.W.

wrote the manuscript with support from H.F. All authors discussed the results and

commented on the manuscript.

354 References:

1. Balsitis, S. J., K. L. Williams, et al. (2010). Lethal antibody enhancement of dengue

disease in mice is prevented by Fc modification. *PLoS Pathog.,* **6**(2): e1000790.

2. Fang, X. T., D. Sehlin, et al. (2017). Efficient and inexpensive transient expression of

multispecific multivalent antibodies in Expi293 cells. *Biol. Proced. Online,* **19**(1): 1–9.

3. Köhler, G. and C. Milstein (1975). Continuous cultures of fused cells secreting antibody

of predefined specificity. *Nature*, **256**(5517): 495–497.

4. Traggiai, E., S. Becker, et al. (2004). An efficient method to make human monoclonal

antibodies from memory B cells: potent neutralization of SARS coronavirus. *Nat. Med*.,

10(8): 871–875.

5.   Kwakkenbos, M. J., S. A. Diehl, et al. (2010). Generation of stable monoclonal

antibody-Producing B cell receptor-positive human memory B cells by genetic

programming. *Nat. Med.*, **16**(1): 123–128.

6.   Wrammert, J., K. Smith, et al. (2008). Rapid cloning of high-affinity human monoclonal

antibodies against influenza virus. *Nature*, **453**(7195): 667–671.

7.   Meijer, P. J., P. S. Andersen, et al. (2006). Isolation of human antibody repertoires

with preservation of the natural heavy and light chain pairing. *J. Mol. Biol.*, **358**(3):

764–772.

8.   Kuraoka, M., A. G. Schmidt, et al. (2016). Complex antigens drive permissive clonal

selection in germinal centers. *Immunity*, **44**(3): 542–552.

9.   Clackson, T., H. R. Hoogenboom, et al. (1991). Making antibody fragments using

phage display libraries. *Nature*, **352**(6336): 624–628.

10. Feldhaus, M. J., R. W. Siegel, et al. (2003). Flow-cytometric isolation of human

antibodies from a nonimmune Saccharomyces cerevisiae surface display library. *Nat.*

Biotechnol., **21**(2): 163–170.

11. Harvey, B. R., G. Georgiou, et al. (2004). Anchored periplasmic expression, a versatile

technology for the isolation of high-affinity antibodies from Escherichia coli-expressed

libraries. *Proc. Natl. Acad. Sci. U. S. A.*, **101**(25): 9193–9198.

12. Schaffitzel, C., J. Hanes, et al. (1999). Ribosome display: an in vitro method for

selection and evolution of antibodies from libraries. *J. Immunol. Methods*, **231**(1–2):

119–135.

13. Mazor, Y., T. Van Blarcom, et al. (2007). Isolation of engineered, full-length antibodies

from libraries expressed in Escherichia coli. *Nat. Biotechnol.*, **25**(5): 563–565.

14. Kubota-Koketsu, R., H. Mizuta, et al. (2009). Broad neutralizing human monoclonal

antibodies against influenza virus from vaccinated healthy donors. *Biochem. Biophys.*

Res. Commun., **387**(1): 180–185.

15. Setliff, I., A. R. Shiakolas, et al. (2019). High-throughput mapping of B cell receptor

sequences to antigen specificity. *Cell*, **179**(7): 1636–1646. e15.

16. Kirchmaier, S., K. Lust, et al. (2013). Golden GATEway cloning--a combinatorial

approach to generate fusion and recombination constructs. *PloS One*, **8**(10): e76117.

17. Okuno, Y., Y. Isegawa, et al. (1993). A common neutralizing epitope conserved

between the hemagglutinins of influenza A virus H1 and H2 strains. *J. Virol.*, **67**(5):

2552–2558.

18. Ekiert, D. C., A. K. Kashyap, et al. (2012). Cross-neutralization of influenza A viruses

mediated by a single antibody loop. *Nature*, **489**(7417): 526–532.

19. Impagliazzo, A., F. Milder, et al. (2015). A stable trimeric influenza hemagglutinin stem

as a broadly protective immunogen. *Science*, **349**(6254): 1301–1306.

20. D'Angelo S, Ferrara F, et al. (2018). Many Routes to an Antibody Heavy-Chain CDR3:

Necessary, Yet Insufficient, for Specific Binding. *Front. Immunol.*, **9**(395)

21. Xu, J. L. and M. M. Davis (2000). Diversity in the CDR3 region of VH is sufficient for

most antibody specificities. *Immunity*, **13**(1): 37–45.

22. Gamblin, S. J., and J.J. Skehel. (2010). Influenza hemagglutinin and neuraminidase

membrane glycoproteins. *J. Biol. Chem.*, **285**(37): 28403–28409.

23. Adachi, Y., T. Onodera, et al. (2015). Distinct germinal center selection at local sites

shapes memory B cell response to viral escape. *J. Exp. Med.*, **212** (10): 1709–1723.

24. Kaplon H, Crescioli S, et al. (2023). Antibodies to watch in 2023. *Mabs*, **15**(1):

2153410

25. P Soni, A Yasuhara, et al. (2018). Evaluation of the fusion partner cell line SPYMEG for

obtaining human monoclonal antibodies against influenza B virus. *J. Vet. Med. Sci.*,

80(6): 1020–1024

26. Petropoulos, K., (2012) Phage display. *Methods Mol. Biol.*, **901**: 33–51.

27. Turchaninova, M. A., A. Davydov, et al. (2016). High-quality full-length

immunoglobulin profiling with unique molecular barcoding. *Nat. Protoc.*, **11**(9): 1599–

1616.

28. Alamyar, E., P. Duroux, et al. (2012). IMGT® tools for the nucleotide analysis of

immunoglobulin (IG) and T cell receptor (TR) V-(D)-J repertoires, polymorphisms, and

IG mutations: IMGT/V-QUEST and IMGT/HighV-QUEST for NGS. *Methods Mol. Biol.,*

882: 569–604.

Figures:

Figure 1. System to link the antigen-binding feature with the Ig repertoire genetic

information. **A.** Single cell-sorted B cells were subjected to two processes: Ig-seq database

building and Ig-expressing transformant library preparation. Antigen-binding Ig

transformants were collected by sorting in bulk and the sequences of the unique CDR3

regions and clones of interest were identified by referring to the Ig-seq database. **B.** To

express both Ig heavy and kappa/lambda chains in a single-expression vector, we

generated a dual-expression vector. Four gene fragments were assembled by the Golden

Gate method using the BsaI restriction enzyme. They included: 1) the destination vector

containing the Ig kappa constant gene and the ccdB Chloramphenicol cassette for negative

selection, 2) fragments containing the Ig heavy constant gene fused to *Venus* derived

from the donor vector, 3) the Ig heavy variable gene fragment, and 4) the Ig light variable

gene fragment. **C.** Individually purified plasmids were transiently transfected to the

floating human FreeStyle 293 cell line. Igs were expressed on the cell surface in 2 days,

and expression levels were confirmed and normalized by *Venus* expression, since the Ig

heavy chain was fused to *Venus* at the cytoplasmic domain tail.

Figure 2. Isolation of H2 and H1-reactive B cells. **A.** Mixture of Ig-expressing transformant

libraries stained with HA probes. **B.** Three prominent populations, H1$^+$H2$^+$ (cross), H2$^+$

(H2), and H1$^+$ (H1), were selected and collected in a bulk fashion. Then, the three

prominent populations were sorted and sequenced. **C.** The bulk Ig-seq data were referred

to as the single-cell Ig-seq data. Wells containing crossed antibodies were determined by

comparing the sequence of the heavy chains of the populations included in H1$^+$H2$^+$ (cross)

449   with the Ig-seq data of single cells. **D.** Cross antibodies obtained by bulk sorting were

450   examined by single-cell presentation. **E.** A summary of the results to check whether a

451   candidate mAb was cross-reactive.

Figure 3. Competition assay using various mAbs for binding to H2 and PR8 was

performed, including two previously reported antibodies, C179, a broadly-reactive mAb

binding to the Stem region of HA, and NSP2 (Narita strain-specific mAb), a strain-specific

mAb that binds to the head region of HA.

Table 1

Surface plasmon resonance (Biacore)   KD (M)

Group	1	1	1	2	1	1
	H2	PR8	Cal	H3	H5	Stem
A6p4	4.51E-08	1.41E-06	1.14E-05		2.91E-05	1.59E-08
B10p2	6.75E-09	2.3E-09	3.29E-09			
C10p2	1.66E-09	1.29E-09	5.66E-10			4.26E-08
A2p1	2.49E-06	1.42E-07	1.14E-05	1.25E-04	1.91E-05	
D11p4	7.93E-07	1.72E-06	1.04E-06	1.36E-05	1.05E-11*	
E11p2	4.4E-08	1.86E-08	1.67E-06	2.92E-07	2.37E-07	
G6p2	3.51E-08	3.53E-08	1.44E-08	1.78E-05	3.05E-08	
D4p4	1.1E-05	1.78E-05	6.4E-05	4.38E-05	2.47E-05	
G12p4	1.45E-07	1.52E-07	5.7E-07	9.44E-07	1.58E-07	

High
Middle
Low

Table 1. Affinities (K_d) of the antibodies were determined to 6 HA antigens from strains

A/Okuda/1957(H2N2) named for H2, A/Puerto Rico/8/1934(H1N1) for PR8,

A/California/2009 (X-179A) [H1N1] Pdm09 for Cal, A/Texas/50/2012 (X-223) [H3N2] for

H3, A/Egypt/N03072/2010(H5N1) for H5, and A/Brisbane/59/2007 (H1N1) for Stem using

surface plasmon resonance analysis. H2, PR8, Cal, and H5 belong to group 1, and H3

belongs to group 2.

* The affinity of D11p4 for H5 was low and inaccurate (Supplementary Figure 2).

Supplementary Tables

Supplementary Table 1. Oligonucleotides

Primer	Application	Sequence
First-strand cDNA synthesis		
Oligo(dT)18	Primer for cDNA synthesis	TTTTTTTTTTTTTTTTTT
Second-strand cDNA synthesis		
SMART_dT	TdT oligo	AAGCAGTGGTATCAACGCAGAGTAC(T)$_{30}$VN
Mouse IGH/IGK primer		
TdT_L_BsaI_heavy	mouse IgG heavy-chain cDNA	CTAGGGTCTCAAGCAGTGGTATCAACGCAGAGTAC
msVHE		GAGGTGCAGCTGCAGGAGTCTGG
W-gamma-outer		AGAAGGTGTGCACACCGCTGGAC
mC_G new2		ATCTCCACACACAGGGGCCAGTGGATAGAC
BsaI_IL6sp_L		CTAGGGTCTCAAGCAGATGAACTCCTTCTCCACAAGCG
mC_G_new2_BsaI		TCCTAGGTCTCCCACACACAGGGGCCAGTGGATAGAC
TdT_L_BsaI_light	mouse IgK light-chain cDNA	CTAGGGTCTCAGTGGTATCAACGCAGAGTAC
mvkappa		GAYATTGTGMTSACMCARWCTMCA
mCkappa-outer		ACTGAGGCACCTCCAGATGTT
mC_K_new		AGGCACCTCCAGATGTTAACTGCTCACTGG
BsaI_IL6sp_kappa_L		CTAGGGTCTCAGTGGTATGAACTCCTTCTCCACAAGCG
mC_K_new_BsaI		TCCTAGGTCTCCTCCAGATGTTAACTGCTCACTGG

478 **Supplementary Table 2. Oligonucleotides**

Primer	Application	Sequence
Next Generation sequencing		
M13fwd_i5_mC_G_new2[a]	Well index	TGTAAAACGACGGCCxxxxxxxxATCTCCACACACAGGGGCCAGTGGATAGAC
M13fwd_i5_mC_K[a]		TGTAAAACGACGGCCxxxxxxxxGCACCTCCAGATGTTAACTG
M13fwd_i5_mC_L[a]		TGTAAAACGACGGCCxxxxxxxxAGCTCCTCAGRGGAAGGTG
M13rev_i7_msVHE[b]		GGAAACAGCTATGACyyyyyyyyGAGGTGCAGCTGCAGGAGTCTGG
M13rev_i7_mvkappa[b]		GGAAACAGCTATGACyyyyyyyyGAYATTGTGMTSACMCARWCTMCA
i5-Fwd[c]	Plate index	AATGATACGGCGACCACCGAGATCTACACxxxxxxxxACACTCTTTCCCTACACGACGCTCTTCCGATCTTGTAAAACGACGGCC
i7-Rev[d]		CAAGCAGAAGACGGCATACGAGATyyyyyyyyGTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTGGAAACAGCTATGAC
i5-Rev[c]		AATGATACGGCGACCACCGAGATCTACACxxxxxxxxACACTCTTTCCCTACACGACGCTCTTCCGATCTGGAAACAGCTATGAC
i7-Fwd[d]		CAAGCAGAAGACGGCATACGAGATyyyyyyyyGTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTTGTAAAACGACGGCC

479 [a]xxxxxxxx: sequence showing the well index from D501 to D508. [b]yyyyyyyy: sequence

480 showing the well index from D701 to D712. [c]xxxxxxxx: sequence showing the plate index

481 from D501 to D508. [d]yyyyyyyy: sequence showing the plate index from D701 to D712.

Supplementary Figures

Supplementary Figure 1.

The Ig repertoires were visualized using the in-house software. **A,** Overview of heavy (V-

D-J) and light (V-J) chain usages. **B,** the repertoire clonality from single cells. **C,** the

mutation rates of heavy chains in three cell populations (PR8+, H2+, PR8+H2+) were

compared. **D,** the lengths of CDR3 of heavy chains of three cell populations (PR8+, H2+,

PR8+H2+) were compared.

492
493

Supplementary Figure 2.

Kinetics analysis was performed using a BIAcore 3000 machine (GE Healthcare). A6p4, B10p2, C10p2, A2p1, D11p4, E11p2, G6p2, D4p4 or G12p4 antibodies were immobilized on a CM5 sensor chip (GE Healthcare) using an amine-coupling kit according to the supplier's instructions. HA probes were serially diluted at five different concentrations in HBS-EP buffer.

500
501
502 **Supplementary Figure 3.**
503 Two hundred and eighty-four clones are plotted in a two-dimensional map of phylogeny
504 trees for heavy and light chain genes. Nine mAb clones are overlaid in green. Six "cross"
505 clones are located in a major cluster. B10p2 and C10p2 are co-localized and distant from
506 the cluster. A6p4 is uniquely located in the map.
507

510 **Supplementary Figure 4.**

511 The affinity of antigen-antibody binding (in this case, probe and membrane-bound

512 antibody expressing cell) was inferred from the fluorescence intensity (flow cytometry

513 analysis).

Blocking assay

Type binding
- cross,
- PR8+,
- H2+,