Endocannabinoid biomarkers of vulnerability to the development of trauma-induced generalized fear responses

Laszlo Szente[1,2], Gyula Y Balla[1], Zoltan K Varga[1], Blanka Toth[3], Laszlo Biro[1], Zoltan Balogh[1], Matthew N Hill[4], Mate Toth[1], Eva Mikics[1,5], Mano Aliczki[1,5,*]

[1]Laboratory of Translational Behavioural Neuroscience, Institute of Experimental Medicine, Hungarian Research Network, Budapest, Hungary

[2]János Szentágothai Doctoral School of Neurosciences, Semmelweis University, Budapest, Hungary

[3]Department of Inorganic and Analytical Chemistry, University of Technology and Economics, Budapest, Hungary

[4]Hotchkiss Brain Institute, University of Calgary, Calgary, Alberta, Canada

[5]equal contribution

[*]corresponding author: Mano Aliczki, aliczki.mano@koki.hun-ren.hu

Abstract

Traumatic experiences result in the development of posttraumatic stress disorder (PTSD) in 10-25% of exposed individuals. While human clinical studies suggest that vulnerability is potentially linked to endocannabinoid (eCB) signaling, neurobiological PTSD vulnerability factors are poorly understood. Employing a rat model of PTSD, we characterized distinct resilient and vulnerable subpopulations based on generalized fear, a core symptom of PTSD. Vulnerable subjects showed lower prelimbic and higher ventral hippocampal levels of eCB 2-arachidonoyl-glycerol (2-AG). Ventral hippocampal 2-AG content positively correlated with the strength of fear generalization. Furthermore, vulnerability was associated with downregulation of neuronal activity and neuroplasticity markers, and altered expression of eCB-related genes. Unsupervised and semi-supervised statistical approaches highlighted that hippocampal gene expression patterns possess strong predictive power regarding vulnerability. Taken together, the marked eCB and neuroplasticity changes in vulnerable individuals associated with abnormal activity patterns in the fear circuitry possibly contribute to context coding deficits, resulting in generalized fear.

Abbreviations

2-AG, 2-arachidonoylglycerol

AEA, N-arachidonoylethanolamine, anandamide

AMY, amygdala

BLA, basolateral amygdala

CB1R, cannabinoid receptor type-1

DAGLα, diacylglycerol lipase alpha

DAGLβ, diacylglycerol lipase beta

eCB, endocannabinoid

HC, hippocampus

IL, infralimbic cortex

mPFC, medial prefrontal cortex

Npas4, neuronal PAS domain protein 4

PrL, prelimbic cortex

PTSD, posttraumatic stress disorder

vHC, ventral hippocampus

Introduction

Experiencing traumatic events can result in the development of posttraumatic stress disorder (PTSD), a severe psychiatric condition characterized by complex symptomatology, including re-experiencing the trauma, avoidance of trauma-related stimuli, emotional and arousal alterations, which symptoms last and fundamentally decrease life quality[1]. In addition, trauma-associated learned fear often becomes generalized, i.e. fear responses are transferred to stimuli unrelated to the previous traumatic event, thus allowing fear responses to occur in safe situations as well[2,3]. As the lifetime prevalence of the disorder is relatively high[4] and its therapy is currently insufficiently resolved[5], its pathophysiology is intensively studied with the ultimate goal of developing effective pharmacotherapeutic interventions. A crucial question in PTSD research is why only 10-25% of trauma-exposed individuals develop PTSD following a traumatic experience, while the majority of the population recover without long-term consequences[6,7]. Identification of the neurobiological basis of risk factors that increase vulnerability to developing PTSD following a traumatic event represents a major clinical challenge with high therapeutic potential.

The endocannabinoid (eCB) signaling system – consisting of cannabinoid receptors. eCBs 2-arachidonoylglycerol (2-AG) and N-arachidonoylethanolamine (anandamide. AEA) and their metabolic enzymes[8–11] – is a major modulator of neuronal plasticity in the central nervous system[12]. The eCB system has recently emerged as a potential target in the therapy of PTSD[13] as i.) its components are abundantly expressed within the neuronal circuits relevant to fear learning and anxiety, such as the medial prefrontal cortex (mPFC), amygdala (AMY) and hippocampus (HC)[14,15]; ii.) it substantially contributes to the regulation of cognitive and emotional functions[16], both significantly altered in PTSD; iii.) its neuropharmacological manipulation in rodent models of PTSD have been shown to affect the behavioral consequences of trauma exposure[17–19]; and iv.) reports on PTSD patients self-medicating with cannabis suggest that symptoms can be ameliorated through activation of the eCB system[20].

While several studies aimed to clarify the involvement of eCB function in different PTSD symptoms by employing means of neuropharmacological approaches in laboratory rodents, our knowledge on the specific trauma-induced changes in the components of the eCB system and their direct contribution to the development of PTSD symptoms is relatively scarce and often contradicting. There is limited information available on the expression patterns of cannabinoid receptor type-1 (CB_1R) in PTSD but the studies addressing the issue conclusively showed that expression is elevated in PTSD patients[21,22]. There are significantly more studies on the levels of eCBs in PTSD, however, findings on this issue are inconclusive

and very few studies addressed the potential link between eCB levels and PTSD vulnerability. Several reports investigating circulating plasma eCB levels in connection with PTSD showed that AEA levels are chronically decreased in human PTSD patients and animal models of trauma exposure when compared to healthy, non-traumatized controls[22–24], although higher levels[25–27] or no detectable changes[28] were also reported. AEA content of PTSD patients (i.e. the vulnerable subpopulation), when compared to trauma-exposed healthy individuals (i.e. the resilient subpopulation) was showed to be lower[22] suggesting that AEA content might predict vulnerability to develop PTSD following trauma exposure. Furthermore, as AEA content negatively correlated with symptom severity in PTSD patients[23,29,30] one might hypothesize that low AEA levels are potentially involved in the pathophysiology of PTSD. In the case of posttraumatic 2-AG levels, studies are rather inconclusive as elevated[26,27,31], decreased[23] and unchanged levels[22,28] were all reported in PTSD patients or traumatized laboratory rodents compared to non-traumatized healthy controls. Regarding vulnerability, the link between 2-AG content and PTSD vulnerability seems similarly inconclusive, as both elevated[26] and decreased[23] serum 2-AG levels were reported in PTSD patients compared to trauma-exposed healthy individuals. Interestingly. deRoon-Cassini and colleagues suggested that 2-AG is differentially involved in PTSD pathophysiology than AEA, as its circulating content is positively correlated with symptom severity[30]. Taken together, while eCB content was shown to be altered in PTSD, the trauma-induced alterations seem to be discrepant, especially when vulnerability issues are not taken into account and PTSD patients are compared with non-traumatized controls. As eCB levels and vulnerability were shown to be linked by several studies and a potential causal connection was suggested by correlations of eCB levels and symptom strength, one might hypothesize that individual differences in eCB function can determine vulnerability to develop trauma-induced behavioral symptoms. The putative connection between eCB function and PTSD vulnerability was suggested by the study of Danan and colleagues on laboratory rodents as well, showing that abnormal eCB reactivity to trauma contributed to subsequent susceptibility to develop a PTSD-like phenotype[32].

In the present study, we employed a rat model of PTSD to assess the putative link between eCB function and PTSD vulnerability in specific regions of the fear circuitry. We focused on generalized fear responses as a core symptom that allows sufficient variability to identify resilient and vulnerable subpopulations. Through this model, our aims were twofold: i) to evaluate disparities in eCB levels among these subgroups, and ii) to scrutinize alterations in gene expression related to the eCB system and their potential association with susceptibility to trauma-induced adverse behaviors. Additionally to eCB-system related genes, we examined

iii.) expression of several immediate early genes, as markers of neuronal activity[33], and modulators of neuroplasticity[34]. Expression of such genes directly contributes to fear learning[35], are involved in stress vulnerability[36] and their expression changes are associated with stress-related mental disorders[37]. Furthermore, they interact with eCB signaling in the modulation of synaptic activity[38]. Finally, leveraging our gene expression data, we employed unsupervised and semi-supervised machine learning-based statistical analysis iv.) to identify patterns of gene expression changes with strongest predictive capacity concerning PTSD vulnerability.

Results

Basal eCB content does not reflect previous trauma exposure in the whole population

To study the short- and long-term effects of trauma exposure on basal levels of 2-AG and AEA in brain regions involved in the regulation of fear responses and memory, we obtained brain samples at either 3 hours or 28 days following fear conditioning (Figure 1A). No significant differences in eCB concentration were observed between groups at the studied time points and brain areas. These findings suggest that basal eCB levels in a particular brain subregion do not reflect previous exposure to a traumatic event at the studied time points when assessed in the whole population (Figure 1B-E). For data of statistical analyses, see Table 1.

Brain subregion-specific eCB content is linked to vulnerability to developing trauma-induced symptoms

In our second experiment, we evaluated the levels of 2-AG and AEA in resilient and vulnerable subpopulations along with non-shocked control subjects, following the testing of fear generalization (Figure 2A). Resilient and vulnerable subpopulations were classified based on the duration spent with freezing behavior in the neutral context (*Context B,* i.e., generalized fear). Subjects in the upper quartile were identified as vulnerable, while those in the lower quartile were categorized resilient. During *Context B*. we observed significant differences in freezing duration between these groups. Post-hoc comparisons revealed that vulnerable individuals froze significantly longer than the resilient and non-shocked groups. In contrast, resilient subjects exhibited no difference in freezing duration compared to non-shocked control subjects (Figure 2D). We found no difference in freezing duration between resilient and vulnerable subjects during contextual fear conditioning (Figure 2B) or contextual reminder (*Context A,* Figure 2C). However, both subpopulations exhibited increased freezing duration compared to non-shocked controls. When we compared all shocked animals

(including the intermediate quartiles) to non-shocked controls, we observed a significant increase in freezing duration during both contextual fear conditioning and contextual reminder (*Context A,* Supplemental Figure 1A and 1B), but interestingly, we detected no significant difference in freezing duration in *Context B* (Supplemental Figure 1C). This lack of difference is possibly due to high variability in generalized fear among subjects previously exposed to footshocks.

Our analysis of eCB content in specific brain regions demonstrated a significant effect of the experimental group on 2-AG levels in the PrL. Post-hoc analysis revealed that resilient subjects exhibited significantly higher 2-AG levels compared to the non-shocked and vulnerable groups. However, we found no differences in AEA content in this region (Figure 2E). In IL, we observed no significant group differences in 2-AG or AEA content (Figure 2F). In the ventral hippocampus (vHC), there was a strong trend towards reduced 2-AG levels in resilient animals (Figure 2G), but AEA levels showed no differences in this brain region (Figure 2G). For data of statistical analyses see Table 2. We found no group differences in the BLA regarding 2-AG or AEA levels (Figure 2H).

We also investigated variations in 2-AG or AEA levels by comparing all subjects who received footshocks (with all shocked subjects pooled together) to non-shocked controls. This investigation did not reveal significant differences at the studied brain sites (Supplemental Figure 1E-H, for data on statistical analyses, see Supplemental Table 1).

As we showed that eCB content reflects vulnerability in our model, we assessed whether populational variability of eCB levels correlate with the variability of generalized fear. We observed a significant positive correlation between total freezing duration and 2-AG concentration in the vHC of subjects who received footshocks (i.e. the whole sample, including 'intermediate' animals). (Figure 3. for data of statistical analyses. see Table 3).

Vulnerability to generalize learned fear is associated with fear circuitry neural hypoactivity and altered neuroplasticity- and eCB system-related gene expression

In our subsequent experiment, we employed RT-qPCR analysis to evaluate the gene expression patterns related to the eCB system in key areas of regulating fear responses in subjects classified as resilient and vulnerable. We targeted 40 genes essential to the eCB system, encompassing receptors, enzymes implicated in eCB synthesis and metabolism, transporters, and proteins involved in eCB signaling pathways. Additionally, we examined the expression of 5 markers of neuronal activation and neuroplasticity to assess the neuronal activity patterns within the circuitry (Figure 5A). The complete list of the 45 evaluated genes is provided in Supplemental Table 2.

Considering the extensive number of parameters from various brain areas that were assessed to elucidate the resilient and vulnerable phenotype, we employed Principal Component Analysis (PCA) for dimension reduction. We found that the first three components explained 57% of the total variance, while each subsequent component accounted for less than 10% (Figure 4A). Therefore, we concentrated on these primary dimensions. The variable loadings of the first three components were distributed in a highly area-specific manner. Specifically, hippocampal genes positively loaded on the first component, amygdalar genes negatively loaded on the second component, and prefrontal genes negatively loaded on the third component of the total variability (Figure 4B). Given that gene expression data from different areas appeared uncorrelated, we implemented additional approaches to further reduce the dimensionality of our data and identify predictors of the resilient and vulnerable phenotype. We applied the machine-learning-based Random Forest (RF) classification to categorize subjects into resilient and vulnerable subpopulations. This allowed us to determine which gene products are the most potent predictors of such a phenomenon. HC *Npas4* expression emerged as the strongest predictor of the phenotype, exhibiting more than four-fold greater variable importance compared to the second most important parameter (Figure 4C top). To address potential bias, we conducted a supplementary analysis excluding Npas4 (AMY) from consideration, revealing additional potential predictors: Abhd12 (mPFC), Daglb (HC), Fos (HC), Fosb (HC), Arc (mPFC), Faah (mPFC), and Slc17a8 (AMY) (refer to Figure 4C bottom). Utilizing these variables, we constructed a model incorporating all subregions, as well as specific models for individual subregions. Upon comparison using receiver operating curve analysis, gene expression analysis in the amygdala, prefrontal and hippocampal demonstrated varying levels of predictive power, with the hippocampal model exhibiting superior performance (Figure 4D). Hypothesis testing methods validated most of these predictors, as our analyses revealed a substantial downregulation of markers of neuronal activity and neuroplasticity associated with altered eCB-related gene expressions in vulnerable compared to resilient subjects across the studied regions (Figure 5B-D). For detailed statistical data, please refer to Table 4.

Discussion

Our findings showed that basal 2-AG and AEA levels in the fear circuitry do not inherently reflect previous traumatic experiences when examined across the entire population. However, upon individual classification into vulnerable or resilient groups regarding the development of trauma-induced fear generalization, distinct trauma-related disparities in eCB content within the PrL and vHC emerged. Specifically, vulnerability was associated with

higher 2-AG content in the PrL and lower in the vHC. Notably, 2-AG levels in the vHC exhibited a positive correlation with the intensity of fear generalization. Moreover, vulnerability to trauma-induced behavioral alterations is associated with marked neuronal hypoactivity in the circuitry involved in the regulation of fear responses, alongside altered expression profiles of genes encoding proteins involved in the modulation of neuroplasticity and eCB signaling. Specifically, these alterations were observed in genes governing eCB metabolism enzymes (*Daglb*, *Faah*, *Abhd12*), receptors (*Grm5*), and neuronal cell surface proteins (*Nlgn3*). Employing unsupervised and semi-supervised machine learning-based statistical approaches, we found that hippocampal gene expression patterns have marked predictive power regarding vulnerability to trauma-induced behavioral alterations.

Our findings underscore the inadequacy of assessing the entire population alone, as it may mask trauma-induced variations in eCB levels in the fear circuitry. Specifically, we have shown that basal levels of 2-AG and AEA in the PrL, IL, vHC, and BLA were unaffected in the short or long-term following a traumatic experience. Though we observed that traumatization resulted in significant long-term contextual fear responses, fear generalization did not differ from the control group, and previous shock exposure did not affect 2-AG or AEA levels in the aforementioned brain regions. It is crucial to avoid misinterpreting these results as indicative of no impact of fear conditioning on eCB signaling. However, this perspective overlooks the considerable variability observed in fear generalization among shocked subjects, a feature previously shown in trauma-exposed individuals to differentiate people with or without subsequent development of PTSD[39]. Our results reflect this phenomenon indicating the heterogeneity of individual responses to the effects of trauma experience and emphasizing the importance of utilizing a nuanced model capable of distinguishing between resilient and vulnerable individuals, an approach gaining more recognition in the field[40,41].

Employing our model of trauma exposure, we were able to generate the necessary variability in trauma-induced generalized fear responses to be able to identify vulnerable and resilient individuals. As the levels of generalized fear showed high variability, these subpopulations markedly differ in this behavioral response. In contrast, neither acute fear responses during traumatic stress exposure, nor long-term contextual fear conditioning responses showed differences between the identified subpopulations. As the ratio of vulnerable subjects in our sample was determined to reflect the proportion of trauma survivors subsequently diagnosed with PTSD (i.e. vulnerable individuals) in the human population[6,7],

we developed a model in which PTSD vulnerability and all its behavioral and neuronal correlates can be studied with high translational validity.

Surprisingly, we detected no discernible differences in AEA content associated with vulnerability in our model. This finding contrasts with studies reporting lower circulating AEA levels in human PTSD patients compared to both healthy individuals and trauma-exposed healthy controls[22]. To reconcile this discrepancy, it is important to consider that circulating endocannabinoid (eCB) levels may not necessarily reflect brain eCB content following release from neurons. Notably, eCBs undergo rapid reuptake processes[42]. Furthermore, it is pertinent to recognize that alterations in AEA levels are not uniformly associated with PTSD, as evidenced by reports of unaltered AEA levels in PTSD patients [23]. Thus, while circulating AEA levels may offer insights in certain contexts, their correlation with PTSD pathology and vulnerability remains an area necessitating further exploration and elucidation.

While our findings seemingly contradict studies in laboratory animals of which the majority reported footshock-induced elevated AEA levels, these works studied the effects immediately after footshock exposure[43], following a reminder to the traumatic context[27] or after extinction training[25]. Furthermore, in the human clinical studies AEA levels were assessed in subjects with the complete PTSD symptom scale while we specifically assessed generalized fear responses as they were the base of our separation of vulnerable subpopulations. Interestingly, while changes in AEA content could not be detected concerning vulnerability, there was significant overexpression of *Faah*, the gene of fatty acid amid hydrolase (FAAH), the main degrading enzyme of AEA in the PrL of vulnerable subjects in comparison to resilient individuals, suggesting that elevated FAAH activity is linked with a tendency to show trauma-induced fear generalization. This finding confirms that FAAH is involved in the manifestation of PTSD-related symptoms as carriers of C385A single nucleotide polymorphism in the *FAAH* gene resulting in less stable FAAH protein with decreased AEA degrading potency[44] showed less intense symptoms[45–47]. Taken together, while our findings regarding no differences between resilient and vulnerable subpopulations in brain AEA content are in contrast with literature findings, this is possibly explained by the source and the time point of sampling. Nevertheless, changes in the expression of the *Faah* gene suggest that AEA signaling might be involved in the basis of vulnerability.

In our investigation of 2-AG content and its association with vulnerability, we have demonstrated that the levels of this eCB were elevated in the PrL of resilient subjects compared to non-shocked or vulnerable individuals. This is particularly interesting in light of

recent work showing that depletion of 2-AG increases fear generalization and neural correlates related to fear generalization within the PrL[48]. As such, the elevated 2-AG signaling we found within the PrL of resilient rats, may enhance neural specificity to fear cues and restrict the development of generalization to promote resiliency. In the vHC, we observed a significant experimental group effect on 2-AG content and a strong, albeit statistically non-significant, trend of lower levels in resilient individuals. Furthermore, we discovered a robust positive correlation between vHC 2-AG content and the strength of generalized fear, suggesting that 2-AG signaling in the vHC is indeed associated with vulnerability to develop trauma-induced generalized fear responses. . This relationship is in opposition to the PrL, reinforcing the importance of examining regional specificity in the manner through which eCB signaling may influence fear processing. This finding is in line with human clinical studies have reported correlations between PTSD symptom severity and circulating eCB content[23,29]. A possible explanation for the elevated vHC 2-AG levels in vulnerable individuals could be the elevated hippocampal expression of *Daglb*, the gene encoding diacylglycerol lipase beta (DAGLβ), one of the synthesizing enzymes of 2-AG. Interestingly, the other 2-AG synthesizing enzyme. diacylglycerol lipase alpha (DAGLα), is also directly linked to stress resilience. Conditional deletion of the *Dagla* in vHC has been shown to increase resilience[49]. Our finding might be intriguing as DAGLβ has long been proposed not to be involved in the retrograde 2-AG signaling route suppressing synaptic transmission[50–52]. However, there is evidence, at least in specific cases, that it indeed contributes to depolarization-induced suppression of excitation mediated by 2-AG[53]. Moreover, *Daglb* expression is reactive to stress, further validating our findings[54].

Intriguingly, in both PrL and vHC, 2-AG levels exhibited a trauma-related pattern. Non-shocked controls and vulnerable individuals showed similar 2-AG content while resilient subjects displayed either elevated or decreased levels in a brain site-dependent manner when compared to these groups. While such a pattern might seem difficult to interpret in terms of vulnerability, as 2-AG levels do not necessarily align with the levels of generalized fear across groups, similar phenomena have been observed regarding vulnerability to stress. For instance, both vulnerable and non-stressed animals show similar expression levels of sphingosine-1-phosphate receptor 3 in the mPFC following chronic social defeat stress, while resilient animals show elevated levels[55]. There are several possible explanations for such a seemingly discrepant pattern. Firstly, 2-AG levels might be differentially involved in the regulation of behavioral responses under physiological and stressed conditions, resulting in similar behavioral phenotypes with different eCB backgrounds. Secondly, vulnerable

individuals might exhibit a deficit in the neuronal regulation of emotional response to a traumatic experience, which cannot suppress further generalized fear response. Lastly, it is important to note that the direction of the difference found in vulnerable compared to resilient subjects is dependent on the brain region, suggesting that the compensatory eCB signaling necessary for healthy behavior observed in resilient subjects has a more general deficit in vulnerable individuals throughout the fear circuitry.

Our unsupervised and semi-supervised machine learning-based statistical analyses of gene expression data in resilient and vulnerable subjects have underscored the substantial role of the HC in vulnerability. Both approaches confirmed that gene expression patterns in this region possess superior predictive power compared to other assessed areas. Notably, the HC expression of *Npas4*, the gene of transcription factor neuronal PAS domain protein 4 (Npas4). a modulator of neuronal plasticity and marker of neuronal activity, had a predictive power four times greater than any other assessed gene. This gene exhibited a marked downregulation in the vulnerable compared to resilient subpopulations across all assessed areas, particularly in the HC. Such downregulation of *Npas4* suggests HC hypoactivity, closely linked to vulnerability. This finding aligns with human clinical data showing that lower HC activation can predict PTSD symptom severity in patients[56]. Furthermore. *Npas4* expression in the HC was shown to be crucial in context coding in a rodent model of PTSD[57], suggesting that downregulation of the gene might lead to a deficit in context coding and contribute to the overgeneralization of fearful responses in contexts not associated with the previous traumatic experience. Indeed. *Npas4* downregulation in several brain regions in the fear circuitry was shown to be associated with stress-related psychopathologies[37]. Decreased expression of *Arc*, another immediate early gene that similarly to *Npas4* showed downregulation in our model, was shown to specifically increase susceptibility to trauma-induced adverse behavioral changes[36]. Most importantly, Npas4 was shown to directly interact with eCB signaling in the HC, promoting eCB-mediated depolarization-induced suppression of inhibition in the regulation of neuronal plasticity[38]. Taken together, one might hypothesize that a substantial deficit in Npas4-mediated mechanisms linked to changes in eCB signaling fundamentally alters neuronal plasticity, leading to abnormal processing of traumatic experiences, impaired contextual coding, and ultimately, generalized fear responses.

Our findings reveal that subpopulations differing in their vulnerability to developing PTSD-like symptoms show anatomically specific differences in eCB content. Specifically, we have highlighted that eCB changes in the vHC are crucial for vulnerability, as 2-AG levels in this region showed significant correlations with the strength of generalized fear and are

associated with neuronal hypoactivity and decreased expression of crucial mediators of neuronal plasticity. Patterns of gene expression related to neuroplasticity and eCB signaling in the HC had substantial predictive power for PTSD vulnerability. Our results describe a novel neuropharmacological marker that outlines vulnerability to developing PTSD following trauma exposure and may contribute to the identification of clinically relevant subpopulations.

Limitations of the study

Although, our laboratory has previously showed that several symptom domains of PTSD can be modeled employing electric footshocks in rodents[58,60,62], our current work only assessed generalized fear and separation of resilient and vulnerable subpopulations was based on this core symptom of PTSD. Further studies are needed to evaluate whether other aspects of trauma-induced behavioral changes reflect vulnerability. Furthermore, only male subjects were used in our experiments.

Author contributions

Conceptualization, M.A., L.Sz., E.M., M.T., and M.N.H.; Methodology, L.Sz., M.A.; Investigation, L.Sz., M.A., Gy.Y.B., B.T., L.B., and Z.B.; Data Curation, L.Sz.; Formal Analysis, L.Sz., V.Z.K.; Writing – Original Draft Preparation, M.A., L.Sz.; Writing – Review & Editing Preparation, L.Sz., Gy.Y.B., Z.K.V., B.T., L.B., Z.B., M.N.H., M.T., E.M, and M.A.; Visualization Preparation, L.Sz., M.A., and V.Z.K.; Supervision, M.A. and E.M.; Project Administration, M.A.; Funding Acquisition, M.A., E.M., M.T., and M.N.H.

Acknowledgments

The project was funded by National Research, Development and Innovation Office grant FK128191 (M.A.), Hungarian Brain Research Program grant 2017-1.2.1-NKP-2017-00002 (E.M.), Project RRF-2.3.1-21-2022-00011, titled National Laboratory of Translational Neuroscience has been implemented with the support provided by the Recovery and Resilience Facility of the European Union within the framework of Programme Széchenyi Plan Plus (E.M.) and the Visiting fellowship Program of the Hungarian Academy of Sciences (M.A. and M.N.H.). We thank the Behavioral Studies Unit core facility of the Institute of Experimental Medicine Hungarian Research Network and the technical assistance of D. Ömböliné and B. Molnar.

Declaration of interests

The authors declare no competing interests.

Figure legends

Figure 1. Basal endocannabinoid content does not reflect previous trauma exposure in the whole population. **A**) Experimental design schematic showing experimental groups: control subjects left undisturbed in their homecage without undergoing behavioral testing, non-shocked control subjects exposed to the footshock context without delivery of footshocks, and subjects exposed to footshocks. Subjects were sacrificed under baseline conditions from their home cage at either 3 h or 28 days after footshock/context exposure. Brain samples from prelimbic cortex (PrL), infralimbic cortex (IL), ventral hippocampus (vHC) and basolateral amygdala (BLA) were collected for HPLC/MS/MS analysis of 2-arachidonoylglycerol (2-AG) and N-arachidonoylethanolamin (AEA) levels. **B-E**) Neither footshock delivery, nor context exposure led to changes in 2-AG or AEA levels in the assessed regions. Data are presented as mean ± SEM. One-way ANOVA.

Figure 2. Brain subregion-specific endocannabinoid content is linked to vulnerability to developing trauma-induced symptoms. A) Experimental design schematic showing our model of trauma exposure. Following contextual fear conditioning (Day 0), contextual fear memory expression was assessed 28 days later in the traumatic context (*Context A*. Day 28). The next day, fear generalization was studied in a novel, safe context (*Context B*. Day 29). Resilient and vulnerable subjects were categorized based on the upper and lower quartiles of generalized fear characterized by freezing duration in *Context B*. Subjects were immediately sacrificed after *Context B* testing, and brain samples from prelimbic cortex (PrL), infralimbic cortex (IL), ventral hippocampus (vHC) and basolateral amygdala (BLA) were collected for HPLC/MS/MS analysis of 2-arachidonoylglycerol (2-AG) and N-arachidonoylethanolamin (AEA) levels. Non-shocked controls underwent the same procedure without the delivery of footshocks. **B**) Contextual fear conditioning induced freezing behavior during the conditioning session. (**C**) which remained high during *Context A* contextual reminder in resilient and vulnerable groups compared to the non-shocked control. **D**) The vulnerable group showed markedly high fear generalization, indicated by the higher freezing responses compared to resilient or non-shocked control groups. **E**) The resilient group exhibited higher levels of 2-AG in the PrL than vulnerable and non-shocked control groups, without changes in AEA levels. **F**) 2-AG and AEA levels did not show differences in the IL. **G**) In the vHC, there was a significant main effect in the 2-AG levels without significant differences in pairwise

comparisons, although 2-AG showed a tendency to decrease in resilient subjects. AEA levels did not differ between groups in this region. **H**) 2-AG and AEA levels in the BLA did not show differences. *: $p < 0.05$. **: $p < 0.01$. ***: $p < 0.001$; Data are presented as mean ± SEM. One-way ANOVA.

Figure 3. 2-arachidonoylglycerol (2-AG) levels in the ventral hippocampus correlate with the strength of footshock-induced generalized fear responses. In the ventral hippocampus (vHC), there was a significant positive correlation between 2-AG levels and the time spent freezing in *Context B*. Pearson correlation test.

Figure 4. Expression of neuronal activity gene markers, neuroplasticity and endocannabinoid system-related genes predict vulnerability to generalize learned fear. **A**) Scree plot of the principal component analysis (PCA) involving all assessed 40 gene expression data of all 4 investigated brain areas. Eigen values show the proportion of variance of generalized fear in *Context B* explained by each component. Red dashed line indicates the last component that explains more than 10% of the total variability. **B**) Variable loadings on the main components (PC1-3, principal components 1-3). Variable-component correlations show an area-specific expression pattern. C) Variable importance levels in two saturated Random Forest models with (top, model A) and without (bottom, model B) the hippocampal *Npas4* expression data. D) ROC curves and the area under the curve (AUC) values of the reduced models involving the significant predictors of the B model of all (black) or in a given brain subregion (red: medial prefrontal cortex (mPFC); green: hippocampus (HC); blue: amygdala (AMY).

Figure 5. Vulnerability to generalize learned fear is associated with fear circuitry neural hypoactivity and altered neuroplasticity- and endocannabinoid system-related gene expression. A) Experimental design schematic showing our model of trauma exposure. Following contextual fear conditioning (Day 0), expression of contextual fear memory was assessed 28 days later in the traumatic context (*Context A*, Day 28). The next day, fear generalization was studied in a novel, safe context (*Context B*, Day 29). Resilient and vulnerable subjects were categorized based on the upper and lower quartiles of the generalized fear spectrum characterized by freezing duration in *Context B*. Subjects were immediately sacrificed after *Context B* testing, and brain samples from the medial prefrontal cortex (mPFC), hippocampus (HC), and amygdala (AMY) were collected to analyze

endocannabinoid (eCB) system-related gene expression using RT-qPCR. **B-D**) Gene expression differences and statistical significance levels of the vulnerable group compared to the resilient group displayed by volcano plots in the mPFC (**B**), HC (**C**), and AMY (**D**). The horizontal blue lines represent the criteria of statistical significance $p = 0.05$. Genes that exhibit a fold change greater than 1.2 are denoted by green dots for downregulation and red dots for upregulation. Vertical dashed lines represent the margin of 1.2 fold change.

Tables and legends

Table 1. Statistical data of *Experiment 1*. 2-AG: 2-arachidonoylglycerol; AEA: N-arachidonoylethanolamine; PrL: prelimbic cortex; IL: infralimbic cortex; vHC: ventral hippocampus; BLA: basolateral amygdala; ANOVA.

sampling time point	area	2-AG F(DFn, DFd)	p-value	AEA F(DFn, DFd)	p-value
3h	PrL	F(2, 31) = 2.972	0.066	F(2, 31) = 0.604	0.552
	IL	F(2, 31) = 1.056	0.359	F(2, 33) = 0.215	0.807
	BLA	F(2, 32) = 0.0677	0.934	F(2, 32) = 1.154	0.328
	vHC	F(2, 33) = 1.933	1.933	F(2, 33) = 1.187	0.317
28 days	PrL	F(2, 33) = 1.187	0.461	F(2, 26) = 1.569	0.227
	IL	F(2, 24) = 1.176	0.325	F(2, 24) = 1.241	0.307
	BLA	F(2, 24) = 0.710	0.501	F(2, 24) = 0.964	0.395
	vHC	F(2, 26) = 0.810	0.455	F(2, 26) = 0.222	0.801

Table 2. Statistical data of *Experiment 2*. 2-AG: 2-arachidonoylglycerol; AEA: N-arachidonoylethanolamine; PrL: prelimbic cortex; IL: infralimbic cortex; vHC: ventral hippocampus; BLA: basolateral amygdala; *: significant group effect. ANOVA.

	test / area	F(2, 11)	p-value
freezing duration	**conditioning**	**57.97**	**< 0.0001***
	Context A	**67.99**	**< 0.0001***
	Context B	**74.25**	**< 0.0001***
2-AG	**PrL**	**7.099**	**0.011***
	IL	0.409	0.674
	vHC	**4.026**	**0.048***
	BLA	0.564	0.584
AEA	PrL	1.135	0.356
	IL	0.176	0.840
	vHC	0.176	0.840
	BLA	1.021	0.391

Table 3. Statistical data of correlation analyses between generalized fear and endocannabinoid content in *Experiment 2*. 2-AG: 2-arachidonoylglycerol; AEA: N-arachidonoylethanolamine; PrL: prelimbic cortex; IL: infralimbic cortex; vHC: ventral hippocampus; BLA: basolateral amygdala; *: significant correlation. Pearson correlation coefficients.

	area	Pearson r	p-value
freezing duration in context B vs. 2-AG cc.	PrL	-0.165	0.498
	IL	-0.128	0.601
	vHC	**0.605**	**0.006***
	BLA	-0.304	0.205
freezing duration in context B vs. AEA cc.	PrL	-0.301	0.210
	IL	-0.098	0.689
	vHC	-0.195	0.423
	BLA	-0.164	0.500

Table 4. Statistical data of gene expression analysis in *Experiment 3*. *Cnr2* showed consistently low expression at all assessed regions, therefore this gene was excluded from analysis. mPFC: medial prefrontal; HC: hippocampus; AMY: amygdala; *: significant group effect. Multiple t-test.

	mPFC		HC		AMY	
gene	t(13)	p-value	t(13)	p-value	t(13)	p-value
Abhd2	1.955	0.072	0.223	0.827	0.081	0.937
Abhd6	0.518	0.613	0.660	0.521	0.319	0.755
Abhd12	**2.198**	**0.047***	1.783	0.098	0.472	0.645
Adam17	0.467	0.649	0.582	0.571	0.879	0.395
Alox5	0.440	0.667	0.112	0.912	0.906	0.381
Alox15	0.917	0.376	0.468	0.647	0.052	0.959
Arc	**2.773**	**0.016***	0.984	0.343	0.797	0.440
Cacna1a	1.411	0.182	1.616	0.130	0.427	0.676
Cacna1b	1.528	0.151	0.352	0.730	0.023	0.982
Cnr1	2.034	0.063	1.382	0.190	0.233	0.819
Cnrip1	0.120	0.907	1.001	0.335	0.572	0.577
Dag1	1.307	0.214	0.797	0.440	0.632	0.538
Dagla	1.421	0.179	1.173	0.262	0.215	0.833
Daglb	0.387	0.705	**2.584**	**0.023***	1.667	0.119
Faah	**2.216**	**0.045***	0.864	0.403	0.012	0.990
Fabp5	1.619	0.130	0.558	0.587	0.076	0.941
Fos	0.934	0.368	1.962	0.071	1.878	0.083
Fosb	**2.264**	**0.041***	1.785	0.098	0.373	0.715
Gnai1	0.700	0.496	0.484	0.636	0.297	0.771
Gnai2	1.409	0.182	0.519	0.613	0.587	0.567
Gpr3	0.608	0.554	0.275	0.787	0.346	0.735
Gpr6	1.108	0.288	0.131	0.898	1.350	0.200
Gpr12	0.624	0.544	0.900	0.384	0.531	0.604
Gpr55	0.598	0.560	NA	NA	1.523	0.154
Grm5	**2.185**	**0.048***	0.609	0.553	0.520	0.612
Homer1	1.031	0.321	0.984	0.343	0.769	0.456
Homer2	1.830	0.090	0.346	0.735	0.313	0.760
Mgll	1.230	0.240	0.769	0.456	0.150	0.883
Mmp9	0.417	0.683	1.326	0.208	0.606	0.555
Napepld	0.855	0.408	1.504	0.156	0.069	0.946
Nat1	1.300	0.216	0.099	0.923	0.952	0.358
Nlgn3	0.424	0.679	0.062	0.951	**2.338**	**0.036***
Npas4	**2.388**	**0.033***	**3.362**	**0.005***	**4.582**	**0.001***
Nptx2	1.134	0.277	0.409	0.689	0.804	0.436
Nrxn1	1.257	0.231	0.889	0.390	0.031	0.975
Nucb1	1.893	0.081	0.686	0.505	0.854	0.409
Plcb1	0.979	0.346	0.969	0.350	1.294	0.218
Plcg1	0.358	0.726	0.344	0.736	0.016	0.988
PPara	0.223	0.827	0.348	0.734	0.097	0.924
Ptgs2	1.153	0.270	0.132	0.897	0.151	0.882
Ptpn22	2.084	0.057	0.032	0.975	0.298	0.770
Slc17a8	1.015	0.329	0.390	0.703	1.944	0.074
Trpv1	1.726	0.108	0.587	0.567	0.327	0.749

STAR Methods

Subjects

Subjects were adult male Long-Evans rats obtained from a breeding colony at the Medical Gene Technology unit of the Institute of Experimental Medicine established from rats from Charles River Laboratories (Italy). Animals were kept in Eurostandard Type IV cages measuring 59.8 cm x 38 cm x 20 cm. in groups of 3-4 rats per cage. In order to avoid

stress contagion (stressed subject affecting the behavior of non-stressed controls) and social buffering (non-stressed control subjects affecting the behavior of stressed subjects) all subjects within the same cage underwent identical treatment (either fear conditioning or exposure to the context without traumatization. see Behavioral Procedures for details). In the animal rooms, temperature was maintained at 22±1°C, and relative humidity was kept at 20±10%. Laboratory chow and water were provided *ad libitum*. Rats were kept on a reverse 12-hour light/dark cycle with lights turning on at 8 p.m. and turning off at 8 a.m. The experiments followed the guidelines of the European Communities Council Directive of November 24. 1986 (86/609/EEC) and were conducted under license PE/EA/141-2/2020 issued by the Pest County Government Office.

Behavioral procedures

All behavioral procedures were carried out in the first part of the dark phase in dedicated experimental rooms by experimenters blind to experimental groups. Behavioral procedures were video recorded with Sony FDR-AX33 digital camcorders for later analysis.

Contextual fear conditioning

Subjects were transported to a brightly lit experimental room (300 lux) and placed in a transparent Plexiglas chamber (measuring 40 x 40 x 40 cm) with a stainless steel electric grid floor. Following a 2.5 minute habituation period, a series of 10 scrambled electric footshocks were administered through the grid floor, each lasting 2 seconds and with an amperage of 2.4 mA, and separated by 30 second long periods between each shock. The conditioning session lasted for a total of 7 minutes. Non-shocked control subjects were introduced to the test chamber for 7 minutes without footshock delivery. Chambers were cleaned with 20% ethanol and dried with paper towel after each subject.

Testing for contextual conditioned fear

28 days after footshock delivery. subjects were exposed to a chamber identical to the one used for conditioning (*Context A*) under the same experimental conditions without further footshock delivery for 5 minutes to assess contextual conditioned fear responses. The 28th day as the time point when behavior was first assessed was selected in order to model the lasting trauma-induced adverse behavioral changes in PTSD[1].

Testing for generalization of contextual fear

29 days after footshock delivery (24 hours after *Context A* exposure), subjects were placed in a novel neutral context (*Context B*) for 20 minutes. *Context B* was carefully designed to differ entirely from *Context A*. utilizing different visual, olfactory, and tactile

stimuli: i.) the experimenter was different from the one conducting conditioning and *Context A* exposure. ii.) a different experimental room was used, illuminated with dim red light (5 lux). iii.) plastic inserts were placed over the grid floor and the walls to modify shape, floor and wall color of the chambers. iv.) chambers were cleaned between subjects using soapy water.

Behavioral analysis

Video recordings taken during behavioral procedures were analyzed using the Noldus Ethovision XT15 (Noldus Information Technology. The Netherlands) software, in which the activity monitoring feature was used to quantify fear responses by calculating the percentage of time spent freezing, a typical fear response in rodents[59]. This feature is based on an automated, frame-by-frame analysis of behavior video recordings during which the algorithm compared the pixel differences between subsequent frames. Pixel differences below a specific *a priori* set threshold represented the immobility of the subject (i.e., freezing behavior). Thresholds were constant across subjects within an experiment and were compared to and validated by previous manual behavior analyses of randomly selected subjects conducted by trained experimenters blind to treatment groups.

Identification of resilient and vulnerable subpopulations

Generalized fear responses in a neutral context measured by the percentage of time spent freezing during the *Context B* test were used to identify resilient and vulnerable individuals. Considering the estimated proportion of patients diagnosed with PTSD and healthy trauma-exposed individuals in the populations[6,7], we characterized a resilient and vulnerable subpopulation as the upper and lower quartile of the population based on their fear generalization. This approach allowed for a clear distinction between the two groups and aided in further analysis of their behavior and physiological functions. Animals that fall between vulnerable and resilient categories were classified as intermediate individuals.

High-performance liquid chromatography-tandem mass spectrometry (HPLC/MS/MS)

Subjects were rapidly decapitated, and their brain was manually removed. Brains were cooled on wet ice and were dissected into 2 mm thick coronal slices on a stainless steel brain mold. Tissue blocks containing relevant areas in the neurobiological basis of PTSD, the prelimbic (PrL) and infralimbic (IL) cortices of the mPFC, basolateral amygdala (BLA). ventral hippocampus (vHC) were identified according to the atlas of Paxinos and Watson[61] based on clearly visible anatomical reference points (e.g. forceps minor and piriform cortex

for the mPFC; the internal and external capsule as well as the optic tract, and the commissure of the stria terminalis for the BLA). In addition to these landmarks, subtle color patterns that circumscribed particular brain regions also guided dissection. The sampled regions were dissected bilaterally and pooled within each subject. Samples were moved into sterile Eppendorf tubes and snap-frozen on dry ice. Tissue samples were kept at -80°C until later analysis. Sample weight was calculated from the difference in tube weight measured before and after sampling using an analytical balance (Kern ABJ 80-4NM. Kern-Sohn GmbH., Germany). Sample preparation and eCB measurement was conducted as described earlier by Lee and colleagues[63]. First, tissue samples were incubated on ice for 30 min in 1 ml of HPLC-gradient grade methanol (Merck KGaA, Darmstadt, Germany) containing the deuterated internal standards 2-arachidonoylglycerol-d5 (100 ng/ml) and arachidonoylethanolamide-d (1 ng/ml) (Cayman Chemical, Ann Arbor, MI, USA). Samples were homogenized and centrifuged on an Eppendorf miniSpin microtube centrifuge at 13,400 rpm for 15 min. The supernatant was transferred into a plastic tube and was diluted with 3ml of ultrapure water. The samples were subjected to solid-phase extraction (SPE) sample clean-up according to the following protocol. First, the SPE cartridges (SUPELCO Discovery DSC-18 1 ml tubes, 100 mg) were conditioned with 2 ml of methanol and 2 ml of ultrapure water. Diluted samples were loaded at approximately 0.5 ml/min flow rate to apply a gentle vacuum. In the consecutive washing step, 2 ml of ultrapure water and 2 ml of methanol/water (50:50 v/v%) were percolated through the sorbent material. Elution was performed with 0.7 ml of methanol. Eluates were diluted to initial HPLC eluent composition with 10 mM ammonium formate (Honeywell Riedel-de Haen GmbH, Seelze, Germany) solution. To measure eCB levels, a PerkinElmer Life and Analytical Sciences HPLC Series 200 system was employed coupled to an Applied Biosystems/Sciex 4000 QTRAP triple quadrupole/linear ion trap tandem mass spectrometer operated in positive electrospray ionization mode. The electrospray ionization ion source parameters were set as follows: curtain gas, 10; ion spray voltage, 5000 V; temperature, 500°C; collisionally activated dissociation gas, medium; gas 1, 50; gas 2, 40. Chromatographic separation was achieved with a Phenomenex Kinetex C18 column (50mm x 3.00 mm, 2.6 x m, 100 Å) using methanol and 10 mM ammonium formate as elution solvents at a flow rate of 500 µl/min and injection volume of 50 µl. The initial eluent condition was 80% methanol/20% buffer that was increased to 85% organic phase during 3 min and then further elevated to 95% during 2 min and was kept at this condition for 2 min. Afterward, the column was equilibrated to the initial condition. Analytes were detected in multiple reaction monitoring (MRM) mode at the following ion transitions and parameter settings: (1)2-AG,

MRM transition [mass/charge ratio (m/z), 379.4 □ 287.2, 379.4 □ 91.1], declustering potential (81 V), collision energy (23 V, 81 V), collision cell exit potential (10 V, 8 V); (2) 2-arachidonoylglycerol-d5, MRM transition (m/z, 384.4 □ 287.2, 384.4 □ 91.1), declustering potential (81 V), collision energy (23 V, 81 V), collision cell exit potential (10 V, 8 V); (3) arachidonoylethanolamide, MRM transition (m/z, 348.4 □ 62.1, 348.4 □ 90.9), declustering potential (51 V), collision energy (43 V, 63 V), collision cell exit potential (4 V, 8 V); and (4) arachidonoylethanolamide-d4, MRM transition (m/z, 352.4 □ 66.0, 352.4 □ 91.2), declustering potential (81 V), collision energy (41 V, 77 V), collision cell exit potential (6 V). The peak areas were determined with Analyst 1.4.2. software. The quantity of the analytes was calculated by comparing their peak areas with those of the deuterated internal standards, and normalized to sample weight.

Real-Time quantitative PCR (RT-qPCR)

Rats were rapidly decapitated, and the brains were promptly excised from the skull, cooled using ice, and maintained in an RNase-free environment. Based on the atlas of Paxinos and Watson[61], a 2 mm thick coronal section containing the respective mPFC and the AMY region was sliced on a stainless steel mold. Whole HC samples were collected using ophthalmic tweezers. The mPFC, AMY, and HC samples were dissected promptly, placed in sterile Eppendorf tubes, and snap-frozen on dry ice. The tissue samples were stored at -80°C until later analysis.

The samples were moved into QIAzol lysis reagent (Qiagen. The Netherlands) and homogenized using a Tissue-Tearor (BioSpec Products. USA). The total RNA extraction was carried out using the RNeasy Lipid Tissue Mini Kit (Qiagen. The Netherlands) according to the manufacturer's protocol. All samples were treated with RNase Free DNase Set (Qiagen. The Netherlands) to remove genomic DNA contamination. The Qubit 4 Fluorometer (Invitrogen. USA) was used to obtain information about the quantity (Qubit RNA BR Assay Kit; Invitrogen. USA) and quality (Qubit RNA IQ Assay Kit; Invitrogen. USA) of the RNA in the samples. The total RNA samples were diluted in nuclease-free water to a concentration of 0.1 µg/µl. using 1 µg of RNA to perform reverse transcription.

Each sample underwent reverse transcription of 1 µg of total RNA using random primers and the High Capacity cDNA reverse transcription kit (Thermo Fisher Scientific. USA). Reverse transcription was performed on a VWR Ristretto thermal cycler (VWR International GmbH. Germany). The cDNA concentrations were measured using a Qubit ssDNA assay kit (Invitrogen. USA) on a Qubit 4 Fluorometer.

The cDNA samples were diluted in nuclease-free water and then combined with TaqMan® Gene Expression Master Mix to create a PCR reaction mix, resulting in a final concentration of 1 ng/µl of cDNA. The expression of 45 genes associated with either the eCB system or neuronal activity and 3 housekeeping genes were analyzed using Custom 384-well microfluidic TaqMan® Gene Expression Array Cards (Applied Biosystems. USA). The array cards consisted of eight fill reservoirs linked to a series of 48 reaction wells. These wells contained dried-down TaqMan® assays for 48 unique genes. PCR reaction mixes were loaded and centrifuged twice at 331 x g for 1 minute. The main distribution channels of the array card were then sealed with a TaqMan® Array Card Sealer to preserve the mixtures. A ViiA™ 7 Real-Time PCR System (Applied Biosystems. USA) was used to perform RT-qPCR. The cycle threshold (Ct) values and other data related to PCR reactions were collected using the QuantStudio software (Applied Biosystems. USA). The gene expression levels for the candidate genes were normalized to the geometric mean of expression levels of the three endogenous control genes (*18S*, *Actb*, and *Gapdh*). The relative quantity was then calculated using the $2^{-\Delta\Delta Ct}$ method[64].

Experimental design

In *Experiment 1*, we evaluated the variations in AEA and 2-AG levels over time after a traumatic experience by assessing the short- (3 hours after footshocks) and long-term (28 days after footshock) effects of shock exposure. Subjects were assigned to experimental groups *home cage control*, *non-shocked control,* and *shocked*. Following footshock delivery, shocked and non-shocked control subjects were returned to their home cages and sacrificed either 3 hours or 28 days later under baseline conditions. Home cage control subjects not exposed to the footshock apparatus were sacrificed at each time point as well. Brain punch samples from PrL, IL, vHC, and BLA were collected for eCB content assessment by HPLC/MS/MS.

In *Experiment 2*, we assessed the long-term effects of traumatic experiences on AEA and 2-AG levels in connection with resilience/vulnerability to develop trauma-induced generalized fear responses, a core symptom of PTSD. We subjected rats to contextual fear conditioning, and then 28 days later, a contextual reminder (*Context A*) was performed to assess conditioned fear responses. On the 29th day post-conditioning, subjects were exposed to a different context (*Context B*) to evaluate fear generalization. After *Context B* exposure, subjects were immediately sacrificed, and brain punch samples from PrL, IL, vHC, and BLA were collected. AEA and 2-AG levels were assessed by HPLC/MS/MS in resilient and vulnerable subpopulations that were identified *a posteriori* based on generalized fear

responses in *Context B*. Additionally, a cohort of non-shocked control subjects underwent exposure to Contexts A and B without the presentation of footshocks on the initial day of conditioning.

In *Experiment 3*, we assessed trauma-induced expression patterns in eCB system-associated genes in connection with resilience/vulnerability to develop trauma-induced generalized fear responses. Following contextual fear conditioning and subsequent exposure to *Context A* and *Context B*, subjects were classified into vulnerable and resilient groups, as outlined in *Experiment 2*. Rats were sacrificed immediately following *Context B* exposure, and brain punch samples were collected from the mPFC, HC, and AMY regions. Punch samples were used for RT-qPCR analysis of the expression of eCB system-related genes.

Statistical analysis

Descriptive statistics and hypothesis testing

Behavioral and eCB level data are shown as mean ± standard error of the mean. eCB content in *Experiment 1* was evaluated by one-factor analysis of variance (ANOVA) (factor: *experimental group*). Time spent with freezing during conditioning and contextual reminders as well as brain eCB content in *Experiment 2* were evaluated by one-factor ANOVA (factor: *experimental group*). Tukey's multiple comparisons tests were performed for post-hoc analyses when a main effect was significant. Correlations between brain eCB levels and time spent freezing during *Context B* exposure were evaluated by the Pearson correlation coefficient. For the analyses including all individuals in *Experiment 2*, the Mann-Whitney U tests were performed. For the analyses of PCR data in *Experiment 3*, we used multiple unpaired t-tests. P-values lower than 0.05 were considered statistically significant. All statistical analyses above were conducted with GraphPad Prism software version 9.5.1 (MA. USA).

Principal component analysis

To assess the structure and correlatedness of the expression data of different brain regions, we conducted a principal component analysis (PCA) using the varcomp R package. We calculated the proportion of the variance explained by each principle component to see the dimensionality of the expression data. We also calculated the loadings of gene expression variables on each principle component to reveal their correlatedness.

Random Forest classification

To decrease the dimensions of our multivariate data and identify the gene expression predictors of resilient and vulnerable subpopulations, we applied the semi-supervised machine

learning method. Random Forest (RF)[65]. RF ranks the importance of variables in the classification of subjects to subpopulations, in this particular case, to resilient and vulnerable subgroups. It can create thousands of decision trees and splits subjects into groups according to the expression of a random subset of the genes. This way RF is suitable for classifying samples that have many features, even with limited sample sizes. Here. i) we developed a saturated model using all gene expression data of all investigated brain subregions and ii) ranked the genes according to their importance. Then we iii) created subset models with the most important variables, one in which all areas were included and three in a subregion-specific manner. Finally. iv) we compared these subset models according to their classification accuracy. Note that, due to the outstanding prediction accuracy of the gene *Npas4* expression in the HC and that RF can be biased by features with the impact of this level, we conducted two separate analyses, one with and one without the inclusion of HC *Npas4* data. All models produced 5.000 decision trees and considered 9 genes at a time. Sampling was balanced to 7-7 subjects per group in the RF analysis. Variable importance was ranked based on permutation importance, which shows the mean decrease in prediction accuracy of a feature in response to randomly permuting its values.

Supplemental information legends

Supplemental Figure 1. Endocannabinoid levels show no trauma-induced changes in the fear circuitry when assessed in the whole sample regardless of vulnerabilities. A) Experimental design schematic showing our model of trauma exposure. Following contextual fear conditioning (Day 0), contextual fear memory expression was assessed 28 days later in the traumatic context (*Context A*. Day 28). The next day, fear generalization was studied in a novel, safe context (*Context B*. Day 29). Subjects were immediately sacrificed after *Context B* testing, and brain samples from prelimbic cortex (PrL), infralimbic cortex (IL), ventral hippocampus (vHC) and basolateral amygdala (BLA) were collected for HPLC/MS/MS analysis of 2-arachidonoylglycerol (2-AG) and N-arachidonoylethanolamin (AEA) levels. Non-shocked controls underwent the same procedure without the delivery of footshocks. **B**) Contextual fear conditioning induced freezing behavior during the conditioning session. (**C**) which remained high during Context A contextual reminder. **D**) The shocked group showed no difference in fear generalization compared to non-shocked control groups. **E-H**) No differences were detected in eCB levels at the assessed areas between shocked and non-shocked groups. ***: $p < 0.001$; Data are presented as mean ± SEM. One-way ANOVA.

References

1. American Psychiatric Association (2022). Diagnostic and Statistical Manual of Mental Disorders (American Psychiatric Association Publishing) 10.1176/appi.books.9780890425787.

2. Lis, S., Thome, J., Kleindienst, N., Mueller-Engelmann, M., Steil, R., Priebe, K., Schmahl, C., Hermans, D., and Bohus, M. (2020). Generalization of fear in post-traumatic stress disorder. Psychophysiology *57*, e13422. 10.1111/psyp.13422.

3. Sangha, S., Diehl, M.M., Bergstrom, H.C., and Drew, M.R. (2020). Know safety, no fear. Neurosci Biobehav Rev *108*, 218–230. 10.1016/j.neubiorev.2019.11.006.

4. Kessler, R.C., Chiu, W.T., Demler, O., Merikangas, K.R., and Walters, E.E. (2005). Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. Arch Gen Psychiatry *62*, 617–627. 10.1001/archpsyc.62.6.617.

5. Davis, L.L., Frazier, E.C., Williford, R.B., and Newell, J.M. (2006). Long-term pharmacotherapy for post-traumatic stress disorder. CNS Drugs *20*, 465–476. 10.2165/00023210-200620060-00003.

6. Breslau, N., Davis, G.C., Andreski, P., and Peterson, E. (1991). Traumatic events and posttraumatic stress disorder in an urban population of young adults. Arch Gen Psychiatry *48*, 216–222. 10.1001/archpsyc.1991.01810270028003.

7. Breslau, N., Kessler, R.C., Chilcoat, H.D., Schultz, L.R., Davis, G.C., and Andreski, P. (1998). Trauma and posttraumatic stress disorder in the community: the 1996 Detroit Area Survey of Trauma. Arch Gen Psychiatry *55*, 626–632. 10.1001/archpsyc.55.7.626.

8. Devane, W.A., Breuer, A., Sheskin, T., Järbe, T.U., Eisen, M.S., and Mechoulam, R. (1992). A novel probe for the cannabinoid receptor. J Med Chem *35*, 2065–2069. 10.1021/jm00089a018.

9. Di Marzo, V., De Petrocellis, L., Bisogno, T., and Melck, D. (1999). Metabolism of anandamide and 2-arachidonoylglycerol: an historical overview and some recent developments. Lipids *34 Suppl*, S319-25. 10.1007/BF02562332.

10. Mechoulam, R., Ben-Shabat, S., Hanus, L., Ligumsky, M., Kaminski, N.E., Schatz, A.R., Gopher, A., Almog, S., Martin, B.R., and Compton, D.R. (1995). Identification of an endogenous 2-monoglyceride, present in canine gut, that binds to cannabinoid receptors. Biochem Pharmacol *50*, 83–90. 10.1016/0006-2952(95)00109-d.

11. Munro, S., Thomas, K.L., and Abu-Shaar, M. (1993). Molecular characterization of a peripheral receptor for cannabinoids. Nature *365*, 61–65. 10.1038/365061a0.

12. Freund, T.F., Katona, I., and Piomelli, D. (2003). Role of endogenous cannabinoids in synaptic signaling. Physiol Rev *83*, 1017–1066. 10.1152/physrev.00004.2003.

13. Hill, M.N., Campolongo, P., Yehuda, R., and Patel, S. (2018). Integrating Endocannabinoid Signaling and Cannabinoids into the Biology and Treatment of

Posttraumatic Stress Disorder. Neuropsychopharmacology *43*, 80–102. 10.1038/npp.2017.162.

14. Herkenham, M., Lynn, A.B., Little, M.D., Johnson, M.R., Melvin, L.S., de Costa, B.R., and Rice, K.C. (1990). Cannabinoid receptor localization in brain. Proc Natl Acad Sci U S A *87*, 1932–1936. 10.1073/pnas.87.5.1932.

15. Matsuda, L.A., Lolait, S.J., Brownstein, M.J., Young, A.C., and Bonner, T.I. (1990). Structure of a cannabinoid receptor and functional expression of the cloned cDNA. Nature *346*, 561–564. 10.1038/346561a0.

16. Zanettini, C., Panlilio, L. V, Alicki, M., Goldberg, S.R., Haller, J., and Yasar, S. (2011). Effects of endocannabinoid system modulation on cognitive and emotional behavior. Front Behav Neurosci *5*, 57. 10.3389/fnbeh.2011.00057.

17. Aliczki, M., and Haller, J. (2015). Electric Shock as Model of Post-traumatic Stress Disorder in Rodents. In Comprehensive Guide to Post-Traumatic Stress Disorder (Springer International Publishing), pp. 1–16. 10.1007/978-3-319-08613-2_132-1.

18. Danandeh, A., Vozella, V., Lim, J., Oveisi, F., Ramirez, G.L., Mears, D., Wynn, G., and Piomelli, D. (2018). Effects of fatty acid amide hydrolase inhibitor URB597 in a rat model of trauma-induced long-term anxiety. Psychopharmacology (Berl) *235*, 3211–3221. 10.1007/s00213-018-5020-7.

19. Fotio, Y., Mabou Tagne, A., Jung, K.-M., and Piomelli, D. (2023). Fatty acid amide hydrolase inhibition alleviates anxiety-like symptoms in a rat model used to study post-traumatic stress disorder. Psychopharmacology (Berl). 10.1007/s00213-023-06358-y.

20. McNabb, M., Durante, K.A., Trocchio, S., Ritter, D.J., MacCaffrie, R., Brum, A., Mandile, S., and White, S. (2023). Self-reported Medicinal Cannabis Use as an Alternative to Prescription and Over-the-counter Medication Use Among US Military Veterans. Clin Ther *45*, 562–577. 10.1016/j.clinthera.2023.04.003.

21. Neumeister, A., Seidel, J., Ragen, B.J., and Pietrzak, R.H. (2015). Translational evidence for a role of endocannabinoids in the etiology and treatment of posttraumatic stress disorder. Psychoneuroendocrinology *51*, 577–584. 10.1016/j.psyneuen.2014.10.012.

22. Neumeister, A., Normandin, M.D., Pietrzak, R.H., Piomelli, D., Zheng, M.Q., Gujarro-Anton, A., Potenza, M.N., Bailey, C.R., Lin, S.F., Najafzadeh, S., et al. (2013). Elevated brain cannabinoid CB1 receptor availability in post-traumatic stress disorder: a positron emission tomography study. Mol Psychiatry *18*, 1034–1040. 10.1038/mp.2013.61.

23. Hill, M.N., Bierer, L.M., Makotkine, I., Golier, J.A., Galea, S., McEwen, B.S., Hillard, C.J., and Yehuda, R. (2013). Reductions in circulating endocannabinoid levels in individuals with post-traumatic stress disorder following exposure to the World Trade Center attacks. Psychoneuroendocrinology *38*, 2952–2961. 10.1016/j.psyneuen.2013.08.004.

24. Morena, M., Berardi, A., Colucci, P., Palmery, M., Trezza, V., Hill, M.N., and Campolongo, P. (2018). Enhancing Endocannabinoid Neurotransmission Augments

The Efficacy of Extinction Training and Ameliorates Traumatic Stress-Induced Behavioral Alterations in Rats. Neuropsychopharmacology *43*, 1284–1296. 10.1038/npp.2017.305.

25. Gunduz-Cinar, O., MacPherson, K.P., Cinar, R., Gamble-George, J., Sugden, K., Williams, B., Godlewski, G., Ramikie, T.S., Gorka, A.X., Alapafuja, S.O., et al. (2013). Convergent translational evidence of a role for anandamide in amygdala-mediated fear extinction, threat processing and stress-reactivity. Mol Psychiatry *18*, 813–823. 10.1038/mp.2012.72.

26. Hauer, D., Schelling, G., Gola, H., Campolongo, P., Morath, J., Roozendaal, B., Hamuni, G., Karabatsiakis, A., Atsak, P., Vogeser, M., et al. (2013). Plasma concentrations of endocannabinoids and related primary fatty acid amides in patients with post-traumatic stress disorder. PLoS One *8*, e62741. 10.1371/journal.pone.0062741.

27. Olango, W.M., Roche, M., Ford, G.K., Harhen, B., and Finn, D.P. (2012). The endocannabinoid system in the rat dorsolateral periaqueductal grey mediates fear-conditioned analgesia and controls fear expression in the presence of nociceptive tone. Br J Pharmacol *165*, 2549–2560. 10.1111/j.1476-5381.2011.01478.x.

28. Schaefer, C., Enning, F., Mueller, J.K., Bumb, J.M., Rohleder, C., Odorfer, T.M., Klosterkötter, J., Hellmich, M., Koethe, D., Schmahl, C., et al. (2014). Fatty acid ethanolamide levels are altered in borderline personality and complex posttraumatic stress disorders. Eur Arch Psychiatry Clin Neurosci *264*, 459–463. 10.1007/s00406-013-0470-8.

29. Bluett, R.J., Gamble-George, J.C., Hermanson, D.J., Hartley, N.D., Marnett, L.J., and Patel, S. (2014). Central anandamide deficiency predicts stress-induced anxiety: behavioral reversal through endocannabinoid augmentation. Transl Psychiatry *4*, e408. 10.1038/tp.2014.53.

30. deRoon-Cassini, T.A., Bergner, C.L., Chesney, S.A., Schumann, N.R., Lee, T.S., Brasel, K.J., and Hillard, C.J. (2022). Circulating endocannabinoids and genetic polymorphisms as predictors of posttraumatic stress disorder symptom severity: heterogeneity in a community-based cohort. Transl Psychiatry *12*, 48. 10.1038/s41398-022-01808-1.

31. Lim, J., Igarashi, M., Jung, K.-M., Butini, S., Campiani, G., and Piomelli, D. (2016). Endocannabinoid Modulation of Predator Stress-Induced Long-Term Anxiety in Rats. Neuropsychopharmacology *41*, 1329–1339. 10.1038/npp.2015.284.

32. Danan, D., Todder, D., Zohar, J., and Cohen, H. (2021). Is PTSD-Phenotype Associated with HPA-Axis Sensitivity?: The Endocannabinoid System in Modulating Stress Response in Rats. Int J Mol Sci *22*, 6416. 10.3390/ijms22126416.

33. Hoffman, G.E., Smith, M.S., and Verbalis, J.G. (1993). c-Fos and related immediate early gene products as markers of activity in neuroendocrine systems. Front Neuroendocrinol *14*, 173–213. 10.1006/frne.1993.1006.

34. Minatohara, K., Akiyoshi, M., and Okuno, H. (2015). Role of Immediate-Early Genes in Synaptic Plasticity and Neuronal Ensembles Underlying the Memory Trace. Front Mol Neurosci *8*, 78. 10.3389/fnmol.2015.00078.

35. Ploski, J.E., Monsey, M.S., Nguyen, T., DiLeone, R.J., and Schafe, G.E. (2011). The neuronal PAS domain protein 4 (Npas4) is required for new and reactivated fear memories. PLoS One *6*, e23760. 10.1371/journal.pone.0023760.

36. Kozlovsky, N., Matar, M.A., Kaplan, Z., Kotler, M., Zohar, J., and Cohen, H. (2008). The immediate early gene Arc is associated with behavioral resilience to stress exposure in an animal model of posttraumatic stress disorder. Eur Neuropsychopharmacol *18*, 107–116. 10.1016/j.euroneuro.2007.04.009.

37. Selçuk, B., Aksu, T., Dereli, O., and Adebali, O. (2023). Downregulated NPAS4 in multiple brain regions is associated with major depressive disorder. Sci Rep *13*, 21596. 10.1038/s41598-023-48646-9.

38. Hartzell, A.L., Martyniuk, K.M., Brigidi, G.S., Heinz, D.A., Djaja, N.A., Payne, A., and Bloodgood, B.L. (2018). NPAS4 recruits CCK basket cell synapses and enhances cannabinoid-sensitive inhibition in the mouse hippocampus. Elife *7*. 10.7554/eLife.35927.

39. Kaczkurkin, A.N., Burton, P.C., Chazin, S.M., Manbeck, A.B., Espensen-Sturges, T., Cooper, S.E., Sponheim, S.R., and Lissek, S. (2017). Neural Substrates of Overgeneralized Conditioned Fear in PTSD. Am J Psychiatry *174*, 125–134. 10.1176/appi.ajp.2016.15121549.

40. Colucci, P., Marchetta, E., Mancini, G.F., Alva, P., Chiarotti, F., Hasan, M.T., and Campolongo, P. (2020). Predicting susceptibility and resilience in an animal model of post-traumatic stress disorder (PTSD). Transl Psychiatry *10*, 243. 10.1038/s41398-020-00929-9.

41. Pascual Cuadrado, D., Todorov, H., Lerner, R., Islami, L., Bindila, L., Gerber, S., and Lutz, B. (2022). Long-term molecular differences between resilient and susceptible mice after a single traumatic exposure. Br J Pharmacol *179*, 4161–4180. 10.1111/bph.15697.

42. Fowler, C.J. (2013). Transport of endocannabinoids across the plasma membrane and within the cell. FEBS J *280*, 1895–1904. 10.1111/febs.12212.

43. Morena, M., Roozendaal, B., Trezza, V., Ratano, P., Peloso, A., Hauer, D., Atsak, P., Trabace, L., Cuomo, V., McGaugh, J.L., et al. (2014). Endogenous cannabinoid release within prefrontal-limbic pathways affects memory consolidation of emotional training. Proc Natl Acad Sci U S A *111*, 18333–18338. 10.1073/pnas.1420285111.

44. Sipe, J.C., Chiang, K., Gerber, A.L., Beutler, E., and Cravatt, B.F. (2002). A missense mutation in human fatty acid amide hydrolase associated with problem drug use. Proceedings of the National Academy of Sciences *99*, 8394–8399. 10.1073/pnas.082235799.

45. Crombie, K.M., Privratsky, A.A., Schomaker, C.M., Heilicher, M., Ross, M.C., Sartin-Tarm, A., Sellnow, K., Binder, E.B., Andrew James, G., and Cisler, J.M. (2022). The

influence of FAAH genetic variation on physiological, cognitive, and neural signatures of fear acquisition and extinction learning in women with PTSD. Neuroimage Clin *33*, 102922. 10.1016/j.nicl.2021.102922.

46. Spagnolo, P.A., Ramchandani, V.A., Schwandt, M.L., Kwako, L.E., George, D.T., Mayo, L.M., Hillard, C.J., and Heilig, M. (2016). FAAH Gene Variation Moderates Stress Response and Symptom Severity in Patients with Posttraumatic Stress Disorder and Comorbid Alcohol Dependence. Alcohol Clin Exp Res *40*, 2426–2434. 10.1111/acer.13210.

47. Mayo, L.M., Asratian, A., Lindé, J., Morena, M., Haataja, R., Hammar, V., Augier, G., Hill, M.N., and Heilig, M. (2020). Elevated Anandamide, Enhanced Recall of Fear Extinction, and Attenuated Stress Responses Following Inhibition of Fatty Acid Amide Hydrolase: A Randomized, Controlled Experimental Medicine Trial. Biol Psychiatry *87*, 538–547. 10.1016/j.biopsych.2019.07.034.

48. Rosas-Vidal, L.E., Naskar, S., Mayo, L.M., Perini, I., Altemus, M., Engelbrektsson, H., Jagasia, P., Heilig, M., and Patel, S. (2024). Prefrontal correlates of fear generalization during endocannabinoid depletion. Preprint.

49. Kondev, V., Bluett, R., Najeed, M., Rosas-Vidal, L.E., Grueter, B.A., and Patel, S. (2023). Ventral hippocampal diacylglycerol lipase-alpha deletion decreases avoidance behaviors and alters excitation-inhibition balance. Neurobiol Stress *22*, 100510. 10.1016/j.ynstr.2022.100510.

50. Gao, Y., Vasilyev, D. V., Goncalves, M.B., Howell, F. V., Hobbs, C., Reisenberg, M., Shen, R., Zhang, M.-Y., Strassle, B.W., Lu, P., et al. (2010). Loss of Retrograde Endocannabinoid Signaling and Reduced Adult Neurogenesis in Diacylglycerol Lipase Knock-out Mice. The Journal of Neuroscience *30*, 2017–2024. 10.1523/JNEUROSCI.5693-09.2010.

51. Tanimura, A., Yamazaki, M., Hashimotodani, Y., Uchigashima, M., Kawata, S., Abe, M., Kita, Y., Hashimoto, K., Shimizu, T., Watanabe, M., et al. (2010). The Endocannabinoid 2-Arachidonoylglycerol Produced by Diacylglycerol Lipase α Mediates Retrograde Suppression of Synaptic Transmission. Neuron *65*, 320–327. 10.1016/j.neuron.2010.01.021.

52. Yoshino, H., Miyamae, T., Hansen, G., Zambrowicz, B., Flynn, M., Pedicord, D., Blat, Y., Westphal, R.S., Zaczek, R., Lewis, D.A., et al. (2011). Postsynaptic diacylglycerol lipase α mediates retrograde endocannabinoid suppression of inhibition in mouse prefrontal cortex. J Physiol *589*, 4857–4884. 10.1113/jphysiol.2011.212225.

53. Jain, T., Wager-Miller, J., Mackie, K., and Straiker, A. (2013). Diacylglycerol Lipase *α* (DAGL *α*) and DAGL *β* Cooperatively Regulate the Production of 2-Arachidonoyl Glycerol in Autaptic Hippocampal Neurons. Mol Pharmacol *84*, 296–302. 10.1124/mol.113.085217.

54. Marco, E.M., Echeverry-Alzate, V., López-Moreno, J.A., Giné, E., Peñasco, S., and Viveros, M.P. (2014). Consequences of early life stress on the expression of endocannabinoid-related genes in the rat brain. Behavioural Pharmacology *25*, 547–556. 10.1097/FBP.0000000000000068.

55. Corbett, B.F., Luz, S., Arner, J., Pearson-Leary, J., Sengupta, A., Taylor, D., Gehrman, P., Ross, R., and Bhatnagar, S. (2019). Sphingosine-1-phosphate receptor 3 in the medial prefrontal cortex promotes stress resilience by reducing inflammatory processes. Nat Commun *10*, 3146. 10.1038/s41467-019-10904-8.

56. van Rooij, S.J.H., Stevens, J.S., Ely, T.D., Hinrichs, R., Michopoulos, V., Winters, S.J., Ogbonmwan, Y.E., Shin, J., Nugent, N.R., Hudak, L.A., et al. (2018). The Role of the Hippocampus in Predicting Future Posttraumatic Stress Disorder Symptoms in Recently Traumatized Civilians. Biol Psychiatry *84*, 106–115. 10.1016/j.biopsych.2017.09.005.

57. Weng, F.-J., Garcia, R.I., Lutzu, S., Alviña, K., Zhang, Y., Dushko, M., Ku, T., Zemoura, K., Rich, D., Garcia-Dominguez, D., et al. (2018). Npas4 Is a Critical Regulator of Learning-Induced Plasticity at Mossy Fiber-CA3 Synapses during Contextual Memory Formation. Neuron *97*, 1137-1152.e5. 10.1016/j.neuron.2018.01.026.

58. Haller, J., Aliczki, M., Pelczer, K.G., Spitzer, K., Balogh, Z., and Kantor, S. (2014). Effects of the fatty acid amide hydrolase inhibitor URB597 on coping behavior under challenging conditions in mice. Psychopharmacology (Berl) *231*, 593–601. 10.1007/s00213-013-3273-8.

59. Fanselow, M.S. (1980). Conditioned and unconditional components of post-shock freezing. Pavlov J Biol Sci *15*, 177–182. 10.1007/BF03001163.

60. Barsy, B., Mikics, É., Barsvári, B., and Haller, J. (2011). The long-term impact of footshock stress on addiction-related behaviors in rats. Neuropharmacology *60*, 267–273. 10.1016/j.neuropharm.2010.09.009.

61. Paxinos, G., and Watson, C. (2006). The Rat Brain in Stereotaxic Coordinates 6th edition. (Academic Press).

62. Mikics, É., Tóth, M., Varjú, P., Gereben, B., Liposits, Z., Ashaber, M., Halász, J., Barna, I., Farkas, I., and Haller, J. (2008). Lasting changes in social behavior and amygdala function following traumatic experience induced by a single series of foot-shocks. Psychoneuroendocrinology *33*, 1198–1210. 10.1016/j.psyneuen.2008.06.006.

63. Lee, S.-H., Ledri, M., Tóth, B., Marchionni, I., Henstridge, C.M., Dudok, B., Kenesei, K., Barna, L., Szabó, S.I., Renkecz, T., et al. (2015). Multiple Forms of Endocannabinoid and Endovanilloid Signaling Regulate the Tonic Control of GABA Release. J Neurosci *35*, 10039–10057. 10.1523/JNEUROSCI.4112-14.2015.

64. Livak, K.J., and Schmittgen, T.D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods *25*, 402–408. 10.1006/meth.2001.1262.

65. Breiman, L. (2001). Random Forests. Mach Learn *45*, 5–32. 10.1023/A:1010933404324.

Figure 1

Figure 2