Forskolin reverses the O-GlcNAcylation dependent decrease in GABA$_A$R current amplitude at hippocampal synapses possibly at a neurosteroid site on GABA$_A$Rs.

Shekinah Phillips[1,3], John Chatham[2], Lori L. McMahon*[1,3]

1. Department of Cell, Developmental and Integrative Biology, University of Alabama at Birmingham, Birmingham, Alabama 35294.
2. Division of Molecular and Cellular Pathology, Department of Pathology, University of Alabama at Birmingham, Birmingham, Alabama 35294.
3. Department of Neuroscience, Medical University of South Carolina, Charleston, SC 29403

*Corresponding author: Lori L. McMahon, PhD, Department of Neuroscience, Medical University of South Carolina, 173 Ashley Ave, Charleston, SC 29403

Acknowledgements: This research was supported by NIH NINDS R01AG066489 to LLM and JCC, 1F99NS134163-01 to SMP.

Author Contributions Statement: S.P. and L.L.M. designed the experiments; S.P. conducted experiments; S.P. and L.L.M. analyzed data; S.P, J.C.C., and L.L.M. wrote and revised the manuscript.

Data Availability Statement: The datasets generated and analyzed during the current study are available from the corresponding author on reasonable request.

Competing Interests Statement: The authors declare no competing interests.

Abstract

GABAergic transmission is influenced by post-translational modifications, like phosphorylation, impacting channel conductance, allosteric modulator sensitivity, and membrane trafficking. O-GlcNAcylation is a post-translational modification involving the O-linked attachment of β–N-acetylglucosamine on serine/threonine residues. Previously we reported an acute increase in O-GlcNAcylation elicits a long-term depression of evoked $GABA_AR$ inhibitory post synaptic currents (eIPSCs) onto hippocampal principal cells. Importantly O-GlcNAcylation and phosphorylation can co-occur or compete for the same residue; whether they interact in modulating GABAergic IPSCs is unknown. We tested this by recording IPSCs from hippocampal principal cells and pharmacologically increased O-GlcNAcylation, before or after increasing serine phosphorylation using the adenylate cyclase activator, forskolin. Although forskolin had no significant effect on baseline eIPSC amplitude, we found that a prior increase in O-GlcNAcylation unmasks a forskolin-dependent increase in eIPSC amplitude, reversing the O-GlcNAc-induced eIPSC depression. Inhibition of adenylate cyclase or protein kinase A did not prevent the potentiating effect of forskolin, indicating serine phosphorylation is not the mechanism. Surprisingly, increasing O-GlcNAcylation also unmasked a potentiating effect of the neurosteroids 5α-pregnane-3α,21-diol-20-one (THDOC) and progesterone on eIPSC amplitude, mimicking forskolin. Our findings show under conditions of heightened O-GlcNAcylation, the neurosteroid site on synaptic $GABA_ARs$ is accessible to agonists, permitting strengthening of synaptic inhibition.

Introduction

Gamma-aminobutyric acid type A receptors (GABA$_A$Rs) are heteropentameric ligand-gated chloride channels composed of α, β, γ, and sometimes δ subunits that mediate both fast synaptic and tonic inhibition, depending on their synaptic vs extrasynaptic location, respectively[1]. Mutations in specific subunits are linked to epilepsy syndromes [2], and gene polymorphisms in specific GABA$_A$R subunits associate with neuropsychiatric disorders, including alcohol use disorder [3], anxiety [4], schizophrenia [5], bipolar disorder [5], and even major depressive disorder [6], including postpartum depression [7].

For decades it has been appreciated that neurosteroids mediate their sedative hypnotic and anxiolytic effects via positive allosteric binding to specific GABA$_A$R subunits, particularly extrasynaptic GABA$_A$Rs containing α5, α4 or δ subunits [8,9]. GABA$_A$R function is also potently modulated by serine phosphorylation [10]. For example, protein kinase A (PKA) mediated phosphorylation of specific serines on synaptic GABA$_A$Rs induced by application of the adenylate cyclase activator, forskolin, bidirectionally modulates GABA-gated current amplitude and induces endocytosis depending on the neuron type and phosphorylated residue[8,11–14]. Importantly, the phosphorylation state has direct consequences on potency of allosteric modulation of GABA$_A$Rs by neurosteriods, barbiturates, and benzodiazepines in a subunit-specific manner [10].

The post-translational modification of proteins by β-N-acetylglucosamine via an O-linkage on serine and threonine residues (O-GlcNAcylation), can modulate protein phosphorylation by competing directly with phosphorylation for the same residues, or indirectly via modification of other sites thereby changing protein structure and protein-protein interactions. In addition, many kinases are modified by O-GlcNAc and this can regulate their function [15]. Both O-GlcNAcylation and phosphorylation are dynamic, reversible, and ubiquitous. While many kinases and phosphatases exist, O-GlcNAc is tightly regulated by a single enzyme pair, OGT (O-GlcNAc transferase) and OGA (O-GlcNAcase), which add and remove O-GlcNAc from serine/threonine residues, respectively. Of note, these enzymes are highly expressed in hippocampus [16,17]. The OGT substrate, UDP-GlcNAc, is generated by the hexosamine biosynthetic pathway (HBP) via glucose metabolism potentially linking this modification to nutrient availability [18,19].

Our lab recently reported that O-GlcNAcylation can be increased within minutes by exposing hippocampal brain slices to the HBP substrate glucosamine (GlcN), the OGA inhibitor thiamet-G (TMG) or in combination and this leads to depression of excitatory synaptic transmission [20,21]. More recently, we reported that pharmacologically increasing O-GlcNAc using GlcN and TMG in combination or GlcN alone induces a long-lasting depression of GABA$_A$R-mediated inhibitory postsynaptic currents (IPSCs) at hippocampal synapses within minutes, and decreases the amplitude and frequency of spontaneous IPSCs (sIPSCs) [22]. In *Oga +/-* mice, where O-GlcNAc is chronically elevated, inhibitory synaptic transmission was reduced in medial prefrontal cortex, and this was rescued by OGA overexpression via adeno-associated viral injection [23]. Additionally, the *Oga +/-* mice exhibited an antidepressant-like behavior, which was also reversed by viral OGA overexpression. Collectively, acute and chronic elevation in O-GlcNAcylation *in vitro* and *in vivo* depresses GABA$_A$R- mediated synaptic inhibition. These findings highlight O-GlcNAcylation as a critical regulator of the efficacy of synaptic neuronal inhibition, in both physiological and pathophysiological conditions of elevated O-GlcNAcylation.

Although serine phosphorylation [10,11] and O-GlcNAcylation [22] are fundamental modulators of the strength of GABA$_A$R-mediated synaptic inhibition, no study has examined whether these modifications interact in modifying the efficacy of synaptic inhibition. We investigated this possibility using electrophysiology in hippocampal slices and pharmacologically increased O-GlcNAc using glucosamine (GlcN) and thiamet-G (TMG) in combination. We also increased serine phosphorylation using the adenylate cyclase activator forskolin. Unexpectedly, we found that a prior increase in O-GlcNAcylation, which induces depression of GABA$_A$R-mediated eIPSC amplitude, unmasks a forskolin-dependent increase in eIPSC amplitude, even though forskolin had no effects when applied in the absence of increased O-GlcNAcylation. Surprisingly, inhibition of adenylate cyclase or PKA did not prevent the potentiating effect of forskolin on eIPSC amplitude, indicating that serine phosphorylation is not the mechanism. Similar to findings in a study in carp amacrine-like cells showing that forskolin binds to a GABA$_A$R neurosteroid site [24], we found that the neurosteroids THDOC and progesterone potentiate the IPSC amplitude following a prior increase in O-GlcNAcylation, mimicking the effect of forskolin. These findings suggest that O-GlcNAcylation promotes neurosteroid site accessibility on GABA$_A$Rs thereby reversing the depressive effect of O-GlcNAcylation and strengthening synaptic inhibition.

Results

An acute increase in O-GlcNAcylation reduces inhibitory post-synaptic currents on CA1 pyramidal cells.

Using hippocampal slices and Western blot analysis, we previously reported that an acute 10 min exposure to the HBP substrate GlcN (100 µM or 5 mM) or to the OGA inhibitor TMG (1 µM) induces a significant, global increase in protein O-GlcNAcylation (O-GlcNAc) [20,21]. This same 10 min exposure induces a long-term depression (LTD) at both excitatory CA3-CA1 synapses [20,21], and inhibitory synapses[22] during electrophysiological recordings. Additionally, increasing protein O-GlcNAc reduces the amplitude and frequency of spontaneous IPSC (sIPSCs) recorded from CA1 pyramidal cells and dentate granule cells (DGCs). Here, we confirm that a 10 min bath application of GlcN (5mM) and TMG (1µM) (GlcN+TMG) to pharmacologically increase O-GlcNAcylation during whole-cell voltage clamp recordings from CA1 pyramidal cells significantly depresses the sIPSC amplitude (Figure 1Aii, cumulative probability distribution, $p < 0.0001$, KS D value =0.17, Kolmogorov-Smirnov test; inset: $p < 0.0001$, Wilcoxon matched-pairs signed rank test) and increases the interevent interval, reflecting a decrease in sIPSC frequency (Figure 1Aiii, cumulative probability distribution, $p < 0.0001$, KS D value = 0.23, Kolmogorov-Smirnov test; inset: $p < 0.0001$, Wilcoxon matched-pairs signed rank test). Our previous work also showed that the amplitude of miniature IPSCs (mIPSCs) was significantly decreased by increasing O-GlcNAcylation[22], suggesting this synaptic depression is through a postsynaptic mechanism.

Figure 1. Acute increase in O-GlcNAcylation reduces spontaneous IPSCs in hippocampal CA1 pyramidal cells. Ai. (left) Schematic illustrating whole-cell recording of CA1 pyramidal cells (right) representative sIPSC trace showing baseline and GlcN + TMG application (top)and expanded time scale (bottom) during baseline (black) and GlcN + TMG application (orange). **Aii.** Cumulative probability

distribution of sIPSC amplitude (p < 0.0001, KS D value = 0.17, Kolmogorov-Smirnov test); *inset*: (left) average sIPSC trace before (black) and after (orange) GlcN + TMG;scaled trace (dashed) shows no change in rise or decay time of depressed sIPSC. *inset*: (right) bar chart showing average (±SEM) sIPSC amplitude. Baseline: 69.7 ± 0.8 pA, GlcN + TMG: 43.5 ± 0.5 pA (p < 0.0001, Wilcoxon matched-pairs signed rank test, n = 9 cells, 6 rats). **Aiii.** Cumulative probability distribution of sIPSC interevent interval (p < 0.0001, KS D value = 0.225, Kolmogorov-Smirnov test). Baseline: 344.5 ± 4.7 ms, GlcN + TMG: 767.3 ± 16.6 ms, *inset*: (p < 0.0001, Wilcoxon matched-pairs signed rank test, n = 9 cells, 6 rats).

The O-GlcNAc-induced depression of synaptic inhibition is not prevented by an actin stabilizer, but is partially dependent upon dynamin-mediated GABA$_A$R endocytosis.

Because serine phosphorylation can cause GABA$_A$R endocytosis in a subunit and serine specific manor [10,25,26] by analogy, we speculated that the O-GlcNAc-induced LTD of synaptic inhibition (or O-GlcNAc iLTD) we previously reported[22] is occurring via GABA$_A$R endocytosis. This possibility is supported by findings from our lab and others that O-GlcNAcylation of the GluA2 AMPAR subunit leads to long-term depression of excitatory transmission (O-GlcNAc LTD) at hippocampal CA3-CA1 synapses[20] and causes AMPAR endocytosis [27].

Therefore, to determine if increasing O-GlcNAc induces GABA$_A$R endocytosis during expression of O-GlcNAc iLTD, two experiments were performed. First, we tested whether interfering with actin prevents O-GlcNAc iLTD. We recorded CA1 pyramidal cells (Cs Gluconate pipette solution; E$_{Cl-}$ = −60 mV) and included the actin stabilizer, jasplakinolide (jasp) [28], in the pipette solution during whole-cell voltage clamp recordings. We interleaved experiments using pipette solution without jasplakinolide to ensure successful expression of O-GlcNAc iLTD. After a 15-min baseline recording, GlcN+TMG was applied for 10 min to induce O-GlcNAc iLTD. With or without jasplakinolide, successful iLTD was induced (Fig. 2Ai, O-GlcNAc iLTD – jasp: 69.2 ± 5.4% of baseline transmission, p=0.002, paired t-test; Fig. 2Aii, O-GlcNAc iLTD + jasp: 75.2 ± 3.4% of baseline transmission, p=0.0002, paired t-test). However, no significant difference was found between groups (Fig. 2Aiii, p=0.59, One-way ANOVA), indicating that the O-GlcNAc iLTD is not caused by actin mediated GABA$_A$R endocytosis.

Figure 2. Blocking actin mediated GABA_AR endocytosis does not prevent O-GlcNAciLTD in CA1 pyramidal cells. Ai. Normalized data showing average eIPSC amplitude over time exposed to GlcN+TMG (10 min, orange) after a 15-minute baseline without jasp, (p= 0.005, paired t-test, n = 6 cells, 6 rats). *Inset*: representative eIPSC traces before (black) and after GlcN + TMG (orange) application and scaled average without jasp shows no change in rise or decay time of the eIPSC. **Aii.** Normalized data showing average eIPSC amplitude over time exposed to GlcN+TMG (10 min, orange) after a 15-minute baseline with jasplakinolide included in the pipette solution (jasp) (p= 0.0005, paired t-test, n = 8 cells, 8 rats*). Inset*: representative eIPSC traces before (black) and after GlcN + TMG (orange) application and scaled average with jasp shows no change in rise time or decay of the eIPSC.. **Aiii.** Overlay of the experimental groups showing no significant difference in the average (±SEM) eIPSC amplitude over time (p<0.0001, One-way ANOVA). Gray horizontal bars represent the mean ± SEM.

Next, we tested whether expression of O-GlcNAc iLTD requires dynamin-dependent GABA_AR endocytosis. To accomplish this, we incubated slices in dynasore (80 µM, 30 min) or DMSO (vehicle) and performed whole-cell recordings from CA1 pyramidal cells in an interleaved fashion. After a 5 min baseline, GlcN+TMG was applied for 10 min to induce iLTD. With or without dynasore, successful iLTD was induced (Fig. 3Ai, O-GlcNAc iLTD – dynasore: 79.3 ± 4.5% of baseline transmission, p= 0.005,

paired t-test; Fig. 3Aii, O-GlcNAc iLTD + dynasore: 81.1 ± 13.2% of baseline transmission, p=0.004, paired t-test). However, there were no significant differences between groups (Fig.3Aiii, p=0.08, One-Way ANOVA). Importantly, it was noted that the dataset with dynasore had greater variability during iLTD expression as indicated by the larger error bars between 20-25 mins (Fig. 3Aii). Upon further inspection of individual experiments, we recognized that in some cells (n=4/8), the eIPSC amplitude was potentiated following GlcN+TMG, which would be expected if endocytosis is prevented. Therefore, when the dynasore dataset was separated into those with potentiation and those without, a clear population of cells was revealed that exhibited significant potentiation of the eIPSC amplitude (Fig. 3Aiv, 136.3 ± 10.2% of normalized eIPSC amplitudes following GlcN+TMG, application n= 4 cells, p=0.039, paired t-test), while the remaining population exhibited no change in eIPSC amplitude from the previously depressed level following GlcN+TMG application (Fig. 3Aiv, 73.9 ± 13.5% of normalized eIPSC amplitudes following GlcN+TMG, application n= 4 cells, p=0.13, paired t-test) and from the O-GlcNAC iLTD without dynasore (Fig3Ai) (p=0.131, unpaired t-test). Additionally, there was a significant difference between the potentiated versus the non-potentiated population of normalized eIPSC amplitudes following GlcN+TMG (p=0.03, paired t-test). This result is consistent with the interpretation that in some cells increasing O-GlcNAc induced a dynamin-dependent endocytosis of GABA$_A$Rs that could underlie the synaptic depression and in others a different mechanism exists. Furthermore, these findings reaffirm the heterogeneity in GABA$_A$Rs that exist at synapses in hippocampus. Additional experiments are needed to fully understand how O-GlcNAc impacts GABA$_A$R trafficking.

Figure 3. O-GlcNAc iLTD is partially dependent upon dynamin-mediated GABA$_A$R endocytosis. **Ai.** Normalized data showing average eIPSC amplitude over time exposed to GlcN+TMG (10 min, orange) after a 15-minute baseline without dynasore (DMSO-vehicle) (p= 0.005, paired t-test, n = 8 cells, 7 rats). *Inset:* representative eIPSC traces before (black) and after GlcN + TMG (orange) application and scaled average shows no difference in rise or decay time of the eIPSC. **Aii.** Normalized data showing average eIPSC amplitude over time exposed to GlcN+TMG (10 min, orange) after a 5-minute baseline in slices incubated in dynasore for 30 min (p= 0.004, paired-test, n = 8 cells, 7 rats*). Inset*: representative eIPSC traces before (black) and after GlcN + TMG (orange) application and scaled average with dynasore. **Aiii.** Overlay of experimental groups showing no significant difference in the average (±SEM) eIPSC amplitude over time (p=0.08, One-way ANOVA). **Aiii.** Separation of two

populations in the presence of dynasore where the eIPSC amplitude was significantly increased in one population (white, p=0.0392, paired t-test, n= 4 cells, 4 rats) and with no effect in the other population (red, p=0.13, paired t-test, n= 4 cells, 4 rats). Gray horizontal bars represent the mean ± SEM.

Possible interactions between phosphorylation and O-GlcNAcylation

Next, we wanted to determine if serine phosphorylation and O-GlcNAcylation interact to affect GABA$_A$R-mediated synaptic inhibition and whether an order effect exists. For decades, forskolin has been used to activate adenylate cyclase to drive protein kinase A (PKA) dependent phosphorylation of AMPARs at excitatory synapses in hippocampus, leading to synaptic potentiation [29–31]. PKA-dependent serine phosphorylation also modulates synaptic inhibition, but the effect is variable depending on the GABA$_A$R subunit confirmation [10]. Therefore, to test whether serine phosphorylation has an impact on subsequent induction of O-GlcNAc iLTD, we recorded from CA1 pyramidal cells (Cs Gluconate pipette solution; E$_{Cl-}$ = −60 mV), and bath applied forskolin (50µM) for 10 min to drive activation of adenylate cyclase and PKA followed by 10 min application of GlcN+TMG to induce O-GlcNAc iLTD. The eIPSC amplitudes during forskolin and GlcN+TMG were normalized to baseline and statistically compared by averaging 30 events during (a) baseline, (b) forskolin and (c) GlcN+TMG using repeated measures (RM) RM-ANOVA and Šídák's multiple comparisons post hoc test (Fig. 4Ai-iii, p=0.0001, RM ANOVA). We found no significant effect of forskolin compared to baseline transmission (Fig. 4Aii, 89.2 ± 6.9% of baseline transmission, p=0.39), and subsequent application of GlcN+TMG induced significant iLTD (Fig. 4Aii, 64.2 ± 4.4% of baseline transmission, p<0.0001). Despite no significant effect of forskolin on the eIPSC amplitude in the averaged dataset compared to baseline, we want to ensure there was no effect on the magnitude of subsequently induced O-GlcNAc iLTD. Therefore, we normalized the eIPSC amplitudes at the end of the 10 min forskolin application, thereby establishing new baseline from which to measure the O-GlcNAc iLTD magnitude. We found that from this new baseline, subsequent application of GlcN+TMG induced a significant iLTD (74.9 ± 3.8% of new baseline transmission (b-c comparison), p<0.0001, paired t-test) that is not different from the iLTD magnitude under control conditions obtained in Fig. 2Ai in the absence of jasp (Fig. 2Ai, 67.6 ± 5.6% of baseline transmission versus 73.5 ± 4.2% of new baseline transmission p= 0.53, unpaired t-test). In reviewing the data, it is important to note that there was high cell-to-cell variability in eIPSC amplitude during forskolin application, with some cells displaying potentiation and some depression of the eIPSC amplitude, as can be seen by inspection of the individual data points in the bar chart in Fig. 4Aii.

To determine if this finding in CA1 is generalizable, we performed the same experiment in dentate granule cells (DGCs). Similar to CA1, we recorded from DGCs (Cs Gluconate pipette solution; E_{Cl-} = −60 mV), and bath applied forskolin (50µM) for 10 min followed by 10 min of GlcN+TMG to induce O-GlcNAc iLTD. Again, the eIPSC amplitudes during forskolin and GlcN+TMG were normalized to baseline and statistically compared by averaging 30 events during (a) baseline, (b) forskolin and (c) GlcN+TMG (Fig. 4Bi, p=0.0041, RM-ANOVA). Using Šídák's multiple comparisons post hoc test, we found no significant change in eIPSC amplitude during forskolin application compared to baseline (Fig. 4Bi, 98.8 ± 4.6% of baseline transmission, p = 0.98), with subsequent GlcN+TMG application inducing significant iLTD (Fig. 4Bii, 75.2 ± 6.0% of baseline, p=0.013). In addition, there was a significant difference between baseline vs. GlcN+ TMG (Fig. 4Biii, p=0.029) and between forskolin vs GlcN+TMG (Fig. 4Biii, p=0.040). Also, as before, to measure the magnitude of the O-GlcNAc iLTD after forskolin, we re-normalized the eIPSC amplitudes at the end of the 10 min forskolin application to establish a new baseline and find significant O-GlcNAc iLTD (81.7 ± 9.0% of new baseline (b-c comparison), p=0.005, paired t-test).

To determine if there is any effect of forskolin or GlcN+TMG on presynaptic release probability, we analyzed the paired-pulse ratio (PPR), an indirect measure of presynaptic release probability, during baseline, GlcN+TMG and forskolin. No significant differences were detected in CA1 (Fig. 4Aiv, p=0.69, RM-ANOVA) or in the dentate gyrus (Fig. 4Biv, p=0.46, RM-ANOVA), indicating that a presynaptic mechanism is not involved.

Figure 4. Magnitude of the O-GlcNAc iLTD is unaffected by prior application of forskolin. Ai. CA1: Normalized data showing average eIPSC amplitude over time during baseline (5 min), bath application of forskolin (10 min, blue) and GlcN + TMG (10 min, orange). *Inset:* overlaid averaged representative eIPSC traces during baseline (black), forskolin (blue), and GlcN + TMG (orange) (left) and scaled (right) **Aii.** Bar chart illustrates the average affect of forskolin and GlcN+TMG on eIPSC amplitude from the original baseline (dotted line): 89.2% ± 6.9%, p=0.39 (forskolin), 64.2% ± 4.4%, p<0.0001 (GlcN+TMG) and each other (p=0.0023); (p=0.0001, RM-ANOVA). **Aiii.** For each individual experiment in Ai, line graphs show the average eIPSC amplitude during the last the 5 min in each condition corresponding to a,b,c in Ai (p=0.0002, RM-ANOVA, n = 10 cells, 9 rats), Šídák's: a-c (p=0.002) and b-c (p=0.01). Baseline: 404.9 ± 47.9 pA, forskolin: 366.5 ± 50.2 pA, GlcN+ TMG: 256.6 ± 25.0 pA (mean ± SEM). **Aiv.** Paired-Pulse Ratio (100 ms inter-pulse interval) between baseline, forskolin and GlcN+TMG showed no significant difference (p=0.69, RM-ANOVA). **Bi.** Dentate: Normalized data showing average eIPSC amplitude over time during baseline, bath application of forskolin (blue), and GlcN+TMG (orange). *Inset*: overlaid averaged representative eIPSC traces during baseline (black) forskolin (blue), GlcN + TMG (orange) (left) and scaled (right). **Bii.** Bar chart illustrates the average affect of forskolin and GlcN+TMG from the original baseline (dotted line): 98.8% ± 4.6%, p=0.98 (forskolin) and 75.2% ± 6.0%, p=0.0128 (GlcN+TMG) and each other (p=0.0453); (p=0.004, RM-ANOVA). **Biii.** For each individual experiment in Bi, line graphs show the average amplitudes for last 5 min in each condition corresponding to a,b,c in Bi (p = 0.005, RM-ANOVA, n = 8 cells, 8 rats), Šídák's: a-c (p=0.029) and b-c (p=0.040). Baseline: 523.7 ± 51.7 pA, forskolin: 514.9 ± 50.5 pA, GlcN+ TMG: 360.2 ± 28.3 pA (mean ± SEM). **Biv.** Paired- Pulse Ratio between baseline, forskolin and GlcN+TMG showed no significant difference (p=0.46, RM-ANOVA). Gray horizontal bars represent the mean ± SEM. Šídák's post hoc depicted with asterisks.

Next, we performed the experiment in reverse order, increasing O-GlcNAc with GlcN+TMG prior to driving phosphorylation with forskolin. We recorded from both from CA1 pyramidal cells and DGCs and applied GlcN+TMG for 10 min followed by forskolin for 10 min. Similar to above, eIPSC amplitudes

during forskolin and GlcN+TMG were normalized to baseline and compared (Fig. 5Ai, p<0.0001, RM-ANOVA; 5Bi; p=0.009 RM-ANOVA). A 10 min exposure to GlcN+TMG induced O-GlcNAc iLTD in CA1 pyramidal cells (Fig. 5Aii: 65.4 ± 5.2% of baseline transmission, p=0.0002; Šídák's post hoc test) and in dentate granule cells (Fig. 5Bi, Bii: 82.3 ± 2.6% of baseline transmission, p=0.002, Šídák's post hoc test). Surprisingly, subsequent application of forskolin reversed the O-GlcNAc iLTD and elicited a potentiation of the eIPSC amplitude in recordings from both CA1 pyramidal cells and DGCs (Figs. 5Ai-iii and 5Bi-iii). To analyze the magnitude of the forskolin-induced eIPSC potentiation, we re-normalized eIPSC amplitudes at the end of the 10 min GlcN+TMG application, establishing a new baseline, and then normalized forskolin values to the new baseline. We found a significant potentiation in CA1 (138.5 ± 7.8% of new baseline (b-c comparison), p= 0.006, paired t-test) and in dentate (143.4 ± 8.8% of new baseline (b-c comparison), p= 0.003, paired t-test) that reverses O-GlcNAc iLTD and in dentate, the potentiation overshoots the original baseline. Furthermore, these results suggest that a prior increase in O-GlcNAc unmasks a possible PKA dependent potentiation of synaptic inhibition that is absent under control conditions.

Figure 5. Forskolin reverses O-GlcNAc iLTD in CA1 and dentate. CA1: **Ai.** Normalized data showing average eIPSC amplitude over time during baseline (5 min), bath application of GlcN+TMG (10 min, orange), and forskolin (10 min, blue). *Inset:* overlaid averaged representative eIPSC traces during baseline (black), GlcN + TMG (orange) and forskolin (blue) (left) and scaled (right). **Aii.** Bar chart illustrates the average affect of GlcN+TMG and forskolin on eIPSC amplitude from the original baseline (dotted line): 65.4 ± 5.2%, p=0.0002 (GlcN+TMG) and 86.0% ± 6.3%, p=0.1561 (forskolin) of baseline and each other (p=0.0011); (p<0.0001, RM-ANOVA). **Aiii.** For each individual experiment in Ai, line

graphs show the average eIPSC amplitudes during the last 5 min of each condition corresponding to a,b,c in Ai (p < 0.0001, RM-ANOVA, n = 11 cells, 10 rats), Šídák's: a-b (p=0.0007) and b-c (p=0.028), Baseline: 471.2 ± 64.6 pA, GlcN+ TMG: 304.6 ± 52.0 pA, Forskolin: 397.6± 77.2 pA (mean ± SEM). **Aiv.** Paired-Pulse Ratio between baseline, GlcN+TMG and forskolin showed no significant difference (p=0.06, RM-ANOVA). Dentate: **Bi.** Normalized data showing average eIPSC amplitude over time during baselinebath application of GlcN +TMG (10 min, orange) and forskolin (10 min, blue). *Inset*: overlaid averaged representative eIPSC traces during baseline(black), GlcN + TMG (orange), and forskolin (blue) (left) and scaled (right). **Bii.** Bar chart illustrates the average affect of GlcN+TMG and forskolin on eIPSC amplitude from the original baseline (dotted line): 82.3% ± 2.6%, p=0.002 (GlcN+TMG) and 118.8% ± 9.3%, p=0.2500 (forskolin) and each other (p=0.008); (p=0.009, RM-ANOVA). **Biii.** For each individual experiment in Bi, line graphs show the averaged eIPSC amplitudes for last 5 min of each condition corresponding to a,b,c in **Bi**. (p=0.015, RM-ANOVA, n = 7 cells, 6 rats), Šídák's: a-b (p=0.0340) and b-c (p=0.038), Baseline: 376.3 ± 58.7 pA, GlcN+ TMG: 306.5 ± 44.2 pA, Forskolin: 443.0± 77.1 pA (mean ± SEM). **Biv.** Paired-Pulse Ratio between baseline, GlcN+TMG and forskolin showed no significant difference (p=0.24, RM-ANOVA). Gray horizontal bars represent the mean ± SEM. Šídák's post hoc test depicted with asterisks.

The forskolin dependent increase in eIPSC amplitude is not PKA dependent.

To confirm that the forskolin induced potentiation involves PKA dependent phosphorylation (Fig. 6A), we focused our experiments only on CA1 pyramidal cells. We first asked if the PKA inhibitor KT5720 (3 µM) applied for 10 mins before and during forskolin application was able to prevent the forskolin-induced potentiation that reverses the O-GlcNAc iLTD. Experiments with and without KT5720 were interleaved. eIPSC amplitudes during GlcN+TMG and forskolin were normalized to baseline (Fig. 6Bi-Biii) and statistically compared. GlcN+TMG induced O-GlcNAc iLTD and subsequent application of forskolin reversed the O-GlcNAc iLTD and elicited a potentiation of the eIPSC amplitude with (Fig. 6Bii, p=0.004, RM-ANOVA) and without (Fig. 6Biii, p=0.008, RM-ANOVA) KT5720. In the dataset containing KT5720, post hoc Šídák's multiple comparisons test showed a significant difference between baseline vs. GlcN+ TMG (Fig. 6Bii, p=0.002) and between GlcN+TMG vs. forskolin (Fig. 6Bii, p=0.011). In the dataset without KT5720, post hoc Šídák's multiple comparisons test showed a significant difference between baseline vs. GlcN+ TMG (Fig. 6Biii, p=0.0011) and between GlcN+TMG vs. forskolin (Fig. 6Biii, p=0.013). To measure the magnitude of the forskolin-induced potentiation, we re-normalized the eIPSC amplitudes at the end of the 10 min GlcN+TMG application to establish a new baseline, and found significant potentiation with (139.7 ± 9.4%, p= 0.004, paired t-test) and without (169.9 ±18.3, p= 0.005, paired t-test) KT5720, and no significant difference between groups (Fig. 6Bi, p= 0.18, unpaired t-test).

Since we were unable to block the forskolin mediated potentiation via PKA inhibition, we next targeted adenylate cyclase using the inhibitor, SQ22536 (100 μM). SQ2253 was bath applied for 10 mins before and during forskolin application, and experiments with and without SQ2253 were interleaved. eIPSC amplitudes during GlcN+TMG and forskolin amplitude were normalized to baseline (Fig. 6Ci) and statistically compared (Fig. 6Ci-Ciii). GlcN+TMG induced O-GlcNAc iLTD and subsequent application of forskolin reversed the O-GlcNAc iLTD and elicited a potentiation of the eIPSC amplitude with (Fig. 6Cii, p=0.031, RM ANOVA) and without SQ22536 (Fig. 6Ciii, p=0.03, RM ANOVA). In the dataset with SQ22536, post hoc Šídák's multiple comparisons test showed a significant difference between baseline vs. GlcN+ TMG (Fig. 6Cii, p=0.031). In the dataset without SQ22536, Šídák's multiple comparisons test showed a significant difference between baseline vs. GlcN+TMG (Fig. 6Ciii, p=0.020). To measure the magnitude of the forskolin-induced potentiation, we re-normalized the eIPSC amplitudes at the end of the 10 min GlcN+TMG application to establish a new baseline, and found significant potentiation with (122.7± 7.1%, p=0.017, paired t-test) and without (134.8 ± 13.3%, p=0.047, paired t-test) SQ22536, but similar to CA1, there was no significant difference between groups (Fig. 6Ci, p=0.42, unpaired t-test). Being that neither the adenylate cyclase nor PKA inhibitor prevented the forskolin dependent increase in eIPSC following a prior increase in O-GlcNAc, we concluded that this potentiation occurs through another mechanism.

The inactive adenylate cyclase activator, 1,9 dideoxyforskolin, partially mimics forskolin

While forskolin is known to activate adenylate cyclase [32–34], it has many other 'off-target' effects [32,35] that could be mediating the eIPSC potentiation following O-GlcNAc iLTD. Therefore, we asked whether the adenylate cyclase inactive forskolin analog, 1,9-dideoxyforskolin (50 μM) would mimic the effect of forskolin. Experiments with 1,9-dideoxyforskolin were interleaved with forskolin. eIPSC amplitudes during GlcN+TMG and subsequent application of 1,9 dideoxyforskolin or forskolin were normalized to baseline and statistically compared. GlcN+TMG induced O-GlcNAc iLTD and subsequent application of 1,9 dideoxyforskolin (Fig. 6Dii, p= 0.028, RM-ANOVA) and forskolin (Fig. 6Diii, forskolin p=0.037, RM-ANOVA) reversed the O-GlcNAc iLTD and elicited a potentiation of the eIPSC amplitude. In the 1,9 dideoxyforskolin dataset, Šídák's multiple comparisons test showed a significant difference between baseline vs. GlcN+ TMG (Fig. 6Dii, p= 0.011), indicating O-GlcNAc iLTD, followed by a significant potentiation upon 1,9 dideoxyforksolin exposure (Fig. 6Dii, p=0.037). In the forskolin dataset, Šídák's multiple comparisons test showed after O-GlcNAc iLTD (Fig. 6Diii, P=0.041), forskolin exposure induced a significant potentiation (Fig. 6Diii, p=0.035). When comparing the magnitude of the

potentiation induced by 1,9 dideoxyforskolin vs forskolin, eIPSC amplitudes were re-normalized at the end of the GlcN+TMG, application to re-establish a new baseline, and found significant potentiation with 1,9 dideoxyforskolin (121.3± 6.3%, p=0.0115, paired t-test) and forskolin (146.5 ± 11.8%, p=0.0075, paired t-test) and no significant difference between groups (Fig. 6Di, p= 0.029, unpaired t-test).

Figure 6. Forskolin induced potentiation of the eIPSC amplitude following an increase in O-GlcNAc is resistant to inhibitors of adenylate cyclase and PKA, suggesting serine phosphorylation is not the mechanism. A. Schematic illustrating possible interaction between serine O-GlcNAcylation and PKA-dependent serine phosphorylation of the GABA$_A$ receptor with pharmacological activators and inhibitors. **Bi**. Normalized data showing average (±SEM) eIPSC

amplitude over time from CA1 pyramidal cells with and without the PKA inhibitor KT5720 (3 µM) during baseline (5 min), GlcN+TMG (10 min, orange) and forskolin application (10 min, blue). KT5720 was applied for 10 mins before and during forskolin. **Bii**,**Biii**. For each individual experiment, line graphs show average eIPSC amplitudes for the last 5 min of each condition corresponding to a,b,c with KT5720 (white) (p=0.004, RM-ANOVA, n= 8 cells, 6 rats), Šídák's: a-b (p=0.002) and b-c (p=0.011) and without KT5720 (black) (p=0.007, RM-ANOVA, n= 9 cells, 6 rats), Šídák's: a-b (p=0.0011) and b-c (p=0.0126). **Ci**. Normalized data showing average (±SEM) eIPSC amplitude over time with and without the adenylate cyclase inhibitor, SQ22536 (100 µM) during baseline, GlcN+TMG (10 min, orange) and forskolin application (10 min, blue) **Cii, Ciii** For each individual experiment, line graphs show average eIPSC amplitudes for the last 5 min of each condition corresponding to a,b,c with SQ22536 (white) (p=0.031, RM-ANOVA, n= 7 cells, 6 rats), Šídák's: a-b (p=0.031) and without SQ22536 (black) (p=0.03, RM-ANOVA, n= 6 cells, 5 rats), Šídák's: a-b (p=0.020). **Di.** Normalized data showing average (±SEM) eIPSC amplitude over time during baseline, GlcN+TMG (10 min, orange) and application of forskolin or adenylate cyclase inactive analog 1,9 Dideoxyforskolin (10 min, blue). **Dii.** The average amplitudes for last 5 minutes of each condition corresponding to a,b,c with 1,9 Dideoxyforskolin (p= 0.028 RM-ANOVA, n=7 cells, 5 rats), Šídák's: a-b (p=0.011) and b-c (p=0.037). **Diii.** The average amplitudes for last 5 minutes of each condition with forskolin (p=0.037, RM-ANOVA, n= 6 cells, 4 rats), Šídák's: a-b (p=0.041) and b-c (p=0.035). Gray horizontal bars represent the mean ± SEM. Šídák's post-hoc depicted with asterisks.

5α-pregnane-3α,21-diol-20-one (THDOC) and progesterone reverse the O-GlcNAc-mediated depression of evoked IPSCs, mimicking forskolin.

The inability to prevent the forskolin-induced eIPSC potentiation following O-GlcNAc iLTD with adenylate cyclase and PKA inhibitors, and the partial mimic of forskolin's effect by the inactive adenylate cyclase analog 1,9 dideoxyforksolin, was very puzzling. In searching for a possible explanation, we were intrigued by a report where both forskolin and 1,9-dideoxyforskolin accelerated desensitization of GABA$_A$R currents in recordings from amacrine-like cells in carp (Carassius auratus) retina that was resistant to PKA inhibition [24]. Surprisingly, the neurosteroid, 5α-pregnane-3α,21-diol-20-one (THDOC), which is a structural analog to forskolin, also accelerated GABA$_A$R desensitization, mimicking the effect of forskolin [24]. Further experiments led to the conclusion that forskolin is acting at an allosteric neurosteroid site on GABA$_A$Rs. Because GABA$_A$Rs in mammalian hippocampus are potently modulated by neurosteroids containing specific subunit combinations [9], we sought to determine if the hippocampal neurosteroid, THDOC, also mimics the forskolin-induced eIPSC potentation following O-GlcNAc iLTD.

To test this, we recorded eIPSC from CA1 pyramidal cells and bath applied THDOC (10 µM) before or after GlcN+TMG. The eIPSC amplitudes during THDOC and GlcN+TMG were normalized to baseline and statistically compared (Fig. 7Ai-iii, p=0.003 RM-ANOVA). THDOC application (10 min) led to a

slight but not significant depression in eIPSC amplitude (Fig. 7Aii: 83.4 ± 6.9% of baseline transmission, p=0.153) similar to the lack of significant effect of forskolin on baseline transmission (see Fig. 4Ai, Bi). Subsequent application of GlcN+TMG induced O-GlcNAc iLTD (Fig. 7Aii: 56.9 ± 10.3% of baseline transmission, p=0.016).

Next, we performed the experiment in the reverse order to specifically ask whether THDOC mimics forskolin by eliciting a potentiation of the eIPSC amplitude when applied after GlcN+TMG (Fig.7Bi). The eIPSC amplitudes recorded from CA1 pyramidal cells during GlcN+TMG and THDOC were normalized to baseline and statistically compared (Fig. 7Bi-iii, p<0.0001, RM-ANOVA). Bath application of GlcN+TMG induced O-GlcNAc iLTD (Fig. 7Bii: 72.1 ± 3.3% of baseline transmission, p<0.0001), but subsequent application of THDOC yields no additional change in eIPSC amplitude (Fig. 7Bii: 76.7 ± 5.5% of baseline transmission, p=0.003), and is not different from eIPSC amplitude following GlcN+TMG (p=0.2738, paired t-test). There was a significant difference between baseline vs. GlcN+TMG (Fig. 7Biii, p<0.0001) and baseline vs. THDOC (Fig. 7Biii, p=0.005).

However, similar to the cell-to-cell variability we observed with forskolin (e.g., Fig. 4Aii), we noted variability in the response to THDOC (Fig. 7Bii), with eIPSCs recorded from some cells having a clear potentiation (Fig. 7Ci-iii, n=7/12), while others had no change (Fig.7 Di-iii, n=5/12 cells). To measure the effect of THDOC in these two populations, we re-normalized eIPSC amplitudes at the end of the 10 min GlcN+TMG application to establish a new baseline, then THDOC values were normalized to the new baseline. Similar to forskolin (Fig.4Ai-iii), we found a significant potentiation of the eIPSC amplitude in this subset of cells (116.4 ± 5.3%, p= 0.021, paired t-test, n=7/12). In the remaining cells, (Fig. 7Di-iii), we found no further change in eIPSC amplitude (90.0 ± 5.5%, p= 0.15, paired t-test) beyond what occurred following GlcN+TMG application.

Figure 7. 5α-pregnane-3α,21-diol-20-one (THDOC) reverses the O-GlcNAc iLTD, mimicking forskolin in CA1 pyramidal cells. Ai. Normalized data showing average eIPSC amplitude during THDOC (10 min, green) followed by GlcN+TMG (10 min, orange). **Aii.** Average THDOC and GlcN+TMG eIPSC amplitude from the original baseline (gray bar, dotted line): 83.4 ± 6.9%, p=0.153 (THDOC) and 56.9%± 10.3%, p=0.016 (GlcN+TMG) and each other (p=0.093); (p=0.0032, RM-ANOVA). **Aiii.** The average eIPSC amplitudes during the last 5 min of each condition corresponding to a,b,c in Ai (p= 0.0059, RM-ANOVA, n = 7 cells, 7 rats), Šídák's: a-c (p=0.032). **Bi.** Normalized data showing average eIPSC amplitude during GlcN+TMG (10 min, orange) followed by THDOC (10 min, green). **Bii.** Average GlcN+TMG and THDOC eIPSC amplitude from the original baseline (gray bar, dotted line): 72.1% ± 3.3% (GlcN+TMG), p< 0.0001 and 76.7% ± 5.5% (THDOC), p=0.003 and each other (p=0.617); (p<0.0001, RM-ANOVA). **Biii.** The average eIPSC amplitudes during the last 5 min of each condition corresponding to a,b,c in Bi (p <0.0001, RM-ANOVA, n = 12 cells, 12 rats), Šídák's: a-b (p<0.0001), a- (p=0.005). *Potentiated:* **Ci.** Normalized data showing average eIPSC amplitude during GlcN+TMG followed by THDOC. **Cii.** Average GlcN+TMG and THDOC eIPSC amplitude from the original baseline (gray bar, dotted line): 75.8% ± 2.6% (GlcN+TMG), p= 0.0002 and 88.6% ± 4.0%

(THDOC), p=0.089 and each other (p=0.034); (p=0.0002, RM-ANOVA). **Ciii.** Last 5 min average amplitudes corresponding to a,b,c (p=0.0003, RM-ANOVA, n = 7 cells, 7 rats), Šídák's: a-b (p=0.0002), b-c (p=0.036).   *Non-potentiated:* **Di.** Normalized data showing average eIPSC amplitude during GlcN+TMG followed by THDOC. **Dii.** Average GlcN+TMG and THDOC eIPSC amplitude from the original baseline (gray bar, dotted line): 66.9% ± 6.9% (GlcN+TMG), p= 0.026 and 59.9% ± 6.8% (THDOC), p=0.012 and each other (p=0.422); (p=0.0014, RM-ANOVA). **Diii.** Last 5 min average amplitudes corresponding to a,b,c (p=0.0099, RM-ANOVA, n = 5 cells, 5 rats), Šídák's: a-b (p=0.041), a-c (p=0.025). *Inset:* overlaid averaged representative eIPSC traces during baseline (black), GlcN + TMG (orange), THDOC (green) (left) and scaled (right). Gray horizontal bars represent mean ± SEM. Šídák's post-hoc depicted with asterisks.
.

To firm up the above results, we felt it important to test a second steroid, progesterone, known to modulate GABA_ARs, to see if the potentiation following GlcN+TMG is again mimicked. The eIPSC amplitudes recorded from CA1 pyramidal cells during progesterone and GlcN+TMG were normalized to baseline and statistically compared (Fig. 8Ai-iii, p=0.047, RM-ANOVA). Similar to forskolin and THDOC, progesterone exposure induced no significant change from baseline (Fig. 8Aii: 92.2 ± 7.3% of baseline, p=0.701) and subsequent GlcN+TMG induced O-GlcNAc iLTD (Fig. 8Aii: 76.9% ± 9.1% of baseline, p=0.126). A significant difference also existed between progesterone vs. GlcN+TMG (Fig. 8Aiii, p=0.008). When performing the experiment in reverse, the eIPSC amplitudes recorded from CA1 pyramidal cells during GlcN+TMG and progesterone were normalized to baseline and statistically compared (Fig 8Bi, p<0.0001, RM-ANOVA). GlcN+TMG induced O-GlcNAc iLTD (Fig. 8Bii: 68.7% ± 3.8% of baseline transmission, p<0.001,) but subsequent application of progesterone yields no additional change in eIPSC amplitude (Fig. 8Bii: 75.3% ± 4.9% of baseline transmission, p=0.0011), and is not different from eIPSC amplitude following GlcN+TMG (p=0.1705, paired t-test). There was a significant difference between baseline vs. GlcN+ TMG (Fig. 8Biii, p=0.0004).

Similar to THDOC, we noted variability in the progesterone response (Fig. 8Bii), eIPSCs recorded from some cells having a potentiation (Fig. 8Ci-Ciii, n = 7/13), and others, no potentiation (Fig. 8Di-Diii, n =6/13). To measure the effect of progesterone in these two populations, we re-normalized eIPSC amplitudes at the end of the 10 min GlcN+TMG application to establish a new baseline, then progesterone values were normalized to the new baseline. Similar to forskolin (Fig.4Ai-iii) and THDOC (Fig.7Ci-iii), we found a significant potentiation of the eIPSC amplitude in this subset of cells (134.4 ± 7.6%, p= 0.0041, paired t-test, n=7/13). In the remaining cells, (Fig. 8Di-iii), we found a slight but significant decrease (96.4 ± 1.3%, p=0.0443, paired t-test).

Figure 8. Progesterone also reverses O-GlcNAc iLTD, mimicking forskolin in CA1 pyramidal cells. Ai. Normalized data showing average eIPSC amplitude during progesterone (10 min, pink) followed by GlcN+TMG (10 min, orange). **Aii.** Average progesterone and GlcN+TMG eIPSC amplitude from the original baseline (gray bar, dotted line): 92.2% ± 7.3% (Progesterone), p=0.701 and 76.9% ± 9.1% (GlcN+TMG), p=0.126 (GlcN+TMG) and each other (p=0.008); (p=0.047, RM-ANOVA). **Aiii.** The average eIPSC amplitudes during the last 5 min of each condition corresponding to a,b,c in Ai (p= 0.0328, RM-ANOVA, n = 7 cells, 7 rats). Šídák's: b-c (p=0.008). **Bi.** Normalized data showing average eIPSC amplitude during GlcN+TMG (10 min, orange) followed by progesterone (10 min, pink). **Bii.** Average GlcN+TMG and progesterone eIPSC amplitude from the original baseline (gray bar, dotted line): 68.7% ± 3.8%, p<0.001 (GlcN+TMG), and 75.3% ± 4.9% (progesterone), p=0.0011 and each other (p=0.428); (p<0.0001, RM-ANOVA). **Biii.** The average eIPSC amplitudes during the last 5 min of each condition corresponding to a,b,c in Bi (p= 0.0007, RM-ANOVA, n= 13 cells, 10 rats) Šídák's: a-b (p=0.0004). *Potentiated:* **Ci.** Normalized data showing average eIPSC amplitude during GlcN+TMG

followed by progesterone. **Cii.** Average GlcN+TMG and progesterone eIPSC amplitude from the original baseline (gray bar, dotted line): 65.9% ± 6.2% (GlcN+TMG), p= 0.0045 and 83.3% ± 7.1% (progesterone), p=0.175 and each other (p=0.019); (p=0.002 RM-ANOVA). **Ciii.** Last 5 min average amplitudes corresponding to a,b,c (p=0.002, RM-ANOVA, n = 7 cells, 7 rats) Šídák's: a-b (p=0.0019), b-c (p=0.037). *Non-potentiated:* **Di.** Normalized data showing average eIPSC amplitude during GlcN+TMG followed by progesterone. **Dii.** Average GlcN+TMG and progesterone on eIPSC amplitude from the original baseline (gray bar, dotted line): 71.8% ± 4.7% (GlcN+TMG), p= 0.006 and 65.9% ± 5.0% (progesterone), p=0.0031 and each other (p=0.591); (p<0.0001, RM-ANOVA). **Diii.** Last 5 min average amplitudes corresponding to a,b,c (p=0.004, RM-ANOVA, n = 6 cells, 6 rats), Šídák's: a-b (p=0.015), a-c (p=0.0096). *Inset:* overlaid averaged representative eIPSC traces during baseline (black), GlcN + TMG (orange), progesterone (pink) (left) and scaled (right). Gray horizontal bars represent mean ± SEM. Šídák's post hoc depicted with asterisks.

Discussion

We previously reported that pharmacologically increasing protein O-GlcNAcylation rapidly depresses spontaneous and evoked GABA$_A$R-mediated synaptic inhibition[22]. The depression of evoked IPSCs is long-lasting, representing a novel form of LTD of synaptic inhibition, referred to here as O-GlcNAc iLTD. In the current study, we extend those initial findings by investigating whether increasing O-GlcNAcylation triggers GABA$_A$R endocytosis and how O-GlcNAcylation and serine phosphorylation interact in modulating the strength of GABAergic inhibition on principle cells in hippocampus.

This finding suggests that O-GlcNAcylation has a multiprong approach where manipulates the conductance of the GABA$_A$R, but also triggers a form of endocytosis in a subunit specific manner, similar to reports by [37]. Second, we found that forskolin reverses the O-GlcNAcylation dependent decrease in GABA$_A$R current amplitude and this modulation in polarity is determined by a co-occurring O-GlcNAc modification. Third, we sought to examine whether O-GlcNAcylation and phosphorylation interact to affect GABA$_A$R function through forskolin, and we found that the forskolin dependent increase in GABA$_A$R current amplitude following a prior acute increase in O-GlcNAcylation is not PKA dependent, but rather through a different mechanism. Finally, we show that THDOC and progesterone mimic forskolin upon prior increase of O-GlcNAcylation, supporting previous findings that forskolin may be acting on a neurosteroid site on GABA$_A$Rs [25] and discovering that O-GlcNAcylation enhances the mechanism of action of neurosteroids in a possible population or subunit dependent manner by modulating GABA$_A$R synaptic transmission and reversing O-GlcNAc LTD. These results further support our findings that protein O-GlcNAcylation serves as a novel regulator of both synaptic inhibition and excitation[20–22], but also provides highly novel information that O-GlcNAc dictates the polarity of the

change in GABA$_A$R synaptic current amplitude mediated by forskolin, THDOC and progesterone, highlighting O-GlcNAcylation's ability to modify the effectiveness of allosteric modulators on GABAergic transmission.

Serine phosphorylation has also been shown to regulate trafficking of GABA$_A$Rs currents depending on the kinase and neuron type [12]. This study discovered that in CA1 pyramidal cells only, PKA activation through forskolin caused a decrease in synaptic GABA$_A$R transmission, while PKC activation only increased the peak amplitude of miniature IPSCs in dentate granule cells. Hence, we suspected that if phosphorylation is triggering endocytosis in a kinase-and region-specific manner, O-GlcNAcylation triggers endocytosis of the GABA$_A$R, and impacts the mechanism of action of phosphorylation on GABA$_A$R transmission. Because our findings show that the O-GlcNAc mediated depression of GABAergic transmission is not due to the internalization of GABA$_A$Rs, several mechanisms may be involved that could be changing the conductance of individual GABA$_A$Rs or destabilizing scaffolding proteins that anchor GABA$_A$Rs to the membrane. Many studies have shown that phosphorylation of recombinant GABA$_A$Rs cause diverse functional effects ranging from enhancements to inhibitions depending on the identity of the expressed subunits [38] and the phosphorylation of specific serine/threonine residues on GABA$_A$Rs [10] , which may be the case for O-GlcNAcylation. Additionally, experimental GABAergic research has uncovered that GABA$_A$R subunit composition also directly influences single channel conductance, gating responses and their response to allosteric modulators[39,40], which may play a factor in the GABAergic response upon increased O-GlcNAcylation. Future experiments to determine how O-GlcNAcylation modulates various GABA$_A$Rs subunit confirmations would shed more light on whether it plays a role in the observed reduction in GABA$_A$R currents.

Notably, published literature details how the crosstalk between O-GlcNAcylation and phosphorylation [41,42] affects the regulation of various proteins involved in transcription, translation, and trafficking [43,44]. For example, O-GlcNAc and phosphorylation compete for the same serines on Tau, such that increasing O-GlcNAc can prevent Tau hyperphosphorylation, and thus accumulation of pTau [44]. We sought to elucidate whether there was a positive or negative interplay between O-GlcNAcylation and phosphorylation via Forskolin. However, we show for the first time that, though forskolin reverses the polarity of eIPSC amplitude following O-GlcNAcylation, this modulation is not PKA dependent and occurs through another mechanism, supported by our 1,9 Dideoxyforskolin results. Upon these

findings, we sought to determine the mechanism in which forskolin is triggering a polarity reversal because it is a common modulator of PKA through adenylate cyclase.

As it pertains to GABA$_A$R amplitude changes upon forskolin application, there are inconsistent results upon electrophysiological recordings whereby it can potentiate or suppress GABA-activated currents depending on the expression system [45–48]. Additionally, Li & Yang, 2001 outlines how the forskolin mediated potentiation or suppression on current amplitude depends on cAMP [27,45,49–51] and on the presence of different GABA$_A$ isoforms in different regions of the central nervous system [52,53]. However, the caveat to these findings it that forskolin modulation of receptor desensitization is not due to cAMP-dependent phosphorylation, but rather acts directly on the extracellular binding site on GABA receptors shared with neurosteroids like THDOC [25]

Neurosteroids and metabolites of progesterone positively modulate GABA$_A$Rs in a dose dependent manner by acting on synaptic and extrasynaptic GABA$_A$Rs, and their mechanism of action can be enhanced or diminished depending on the activation or inhibition of serine phosphorylation of GABAARs as well as GABA$_A$R subunit expression [10,14,54]. Our data supports these findings because while THDOC and progesterone reversed the O-GlcNAc-mediated depression of evoked IPSCs, mimicking forskolin, some cells did not exhibit this polarity reversal, suggesting that there may be a GABA$_A$R subunit confirmation preference and subunit expression, being that the subunit composition of GABA$_A$Rs can vary across the same cell types, leading to different responses upon drug exposure. In addition, our findings indicate that while forskolin can be binding to a different GABA$_A$R site and allosterically modulating the neurosteroid binding site sensitivity, we can conclude that O-GlcNAcylation somehow enhances access to the neurosteroid site on GABA$_A$Rs for forskolin and allosteric modulators to act. Because GABA$_A$Rs are therapeutic targets for drugs used in the treatment of neurological and neuropsychiatric conditions, understanding how they are modulated by O-GlcNAcylation has clinical implications, particularly if O-GlcNAc interferes with or enhances their efficacy. Collectively, our published studies have solidified O-GlcNAcylation as a critical sensor that impacts the strength of synaptic inhibition and excitation, and our current findings uncovers how protein O-GlcNAcylation can possibly serve as a gauge for the potency of allosteric modulators in a subunit-specific manner and a target for novel therapies.

Materials and Methods:

All experimental procedures were approved by the Medical University of South Carolina Institutional Animal Care and Use Committee and follow the National Institutes of Health experimental guidelines.

Hippocampal slice preparation

Male and female Sprague Dawley rats (age 3–7 weeks; Charles River Laboratories) were anesthetized with isoflurane, decapitated, and brains removed; 400 μm coronal slices from dorsal hippocampus were prepared on a VT1200S vibratome (Leica Biosystems) in oxygenated (95%O2/5%CO2) ice-cold, high sucrose cutting solution (in mM as follows: 85.0 NaCl, 2.5 KCl, 4.0 MgSO4, 0.5 CaCl2, 1.25 NaPO4, 25.0 glucose, 75.0 sucrose). After cutting, slices were held at room temperature for 1 to 5 hr in a submersion chamber with continuously oxygenated standard ACSF (in mM as follows: 119.0 NaCl, 2.5 KCl, 1.3 MgSO4, 2.5 CaCl2, 1.0 NaH2PO4, 26.0 NaHCO3, 11.0 glucose).

Electrophysiology

All recordings were performed in a submersion chamber with continuous perfusion of oxygenated standard ACSF. The blind patch technique was used to acquire interleaved whole-cell recordings from CA1 pyramidal neurons and dentate granule cells. Neuronal activity was recorded using an Axopatch 200B amplifier and pClamp10.7 acquisition software (Molecular Devices, Sunnyvale, CA). Signals were filtered at 5 kHz and digitized at 10 kHz (Digidata 1440). Patch pipettes (BF150-110 HP; Sutter Instruments, Novato, CA) were pulled on a Sutter P-97 (Sutter Instruments, Novato, CA) horizontal puller to a resistance between 3-5 MΩ. Spontaneous IPSCs were recorded using CsCl internal solution (in mM: 140.0 CsCl, 10.0 EGTA, 5.0 MgCl2, 2.0 Na-ATP, 0.3 Na-GTP, 10.0 HEPES; $E_{Cl} = 0$ mV). Evoked GABA$_A$R currents were recorded using Cs-gluconate internal solution (in mM: 100.0 Cs-gluconate, 0.6 EGTA, 5.0 MgCl2, 2.0 Na-ATP, 0.3 Na-GTP, 40.0 HEPES; $E_{Cl} = -60$ mV) with a twisted nichrome wire bipolar electrode positioned in stratum radiatum to activate Schaffer collateral axons (0.1 Hz, 100 μs duration) when recording in CA1 and positioned in the medial performant path when recording in the dentate gyrus. GABA$_A$R currents were pharmacologically isolated with bath perfusion of DNQX (10μM; Hello Bio) and R-CPP (5 μM; Hello Bio). Recordings were performed at a 10 mV test pulse at the end of each sweep to monitor series resistance and was excluded if there was more than a 20% change during the experiment. Stability of series resistance was verified using post-hoc scaling of averaged waveforms before and after pharmacologically increasing O-GlcNAcylation and after Forskolin, KT5720, SQ22536, 1,9 Dideoxyforskolin, Jasplakinolide and THDOC exposure.

Chemicals

Forskolin (Hello Bio) and 1,9 Dideoxyforskolin (Sigma-Aldrich) were prepared as a 50 mM stock in DMSO and stock was added to external solution for a final concentration of 50 µM. KT5720 (Hello Bio) was prepared as a 25 mM stock in DMSO and stock was added to external solution for a final concentration of 3 µM. SQ22536 (Tocris), 5α,21-pregnane-3α,21-diol-20-one (THDOC) (Sigma-Aldrich, mixed and sonicated), Progesterone (Sigma-Aldrich) was prepared as a 100 mM stock in DMSO and the stock was added to external solution for a final concentration of 100 µM, 10 µM, and 1 µM respectively. Jasplakinolide (Hello Bio) was prepared as a 1 mM stock in DMSO and stock was added to Cs-gluconate internal solution for a final concentration of 2 µM. Dynasore (Sigma-Aldrich) was prepared as a 100 mM stock in DMSO and stock was added to external solution for a final concentration of 80 µM. The actin stabilizer jasplakinolide (2µM, Hello Bio)[61,62] was included in the pipet solution to prevent $GABA_AR$ endocytosis. PKA dependent serine phosphorylation was triggered by bath application of the adenylate cyclase activator, forskolin (50µM, Hello Bio) [31] determine if the O-GlcNAc induced synaptic depression is prevented.

Alterations in the levels of O-GlcNAcylation

As in our previously published papers[20–22], the HBP substrate glucosamine (GlcN, 5 mM; Sigma) and OGA inhibitor, Thiamet-G (TMG) (10 min) (1 µM; Chem Molecules) were combined to acutely and robustly increase O-GlcNAcylation via bath application *in vitro*, ensuring a long-lasting effect.

Statistical analysis

Recordings were analyzed using Clampfit 11.2. GraphPad Prism 10.0.2, La Jolla, CA, was used for all analyses and outliers were excluded. Statistical significance was detected via two-tailed paired or unpaired Student's t-tests (parametric), or Wilcoxon matched-pairs signed rank test (non-parametric, paired) when comparing two-groups. For percent baseline calculations, the amplitudes of the last five minutes of each drug treatment for each cell were normalized to baseline or GlcN+TMG, as the new baseline, averaged and converted into percentages. Multi-groups were analyzed using RM-ANOVA with post hoc Šídák's multiple comparisons test (nonparametric). Data are displayed as mean ± SEM and p values considered statistically significant were as follows: $p \leq 0.05$, **$p \leq 0.01$,***$p \leq 0.001$,***$p \leq 0.0001$,****$p \leq 0.00001$.

Acknowledgements: This research was supported by NIH NINDS R01AG066489 to LLM and JCC, 1F99NS134163-01 to SMP

Author Contributions Statement: S.P. and L.L.M. designed the experiments; S.P. conducted experiments; S.P. and L.L.M. analyzed data; S.P, J.C.C., and L.L.M. wrote and revised the manuscript.

Data Availability Statement: The datasets generated and analyzed during the current study are available from the corresponding author on reasonable request.

Competing Interests Statement: The authors declare no competing interests.

1. Jacob, T. C., Moss, S. J. & Jurd, R. GABAA receptor trafficking and its role in the dynamic modulation of neuronal inhibition. *Nat Rev Neurosci* **9**, 331–343 (2008).
2. Macdonald, R. L., Kang, J.-Q. & Gallagher, M. J. Mutations in GABAA receptor subunits associated with genetic epilepsies. *J Physiol* **588**, 1861–1869 (2010).
3. Liang, J. & Olsen, R. W. Alcohol use disorders and current pharmacological therapies: the role of GABAA receptors. *Acta Pharmacol Sin* **35**, 981–993 (2014).
4. Mele, M., Costa, R. O. & Duarte, C. B. Alterations in GABAA-Receptor Trafficking and Synaptic Dysfunction in Brain Disorders. *Front Cell Neurosci* **13**, (2019).
5. Charych, E. I., Liu, F., Moss, S. J. & Brandon, N. J. GABAA receptors and their associated proteins: Implications in the etiology and treatment of schizophrenia and related disorders. *Neuropharmacology* **57**, 481–495 (2009).
6. Luscher, B., Shen, Q. & Sahir, N. The GABAergic deficit hypothesis of major depressive disorder. *Mol Psychiatry* **16**, 383–406 (2011).
7. Edinoff, A. N. *et al.* Brexanolone, a GABAA Modulator, in the Treatment of Postpartum Depression in Adults: A Comprehensive Review. *Front Psychiatry* **12**, (2021).
8. Abramian, A. M. *et al.* Neurosteroids promote phosphorylation and membrane insertion of extrasynaptic GABAA receptors. *Proceedings of the National Academy of Sciences* **111**, 7132–7137 (2014).
9. Wang, M. Neurosteroids and GABA-A Receptor Function. *Front Endocrinol (Lausanne)* **2**, (2011).
10. Nakamura, Y., Darnieder, L. M., Deeb, T. Z. & Moss, S. J. Regulation of GABAARs by Phosphorylation. in 97–146 (2015). doi:10.1016/bs.apha.2014.11.008.
11. Chuang, S.-H. & Reddy, D. S. Genetic and Molecular Regulation of Extrasynaptic GABA-A Receptors in the Brain: Therapeutic Insights for Epilepsy. *Journal of Pharmacology and Experimental Therapeutics* **364**, 180–197 (2018).
12. Comenencia-Ortiz, E., Moss, S. J. & Davies, P. A. Phosphorylation of GABAA receptors influences receptor trafficking and neurosteroid actions. *Psychopharmacology (Berl)* **231**, 3453–3465 (2014).
13. Hausrat, T. J. *et al.* Radixin regulates synaptic GABAA receptor density and is essential for reversal learning and short-term memory. *Nat Commun* **6**, 6872 (2015).
14. Poisbeau, P., Cheney, M. C., Browning, M. D. & Mody, I. Modulation of Synaptic GABAA Receptor Function by PKA and PKC in Adult Hippocampal Neurons. *The Journal of Neuroscience* **19**, 674–683 (1999).
15. Zeidan, Q. & Hart, G. W. The intersections between O-GlcNAcylation and phosphorylation: implications for multiple signaling pathways. *J Cell Sci* **123**, 13–22 (2010).
16. Wani, W. Y., Chatham, J. C., Darley-Usmar, V., McMahon, L. L. & Zhang, J. O-GlcNAcylation and neurodegeneration. *Brain Res Bull* **133**, 80–87 (2017).
17. Liu, Y. *et al.* Developmental Regulation of Protein O-GlcNAcylation, O-GlcNAc Transferase, and O-GlcNAcase in Mammalian Brain. *PLoS One* **7**, e43724 (2012).
18. Yang, X. & Qian, K. Protein O-GlcNAcylation: emerging mechanisms and functions. *Nat Rev Mol Cell Biol* **18**, 452–465 (2017).
19. Ruan, H.-B., Singh, J. P., Li, M.-D., Wu, J. & Yang, X. Cracking the O-GlcNAc code in metabolism. *Trends in Endocrinology & Metabolism* **24**, 301–309 (2013).
20. Taylor, E. W. *et al.* O-GlcNAcylation of AMPA Receptor GluA2 Is Associated with a Novel Form of Long-Term Depression at Hippocampal Synapses. *The Journal of Neuroscience* **34**, 10–21 (2014).

21. Stewart, L. T. *et al.* Acute Increases in Protein O-GlcNAcylation Dampen Epileptiform Activity in Hippocampus. *The Journal of Neuroscience* **37**, 8207–8215 (2017).

22. Stewart, L. T., Abiraman, K., Chatham, J. C. & McMahon, L. L. Increased O-GlcNAcylation rapidly decreases GABAAR currents in hippocampus but depresses neuronal output. *Sci Rep* **10**, 7494 (2020).

23. Cho, Y., Hwang, H., Rahman, Md. A., Chung, C. & Rhim, H. Elevated O-GlcNAcylation induces an antidepressant-like phenotype and decreased inhibitory transmission in medial prefrontal cortex. *Sci Rep* **10**, 6924 (2020).

24. Li, P. & Yang, X.-L. Forskolin modulation of desensitization at GABA A and glycine receptors is not mediated by cAMP-dependent protein kinase in isolated carp amacrine-like cells. *Pflugers Arch* **441**, 739–745 (2001).

25. McDonald, B. J. *et al.* Adjacent phosphorylation sites on GABAA receptor β subunits determine regulation by cAMP-dependent protein kinase. *Nat Neurosci* **1**, 23–28 (1998).

26. Moss, S. J., Smart, T. G., Blackstone, C. D. & Huganir, R. L. Functional Modulation of GABAA Receptors by cAMP-Dependent Protein Phosphorylation. *Science (1979)* **257**, 661–665 (1992).

27. Hwang, H. & Rhim, H. Acutely elevated O-GlcNAcylation suppresses hippocampal activity by modulating both intrinsic and synaptic excitability factors. *Sci Rep* **9**, 7287 (2019).

28. Wang, X., Yang, Y. & Zhou, Q. Independent Expression of Synaptic and Morphological Plasticity Associated with Long-Term Depression. *The Journal of Neuroscience* **27**, 12419–12429 (2007).

29. Luchkina, N. V. *et al.* Developmental switch in the kinase dependency of long-term potentiation depends on expression of GluA4 subunit-containing AMPA receptors. *Proceedings of the National Academy of Sciences* **111**, 4321–4326 (2014).

30. Otmakhov, N. *et al.* Forskolin-Induced LTP in the CA1 Hippocampal Region Is NMDA Receptor Dependent. *J Neurophysiol* **91**, 1955–1962 (2004).

31. Sokolova, I. V., Lester, H. A. & Davidson, N. Postsynaptic Mechanisms Are Essential for Forskolin-Induced Potentiation of Synaptic Transmission. *J Neurophysiol* **95**, 2570–2579 (2006).

32. Insel, P. A. Forskolin as a tool for examining adenylyl cyclase expression, regulation, and G protein signaling. *Cell Mol Neurobiol* **23**, 305–314 (2003).

33. Takeda, J. *et al.* Forskolin Activates Adenylate Cyclase Activity and Inhibits Mitosis in In Vitro in Pig Epidermis. *Journal of Investigative Dermatology* **81**, 236–240 (1983).

34. Seamon, K. B., Padgett, W. & Daly, J. W. Forskolin: unique diterpene activator of adenylate cyclase in membranes and in intact cells. *Proceedings of the National Academy of Sciences* **78**, 3363–3367 (1981).

35. Riddell, M. R., Winkler-Lowen, B., Jiang, Y., Davidge, S. T. & Guilbert, L. J. Pleiotropic Actions of Forskolin Result in Phosphatidylserine Exposure in Primary Trophoblasts. *PLoS One* **8**, e81273 (2013).

36. Stelzer, A., Slater, N. T. & ten Bruggencate, G. Activation of NMDA receptors blocks GABAergic inhibition in an in vitro model of epilepsy. *Nature* **326**, 698–701 (1987).

37. Muir, J. *et al.* NMDA receptors regulate GABAA receptor lateral mobility and clustering at inhibitory synapses through serine 327 on the γ2 subunit. *Proceedings of the National Academy of Sciences* **107**, 16679–16684 (2010).

38. Lu, Y. M., Mansuy, I. M., Kandel, E. R. & Roder, J. Calcineurin-Mediated LTD of GABAergic Inhibition Underlies the Increased Excitability of CA1 Neurons Associated with LTP. *Neuron* **26**, 197–205 (2000).

39. Wang, J. *et al.* Interaction of Calcineurin and Type-A GABA Receptor γ2 Subunits Produces Long-Term Depression at CA1 Inhibitory Synapses. *The Journal of Neuroscience* **23**, 826–836 (2003).

40. Kurotani, T., Yamada, K., Yoshimura, Y., Crair, M. C. & Komatsu, Y. State-Dependent Bidirectional Modification of Somatic Inhibition in Neocortical Pyramidal Cells. *Neuron* **57**, 905–916 (2008).

41. Saliba, R. S., Gu, Z., Yan, Z. & Moss, S. J. Blocking L-type Voltage-gated Ca2+ Channels with Dihydropyridines Reduces γ-Aminobutyric Acid Type A Receptor Expression and Synaptic Inhibition. *Journal of Biological Chemistry* **284**, 32544–32550 (2009).

42. Kittler, J. T. *et al.* Constitutive Endocytosis of GABAA Receptors by an Association with the Adaptin AP2 Complex Modulates Inhibitory Synaptic Currents in Hippocampal Neurons. *The Journal of Neuroscience* **20**, 7972–7977 (2000).

43. Vithlani, M., Terunuma, M. & Moss, S. J. The Dynamic Modulation of GABAA Receptor Trafficking and Its Role in Regulating the Plasticity of Inhibitory Synapses. *Physiol Rev* **91**, 1009–1022 (2011).

44. Zhou, C. *et al.* Altered Cortical GABAA Receptor Composition, Physiology, and Endocytosis in a Mouse Model of a Human Genetic Absence Epilepsy Syndrome. *Journal of Biological Chemistry* **288**, 21458–21472 (2013).

45. Brandon, N. J. *et al.* GABAA Receptor Phosphorylation and Functional Modulation in Cortical Neurons by a Protein Kinase C-dependent Pathway. *Journal of Biological Chemistry* **275**, 38856–38862 (2000).

46. Goetz, T., Arslan, A., Wisden, W. & Wulff, P. GABAA receptors: structure and function in the basal ganglia. in 21–41 (2007). doi:10.1016/S0079-6123(06)60003-4.

47. Mody, I. & Pearce, R. A. Diversity of inhibitory neurotransmission through GABAA receptors. *Trends Neurosci* **27**, 569–575 (2004).

48. Leney, A. C., El Atmioui, D., Wu, W., Ovaa, H. & Heck, A. J. R. Elucidating crosstalk mechanisms between phosphorylation and O-GlcNAcylation. *Proceedings of the National Academy of Sciences* **114**, (2017).

49. Song, H. *et al.* Global profiling of O-GlcNAcylated and/or phosphorylated proteins in hepatoblastoma. *Signal Transduct Target Ther* **4**, 40 (2019).

50. Bourré, G. *et al.* Direct Crosstalk Between O-GlcNAcylation and Phosphorylation of Tau Protein Investigated by NMR Spectroscopy. *Front Endocrinol (Lausanne)* **9**, (2018).

51. Veruki, M. L. & Yeh, H. H. Vasoactive intestinal polypeptide modulates GABAA receptor function through activation of cyclic AMP. *Vis Neurosci* **11**, 899–908 (1994).

52. Robello, M., Amico, C. & Cupello, A. Regulation of gabaa receptor in cerebellar granule cells in culture: Differential involvement of kinase activities. *Neuroscience* **53**, 131–138 (1993).

53. Heuschneider, G. & Schwartz, R. D. cAMP and forskolin decrease gamma-aminobutyric acid-gated chloride flux in rat brain synaptoneurosomes. *Proceedings of the National Academy of Sciences* **86**, 2938–2942 (1989).

54. Gillette, M. A. & Dacheux, R. F. Protein kinase modulation of GABAA currents in rabbit retinal rod bipolar cells. *J Neurophysiol* **76**, 3070–3086 (1996).

55. Feigenspan, A. & Bormann, J. Facilitation of GABAergic signaling in the retina by receptors stimulating adenylate cyclase. *Proceedings of the National Academy of Sciences* **91**, 10893–10897 (1994).

56. Han, Y. & Slaughter, M. M. Protein kinases modulate two glycine currents in salamander retinal ganglion cells. *J Physiol* **508**, 681–690 (1998).

57. Tehrani, M. H. J., Hablitz, J. J. & Barnes, E. M. cAMP increases the rate of GABAA receptor desensitization in chick cortical neurons. *Synapse* **4**, 126–131 (1989).
58. Zafra, F., Aragón, C. & Giménez, C. Molecular biology of glycinergic neurotransmission. *Mol Neurobiol* **14**, 117–142 (1997).
59. McKernan, R. M. & Whiting, P. J. Which GABAA-receptor subtypes really occur in the brain? *Trends Neurosci* **19**, 139–143 (1996).
60. Belelli, D. & Lambert, J. J. Neurosteroids: endogenous regulators of the GABAA receptor. *Nat Rev Neurosci* **6**, 565–575 (2005).
61. Kleschevnikov, A. M. *et al.* Hippocampal Long-Term Potentiation Suppressed by Increased Inhibition in the Ts65Dn Mouse, a Genetic Model of Down Syndrome. *The Journal of Neuroscience* **24**, 8153–8160 (2004).
62. Brown, M. T. C. *et al.* Drug-Driven AMPA Receptor Redistribution Mimicked by Selective Dopamine Neuron Stimulation. *PLoS One* **5**, e15870 (2010).

Figure 1. Acute increase in O-GlcNAcylation reduces spontaneous IPSCs in hippocampal CA1 pyramidal cells. Ai. (left) Schematic illustrating whole-cell recording of CA1 pyramidal cells (right) representative sIPSC trace showing baseline and GlcN + TMG application (top)and expanded time scale (bottom) during baseline (black) and GlcN + TMG application (orange). **Aii.** Cumulative probability distribution of sIPSC amplitude (p < 0.0001, KS D value = 0.17, Kolmogorov-Smirnov test); *inset*: (left) average sIPSC trace before (black) and after (orange) GlcN + TMG;scaled trace (dashed) shows no change in rise or decay time of depressed sIPSC. *inset*: (right) bar chart showing average (±SEM) sIPSC amplitude. Baseline: 69.7 ± 0.8 pA, GlcN + TMG: 43.5 ± 0.5 pA (p < 0.0001, Wilcoxon matched-pairs signed rank test, n = 9 cells, 6 rats). **Aiii.** Cumulative probability distribution of sIPSC interevent interval (p < 0.0001, KS D value = 0.225, Kolmogorov-Smirnov test). Baseline: 344.5 ± 4.7 ms, GlcN + TMG: 767.3 ± 16.6 ms, *inset*: (p < 0.0001, Wilcoxon matched-pairs signed rank test, n = 9 cells, 6 rats).

Figure 2. Blocking actin mediated GABA$_A$R endocytosis does not prevent O-GlcNAciLTD in CA1 pyramidal cells. Ai. Normalized data showing average eIPSC amplitude over time exposed to GlcN+TMG (10 min, orange) after a 15-minute baseline without jasp, (p= 0.005, paired t-test, n = 6 cells, 6 rats). *Inset*: representative eIPSC traces before (black) and after GlcN + TMG (orange) application and scaled average without jasp shows no change in rise or decay time of the eIPSC. **Aii.** Normalized data showing average eIPSC amplitude over time exposed to GlcN+TMG (10 min, orange) after a 15-minute baseline with jasplakinolide included in the pipette solution (jasp) (p= 0.0005, paired t-test, n = 8 cells, 8 rats*). Inset*: representative eIPSC traces before (black) and after GlcN + TMG (orange) application and scaled average with jasp shows no change in rise time or decay of the eIPSC.. **Aiii.** Overlay of the experimental groups showing no significant difference in the average (±SEM) eIPSC amplitude over time (p<0.0001, One-way ANOVA). Gray horizontal bars represent the mean ± SEM.

Figure 3. O-GlcNAc iLTD is partially dependent upon dynamin-mediated GABA$_A$R endocytosis.
Ai. Normalized data showing average eIPSC amplitude over time exposed to GlcN+TMG (10 min, orange) after a 15-minute baseline without dynasore (DMSO-vehicle) (p= 0.005, paired t-test, n = 8 cells, 7 rats). *Inset:* representative eIPSC traces before (black) and after GlcN + TMG (orange) application and scaled average shows no difference in rise or decay time of the eIPSC. **Aii.** Normalized data showing average eIPSC amplitude over time exposed to GlcN+TMG (10 min, orange) after a 5-minute baseline in slices incubated in dynasore for 30 min (p= 0.004, paired-test, n = 8 cells, 7 rats*).* *Inset*: representative eIPSC traces before (black) and after GlcN + TMG (orange) application and scaled average with dynasore. **Aiii.** Overlay of experimental groups showing no significant difference in the average (±SEM) eIPSC amplitude over time (p=0.08, One-way ANOVA). **Aiii.** Separation of two populations in the presence of dynasore where the eIPSC amplitude was significantly increased in one population (white, p=0.0392, paired t-test, n= 4 cells, 4 rats) and with no effect in the other population (red, p=0.13, paired t-test, n= 4 cells, 4 rats). Gray horizontal bars represent the mean ± SEM.

Figure 4. Magnitude of the O-GlcNAc iLTD is unaffected by prior application of forskolin. Ai. CA1: Normalized data showing average eIPSC amplitude over time during baseline (5 min), bath application of forskolin (10 min, blue) and GlcN + TMG (10 min, orange). *Inset:* overlaid averaged representative eIPSC traces during baseline (black), forskolin (blue), and GlcN + TMG (orange) (left) and scaled (right) **Aii.** Bar chart illustrates the average affect of forskolin and GlcN+TMG on eIPSC amplitude from the original baseline (dotted line): 89.2% ± 6.9%, p=0.39 (forskolin), 64.2% ± 4.4%, p<0.0001 (GlcN+TMG) and each other (p=0.0023); (p=0.0001, RM-ANOVA). **Aiii.** For each individual experiment in Ai, line graphs show the average eIPSC amplitude during the last the 5 min in each condition corresponding to a,b,c in Ai (p=0.0002, RM-ANOVA, n = 10 cells, 9 rats), Šídák's: a-c (p=0.002) and b-c (p=0.01). Baseline: 404.9 ± 47.9 pA, forskolin: 366.5 ± 50.2 pA, GlcN+ TMG: 256.6 ± 25.0 pA (mean ± SEM). **Aiv.** Paired-Pulse Ratio (100 ms inter-pulse interval) between baseline, forskolin and GlcN+TMG showed no significant difference (p=0.69, RM-ANOVA). **Bi.** Dentate: Normalized data showing average eIPSC amplitude over time during baseline, bath application of forskolin (blue), and GlcN+TMG (orange). *Inset*: overlaid averaged representative eIPSC traces during baseline (black) forskolin (blue), GlcN + TMG (orange) (left) and scaled (right). **Bii.** Bar chart illustrates the average affect of forskolin and GlcN+TMG from the original baseline (dotted line): 98.8% ± 4.6%, p=0.98 (forskolin) and 75.2% ± 6.0%, p=0.0128 (GlcN+TMG) and each other (p=0.0453); (p=0.004, RM-ANOVA). **Biii.** For each individual experiment in Bi, line graphs show the average amplitudes for last 5 min in each condition corresponding to a,b,c in Bi (p = 0.005, RM-ANOVA, n = 8 cells, 8 rats), Šídák's: a-c (p=0.029) and b-c (p=0.040). Baseline: 523.7 ± 51.7 pA, forskolin: 514.9 ± 50.5 pA, GlcN+ TMG: 360.2 ± 28.3 pA (mean ± SEM). **Biv.** Paired- Pulse Ratio between baseline, forskolin and GlcN+TMG showed no significant difference (p=0.46, RM-ANOVA). Gray horizontal bars represent the mean ± SEM. Šídák's post hoc depicted with asterisks.

Figure 5. Forskolin reverses O-GlcNAc iLTD in CA1 and dentate. CA1: **Ai.** Normalized data showing average eIPSC amplitude over time during baseline (5 min), bath application of GlcN+TMG (10 min, orange), and forskolin (10 min, blue). *Inset:* overlaid averaged representative eIPSC traces during baseline (black), GlcN + TMG (orange) and forskolin (blue) (left) and scaled (right). **Aii.** Bar chart illustrates the average affect of GlcN+TMG and forskolin on eIPSC amplitude from the original baseline (dotted line): 65.4 ± 5.2%, p=0.0002 (GlcN+TMG) and 86.0% ± 6.3%, p=0.1561 (forskolin) of baseline and each other (p=0.0011); (p<0.0001, RM-ANOVA). **Aiii.** For each individual experiment in Ai, line graphs show the average eIPSC amplitudes during the last 5 min of each condition corresponding to a,b,c in Ai (p < 0.0001, RM-ANOVA, n = 11 cells, 10 rats), Šídák's: a-b (p=0.0007) and b-c (p=0.028), Baseline: 471.2 ± 64.6 pA, GlcN+ TMG: 304.6 ± 52.0 pA, Forskolin: 397.6± 77.2 pA (mean ± SEM). **Aiv.** Paired-Pulse Ratio between baseline, GlcN+TMG and forskolin showed no significant difference (p=0.06, RM-ANOVA). Dentate: **Bi.** Normalized data showing average eIPSC amplitude over time during baselinebath application of GlcN +TMG (10 min, orange) and forskolin (10 min, blue). *Inset*: overlaid averaged representative eIPSC traces during baseline(black), GlcN + TMG (orange), and forskolin (blue) (left) and scaled (right). **Bii.** Bar chart illustrates the average affect of GlcN+TMG and forskolin on eIPSC amplitude from the original baseline (dotted line): 82.3% ± 2.6%, p=0.002 (GlcN+TMG) and 118.8% ± 9.3%, p=0.2500 (forskolin) and each other (p=0.008); (p=0.009, RM-ANOVA). **Biii.** For each individual experiment in Bi, line graphs show the averaged eIPSC amplitudes for last 5 min of each condition corresponding to a,b,c in **Bi**. (p=0.015, RM-ANOVA, n = 7 cells, 6 rats), Šídák's: a-b (p=0.0340) and b-c (p=0.038), Baseline: 376.3 ± 58.7 pA, GlcN+ TMG: 306.5 ± 44.2 pA, Forskolin: 443.0± 77.1 pA (mean ± SEM). **Biv.** Paired-Pulse Ratio between baseline, GlcN+TMG and forskolin showed no significant difference (p=0.24, RM-ANOVA). Gray horizontal bars represent the mean ± SEM. Šídák's post hoc test depicted with asterisks.

Figure 6. Forskolin induced potentiation of the eIPSC amplitude following an increase in O-GlcNAc is resistant to inhibitors of adenylate cyclase and PKA, suggesting serine phosphorylation is not the mechanism. A. Schematic illustrating possible interaction between serine O-GlcNAcylation and PKA-dependent serine phosphorylation of the GABA$_A$ receptor with pharmacological activators and inhibitors. **Bi**. Normalized data showing average (±SEM) eIPSC amplitude over time from CA1 pyramidal cells with and without the PKA inhibitor KT5720 (3 μM) during baseline (5 min), GlcN+TMG (10 min, orange) and forskolin application (10 min, blue). KT5720 was applied for 10 mins before and during forskolin. **Bii,Biii**. For each individual experiment, line graphs show average eIPSC amplitudes for the last 5 min of each condition corresponding to a,b,c with KT5720 (white) (p=0.004, RM-ANOVA, n= 8 cells, 6 rats), Šídák's: a-b (p=0.002) and b-c (p=0.011) and without KT5720 (black) (p=0.007, RM-ANOVA, n= 9 cells, 6 rats), Šídák's: a-b (p=0.0011) and b-c (p=0.0126). **Ci.** Normalized data showing average (±SEM) eIPSC amplitude over time with and without the adenylate cyclase inhibitor, SQ22536 (100 μM) during baseline, GlcN+TMG (10 min, orange) and forskolin application (10 min, blue) **Cii, Ciii** For each individual experiment, line graphs show average eIPSC amplitudes for the last 5 min of each condition corresponding to a,b,c with SQ22536 (white)

(p=0.031, RM-ANOVA, n= 7 cells, 6 rats), Šídák's: a-b (p=0.031) and without SQ22536 (black) (p=0.03, RM-ANOVA, n= 6 cells, 5 rats), Šídák's: a-b (p=0.020). **Di.** Normalized data showing average (±SEM) eIPSC amplitude over time during baseline, GlcN+TMG (10 min, orange) and application of forskolin or adenylate cyclase inactive analog 1,9 Dideoxyforskolin (10 min, blue). **Dii.** The average amplitudes for last 5 minutes of each condition corresponding to a,b,c with 1,9 Dideoxyforskolin (p= 0.028 RM-ANOVA, n=7 cells, 5 rats), Šídák's: a-b (p=0.011) and b-c (p=0.037). **Diii.** The average amplitudes for last 5 minutes of each condition with forskolin (p=0.037, RM-ANOVA, n= 6 cells, 4 rats), Šídák's: a-b (p=0.041) and b-c (p=0.035). Gray horizontal bars represent the mean ± SEM. Šídák's post-hoc depicted with asterisks.

Figure 7. 5α-pregnane-3α,21-diol-20-one (THDOC) reverses the O-GlcNAc iLTD, mimicking forskolin in CA1 pyramidal cells. Ai. Normalized data showing average eIPSC amplitude during THDOC (10 min, green) followed by GlcN+TMG (10 min, orange). **Aii.** Average THDOC and GlcN+TMG eIPSC amplitude from the original baseline (gray bar, dotted line): 83.4 ± 6.9%, p=0.153 (THDOC) and 56.9%± 10.3%, p=0.016 (GlcN+TMG) and each other (p=0.093); (p=0.0032, RM-ANOVA). **Aiii.** The average eIPSC amplitudes during the last 5 min of each condition corresponding to a,b,c in Ai (p= 0.0059, RM-ANOVA, n = 7 cells, 7 rats), Šídák's: a-c (p=0.032). **Bi.** Normalized data showing average eIPSC amplitude during GlcN+TMG (10 min, orange) followed by THDOC (10 min, green). **Bii.** Average GlcN+TMG and THDOC eIPSC amplitude from the original baseline (gray bar, dotted line): 72.1% ± 3.3% (GlcN+TMG), p< 0.0001 and 76.7% ± 5.5% (THDOC), p=0.003 and each other (p=0.617); (p<0.0001, RM-ANOVA). **Biii.** The average eIPSC amplitudes during the last 5 min of each condition corresponding to a,b,c in Bi (p <0.0001, RM-ANOVA, n = 12 cells, 12 rats), Šídák's: a-b (p<0.0001), a- (p=0.005). *Potentiated:* **Ci.** Normalized data showing average eIPSC amplitude during GlcN+TMG followed by THDOC. **Cii.** Average GlcN+TMG and THDOC eIPSC amplitude from the original baseline (gray bar, dotted line): 75.8% ± 2.6% (GlcN+TMG), p= 0.0002 and 88.6% ± 4.0% (THDOC), p=0.089 and each other (p=0.034); (p=0.0002, RM-ANOVA). **Ciii.** Last 5 min average amplitudes corresponding to a,b,c (p=0.0003, RM-ANOVA, n = 7 cells, 7 rats), Šídák's: a-b (p=0.0002),

b-c (p=0.036). *Non-potentiated:* **Di.** Normalized data showing average eIPSC amplitude during GlcN+TMG followed by THDOC. **Dii.** Average GlcN+TMG and THDOC eIPSC amplitude from the original baseline (gray bar, dotted line): 66.9% ± 6.9% (GlcN+TMG), p= 0.026 and 59.9% ± 6.8% (THDOC), p=0.012 and each other (p=0.422); (p=0.0014, RM-ANOVA). **Diii.** Last 5 min average amplitudes corresponding to a,b,c (p=0.0099, RM-ANOVA, n = 5 cells, 5 rats), Šídák's: a-b (p=0.041), a-c (p=0.025). *Inset:* overlaid averaged representative eIPSC traces during baseline (black), GlcN + TMG (orange), THDOC (green) (left) and scaled (right). Gray horizontal bars represent mean ± SEM. Šídák's post-hoc depicted with asterisks.

Figure 8. Progesterone also reverses O-GlcNAc iLTD, mimicking forskolin in CA1 pyramidal cells. Ai. Normalized data showing average eIPSC amplitude during progesterone (10 min, pink) followed by GlcN+TMG (10 min, orange). **Aii.** Average progesterone and GlcN+TMG eIPSC amplitude from the original baseline (gray bar, dotted line): 92.2% ± 7.3% (Progesterone), p=0.701 and 76.9% ± 9.1% (GlcN+TMG), p=0.126 (GlcN+TMG) and each other (p=0.008); (p=0.047, RM-ANOVA). **Aiii.** The average eIPSC amplitudes during the last 5 min of each condition corresponding to a,b,c in Ai (p= 0.0328, RM-ANOVA, n = 7 cells, 7 rats). Šídák's: b-c (p=0.008). **Bi.** Normalized data showing average eIPSC amplitude during GlcN+TMG (10 min, orange) followed by progesterone (10 min, pink). **Bii.** Average GlcN+TMG and progesterone eIPSC amplitude from the original baseline (gray bar, dotted line): 68.7% ± 3.8%, p<0.001 (GlcN+TMG), and 75.3% ± 4.9% (progesterone), p=0.0011 and each other (p=0.428); (p<0.0001, RM-ANOVA). **Biii.** The average eIPSC amplitudes during the last 5 min of each condition corresponding to a,b,c in Bi (p= 0.0007, RM-ANOVA, n= 13 cells, 10 rats) Šídák's: a-b (p=0.0004). *Potentiated:* **Ci.** Normalized data showing average eIPSC amplitude during GlcN+TMG followed by progesterone. **Cii.** Average GlcN+TMG and progesterone eIPSC amplitude from the original baseline (gray bar, dotted line): 65.9% ± 6.2% (GlcN+TMG), p= 0.0045 and 83.3% ± 7.1% (progesterone), p=0.175 and each other (p=0.019); (p=0.002 RM-ANOVA). **Ciii.** Last 5 min average

amplitudes corresponding to a,b,c (p=0.002, RM-ANOVA, n = 7 cells, 7 rats) Šídák's: a-b (p=0.0019), b-c (p=0.037). *Non-potentiated:* **Di.** Normalized data showing average eIPSC amplitude during GlcN+TMG followed by progesterone. **Dii.** Average GlcN+TMG and progesterone on eIPSC amplitude from the original baseline (gray bar, dotted line): 71.8% ± 4.7% (GlcN+TMG), p= 0.006 and 65.9% ± 5.0% (progesterone), p=0.0031 and each other (p=0.591); (p<0.0001, RM-ANOVA). **Diii.** Last 5 min average amplitudes corresponding to a,b,c (p=0.004, RM-ANOVA, n = 6 cells, 6 rats), Šídák's: a-b (p=0.015), a-c (p=0.0096). *Inset:* overlaid averaged representative eIPSC traces during baseline (black), GlcN + TMG (orange), progesterone (pink) (left) and scaled (right). Gray horizontal bars represent mean ± SEM. Šídák's post hoc depicted with asterisks.