# Neurodevelopmental defects in Dravet syndrome *Scn1a*$^{+/-}$ mice: targeting GABA-switch rescues behavioral dysfunctions but not seizures and mortality.

Lara Pizzamiglio[1,2,3], Fabrizio Capitano[1,2,3], Evgeniia Rusina[1,2,3], Giuliana Fossati[4], Elisabetta Menna[4,5], Isabelle Léna[1,2,3], Flavia Antonucci[5,6*], Massimo Mantegazza[1,2,3,*]

1. Université Côte d'Azur, Valbonne-Sophia Antipolis, France.

2. CNRS UMR7275, Institute of Molecular and Cellular Pharmacology (IPMC), Valbonne-Sophia Antipolis, France.

3. Inserm U1323, Valbonne-Sophia Antipolis, France

4. IRCCS Humanitas Research Hospital, Rozzano, Milan, Italy

5. Institute of Neuroscience - National Research Council of Italy (CNR) c/o Humanitas Mirasole S.p.A, Rozzano, Milan, Italy.

6. Department of Biotechnology and Translational Medicine, University of Milan, Milan, Italy.

* Correspondence to:

Massimo Mantegazza, Institute of Molecular and Cellular Pharmacology (IPMC), University Côte d'Azur-CNRS UMR7275-Inserm U1323, 660 Route des Lucioles, 06560 Valbonne-Sophia Antipolis, France, Tel. +33 493953425, e-mail mantegazza@ipmc.cnrs.fr

Flavia Antonucci, Department of Biotechnology and Translational Medicine, University of Milan, Via Vanvitelli 32, 20133 Milan, Italy, Tel. Tel. +39 0250317095, e-mail flavia.antonucci@unimi.it;

**Running title**: Neurodevelopmental defects in Dravet syndrome mice.

**Keywords**: developmental and epileptic encephalopathy; GABA; epilepsy; autism; Nav1.1; sodium channel.

## Abstract

Dravet syndrome (DS) is a developmental and epileptic encephalopathy (DEE) caused by mutations

of the *SCN1A* gene (Na$_V$1.1 sodium channel) and characterized by seizures, motor disabilities and

cognitive/behavioral deficits, including autistic traits. The relative role of seizures and

neurodevelopmental defects in disease progression, as well as the role of the mutation in inducing

early neurodevelopmental defects before symptoms' onset, are not clear yet. A delayed switch of

GABAergic transmission from excitatory to inhibitory (GABA-switch) was reported in models of DS,

but its effects on the phenotype have not been investigated.

Using a multi-scale approach, here we show that targeting GABA-switch with the drugs KU55933

(KU) or bumetanide (which upregulate KCC2 or inhibits NKCC1 chloride transporters, respectively)

rescues social interaction deficits and reduces hyperactivity observed in P21 *Scn1a$^{+/-}$* DS mouse

model. Bumetanide also improves spatial working memory defects. Importantly, neither KU nor

bumetanide have effect on seizures or mortality rate. Also, we disclose early behavioral defects and

delayed neurodevelopmental milestones well before seizure onset, at the beginning of Na$_V$1.1

expression.

We thus reveal that neurodevelopmental components in DS, in particular GABA switch,  selectively

underlie some cognitive/behavioral defects, but not seizures. Our work provides further evidence

that seizures and neuropsychiatric dysfunctions in DEEs can be uncoupled and can have differential

pathological mechanisms. They could be treated separately with targeted pharmacological

strategies.

# Introduction

Developmental and epileptic encephalopathies (DEEs) are characterized by severe epileptic seizures
with early-onset and a variable amount of underlying neurodevelopmental impairment that tends
to worsen as a consequence of epilepsy, but it is difficult to discriminate the neurodevelopmental
and epileptic components of the disease [1]. Dravet syndrome (DS) is a severe DEE [2,3] caused by
heterozygous mutations of the *SCN1A* gene, which lead to loss of function of the $Na_V1.1$ voltage-
gated sodium channel [4]. Disease onset occurs during the first year of life with seizures triggered by
fever or hyperthermia, which persist also in adulthood, and subsequent appearance of various other
types of seizures with high rate of sudden unexpected death in epilepsy (SUDEP) [3,5]. Beyond the
severe epileptic phenotype, DS is characterized by developmental plateauing with motor
disabilities, cognitive deficits and neuropsychiatric symptoms that often overlap with symptoms of
autism spectrum disorder (ASD), which are permanent and greatly contribute to the burden of the
disease [6–9,3]. Three stages have been proposed in the natural history of DS: febrile, worsening and
stabilization [5]. The available drugs are symptomatic treatments with limited efficacy and insufficient
control of convulsive seizures, whereas there is not yet a treatment reported as clearly effective on
cognitive/behavioral dysfunctions [3,10]. Heterozygous knock-out $Na_V1.1$ (*Scn1a*$^{+/-}$) mice recapitulate
the human DS phenotype [11–13]. Hyperthermia-induced seizures can be induced from around
postnatal-day 15 (P15), equivalent to the febrile stage. Onset of spontaneous recurrent seizures is
at around P20 with high SUDEP rate, as well as cognitive/behavioral defects, including autistic-like
traits; in this period, which is equivalent to the human worsening stage, seizures are particularly
frequent and mortality particularly high. A stabilization of the phenotype is observed after around
P35, but cognitive/behavioral defects are permanent, as in patients. Mouse models have been
instrumental for showing that $Na_V1.1$ loss-of-function, caused by DS mutations, induces reduction
of sodium current and of excitability in GABAergic interneurons, leading to impaired GABAergic
synaptic transmission [12,14]. This mechanism has been confirmed by numerous studies in different
animal models and in patients, showing the pivotal role of GABAergic interneurons' dysfunctions in
DS [11,15–18].

The relative contribution of seizures and underlying neurodevelopmental impairments to the
encephalopathy, in particular to neuropsychiatric features, is still debated in DS, as for other DEEs.
In fact, some clinical studies highlighted mainly epileptic encephalopathy features, whereas others
showed developmental alterations even before the onset of epilepsy and reported that in some
cases the severity of epilepsy and cognitive defects were not correlated [19–23].

Although hypoexcitability of GABAergic neurons is the most convincing pathological mechanism and

therapeutic target in DS, the detailed molecular mechanisms underlying epilepsy and comorbidities,

their causal relationships and the precise time window at which the different pathological features

develop are still poorly understood, limiting the development of targeted therapeutic strategies. A

study has shown that in a DS *Scn1a* knock-out mouse model the reversal potential of GABA-evoked

currents is depolarized compared to WT mice between P13 and P21, consistent with a delay in the

switch of GABAergic transmission from excitatory to inhibitory (i.e. GABA-switch) [24]. Also, a

depolarized reversal potential for GABA-evoked currents was observed in *Xenopus* oocytes

transplanted with cell membranes prepared from autopsy brain tissues of DS patients [25]. However,

the effects of this dysfunction on the DS phenotype were not investigated, although delayed

mortality rescuing GABA-switch was reported in a model of sodium channel β1 subunit DEE [24], which

is a different clinical entity with different phenotype and pathological mechanisms compared to DS

[1,26,27].

The GABA-switch is a fundamental step of brain development that reflects the maturation time

course of GABAergic signaling; it is in general completed by P15 and depends on the electrochemical

$Cl^-$ gradient, which is set by the relative expression of cation–chloride transporters NKCC1 ($Na^+$-$K^+$-

$Cl^-$ cotransporter isoform 1) and KCC2 ($K^+$-$Cl^-$ cotransporter isoform 2)[28–30]. Importantly, a delay in

GABA-switch has been found in numerous models of neurodevelopmental disorders and it has been

proposed as a therapeutic target in patients [31–33]. Overall brain expression of $Na_V1.1$ starts around

P8 in mice and homozygous $Scn1a^{-/-}$ mice die by P15 [12,16], showing that $Na_V1.1$ loss of function can

induce early effects. Thus, GABA-switch may be a target in therapeutic approaches for DS, and early

neurodevelopmental defects may be implicated in DS.

Here, we investigated in DS $Scn1a^{+/-}$ mice overall neurodevelopmental defects in the period before

the onset of seizures, and we targeted GABA-switch in DS $Scn1a^{+/-}$ mice by performing specific

pharmacological interventions and assessing their effect on both epileptic and behavioral

phenotypes.

**Methods**

**Animals.**

Mice were group-housed in standard mouse cages, on a 12-h day/12-h night cycle and allowed free

access to food and water. We used the exon 25-floxed (Fl) *Scn1a* mice (purchased from MMRRC,

Stock No: 041829-UCD) [15,34], which was maintained on a C57BL/6J background. They were

genotyped between post-natal day 5 (P5) and P7 for the *Scn1a* Fl allele using the primers 5'-

CTTGATGTGTTGAAATTCAC-3' and 5'-TATAGAGTGTTTAATCTCAAC-3'. Meox2-Cre mice were

purchased from the Jackson Laboratory (Stock No: 003755) and maintained on a C57BL/6J

background. They were genotyped at P5-P7 for the presence of Cre using the primers 5'-

GGTTTCCCGCAGAACCTGAA-3' and 5'-CCATCGCTCGACCAGTTTAGT-3'. In the Meox2-Cre line, Cre

recombinase is expressed in all the tissues: it is observed in the epiblast as early as embryonic day

5 and subsequently in all the derivatives of the epiblast (Endoderm, Mesoderm and Ectoderm). For

the experiments, we used F1 mice generated mating Fl *Scn1a* homozygous females and Meox2-Cre

heterozygous males, comparing *Scn1a*$^{Fl/+}$:Meox2-Cre$^{+/-}$ mice as DS model (*Scn1a*$^{+/-}$) and

*Scn1a*$^{Fl/+}$:Meox2-Cre$^{-/-}$ mice as wild-type controls (WT). This strategy has been already used in

previous studies and *Scn1a*$^{Fl/+}$:Meox2-Cre$^{+/-}$ mice show a typical DS-like phenotype[34,35]. Additionally,

comparing C57Bl/6J WT and Meox2-Cre$^{+/-}$ mice (Suppl. Fig. 2), we showed that Meox2-Cre$^{+/-}$ mice

do not have delay in the neurodevelopmental features that we have tested. All experiments were

performed according to policies on the care and use of laboratory animals of the European Directive

2010/63/EU; animal experimental protocols have been validated by ethics committees (CIEPAL

Azur) and the French Ministry of research under the agreement number APAFIS#9901-

2017040616024590v6-PEA216/408 and APAFIS#15665-201805301624157v2-PEA499. Animals'

health was evaluated daily with a scale established a priori for signs of discomfort and pain

(disclosed in the approved animal experimentation protocols). We have calculated a priori the

number of mice to use per group with the software G-Power, evaluating the variability observed in

the experiments according to previous experience of the team in similar projects, preliminary

experiments and reports in the literature, and taking into account the average mortality rate of

*Scn1a*$^{+/-}$ mice. Animals were randomized for allocation within each experimental group and the

order of treatments and measurements were randomized with the online tool Randomizer.

**Pharmacological treatments.**

KU55933 (Tocris, S1092) at the indicated concentration (7,5 mg/kg) diluted in the vehicle solution

(DMSO) was delivered by intranasal administrations in a volume of 1 ml/kg, as in [36]. Animals were

treated every 72 hours from P15 to P21 (short treatment, two administrations) or until P30 (long

treatment) (Fig. 1C, 3A, 4A). Bumetanide (Sigma, B3023) at the indicated concentration (0,2 mg/kg)

diluted in the vehicle solution (PBS with 1% DMSO) was delivered with intraperitoneal daily

injections in a volume of 0,02 ml/g from P15 for one or two weeks depending on the experiment

(Fig. 3A & 4A). Similar Bumetanide treatments have been used in several in vivo studies of mouse

models of brain diseases [37–39]. For *ex-vivo* electrophysiology experiments, for which a single animal

per day is used, we treated animals in sliding windows that allowed us to compare treated and

controls animals from the same litter and to minimize the number of animals used for the study.

We did not observe differences comparing recordings within the sliding window.

**Preparation of brain slices and electrophysiological recordings.**

Procedures were similar to those reported in [40,41]. Mice were deeply anesthetized with Isoflurane

and decapitated. The brain was quickly removed and placed in ice-cold standard artificial

cerebrospinal fluid (s-ACSF), which contained (in mM): 125 NaCl, 2.5 KCl, 2 $CaCl_2$, 1 $MgCl_2$, 1.25

$NaH_2PO_4$, 25 $NaHCO_3$ and 25 glucose, saturated with 95 % O2 - 5 % $CO_2$. Acute coronal slices (300

μm thick) were prepared with a vibratome (HM650V, MicroM, Germany) in ice-cold s-ACSF. Slices

were then stored in a submerged chamber with s-ACSF at 34°C for at least one hour before the

beginning of the recordings. Slices were visualized using infra-red differential interference contrast

(DIC) microscopy (Nikon Eclipse FN1, Japan) equipped with a CCD camera (CoolSnap ES2,

Photometrics, USA). Electrophysiological signals were recorded with a Multiclamp 700B amplifier

(CV-7B headstage), a Digidata 1440A acquisition board and pClamp 10.3 software (Molecular

Devices, USA). Recordings were performed at 34°C using an inline solution heater and a

temperature controller (TC-344B, Warner Instruments, Hamden, CT, USA), while continuously

perfusing the slices with s-ACSF bubbled with 95% $O_2$/5% $CO_2$ (pH 7.4).

Whole-cell patch-clamp recordings were performed in CA1 pyramidal neurons of the hippocampus

using borosilicate glass pipettes containing (in mM): 120 K-gluconate, 15 KCl, 2 $MgCl_2$, 0.2 EGTA, 10

HEPES, 20 P-Creatine, $Na_2$-0.2 GTP, 2 $Na_2$-ATP, 0.1 leupeptin, adjusted to pH 7.25 with KOH (3–5 MΩ

resistance). For recordings of post synaptic currents (PSCs) QX314 1.5 mM was added to the internal

solution in order to stabilize the membrane potential. Whole-cell access resistance (10-25 MΩ) was

1 monitored and cells showing unstable access resistance (> 20 MΩ) were discarded. Spontaneous

2 glutamatergic currents (sEPSCs) were measured at the reversal potential for GABA-A receptor

3 mediated events (-60 mV) and spontaneous GABA-A receptor mediated currents (sIPSCs) were

4 measured at the reversal potential of glutamatergic events (+10 mV). To be considered,

5 sEPSCs/sIPSCs had to exceed a threshold of 4-times the standard deviation of the RMS noise

6 evaluated on an eventless window of the trace. The E/I ratio was calculated by dividing sEPSCs and

7 sIPSCs frequencies measured in the same neuron.

8 Juxtacellular-loose patch recordings were performed with the same pipettes used for whole cell

9 experiments, but filled with s-ACSF, and action potentials were recorded in the I=0 mode. Cells were

10 recorded for 3/5 minutes to obtain a stable baseline, isoguvacine (10 µM) was bath applied for 3

11 minutes and washed out for at least 10 minutes.

12 Data were not corrected for the liquid junction potential. Electrophysiological signals were filtered

13 at 2 kHz and sampled at 10 kHz. Data were analyzed off-line (pClamp-10 software, Axon

14 Instruments).

16 **Golgi staining.**

17 WT and *Scn1a*$^{+/-}$ mice (males and females; P11 or P21) were deeply anesthetized with intra-

18 peritoneal injections of Kétamine (100 mg/kg) et Xylasine (16 mg/kg) and intracardiacally perfused

19 with saline (0,9% NaCl) to remove blood followed by 4% paraformaldehyde. The brain was removed

20 and postfixed in 4% paraformaldehyde for 24 hours, then it was stained with Golgi-Cox solution as

21 described in [42,43] with slight modifications. For the impregnation the brains were stored at room

22 temperature in darkness for 14 days. After this period the Golgi-Cox solution was removed and

23 replaced by 25% sucrose solution for at least 48 hours. Coronal sections of 100 µm thickness were

24 obtained from the hippocampus using a vibratome (VT1000S, Leica, Germany) and collected in TBS.

25 Sections were mounted onto gelatin treated or Superfrost Plus Micro slides using TBS and let them

26 air dry for 24 hours. The sections were processed with ammonium hydroxide for 10 min, rinsed with

27 distilled water, dehydrated with descending ethanol series, and mounted with a xylene-based

28 medium. Spine density and morphology were evaluated on the secondary branches of apical

29 dendrite of pyramidal neurons located in the CA1 subfield of the dorsal hippocampus. At least 20

30 neurons per mice were evaluated.

**Western Blotting.**

The general procedure was described in [36]. Proteins were obtained starting from explanted brain

tissues (hippocampus) homogenized in sample buffer (sodium dodecyl sulphate 1% (SDS), 62.5 mM

Tris-HCl (pH 6.8), 290 mM sucrose) with Protease Inhibitor Cocktail and PhosSTOP (Roche). Protein

content was assessed by Coomassie Protein Assay Kit (Thermo Fisher). Sample absorbance was read

through a spectrophotometer (Eppendorf BioPhotometer plus) set to 550 nm. Protein content was

assessed through a bovine serum albumin-based standard curve. Proteins were separated by SDS-

PAGE electrophoresis, blotted, and incubated with primary antibody followed by HRP-conjugated

secondary antibody (Jackson ImmunoResearch) and revealed by Pierce ECL Western Blotting

Substrate (Thermo Fisher Scientific) through Fusion FX Vilber. βIII tubulin was used as normalizer.

The following primary antibodies were used: rabbit anti-KCC2 (Millipore), mouse anti-NKCC1 clone

T4 (Developmental Studies Hybridoma Bank) and mouse anti- βIII tubulin (Sigma-Aldrich).

**Seizure induction by hyperthermia.**

The animal was placed in a small incubator and the core temperature was continuously monitored

using a rectal probe. Body temperature was increased 0.5 °C every 2 min as previously described [44]

until either a behavioral seizure occurred, or a core body temperature of 42.5 °C was reached. The

severity of seizures was quantified using a modified Racine's scale [45]: 0 - No behavioral response; 1-

motionless starring (with orofacial automatism); 2 – Head nodding; 3 – Unilateral forelimb clonus;

4 - Bilateral forelimb clonus; 5 – Rearing and falling; 6 - Generalized tonic-clonic seizures. Thermal

induction was performed in P21 Scn1a$^{+/-}$ animals treated with KU or BUM or with their respective

controls.

**Video analysis of spontaneous seizures.**

Starting from P15, the first day of treatment with KU or BUM or the respective controls, littermates

were placed in standard cages with the mother and continuously monitored by CMOS video

cameras (HDCVI, 2 Mpx, Sony exmor) equipped with infrared light and connected to a digital video

recorder (DVR, HDVCI 1080P). Seizures frequency and clusters were measured with a noninvasive

semi-automated detection of convulsive seizures from videos, based on an open-source package of

two software programs that we developed for automated video acquisition (VASD) and semi-

automated detection of seizures (SASDI) [46].

**Behavioral tests (performed at P21).**

**Social Interaction.** The assay was performed as previously described [47]. The three-chamber

apparatus, slightly illuminated (30 lux), is a non-transparent Plexiglas box (30×60 cm) with two

transparent partitions that make left, center, and right chambers (30×20 cm). In the first 10-min

session, the test mouse was placed in the center of the three-chamber apparatus for habituation

purposes. The mouse was allowed to freely explore each chamber containing one empty wire cage

(11 cm high x10cm diameter). In the second 10-min session, the sociability phase, an age- and sex-

matched mouse of the same strain that had never been exposed to the test mouse was placed in

one of two wire cages in the side chambers. The location of the stranger mouse in the right or left

chamber was alternated between trials. The empty wire cage in the opposite chamber served as the

inanimate object. The test mouse was placed again in the center, and allowed to freely explore the

chambers for 10 min. The time spent by the test mouse sniffing and interacting with each of the two

wire cages was manually scored. The sociability index was calculated as the difference between the

time spent exploring the stranger mouse and the empty cage divided by the total time of exploration

of both cages.

**Y maze spontaneous alterations.** Spontaneous alternation was measured using a Y-shaped maze

constructed with three symmetrical grey solid plastic arms at a 120-degree angle (26 cm length, 10

cm width, and 15 cm height) as previously described [48]. Mice were individually placed at the end of

one arm and were allowed to freely explore the three arms for 8 minutes. Arm entry was defined

when all four limbs were within the arm. A triad was defined as a set of three arm entries, when

each entry was in a different arm of the maze. Videos were analyzed with the ANY-maze video-

tracking system (Stoelting Europe, Dublin, Ireland). The number of arm entries and the number of

triads were recorded to calculate the alternation percentage (generated by dividing the number of

triads by the number of possible alternations, total arms entries -2, expressed as percentage).

**Neurodevelopmental tests (performed from P7 to P14).**

**Righting reflex.** The pup was placed on its back and the time required to recover to a normal position

with the 4 paws touching the surface was measured manually. This reflex was evaluated at P7 and

P10-11 using a cut-off of 30 seconds. The test reflects both motor coordination and vestibular

function [49].

**Forepaw grasping reflex.** When a thin rod is placed against the palm of one paw with a light

pressure, the paw flexed to grasp the rod. We assessed if this motor reflex was acquired for one or

both forepaws on P7 and P10-11.

**Negative geotaxis reflex.** The pup was placed, head down, on a rough surface inclined at 15° [49]. The

time taken to turn 180° was measured manually on P10-P11 using a cut-off of 60s. As the righting

reflex, geotaxis reflex assesses both motor coordination and vestibular function.

**Gait.** The pups were placed in the center of a 15 cm diameter circle and the time needed to move

outside the circle task was measured on P10-P11 [50].

**Olfactory motivation (nest-bedding/homing test).** WT and $Scn1a^{+/-}$ mice were tested for olfactory

motivation at P10-11 by placing them in the center of a square (7 cm × 7 cm) plastic box as in [51].

One corner of the tray contained bedding from the home cage, while the opposite one contained

new clean bedding. Mice were placed with their heads pointed toward an empty corner of the tray,

forced to turn left or right to orientate toward the bedding corners. Mice were allowed up to one

minute to reach a corner, before the trial was stopped. Trials in which mice failed to move, or to

arrive at a bedding corner, were considered uncompleted trials, whereas those in which mice

arrived at a bedding corner within one minute were considered completed. The latency to reach

one of the bedding corners and the type of bedding corner reached were recorded for each of the

3 trials. In each trials the beddings were in different orientations relative to the mouse. The

percentage of success was calculated as the percentage of trials in which the mice reached the home

cage bedding on the total of 3 trials.

**Eye opening.** The number of open eyelids was checked at P13-P14.

**Data analysis and statistics.**

Data analysis was performed with pClamp 10 (Molecular Devices, USA), Excel (Microsoft Office,

USA), ImageJ, Prism9 (GraphPad, USA) and Origin2021 (OriginLab, USA). Statistical tests were

performed with Prism9 or Origin2021. The Fisher Exact test or the chi-square test were used for

comparing categorical data. Other comparisons were performed with the Student's t-test (two

groups, paired or unpaired) and one-way ANOVA followed by Fisher's LSD post-hoc test (three

groups) (19) for normally distributed data, the Mann-Whitney test (two unpaired groups) and the

Wilcoxon signed rank test (two paired groups) for non-normally distributed data. Normality was

tested with the Kolmogorov-Smirnov (KS) test. The KS test was used also for comparing two

empirical distributions. Differences were considered significant when P values where smaller than

0.05 (*P<0.05, **P<0.01, ***P<0.001, ****P<0.0001).

## Results

**GABA-switch is delayed in DS *Scn1a$^{+/-}$* mice.**

As DS model, we used F1 mice generated mating floxed homozygous *Scn1a* C57BL/6J females [15,34]

and heterozygote Meox2-Cre C57BL/6J males (Jackson Laboratories, stock No: 003755) to obtain

*Scn1a$^{+/-}$* mice. This strategy has been already used in previous studies and these *Scn1a$^{+/-}$* mice show

a typical DS-like phenotype [34,35].

We evaluated the effect induced by the activation of GABA-A receptors on the spontaneous

excitability of CA3 excitatory hippocampal pyramidal neurons in brain slices of P18-P21 *Scn1a$^{+/-}$*

mice, an age at which the GABA-switch should be accomplished and before onset of spontaneous

seizures in this model. Hippocampal CA3 neurons have been used in numerous studies that have

evaluated the kinetics of GABA-switch [29–33]. We performed current clamp (I=0) electrophysiological

recordings of spontaneous spiking activity in loose patch configuration, quantifying the

modifications of spiking activity induced by the application of the GABA-A receptor agonist

isoguvacine (10 μM) [52]. We found a preponderant depolarizing response leading to increased

excitability in pyramidal neurons of *Scn1a$^{+/-}$* mice, whereas in WT littermates most of the cells

showed a reduction of spike frequency, although the reduction did not reach statistical significance

(Fig. 1A,B). These results confirm that the GABA-switch is delayed in DS mouse models, as it has

been reported in a different DS *Scn1a* knock-out model measuring the reversal potential of GABA-

evoked currents [24].

**KU55933 normalizes GABA-switch and decreases the excitatory/inhibitory ratio in *Scn1a$^{+/-}$* mice**

A delayed GABA-switch has been implicated in numerous neurodevelopmental diseases [31–33,53]. We

investigated its consequences on the overall phenotype and its importance as pathological

mechanism in *Scn1a$^{+/-}$* DS mice. We used two different pharmacological strategies, targeting the

two chloride transporters that are responsible of the intracellular chloride homeostasis in neurons:

KCC2, involved in chloride efflux, and NKCC1, involved in chloride influx. We exploited KU55933 (KU)

which, blocking ATM kinase activity, acts on the GABA-switch by increasing the expression of KCC2

[36,54], and bumetanide, a FDA-approved NKCC1 specific antagonist [55–57]. The action of bumetanide

has been reported in numerous studies, whereas KU is less characterized. Therefore, we

investigated the effects of KU in *Scn1a$^{+/-}$* mice performing *ex-vivo* electrophysiological experiments.

We treated mice in the period preceding the onset of spontaneous seizures (equivalent to the

febrile stage in patients) using intranasal delivery of KU (7,5 mg/kg) to efficiently cross the BBB and

to target specifically and directly the central nervous system, as in [36,58]. We performed the treatment

(two administrations every 72 hours) with sliding windows starting from P12-P15 and the recordings

at P18-21 (three days after the last dose of KU or DMSO); biochemical analysis was performed at

P21 starting the treatment at P15 (Fig. 1C). The sliding windows allowed us to compare treated and

control mice from the same litter and to minimize the number of animal used for

electrophysiological recordings, which are performed with a single animal per day. The

electrophysiological evaluation of the effect of the treatment was obtained measuring the effect

induced by the application of the GABA-A receptor agonist isoguvacine on the spontaneous spiking

activity of CA3 pyramidal cells in hippocampal slices, which showed a rescue of the delayed GABA-

switch that we observed in control conditions in $Scn1a^{+/-}$ mice (Fig. 1D-E). Western blot experiments

showed that the dysregulation of the GABA-switch in $Scn1a^{+/-}$ mice was correlated to decreased

expression of KCC2, which was normalized by the treatment with KU (Fig. 1F). Conversely, we found

no changes in NKCC1 expression, neither in control nor upon KU treatment (Fig. 1G).

Notably, besides the GABA-switch (which is a postsynaptic feature), KU acts also on the overall

kinetics of maturation of synaptic transmission and thus on the overall features of the network [36,54].

Thus, we evaluated the effect of KU on spontaneous GABAergic (sIPSCs) and glutamatergic (sEPSCs)

post-synaptic currents in hippocampal CA1 pyramidal neurons of KU- or vehicle-treated $Scn1a^{+/-}$

mice, compared to untreated WT littermates (Fig. 2A). The properties of sIPSCs and sEPSCs depend

on both synaptic properties and network excitability, allowing to evaluate the overall features of

the network. We analyzed hippocampal CA1 pyramidal neurons because they have been

investigated in numerous previous studies and are clearly involved in the pathophysiology of DS

[59,60]. In vehicle-treated $Scn1a^{+/-}$ mice, we observed a reduced frequency of sIPSCs (Fig. 2B), as

already reported in numerous studies of $Scn1a$ loss of function. We also observed an increased

frequency of sEPSCs (Fig. 2C), possibly induced by the reduction of IPSCs, leading to an increase of

the excitatory/inhibitory ratio (Fig. 2D). We observed a small increase of sIPSCs and sEPSCs

amplitude analyzing cumulative distributions, which was not significant comparing mean values

(Fig2E, F). The KU treatment decreased the frequency and the amplitude of sEPSCs (Fig. 2C and 2F),

whereas we observed only minor effects on sIPSCs (Fig. 2B and 2E). Despite the small effect on

sIPSCs, the KU treatment was sufficient to rescue the E/I ratio of the cells (Fig. 2D). Although this is

an interesting result, it should be kept in mind that the E/I ratio is a general parameter, obtained

from mean values, which does not reflect the actual dynamical relationship between excitation and

inhibition in the neuronal network.

Overall, the KU treatment can rescue some of the dysfunctions that we observed, in particular the

GABA-switch delay.

**Targeting GABA switch in *Scn1a^+/-* mice has no effect on induced or spontaneous seizures**.

We performed *in vivo* experiments to evaluate the effect of KU and bumetanide on the epileptic

phenotype of *Scn1a^+/-* DS mice, because the two treatments can rescue GABA-switch targeting

different mechanisms. We treated animals starting from P15 (Fig. 3A) with intranasal

administrations of KU (7,5 mg/kg every 72 hours) as in [36], or with bumetanide (2 mg/kg IP daily),

method of administration that has been effectively used in numerous studies on different animal

models of brain diseases [37–39]. As for the *ex vivo* experiments, we began treatments at P15 to

simulate clinical treatments, which do not start before the onset of the disease at the febrile stage.

We initially evaluated the effect on hyperthermia-induced seizures at P21. Both treatments were

ineffective in modifying the threshold temperature for seizure induction (Fig. 3B, D) or the severity

of seizures, quantified using a modified Racine scale (Fig. 3C, E; see methods), which were similar to

the values that we observed in control *Scn1a^+/-* mice treated with vehicle. To further characterize

the contribution of the developmental delay linked to GABA-switch on the epileptic phenotype in

*Scn1a^+/-* DS mice, we quantified in separated animal cohorts spontaneous convulsive seizures

identified by semi-automatic analysis of high-resolution video recordings [46]. This method allowed to

avoid the surgical procedure for implanting electrodes for chronic recordings, which are difficult to

implement at early ages, and permitted to chronically evaluate the animals for a long period starting

from P15, the first day of the treatment. We performed video recordings for fifteen days (24/24h,

7/7d), from P15 to P30, on animals chronically treated with KU or bumetanide (Fig. 3A). Neither KU

nor bumetanide reduced the frequency of seizures (Fig. 3F, I). We also analyzed the number of

seizure clusters (we considered a cluster as two or more seizures with a delay of less than fifteen

minutes) and found no changes in animals treated with KU or bumetanide compared to controls

(Fig. 3G, J). Also, the two compounds had no effect on the survival rate of DS mice (Fig. 3H, K). Thus,

neither KU nor bumetanide had effect on hyperthermia-induced or spontaneous seizures, showing

that the long-lasting depolarizing effect of GABA does not play a crucial role on epileptogenesis or

seizures generation in *Scn1a^+/-* DS mice.

**Rescuing the GABA-switch delay ameliorates sociability, hyperactivity, and cognitive deficits in**

**juvenile *Scn1a^+/-* mice.**

Since cognitive/behavioral dysfunctions are a major issue in DS, we investigated whether the

pharmacological rescue by KU or bumetanide of the abnormal GABA-switch could ameliorate the

cognitive and behavioral deficits of $Scn1a^{+/-}$ mice. We began the treatment at P15, and we

performed behavioral tests at P21 (Fig. 4A). We analyzed juvenile animals before the establishment

of a strong epileptic phenotype to discriminate precisely the neurodevelopmental component

involved in behavioral features from secondary effects induced by seizures.

We initially focused on autism-like features, assessing deficits in social interactions, which have

been already reported in adult $Scn1a^{+/-}$ mice [11,61]. In the three-chamber test, $Scn1a^{+/-}$ male mice had

profound defects in social interactions already at P21. Notably, targeting GABA-switch with KU or

bumetanide rescued this behavioral defect. Both WT and $Scn1a^{+/-}$ displayed no preference for the

left or right chamber during the habituation period (Supplementary Fig. 1). However, when a

stranger mouse was placed in one of the chambers, WT mice spent more time in close interaction

with the stranger mouse than in the empty cage. Conversely, $Scn1a^{+/-}$ mice had no preference for

the stranger mouse. Remarkably, both KU and bumetanide treatments rescued the impairment in

social interaction (Fig. 4B, D). As already reported for adult mice [34], juvenile $Scn1a^{+/-}$ female mice

did not show impairments in the three-chamber test (Fig. 4C).

Then, we tested cognitive features in juvenile $Scn1a^{+/-}$ mice using the Y maze spontaneous

alternation test, which can be used to assess spatial working memory. Spontaneous alternation is a

behavior driven by the innate curiosity of rodents for exploring previously unvisited areas and can

be evaluated by allowing mice to explore freely all three arms of the maze. Moreover, the number

of entries in each arm and the total distance travelled can be used for evaluating locomotor (hyper-

) activity. We tested P21 $Scn1a^{+/-}$ mice treated with KU or bumetanide and compared them with WT

littermates, considering males and females separately (Fig. 5). Both groups of control $Scn1a^{+/-}$ male

mice displayed a statistically significant decrease in the percentage of spontaneous alternations

compared to WT males (Fig. 5A, D). In contrast, performances of $Scn1a^{+/-}$ females were not different

compared to WT female mice (Fig. 5G, J). Both male and female $Scn1a^{+/-}$ DS mice showed a higher

number of entries in the arms of the Y maze (Fig. 5B, E, H, K) and covered significantly longer

distances during exploration (Fig. 5C, F, I, L). These data are in line with previous reports of

hyperactive behavior in DS mouse models (52,51,70), and show that hyperactivity can be already

observed in the pre-epileptic period. As shown in Fig. 5A, treatment with KU did not improve the

dysfunctions of $Scn1a^{+/-}$ males in spontaneous alternations. Although it did not reach statistical

significance, KU treatment showed a loose trend towards the rescue of total distance travelled by

males (5C) and a strong trend towards reduction of both the number of entries and the total

distance travelled by females (Fig 5H, I). Notably, bumetanide rescued the reduced alternations, the

number of entries and the total distance travelled of *Scn1a*[+/-] males (Fig. 5D, E, F). In *Scn1a*[+/-] female

mice, bumetanide rescued the number of entries and partially improved the total distance travelled

(Fig. 5K, L).

Therefore, targeting GABA-switch has an effect on autism-like features, spatial memory and

hyperactive behavior in *Scn1a*[+/-] mice.

*Scn1a*[+/-] **mice display early neurodevelopmental defects before seizures appearance**

The timing of the GABA-switch is one of the features of GABA in neurodevelopment. In fact, GABA

has trophic actions during early development before GABA-switch onset; its reduced release caused

by Na$_V$1.1 loss of-function may decrease depolarizing actions, which are important in these

developmental stages (also for inducing the GABA-switch), and cause an overall delay in maturation

[31,32,63,64]. Thus, we investigated whether behavioral deficits in *Scn1a*[+/-] mice can be observed at early

postnatal stages, when Na$_V$1.1 is already expressed but before seizure onset. We quantified several

developmental milestones, in particular those frequently used for characterizing dysfunctions in

animal models of ASD [51,65]. We analyzed motor ability and coordination at P7 and P10-11 and found

no differences in righting reflex, grasping reflex, negative geotaxis, and gait (Fig. 6A-D). Olfactory

detection and motivation to reach the nest and mother is one of the earliest behavioral correlates

of social information processing, and a poor interest in reaching the home-cage bedding has been

reported in mouse models of autism and socially anhedonic syndromes [51,65,66]. We performed the

nest bedding olfactory motivation test at P10-11. As shown in Fig. 6E, *Scn1a*[+/-] mice displayed

significant impairments in reaching the home cage bedding compared to WT, with no differences in

the average time taken to reach the nest, consistent with no dysfunctions in sensory and motor

abilities. Moreover, eye opening was delayed in *Scn1a*[+/-] mice compared to WT at P13-14 (Fig. 6F), a

feature that has been related to slowed and abnormal development of neuronal networks, also

linked to anomalies in the development of the GABAergic system [67,68]. To evaluate if Na$_V$1.1 loss of

function can reduce network activities also at these stages, we recorded GABAergic (sIPSCs) and

glutamatergic (sEPSCs) post-synaptic currents in pyramidal cells of the CA1-area of hippocampal

slices from P11-12 *Scn1a*[+/-] mice (Fig.7A). We found a clear reduction in frequency (Fig.7C) and a

mild reduction in amplitude of sIPSCs (Fig.7E) already at this early stage of development.

Furthermore, we observed a mild reduction in the amplitude of sEPSCs, although their frequency

was not modified (Fig. 2F).

Features of dendritic spines are hallmarks of neurodevelopment and KCC2, which we found

downregulated in *Scn1a*[+/-] mice, is implicated in spine formation and maturation [69,70]. We evaluated

the features of dendritic spines of pyramidal neurons in the CA1 area of the hippocampus by using

Golgi staining (Fig.8 A-B) and indeed observed a decreased spine head width at both P11 and P21 in

*Scn1a*[+/-] mice (Fig.8C), whereas spine density was not modified (Fig.8D). This is consistent with an

impairment of dendritic spine maturation.

Taken together these data show a neurodevelopmental component in *Scn1a*[+/-] DS mice, with

alterations appearing at early developmental stages at the beginning of $Na_V1.1$ expression and

before the onset of seizures, in which the decreased release of GABA and the consequences on its

trophic actions may be involved.

# Discussion

The disease onset in DEEs is often considered as the onset of seizures, which is the initial most

evident symptom of the pathology, but other dysfunctions, in particular cognitive/behavioral ones,

may begin to develop earlier, and they are a persistent major burden in DS. Although numerous

studies have reported in DS mouse models hypoexcitability of GABAergic neurons and reduced

GABAergic synaptic transmission in the period preceding the onset of spontaneous seizures [11–

13,14(p1),15,18,71], little is known about the link between these early dysfunctions of neuronal networks

and of development of disease features, in particular of cognitive/behavioral defects. Notably, early

pre-epileptic dysfunctions in the GABAergic system may lead to neurodevelopmental delay and

contribute to the generation of neuropsychiatric symptoms [30,64,72]. Interestingly, a delayed GABA-

switch has been reported in models of DS [24,25], but its importance as a pathological mechanism in

DS has not been investigated yet. A study showed that its pharmacological rescue with bumetanide

was able to induce a slight delay of mortality in the $Scn1b^{-/-}$ mouse, which is however a model of

sodium channel β1 subunit DEE [24], a different clinical entity with specific pathologic mechanisms

and phenotype than DS [1,26,27].

Here we assessed the importance of the delay in the GABA-switch as a pathological mechanism in

$Scn1a^{+/-}$ mice, rescuing it by targeting either KCC2 or NKCC1 with KU55933 (KU) or bumetanide,

respectively, and evaluating the effect on both epileptic and cognitive/behavioral features. For

increasing the translational value of the results, we started the treatments at P15, an age that

corresponds to the onset of the febrile stage in patients. Both drugs rescued social interaction

deficits observed in P21 $Scn1a^{+/-}$ mice and ameliorated hyperactivity in the Y maze test; bumetanide

also improved spatial working memory deficits. Although its blood-brain barrier permeability is low,

similar bumetanide treatments have been effectively used in several *in vivo* studies of mouse

models of neurodevelopmental diseases [37–39,73]. In particular, our results are in agreement with

previous studies showing beneficial effects of bumetanide treatment on hyperactivity and social

behavior in other neuropsychiatric diseases: on hyperactivity in mice with dysfunctional $K_V7$ voltage-

gated $K^+$ channels [73] and on both hyperactivity and social behavior in patients with tuberous sclerosis

complex (TSC) [74]. Thus, bumetanide, as well as other drugs targeting GABA-switch that may be

approved in the future for clinical use (possibly including KU), could be beneficial also for DS

patients. Conversely, neither KU nor bumetanide ameliorated the epileptic phenotype

(hyperthermia-induced seizures, spontaneous seizures or mortality), although alterations of GABA-

switch and KCC2 expression level/activity have been identified in epilepsy [75] and in our *ex vivo*

experiments the KU treatment also decreased the frequency of spontaneous excitatory events in

*Scn1a[+/−]* mice . This may be linked to the lack of effect of the treatment that we used on the reduced

GABAergic synaptic transmission of *Scn1a[+/-]* mice, which characterizes DS models and that is

thought to be implicated in DS epileptogenesis. Taken together, our data suggest that GABA-switch

delay is implicated in neuropsychiatric comorbidities of DS, in particular sociability defects (a

prominent feature of ASD), but not in the epileptic phenotype.

Importantly, in addition to the delayed GABA-switch, we showed that DS mice display behavioral

defects and delayed neurodevelopmental milestones at very early stages before the age of

physiological GABA-switch onset, including olfactory discrimination deficits of nest bedding, which

we observed at P10, and delayed eye opening. This is consistent with an early effect of the *Scn1a*

mutation. In fact, we have shown that DS mice display a reduction of spontaneous GABAergic

synaptic currents in CA1 pyramidal hippocampal neurons already at P11-12, an early developmental

stage (well before seizure onset) in the first days after onset of $Na_V1.1$ expression, which in mice is

at around P8 [16]. As already proposed in numerous previous studies [11,12,14,76], the reduced frequency

of sIPSCs can directly depend on the $Na_V1.1$ loss-of-function induced by DS mutations, which leads

to a reduction of sodium current and excitability of GABAergic interneurons. However, this initial

dysfunction of the GABAergic system could induce also a more general maturation delay. In fact,

GABA exerts a trophic action at early stages of neurodevelopment promoting cell survival

differentiation and migration, expression of brain-derived neurotrophic factor (BDNF), and spine

maturation [69,77,78]. Notably, GABA itself can promote GABA-switch [79] and reduced GABAergic

transmission in *Scn1a[+/-]* mice may interfere with it. Delayed GABA-switch might also be interpreted

as a homeostatic response triggered by a reduction of the depolarizing action of GABA in DS models

(because of hypoexcitability of GABAergic neurons) that leads to an overcompensation. However,

delayed GABA-switch has been observed in numerous models of neuropsychiatric diseases and

should not be specifically linked to the reduction in GABAergic transmission observed in DS models.

Overall, because of its trophic role in neurodevelopment, the decreased GABA release in *Scn1a[+/-]*

mice might be implicated in the behavioral defects, reduced amplitude of sEPSCs, and modifications

of spine morphology that we have observed already at early developmental stages (P11-12).

Interestingly, beside reducing GABAergic transmission, $Na_V1.1$ loss-of-function can modify

developmental activity patterns of neuronal networks possibly interfering with neuronal maturation

[63,80]. Overall, in the disease trajectory of DS, the GABAergic interneuron hypo-activity induced by

Na$_V$1.1 loss-of-function, which also reduces GABA release, should be the initial main event upstream

to a series of further pathological dysfunctions, including the GABA-switch delay.

Our functional data correlates with the results of our western blot experiments, which showed

decreased expression of KCC2 in the hippocampus of *Scn1a$^{+/-}$* mice. This is at odds with the results

reported for another Scn1a$^{+/-}$ model that showed no modification of KCC2 expression, although a

delay in GABA-switch was observed [24]. However, that study used total brain membranes for western

blot and a mixed 129×C57BL/6J genetic background that gives milder cellular dysfunctions and

phenotype compared to the pure C57BL/6J mice that we used [12,81], which may explain the different

results.

Features of dendritic spines are a hallmark of neurodevelopment and KCC2, independently of its

chloride transport function, plays an important role in the development and morphology of

dendritic spines through interaction with post-synaptic proteins and signaling pathways, leading to

a remodeling of the spine's actin cytoskeleton [32,69,70]. This function could be implicated in the

reduced spine head size without reduction of spine density that we observed in *Scn1a$^{+/-}$* mice.

Consistently, KCC2 knock-out mice display spines with immature morphology (no obvious spine

heads), whereas the density of the dendritic protrusions is not affected [82]. Thin filopodia-like spine

heads are considered less mature, indicating early stages of synapse formation or a lack of synaptic

stabilization. Notably, altered spine morphology, including reduced head width, has been associated

with pathological conditions, in particular neurodevelopmental diseases [83]. Notably, also other

dysfunctions that we have observed in *Scn1a$^{+/-}$* mice (i.e. E/I imbalance, defective GABAergic

transmission, delay in neurodevelopmental milestones such as eye opening and GABA-switch) are

key features observed in animal models of neurodevelopmental disorders, for example Down

syndrome, Rett syndrome, Fragile X syndrome and the VPA model of autism spectrum disorder

[31,36,84–86].

Our data showing the effects on behavioral dysfunctions, but not on seizures, of treatments

targeting the delayed GABA-switch, as well as those showing early neurodevelopmental defects

before seizure onset, demonstrate that DS mice display a neurodevelopmental delay that is

specifically linked to the cognitive/behavioral phenotype, but not the epileptic phenotype. Thus, we

disclose a neurodevelopmental component in DS which selectively impacts the behavioral

phenotype, in particular autistic-like traits. We provide here new experimental evidence to the

hypothesis that cognitive/behavioral defects in DS can develop before the onset of spontaneous

seizures (as observed for some motor features and hyperactivity in the Scn1a$^{A1783V/+}$ knock-in mouse

model [87]), and that they can be uncoupled from seizures and have specific pathological mechanisms,

at least at the early stages of the disease. This can be relevant in a translational perspective. In fact,

DS treatments are currently started after seizure onset and are mainly targeted to alleviation of

seizures, whereas there are still no effective treatments against the cognitive/behavioral

comorbidities, which are among the most severe phenotypic features, especially at later stages of

the disease. Our results suggest novel therapeutic options for DS that can target

cognitive/behavioral defects separately from seizures with specific pharmacological interventions

that should be started as early as possible. In particular, targeting the GABA-switch may be an

innovative pharmacological strategy to ameliorate DS behavioral defects.

Overall, our results are consistent with partially specific pathological mechanisms for seizures and

some cognitive/behavioral defects in DS. This is a novel major finding for DS and DEEs. The early

reduction of activity of GABAergic interneurons and the consequent reduction of GABA release in

DS could determine a cascade of events resulting in abnormal neurodevelopmental trajectories,

including delayed GABA-switch, because of the role of neuronal activity and GABA in

neurodevelopment. These dysfunctions may be specifically targeted in therapy.

# Acknowledgments

We thank Emilie Bonnet (Institute of Molecular and Cellular Pharmacology, CNRS UMR 7275 and University Cote d'Azur) for her skillful technical support.

This work was funded by UCAJEDI (https://univ-cotedazur.fr/ucajedi-lidex-duniversite-cote-dazur, ANR-15-IDEX-01, France), Laboratory of Excellence "Ion Channel Science and Therapeutics" - LabEx ICST (https://www.labex-icst.fr/en, ANR-11-LABX-0015-01, France) and EJPRD *SCN1A*-UP! (ANR-20-RAR4-0001, France) to MM. LP received a postdoc fellowship from the Fondazione Umberto Veronesi, Italy. FA was supported by Italian Ministry of University and Research, PRIN: Progetti di ricerca di rilevanza nazionale, 2022 (https://prin.mur.gov.it/Pages/Index/119) Prot.20229XWM5L, and PRIN: Progetti di ricerca di rilevanza nazionale, 2017 Prot. 20175C22WM.

# Author contributions

LP: designing research studies, conducting experiments, acquiring data, analyzing data, interpreting data, writing the manuscript.

FC: conducting experiments, acquiring data, analyzing data.

ER: conducting experiments, acquiring data, analyzing data.

GF: conducting experiments, acquiring data, analyzing data.

EM: analyzing data, interpreting data.

IL: conducting experiments, acquiring data, analyzing data, interpreting data.

FA designing research studies, interpreting data.

MM: designing research studies, analyzing data, interpreting data, writing the manuscript, supervising the project, acquiring funding.

# Declaration of interests

The authors declare no competing interests.

# References

1. Guerrini R, Conti V, Mantegazza M, Balestrini S, Galanopoulou AS, Benfenati F. Developmental and epileptic encephalopathies: from genetic heterogeneity to phenotypic continuum. *Physiological Reviews*. 2023;103(1):433-513. doi:10.1152/physrev.00063.2021

2. Dravet C, Bureau M, Oguni H, Fukuyama Y, Cokar O. Severe myoclonic epilepsy in infancy: Dravet syndrome. *Adv Neurol*. 2005;95:71-102.

3. Wirrell EC, Hood V, Knupp KG, et al. International consensus on diagnosis and management of Dravet syndrome. *Epilepsia*. 2022;63(7):1761-1777. doi:10.1111/epi.17274

4. Claes L, Del-Favero J, Ceulemans B, Lagae L, Van Broeckhoven C, De Jonghe P. De novo mutations in the sodium-channel gene SCN1A cause severe myoclonic epilepsy of infancy. *Am J Hum Genet*. 2001;68(6):1327-1332. doi:10.1086/320609

5. Gataullina S, Dulac O. From genotype to phenotype in Dravet disease. *Seizure - European Journal of Epilepsy*. 2017;44:58-64. doi:10.1016/j.seizure.2016.10.014

6. Li BM, Liu XR, Yi YH, et al. Autism in Dravet syndrome: Prevalence, features, and relationship to the clinical characteristics of epilepsy and mental retardation. *Epilepsy & Behavior*. 2011;21(3):291-295. doi:10.1016/j.yebeh.2011.04.060

7. Brunklaus A, Ellis R, Reavey E, Forbes GH, Zuberi SM. Prognostic, clinical and demographic features in SCN1A mutation-positive Dravet syndrome. *Brain*. 2012;135(Pt 8):2329-2336. doi:10.1093/brain/aws151

8. Sinoo C, de Lange IML, Westers P, Gunning WB, Jongmans MJ, Brilstra EH. Behavior problems and health-related quality of life in Dravet syndrome. *Epilepsy & Behavior*. 2019;90:217-227. doi:10.1016/j.yebeh.2018.11.029

9. Ouss L, Leunen D, Laschet J, et al. Autism spectrum disorder and cognitive profile in children with Dravet syndrome: Delineation of a specific phenotype. *Epilepsia Open*. 2019;4(1):40-53. doi:10.1002/epi4.12281

10. Wirrell EC. Treatment of Dravet Syndrome. *Canadian Journal of Neurological Sciences*. 2016;43(S3):S13-S18. doi:10.1017/cjn.2016.249

11. Han S, Tai C, Westenbroek RE, et al. Autistic-like behaviour in Scn1a+/− mice and rescue by enhanced GABA-mediated neurotransmission. *Nature*. 2012;489(7416):385-390. doi:10.1038/nature11356

12. Yu FH, Mantegazza M, Westenbroek RE, et al. Reduced sodium current in GABAergic interneurons in a mouse model of severe myoclonic epilepsy in infancy. *Nat Neurosci*. 2006;9(9):1142-1149. doi:10.1038/nn1754

13. Mantegazza M, Cestèle S, Catterall WA. Sodium channelopathies of skeletal muscle and brain. *Physiological Reviews*. 2021;101(4):1633-1689. doi:10.1152/physrev.00025.2020

14. Mantegazza M, Broccoli V. SCN1A/NaV1.1 channelopathies: Mechanisms in expression systems, animal models, and human iPSC models. *Epilepsia*. 2019;60(S3):S25-S38. doi:10.1111/epi.14700

15. Cheah CS, Yu FH, Westenbroek RE, et al. Specific deletion of Na$_v$1.1 sodium channels in inhibitory interneurons causes seizures and premature death in a mouse model of Dravet syndrome. *Proc Natl Acad Sci USA*. 2012;109(36):14646-14651. doi:10.1073/pnas.1211591109

16. Ogiwara I, Miyamoto H, Morita N, et al. Nav1.1 Localizes to Axons of Parvalbumin-Positive Inhibitory Interneurons: A Circuit Basis for Epileptic Seizures in Mice Carrying an Scn1a Gene Mutation. *Journal of Neuroscience*. 2007;27(22):5903-5914. doi:10.1523/JNEUROSCI.5270-06.2007

17. Stern WM, Sander JW, Rothwell JC, Sisodiya SM. Impaired intracortical inhibition demonstrated in vivo in people with Dravet syndrome. *Neurology*. 2017;88(17):1659-1665. doi:10.1212/WNL.0000000000003868

18. Kaneko K, Currin CB, Goff KM, et al. Developmentally regulated impairment of parvalbumin interneuron synaptic transmission in an experimental model of Dravet syndrome. *Cell Reports*. 2022;38(13):110580. doi:10.1016/j.celrep.2022.110580

19. Riva D, Vago C, Pantaleoni C, Bulgheroni S, Mantegazza M, Franceschetti S. Progressive neurocognitive decline in two children with Dravet syndrome, de novo SCN1A truncations and different epileptic phenotypes. *Am J Med Genet A*. 2009;149A(10):2339-2345. doi:10.1002/ajmg.a.33029

20. Guzzetta F. Cognitive and behavioral characteristics of children with Dravet syndrome: an overview. *Epilepsia*. 2011;52 Suppl 2:35-38. doi:10.1111/j.1528-1167.2011.02999.x

21. Catarino CB, Liu JYW, Liagkouras I, et al. Dravet syndrome as epileptic encephalopathy: evidence from long-term course and neuropathology. *Brain*. 2011;134(Pt 10):2982-3010. doi:10.1093/brain/awr129

22. Nabbout R, Chemaly N, Chipaux M, et al. Encephalopathy in children with Dravet syndrome is not a pure consequence of epilepsy. *Orphanet Journal of Rare Diseases*. 2013;8(1):176. doi:10.1186/1750-1172-8-176

23. Li W, Schneider AL, Scheffer IE. Defining Dravet syndrome: An essential pre-requisite for precision medicine trials. *Epilepsia*. 2021;62(9):2205-2217. doi:10.1111/epi.17015

24. Yuan Y, O'Malley HA, Smaldino MA, Bouza AA, Hull JM, Isom LL. Delayed maturation of GABAergic signaling in the Scn1a and Scn1b mouse models of Dravet Syndrome. *Sci Rep*. 2019;9(1):6210. doi:10.1038/s41598-019-42191-0

25. Ruffolo G, Cifelli P, Roseti C, et al. A novel GABAergic dysfunction in human Dravet syndrome. *Epilepsia*. 2018;59(11):2106-2117. doi:10.1111/epi.14574

26. Aeby A, Sculier C, Bouza AA, et al. SCN1B-linked early infantile developmental and epileptic encephalopathy. *Annals of Clinical and Translational Neurology*. 2019;6(12):2354-2367. doi:10.1002/acn3.50921

27. Scala M, Efthymiou S, Sultan T, et al. Homozygous SCN1B variants causing early infantile epileptic encephalopathy 52 affect voltage-gated sodium channel function. *Epilepsia*. 2021;62(6):e82-e87. doi:10.1111/epi.16913

28. Watanabe M, Fukuda A. Development and regulation of chloride homeostasis in the central nervous system. *Front Cell Neurosci*. 2015;9. doi:10.3389/fncel.2015.00371

29. Owens DF, Kriegstein AR. Is there more to gaba than synaptic inhibition? *Nat Rev Neurosci*. 2002;3(9):715-727. doi:10.1038/nrn919

30. Ben-Ari Y. Excitatory actions of gaba during development: the nature of the nurture. *Nat Rev Neurosci*. 2002;3(9):728-739. doi:10.1038/nrn920

31. Deidda G, Bozarth IF, Cancedda L. Modulation of GABAergic transmission in development and neurodevelopmental disorders: investigating physiology and pathology to gain therapeutic perspectives. *Frontiers in Cellular Neuroscience*. 2014;8. Accessed June 17, 2022. https://www.frontiersin.org/article/10.3389/fncel.2014.00119

32. Kaila K, Price TJ, Payne JA, Puskarjov M, Voipio J. Cation-chloride cotransporters in neuronal development, plasticity and disease. *Nat Rev Neurosci*. 2014;15(10):637-654. doi:10.1038/nrn3819

33. Khazipov R, Valeeva G, Khalilov I. Depolarizing GABA and Developmental Epilepsies. *CNS Neuroscience & Therapeutics*. 2015;21(2):83-91. doi:10.1111/cns.12353

34. Williams AD, Kalume F, Westenbroek RE, Catterall WA. A more efficient conditional mouse model of Dravet syndrome: Implications for epigenetic selection and sex-dependent behaviors. *Journal of Neuroscience Methods*. 2019;325:108315. doi:10.1016/j.jneumeth.2019.108315

35. Capitano F, Kuchenbuch M, Lavigne J, et al. Preictal dysfunctions of inhibitory interneurons paradoxically lead to their rebound hyperactivity and to low-voltage-fast onset seizures in Dravet syndrome. *Proceedings of the National Academy of Sciences*. 2024;121(23):e2316364121. doi:10.1073/pnas.2316364121

36. Pizzamiglio L, Focchi E, Cambria C, et al. The DNA repair protein ATM as a target in autism spectrum disorder. *JCI Insight*. 6(3):e133654. doi:10.1172/jci.insight.133654

37. Deidda G, Parrini M, Naskar S, Bozarth IF, Contestabile A, Cancedda L. Reversing excitatory GABAAR signaling restores synaptic plasticity and memory in a mouse model of Down syndrome. *Nat Med*. 2015;21(4):318-326. doi:10.1038/nm.3827

38. Sivakumaran S, Maguire J. Bumetanide reduces seizure progression and the development of pharmacoresistant status epilepticus. *Epilepsia*. 2016;57(2):222-232. doi:10.1111/epi.13270

39. Dargaei Z, Bang JY, Mahadevan V, et al. Restoring GABAergic inhibition rescues memory deficits in a Huntington's disease mouse model. *Proc Natl Acad Sci U S A*. 2018;115(7):E1618-E1626. doi:10.1073/pnas.1716871115

40. Chever O, Zerimech S, Scalmani P, et al. Initiation of migraine-related cortical spreading depolarization by hyperactivity of GABAergic neurons and NaV1.1 channels. *J Clin Invest*. 131(21):e142203. doi:10.1172/JCI142203

41. Lemaire L, Desroches M, Krupa M, Pizzamiglio L, Scalmani P, Mantegazza M. Modeling NaV1.1/SCN1A sodium channel mutations in a microcircuit with realistic ion concentration dynamics suggests differential GABAergic mechanisms leading to hyperexcitability in epilepsy and hemiplegic migraine. *PLoS Comput Biol*. 2021;17(7):e1009239. doi:10.1371/journal.pcbi.1009239

42. Menna E, Zambetti S, Morini R, et al. Eps8 controls dendritic spine density and synaptic plasticity through its actin-capping activity. *EMBO J*. 2013;32(12):1730-1744. doi:10.1038/emboj.2013.107

43. Bayram-Weston Z, Olsen E, Harrison DJ, Dunnett SB, Brooks SP. Optimising Golgi-Cox staining for use with perfusion-fixed brain tissue validated in the zQ175 mouse model of Huntington's disease. *J Neurosci Methods*. 2016;265:81-88. doi:10.1016/j.jneumeth.2015.09.033

44. Salgueiro-Pereira AR, Duprat F, Pousinha PA, et al. A two-hit story: Seizures and genetic mutation interaction sets phenotype severity in SCN1A epilepsies. *Neurobiology of Disease*. 2019;125:31-44. doi:10.1016/j.nbd.2019.01.006

45. Racine RJ. Modification of seizure activity by electrical stimulation: II. Motor seizure.
*Electroencephalography and Clinical Neurophysiology*. 1972;32(3):281-294. doi:10.1016/0013-
4694(72)90177-0

46. Diaz-Arce D, Ghouma A, Scalmani P, Mantegazza M, Duprat F. A python-based package for long-lasting
video acquisition and semi-automated detection of convulsive seizures in rodents. Published online
April 16, 2022:2022.04.15.488472. doi:10.1101/2022.04.15.488472

47. Sala M, Braida D, Lentini D, et al. Pharmacologic rescue of impaired cognitive flexibility, social deficits,
increased aggression, and seizure susceptibility in oxytocin receptor null mice: a neurobehavioral
model of autism. *Biol Psychiatry*. 2011;69(9):875-882. doi:10.1016/j.biopsych.2010.12.022

48. Begenisic T, Baroncelli L, Sansevero G, et al. Fluoxetine in adulthood normalizes GABA release and
rescues hippocampal synaptic plasticity and spatial memory in a mouse model of Down Syndrome.
*Neurobiology of Disease*. 2014;63:12-19. doi:10.1016/j.nbd.2013.11.010

49. McFadyen-Leussis MP, Heinrichs SC. Seizure-prone EL/Suz mice exhibit physical and motor delays and
heightened locomotor activity in response to novelty during development. *Epilepsy & Behavior*.
2005;6(3):312-319. doi:10.1016/j.yebeh.2005.01.010

50. Nguyen AT, Armstrong EA, Yager JY. Neurodevelopmental Reflex Testing in Neonatal Rat Pups. *J Vis
Exp*. 2017;(122). doi:10.3791/55261

51. Moldrich RX, Leanage G, She D, et al. Inhibition of histone deacetylase in utero causes sociability
deficits in postnatal mice. *Behavioural Brain Research*. 2013;257:253-264.
doi:10.1016/j.bbr.2013.09.049

52. Tyzio R, Nardou R, Ferrari DC, et al. Oxytocin-Mediated GABA Inhibition During Delivery Attenuates
Autism Pathogenesis in Rodent Offspring. *Science*. 2014;343(6171):675-679.
doi:10.1126/science.1247190

53. Schulte JT, Wierenga CJ, Bruining H. Chloride transporters and GABA polarity in developmental,
neurological and psychiatric conditions. *Neuroscience & Biobehavioral Reviews*. 2018;90:260-271.
doi:10.1016/j.neubiorev.2018.05.001

54. Focchi E, Cambria C, Pizzamiglio L, et al. ATM rules neurodevelopment and glutamatergic transmission
in the hippocampus but not in the cortex. *Cell Death Dis*. 2022;13(7):616. doi:10.1038/s41419-022-
05038-7

55. Ward OC, Lam LK. Bumetanide in heart failure in infancy. *Arch Dis Child*. 1977;52(11):877-882.
doi:10.1136/adc.52.11.877

56. Lemonnier E, Degrez C, Phelep M, et al. A randomised controlled trial of bumetanide in the treatment
of autism in children. *Transl Psychiatry*. 2012;2(12):e202-e202. doi:10.1038/tp.2012.124

57. Fernell E, Gustafsson P, Gillberg C. Bumetanide for autism: Open-label trial in six children. *Acta
Paediatr*. 2021;110(5):1548-1553. doi:10.1111/apa.15723

58. Lochhead JJ, Thorne RG. Intranasal delivery of biologics to the central nervous system. *Adv Drug Deliv
Rev*. 2012;64(7):614-628. doi:10.1016/j.addr.2011.11.002

59. Liautard C, Scalmani P, Carriero G, de Curtis M, Franceschetti S, Mantegazza M. Hippocampal
hyperexcitability and specific epileptiform activity in a mouse model of Dravet syndrome. *Epilepsia*.
2013;54(7):1251-1261. doi:10.1111/epi.12213

60. Jansen NA, Dehghani A, Breukel C, Tolner EA, van den Maagdenberg AMJM. Focal and generalized seizure activity after local hippocampal or cortical ablation of NaV 1.1 channels in mice. *Epilepsia*. 2020;61(4):e30-e36. doi:10.1111/epi.16482

61. Ito S, Ogiwara I, Yamada K, et al. Mouse with Nav1.1 haploinsufficiency, a model for Dravet syndrome, exhibits lowered sociability and learning impairment. *Neurobiology of Disease*. 2013;49:29-40. doi:10.1016/j.nbd.2012.08.003

62. Rubinstein M, Han S, Tai C, et al. Dissecting the phenotypes of Dravet syndrome by gene deletion. *Brain*. 2015;138(Pt 8):2219-2233. doi:10.1093/brain/awv142

63. Warm D, Schroer J, Sinning A. Gabaergic Interneurons in Early Brain Development: Conducting and Orchestrated by Cortical Network Activity. *Front Mol Neurosci*. 2022;14. doi:10.3389/fnmol.2021.807969

64. Peerboom C, Wierenga CJ. The postnatal GABA shift: A developmental perspective. *Neuroscience & Biobehavioral Reviews*. 2021;124:179-192. doi:10.1016/j.neubiorev.2021.01.024

65. Roullet FI, Wollaston L, deCatanzaro D, Foster JA. Behavioral and molecular changes in the mouse in response to prenatal exposure to the anti-epileptic drug valproic acid. *Neuroscience*. 2010;170(2):514-522. doi:10.1016/j.neuroscience.2010.06.069

66. Cinque C, Pondiki S, Oddi D, et al. Modeling socially anhedonic syndromes: genetic and pharmacological manipulation of opioid neurotransmission in mice. *Transl Psychiatry*. 2012;2(8):e155-e155. doi:10.1038/tp.2012.83

67. Chattopadhyaya B, Cristo GD, Higashiyama H, et al. Experience and Activity-Dependent Maturation of Perisomatic GABAergic Innervation in Primary Visual Cortex during a Postnatal Critical Period. *J Neurosci*. 2004;24(43):9598-9611. doi:10.1523/JNEUROSCI.1851-04.2004

68. Guan W, Cao JW, Liu LY, Zhao ZH, Fu Y, Yu YC. Eye opening differentially modulates inhibitory synaptic transmission in the developing visual cortex. Bartos M, ed. *eLife*. 2017;6:e32337. doi:10.7554/eLife.32337

69. Awad PN, Amegandjin CA, Szczurkowska J, et al. KCC2 Regulates Dendritic Spine Formation in a Brain-Region Specific and BDNF Dependent Manner. *Cerebral Cortex*. 2018;28(11):4049-4062. doi:10.1093/cercor/bhy198

70. Virtanen MA, Uvarov P, Mavrovic M, Poncer JC, Kaila K. The Multifaceted Roles of KCC2 in Cortical Development. *Trends in Neurosciences*. 2021;44(5):378-392. doi:10.1016/j.tins.2021.01.004

71. Almog Y, Fadila S, Brusel M, Mavashov A, Anderson K, Rubinstein M. Developmental alterations in firing properties of hippocampal CA1 inhibitory and excitatory neurons in a mouse model of Dravet syndrome. *Neurobiology of Disease*. 2021;148:105209. doi:10.1016/j.nbd.2020.105209

72. Tang X, Jaenisch R, Sur M. The role of GABAergic signalling in neurodevelopmental disorders. *Nat Rev Neurosci*. 2021;22(5):290-307. doi:10.1038/s41583-021-00443-x

73. Marguet SL, Le-Schulte VTQ, Merseburg A, et al. Treatment during a vulnerable developmental period rescues a genetic epilepsy. *Nat Med*. 2015;21(12):1436-1444. doi:10.1038/nm.3987

74. van Andel DM, Sprengers JJ, Oranje B, Scheepers FE, Jansen FE, Bruining H. Effects of bumetanide on neurodevelopmental impairments in patients with tuberous sclerosis complex: an open-label pilot study. *Mol Autism*. 2020;11(1):30. doi:10.1186/s13229-020-00335-4

75. Di Cristo G, Awad PN, Hamidi S, Avoli M. KCC2, epileptiform synchronization, and epileptic disorders. *Progress in Neurobiology*. 2018;162:1-16. doi:10.1016/j.pneurobio.2017.11.002

76. Almog Y, Fadila S, Brusel M, Mavashov A, Anderson K, Rubinstein M. Developmental alterations in firing properties of hippocampal CA1 inhibitory and excitatory neurons in a mouse model of Dravet syndrome. *Neurobiology of Disease*. 2021;148:105209. doi:10.1016/j.nbd.2020.105209

77. Cancedda L, Fiumelli H, Chen K, Poo M ming. Excitatory GABA Action Is Essential for Morphological Maturation of Cortical Neurons In Vivo. *J Neurosci*. 2007;27(19):5224-5235. doi:10.1523/JNEUROSCI.5169-06.2007

78. Obrietan K, Gao XB, van den Pol AN. Excitatory Actions of GABA Increase BDNF Expression via a MAPK-CREB–Dependent Mechanism—A Positive Feedback Circuit in Developing Neurons. *Journal of Neurophysiology*. 2002;88(2):1005-1015. doi:10.1152/jn.2002.88.2.1005

79. Ganguly K, Schinder AF, Wong ST, Poo M ming. GABA Itself Promotes the Developmental Switch of Neuronal GABAergic Responses from Excitation to Inhibition. *Cell*. 2001;105(4):521-532. doi:10.1016/S0092-8674(01)00341-5

80. De Marco García NV, Karayannis T, Fishell G. Neuronal activity is required for the development of specific cortical interneuron subtypes. *Nature*. 2011;472(7343):351-355. doi:10.1038/nature09865

81. Rubinstein M, Westenbroek RE, Yu FH, Jones CJ, Scheuer T, Catterall WA. Genetic background modulates impaired excitability of inhibitory neurons in a mouse model of Dravet syndrome. *Neurobiology of Disease*. 2015;73:106-117. doi:10.1016/j.nbd.2014.09.017

82. Li H, Khirug S, Cai C, et al. KCC2 Interacts with the Dendritic Cytoskeleton to Promote Spine Development. *Neuron*. 2007;56(6):1019-1033. doi:10.1016/j.neuron.2007.10.039

83. Martínez-Cerdeño V. Dendrite and spine modifications in autism and related neurodevelopmental disorders in patients and animal models. *Developmental Neurobiology*. 2017;77(4):393-404. doi:10.1002/dneu.22417

84. Chao HT, Chen H, Samaco RC, et al. Dysfunction in GABA signalling mediates autism-like stereotypies and Rett syndrome phenotypes. *Nature*. 2010;468(7321):263-269. doi:10.1038/nature09582

85. Lozovaya N, Nardou R, Tyzio R, et al. Early alterations in a mouse model of Rett syndrome: the GABA developmental shift is abolished at birth. *Sci Rep*. 2019;9(1):9276. doi:10.1038/s41598-019-45635-9

86. He Q, Nomura T, Xu J, Contractor A. The Developmental Switch in GABA Polarity Is Delayed in Fragile X Mice. *J Neurosci*. 2014;34(2):446-450. doi:10.1523/JNEUROSCI.4447-13.2014

87. Fadila S, Quinn S, Turchetti Maia A, et al. Convulsive seizures and some behavioral comorbidities are uncoupled in the Scn1aA1783V Dravet syndrome mouse model. *Epilepsia*. 2020;61(10):2289-2300. doi:10.1111/epi.16662

**Figures**

**Figure 1. KU rescues the altered GABA switch in *Scn1a+/-* mice.**

**A.** Representative traces of CA3 pyramidal cells' spontaneous spiking activity measured by current

clamp (I=0) recordings in loose patch configuration. Cells were recorded for 3/5 minutes to obtain

a stable control baseline, isoguvacine (10 μM) was bath applied for 3 minutes (bars above the traces)

and washed out for at least 10 minutes. **B.** Analysis of mean spike frequency in control and upon

isoguvacine application in WT or *Scn1a+/-* CA3 pyramidal neurons revealed an excitatory effect of

the GABA agonist in *Scn1a+/-* mice (paired t-test, p-value <0.05) whereas an inhibitory effect was

observed in WT mice (paired t-test, p-value <0.05). **C.** Schematic representation of KU55933 (KU)

treatment and experimental timeline. For comparing littermates in *ex-vivo* electrophysiological

recordings, we used sliding windows: mice received two intranasal administrations (every 72 hours)

of KU or vehicle (DMSO), the first in the window P12-P15 and the second in the window P15-18, and

recordings were performed in the window P18-P21. For western blotting analysis, mice received

two intranasal administrations (every 72 hours) at 15 and P18, and western blot was performed at

P21. **D.** Representative traces of P18-21 CA3 pyramidal cells' spiking activity and effect of

isoguvacine (10 µM; bar above the traces) measured as in A. **E.** Analysis of mean spike frequency of

baseline and upon isoguvacine application (bars above the traces) of $Scn1a^{+/-}$-DMSO and $Scn1a^{+/-}$-

KU CA3 pyramidal neurons revealed an excitatory effect of GABA in vehicle-treated $Scn1a^{+/-}$ mice

(paired t-test, p-value <0.01), whereas no significant modification was found in KU-treated animals

between baseline and isoguvacine (paired t-test, p-value=0.4). **F.** Western blotting quantification of

P21 hippocampal tissues showed a reduced expression of KCC2 chloride transporter in $Scn1a^{+/-}$

mice, which is related to GABA switch delay. KU treatment rescued this alteration increasing KCC2

expression. **G.** There was no difference in NKCC2 levels in the three conditions. See Supplementary

material for values, n and statistical tests.

**Figure 2. KU ex-vivo characterization in P18-21 *Scn1a*⁺/⁻ mice: effect on spontaneous synaptic activity.**

**A.** Representative traces of spontaneous GABAergic post-synaptic currents (sIPSCs) and spontaneous glutamatergic post-synaptic currents (sEPSCs) recorded at +10 mV and -60 mV, respectively, in pyramidal neurons from hippocampal CA1 slices of vehicle-treated (DMSO) or KU-treated *Scn1a*⁺/⁻ mice compared to WT littermates. **B.** The cumulative distribution showed an increased inter-event interval (IEI) of sIPSCs in vehicle-treated *Scn1a*⁺/⁻ mice compared to WT. KU treatment decreased sIPSCs IEI in *Scn1a*⁺/⁻ treated animals (KS test p-value <0.0001). Consistently, the comparison of mean sIPSCs frequency showed a decrease in vehicle-treated *Scn1a*⁺/⁻ mice compared to WT, which was partially rescued in KU-treated *Scn1a*⁺/⁻ mice. **C.** The cumulative distribution of sEPSCs IEIs displayed a decrease in vehicle-treated *Scn1a*⁺/⁻ mice and KU treatment rescued this defect, but increasing the IEI above WT level (KS test p-value <0.0001). Consistently, the comparison of mean sEPSCs frequency showed an increase in *Scn1a*⁺/⁻ compared to WT and KU

administration rescued this defect, but reducing the mean sEPSCs frequency below that of WT. **D.**

The excitatory/inhibitory ratio (E/I) was obtained dividing mean sEPSCs and sIPSCs frequencies

measured of the same neuron. **E.** The cumulative distribution of sIPSCs amplitude showed a slight

difference between the distributions (KS test p-value <0.0001), which did not reach statistical

significance comparing mean values. **F.** The cumulative distribution of sEPSCs amplitudes showed

an increase in vehicle-treated *Scn1a*$^{+/-}$ mice and a decrease in KU-treated ones compared to WT (KS

test p-value <0.0001). The decrease in vehicle-treated *Scn1a*$^{+/-}$ mice was not observed comparing

mean values, whereas KU treatment decreased sEPSCs amplitude below WT level also considering

mean values. See Supplementary material for values, n and statistical tests.

**Figure 3. KU and bumetanide treatments do not affect the epileptic phenotype of *Scn1a*⁺/⁻ mice.**

**A.** Representative diagrams of KU55933 (KU) and Bumetanide (BUM) treatments and experimental timelines. Mice received KU 7.5 mg/kg intranasally every 72 hours or bumetanide 2 mg/kg IP daily starting from P15. The effect on hyperthermia induced seizures were assessed at P21 after a 1-week treatment. The effect on spontaneous seizures was assessed in separated animal cohorts treated chronically for two weeks. **B, C.** Hyperthermia-induced seizures did not show any differences upon KU treatment, neither in the threshold temperature nor in the seizures' severity score. **D, E**. Similar

to KU, BUM did not affect neither the threshold temperature nor the severity score. **F-G.** Video-

analysis of spontaneous convulsive seizures showed no changes in KU-treated vs DMSO-treated

mice in seizures' frequency (F) or number of clusters of seizures (G). **H.** KU did not improve the

mortality rate (p-value = 0.9755 Kaplan-Meier). **I-J.** Similar to KU, Bumetanide treatment did not

improve seizures' frequency (I) or clusters of seizures (J). **K.** BUM treatment did not modify the

mortality rate (p-value = 0.2356; Kaplan-Meier). See Supplementary material for values, n and

statistical tests.

**Figure 4. KU and bumetanide treatments ameliorate sociability of *Scn1a^{+/-}* mice.**

**A.** Diagram of the experimental timeline to test effects of KU55933 (KU) and bumetanide (BUM) on

*Scn1a^{+/-}* autistic-like phenotype. Animals received KU treatment (two intranasal administrations, at

P15 and P18, spaced of 72 hours) or BUM treatment (one-week daily IP injections from P15). Three-

chamber sociability test and Y-maze were performed at P21. **B.** *Scn1a^{+/-}* male mice showed a defect

in social preference compared to WT at P21, as indicated by a lower sociability index, and KU

treatment rescued this alteration. **C.** The three-chamber test showed no defects in social interaction

in *Scn1a^{+/-}* females. **D.** Like KU, BUM treatment rescued defective sociability in *Scn1a^{+/-}* male mice.

See Supplementary material for values, n and statistical tests.

**Figure 5. Bumetanide, but not KU, improves Y-maze performance and decreases hyperactivity in**

*Scn1a<sup>+/-</sup>* **mice.**

**A.** The Y-maze test showed a defect in spontaneous alternation in *Scn1a<sup>+/-</sup>* male mice compared to

WT at P21. KU treatment did not show a clear rescue of this defect. **B.** KU-treated and control DMSO-

treated *Scn1a<sup>+/-</sup>* male mice displayed an increased number of entries in the maze during the test. **C.**

*Scn1a<sup>+/-</sup>* male mice showed increased total distance compared to WT, and KU partially ameliorated

this feature. **D.** The control group of *Scn1a*$^{+/-}$ male mice used for the bumetanide (BUM) treatment

showed, similarly to the DMSO-control ones used for testing KU, a defect in spontaneous alternation

compared to WT, which was rescued by BUM. **E.** Control *Scn1a*$^{+/-}$ male mice showed an increased

number of entries which was rescued by BUM.  **F.** Control *Scn1a*$^{+/-}$ male mice showed an increased

distance travelled, which was rescued by BUM. **G.** WT-DMSO, *Scn1a*$^{+/-}$- DMSO and *Scn1a*$^{+/-}$- KU

female mice did not show differences in spontaneous alternation in the Y-maze test. **H.** Female

*Scn1a*$^{+/-}$ mice, as male ones, showed an increased number of entries in the Y-maze, with a trend

towards rescue upon KU treatment. **I.** Female *Scn1a*$^{+/-}$ mice, as male ones, showed increased total

distance travelled during the Y-maze test, with a trend towards rescue upon KU treatment. **J.** As in

the KU experimental cohort, Y-maze test showed no differences in Y-maze performance in WT,

*Scn1a*$^{+/-}$- CTRL and *Scn1a*$^{+/-}$- CTRL female mice. **K.** BUM rescued the increased number of entries

showed by *Scn1a*$^{+/-}$ mice during the Y-maze test. **L.** The increased total distance travelled by *Scn1a*$^{+/-}$

mice during the Y-maze test was partially reduced by BUM, but it was still different from WT. See

Supplementary material for values, n and statistical tests.

**Figure 6. Early neurodevelopmental characterization of *Scn1a^+/-* mice.**

**A.** Righting reflex analysis (cut-off 30 s) did not show any differences between WT and *Scn1a^+/-* mice

at P7 or P10-11. **B.** Grasping reflex test (forepaws) did not show any differences between WT and

*Scn1a^+/-* mice at P7 and P10-11. **C.** The negative geotaxis test at P10-11 gave similar results for

*Scn1a^+/-* and WT mice. **D.** The gait test showed no differences between *Scn1a^+/-* and WT mice at P10-

11. **E.** *Scn1a^+/-* mice showed dysfunctions in the nest bedding test (olfactory motivation) compared

with WT at P10-11. Three trials for each animal were performed and considered successful if the

animal reached the nest bedding in less than 60 s. The time needed to reach the nest was similar

for the two genotypes. **F.** Eye opening is delayed in *Scn1a^+/-* compared with WT mice. See

Supplementary material for values, n and statistical tests.

**Figure 7. *Scn1a⁺/⁻* mice show defects in spontaneous synaptic activity before seizures appearance.**

**A.** Representative traces of spontaneous GABAergic post-synaptic currents (sIPSCs) and

glutamatergic post-synaptic currents (sEPSCs) recorded in P10-12 hippocampal CA1 pyramidal

neurons clamped at +10 mV and -60 mV, respectively. **B.** The cumulative distribution of sIPSCs inter

event interval (IEI) revealed an increase in *Scn1a⁺/⁻* mice compared to WT, KS test p-value <0.0001.

Consistently, the analysis of mean synaptic events showed a decreased frequency of sIPSCs in

*Scn1a⁺/⁻* mice. **C.** Cumulative distribution of sEPSCs IEI revealed no changes in *Scn1a⁺/⁻* compared to

WT mice, KS test p-value = 0.3141. Similarly, there were no changes in sEPSCs mean frequency in

*Scn1a⁺/⁻* mice. **D.** The excitatory/inhibitory ratio (E/I) has been obtained dividing sEPSCs and sIPSCs

frequencies measured in the same neuron. **E.** The cumulative distribution of sIPSCs amplitudes

revealed a slight decrease in *Scn1a⁺/⁻* mice, KS test p-value <0.0001, which however did not reach

statistical significance comparing mean values. **F.** Cumulative distribution of sEPSCs amplitude

revealed a decrease in *Scn1a⁺/⁻* compared to WT mice, KS test p-value <0.0001, which did not reach

1    statistical significance comparing mean values. See Supplementary material for values, n and

2    statistical tests.

**Figure 8. *Scn1a^+/-* mice show defects in spine maturation.**

**A.** Representation of dendritic spines in CA1 pyramidal cell and illustration of how the morphometric

parameter "head width" was quantified (black segments in the spine heads). **B.** Representative

images of dendritic spines of pyramidal neurons in the CA1 area of the hippocampus of P11 (upper

panels) and P21 (lower panels) in WT and *Scn1a^+/-* mice (Golgi staining). **C(a).** No change in spine

density between WT and *Scn1a^+/-* P11 mice. **C(b).** Decreased spine head width in *Scn1a^+/-* mice at

P11. WT and *Scn1a^+/-*. **D(a).** No change in spine density in P21 *Scn1a^+/-* mice compared with WT.

**D(b).** Decreased spine head width in *Scn1a^+/-* mice at P21. WT and *Scn1a^+/-*. See Supplementary

material for values, n and statistical tests.