TITLE:

Effect of 10 days of unilateral lower limb suspension on knee extensors neuromuscular function and spinal excitability

SHORT TITLE:

Muscle disuse and neuromuscular function

AUTHORS:

Loïc Lebesque[1], Marco V. Narici[2,3], Alain Martin[1], Giuseppe De Vito[2,3], Fabio Sarto[2] and Gil Scaglioni[1]

[1] INSERM UMR1093-CAPS, Université Bourgogne Franche-Comté, UFR des Sciences du Sport, F-21000, Dijon.

[2] Department of Biomedical Sciences, University of Padova, Padova, 35131, Italy.

[3] CIR-MYO Myology Center, University of Padua, 35131, Italy.

CORRESPONDING AUTHOR: Address for reprint requests and other correspondence:

Gil Scaglioni

INSERM UMR1093 "Cognition, Action et Plasticité Sensorimotrice »

Université de Bourgogne - UFR STAPS

Campus Universitaire, BP 27887 DIJON

Phone: +33 3 80 39 67 58

E-mail : gil.scaglioni@u-bourgogne.fr

27   Keywords : Force control; Disuse; Atrophy; H-reflex; T-reflex; Unloading; Vastus

28   medialis

ABSTRACT

Introduction. The reduction in mechanical loading applied on the lower limb has numerous detrimental consequences on neuromuscular function. While muscle mass and strength are known to be altered, changes in motoneuronal function remain poorly investigated.

Methods. Ten young healthy males (19-28 years old) underwent 10 days of unilateral lower limb suspension (ULLS) to simulate muscle disuse. Modulation by unloading of knee extensors' function (muscle morphology and strength, activation capacity and contractile properties) and spinal reflexes were explored before and after the ULLS.

Results. The knee extensors' cross-sectional area (-4%, $p = 0.007$), maximal strength (-27%, $p < 0.001$) and activation capacity (-3%, $p = 0.006$) were reduced after 10 days of ULLS. Vastus medialis H-reflex amplitude was enhanced both at rest (+33%, $p = 0.038$) and during a low-intensity contraction set at 10% of maximal strength (+103%, $p = 0.038$). No change in nerve conduction velocity was observed after the ULLS.

Conclusion. The present study suggests that both muscular and neural impairments may contribute to the de crease in knee extensors' function induced by short-term ULLS. The decrease in muscle activation after a short period of ULLS was accompanied by an increase in spinal excitability. However, the latter adaptation was not sufficient to counteract the reduced activation capacity resulting from muscle disuse. These adaptations to short-term ULLS should be considered when aiming at improving the neuromuscular function of people experiencing muscle hypoactivity.

INTRODUCTION

Segmental limb reduction in mechanical loading may occur as a consequence of surgery, injury and illness (1–3), but also as a result of reduced daily physical activity (4,5) or gravitational input, as during spaceflight (6). The direct outcome is a loss of muscle mass and function, associated with changes in motor control (7,8).

Several models of immobilization allow to explore the neuromuscular impairment in response to disuse (8). However, unilateral lower limb suspension (ULLS) enables to restrain atrophy to the explored musculoskeletal segment, easily mimicking the joint unloading that follows injuries or surgery (7,9) and, compared to casting, has the advantage of being less binding.

Extensive literature shows that the extent of muscle weakness is greater than the decrease in muscle mass since the first few days of ULLS, meaning that neuromuscular function is rapidly impaired by disuse (7,9–12). Among the antigravity muscles, the knee extensors appear to be the most affected by ULLS (13–15), showing a strength loss considerably larger than that of the plantar flexors (-1.0% · day^{-1} *vs.* 0,7% · day^{-1}, for durations between 10 and 35 days of ULLS; 8,11). Therefore, although it is reasonable to assume that for unloading, as for overloading, changes in neural drive are the first to occur (8), their involvement in the weakening of antigravity muscles, associated with disuse, is still an unresolved issue. Although studies have shown that ULLS (21-30 days) does not interfere with the central motor drive of knee extensors muscles (9,11,15,16), others have found a decrease (17,18) or even an increase (range + 0.1%.day^{-1} increase to – 0.2%.day^{-1} decrease; median – 0.2%.day^{-1}) of muscle voluntary activation (8). This divergence was also observed

in surface electromyographic (EMG) data, recorded during maximal voluntary

contractions, with studies showing a reduction of the vastii activity after ULLS (7-42

days) (19–23), and others finding no change for the vastii and RF (ULLS 7-35 days)

(11,22,24). On the other hand, the literature agrees on the EMG data recorded during

submaximal contractions. Interestingly, in this case, all studies showed that vastii

EMG activity increased following ULLS (7,23–25), suggesting that additional motor

units are necessary to produce the same absolute force output after unloading. In

light of these considerations, it appears that changes at neural level have not yet

been fully understood, and neither have the underlying mechanisms. Previous

studies, on plantar flexors, have shown an unequivocal enhancement in the

amplitude of the Hoffman (H)-reflex (the EMG response evoked by electrically

stimulating the Ia fibers of a peripheral nerve and considered as a comprehensive

index of spinal excitability) in response to unloading (7,8,12–14,26–30). The

enhancement of H-reflex amplitude with unloading may be ascribed to an increased

synaptic efficiency and/or to a greater motoneuronal excitability (31), however, the

improved synaptic efficiency seems to be the predominant adaptation in all forms of

disuse (13). This mechanism may interfere with the reduction in motoneuronal

excitability that might occur together with an increased proportion of fast fibers in

response to unloading (27). These adjustments should equally affect H and T-

reflexes. The T-reflex is an electrophysiological variant of the stretch reflex caused by

a mechanical tap applied on the tendon that, generating a quick stretch of the tendon

itself, triggers muscle spindles discharge to Ia afferents. Both reflexes originate from

the monosynaptic projections of the Ia afferences to alpha-motor neurones, but the T-

reflex involves all the elements of the reflex pathway, including muscle spindles

sensitivity (32,33). The literature shows that the T-reflex may be further modulated by

conflicting changes arising in peripheral components of the stretch reflex pathway.

Actually, Anderson *et al.* (27) found, after 3 weeks of hind limb suspension in rat, a

decline of the maximal soleus T-reflex but a rise in the maximal H-reflex (H_{MAX}).

Seynnes *et al.* (29) observed no change in soleus T-reflex, following 23 days of

ULLS, but an increase in H-reflex and a decrease in tendon stiffness. The different

modulations of the T-reflex response with unloading may be ascribed i) to a lower

spindles' sensitivity, due to their reduced solicitation during immobilization (34),

and/or ii) to a decreased stiffness of the musculo-tendinous elements in series with

the muscle spindles (27,29).

In view of these considerations and since the literature seems to show that few

days of immobilization induce significant changes in neuromuscular function (8), this

study sought to investigate the role of changes in spinal reflex excitability, muscle

contractile characteristics and muscle morphology in the loss of knee extensors'

strength following 10 days of muscle unloading. Since, in plantar flexors, H-reflex is

always modulated by ULLS and considering that knee extensors are among the

antigravity muscles most affected by inactivity, we hypothesized that both spinal

excitability and muscle morphology may be altered by unloading. However, given the

short period of immobilization, it could also be assumed that neural adaptations might

be preponderant over muscular changes. Therefore, apart from improving knowledge

on mechanisms responsible for the decline of the neuromuscular function due to

unloading, the outcomes of the present investigation may be useful for refining the

rehabilitation strategies of patients following periods of muscle disuse.

MATERIALS AND METHODS

Ethical approval.

The present study was approved by the Ethics Committee of the Department of Biomedical Sciences of the University of Padova (Italy) (reference number HEC-DSB/01-18) and conducted in accordance with the standards set by the latest revision of the Declaration of Helsinki. Participants were informed about all experimental procedures through an interview and an information sheet, and could ask questions to the selection committee. After signing the written consent form, participants were enrolled in the study with the possibility to withdraw at any stage of the experiment. This study was part of a larger investigation aiming to study the neuromuscular alterations with short-term disuse (18,35).

Participants

Ten healthy and recreationally active young male volunteers (age: 22.9 ± 4.0 years old; body mass: 72.2 ± 7.2 kg; height: 177.6 ± 3.1 cm; body mass index: 22.9 ± 2.1 $kg \cdot m^{-2}$) took part to this study. Because unilateral lower limb suspension (ULLS) is associated with an increased risk of deep venous thrombosis (36), more common in females, only males were recruited in this study. Inclusion criteria were: age comprised between 18 and 35 years, body mass index comprised between 20 and 28 $kg \cdot m^{-2}$ and involvement in recreational physical activities 1-3 times per week.

Exclusion criteria were: sedentary lifestyle, professional sport practice, smoking,

history of deep venous thrombosis and acute or chronic musculoskeletal, metabolic

and cardiovascular disorders. Using the Global Physical Activity Questionnaire (37),

participants demonstrated a high level of physical activity (3268 ± 1468

MET·min·week^{-1}) in the period leading up to the ULLS experiment. Volunteers were

asked to maintain their habitual diet throughout the intervention period and refrain

from intense exercise, coffee and alcohol intake during the 24 hours preceding the

data collection at each time point.

Protocol

Experimental design. Data were collected the day before (LS0) and after 10 days

(LS10) of unilateral lower limb suspension (ULLS). For each subject, sessions lasted

2 hours and were planned at the same time of the day to avoid any circadian

disturbance (38,39). At both LS0 and LS10, the experimental procedure was

performed beginning with the evaluation of the maximal voluntary isometric

contraction (MVC) of knee extensors and the maximal voluntary activation level

(VAL). Then, vastus medialis (VM) spinal excitability parameters were assessed

through electrical stimulations and patella tendon striking at rest and during a weak

contraction of knee extensors (10% MVC assessed for each session). In addition,

quadriceps and VM morphology was assessed through ultrasonography, on the

same day of the ULLS beginning (baseline data collection, BDC) and the day before

the end of the ULLS (LS9).

Maximal force production capacity. To start the maximal force production evaluation,

subjects performed a knee extensors warm-up composed of 10 submaximal

contractions, lasting 4 s each. Then, three knee extensors MVCs (lasting 4 s, with 1-

min rest in-between) were performed with maximal electrical doublet (100 Hz)

delivered over the peak force.

Spinal excitability.   Since it is more challenging to evoke rest H-reflex in the

quadriceps muscles compared to the soleus muscle, several studies suggest eliciting

it during a weak muscle contraction (40–43). The reflex response is indeed enhanced

by the muscle contraction because motoneurons are brought closer to their firing

threshold by the voluntary drive (44). In addition, muscle activation allows to keep

constant the "background" level of α-motoneuron depolarisation and limits post-

synaptic modulations (41,45). Therefore, the assessment of spinal excitability during

muscle contraction has the advantage of increasing amplitude and reducing

variability in both latency and amplitude of the H-reflex (44,46,47). Consequently, VM

H-reflex and M-wave recruitment curves were built by applying electrical stimulations

first during weak knee extensors contractions (10% MVC) and then at rest. To build

the recruitment curves, stimulations started at 5 mA and were progressively

increased with a 5-mA increment until the maximal M-wave (M_{MAX}) was reached. To

ensure that the M-wave lied in the plateau of its maximal value throughout the

experiment, the maximal stimulation intensity was increased by 20%. Subsequently,

additional stimulations were applied, in 2 mA increments, around the intensity that

allowed to record the highest H-reflex, in order to accurately estimate the intensity

needed to elicit the maximal H reflex (H_{MAX}). Four stimuli were delivered at each

intensity interspaced by 10 s intervals. Then, 6 out of 10 subjects underwent

complementary evaluation of spinal motoneuron excitability at rest through the

assessment of tendinous reflex (i.e., T-reflex, for the method see below mechanical

stimulation).

Unilateral lower limb suspension. The ULLS model was employed for 10 days. The

non-dominant leg (left leg for all participants) of the participants was fitted with a shoe

having an elevated sole (50 mm), while the dominant leg (right leg for all participants)

was suspended and kept at a slightly flexed position (approximately 15-20° of knee

flexion) using straps. Volunteers were asked to walk on crutches during the whole

ULLS period and to refrain from loading the suspended leg in any way. A

familiarization session was set up for the participants to practice daily tasks with the

limb suspended (9). Participants were recommended to wear elastic compression

socks on the suspended leg during ULLS period and to perform passive, range of

motion, non-weight-bearing exercises of the ankle, as precautionary measures to

prevent deep venous thrombosis (36). Moreover, we performed an ultrasound-

Doppler examination after 5 days of ULLS. Participants' compliance was evaluated

through daily calls and messages and by comparing calves' temperature and

circumference after 5 and 10 days of ULLS, as previously suggested (9).

Data acquisition.

Ultrasonography. To evaluate morphology of quadriceps and VM muscles, subjects

were lying on their backs on an auscultation table. Muscles cross-sectional area

(CSA) was obtained using extended-field-of-view ultrasonography imaging (Mylab,

Esaote, Genoa, Italy). A 47 mm, 7.5 MHz linear array transducer was used to collect

images at different muscle length percentages. Three regions of interest were

detected at 30%, 50% and 70% of femur length (measured as the distance between

the great trochanter and the mid-patellar point), where 0% represents the mid-patellar

point (distal part) and 100% the greater trochanter (proximal part). An adjustable

guide was used in each acquisition to keep the same transverse path (48). After

applying a generous amount of transmission gel on the participant's skin to improve

the acoustic contact, the transducer was moved slowly in the transverse plane from

the medial border of the VM to the lateral border of the vastus lateralis. During the

transducer displacement, we kept the pressure on the skin as constant as possible.

Two scans were obtained for each site.

Force recordings. Voluntary and electrically-evoked force of knee extensors was

recorded using a custom-made knee dynamometer fitted with a strain gauge. The hip

and knee angles were set at 90° (180° full extension). Subject's dominant leg was

securely strapped to the custom-made knee dynamometer, equipped with a load cell,

at the ankle level. Subject was asked to cross the arms on the chest and keep the

head in a neutral position during the experiment procedure. The force exerted by the

subject was displayed in real-time on a computer screen placed at 1 m in front of him.

Because spinal excitability was further assessed during a weak quadriceps

contraction, a line representing 10% of the session MVC was added on the computer

screen once the MVC was determined. Force data were were recorded at a sampling

frequency of 1 kHz with the Spike2 software (version 8.01, Cambridge Electronics

Design Ltd, Cambridge, UK; RRID: SCR_000903) and stored for offline analysis with

the LabChart software (version 8, AD Instruments®, Australia; RRID: SCR_017551).

Electromyographic recordings. Electromyographic activity (EMG) from the dominant-

leg VM was recorded during the 10% MVC contractions with bipolar active-type Ag-

AgCl electrodes electrodes (44 x 22 mm, Ambu®). To minimize skin impedance (< 5

kΩ), the skin was shaved, rubbed with an abrasive paste and cleaned with alcohol,

before electrode positioning. Electrodes were positioned 10 cm above the patella

upper part and 4 cm medially with an inter-electrode center-to-center distance of 20

mm. The EMG signals were amplified (gain = 1,000) and bandpass filtered at 5 Hz to

5 kHz using a CED 1902 amplifier (Cambridge Electronics Design Ltd, Cambridge,

UK) and digitized with a CED Micro 1401 data acquisition unit (Cambridge

Electronics Design Ltd, Cambridge, UK). EMG data were recorded at a sampling

frequency of 40 kHz with the Spike2 software (version 8.01, Cambridge Electronics

Design Ltd, Cambridge, UK; RRID: SCR_000903) and stored for offline analysis with

the LabChart software (version 8, AD Instruments®, Australia; RRID: SCR_017551).

Nerve electrical stimulations. Nerve electrical stimulations were delivered using a

high-voltage (400 V) constant-current stimulator (DS7AH model, Digitimer,

Hertfordshire, UK). For maximal voluntary activation assessment, electrical doublets

(100 Hz, monophasic rectangular pulses, 1 ms width) were delivered through two

stimulation electrodes (5 x 10 cm, Compex SA, Ecublens, Switzerland) placed in the

proximal and distal region of the knee extensors. Maximal electrical stimulation

intensity was determined as the intensity eliciting the maximal twitch amplitude. For

spinal excitability assessment, we delivered percutaneous electrical stimulations

(monophasic rectangular pulses, 1 ms width) in the femoral trigone over the femoral

nerve to evoke electrophysiological and mechanical responses of the VM muscle.

The self-adhesive anode (5 x 10 cm, Compex SA, Ecublens, Switzerland) was placed

in the gluteal fossa. The optimum stimulation site was located with a hand-held

cathode stylus in order to obtain the greatest H-reflex amplitude of the VM for the

lowest stimulation intensity. Then, the self-adhesive cathode electrode (7 mm

diameter, Contrôle Graphique Medical, Brie-Compte-Robert, France) was positioned

and firmly fixed with a velcro elastic band.

Mecanichal stimulations. During T-reflex procedure, subjects were positioned as

described previously in *Force recordings* section. Twenty T-reflexes were elicited for

each subject by tapping the anterior face of the patellar tendon of the quadriceps with

a manual reflex hammer (neurological hammer, Babinsky model) at various

intensities. The T-reflex on the VM EMG signal and its associated mechanical twitch

were recorded for each strike.

Data analysis.

Ultrasonography. Of the two scans obtained at each site, we analyzed the image with

the best quality. Quadriceps and VM CSAs were measured by tracing the muscle

contours with the image analysis software "ImageJ" (https://imagej.net; RRID:

SCR_003070). Two measurements per parameter were performed on each image,

and their average was considered. Quadriceps CSA (qCSA) was computed as the

average value of the CSAs recorded at 30, 50 and 70% of the femur length for each

subject at both LS0 and LS10. VM CSA was calculated as the average value of the

CSAs recorded at 30 and 50% of the femur length, due to technical issues in clearly

detecting the border between the VM and vastus intermedius at 70% of the femur

length.

Maximal force production. The MVC was determined as the highest force value

reached during the 3 MVCs. The target of 10% MVC was calculated from the MVC of

each session (i.e., LS0 or LS10). The specific force (i.e., maximal force by surface

unit) was calculated as the MVC divided by the quadriceps CSA (MVC/qCSA).

Activation capacity. The voluntary activation associated with the maximal force

production and the weak contraction (10% MVC) was quantified with the central

activation ratio (CAR_{MVC} and $CAR_{10\%}$ respectively) as follows (49):

$$CAR = \frac{Torque}{Torque + superimposed\ twitch}$$

where force was either the MVC (CAR_{MVC}) or the 10% MVC ($CAR_{10\%}$). It should be

noted that, while the CAR_{MVC} was calculated using maximal doublet stimulations (100

Hz) delivered during brief MVC, the $CAR_{10\%}$ was derived from the M-wave

recruitment curve recorded during weak contractions at 10% MVC using maximal

single stimulations (see above *Spinal excitability* section). The ratio between the

$CAR_{10\%}$ and the force produced during the contraction at 10% MVC was calculated.

The root mean square (RMS) of VM EMG activity was computed offline over a 500

ms period preceding H_{MAX} and M_{MAX} stimulations during the weak contractions at

10% MVC, and subsequently normalized to the M_{MAX} amplitude (RMS/M_{MAX}).

Spinal excitability parameters. One subject has been exempted from the spinal

excitability assessment because he did not tolerate electrical stimulations over the

femoral nerve. Peak-to-peak amplitude of both H-reflex and M-wave was measured

and averaged for each stimulation intensity. The highest amplitude of M-wave and H-

reflex were labelled H_{MAX} and M_{MAX} respectively. The H_{MAX}/M_{MAX} ratio was then

calculated. In one subject it was not possible to evoke the H-reflex at rest and was

excluded from H_{MAX}-related parameters analysis at rest. The amplitude of the M-

wave accompanying H_{MAX} (M_{atHMAX}) was also measured and normalized by M_{MAX} to

verify that stimulation conditions were similar between sessions. The RMS of VM

EMG signal was measured over the 500 ms preceding each H_{MAX} (RMS_H) and M_{MAX}

(RMS_M) electrical stimulation during the contraction. We calculated the H_{MAX}/RMS_H

ratio to assess the influence of muscle activity background on spinal excitability (50).

We also measured H_{MAX} and M_{MAX} latencies at rest ($H_{LATENCY}$ and $M_{LATENCY}$) as the

delay between the onset of stimulation artefact and the upshot of electrophysiological

response. While $M_{LATENCY}$ represents the time required for conduction down the

motor units to the terminal branches and propagation across the neuromuscular

junction, $H_{LATENCY}$ reflects the time required for signal propagation through the reflex

arc, including the Ia afferent synaptic delay at the spinal level. Moreover, the

difference between $H_{LATENCY}$ and $M_{LATENCY}$ (H-$M_{LATENCY}$), representing an index of

nerve conduction velocity through the spinal reflex loop, was computed (13). Then,

these latencies were normalized by the subject's femur length (i.e., the distance

between the greater trochanter and the lateral condyle). For one subject, the onset of

the VM H_{MAX} was merged with the associated M-wave, thus this subject was

excluded from H_{MAX} latency analysis. The amplitude of the mechanical twitch

associated with both M_{MAX} (M_{TW}) and H_{MAX} at rest was measured. Since the M_{TW}

represents the mechanical response of the muscle when all the motor units are

recruited, its characteristics were analyzed. For each M_{TW}, we determined the

contraction time (i.e., the time between twitch onset and maximal twitch value; CT),

the rate of force development (i.e., the maximal value of the first derivative of the

force signal using a 1 ms sliding window; RFD) and the half relaxation time (i.e.,

corresponding to the time needed to reach a 50% decrease of twitch force; HRT)

before computing the averages of the four measurements. To assess the

electromechanical efficiency, the M_{TW}/M_{MAX} was calculated. We analyzed the

relationship between the amplitude of the T-reflex and the corresponding twitch since

we did not have the complete T-reflex recruitment curve and therefore were not

certain of obtaining the maximal T-wave response. The relationship between VM T-

reflex amplitudes, expressed as a percentage of M_{MAX}, and the associated twitch

amplitudes were scatter-plotted individually. The slope and the intercept of the linear

regression of these relationships were determined for each subject and averaged at

LS0 and LS10.

Statistical analysis. The normality of each dataset was assessed with the Shapiro-

Wilk test. For all variables that passed normality test, Student's paired t-test was

applied in order to investigate whether difference between LS0 and LS10 was

present within the considered parameters. When normality test was not passed, non-

parametric Wilcoxon's signed-rank test for paired comparison was used. Only for the

RMS/M_{MAX} variable, a two-way repeated measures ANOVA with *time* (LS0 *vs.* LS10)

and *stimulation* (H_{MAX} *vs.* M_{MAX}) factors was performed. If a significant effect was

found, a Tukey's post-hoc analysis was conducted. Correlations between MVC and

qCSA as well as between MVC and CAR_{MVC} were analyzed using repeated

measures correlations (rmcorr R package, 51). Effect sizes (ES) were calculated with

Cohen's d and r respectively for parametric and non-parametric comparison (52).

Small, moderate and large effects are considered for Cohen's d ≥ 0.2, ≥ 0.5, ≥ 0.8,

and Cohen's r ≥ 0.1, ≥ 0.3, ≥ 0.5 respectively. Results are reported as mean ±

standard deviation. Significance level was set at $p < 0.05$ and the confidence interval

at 95% ($CI_{95\%}$). Student's and Wilcoxon's tests were performed using Statistica

software (Statsoft, version 12, Tulsa, OK, USA; RRID: SCR_014213) while Cohen's

effect sizes were calculated with G*Power software (version 3.1.9.2, Universität

Düsseldorf, Germany; RRID: SCR_013726). Repeated measures correlations were

performed with R (version 4.1.3, R Core Team, 2017; RRID: SCR_001905). Graphs

were made with GraphPad Prism software (version 8.4.0, GraphPad software Inc.

San Diego, CA, USA; RRID: SCR_002798).

RESULTS

Muscle morphology, strength and activation capacity.

The CSA of the unloaded VM ($t_{(9)}$ = 5.259, $CI_{95\%}$ = [0.86 : 2.17], p = 0.0005,

ES = 0.619) and quadriceps ($t_{(9)}$ = 3.444, $CI_{95\%}$ = [0.87: 4.19], p = 0.007, ES = 0.397)

decreased (-7.8 and -4.4 %, respectively) after 10 days of ULLS (Table 1). The CSA

losses were accompanied by a 27.1 % reduction in knee extensors MVC ($t_{(9)}$ = 9.464,

$CI_{95\%}$ = [17.01 : 27.70], p < 0.0001, ES = 2.196) (Table 1). The specific force

(MVC/qCSA) was 23.7 % lower after ULLS ($t_{(9)}$ = 8.792, $CI_{95\%}$ = [0.25 : 0.43], p =

0.0001, ES = 2.016) (Table 1).

Activation capacity as measured by CAR_{MVC} decreased by 3.0 % after ULLS

($t_{(9)}$ = 3.557, $CI_{95\%}$ = [0.01 : 0.05], p = 0.006, ES = 1.175) (Figure 1) while the $CAR_{10\%}$

decreased by 15.0 % ($t_{(9)}$ = 4.011, $CI_{95\%}$ = [2.79 : 10.03], p = 0.003, ES = 0.997). The

ratio between the CAR_{MVC} and the MVC increases by 33.4 % following ULLS ($t_{(9)}$ = -

6.839, $CI_{95\%}$ = [-0.01 : -0.00], p < 0.0001, ES = 1.880), while the ratio between the

$CAR_{10\%}$ and the force produced at 10% of MVC increased by 16.5 % ($t_{(9)}$ = -2.540,

$CI_{95\%}$ = [-0.02 : -0.00], p = 0.032, ES = 0.869) (Table 1). Furthermore, the normalized

EMG activity (RMS/M_{MAX}) of the VM during the 10% MVC plateau declined by 34.2%

after ULLS (only *time* effect; $F_{(1,8)}$ = 8.482, p = 0.020, η_p^2 = 0.515 ; Tukey post-hoc : p

= 0.020, ES = 0.839) (Table 1). Based on repeated measures correlation analysis, it

appears that MVC is positively correlated to qCSA (r_{rm} (9) = 0.773, $CI_{95\%}$ = [0.22 :

0.95], p = 0.005) (Figure 2A) and CAR_{MVC} (r_{rm} (9) = 0.624, $CI_{95\%}$ = [-0.07 : 0.91], p =

0.040) (Figure 2B). In other words, the higher the qCSA or CAR_{MVC}, the greater the

production of maximal force.

Muscle Contractile Properties

Mechanical parameters associated with H_{MAX} and M_{MAX} stimulations at rest

were also compared between LS0 and LS10. The ULLS did not modify the twitch

amplitude associated with M_{MAX} (M_{TW}; $t_{(8)}$ = 1.593, $CI_{95\%}$ = [-0.69 : 3.75], p = 0.150,

ES = 0.599) (Figure 3). Specifically, the RFD of M_{TW} was not altered after 10 days of

ULLS ($t_{(8)}$ = 1.685, $CI_{95\%}$ = [-0.01 : 0.09], p = 0.130, ES = 0.678), nor were the CT ($t_{(8)}$

= -1.159, $CI_{95\%}$ = [-14.61 : 4.84], p = 0.280, ES = 0.420) and the HRT ($t_{(8)}$ = 0.165,

$CI_{95\%}$ = [-15.05 : 17.36], p = 0.873, ES = 0.070). Also, the M_{TW}/M_{MAX} ratio was not

changed by the ULLS protocol (Z = 0.770, p = 0.441, ES = 0.257). Muscle contractile

properties obtained at rest are provided in Table 2. Regarding the twitch associated

with H_{MAX}, its amplitude was not different between LS0 and LS10 ($t_{(7)}$ = 1.015, $CI_{95\%}$ =

[-0.84 : 2.11], p = 0.344, ES = 0.253) (Figure 3).

H-Reflex responses and M-wave

While the H_{MAX} at rest was increased by 59.0% after 10 days of ULLS (Z =

2.100, p = 0.036, ES = 0.743), H_{MAX} obtained during muscle contractions at 10%

MVC was not modified (Z = 0.770, p = 0.441, ES = 0.257) (Table 3). It should be

noted that the amplitude of the M-wave associated with H_{MAX} (M_{atHMAX}) was not

different between LS0 and LS10 either at rest (Z = 0.700, p = 0.484, ES = 0.247) or

during contraction (Z = 1.362, p = 0.173, ES = 0.454) (Table 3). Therefore, the

stimulation conditions at H_{MAX} may be considered stable, allowing the comparison of

spinal excitability features between the two time points. M_{MAX} did not differ between

LS0 and LS10 either at rest ($t_{(8)}$ = -0.258, $CI_{95\%}$ = [-1.83 : 1.46], p = 0.803, ES =

0.025) or during 10% MVC contraction ($t_{(8)}$ = 1.345, $CI_{95\%}$ = [-0.42 : 1.59], p = 0.215,

ES = 0.074) (Table 3). The ULLS increased the H_{MAX}/M_{MAX} at rest by 33.1 % (LS0 :

0.20 ± 0.15 *vs.* LS10 : 0.27 ± 0.15 ; $t_{(7)}$ = -2.557, $CI_{95\%}$ = [-0.13 : -0.01], p = 0.038, ES

= 0.441) but had no impact on the ratio during contraction at 10% MVC (LS0 : 0.34 ±

0.16 *vs.* LS10: 0.36 ± 0.17; $t_{(8)}$ = -1.947, $CI_{95\%}$ = [-0.04 : 0.00], p = 0.087, ES = 0.121)

(Figure 4A). This result is also depicted in Figure 4B where experimental traces of

M_{MAX} and H_{MAX} obtained at rest, before and after ULLS, from one representative

subject, are presented. In order to overcome differences between sessions in EMG

activity during contractions at 10% MVC, we also normalized H_{MAX} by muscle activity

background preceding H_{MAX} stimulations (RMS_H), and we observed an increase of

this ratio after the ULLS period (+102.9 % ; $t_{(8)}$ = -2.480, $CI_{95\%}$ = [-74.00 : -2.70], p =

0.038, ES = 1.097) (Table 3 and Figure 4C).

Moreover, the delay between electrical stimulation and electrophysiological

responses was determined for the M_{MAX} and H_{MAX} waves. The latency of M_{MAX}

($M_{LATENCY}$) and H_{MAX} ($H_{LATENCY}$) was not affected by ULLS ($M_{LATENCY}$: $t_{(8)}$ = -0.527,

$CI_{95\%}$ = [-1.75 : 1.10], p = 0,613, ES = 0.192; $H_{LATENCY}$: $t_{(6)}$ = -0.913, $CI_{95\%}$ = [-2.53 :

1.15], p = 0,396, ES = 0.134) (Table 3). The difference between $H_{LATENCY}$ and

$M_{LATENCY}$ (H-$M_{LATENCY}$) was not changed after 10 days of ULLS ($t_{(6)}$ = 0.289, $CI_{95\%}$ = [-

1.73 : 2.20], p = 0.782, ES = 0.049).

In addition, the ULLS did not modulate the relationship between the VM T-

reflex and the associated twitch. Indeed, the unloading period did not alter the slope

($t_{(5)}$ = -0.874, $CI_{95\%}$ = [-0.31 : 0.15], p = 0.422, ES = 0.381) and the intercept ($t_{(5)}$ = -

1.397, $CI_{95\%}$ = [-1.91 : 0.57], p = 0.221, ES = 0.946) of this linear regression (Table

3).

DISCUSSION

The present study aimed to investigate neural, contractile and morphological

changes associated with muscle weakness following 10 days of ULLS of the main

antigravity muscle group of the lower limbs. We showed that knee extensors' disuse

led to a reduction in force production capacity explained both by muscle atrophy and

by an impairment in central motor drive. For the first time, spinal excitability was

investigated in the quadriceps muscle group to deepen knowledge of neural

adaptation to unloading in antigravity muscles. Our main results show that spinal

excitability was enhanced at rest and during muscle contraction following 10 days of

ULLS. These findings seem particularly relevant as they provide new insights into the

neurophysiological adjustments that occur in human muscle disuse.

Muscle morphology and function.

The present investigation shows that 10 days of ULLS reduced the maximal

force developed by the knee extensors (-27 %, -2.7 %·day^{-1}). This result is consistent

with previous investigations that reported an impairment in force production capacity

of the knee extensors following lower limb unloading (11,14–16,23,25,53), even

though the rate of decrease we observed was greater than in the aforementioned

studies (-0.5 to -1.3 %·day^{-1}). The faster rate of force loss observed in the present

study may be related to particular efforts made to maximise adherence to the disuse

protocol (see *Materials and Methods* section). The alteration in MVC following

unloading was associated with an overall reduction in qCSA of 0.44 %·day^{-1} (-4.4 %

in 10 days). This result is in accordance with previous studies that reported a

decrease of 0.3 to 0.5 %·day^{-1} in knee extensors size after ULLS (8,11,16,23). As

reported in the literature, muscle atrophy could be explained by decreases in protein

synthesis rate after lower limb muscle disuse (54,55) affecting the muscle protein

synthesis/breakdown balance and thus, the myofibrillar content (56). The present
study corroborated, in agreement with earlier investigations (8,16,57–59), that
strength loss, after unloading, was greater than the rate of muscle size loss, therefore
resulting in an alteration of specific force (-24 %). The mechanisms underlying the
impairment in specific force, under these conditions, remain unclear. It is thought that
muscular and tendinous alterations are involved, such as a reduction in myosin
content (16,60) and tendon stiffness (11), and an impairment of the excitation-
contraction coupling (59). Indeed, Monti *et al.* (59) notably highlighted a reduced
contractile response of *vastus lateralis myo*fibers for the same excitatory stimulus
after 10 days of bed rest. Neural alterations, inclunding a reduction in voluntary
activation and coordination of antagonistic muscles (61), could also be potential
explanatory factors contributing to the impairment of specific force during muscle
disuse. In the current study, we observed an impairment of both maximal and
submaximal central motor drive, as revealed by the decrease in CAR_{MVC} (-3 %),
$CAR_{10\%}$ (-15 %) and RMS/M_{MAX} (-34 %). Particularly, the alteration of the CAR_{MVC}
was associated with the reduction in maximal force production capacity. In addition,
our data showed a rise in CAR_{MVC}/MVC and $CAR_{10\%}/10\%$ MVC ratios after the ULLS
protocol, suggesting an impairment in electromechanical efficiency after knee
extensors unloading. Though our findings are consistent with some previous
investigations on knee extensors and plantar flexors after 21-30 days of limb
suspension (23,29,53,62), some others did not observed any change in maximal
muscle activation, for the same muscle groups, after 10-30 days of limb unloading
(11,13,16,25,30,63). Despite these conflicting results, strength loss after unloading
seems to be more related to alterations in neural than muscular factors (13,64). As a
matter of fact, among multiple predictive variables, including muscle CSA, twitch RFD

and HRT or the amplitude of the H-reflex and M-wave for instance, it appears that

central activation is the most influential variable in mediating strength loss after

muscle disuse (13). This statement is also supported by the study of Kawakami *et al.*

(64) which showed that 55% of the variance in knee extensor maximal strength after

20 days of bed rest was explained by relative changes in voluntary activation.

However, it should be noted that their calculation was based only on a correlation

analysis without accounting for shared variance with other variables such as muscle

size.

Regarding mechanical twitch evoked by H_{MAX} stimulation at rest, no change

was reported following 10 days of ULLS in the current study. The present observation

contrasts with the result of Seynnes *et al.* (29) which reported a 40 and 54% increase

in H_{MAX} twitch amplitude for the plantar flexors after 14 and 23 days of ULLS

respectively. This discrepancy might depend on the muscle group investigated and

the different duration of exposure to limb unloading. On the other hand, in agreement

with data from Campbell *et al.* (16) following 21 days of ULLS, the current analysis,

did not find any change in M_{TW}. However, it should be noted that there is no

consensus in the literature about the evolution of M_{TW} following muscle disuse, since

studies have shown that ULLS decreased (15,53,57) or even increased (30) the

maximal twitch amplitude on lower limb muscles. In the current study, the M_{TW}

parameters (CT, RFD and HRT) were unaltered after the knee extensors' unloading.

These outcomes are in accordance with the literature focusing on the muscle

contractility of knee extensors following 21-30 days of ULLS (16,30,53). Specifically,

lower limb suspension did not result in the alteration of M_{TW} CT (16), RFD (16,30,53)

and HRT (16,30,53). Therefore, the present model of disuse does not seem to impair

knee extensors' contractility and relaxation processes, although it is effective in

inducing strength loss through muscle atrophy and impairments in muscle activation.

H-Reflex responses and M-wave

As observed at rest by de Boer *et al.* (11) following 23 days of ULLS, our

protocol did not modify the maximal M-wave obtained at rest, and during a weak

knee extensor contraction (10% MVC). This result indicates that sarcolemmal

excitability and potential action propagation were not altered by limb unloading

(65,66).

In the present study, unloading increased the spinal excitability of the VM at

rest, as revealed by the rise of 33% in H_{MAX}/M_{MAX}. Although there is no available data

on knee extensors, this finding is consistent with numerous earlier studies that have

shown an increase in spinal excitability of plantar flexors after 23-28 days of ULLS

(7,8,12–14,29,30). Our results showed that H_{MAX}/M_{MAX} obtained during a weak

contraction (10% MVC) was not altered by ULLS. However, it should be noted that

the background EMG activity (i.e., the RMS of VM EMG preceding H_{MAX} stimulation,

RMS/M_{MAX}) was different between LS0 and LS10. To overcome that limit, the H_{MAX}

was normalized by the RMS_H (H_{MAX}/RMS_H). The latter parameter appeared to be 2-

fold greater after the 10 days of ULLS, meaning that, for a similar muscle electrical

activity, the amplitude of the maximal H-reflex was increased. In light of these results,

it appears that unloading induced an enhancement of H-reflex amplitude both at rest

and during contraction. As shown by Delwaide (31), this enhancement could be

ascribed to an increase in motoneuronal excitability and/or a decrease in presynaptic

inhibition. Particularly, the latter explanation has been confirmed in an immobilization

model of disuse (28). In their study, Lundbye-Jensen & Nielsen (28) showed that

after immobilization of the left foot and ankle joint by a cast for 14 days, the increase

in *soleus* H-reflex results from decreased presynaptic inhibition of the Ia afferents.

More specifically, the authors reported a reduction in the primary afferent

depolarisation mechanism (i.e., a decrease in the inhibitory activity of a spinal

GABAergic interneuron) and a reduction in homosynaptic post-activation depression

(i.e., a decreased quantity of neurotransmitters available in the synapse following

repeated solicitation of the Ia afferents).

The present analysis showed that the decrease in muscle activation after a

short period of ULLS is accompanied by an increased spinal excitability. The latter

could be viewed as a compensatory adaptation, implemented by the nervous system

to counterbalance the reduced capacity to activate knee extensors. However, this

adaptation is not sufficient to counteract the entire cortical changes that occur during

muscle disuse. Additionally, it seems that the increase in the H-reflex following ULLS

is not related to the activation capacity, as reported by Seynnes *et al.* (30) on the

plantar flexors after 24 days of limb unloading. Indeed, the authors did not observe a

significant relationship between relative changes in amplitude of resting H-reflex and

V-wave obtained during MVC over the ULLS period, indicating that the factors

modulating the former had a limited effect on the latter (30). The monosynaptic T-

reflex of knee extensors, evoked by patellar tendon striking, was also used as a

supplementary tool to investigate the modulation of spinal circuitry following knee

extensors unloading. We observed no change in the relationship between the T-

reflex of the VM and the associated twitch, even if the spinal excitability was

enhanced. This result suggests that other factors that affect T-reflex, including the

mechanical properties of the tendon-aponeurosis and the sensitivity of the muscle

spindle, have probably been altered in such a way as to mask the effect of increased

spinal excitability on T-reflex (29). In contrast to our results, Seynnes *et al.* (29)

reported an increase in the slope and X-intercept of the T-reflex recruitment curve.

Likely, the discrepancy between the present and earlier findings may likely be

ascribed to the muscle group investigated (knee extensors *vs.* plantar flexors) and

the ULLS duration (10 *vs.* 23 days). However, these interpretations should be taken

with caution because of the limited number of subjects ($n = 6$) that underwent the T-

reflex testing procedure during the present investigation.

In the current study, neither $H_{LATENCY}$, $M_{LATENCY}$ nor $H-M_{LATENCY}$ were affected

by 10 days of lower limb suspension, suggesting that nerve conduction velocity

through the reflex arc and synaptic delay at the spinal level were not impaired by

unloading. This outcome is in contrast to those of Clark *et al.* (13) and Seynnes *et al.*

(30) who found an increase in soleus M_{MAX} and H_{MAX} latencies after 28 and 24 days

of ULLS respectively. More precisely, since the $H-M_{LATENCY}$, an index of conduction

velocity through the spinal reflex loop, was not changed in their studies (13,30),

authors primarily attributed the slowed conduction of H-reflex and M-wave responses

to the conduction velocity in the supply nerves, the branching axon terminals, and/or

transmission across the neuromuscular junction (67). No change in nerve conduction

through the reflex arc occurred after 10 days of ULLS for the lower limb muscles, and

confirm similar evidences from Manganotti *et al.* (68) who used the bed rest model

during an identical period of disuse. Further studies using ULLS protocols exceeding

10 days are needed to better understand the modulation kinetics of nerve conduction

in knee extensors.

In conclusion, the present study was the first to investigate the influence of

short-duration (10 days) ULLS on the neuromuscular function and spinal excitability

of knee extensors. The strength loss of knee extensors, observed after 10 days of

lower limb unloading, was associated with muscular and neural impairments, such as a reduction in quadriceps CSA and central activation. To compensate for reduced peripheral neurosensory inputs during the ULLS protocol, the spinal excitability of the VM was enhanced at rest and during a weak knee extension (10% MVC) after the ULLS protocol. These findings are paramount since they bring out new insights into the neural adaptations that occur in the major antigravity muscle in response to muscle disuse. Understanding the physiological adaptations during ULLS can guide the development of effective interventions to improve overall health outcomes for individuals experiencing muscle inactivity.

ACKNOWLEDGMENTS

The authors thank all volunteers who participated in the study.

FUNDING

The present study was funded by the Italian Space Agency (ASI), MARS-PRE, Project, n. DC-VUM-2017-006.

DISCLOSURES

No conflicts of interest, financial or otherwise, are declared by the authors. The results of the present study are presented clearly, honestly, and without fabrication, falsification, or inappropriate data manipulate. They do not constitute endorsement by ACSM.

DATA AVAILABILITY

Data will be made available on reasonable request.

REFERENCES

1. Nash CE, Mickan SM, Del Mar CB, Glasziou PP. Resting injured limbs delays recovery: a systematic review. J Fam Pract. 2004 Sep;53(9):706–12.

2. Lin CWC, Donkers NAJ, Refshauge KM, Beckenkamp PR, Khera K, Moseley AM. Rehabilitation for ankle fractures in adults. Cochrane Database Syst Rev. 2012 Nov 14;11:CD005595.

3. Graham ZA, Lavin KM, O'Bryan SM, Thalacker-Mercer AE, Buford TW, Ford KM, et al. Mechanisms of exercise as a preventative measure to muscle wasting. Am J Physiol Cell Physiol. 2021 Jul 1;321(1):C40–57.

4. Evans WJ. Skeletal muscle loss: cachexia, sarcopenia, and inactivity. Am J Clin Nutr. 2010 Apr;91(4):1123S-1127S.

5. Sarto F, Bottinelli R, Franchi MV, Porcelli S, Simunič B, Pišot R, et al. Pathophysiological mechanisms of reduced physical activity: Insights from the human step reduction model and animal analogues. Acta Physiologica. 2023;238(3):e13986.

6. Lee PHU, Chung M, Ren Z, Mair DB, Kim DH. Factors mediating spaceflight-induced skeletal muscle atrophy. American Journal of Physiology-Cell Physiology. 2022 Mar;322(3):C567–80.

7. Hackney KJ, Ploutz-Snyder LL. Unilateral lower limb suspension: integrative physiological knowledge from the past 20 years (1991-2011). Eur J Appl Physiol. 2012 Jan;112(1):9–22.

8. Campbell M, Varley-Campbell J, Fulford J, Taylor B, Mileva KN, Bowtell JL. Effect of Immobilisation on Neuromuscular Function In Vivo in Humans: A Systematic Review. Sports Med. 2019 Jun;49(6):931–50.

9. Tesch PA, Lundberg TR, Fernandez-Gonzalo R. Unilateral lower limb suspension: From subject selection to "omic" responses. J Appl Physiol (1985). 2016 May 15;120(10):1207–14.

10. Booth FW. Effect of limb immobilization on skeletal muscle. J Appl Physiol Respir Environ Exerc Physiol. 1982 May;52(5):1113–8.

11. de Boer MD, Maganaris CN, Seynnes OR, Rennie MJ, Narici MV. Time course of muscular, neural and tendinous adaptations to 23 day unilateral lower-limb suspension in young men. J Physiol. 2007 Sep 15;583(Pt 3):1079–91.

12. Clark BC. In vivo alterations in skeletal muscle form and function after disuse atrophy. Med Sci Sports Exerc. 2009 Oct;41(10):1869–75.

13. Clark BC, Manini TM, Bolanowski SJ, Ploutz-Snyder LL. Adaptations in human neuromuscular function following prolonged unweighting: II. Neurological properties and motor imagery efficacy. J Appl Physiol (1985). 2006 Jul;101(1):264–72.

14. Clark BC, Pierce JR, Manini TM, Ploutz-Snyder LL. Effect of prolonged unweighting of human skeletal muscle on neuromotor force control. Eur J Appl Physiol. 2007 Apr 12;100(1):53–62.

15. Horstman AM, de Ruiter CJ, van Duijnhoven NTL, Hopman MTE, de Haan A. Changes in
muscle contractile characteristics and jump height following 24 days of unilateral lower
limb suspension. Eur J Appl Physiol. 2012 Jan;112(1):135–44.

16. Campbell EL, Seynnes OR, Bottinelli R, McPhee JS, Atherton PJ, Jones DA, et al.
Skeletal muscle adaptations to physical inactivity and subsequent retraining in young
men. Biogerontology. 2013 Jun;14(3):247–59.

17. Cook SB, Kanaley JA, Ploutz-Snyder LL. Neuromuscular function following muscular
unloading and blood flow restricted exercise. Eur J Appl Physiol. 2014;114(7):1357–65.

18. Sarto F, Stashuk DW, Franchi MV, Monti E, Zampieri S, Valli G, et al. Effects of
short□term unloading and active recovery on human motor unit properties,
neuromuscular junction transmission and transcriptomic profile. The Journal of
Physiology. 2022 Sep 27;JP283381.

19. Dudley GA, Duvoisin MR, Adams GR, Meyer RA, Belew AH, Buchanan P. Adaptations to
unilateral lower limb suspension in humans. Aviat Space Environ Med. 1992
Aug;63(8):678–83.

20. Deschenes MR, Giles JA, McCoy RW, Volek JS, Gomez AL, Kraemer WJ. Neural factors
account for strength decrements observed after short-term muscle unloading. Am J
Physiol Regul Integr Comp Physiol. 2002 Feb;282(2):R578-583.

21. Deschenes MR, Holdren AN, McCoy RW. Adaptations to short-term muscle unloading in
young and aged men. Med Sci Sports Exerc. 2008 May;40(5):856–63.

22. Deschenes MR, McCoy RW, Mangis KA. Factors relating to gender specificity of
unloading-induced declines in strength. Muscle Nerve. 2012 Aug;46(2):210–7.

23. Schulze K, Gallagher P, Trappe S. Resistance training preserves skeletal muscle
function during unloading in humans. Med Sci Sports Exerc. 2002 Feb;34(2):303–13.

24. Tesch PA, Trieschmann JT, Ekberg A. Hypertrophy of chronically unloaded muscle
subjected to resistance exercise. J Appl Physiol (1985). 2004 Apr;96(4):1451–8.

25. Berg HE, Tesch PA. Changes in muscle function in response to 10 days of lower limb
unloading in humans. Acta Physiologica Scandinavica. 1996 May;157(1):63–70.

26. Duchateau J. Bed rest induces neural and contractile adaptations in triceps surae. Med
Sci Sports Exerc. 1995 Dec;27(12):1581–9.

27. Anderson J, Almeida-Silveira MI, Perot C. Reflex and muscular adaptations in rat soleus
muscle after hindlimb suspension. Journal of Experimental Biology. 1999 Oct
1;202(19):2701–7.

28. Lundbye-Jensen J, Nielsen JB. Immobilization induces changes in presynaptic control of
group Ia afferents in healthy humans. J Physiol. 2008 Sep 1;586(Pt 17):4121–35.

29. Seynnes OR, Maffiuletti NA, Maganaris CN, de Boer MD, Pensini M, di Prampero PE, et
al. Soleus T reflex modulation in response to spinal and tendinous adaptations to
unilateral lower limb suspension in humans. Acta Physiol (Oxf). 2008 Nov;194(3):239–
51.

30. Seynnes OR, Maffiuletti NA, Horstman AM, Narici MV. Increased H-reflex excitability is
not accompanied by changes in neural drive following 24 days of unilateral lower limb
suspension. Muscle Nerve. 2010 Nov;42(5):749–55.

31. Delwaide P. Human monosynaptic reflexes and presynaptic inhibition. In: Desmedt JE,
editor. Basel: Karger; 1973. p. 508–22. (New developments in electromyography and
clinical neurophysiology; vol. 3).

32. Paillard J. [Electrophysiologic analysis and comparison in man of Hoffmann's reflex and
myotatic reflex]. Pflugers Arch Gesamte Physiol Menschen Tiere. 1955;260(6):448–79.

33. Paillard J. Functional organization of afferent innervation of muscle studied in man by
monosynaptic testing. Am J Phys Med. 1959 Dec;38:239–47.

34. Józsa L, Kvist M, Kannus P, Järvinen M. The effect of tenotomy and immobilization on
muscle spindles and tendon organs of the rat calf muscles. A histochemical and
morphometrical study. Acta Neuropathol. 1988;76(5):465–70.

35. Valli G, Sarto F, Casolo A, Del Vecchio A, Franchi MV, Narici MV, et al. Lower limb
suspension induces threshold-specific alterations of motor units properties that are
reversed by active recovery. Journal of Sport and Health Science [Internet]. 2023 Jun 17
[cited 2023 Nov 7]; Available from:
https://www.sciencedirect.com/science/article/pii/S2095254623000595

36. Bleeker MWP, Hopman MTE, Rongen GA, Smits P. Unilateral lower limb suspension can
cause deep venous thrombosis. Am J Physiol Regul Integr Comp Physiol. 2004
Jun;286(6):R1176-1177.

37. Wanner M, Hartmann C, Pestoni G, Martin BW, Siegrist M, Martin-Diener E. Validation of
the Global Physical Activity Questionnaire for self-administration in a European context.
BMJ Open Sport & Exercise Medicine. 2017 May 1;3(1):e000206.

38. Guette M, Gondin J, Martin A. Time-of-day effect on the torque and neuromuscular
properties of dominant and non-dominant quadriceps femoris. Chronobiol Int.
2005;22(3):541–58.

39. Douglas CM, Hesketh SJ, Esser KA. Time of Day and Muscle Strength: A Circadian
Output? Physiology (Bethesda). 2021 Jan 1;36(1):44–51.

40. Hopkins JT, Ingersoll CD, Andrew Krause B, Edwards JE, Cordova ML. Effect of knee
joint effusion on quadriceps and soleus motoneuron pool excitability: Medicine and
Science in Sports and Exercise. 2001 Jan;123–6.

41. Zehr E. Considerations for use of the Hoffmann reflex in exercise studies. European
journal of applied physiology. 2002 May 1;86:455–68.

42. Mau-Moeller A, Behrens M, Lindner T, Bader R, Bruhn S. Age-related changes in
neuromuscular function of the quadriceps muscle in physically active adults. Journal of
Electromyography and Kinesiology. 2013 Jun 1;23(3):640–8.

43. Doguet V, Jubeau M. Reliability of H-reflex in vastus lateralis and vastus medialis
muscles during passive and active isometric conditions. Eur J Appl Physiol. 2014
Dec;114(12):2509–19.

44. Burke D, ADAMS RW, SKUSE NF. The effects of voluntary contraction on the H reflex of
human limb muscles. Brain. 1989 Apr 1;112(2):417–33.

45. Knikou M. The H-reflex as a probe: pathways and pitfalls. J Neurosci Methods. 2008 Jun
15;171(1):1–12.

46. Funase K, Miles TS. Observations on the variability of the H reflex in human soleus.
Muscle & Nerve. 1999;22(3):341–6.

47. McNeil CJ, Butler JE, Taylor JL, Gandevia SC. Testing the excitability of human
motoneurons. Frontiers in Human Neuroscience [Internet]. 2013 [cited 2019 Dec 10];7.
Available from: http://journal.frontiersin.org/article/10.3389/fnhum.2013.00152/abstract

48. Monti E, Franchi MV, Badiali F, Quinlan JI, Longo S, Narici MV. The Time-Course of
Changes in Muscle Mass, Architecture and Power During 6 Weeks of Plyometric
Training. Frontiers in Physiology [Internet]. 2020 [cited 2022 Nov 3];11. Available from:
https://www.frontiersin.org/articles/10.3389/fphys.2020.00946

49. Gandevia SC, Allen GM, Butler JE, Taylor JL. Supraspinal factors in human muscle
fatigue: evidence for suboptimal output from the motor cortex. J Physiol. 1996;8.

50. Theodosiadou A, Henry M, Duchateau J, Baudry S. Revisiting the use of Hoffmann reflex
in motor control research on humans. Eur J Appl Physiol. 2023 Apr;123(4):695–710.

51. Bakdash JZ, Marusich LR. Repeated Measures Correlation. Front Psychol [Internet].
2017 [cited 2022 Mar 31];0. Available from:
https://www.frontiersin.org/articles/10.3389/fpsyg.2017.00456/full

52. Fritz CO, Morris PE, Richler JJ. Effect size estimates: Current use, calculations, and
interpretation. Journal of Experimental Psychology: General. 2012;141(1):2–18.

53. Cook SB, Brown KA, Deruisseau K, Kanaley JA, Ploutz-Snyder LL. Skeletal muscle
adaptations following blood flow-restricted training during 30 days of muscular unloading.
J Appl Physiol (1985). 2010 Aug;109(2):341–9.

54. de Boer MD, Selby A, Atherton P, Smith K, Seynnes OR, Maganaris CN, et al. The
temporal responses of protein synthesis, gene expression and cell signalling in human
quadriceps muscle and patellar tendon to disuse. J Physiol. 2007 Nov 15;585(Pt 1):241–
51.

55. Brook MS, Stokes T, Gorissen SHM, Bass JJ, McGlory C, Cegielski J, et al. Declines in
muscle protein synthesis account for short☐term muscle disuse atrophy in humans in the
absence of increased muscle protein breakdown. J Cachexia Sarcopenia Muscle. 2022
Aug;13(4):2005–16.

56. Brocca L, Longa E, Cannavino J, Seynnes O, De Vito G, McPhee J, et al. Human
skeletal muscle fibre contractile properties and proteomic profile: adaptations to 3 weeks
of unilateral lower limb suspension and active recovery: Impact of ULLS and active
rehabilitation on human muscle fibres. J Physiol. 2015 Dec 15;593(24):5361–85.

57. Clark BC, Fernhall B, Ploutz-Snyder LL. Adaptations in human neuromuscular function
following prolonged unweighting: I. Skeletal muscle contractile properties and applied
ischemia efficacy. J Appl Physiol (1985). 2006 Jul;101(1):256–63.

58. Demangel R, Treffel L, Py G, Brioche T, Pagano AF, Bareille MP, et al. Early structural
and functional signature of 3-day human skeletal muscle disuse using the dry immersion
model. J Physiol. 2017 Jul 1;595(13):4301–15.

59. Monti E, Reggiani C, Franchi MV, Toniolo L, Sandri M, Armani A, et al. Neuromuscular
junction instability and altered intracellular calcium handling as early determinants of
force loss during unloading in humans. The Journal of Physiology. 2021;599(12):3037–
61.

60. D'Antona G, Pellegrino MA, Adami R, Rossi R, Carlizzi CN, Canepari M, et al. The effect
of ageing and immobilization on structure and function of human skeletal muscle fibres. J
Physiol. 2003 Oct 15;552(Pt 2):499–511.

61. Maganaris CN, Baltzopoulos V, Ball D, Sargeant AJ. In vivo specific tension of human
skeletal muscle. J Appl Physiol (1985). 2001 Mar;90(3):865–72.

62. Seynnes OR, Maganaris CN, de Boer MD, di Prampero PE, Narici MV. Early structural
adaptations to unloading in the human calf muscles. Acta Physiol (Oxf). 2008
Jul;193(3):265–74.

63. Hotta N, Ishida K, Sato K, Koike T, Katayama K, Akima H. The effect of intense interval
cycle-training on unloading-induced dysfunction and atrophy in the human calf muscle. J
Physiol Anthropol. 2011;30(1):29–35.

64. Kawakami Y, Akima H, Kubo K, Muraoka Y, Hasegawa H, Kouzaki M, et al. Changes in
muscle size, architecture, and neural activation after 20 days of bed rest with and without
resistance exercise. European Journal of Applied Physiology. 2001 Feb 16;84(1–2):7–
12.

65. Fuglevand AJ, Zackowski KM, Huey KA, Enoka RM. Impairment of neuromuscular
propagation during human fatiguing contractions at submaximal forces. J Physiol (Lond).
1993 Jan;460:549–72.

66. Allen DG, Lamb GD, Westerblad H. Skeletal muscle fatigue: cellular mechanisms.
Physiol Rev. 2008 Jan;88(1):287–332.

67. Kim BJ, Date ES, Park BK, Choi BY, Lee SH. Physiologic changes of compound muscle
action potentials related to voluntary contraction and muscle length in carpal tunnel
syndrome. J Electromyogr Kinesiol. 2005 Jun;15(3):275–81.

68. Manganotti P, Buoite Stella A, Ajcevic M, di Girolamo FG, Biolo G, Franchi MV, et al.
Peripheral nerve adaptations to 10 days of horizontal bed rest in healthy young adult
males. American Journal of Physiology-Regulatory, Integrative and Comparative
Physiology. 2021 Sep;321(3):R495–503.

FIGURE CAPTIONS

Figure 1. Maximal force and corresponding central activation. The maximal voluntary contraction (MVC) and corresponding central activation ratio (CAR$_{MVC}$) were individually scatter plotted before (white circles) and after (grey circles) 10 days of unilateral lower limb suspension (n = 10). Black (before limb suspension) and grey (after limb suspension) crosses represent the mean of both scatter plots. ***$p < 0.001$ and $^{\$\$}p < 0.01$ express significant differences in MVC and CAR$_{MVC}$, respectively, between before and after lower limb suspension.

Figure 2. Individual repeated measures correlations (rmcorr) between muscle function parameters and maximal force production (n = 10). Rmcorr was plotted between qCSA and MVC (A), and between CAR and MVC (B). Observations from the same participant are given the same colour, the lines correspond to the rmcorr fit

for each participant. MVC = maximal voluntary contraction; qCSA = quadriceps

cross-sectional area; CAR = central activation ratio.

Figure 3. Twitch amplitude associated with H_{MAX} and M_{MAX}. The twitch amplitude

associated with H_{MAX} (*n* = 8) and M_{MAX} (*n* = 9) was measured at rest before (white

columns) and after (grey columns) 10 days of unilateral lower limb suspension.

Circles and lines represent individual data evolution. H_{MAX} = maximal H-reflex; M_{MAX}

= maximal M-wave.

Figure 4. Changes in spinal excitability after 10 days of ULLS. A. H_{MAX}/M_{MAX} ratio

was calculated before (white columns) and after (grey columns) 10 days of ULLS,

either at rest (*n* = 8) and during 10% MVC contractions (*n* = 9). Circles and lines

represent individual data evolution. B. H_{MAX} and M_{MAX} traces at rest, from a typical

subject, before (red lines) and after (black lines) ULLS. C. Representative data of VM

EMG activity during the 10% MVC contraction with associated H_{MAX} before (red line)

and after (black line) ULLS from a typical subject. VM EMG = electromyographic

activity of vastus medialis; H_{MAX} = maximal H-reflex; M_{MAX} = maximal M-wave. ULLS

= unilateral lower limb suspension. *p < 0.05 express significant difference.

Table 1. Muscle morphology, strength and activation capacity.

	BDC (n = 10)	**LS9** (n = 10)	*p*-value
VM CSA (cm²)	19.53 ± 2.61	18.02 ± 2.41	0.0005
qCSA (cm²)	57.79 ± 6.54	55.26 ± 6.20	0.007
	LS0 *(n = 10)*	**LS10** *(n = 10)*	*p*-value
MVC (kg)	82.54 ± 11.13	60.19 ± 9.13	< 0.0001
10% MVC (kg)	8.25 ± 1.11	6.02 ± 0.91	< 0.0001
MVC/qCSA $(kg \cdot cm^{-2})$	1.43 ± 0.18	1.09 ± 0.16	0.0001
CAR_{MVC}	0.980 ± 0.013	0.950 ± 0.033	0.006
CAR_{MVC}/MVC (kg^{-1})	0.012 ± 0.002	0.016 ± 0.002	< 0.0001
$CAR_{10\%}$	0.428 ± 0.065	0.364 ± 0.063	0.003
$CAR_{10\%}$/10% MVC (kg^{-1})	0.052 ± 0.009	0.061 ± 0.011	0.032
RMS/M_{MAX} at 10% MVC $(x10^{-3})$	9.39 ± 4.05	6.18 ± 3.54	0.020

All variables were measured before (LS0) and after (LS10) 10 days of unilateral lower limb suspension, except CSA measured the day of ULLS beginning (BDC) and the day before the ULLS end (LS9). MVC = knee extensors maximal voluntary contraction; VM = vastus medialis muscle; CSA = muscle cross-sectional area; qCSA = quadriceps CSA; CAR_{MVC} and $CAR_{10\%}$ = central activation ratio calculated during a MVC and during a contraction at 10% MVC; RMS = root mean square; M_{MAX} = maximal M-wave. Values are reported as mean ± standard deviation.

Table 2. Mechanical parameters associated with M_{MAX} before and after ULLS.

	M_{MAX} stimulations		*p*-value
	LS0 *(n = 9)*	**LS10** *(n = 9)*	
M_{TW}/M_{MAX} (kg/mV)	1.38 ± 1.07	1.33 ± 1.25	0.441
CT (ms)	82.44 ± 8.59	87.33 ± 14.03	0.280
RFD (kg/ms)	0.247 ± 0.060	0.210 ± 0.048	0.130
HRT (ms)	83.60 ± 15.15	82.44 ± 17.80	0.873

M_{MAX} = maximal M-wave; M_{TW} = twitch associated with M_{MAX}; CT = twitch contraction time; RFD = twitch rate of force development; HRT = twitch half relaxation time; ULLS = unilateral lower limb suspension; LS0 and LS10 = before and after 10 days of ULLS. Values are reported as mean ± standard deviation.

Table 3. Spinal excitability and nerve conduction parameters before and after 10 days of ULLS.

	Rest			Contraction		
	LS0	LS10	p-value	LS0	LS10	*p*-value
M_{atHMAX} (% M_{MAX})	8.39 ± 7.35	6.64 ± 8.65	0.484	9.02 ± 9.03	5.17 ± 4.63	0.173
H_{MAX} (mV)	2.22 ± 1.94	3.53 ± 3.25	0.036	4.29 ± 3.21	4.37 ± 3.44	0.441
M_{MAX} (mV)	13.12 ± 7.04	13.30 ± 7.41	0.803	13.48 ± 7.97	12.89 ± 7.91	0.215
H_{MAX}/RMS_H	-	-	-	37.27 ± 12.95	75.62 ± 47.69	0.038
$M_{LATENCY}$ (ms/m)	15.22 ± 1.60	15.74 ± 1.81	0.613	-	-	-
$H_{LATENCY}$ (ms/m)	48.04 ± 5.55	48.72 ± 5.55	0.396	-	-	-
H-$M_{LATENCY}$ (ms/m)	34.07 ± 5.34	33.84 ± 4.02	0.782	-	-	-
T-twitch slope	0.253 ± 0.096	0.331 ± 0.275	0.422	-	-	-
T-twitch intercept	0.876 ± 0.585	1.550 ± 0.807	0.221	-	-	-

Data were obtained at rest and/or during 10% MVC contractions. For each parameter, the number of subjects was 9, except for M_{atHMAX} at rest ($n = 8$), $H_{LATENCY}$ and H-$M_{LATENCY}$ ($n = 7$), T-twitch slope and intercept ($n = 6$). LS0 = before leg suspension; LS10 = after 10 days of leg suspension; H_{MAX} = maximal H-reflex; M_{MAX} = maximal M-wave ; M_{atHMAX} = M-wave associated with maximal H-reflex; $H_{LATENCY}$ and $M_{LATENCY}$ = H_{MAX} and M_{MAX} latency respectively; H-$M_{LATENCY}$ = difference between $H_{LATENCY}$ and $M_{LATENCY}$; T = T-reflex; twitch = T-reflex associated twitch; MVC = maximal voluntary contraction; ULLS = unilateral lower limb suspension. Values are reported as mean ± standard deviation. Please note that some parameters are obtained only in one condition (rest or contraction). *$p < 0.05$ and **$p < 0.01$ express significant difference from LS0 for the considered condition. Values are reported as mean ± standard deviation.