TITLE:

Effect of 10 days of unilateral lower limb suspension on knee extensors neuromuscular function

and spinal excitability

RUNNING TITLE:

Muscle disuse and neuromuscular function

AUTHORS:

Loïc Lebesque[1], Marco V. Narici[2,3], Alain Martin[1], Giuseppe De Vito[2,3], Fabio Sarto[2] and Gil

Scaglioni[1]

AFFILIATIONS

[1] INSERM UMR1093-CAPS, Université Bourgogne Franche-Comté, UFR des Sciences du Sport,

F-21000, Dijon.

[2] Department of Biomedical Sciences, University of Padova, Padova, 35131, Italy.

[3] CIR-MYO Myology Center, University of Padua, 35131, Italy.

CORRESPONDING AUTHOR: Address for reprint requests and other correspondence:

Gil Scaglioni

INSERM UMR1093 "Cognition, Action et Plasticité Sensorimotrice »

Université de Bourgogne - UFR STAPS

Campus Universitaire, BP 27887 DIJON

Phone: +33 3 80 39 67 58

E-mail : gil.scaglioni@u-bourgogne.fr

ABSTRACT

The reduction in mechanical loading applied on the lower limb has numerous detrimental consequences on neuromuscular function. The current study aimed to investigate the changes in knee extensors' strength and spinal excitability induced by unilateral lower limb suspension (ULLS), providing new insights into the neuromuscular adaptations to muscle hypoactivity. Ten young healthy males (19-28 years old) underwent 10 days of ULLS to simulate muscle disuse. Modulation by unloading of knee extensors' function (muscle morphology and strength, activation capacity and contractile properties) and spinal reflexes were explored before and after the ULLS. The knee extensors' anatomical cross-sectional area (-4%, $p = 0.007$), maximal strength (-27%, $p < 0.001$) and central activation ratio (-3%, $p = 0.006$) were reduced after 10 days of ULLS. Vastus medialis H-reflex amplitude was enhanced both at rest (+33%, $p = 0.038$) and during a low-intensity contraction set at 10% of maximal strength (+103%, $p = 0.038$). No changes in muscle contractility and nerve conduction velocity were observed after the ULLS. The present study suggests that neural impairments mainly contribute to the decrease in knee extensors' strength induced by short-term ULLS. The decrease in muscle activation after a short period of ULLS was accompanied by an increase in spinal excitability. However, the latter adaptation did not counterbalance the reduction in activation capacity and thus in maximal strength resulting from ULLS. These adaptations to short-term ULLS should be considered when aiming at improving the neuromuscular function of people experiencing muscle hypoactivity.

NEW & NOTEWORTHY (75 words)

This study provides new insights into the effects of muscle hypoactivity on neuromuscular function and spinal excitability in the major antigravity muscle group of the lower limb. Neural impairments primarily contribute to maximal strength loss after short-term unilateral lower

51 limb suspension, while spinal excitability increased. These findings are crucial as they offer

52 valuable understanding for developing effective interventions to improve health outcomes for

53 individuals experiencing muscle inactivity.

55 **KEY WORDS**

56 Atrophy; Disuse; Force control; H-reflex; T-reflex; Unloading; Vastus medialis

INTRODUCTION

Reductions in mechanical loading may occur as a consequence of surgery, injury and illness (1–3), but also as a result of reduced daily physical activity (4,5) or gravitational input, as during spaceflight (6). The direct outcome is a loss of muscle mass and function, associated with changes in motor control (7,8).

Several models of immobilization allow to explore the neuromuscular impairment in response to disuse (8). However, unilateral lower limb suspension (ULLS) enables to restrain atrophy to the explored musculoskeletal segment, easily mimicking the joint unloading that follows injuries or surgery (7,9) and, compared to casting, has the advantage of being less binding.

Extensive literature shows that the extent of muscle weakness is greater than the decrease in muscle mass since the first few days of ULLS, meaning that neuromuscular function is rapidly impaired by disuse (7,9–12). Among the antigravity muscles, the knee extensors appear to be the most affected by ULLS (13–15), showing a strength loss considerably larger than that of the plantar flexors (-1.0% $\cdot$ day^{-1} *vs.* 0,7% $\cdot$ day^{-1}, for durations between 10 and 35 days of ULLS; 8,11). Therefore, although it is reasonable to assume that for unloading, as for overloading, changes in neural drive are the first to occur (8), their involvement in the weakening of antigravity muscles, associated with disuse, is still an unresolved issue. Although studies have shown that ULLS (21-30 days) does not interfere with the central motor drive of knee extensors muscles (9,11,15,16), others have found a decrease (17,18) or even an increase (range + 0.1%.day^{-1} increase to − 0.2%.day^{-1} decrease; median − 0.2%.day^{-1}) of muscle voluntary activation (8). This divergence was also observed in surface electromyographic (EMG) data, recorded during maximal voluntary contractions, with studies showing a reduction of the vastii activity after ULLS (7-42 days) (19–23), and others finding no change for the vastii and RF

(ULLS 7-35 days) (11,22,24). On the other hand, the literature agrees on the EMG data recorded

during submaximal contractions. Interestingly, in this case, all studies showed that vastii EMG

activity increased during submaximal contractions following ULLS (7,23–25), suggesting that

additional motor units are necessary to produce the same absolute force output after

unloading. In light of these considerations, it appears that changes at neural level have not yet

been fully understood, and neither have the underlying mechanisms. Previous studies, on

plantar flexors, have shown an unequivocal enhancement in the amplitude of the Hoffman (H)-

reflex in response to unloading (7,8,12–14,26–30). The H-reflex is the EMG response evoked

by electrically stimulating the Ia fibers of a peripheral nerve and reflects the efficacy of the

synaptic transmission between Ia afferents and the motor neurons (31). Since it is generally

considered as a comprehensive index of spinal excitability (e.g. 32,33), for greater readability

we will use the term 'spinal excitability' to refer to the efficiency of the synaptic transmission

between Ia afferents and motor neurons. The enhancement of H-reflex amplitude with

unloading may be ascribed to an increased synaptic transmission efficiency and/or to a greater

motoneuronal excitability (13,30,34), however, the improved synaptic transmission efficiency

seems to be the predominant adaptation in all forms of disuse (13). This mechanism may

interfere with the reduction in motoneuronal excitability that might occur together with an

increased proportion of fast fibers in response to unloading (27). These adjustments should

equally affect H and T-reflexes. The T-reflex is an electrophysiological variant of the stretch

reflex caused by a mechanical tap applied on the tendon that, generating a quick stretch of the

tendon itself, triggers muscle spindles discharge to Ia afferents notably. Both reflexes originate

partly from the monosynaptic projections of the Ia afferents to alpha-motor neurones, but the

T-reflex involves all the elements of the reflex pathway, including muscle spindles sensitivity

and II afferents (35,36). The literature shows that the T-reflex may be further modulated by

conflicting changes arising in peripheral components of the stretch reflex pathway. Actually,

Anderson *et al.* (27) found, after 3 weeks of hind limb suspension in rat, a decline of the maximal

soleus T-reflex but a rise in the maximal H-reflex (H_{MAX}). Seynnes *et al.* (29) observed no change

in soleus T-reflex, following 23 days of ULLS, but an increase in H-reflex and a decrease in

tendon stiffness. The different modulations of the T-reflex response with unloading may be

ascribed i) to a lower spindles' sensitivity, due to their reduced solicitation during

immobilization (37), and/or ii) to a decreased stiffness of the musculo-tendinous elements in

series with the muscle spindles (27,29).

In view of these considerations and since the literature seems to show that few days of

immobilization induce significant changes in neuromuscular function (8), this study sought to

investigate the role of changes in spinal reflex excitability, muscle contractile characteristics

and muscle morphology in the loss of knee extensors' strength following 10 days of muscle

unloading. Since, in plantar flexors, H-reflex is always modulated by ULLS and considering that

knee extensors are among the antigravity muscles most affected by inactivity, we hypothesized

that spinal excitability would be increased and muscle mass reduced by unloading. However,

given the short period of immobilization, it could also be assumed that neural adaptations

might be preponderant over muscular changes. Therefore, apart from improving knowledge on

mechanisms responsible for the decline of the neuromuscular function due to unloading, the

outcomes of the present investigation may be useful for refining the rehabilitation strategies

of patients following periods of muscle disuse.

MATERIALS & METHODS

Ethical approval.

The present study was approved by the Ethics Committee of the Department of Biomedical Sciences of the University of Padova (Italy) (reference number HEC-DSB/01-18) and conducted in accordance with the standards set by the latest revision of the Declaration of Helsinki. Participants were informed about all experimental procedures through an interview and an information sheet, and could ask questions to the selection committee. After signing the written consent form, participants were enrolled in the study with the possibility to withdraw at any stage of the experiment. This study was part of a larger investigation aiming to study the neuromuscular alterations with short-term disuse (18,38).

Participants

Ten healthy and recreationally active young male volunteers (age: 22.9 ± 4.0 years old; body mass: 72.2 ± 7.2 kg; height: 177.6 ± 3.1 cm; body mass index: 22.9 ± 2.1 kg·m^{-2}) took part to this study. Because unilateral lower limb suspension (ULLS) is associated with an increased risk of deep venous thrombosis (39), more commonly experienced in young females (40), only males were recruted in this study. Inclusion criteria were: age comprised between 18 and 35 years, body mass index comprised between 20 and 28 kg·m^{-2} and involvement in recreational physical activities 1-3 times per week. Exclusion criteria were: sedentary lifestyle, professional sport practice, smoking, history of deep venous thrombosis and acute or chronic musculoskeletal, metabolic and cardiovascular disorders. Using the Global Physical Activity Questionnaire (41), participants demonstrated a high level of physical activity (3268 ± 1468 MET·min·week^{-1}) in the period leading up to the ULLS experiment. Volunteers were asked to

maintain their habitual diet throughout the intervention period and refrain from intense

exercise, coffee and alcohol intake during the 24 hours preceding the data collection at each

time point.

Protocol

Experimental design. Data were collected the day before (LS0) and after 10 days (LS10) of

unilateral lower limb suspension (ULLS). For each subject, sessions lasted 2 hours and were

planned at the same time of the day to avoid any circadian disturbance (42,43). At both LS0

and LS10, the experimental procedure was performed beginning with the evaluation of the

maximal voluntary isometric contraction (MVC) of knee extensors and the maximal voluntary

activation level (VAL). Then, vastus medialis (VM) spinal excitability were assessed through

electrical stimulations and patella tendon striking at rest and during a weak contraction of knee

extensors (10% MVC assessed for each session). In addition, quadriceps and VM morphology

was assessed through ultrasonography, on the same day of the ULLS beginning (baseline data

collection, BDC) and the day before the end of the ULLS (LS9).

Maximal force production capacity. To start the maximal force production evaluation, subjects

performed a knee extensors warm-up composed of 10 submaximal contractions, lasting 4 s

each. Then, three knee extensors MVCs (lasting 4 s, with 1-min rest in-between) were

performed with maximal electrical doublet (100 Hz) delivered over the peak force.

Spinal excitability.　Since it is more challenging to evoke rest H-reflex in the quadriceps muscles

compared to the soleus muscle, several studies suggest eliciting it during a weak muscle

contraction (44–47). The reflex response is indeed enhanced by the muscle contraction

because motoneurons are brought closer to their firing threshold by the voluntary drive (48).

In addition, muscle activation allows to keep constant the "background" level of α-motoneuron

depolarisation and limits post-synaptic modulations (45,49). Therefore, the assessment of

spinal excitability during muscle contraction has the advantage of increasing amplitude and

reducing variability in both latency and amplitude of the H-reflex (48,50,51). Consequently, VM

H-reflex and M-wave recruitment curves were built by applying electrical stimulations first

during weak knee extensors contractions (10% MVC) and then at rest. To build the recruitment

curves, stimulations started at 5 mA and were progressively increased with a 5-mA increment

until the maximal M-wave (M_{MAX}) was reached. To ensure that the M-wave lied in the plateau

of its maximal value throughout the experiment, the maximal stimulation intensity was

increased by 20%. Subsequently, additional stimulations were applied, in 2 mA increments,

around the intensity that allowed to record the highest H-reflex, in order to accurately estimate

the intensity needed to elicit the maximal H reflex (H_{MAX}). Four stimuli were delivered at each

intensity interspaced by 10 s intervals. Then, 6 out of 10 subjects underwent complementary

evaluation of spinal excitability at rest through the assessment of tendinous reflex (i.e., T-reflex,

for the method see below mechanical stimulation).

Unilateral lower limb suspension. The ULLS model was employed for 10 days. The non-dominant

leg (left leg for all participants) of the participants was fitted with a shoe having an elevated

sole (50 mm), while the dominant leg (right leg for all participants) was suspended and kept at

a slightly flexed position (approximately 15-20° of knee flexion) using straps. Volunteers were

asked to walk on crutches during the whole ULLS period and to refrain from loading the

suspended leg in any way. A familiarization session was set up for the participants to practice

daily tasks with the limb suspended (9). Participants were recommended to wear elastic

compression socks on the suspended leg during ULLS period and to perform passive, range of

motion, non-weight-bearing exercises of the ankle, as precautionary measures to prevent deep

venous thrombosis (39). Moreover, we performed an ultrasound-Doppler examination after 5

days of ULLS. Participants' compliance was evaluated through daily calls and messages and by

comparing calves' temperature and circumference after 5 and 10 days of ULLS, as previously

suggested (9).

Data acquisition.

Ultrasonography. The morphology of the quadriceps was assessed by ultrasonography, which

also allowed us to define the morphology of the VM, for which electrophysiological data were

also recorded. To evaluate morphology of quadriceps and VM muscles, subjects were lying on

their backs on an auscultation table. Muscles anatomical cross-sectional area (ACSA) was

obtained using extended-field-of-view ultrasonography imaging (Mylab, Esaote, Genoa, Italy).

A 47 mm, 7.5 MHz linear array transducer was used to collect images at different muscle length

percentages. Three regions of interest were detected at 30%, 50% and 70% of femur length

(measured as the distance between the great trochanter and the mid-patellar point), where 0%

represents the mid-patellar point (distal part) and 100% the greater trochanter (proximal part).

An adjustable guide was used in each acquisition to keep the same transverse path (52). After

applying a generous amount of transmission gel on the participant's skin to improve the

acoustic contact, the transducer was moved slowly in the transverse plane from the medial

border of the VM to the lateral border of the vastus lateralis. During the transducer

displacement, we kept the pressure on the skin as constant as possible. Two scans were

obtained for each site.

Force recordings. Voluntary and electrically-evoked force of knee extensors was recorded using

a custom-made knee dynamometer fitted with a strain gauge. The hip and knee angles were

set at 90° (180° full extension). Subject's dominant leg was securely strapped to the custom-

made knee dynamometer, equipped with a load cell, at the ankle level. Subject was asked to

cross the arms on the chest and keep the head in a neutral position during the experiment

procedure. The force exerted by the subject was displayed in real-time on a computer screen

placed at 1 m in front of him. Because spinal excitability was further assessed during a weak

quadriceps contraction, a line representing 10% of the session MVC was added on the

computer screen once the MVC was determined. Force data were were recorded at a sampling

frequency of 1 kHz with the Spike2 software (version 8.01, Cambridge Electronics Design Ltd,

Cambridge, UK; RRID: SCR_000903) and stored for offline analysis with the LabChart software

(version 8, AD Instruments®, Australia; RRID: SCR_017551).

Electromyographic recordings. Electromyographic activity (EMG) from the dominant-leg VM

was recorded during the 10% MVC contractions with bipolar active-type Ag-AgCl electrodes

electrodes (44 x 22 mm, Ambu®). To minimize skin impedance (< 5 kΩ), the skin was shaved,

rubbed with an abrasive paste and cleaned with alcohol, before electrode positioning.

Electrodes were positioned 10 cm above the patella upper part and 4 cm medially with an inter-

electrode center-to-center distance of 20 mm. The EMG signals were amplified (gain = 1,000)

and bandpass filtered at 5 Hz to 5 kHz using a CED 1902 amplifier (Cambridge Electronics Design

Ltd, Cambridge, UK) and digitized with a CED Micro 1401 data acquisition unit (Cambridge

Electronics Design Ltd, Cambridge, UK). EMG data were recorded at a sampling frequency of

40 kHz with the Spike2 software (version 8.01, Cambridge Electronics Design Ltd, Cambridge,

UK; RRID: SCR_000903) and stored for offline analysis with the LabChart software (version 8,

AD Instruments®, Australia; RRID: SCR_017551).

Nerve electrical stimulations. Nerve electrical stimulations were delivered using a high-voltage

(400 V) constant-current stimulator (DS7AH model, Digitimer, Hertfordshire, UK). For maximal

voluntary activation assessment, electrical doublets (100 Hz, monophasic rectangular pulses, 1

ms width) were delivered through two stimulation electrodes (5 x 10 cm, Compex SA, Ecublens,

Switzerland) placed in the proximal and distal region of the knee extensors. Maximal electrical

stimulation intensity was determined as the intensity eliciting the maximal twitch amplitude.

For spinal excitability assessment, we delivered percutaneous electrical stimulations

(monophasic rectangular pulses, 1 ms width) in the femoral trigone over the femoral nerve to

evoke electrophysiological and mechanical responses of the VM muscle. The self-adhesive

anode (5 x 10 cm, Compex SA, Ecublens, Switzerland) was placed in the gluteal fossa. The

optimum stimulation site was located with a hand-held cathode stylus in order to obtain the

greatest H-reflex amplitude of the VM for the lowest stimulation intensity. Then, the self-

adhesive cathode electrode (7 mm diameter, Contrôle Graphique Medical, Brie-Compte-

Robert, France) was positioned and firmly fixed with a velcro elastic band.

Mecanichal stimulations. During T-reflex procedure, subjects were positioned as described

previously in *Force recordings* section. Twenty T-reflexes were elicited for each subject by

tapping the anterior face of the patellar tendon of the quadriceps with a manual reflex hammer

(neurological hammer, Babinsky model) at various intensities (53). The T-reflex on the VM EMG

signal and its associated mechanical twitch were recorded for each strike.

Data analysis.

Ultrasonography. Of the two scans obtained at each site, we analyzed the image with the best

quality. Quadriceps and VM ACSAs were measured by tracing the muscle contours with the

image analysis software "ImageJ" (https://imagej.net; RRID: SCR_003070). Two measurements

per parameter were performed on each image, and their average was considered. Quadriceps

ACSA (qACSA) was computed as the average value of the ACSAs recorded at 30, 50 and 70% of

the femur length for each subject at both LS0 and LS10. VM ACSA was calculated as the average

value of the ACSAs recorded at 30 and 50% of the femur length, due to technical issues in clearly

detecting the border between the VM and vastus intermedius at 70% of the femur length.

Maximal force production. The MVC was determined as the highest force value reached during

the 3 MVCs. The target of 10% MVC was calculated from the MVC of each session (i.e., LS0 or

LS10). The normalized force was calculated as the MVC divided by the quadriceps ACSA

(MVC/qACSA).

Activation capacity. The voluntary activation associated with the maximal force production and

the weak contraction (10% MVC) was quantified with the central activation ratio (CAR_{MVC} and

$CAR_{10\%}$ respectively) as follows (54):

308
$$CAR = \frac{Torque}{Torque + superimposed\ twitch}$$

where torque was either the MVC (CAR_{MVC}) or the 10% MVC ($CAR_{10\%}$). A higher CAR value

indicates an increase in muscle activation. It should be noted that, while the CAR_{MVC} was

calculated using maximal doublet stimulations (100 Hz) delivered during brief MVC, the $CAR_{10\%}$

was derived from the M-wave recruitment curve recorded during weak contractions at 10%

MVC using maximal single stimulations (see above *Spinal excitability* section). The ratio

between the $CAR_{10\%}$ and the force produced during the contraction at 10% MVC was calculated

to determine the motor unit activity required to achieve this specific target force. The root

mean square (RMS) of VM EMG activity was computed offline over a 500 ms period preceding

H_{MAX} and M_{MAX} stimulations during the weak contractions at 10% MVC, and subsequently

normalized to the M_{MAX} amplitude (RMS/M_{MAX}).

Spinal excitability parameters. One subject has been exempted from the spinal excitability

assessment because he did not tolerate electrical stimulations over the femoral nerve. Peak-

to-peak amplitude of both H-reflex and M-wave was measured and averaged for each

stimulation intensity. The highest amplitude of M-wave and H-reflex were labelled H_{MAX} and

M_{MAX} respectively. The H_{MAX}/M_{MAX} ratio was then calculated. In one subject it was not possible

to evoke the H-reflex at rest and was excluded from H_{MAX}-related parameters analysis at rest.

The amplitude of the M-wave accompanying H_{MAX} (M_{atHMAX}) was also measured and normalized

by M_{MAX} to verify that stimulation conditions were similar between sessions. The RMS of VM

EMG signal was measured over the 500 ms preceding each H_{MAX} (RMS_H) and M_{MAX} (RMS_M)

electrical stimulation during the contraction. We calculated the H_{MAX}/RMS_H ratio to assess the

influence of muscle activity background on spinal excitability (55,56). We also measured H_{MAX}

and M_{MAX} latencies at rest ($H_{LATENCY}$ and $M_{LATENCY}$) as the delay between the onset of stimulation

artefact and the upshot of electrophysiological response. While $M_{LATENCY}$ represents the time

required for conduction down the motor axons to the terminal branches and propagation

across the neuromuscular junction, $H_{LATENCY}$ reflects the time required for signal propagation

through the reflex arc, including the Ia afferent synaptic delay at the spinal level. Moreover, the

difference between $H_{LATENCY}$ and $M_{LATENCY}$ (H-$M_{LATENCY}$), representing an index of nerve

conduction velocity through the spinal reflex loop, was computed (13). Then, these latencies

were normalized by the subject's femur length (i.e., the distance between the greater

trochanter and the lateral condyle). For one subject, the onset of the VM H_{MAX} was merged

with the associated M-wave, thus this subject was excluded from H_{MAX} latency analysis. The

amplitude of the mechanical twitch associated with both M_{MAX} (M_{TW}) and H_{MAX} at rest was

measured. Since the M_{TW} represents the mechanical response of the muscle when the

maximum amount of the motor units is recruited (57), its characteristics were analyzed. For

each M_{TW}, we determined the contraction time (i.e., the time between twitch onset and

maximal twitch value; CT), the rate of force development (i.e., the maximal value of the first

derivative of the force signal using a 1 ms sliding window; RFD) and the half relaxation time (i.e.,

corresponding to the time needed to reach a 50% decrease of twitch force; HRT) before

computing the averages of the four measurements. To assess the electromechanical efficiency,

the M_{TW}/M_{MAX} was calculated. We analyzed the relationship between the amplitude of the T-

reflex and the corresponding twitch since we did not have the complete T-reflex recruitment

curve and therefore were not certain of obtaining the maximal T-wave response. The

relationship between VM T-reflex amplitudes, expressed as a percentage of M_{MAX}, and the

associated twitch amplitudes were scatter-plotted individually. The slope and the intercept of

the linear regression of these relationships were determined for each subject and averaged at

LS0 and LS10.

Statistical analysis. The normality of each dataset was assessed with the Shapiro-Wilk test. For

all variables that passed normality test, Student's paired t-test was applied in order to

investigate whether difference between LS0 and LS10 was present within the considered

parameters. When normality test was not passed, non-parametric Wilcoxon's signed-rank test

for paired comparison was used. Only for the RMS/M_{MAX} variable, a two-way repeated

measures ANOVA with *time* (LS0 *vs*. LS10) and *stimulation* (H_{MAX} *vs.* M_{MAX}) factors was

performed. If a significant effect was found, a Tukey's post-hoc analysis was conducted.

Correlations between MVC and qACSA as well as between MVC and CAR_{MVC} were analyzed

using repeated measures correlations (rmcorr R package, 51). Effect sizes (ES) were calculated

with Cohen's d and r respectively for parametric and non-parametric comparison (59). Small,

moderate and large effects are considered for Cohen's d ≥ 0.2, ≥ 0.5, ≥ 0.8, and Cohen's r ≥ 0.1,

≥ 0.3, ≥ 0.5 respectively. Results are reported as mean ± standard deviation. Significance level

was set at $p < 0.05$ and the confidence interval at 95% ($CI_{95\%}$). Student's and Wilcoxon's tests

were performed using Statistica software (Statsoft, version 12, Tulsa, OK, USA; RRID:

SCR_014213) while Cohen's effect sizes were calculated with G*Power software (version

3.1.9.2, Universität Düsseldorf, Germany; RRID: SCR_013726). Repeated measures correlations

were performed with R (version 4.1.3, R Core Team, 2017; RRID: SCR_001905). Graphs were

made with GraphPad Prism software (version 8.4.0, GraphPad software Inc. San Diego, CA, USA;

RRID: SCR_002798).

RESULTS

Muscle morphology, strength and activation capacity.

The ACSA of the unloaded VM ($t_{(9)}$ = 4.042, $CI_{95\%}$ = [0.48 : 1.69], p = 0.003, ES = 0.426)

and quadriceps ($t_{(9)}$ = 3.444, $CI_{95\%}$ = [0.87: 4.19], p = 0.007, ES = 0.397) decreased (-7.8 and -4.4

%, respectively) after 10 days of ULLS (Table 1). The ACSA losses were accompanied by a 27.1

% reduction in knee extensors MVC ($t_{(9)}$ = 9.464, $CI_{95\%}$ = [17.01 : 27.70], p < 0.0001, ES = 2.196)

(Table 1). The normalized force (MVC/qACSA) was 23.7 % lower after ULLS ($t_{(9)}$ = 8.792, $CI_{95\%}$ =

[0.25 : 0.43], p = 0.0001, ES = 2.016) (Table 1).

Activation capacity as measured by CAR_{MVC} decreased by 3.0 % after ULLS ($t_{(9)}$ = 3.557,

$CI_{95\%}$ = [0.01 : 0.05], p = 0.006, ES = 1.175) (Figure 1) while the $CAR_{10\%}$ decreased by 15.0 % ($t_{(9)}$

= 4.011, $CI_{95\%}$ = [2.79 : 10.03], p = 0.003, ES = 0.997). The ratio between the CAR_{MVC} and the

MVC increases by 33.4 % following ULLS ($t_{(9)}$ = -6.839, $CI_{95\%}$ = [-0.01 : -0.00], p < 0.0001, ES =

1.880), while the ratio between the $CAR_{10\%}$ and the force produced at 10% of MVC increased

by 16.5 % ($t_{(9)}$ = -2.540, $CI_{95\%}$ = [-0.02 : -0.00], p = 0.032, ES = 0.869) (Table 1). Furthermore, the

normalized EMG activity (RMS/M_{MAX}) of the VM during the 10% MVC plateau declined by 34.2%

after ULLS (only *time* effect; $F_{(1,8)}$ = 8.482, p = 0.020, η_p^2 = 0.515 ; Tukey post-hoc : p = 0.020, ES

= 0.839) (Table 1). Based on repeated measures correlation analysis, it appears that MVC is

positively correlated to qACSA (r_{rm} (9) = 0.773, $CI_{95\%}$ = [0.22 : 0.95], p = 0.005) (Figure 2A) and

CAR_{MVC} (r_{rm} (9) = 0.624, $CI_{95\%}$ = [-0.07 : 0.91], p = 0.040) (Figure 2B). In other words, the higher

the qACSA or CAR_{MVC}, the greater the production of maximal force. However, MVC change was

not correlated to changes in CAR_{MVC} (r = -0.382, r^2 = 0.146, p = 0.276) and qACSA (r = 0.189, r^2

= 0.036, p = 0.600) (see Supplemental Figure S1).

Muscle Contractile Properties

Mechanical parameters associated with H_{MAX} and M_{MAX} stimulations at rest were also

compared between LS0 and LS10. The ULLS did not modify the twitch amplitude associated

with M_{MAX} (M_{TW}; $t_{(8)}$ = 1.593, $CI_{95\%}$ = [-0.69 : 3.75], p = 0.150, ES = 0.599) (Figure 3). Specifically,

the RFD of M_{TW} was not altered after 10 days of ULLS ($t_{(8)}$ = 1.685, $CI_{95\%}$ = [-0.01 : 0.09], p =

0.130, ES = 0.678), nor were the CT ($t_{(8)}$ = -1.159, $CI_{95\%}$ = [-14.61 : 4.84], p = 0.280, ES = 0.420)

and the HRT ($t_{(8)}$ = 0.165, $CI_{95\%}$ = [-15.05 : 17.36], p = 0.873, ES = 0.070). Also, the M_{TW}/M_{MAX}

ratio was not changed by the ULLS protocol (Z = 0.770, p = 0.441, ES = 0.257). Muscle contractile

properties obtained at rest are provided in Table 2. Regarding the twitch associated with H_{MAX},

its amplitude was not different between LS0 and LS10 ($t_{(7)}$ = 1.015, $CI_{95\%}$ = [-0.84 : 2.11], p =

0.344, ES = 0.253) (Figure 3).

H-Reflex responses and M-wave

While the H_{MAX} at rest was increased by 59.0% after 10 days of ULLS (Z = 2.100, p =

0.036, ES = 0.743), H_{MAX} obtained during muscle contractions at 10% MVC was not modified (Z

= 0.770, p = 0.441, ES = 0.257) (Table 3). It should be noted that the amplitude of the M-wave

associated with H_{MAX} (M_{atHMAX}) was not different between LS0 and LS10 either at rest (Z = 0.700,

p = 0.484, ES = 0.247) or during contraction (Z = 1.362, p = 0.173, ES = 0.454) (Table 3).

Therefore, the stimulation conditions at H_{MAX} may be considered stable, allowing the

comparison of spinal excitability features between the two time points. M_{MAX} did not differ

between LS0 and LS10 either at rest ($t_{(8)}$ = -0.258, $CI_{95\%}$ = [-1.83 : 1.46], p = 0.803, ES = 0.025)

or during 10% MVC contraction ($t_{(8)}$ = 1.345, $CI_{95\%}$ = [-0.42 : 1.59], p = 0.215, ES = 0.074) (Table

3). The ULLS increased the H_{MAX}/M_{MAX} at rest by 33.1 % (LS0 : 0.20 ± 0.15 *vs.* LS10 : 0.27 ± 0.15

; $t_{(7)}$ = -2.557, $CI_{95\%}$ = [-0.13 : -0.01], p = 0.038, ES = 0.441) but had no impact on the ratio during

contraction at 10% MVC (LS0 : 0.34 ± 0.16 *vs.* LS10: 0.36 ± 0.17; $t_{(8)}$ = -1.947, $CI_{95\%}$ = [-0.04 :

0.00], *p* = 0.087, ES = 0.121) (Figure 4A). This result is also depicted in Figure 4B where

experimental traces of M_{MAX} and H_{MAX} obtained at rest, before and after ULLS, from one

representative subject, are presented. In order to overcome differences between sessions in

EMG activity during contractions at 10% MVC, we also normalized H_{MAX} by muscle activity

background preceding H_{MAX} stimulations (RMS_H), and we observed an increase of this ratio after

the ULLS period (+102.9 % ; $t_{(8)}$ = -2.480, $CI_{95\%}$ = [-74.00 : -2.70], *p* = 0.038, ES = 1.097) (Table 3

and Figure 4C).

Moreover, the delay between electrical stimulation and electrophysiological responses

was determined for the M_{MAX} and H_{MAX} waves. The latency of M_{MAX} ($M_{LATENCY}$) and H_{MAX} ($H_{LATENCY}$)

was not affected by ULLS ($M_{LATENCY}$: $t_{(8)}$ = -0.527, $CI_{95\%}$ = [-1.75 : 1.10], *p* = 0,613, ES = 0.192;

$H_{LATENCY}$: $t_{(6)}$ = -0.913, $CI_{95\%}$ = [-2.53 : 1.15], *p* = 0,396, ES = 0.134) (Table 3). The difference

between $H_{LATENCY}$ and $M_{LATENCY}$ (H-$M_{LATENCY}$) was not changed after 10 days of ULLS ($t_{(6)}$ = 0.289,

$CI_{95\%}$ = [-1.73 : 2.20], *p* = 0.782, ES = 0.049).

In addition, the ULLS did not modulate the relationship between the VM T-reflex and

the associated twitch. Indeed, the unloading period did not alter the slope ($t_{(5)}$ = -0.874, $CI_{95\%}$ =

[-0.31 : 0.15], p = 0.422, ES = 0.381) and the intercept ($t_{(5)}$ = -1.397, $CI_{95\%}$ = [-1.91 : 0.57], *p* =

0.221, ES = 0.946) of this linear regression (Table 3).

DISCUSSION

The present study aimed to investigate neural, contractile and morphological changes associated with muscle weakness following 10 days of ULLS of the main antigravity muscle group of the lower limbs. We showed that knee extensors' disuse led to a reduction in MVC mainly explained by an impairment in central motor drive. For the first time, spinal excitability was investigated in the quadriceps muscle group to deepen knowledge of neural adaptation to unloading in antigravity muscles. Our main results show that spinal excitability was enhanced at rest and during muscle contraction following 10 days of ULLS. These findings seem particularly relevant as they provide new insights into the neurophysiological adjustments that occur in a major antigravity muscle in response to human muscle disuse.

Muscle morphology and function.

The present investigation shows that 10 days of ULLS reduced the maximal force developed by the knee extensors (-27 %, -2.7 %·day^{-1}). This alteration in MVC was associated with a reduction in qACSA of 0.44 %·day^{-1} (-4.4 % in 10 days). In overall our results are consistent with previous investigations that reported an impairment in force production capacity (11,14–16,23,25,60) as well as a decrease in knee extensors size after ULLS (8,11,16,23)of the knee extensors . As reported in the literature, muscle atrophy could be explained by decreases in protein synthesis rate after lower limb muscle disuse (61,62) affecting the muscle protein synthesis/breakdown balance (63) and thus, the myofibrillar content (64). The present study corroborated, in agreement with earlier investigations (8,16,65–67), that strength loss, after unloading, was greater than the rate of muscle size loss, therefore resulting in an alteration of normalized force (-24 %). It is thought that muscular and tendinous alterations are involved, such as a reduction in myosin content (16,68) and tendon stiffness (11), and an impairment of

the excitation-contraction coupling (67). Neural alterations, including a reduction in voluntary

activation and coordination of antagonistic muscles (69), could also be potential explanatory

factors contributing to the impairment of normalized force during muscle disuse. In the current

study, we observed an impairment of both maximal and submaximal central motor drive, as

revealed by the decrease in CAR_{MVC} (-3 %), $CAR_{10\%}$ (-15 %) and RMS/M_{MAX} (-34 %). Particularly,

a decrease in CAR_{MVC} was associated with a reduction in maximal force production capacity. In

addition, our data showed a rise in CAR_{MVC}/MVC and $CAR_{10\%}$/10% MVC ratios after the ULLS

protocol, suggesting an impairment in muscle electromechanical efficiency (i.e. the dissociation

between electrical and mechanical events of muscle function; 68) after knee extensors

unloading. Though our findings are consistent with some previous investigations on knee

extensors and plantar flexors after 21-30 days of limb suspension (23,29,60,71), some others

did not observed any change in maximal muscle activation, for the same muscle groups, after

10-30 days of limb unloading (11,13,16,25,30,72). Despite these conflicting results, strength

loss after unloading seems to be more related to alterations in neural than muscular factors

(13,73). As a matter of fact, among multiple predictive variables, including muscle ACSA, twitch

RFD and HRT or the amplitude of the H-reflex and M-wave for instance, it appears that central

activation is the most influential variable in mediating strength loss after muscle disuse (13).

Regarding mechanical twitch evoked by H_{MAX} stimulation at rest, no change was

reported following 10 days of ULLS in the current study. The present observation contrasts with

the result of Seynnes *et al.* (29) which reported a 40 and 54% increase in H_{MAX} twitch amplitude

for the plantar flexors after 14 and 23 days of ULLS respectively. This discrepancy might depend

on the muscle group investigated and the different duration of exposure to limb unloading. On

the other hand, in agreement with data from Campbell *et al.* (16) following 21 days of ULLS, the

current analysis, did not find any change in M_{TW}. However, it should be noted that there is no

consensus in the literature about the evolution of M_{TW} following muscle disuse, since studies

have shown that ULLS decreased (15,60,65) or even increased (30) the maximal twitch

amplitude on lower limb muscles. In the current study, the M_{TW} parameters (CT, RFD and HRT)

were unaltered after the knee extensors' unloading. These outcomes are in accordance with

the literature focusing on the muscle contractility of knee extensors following 21-30 days of

ULLS (16,30,60). Specifically, lower limb suspension did not result in the alteration of M_{TW} CT

(16), RFD (16,30,60) and HRT (16,30,60). It should be noted that we evoked a twitch to assess

muscular factors of knee extensors. The sensitivity of this index could not be sufficient to detect

changes in muscle contractility (74). Despite this, the present findings underline the

discrepancy between morphological (i.e. reduction in qACSA, maximal strength) and functional

(i.e. no change in quadriceps contractility) data already observed in the literature (8) and

suggest that short-term ULLS-induced strength loss is primarily attributed to neural factors.

Therefore, the present model of disuse does not seem to impair knee extensors' contractility

and relaxation processes, although it is effective in inducing strength loss through impairments

in muscle activation. Due to the relatively small number of data obtained for muscle

contractility outcomes (n = 9), further future studies will be required to confirm these findings.

H-Reflex responses and M-wave

As observed at rest by de Boer *et al.* (11) following 23 days of ULLS, our protocol did not

modify the maximal M-wave obtained at rest, and during a weak knee extensor contraction

(10% MVC). This result indicates that sarcolemmal excitability and potential action propagation

were not altered by limb unloading (75,76).

In the present study, unloading increased the spinal excitability of the VM at rest, as

revealed by the rise of 33% in H_{MAX}/M_{MAX}. Although there is no available data on knee extensors,

this finding is consistent with numerous earlier studies that have shown an increase in spinal

excitability of plantar flexors after 23-28 days of ULLS (7,8,12–14,29,30). Our results showed

that H_{MAX}/M_{MAX} obtained during a weak contraction (10% MVC) was not altered by ULLS. Since

presynaptic inhibitions are already partially released during voluntary contraction, the

possibility of observing an increase in the H_{MAX}/M_{MAX} ratio following the ULLS is limited in this

condition. However, it should be noted that the background EMG activity (i.e., the RMS of VM

EMG preceding H_{MAX} stimulation, RMS/M_{MAX}) was different between LS0 and LS10. To

overcome that limit, the H_{MAX} was normalized by the RMS_H (H_{MAX}/RMS_H). The latter parameter

appeared to be 2-fold greater after the 10 days of ULLS, meaning that, for a similar muscle

electrical activity, the amplitude of the maximal H-reflex was increased. In light of these results,

it appears that unloading induced an enhancement of H-reflex amplitude both at rest and

during contraction. As shown by Delwaide (34), this enhancement could be ascribed to an

increase in motoneuronal excitability and/or a decrease in presynaptic inhibition. Particularly,

the latter explanation has been confirmed in an immobilization model of disuse (28). In their

study, Lundbye-Jensen & Nielsen (28) showed that after immobilization of the left foot and

ankle joint by a cast for 14 days, the increase in *soleus* H-reflex results from decreased

presynaptic inhibition of the Ia afferents. More specifically, the authors reported a reduction in

the primary afferent depolarisation mechanism (i.e., a decrease in the inhibitory activity of a

spinal GABAergic interneuron) and a reduction in homosynaptic post-activation depression

(i.e., a decreased quantity of neurotransmitters available in the synapse following repeated

solicitation of the Ia afferents).

The present analysis showed that the decrease in muscle activation after a short period

of ULLS is accompanied by an increased spinal excitability. The latter could be viewed as a

compensatory adaptation, implemented by the nervous system to counterbalance the reduced

capacity to activate knee extensors. However, this adaptation is not sufficient to counteract the

entire cortical changes that occur during muscle disuse. Additionally, it seems that the increase

in the H-reflex following ULLS is not related to the activation capacity, as reported by Seynnes

et al. (30) on the plantar flexors after 24 days of limb unloading. Indeed, the authors did not

observe a significant relationship between relative changes in amplitude of resting H-reflex and

V-wave obtained during MVC over the ULLS period, indicating that the factors modulating the

former had a limited effect on the latter (30). The monosynaptic T-reflex of knee extensors,

evoked by patellar tendon striking, was also used as a supplementary tool to investigate the

modulation of spinal circuitry following knee extensors unloading. We observed no change in

the relationship between the T-reflex of the VM and the associated twitch, even if the spinal

excitability was enhanced. This result suggests that other factors that affect T-reflex, including

the mechanical properties of the tendon-aponeurosis and the sensitivity of the muscle spindle,

have probably been altered in such a way as to mask the effect of increased spinal excitability

on T-reflex (29). In contrast to our results, Seynnes *et al.* (29) reported an increase in the slope

and X-intercept of the T-reflex recruitment curve. Likely, the discrepancy between the present

and earlier findings may likely be ascribed to the muscle group investigated (knee extensors *vs.*

plantar flexors) and the ULLS duration (10 *vs.* 23 days). However, these interpretations should

be taken with caution because of the limited number of subjects (n = 6) that underwent the T-

reflex testing procedure during the present investigation.

In the current study, neither $H_{LATENCY}$, $M_{LATENCY}$ nor H-$M_{LATENCY}$ were affected by 10 days

of lower limb suspension, suggesting that nerve conduction velocity through the reflex arc and

synaptic delay at the spinal level were not impaired by unloading. This outcome is in contrast

to those of Clark *et al.* (13) and Seynnes *et al.* (30) who found an increase in soleus M_{MAX} and

H_{MAX} latencies after 28 and 24 days of ULLS respectively. More precisely, since the H-$M_{LATENCY}$,

an index of conduction velocity through the spinal reflex loop, was not changed in their studies

(13,30), authors primarily attributed the slowed conduction of H-reflex and M-wave responses

to the conduction velocity in the supply nerves, the branching axon terminals, and/or

transmission across the neuromuscular junction (77). No change in nerve conduction through

the reflex arc occurred after 10 days of ULLS for the lower limb muscles, and confirm similar

evidences from Manganotti *et al*. (78) who used the bed rest model during an identical period

of disuse. Further studies using ULLS protocols exceeding 10 days are needed to better

understand the modulation kinetics of nerve conduction in knee extensors.

In conclusion, the present study was the first to investigate the influence of short-

duration (10 days) ULLS on the neuromuscular function and spinal excitability of knee

extensors. The strength loss of knee extensors, observed after 10 days of lower limb unloading,

was mainly attributed to neural impairments, such as a reduction in central activation. To

compensate for reduced peripheral neurosensory inputs during the ULLS protocol, the spinal

excitability of the VM was enhanced at rest and during a weak knee extension (10% MVC) after

the ULLS protocol. These findings are paramount since they bring out new insights into the

neural adaptations that occur in the major antigravity muscle in response to muscle disuse.

Understanding the physiological adaptations during ULLS can guide the development of

effective interventions to improve overall health outcomes for individuals experiencing muscle

inactivity. For instance, using tendon vibrations (79,80) could be relevant to stimulate Ia

afferents in patients immobilised with a cast to improve spinal regulation of neural inputs (81)

and maintain efficient control of force production (82,83).

ACKNOWLEDGMENTS

The authors thank all volunteers who participated in the study.

GRANTS

The present study was funded by the Italian Space Agency (ASI), MARS-PRE, Project, n. DC-

VUM-2017-006.

DATA AVAILABILITY

Data will be made available on reasonable request.

AUTHOR CONTRIBUTIONS

L.L., M.V.N., A.M., G.D., F.S. and G.S. conceived and designed research; L.L., F.S and G.S

performed experiments; L.L. and F.S. analyzed data; L.L., M.V.N, A.M., G.D., F.S. and G.S.

interpreted results of experiments; L.L. prepared figures; L.L. drafted manuscript; L.L., M.V.N,

A.M., G.D., F.S. and G.S. edited and revised manuscript; L.L., M.V.N, A.M., G.D., F.S. and G.S.

approved final version of manuscript.

SUPPLEMENTARY DATA

Supplementary data: https://doi.org/10.6084/m9.figshare.26779975

REFERENCES

1. Nash CE, Mickan SM, Del Mar CB, Glasziou PP. Resting injured limbs delays recovery: a systematic review. J Fam Pract. 2004 Sep;53(9):706–12.

2. Lin CWC, Donkers NAJ, Refshauge KM, Beckenkamp PR, Khera K, Moseley AM. Rehabilitation for ankle fractures in adults. Cochrane Database Syst Rev. 2012 Nov 14;11:CD005595.

3. Graham ZA, Lavin KM, O'Bryan SM, Thalacker-Mercer AE, Buford TW, Ford KM, et al. Mechanisms of exercise as a preventative measure to muscle wasting. Am J Physiol Cell Physiol. 2021 Jul 1;321(1):C40–57.

4. Evans WJ. Skeletal muscle loss: cachexia, sarcopenia, and inactivity. Am J Clin Nutr. 2010 Apr;91(4):1123S-1127S.

5. Sarto F, Bottinelli R, Franchi MV, Porcelli S, Simunič B, Pišot R, et al. Pathophysiological mechanisms of reduced physical activity: Insights from the human step reduction model and animal analogues. Acta Physiologica. 2023;238(3):e13986.

6. Lee PHU, Chung M, Ren Z, Mair DB, Kim DH. Factors mediating spaceflight-induced skeletal muscle atrophy. American Journal of Physiology-Cell Physiology. 2022 Mar;322(3):C567–80.

7. Hackney KJ, Ploutz-Snyder LL. Unilateral lower limb suspension: integrative physiological knowledge from the past 20 years (1991-2011). Eur J Appl Physiol. 2012 Jan;112(1):9–22.

8. Campbell M, Varley-Campbell J, Fulford J, Taylor B, Mileva KN, Bowtell JL. Effect of Immobilisation on Neuromuscular Function In Vivo in Humans: A Systematic Review. Sports Med. 2019 Jun;49(6):931–50.

9. Tesch PA, Lundberg TR, Fernandez-Gonzalo R. Unilateral lower limb suspension: From subject selection to "omic" responses. J Appl Physiol (1985). 2016 May 15;120(10):1207–14.

10. Booth FW. Effect of limb immobilization on skeletal muscle. J Appl Physiol Respir Environ Exerc Physiol. 1982 May;52(5):1113–8.

11. de Boer MD, Maganaris CN, Seynnes OR, Rennie MJ, Narici MV. Time course of muscular, neural and tendinous adaptations to 23 day unilateral lower-limb suspension in young men. J Physiol. 2007 Sep 15;583(Pt 3):1079–91.

12. Clark BC. In vivo alterations in skeletal muscle form and function after disuse atrophy. Med Sci Sports Exerc. 2009 Oct;41(10):1869–75.

13. Clark BC, Manini TM, Bolanowski SJ, Ploutz-Snyder LL. Adaptations in human neuromuscular function following prolonged unweighting: II. Neurological properties and motor imagery efficacy. J Appl Physiol (1985). 2006 Jul;101(1):264–72.

14. Clark BC, Pierce JR, Manini TM, Ploutz-Snyder LL. Effect of prolonged unweighting of human skeletal muscle on neuromotor force control. Eur J Appl Physiol. 2007 Apr 12;100(1):53–62.

15. Horstman AM, de Ruiter CJ, van Duijnhoven NTL, Hopman MTE, de Haan A. Changes in muscle contractile characteristics and jump height following 24 days of unilateral lower limb suspension. Eur J Appl Physiol. 2012 Jan;112(1):135–44.

16. Campbell EL, Seynnes OR, Bottinelli R, McPhee JS, Atherton PJ, Jones DA, et al. Skeletal muscle adaptations to physical inactivity and subsequent retraining in young men. Biogerontology. 2013 Jun;14(3):247–59.

17. Cook SB, Kanaley JA, Ploutz-Snyder LL. Neuromuscular function following muscular unloading and blood flow restricted exercise. Eur J Appl Physiol. 2014;114(7):1357–65.

18. Sarto F, Stashuk DW, Franchi MV, Monti E, Zampieri S, Valli G, et al. Effects of short-term unloading and active recovery on human motor unit properties, neuromuscular junction transmission and transcriptomic profile. The Journal of Physiology. 2022;600(21):4731–51.

19. Dudley GA, Duvoisin MR, Adams GR, Meyer RA, Belew AH, Buchanan P. Adaptations to unilateral lower limb suspension in humans. Aviat Space Environ Med. 1992 Aug;63(8):678–83.

20. Deschenes MR, Giles JA, McCoy RW, Volek JS, Gomez AL, Kraemer WJ. Neural factors account for strength decrements observed after short-term muscle unloading. Am J Physiol Regul Integr Comp Physiol. 2002 Feb;282(2):R578-583.

21. Deschenes MR, Holdren AN, McCoy RW. Adaptations to short-term muscle unloading in young and aged men. Med Sci Sports Exerc. 2008 May;40(5):856–63.

22. Deschenes MR, McCoy RW, Mangis KA. Factors relating to gender specificity of unloading-induced declines in strength. Muscle Nerve. 2012 Aug;46(2):210–7.

23. Schulze K, Gallagher P, Trappe S. Resistance training preserves skeletal muscle function during unloading in humans. Med Sci Sports Exerc. 2002 Feb;34(2):303–13.

24. Tesch PA, Trieschmann JT, Ekberg A. Hypertrophy of chronically unloaded muscle subjected to resistance exercise. J Appl Physiol (1985). 2004 Apr;96(4):1451–8.

25. Berg HE, Tesch PA. Changes in muscle function in response to 10 days of lower limb unloading in humans. Acta Physiologica Scandinavica. 1996 May;157(1):63–70.

26. Duchateau J. Bed rest induces neural and contractile adaptations in triceps surae. Med Sci Sports Exerc. 1995 Dec;27(12):1581–9.

27. Anderson J, Almeida-Silveira MI, Perot C. Reflex and muscular adaptations in rat soleus muscle after hindlimb suspension. Journal of Experimental Biology. 1999 Oct 1;202(19):2701–7.

28. Lundbye-Jensen J, Nielsen JB. Immobilization induces changes in presynaptic control of
group Ia afferents in healthy humans. J Physiol. 2008 Sep 1;586(Pt 17):4121–35.

29. Seynnes OR, Maffiuletti NA, Maganaris CN, de Boer MD, Pensini M, di Prampero PE, et al.
Soleus T reflex modulation in response to spinal and tendinous adaptations to unilateral
lower limb suspension in humans. Acta Physiol (Oxf). 2008 Nov;194(3):239–51.

30. Seynnes OR, Maffiuletti NA, Horstman AM, Narici MV. Increased H-reflex excitability is not
accompanied by changes in neural drive following 24 days of unilateral lower limb
suspension. Muscle Nerve. 2010 Nov;42(5):749–55.

31. Schieppati M. The Hoffmann reflex: a means of assessing spinal reflex excitability and its
descending control in man. Prog Neurobiol. 1987;28(4):345–76.

32. Duchateau J, Enoka RM. Neural control of lengthening contractions. Lindstedt SL,
Hoppeler HH, editors. Journal of Experimental Biology. 2016 Jan 1;219(2):197–204.

33. Grosprêtre S, Martin A. H reflex and spinal excitability: methodological considerations. J
Neurophysiol. 2012 Mar;107(6):1649–54.

34. Delwaide P. Human monosynaptic reflexes and presynaptic inhibition. In: Desmedt JE,
editor. Basel: Karger; 1973. p. 508–22. (New developments in electromyography and
clinical neurophysiology; vol. 3).

35. Paillard J. [Electrophysiologic analysis and comparison in man of Hoffmann's reflex and
myotatic reflex]. Pflugers Arch Gesamte Physiol Menschen Tiere. 1955;260(6):448–79.

36. Paillard J. Functional organization of afferent innervation of muscle studied in man by
monosynaptic testing. Am J Phys Med. 1959 Dec;38:239–47.

37. Józsa L, Kvist M, Kannus P, Järvinen M. The effect of tenotomy and immobilization on
muscle spindles and tendon organs of the rat calf muscles. A histochemical and
morphometrical study. Acta Neuropathol. 1988;76(5):465–70.

38. Valli G, Sarto F, Casolo A, Del Vecchio A, Franchi MV, Narici MV, et al. Lower limb
suspension induces threshold-specific alterations of motor units properties that are
reversed by active recovery. Journal of Sport and Health Science. 2024 Mar;13(2):264–76.

39. Bleeker MWP, Hopman MTE, Rongen GA, Smits P. Unilateral lower limb suspension can
cause deep venous thrombosis. Am J Physiol Regul Integr Comp Physiol. 2004
Jun;286(6):R1176-1177.

40. Roach REJ, Cannegieter SC, Lijfering WM. Differential risks in men and women for first and
recurrent venous thrombosis: the role of genes and environment. J Thromb Haemost.
2014 Oct;12(10):1593–600.

41. Wanner M, Hartmann C, Pestoni G, Martin BW, Siegrist M, Martin-Diener E. Validation of
the Global Physical Activity Questionnaire for self-administration in a European context.
BMJ Open Sport & Exercise Medicine. 2017 May 1;3(1):e000206.

42. Guette M, Gondin J, Martin A. Time-of-day effect on the torque and neuromuscular
properties of dominant and non-dominant quadriceps femoris. Chronobiol Int.
2005;22(3):541–58.

43. Douglas CM, Hesketh SJ, Esser KA. Time of Day and Muscle Strength: A Circadian Output?
Physiology (Bethesda). 2021 Jan 1;36(1):44–51.

44. Hopkins JT, Ingersoll CD, Andrew Krause B, Edwards JE, Cordova ML. Effect of knee joint
effusion on quadriceps and soleus motoneuron pool excitability: Medicine and Science in
Sports and Exercise. 2001 Jan;123–6.

45. Zehr E. Considerations for use of the Hoffmann reflex in exercise studies. European
journal of applied physiology. 2002 May 1;86:455–68.

46. Mau-Moeller A, Behrens M, Lindner T, Bader R, Bruhn S. Age-related changes in
neuromuscular function of the quadriceps muscle in physically active adults. Journal of
Electromyography and Kinesiology. 2013 Jun 1;23(3):640–8.

47. Doguet V, Jubeau M. Reliability of H-reflex in vastus lateralis and vastus medialis muscles
during passive and active isometric conditions. Eur J Appl Physiol. 2014
Dec;114(12):2509–19.

48. Burke D, ADAMS RW, SKUSE NF. The effects of voluntary contraction on the H reflex of
human limb muscles. Brain. 1989 Apr 1;112(2):417–33.

49. Knikou M. The H-reflex as a probe: pathways and pitfalls. J Neurosci Methods. 2008 Jun
15;171(1):1–12.

50. Funase K, Miles TS. Observations on the variability of the H reflex in human soleus. Muscle
& Nerve. 1999;22(3):341–6.

51. McNeil CJ, Butler JE, Taylor JL, Gandevia SC. Testing the excitability of human
motoneurons. Frontiers in Human Neuroscience [Internet]. 2013 [cited 2019 Dec 10];7.
Available from: http://journal.frontiersin.org/article/10.3389/fnhum.2013.00152/abstract

52. Monti E, Franchi MV, Badiali F, Quinlan JI, Longo S, Narici MV. The Time-Course of
Changes in Muscle Mass, Architecture and Power During 6 Weeks of Plyometric Training.
Frontiers in Physiology [Internet]. 2020 [cited 2022 Nov 3];11. Available from:
https://www.frontiersin.org/articles/10.3389/fphys.2020.00946

53. Maffiuletti NA, Pensini M, Scaglioni G, Ferri A, Ballay Y, Martin A. Effect of
electromyostimulation training on soleus and gastrocnemii H- and T-reflex properties. Eur
J Appl Physiol. 2003 Nov;90(5–6):601–7.

54. Gandevia SC, Allen GM, Butler JE, Taylor JL. Supraspinal factors in human muscle fatigue:
evidence for suboptimal output from the motor cortex. J Physiol. 1996;8.

55. Theodosiadou A, Henry M, Duchateau J, Baudry S. Revisiting the use of Hoffmann reflex in
motor control research on humans. Eur J Appl Physiol. 2023 Apr;123(4):695–710.

56. Palmieri-Smith R, Ingersoll C, Hoffman M. The Hoffmann Reflex: Methodologic
Considerations and Applications for Use in Sports Medicine and Athletic Training
Research. Journal of athletic training. 2004 Feb 1;39:268–77.

57. Nakagawa K, Fok KL, Masani K. Neuromuscular recruitment pattern in motor point
stimulation. Artificial Organs. 2023;47(3):537–46.

58. Bakdash JZ, Marusich LR. Repeated Measures Correlation. Front Psychol [Internet]. 2017
[cited 2022 Mar 31];0. Available from:
https://www.frontiersin.org/articles/10.3389/fpsyg.2017.00456/full

59. Fritz CO, Morris PE, Richler JJ. Effect size estimates: Current use, calculations, and
interpretation. Journal of Experimental Psychology: General. 2012;141(1):2–18.

60. Cook SB, Brown KA, Deruisseau K, Kanaley JA, Ploutz-Snyder LL. Skeletal muscle
adaptations following blood flow-restricted training during 30 days of muscular unloading.
J Appl Physiol (1985). 2010 Aug;109(2):341–9.

61. de Boer MD, Selby A, Atherton P, Smith K, Seynnes OR, Maganaris CN, et al. The temporal
responses of protein synthesis, gene expression and cell signalling in human quadriceps
muscle and patellar tendon to disuse. J Physiol. 2007 Nov 15;585(Pt 1):241–51.

62. Brook MS, Stokes T, Gorissen SHM, Bass JJ, McGlory C, Cegielski J, et al. Declines in muscle
protein synthesis account for short-term muscle disuse atrophy in humans in the absence
of increased muscle protein breakdown. J Cachexia Sarcopenia Muscle. 2022
Aug;13(4):2005–16.

63. Deane CS, Piasecki M, Atherton PJ. Skeletal muscle immobilisation-induced atrophy:
mechanistic insights from human studies. Clinical Science. 2024 Jun 19;138(12):741–56.

64. Brocca L, Longa E, Cannavino J, Seynnes O, De Vito G, McPhee J, et al. Human skeletal
muscle fibre contractile properties and proteomic profile: adaptations to 3 weeks of
unilateral lower limb suspension and active recovery: Impact of ULLS and active
rehabilitation on human muscle fibres. J Physiol. 2015 Dec 15;593(24):5361–85.

65. Clark BC, Fernhall B, Ploutz-Snyder LL. Adaptations in human neuromuscular function
following prolonged unweighting: I. Skeletal muscle contractile properties and applied
ischemia efficacy. J Appl Physiol (1985). 2006 Jul;101(1):256–63.

66. Demangel R, Treffel L, Py G, Brioche T, Pagano AF, Bareille MP, et al. Early structural and
functional signature of 3-day human skeletal muscle disuse using the dry immersion
model. J Physiol. 2017 Jul 1;595(13):4301–15.

67. Monti E, Reggiani C, Franchi MV, Toniolo L, Sandri M, Armani A, et al. Neuromuscular
junction instability and altered intracellular calcium handling as early determinants of
force loss during unloading in humans. The Journal of Physiology. 2021;599(12):3037–61.

68. D'Antona G, Pellegrino MA, Adami R, Rossi R, Carlizzi CN, Canepari M, et al. The effect of
ageing and immobilization on structure and function of human skeletal muscle fibres. J
Physiol. 2003 Oct 15;552(Pt 2):499–511.

69. Maganaris CN, Baltzopoulos V, Ball D, Sargeant AJ. In vivo specific tension of human
skeletal muscle. J Appl Physiol (1985). 2001 Mar;90(3):865–72.

70. Turnsek G, Paravlic AH. Electromechanical efficiency index of skeletal muscle and its
applicability: a systematic review. Front Bioeng Biotechnol. 2024;12:1398047.

71. Seynnes OR, Maganaris CN, de Boer MD, di Prampero PE, Narici MV. Early structural
adaptations to unloading in the human calf muscles. Acta Physiol (Oxf). 2008
Jul;193(3):265–74.

72. Hotta N, Ishida K, Sato K, Koike T, Katayama K, Akima H. The effect of intense interval
cycle-training on unloading-induced dysfunction and atrophy in the human calf muscle. J
Physiol Anthropol. 2011;30(1):29–35.

73. Kawakami Y, Akima H, Kubo K, Muraoka Y, Hasegawa H, Kouzaki M, et al. Changes in
muscle size, architecture, and neural activation after 20 days of bed rest with and without
resistance exercise. European Journal of Applied Physiology. 2001 Feb 16;84(1–2):7–12.

74. Blinks JR, Rüdel R, Taylor SR. Calcium transients in isolated amphibian skeletal muscle
fibres: detection with aequorin. J Physiol. 1978 Apr;277:291–323.

75. Fuglevand AJ, Zackowski KM, Huey KA, Enoka RM. Impairment of neuromuscular
propagation during human fatiguing contractions at submaximal forces. J Physiol (Lond).
1993 Jan;460:549–72.

76. Allen DG, Lamb GD, Westerblad H. Skeletal muscle fatigue: cellular mechanisms. Physiol
Rev. 2008 Jan;88(1):287–332.

77. Kim BJ, Date ES, Park BK, Choi BY, Lee SH. Physiologic changes of compound muscle action
potentials related to voluntary contraction and muscle length in carpal tunnel syndrome. J
Electromyogr Kinesiol. 2005 Jun;15(3):275–81.

78. Manganotti P, Buoite Stella A, Ajcevic M, di Girolamo FG, Biolo G, Franchi MV, et al.
Peripheral nerve adaptations to 10 days of horizontal bed rest in healthy young adult
males. American Journal of Physiology-Regulatory, Integrative and Comparative
Physiology. 2021 Sep;321(3):R495–503.

79. Nito M, Yoshimoto T, Hashizume W, Shindo M, Naito A. Vibration decreases the
responsiveness of Ia afferents and spinal motoneurons in humans. Journal of
Neurophysiology. 2021 Oct;126(4):1137–47.

80. Souron R, Baudry S, Millet GY, Lapole T. Vibration-induced depression in spinal loop
excitability revisited. J Physiol. 2019 Nov;597(21):5179–93.

81. Rudomin P, Romo R, Mendell LM. Presynaptic Inhibition and Neural Control. Oxford
University Press; 1998. 474 p.

82. Cook WA, Cangiano A. Presynaptic and postsynaptic inhibition of spinal motoneurons.
Journal of Neurophysiology. 1972 May;35(3):389–403.

83. Fink AJP, Croce KR, Huang ZJ, Abbott LF, Jessell TM, Azim E. Presynaptic inhibition of
spinal sensory feedback ensures smooth movement. Nature. 2014 May;509(7498):43–8.

FIGURE LEGENDS

Figure 1. Maximal force and corresponding central activation. The maximal voluntary

contraction (MVC) and corresponding central activation ratio (CAR$_{MVC}$) were individually scatter

plotted before (white circles) and after (grey circles) 10 days of unilateral lower limb suspension

(n = 10). Black (before limb suspension) and grey (after limb suspension) crosses represent the

mean of both scatter plots. ***p < 0.001 and $^{\$\$}p$ < 0.01 express significant differences in MVC

and CAR$_{MVC}$, respectively, between before and after lower limb suspension.

Figure 2. Individual repeated measures correlations (rmcorr) between muscle function

parameters and maximal force production (n = 10). Rmcorr was plotted between qACSA and

MVC (A), and between CAR and MVC (B). Observations from the same participant are given the

same colour, the lines correspond to the rmcorr fit for each participant. MVC = maximal

voluntary contraction; qACSA = quadriceps cross-sectional area; CAR = central activation ratio.

Figure 3. Twitch amplitude associated with H$_{MAX}$ and M$_{MAX}$. The twitch amplitude associated

with H$_{MAX}$ (n = 8) and M$_{MAX}$ (n = 9) was measured at rest before (white columns) and after (grey

columns) 10 days of unilateral lower limb suspension. Circles and lines represent individual data

evolution. H$_{MAX}$ = maximal H-reflex; M$_{MAX}$ = maximal M-wave.

Figure 4. Changes in spinal excitability after 10 days of ULLS. A. H_{MAX}/M_{MAX} ratio was calculated

before (white columns) and after (grey columns) 10 days of ULLS, either at rest ($n = 8$) and

during 10% MVC contractions ($n = 9$). Circles and lines represent individual data evolution. B.

H_{MAX} and M_{MAX} traces at rest, from a typical subject, before (red lines) and after (black lines)

ULLS. C. Representative data of VM EMG activity during the 10% MVC contraction with

associated H_{MAX} before (red line) and after (black line) ULLS from a typical subject. VM EMG =

electromyographic activity of vastus medialis; H_{MAX} = maximal H-reflex; M_{MAX} = maximal M-

wave. ULLS = unilateral lower limb suspension. *$p < 0.05$ express significant difference.

TABLES

Table 1. Muscle morphology, strength and activation capacity.

	BDC (n = 10)	LS9 (n = 10)	*p*-value
VM ACSA (cm²)	16.34 ± 2.78	15.26 ± 2.26	0.003
qACSA (cm²)	57.79 ± 6.54	55.26 ± 6.20	0.007
	LS0 *(n = 10)*	LS10 *(n = 10)*	*p*-value
MVC (kg)	82.54 ± 11.13	60.19 ± 9.13	< 0.0001
10% MVC (kg)	8.25 ± 1.11	6.02 ± 0.91	< 0.0001
MVC/qACSA (kg·cm^{-2})	1.43 ± 0.18	1.09 ± 0.16	0.0001
CAR_{MVC}	0.980 ± 0.013	0.950 ± 0.033	0.006
CAR_{MVC}/MVC (kg^{-1})	0.012 ± 0.002	0.016 ± 0.002	< 0.0001
$CAR_{10\%}$	0.428 ± 0.065	0.364 ± 0.063	0.003
$CAR_{10\%}$/10% MVC (kg^{-1})	0.052 ± 0.009	0.061 ± 0.011	0.032
RMS/M_{MAX} at 10% MVC (x10^{-3})	9.39 ± 4.05	6.18 ± 3.54	0.020

All variables were measured before (LS0) and after (LS10) 10 days of unilateral lower limb

suspension, except ACSA measured the day of ULLS beginning (BDC) and the day before the

ULLS end (LS9). MVC = knee extensors maximal voluntary contraction; VM = vastus medialis

muscle; ACSA = muscle anatomical cross-sectional area; qACSA = quadriceps ACSA; CAR_{MVC} and

$CAR_{10\%}$ = central activation ratio calculated during a MVC and during a contraction at 10% MVC;

RMS = root mean square; M_{MAX} = maximal M-wave. Values are reported as mean ± standard

deviation.

Table 2. Mechanical parameters associated with M_{MAX} before and after ULLS.

| | M_{MAX} stimulations | | p-value |
	LS0 (n = 9)	LS10 (n = 9)	
M_{TW}/M_{MAX} (kg/mV)	1.38 ± 1.07	1.33 ± 1.25	0.441
CT (ms)	82.44 ± 8.59	87.33 ± 14.03	0.280
RFD (kg/ms)	0.247 ± 0.060	0.210 ± 0.048	0.130
HRT (ms)	83.60 ± 15.15	82.44 ± 17.80	0.873

886 M_{MAX} = maximal M-wave; M_{TW} = twitch associated with M_{MAX}; CT = twitch contraction time; RFD

887 = twitch rate of force development; HRT = twitch half relaxation time; ULLS = unilateral lower

888 limb suspension; LS0 and LS10 = before and after 10 days of ULLS. Values are reported as mean

889 ± standard deviation.

891 Table 3. Spinal excitability and nerve conduction parameters before and after 10 days of ULLS.

| | Rest | | | Contraction | | |
	LS0	LS10	p-value	LS0	LS10	p-value
M_{atHMAX} (% M_{MAX})	8.39 ± 7.35	6.64 ± 8.65	0.484	9.02 ± 9.03	5.17 ± 4.63	0.173
H_{MAX} (mV)	2.22 ± 1.94	3.53 ± 3.25	0.036	4.29 ± 3.21	4.37 ± 3.44	0.441
M_{MAX} (mV)	13.12 ± 7.04	13.30 ± 7.41	0.803	13.48 ± 7.97	12.89 ± 7.91	0.215
H_{MAX}/RMS_H	-	-	-	37.27 ± 12.95	75.62 ± 47.69	0.038
$M_{LATENCY}$ (ms/m)	15.22 ± 1.60	15.74 ± 1.81	0.613	-	-	-
$H_{LATENCY}$ (ms/m)	48.04 ± 5.55	48.72 ± 5.55	0.396	-	-	-
$H-M_{LATENCY}$ (ms/m)	34.07 ± 5.34	33.84 ± 4.02	0.782	-	-	-
T-twitch slope	0.253 ± 0.096	0.331 ± 0.275	0.422	-	-	-
T-twitch intercept	0.876 ± 0.585	1.550 ± 0.807	0.221	-	-	-

Data were obtained at rest and/or during 10% MVC contractions. For each parameter, the

number of subjects was 9, except for M_{atHMAX} at rest (n = 8), $H_{LATENCY}$ and $H-M_{LATENCY}$ (n = 7), T-

twitch slope and intercept (n = 6). LS0 = before leg suspension; LS10 = after 10 days of leg

suspension; H_{MAX} = maximal H-reflex; M_{MAX} = maximal M-wave ; M_{atHMAX} = M-wave associated

with maximal H-reflex; $H_{LATENCY}$ and $M_{LATENCY}$ = H_{MAX} and M_{MAX} latency respectively; H-$M_{LATENCY}$ =

difference between $H_{LATENCY}$ and $M_{LATENCY}$; T = T-reflex; twitch = T-reflex associated twitch; MVC

= maximal voluntary contraction; ULLS = unilateral lower limb suspension. Values are reported

as mean ± standard deviation. Please note that some parameters are obtained only in one

condition (rest or contraction). *p < 0.05 and **p < 0.01 express significant difference from LS0

for the considered condition. Values are reported as mean ± standard deviation.

A) r$_{rm}$ = 0.773, p = 0.005

B) r$_{rm}$ = 0.624, p = 0.040