TITLE:

Mean exercise torque is a critical factor influencing neuromuscular fatigability induced

by exhausting contractions

RUNNING HEAD:

Mean exercise torque and fatigability

AUTHORS:

Loïc Lebesque[1]*, Gil Scaglioni[1], Patrick Manckoundia[1,2], Alain Martin[1]

[1] INSERM UMR1093-CAPS, Université de Bourgogne, UFR des Sciences du Sport,

F-21000, Dijon.

[2] Geriatrics Internal Medicine Department, University Hospital of Dijon Bourgogne,

Dijon CEDEX, 21079.

*CORRESPONDING AUTHOR:

Loïc Lebesque

loic.lebesque@u-bourgogne.fr

INSERM UMR1093 CAPS, Université de Bourgogne, UFR STAPS, Campus

Universitaire, BP 27877, F-21078, Dijon, France.

+33380396767

ORCID ID:

Loïc Lebesque: 0000-0002-2761-7276

Patrick Manckoundia: 0000-0002-5518-9803

26   Gil Scaglioni: 0000-0001-9049-2421

49   **ABSTRACT (250 words max.)**

PURPOSE: To get a more detailed description of neuromuscular fatigability, maximal torque sustainability (i.e., the ability to maintain a high torque level) can be assessed in addition to the classically used maximal voluntary contraction (MVC). Since this parameter appears to be affected by mean exercise torque (MET), the present study aims to examine the relationship between MET and neuromuscular fatigability induced by exhausting contractions.

METHODS: Thirteen participants sustained a plantar flexors MVC for 1 min (MVC_{1-MIN}) before and after exhausting exercises designed to produce a similar MET (30% MVC), and following a 10-min rest period. Exercises consisted of intermittent (INT), continuous (CON) or variable (continuous contraction alternating between moderate and low intensity, VAR) contractions performed until task failure.

RESULTS: Although the INT resulted in greater exercise duration and torque-time integral than CON and VAR, MVC similarly decreased after all exercises due to neural and muscular impairments. The torque loss during the MVC_{1-MIN} increased after all exercises to a similar extent, mainly because of neural alterations. Contrary to MVC, the torque loss during the MVC_{1-MIN} returned to baseline value after the recovery period.

CONCLUSION: By considering both maximal torque production and sustainability, INT, CON and VAR exercises, performed with identical mean torque and until exhaustion, led to a similar neuromuscular fatigability. Results confirm the independence of maximal torque production from the contraction pattern and support the impact of MET on maximal torque sustainability. The present findings are crucial to consider for the management of neuromuscular fatigability in both athletes and patients.

KEY WORDS:

Maximal torque production; Maximal torque sustainability; Contraction pattern;

Neuromuscular fatigability;

ABBREVIATIONS

ANOVA	Analysis of variance
BF^{OH}_{01}	Overlapping hypothesis Bayes Factor
BF^{NOH}_{01}	Non-overlapping hypothesis Bayes Factor
CON	Continuous exercise
EMG	Electromyography
ED	Exercise duration
INT	Intermittent exercise
LG	Lateral gastrocnemius
MET	Mean exercise torque
MG	Medial gastrocnemius
M_{MAX}	Maximal M-wave
MVC	Maximal voluntary contraction
$MVC_{1\text{-}MIN}$	Maximal voluntary contraction sustained for 1 minute
Pd	Potentiated doublet
Pd_A	Potentiated doublet obtained after the sustained maximal voluntary contraction
Pd_B	Potentiated doublet obtained before the sustained maximal voluntary contraction
PF	Plantar flexors
RMS	Root mean square
RPE	Rate of perceived exertion
SOL	Soleus
TA	Tibialis anterior
TF	Task failure
TTI	Torque-time integral
VAL	Voluntary activation level
VAR	Variable exercise

STATEMENTS AND DECLARATIONS

Grants

This work was supported by the Regional Council of Bourgogne Franche-Comté.

Conflict of interest

The authors declare that the research was conducted in the absence of any

commercial or financial relationship that could be construed as a potential conflict of

interest.

Data Availability

Data will be made available on reasonable request.

Supplementary data

Supplementary data: https://doi.org/10.6084/m9.figshare.24867426

INTRODUCTION

Neuromuscular fatigability corresponds to the decrement in magnitude or rate of change in a performance criterion, relative to a reference value, over a given time of task performance (Skau et al. 2021). The reduction of the capacity to produce the maximal torque during a brief (i.e., a few seconds long) maximal voluntary contraction (MVC) is the most commonly used index to quantify neuromuscular fatigability. Indeed, a substantial body of studies used the decrease in MVC to investigate the neuromuscular fatigability following a continuous (CON) contraction performed until task failure (TF) with various task modalities, such as different exercise feedback types (Hunter et al. 2002, 2005, 2008; Place et al. 2006a, b, 2007; Klass et al. 2008; Baudry et al. 2009, 2011; Rudroff et al. 2010; Williams et al. 2014), muscle lengths (Place et al. 2005), muscle groups (Neyroud et al. 2013), or even kinds of limb support (Yoon et al. 2009). Although the exercise duration (ED) and torque-time integral (TTI) of the CON exercises seem to be modulated by such exercise parameters, the reduction in MVC was of a similar extent in all studies. The present observation suggests that neuromuscular fatigability, quantified by the decrease in maximal torque production, was similar among task modalities when exercises were performed to exhaustion (i.e., until TF). However, little is known about the influence of contraction patterns on the neuromuscular fatigability induced during such exercises. To address this issue, one of our recent studies compared the neuromuscular fatigability induced by CON and intermittent (15 s on, 5 s off; INT) contractions of plantar flexors (PF) performed at 40% MVC until TF, using a 1-min sustained MVC (MVC_{1-MIN}) rather than a brief MVC (Lebesque et al. 2023). As a matter of fact, the MVC_{1-MIN} allows, at the same time, the assessment of two distinct and complementary characteristics of neuromuscular fatigability – the maximal torque

production and sustainability (i.e., the ability to sustain the maximal torque) – thus it

has the advantage of providing a more detailed and comprehensive description of

neuromuscular fatigability (Lebesque et al. 2022, 2023). Although at exhaustion, the

ED of the INT exercise was longer than that of the CON, both exercises led to a

similar MVC decline, while the maximal torque sustainability was affected by the

CON. Based on these results, a CON exercise appears to be more fatiguing than

intermittent contractions, when performed at the same intensity until exhaustion. In

addition, it is worth noting that the change in maximal torque sustainability was

positively correlated to the mean exercise torque (MET). In other words, the greater

the mean torque developed during the exhausting exercise, the more the ability to

sustain the maximal torque is impaired. This finding suggests that exercises

developing an identical MET should lead to similar neuromuscular fatigability.

Combined with MVC, maximal torque sustainability provides a more accurate

assessment of the fatiguing nature of exercises performed until exhaustion. It is

therefore crucial to acquire information on this neuromuscular capacity and in

particular on its relationship with the MET. Thus, the current study aims to further

analyse the impact of MET on maximal torque sustainability by extending this

analysis to additional exercise profiles generating a similar MET. Therefore, we used

CON, INT and variable (VAR) contractions with a MET set at 30% MVC and

performed until TF. Considering exhaustion achieved at the end of each exercise, it is

hypothesized that the decline in MVC is similar for the three contraction patterns.

Since the MET is identical for all exercises, we also expect maximal torque

sustainability to be equally compromised by each of them. Finally, because

neuromuscular fatigability is a transient phenomenon, the evolution of both capacities

was also explored after 10 min of rest. We hypothesize that 10 min of rest after

reaching exhaustion is sufficient for complete recovery in maximal torque

sustainability but not in maximal torque production, as already observed 10 min after

CON and INT contraction at 40% MVC until TF (Lebesque et al. 2023).

METHODS

Participants and ethics

Thirteen healthy male volunteers (age = 25 ± 3 years old; body mass = 75.4 ± 8.5 kg; height = 179.0 ± 5.8 cm, body mass index = 23.5 ± 1.8 kg·m^{-2}) were enrolled in this study after giving their oral and written informed consent. Participants were recreationally active and none of them suffered from neurological or physical disorders. For all participants, the dominant leg (i.e., the leg used to kick a ball on a target; van Melick et al. 2017) was the right one. This study was carried out in accordance with the latest revision of the Declaration of Helsinki. The CPP SOOM III ethics committee approved the protocol (number 2017-A00064-49).

Protocol

Experimental design. Participants took part in three experimental sessions randomly administered. Sessions lasted approximately 1.5 hours. For each subject, sessions were interspaced by a 7-day interval and planned at the same time of the day to avoid any circadian bias (Guette et al. 2005; Tamm et al. 2009; Douglas et al. 2021). Each session included neuromuscular assessments (torque and electromyographic (EMG) recordings) of PF before (Pre) and immediately after (Post) a submaximal fatiguing exercise, as well as after a 10-min recovery period (Post-10). The fatiguing exercise started 10 min after the completion of the pre-exercise neuromuscular evaluation and it consisted of CON, INT or VAR contractions with a mean torque of 30% MVC performed until exhaustion. Figure 1A shows the experimental protocol used in the present study.

Neuromuscular assessment. A standardized warm-up protocol for PF preceded the neuromuscular assessment and consisted of 10-15 PF isometric contractions lasting 5 s each, with an incremental intensity from 25 to 100% of their maximal effort. After warming up, subjects performed two brief PF MVCs (5-s long, with 1-min rest in-between) and 2 s after each contraction an electrical doublet was delivered to obtain the potentiated doublet (Pd). If the difference between the MVCs was > 5%, an additional maximal contraction was performed. Then, 2 min of rest was given to the subjects before the Pre-exercise $MVC_{1\text{-}MIN}$. For each $MVC_{1\text{-}MIN}$, an electrical doublet was delivered at rest and gave subjects, inter alia, the signal to start the $MVC_{1\text{-}MIN}$. Electrical doublets superimposed to the sustained maximal contraction were delivered 3 and 58 s after the $MVC_{1\text{-}MIN}$ start. A final doublet was delivered at rest, 3 s after the end of the $MVC_{1\text{-}MIN}$. Participants were informed of the duration of the sustained MVC and instructed to perform a maximal performance during the entire contraction in order to lessen the adoption of a pacing strategy (Halperin et al. 2014) during the $MVC_{1\text{-}MIN}$. In addition, strong and loud verbal encouragements were given to the subjects to ensure maximal performance during the sustained contraction. The $MVC_{1\text{-}MIN}$ were performed before (Pre) and after (Post) the fatiguing exercise and after the 10-min recovery period (Post-10), while the brief MVC was carried out only at Pre and Post-10 because the neuromuscular fatigability should be assessed as soon as possible after the exercise completion (Mira et al. 2017; Place and Millet 2019).

Electrical stimulation. Electrophysiological and torque responses of the triceps surae muscles were evoked by percutaneously stimulating the posterior tibial nerve, in the popliteal fossa of the dominant leg. Monophasic rectangular pulses (1 ms) were

delivered using a Digitimer stimulator (DS7AH, Digitimer, Hertfordshire, UK), triggered by a commercially available software (Tida, Heka Elektronik, Lambrecht/Pfalz, Germany). The anode (5 x 10 cm, Compex SA, Ecublens, Switzerland) was placed beneath the patella. A hand-held cathode ball (5 mm diameter) was used to locate the optimal stimulation site, namely the site that evoked the greatest soleus (SOL) M-wave amplitude. Once the stimulation site was determined, the cathode (a self-adhesive electrode with a 7 mm diameter, Contrôle Graphique Medical, Brie-Compte-Robert, France) was fixed to this site with tape. A hand-made hard-pad (a cotton cube wrapped in adhesive tape; 4 x 2 x 2 cm) was then positioned over the cathode and firmly fixed with adhesive tape to the limb to keep constant pressure on the electrode throughout the experiment. To determine the intensity at which the M-wave amplitude plateaued (i.e., no further increase in the M-wave amplitude was evident in SOL), the intensity of stimulation was gradually increased by 5 mA from the motor threshold to the maximal M-wave of SOL. This intensity was further increased by 20% to ensure supramaximal stimulations during the entire session and was kept constant throughout the experimental session (mean supramaximal intensity: 72.2 ± 22.0 mA, range: 42 – 144 mA). Only electrical doublets at 100 Hz (1 ms pulse duration, supramaximal intensity) were delivered during the experiment.

Fatiguing exercises. Visual feedback of the torque exerted by the subjects and the target torque were displayed on a computer screen placed 1 m in front of them. Fatiguing exercises consisted in isometric contractions of PF. For the CON session, subjects were asked to sustain the target torque (30% MVC) until TF. For the INT session, subjects were instructed to perform intermittent (15 s on, 5 s off with sound

signals; duty cycle: 75%) contractions at 40% MVC until TF. Finally, for the VAR

session, subjects were required to perform a sustained contraction characterized by

alternating periods of 10 s at 20% MVC and 10 s at 40% MVC (i.e., variable

contraction intensity) until TF. It is noteworthy that, for this contraction pattern, half of

the subjects started the exercise with 10 s at 20% MVC while the other half started at

40% MVC. In each session, the expected MET was 30% MVC. The TF was defined

as a drop of the torque below the target for three consecutive seconds. All subjects

were strongly encouraged throughout the entire exercise and naive about the

criterion leading to exercise interruption. The rate of perceived exertion (RPE),

defined as "the conscious sensation of how hard, heavy, and strenuous a physical

task is" (Marcora 2010; Pageaux 2016) was assessed at TF for all exercises.

Experimental sessions were administered in a randomized order.

Data acquisition

Torque recordings. PF isometric torque was recorded using an isokinetic

dynamometer (Biodex System 4 pro®, Biodex Medical System Inc., Shirley, NY,

USA). Subjects were comfortably seated on the dynamometer chair with the

dominant foot securely strapped to the footplate at the ankle level. The centre of

rotation of the dynamometer shaft was aligned with the lateral malleolus. The ankle,

knee, and hip angles were set at 90°, 120°, and 130° respectively (180° full

extension). To limit trunk movements that could affect torque development, the trunk

and hip were firmly strapped to the seat with belts. The setting configuration was kept

constant throughout the entire experimental protocol and among sessions. Torque

data were recorded at a sampling frequency of 2 kHz with the Biopac acquisition

system (Biopac MP160, Biopac System Inc, USA).

Electromyographic recordings. The EMG activity was recorded from four muscles of the dominant leg: the medial (MG) and lateral (LG) gastrocnemii, the soleus (SOL), and the tibialis anterior (TA). The EMG signals were recorded bipolarly using silver chloride circular surface electrodes (7 mm recording diameter; Contrôle Graphique Medical, Brie-Compte-Robert, France) with an inter-electrode centre-to-centre distance of 20 mm. The skin was shaved, abraded, and cleaned with alcohol before electrode placement to minimize skin impedance (< 5 kΩ). For SOL, electrodes were placed 3 cm below the insertion of the two gastrocnemii over the Achilles tendon. For MG and LG muscles, electrodes were positioned over the muscle belly. For TA, they were placed on the upper third of the distance between the fibula head and the tip of the lateral malleolus. The reference electrode was positioned between the two gastrocnemii bellies of the same leg in a central position. EMG signals were amplified (gain = 1000) and filtered (10 – 500 Hz). EMG data were recorded at a sampling frequency of 2 kHz with the Biopac acquisition system (Biopac MP160, Biopac System Inc, USA; RRID: SCR_014829).

Data analysis

Torque and electrophysiological data were processed using custom programs written in Matlab (version 9.4 R2018a, Mathworks, Natick, MA, USA; RRID: SCR_001622). An experimental torque trace of the main neuromuscular parameters from a representative subject is depicted in Figure 1B. Data processing was similar to our previous study (Lebesque et al. 2023).

Torque analysis. The maximal torque developed in the first 5 s of the MVC_{1-MIN} is a

reproducible and reliable index of maximal torque production (Lebesque et al. 2022,

2023) and was then labelled 'MVC'. The Pre-exercise MVC was used to set the

target torque of the fatiguing exercises. The torque loss during each MVC_{1-MIN}

(ΔMVC_{1-MIN}) was calculated as the relative change between the MVC and the mean

torque during the final 2 s of contraction:

$$\Delta MVC_{1\text{-MIN}}\ (\%) = \left[\left(\frac{final\ torque}{MVC}\right) - 1\right] \times 100$$

313 This index of maximal torque sustainability showed good intra- and inter-individual

314 reproducibility (Lebesque et al. 2023). Change in ΔMVC_{1-MIN} between pre and post-

315 exercise was quantified as the difference between both measures:

$$\Delta MVC_{1\text{-MIN}} \text{ change }\ (\%) = \text{Pre-exercise } \Delta MVC_{1\text{-MIN}}\ - \text{ Post-exercise } \Delta MVC_{1\text{-MIN}}$$

Concerning the potentiated doublets, we measured their amplitude before (Pd_B) and

after (Pd_A) the MVC_{1-MIN}. The Pd_B, at Pre and Post-10, corresponds to the potentiated

doublet elicited just after the brief MVC, while at Post it corresponds to the

potentiated doublet elicited just before the MVC_{1-MIN}. This latter was potentiated by

the fatiguing exercise performed until exhaustion. At each time point (Pre, Post and

Post-10), the ratio between the Pd_A and the Pd_B represents the evolution of the

potentiated doublet following each MVC_{1-MIN} (ΔPd_{1-MIN}), and was expressed as a

percentage of change as follows:

$$\Delta Pd_{1\text{-MIN}}\ (\%) = \left[\left(\frac{Pd_A}{Pd_B}\right) - 1\right] \times 100$$

325 According to the original formula from Allen et al. (1995), the voluntary activation

326 level (VAL) was determined at the beginning (i.e., after 3 s of contraction) and the

327 end (i.e., after 58 s of contraction) of the MVC_{1-MIN}. Strojnik and Komi correction

328   (Strojnik and Komi 1998) was applied when the superimposed doublet was not

329   exactly delivered over the peak torque as follows:

$$VAL~(\%) = \left[1 - \frac{superimposed~doublet \times \left(\frac{T_{STIM}}{MVC}\right)}{Pd}\right] \times 100$$

T_{STIM} represents the voluntary torque averaged over a 500 ms time window prior to

the superimposed doublet stimulation. Note that the VAL at the end of the $MVC_{1\text{-MIN}}$

was corrected using the mean torque of the last 2 s of contraction instead of the MVC

in the VAL equation. A value of 100% indicates a full voluntary activation. To

calculate the VAL at the beginning of the $MVC_{1\text{-MIN}}$, we used the amplitude of the

Pd_B, while at the end of the $MVC_{1\text{-MIN}}$, we used the amplitude of the Pd_A. The ratio

between the VAL at the beginning and the end of the $MVC_{1\text{-MIN}}$ represents the

evolution of the VAL during the $MVC_{1\text{-MIN}}$ ($\Delta VAL_{1\text{-MIN}}$) and was expressed in

percentage of change as follows:

$$\Delta VAL_{1\text{-MIN}}~(\%) = \left[\left(\frac{VAL~end}{VAL~beginning}\right) - 1\right] \times 100$$

The TTI of the fatiguing exercise, namely the effective amount of torque produced by

PF, is computed as the integral of the torque during the entire exercise period minus

the integral of the baseline torque at rest during the same period. INT exercise period

started with the raising phase of the first contraction and ended after the decline

phase of the last contraction. The total ED refers to the duration of the entire exercise

(contractions and rest periods). The effective ED represents the real duration of

contraction (i.e., without rest periods). Thus, the effective ED is equal to the total ED

for CON and VAR exercises. The MET, calculated as the exercise TTI divided by the

total ED, corresponds to the mean torque developed during the entire ED.

$$Mean~Exercise~Torque~(Nm) = \frac{Exercise~TTI}{Total~ED}$$

EMG analysis. We analysed the second M-wave of each Pd (Kawakami et al. 2000;

Matkowski et al. 2011). The peak-to-peak amplitude of the maximal M-wave of the

SOL, MG and LG was measured and then summed to determine a comprehensive

PF maximal M-wave (M_{MAX}). The EMG root mean square (RMS) of the SOL, MG

and LG was calculated and summed at both the beginning and the end of the

fatiguing exercise. For each measure, RMS was averaged over a 5 s time window.

For CON, we consider the RMS between the 5^{th} and the 10^{th} s of the exercise, and

between 10 and 5 s before the exercise end. For INT, we consider the RMS between

the 5^{th} and the 10^{th} s of the first and last contraction. For VAR, we consider the RMS

between 2.5 and 7.5 s of the first and the last contraction of each intensity (i.e., 20

and 40% MVC). For the latter exercise, we averaged the RMS obtained at 20 and

40% MVC at both the beginning and the end of the exercise. Then, the RMS, at the

beginning and the end of each fatiguing task, was normalized by the Pre and Post

M_{MAX} (RMS/M_{MAX}) respectively. The ratio between the RMS/M_{MAX} at the beginning

and the end of the exercise represents the evolution of the muscle activities during

the fatiguing task and was expressed in percentage of change as follows:

$$\Delta RMS/M_{MAX}\ (\%) = \left[\left(\frac{RMS/M_{MAX}\ end}{RMS/M_{MAX}\ beginning}\right) - 1\right] \times 100$$

Psychometric analysis. The RPE was quantified at TF using the CR-100 Scale (Borg

2007).

Statistical analysis. It was estimated that thirteen participants were needed to detect

differences with moderate effect size ($\eta_p^2 = 0.08$) using standard parameters of 1-β =

0.95 and α = 0.05. We used the Shapiro-Wilk test to assess the normality of each

dataset. Because all variables follow the normal distribution, we used only parametric

analyses of variance (ANOVA). Factors used in ANOVAs were *pattern* (INT, CON

and VAR), *time* (Pre, Post and Post-10) and *duration* (beginning *vs.* end of fatiguing

exercise). One-way ANOVAs (*pattern* factor) were performed on total and effective

ED, exercise TTI, MET, RPE at TF and ΔRMS/M_{MAX}. Two-way repeated-measures

ANOVAs (*pattern* and *time* factors) were conducted on MVC, VAL, M_{MAX}, Pd_B,

$\Delta MVC_{1\text{-MIN}}$, $\Delta VAL_{1\text{-MIN}}$ and $\Delta Pd_{1\text{-MIN}}$. A two-way repeated-measures ANOVA (*pattern*

and *duration* factors) was performed on RMS/M_{MAX}. When a significant effect was

observed, we used Tukey's test for post-hoc comparisons. When necessary, we

performed Bayesian equivalence analysis with a region of practical equivalence

ROPE = [-0.1:0.1] and a prior Cauchy scale of 0.707 (Morey and Rouder 2011). The

correlation between muscle activity during the fatiguing exercise (RMS/M_{MAX}) and the

effective ED was analysed using repeated-measures correlation (rmcorr R package,

Bakdash and Marusich 2017). Effect sizes of ANOVAs are reported as partial eta

squared (η_p^2) with small (≥ 0.01), moderate (≥ 0.07) and large effects (≥ 0.14).

Results are reported as mean ± standard deviation. We set the significant level at *p* <

0.050. The sample size was estimated using G*Power software (version 3.1.9.2,

Universität Düsseldorf, Germany; RRID: SCR_013726). The repeated-measures

correlation was performed and plotted with R (R Core Team, 2017, version 4.1.3;

RRID: SCR_001905). Bayesian equivalence tests were conducted with JASP

software (JASP Team, 2023, version 0.16; RRID: SCR_015823). We conducted all

inferential statistical analyses using Statistica software (Statsoft, version 13, Tulsa,

OK, USA; RRID: SCR_014213). Graphs were made with GraphPad Prism software

(GraphPad Software Inc., version 8.4.0, San Diego, CA, USA; RRID: SCR_002798).

RESULTS

Parameters characterizing fatiguing exercise

The MET, a-priori set at 30% MVC, was not different among exercises, whether expressed in absolute ($F_{(2,24)}$ = 0.600, p = 0.557, η_p^2 = 0.048) or relative ($F_{(2,24)}$ = 0.720, p = 0.496, η_p^2 = 0.057) values (Table 1). Bayesian equivalence tests, among exercises MET, showed that the overlapping hypothesis Bayes Factor (BF^{OH}_{01}) was superior to 2.137 and the non-overlapping hypothesis Bayes Factor (BF^{NOH}_{01}) was superior to 2.406, meaning that the data are 2.1 times more likely to validate the null hypothesis than the alternative one and 2.4 times more likely to lie in the equivalence than in the non-equivalence region. For the INT exercise, participants performed 52.5 ± 27.2 contractions. The analysis of variance detected a significant effect of *pattern* on total ED ($F_{(2,24)}$ = 23.686, $p < 0.001$, η_p^2 = 0.664) and

effective ED ($F_{(2,24)}$ = 12.070, p = 0.002, η_p^2 = 0.501). The INT total and effective ED

were longer than CON (+106.0 %, p < 0.001 and +54.7%, p < 0.001, respectively)

and VAR (+93.7 %, p < 0.001 and +45.5%, p = 0.002, respectively) (Table 1). No

difference in ED between CON and VAR was spotted (p = 0.862; Bayesian

equivalence tests: BF^OH_{01} = 3.349, BF^NOH_{01} = 4.354) (Table 1). We found a main

pattern effect on exercise TTI ($F_{(2,24)}$ = 42.404, p < 0.001, η_p^2 = 0.779). The TTI of the

INT exercise was greater compared to that of CON (+104.9%, p < 0.001) and VAR

(+86.8%, p < 0.001) contractions (Table 1), with no difference between CON and

VAR values (p = 0.725). We found a significant *pattern* x *duration* interaction effect

($F_{(2,24)}$ = 4.471, p = 0.022, η_p^2 = 0.271) on RMS/M_MAX. At the beginning of the

fatiguing exercise, the RMS/M_MAX of the triceps surae muscles was greater for INT

compared to CON (p = 0.002) and VAR (p = 0.017) while at the end of the fatiguing

exercise, no more difference was observed among patterns (p-range: 0.871 – 1.000).

Compared to the initial value, this parameter significantly increased at the end of

each fatiguing exercise (INT: +44.0%, p < 0.001; CON: +120.7%, p < 0.001; VAR:

+87.2%, p < 0.001). In addition, we found a significant negative correlation between

$\Delta RMS/M_\text{MAX}$ and the effective ED (r_{rm} (25) = -0.391, p = 0.044, $CI_{95\%}$ = [-0.671 :

0.013]) (Figure 2). Experimental torque and EMG traces during exercises, for a

representative subject, are depicted in Figure 3.

At TF, the RPE did not differ among exercises ($F_{(2,24)}$ = 3.109, p = 0.063, η_p^2 = 0.206)

and ranged between "extremely strong" and "maximal" on the CR-100 scale (Table

1).

Assessments related to maximal torque production

MVC. Only a significant main *time* effect was detected on MVC ($F_{(2,24)}$ = 145.033, $p <$

0.001, η_p^2 = 0.924). Fatiguing exercises resulted in a 29.7% reduction in MVC (127.0

± 20.7 Nm *vs.* 89.8 ± 17.7 Nm, $p <$ 0.001). After the 10-min rest period, the MVC

(115.4 ± 22.3 Nm) remained 10.2% below the pre-exercise value ($p <$ 0.001),

revealing an incomplete recovery from the fatiguing exercise ($p <$ 0.001) (Figure 4A).

Bayesian equivalence tests showed that MVC was likely similar among exercises at

Pre ($BF^{OH}_{01} > 3.111$ and $BF^{NOH}_{01} > 3.925$ in all cases), Post ($BF^{OH}_{01} > 2.510$ and

$BF^{NOH}_{01} > 2.947$ in all cases) and Post-10 ($BF^{OH}_{01} > 3.226$ and $BF^{NOH}_{01} > 4.130$ in all

cases).

VAL. The ANOVA analysis only revealed a main *time* effect on VAL ($F_{(2,24)}$ = 21.140,

$p <$ 0.001, η_p^2 = 0.638). The VAL was decreased by 7.7% after the fatiguing

exercises (98.4 ± 1.6 % *vs.* 90.8 ± 6.8 %, $p <$ 0.001). The VAL fully recovered after

the rest period since at Post-10, no more difference was detected with the pre-

exercise value ($p =$ 0.721) (Figure 4B).

M_{MAX}. A main *time* effect was observed on the M_{MAX} amplitude ($F_{(2,24)}$ = 18.833, $p <$

0.001, η_p^2 = 0.611). The M_{MAX} following the fatiguing exercises (18.8 ± 6.9 mV) was

17.7% lower than before (22.9 ± 7.6 mV, $p <$ 0.001), and fully recovered after the 10-

min rest period (21.5 ± 7.4 mV) since there was no more difference with Pre-exercise

($p =$ 0.102).

Pd_B. Only a main *time* effect was detected on Pd_B ($F_{(2,24)}$ = 12.739, $p <$ 0.001, η_p^2 =

0.515). The Pd_B decreased by 10.9% at Post (33.8 ± 4.7 Nm *vs.* 30.1 ± 4.8 Nm, $p <$

0.001). At Post-10, the Pd_B (31.8 ± 5.0 Nm) did not recover since it was still

significantly lower than before the exercise (5.9%, $p = 0.030$) (Figure 4C).

Assessments related to maximal torque sustainability

ΔMVC$_{1\text{-}MIN}$. The ANOVA revealed only a main *time* effect on $\Delta MVC_{1\text{-}MIN}$ ($F_{(2,24)}$ =

9.542, $p < 0.001$, $\eta_p^2 = 0.443$). Compared to before (-44.9 ± 10.1 % MVC), the

$\Delta MVC_{1\text{-}MIN}$ increased by 13.1% after the fatiguing exercises (-50.8 ± 10.4 % MVC, $p =$

0.001) and was fully recovered at Post-10 (-46.2 ± 9.0 % MVC) since no more

difference was found with Pre-exercise ($p = 0.617$) (Figure 5A). Bayesian

equivalence tests showed that $\Delta MVC_{1\text{-}MIN}$ were likely similar among exercises at Pre

($BF^{OH}_{01} > 3.501$ and $BF^{NOH}_{01} > 4.641$ in all cases), Post ($BF^{OH}_{01} > 1.711$ and BF^{NOH}_{01}

> 1.840 in all cases) and Post-10 ($BF^{OH}_{01} > 1.241$ and $BF^{NOH}_{01} > 1.271$ in all cases).

ΔVAL$_{1\text{-}MIN}$. The statistical analysis showed only a main *time* effect on $\Delta VAL_{1\text{-}MIN}$

($F_{(2,24)} = 31.765$, $p < 0.001$, $\eta_p^2 = 0.726$). $\Delta VAL_{1\text{-}MIN}$ was 250.8% higher after the

exercises (-38.2 ± 22.8 %) compared to before (-15.2 ± 12.6 %, $p < 0.001$). At Post-

10, the $\Delta VAL_{1\text{-}MIN}$ was fully recovered since no more difference was detected with the

Pre value ($p = 0.335$) (Figure 5B).

ΔPd$_{1\text{-}MIN}$. No *pattern* ($F_{2,24} = 0.792$, $p = 0.465$, $\eta_p^2 = 0.062$), *time* ($F_{2,24} = 1.134$, $p =$

0.339, $\eta_p^2 = 0.086$) or *pattern* x *time* interaction ($F_{4,48} = 1.165$, $p = 0.338$, $\eta_p^2 = 0.089$)

effect on $\Delta Pd_{1\text{-}MIN}$ was detected by statistical analysis (Figure 5C). Pre, Post and

Post-10 pooled values were 3.3 ± 13.9 %, 2.0 ± 9.5 % and 5.4 ± 11.5 % respectively.

DISCUSSION

The current study aims, using different contraction patterns, to gain a better

understanding of how the mean torque of exhausting exercises affects

neuromuscular fatigability. Our findings demonstrate that INT, CON and VAR

contractions produce similar levels of neuromuscular fatigability, as evidenced by the

comparable reductions in maximal torque production and sustainability across the

exercises. These results highlight the independence of neuromuscular fatigability

from the contraction pattern during such exercises and support the relationship

between the MET and maximal torque sustainability. The different recovery kinetics

emphasize that maximal torque production and sustainability represent distinct

capacities and, therefore, must be evaluated simultaneously when exploring

neuromuscular fatigue.

Parameters characterizing fatiguing exercise

The inverse relationship between the ED and the intensity of a sustained

contraction has been substantiated by previous studies (Rohmert 1960; Shahidi and

Mathieu 1995; Rudroff et al. 2007). Our results allow the extension of this relationship

to sustained contraction with variable intensity (i.e., VAR). Therefore, it appears that

the ED of a maintained submaximal contraction correlates to MET even if the exerted

torque is modulated during the exercise. This relationship is not valid anymore when

integrating the INT exercise, regardless of whether we consider total or effective ED.

The ED and TTI were greater for INT compared to CON and VAR. As previously

reported by Lebesque et al. (2023), the presence of rest periods during exercise

appears to be crucial to delay exhaustion and thus to stay in exercise longer. Indeed,

rest periods during INT allow the delivery of blood flow to the active muscles, leading

to a greater supply of substrates and oxygen as well as an enhanced metabolite

washout (Sahlin 1992; Degens et al. 1998; Hargreaves 2005; Ament and Verkerke

2009). It is noteworthy that modulating the contraction intensity from 40 to 20% MVC

during the VAR is not sufficient to extend the ED and TTI since, for both parameters,

no difference was observed with CON.

Since, at the beginning of the fatiguing exercise, the torque developed during

the INT (40% MVC) was greater than CON (30% MVC) and VAR (20 and 40% MVC,

mean torque = 30% MVC), for this modality of exercise we also observed a greater

PF electrical activity. The PF EMG activity increased from contraction onset to task

failure in all exercises, denoting a rise in the excitatory output of the motor neurons

(Fuglevand et al. 1993; Søgaard et al. 2006; Klass et al. 2008). This could be

explained by the increased recruitment of motor units and the modulation of their

firing rate to maintain the required torque, despite the development of neuromuscular

fatigability (Garland et al. 1994; Carpentier et al. 2001; Kallenberg and Hermens

2006; Enoka and Duchateau 2008; Klass et al. 2008). Regardless of the contraction

pattern, in our study, it appears that exhaustion was achieved for a similar level of PF

EMG activity, suggesting that TF occurs when the voluntary descending drive

mismatches the requirement to maintain the target torque. This finding is in

accordance with the recent data from Matta et al. (2023). By manipulating the time

perception during exercise, these authors highlight that subjects stopped their CON

contraction of the knee extensors when a critical value of EMG activity was reached,

independently of ED. Furthermore, in the present study, the effective ED negatively

correlates with the change in muscle activity during exhausting exercises, showing

that the greater the increase in muscle activity, the lower the ED. These findings are

in agreement with those observed during submaximal CON contractions sustained

until TF (Maluf et al. 2005; Hunter et al. 2008; Rudroff et al. 2011).

In the current investigation, the RPE quantified at TF was similar among

exercises. This common observation at the end of exhausting exercises (e.g. Hunter

et al. 2005, 2008; Rudroff et al. 2011; Williams et al. 2014) suggests that participants

performed a similar subjective maximal performance during INT, CON and VAR

exercises, irrespective of exercise duration or TTI.

Maximal torque production

In the present investigation, the maximal torque production was quantified as

the maximal torque achieved during the first 5 s of the MVC_{1-MIN}. This methodology

for assessing maximal torque production from a sustained MVC has already been

used by our research group (Lebesque et al. 2022, 2023). Indeed, it has been shown

that MVC from a MVC_{1-MIN} is a reliable and reproducible index of maximal torque

production when compared to the brief-MVC (Lebesque et al. 2022, 2023). As

hypothesized in the present study, the MVC loss after each exhausting exercises was

similar (-30 %), regardless of the ED or the TTI. In one of our recent investigations,

we also found a similar reduction in MVC (-27%) following CON and INT contractions

performed at 40% MVC until TF (Lebesque et al. 2023). Similarly, Smith et al. (2016)

observed an equivalent reduction in knee extensors MVC (-24%) after 50 intermittent

contractions (6 s on, 2 s off) performed at a constant (60% MVC) or variable (i.e.,

alternation of contractions at 40 and 80% MVC) intensity. Altogether, these outcomes

bring out that the reduction of maximal torque production is not sensitive to the

contraction pattern (i.e., CON, INT and VAR).

In the present investigation, we also found a lack of correlation between the

TTI and the MVC loss, for exercises performed until exhaustion with different

contraction patterns. This observation could also be inferred from the results of

previous investigations (Place et al. 2005, 2006a, b, 2007; Hunter et al. 2005, 2008;

Klass et al. 2008; Baudry et al. 2009, 2011; Yoon et al. 2009; Rudroff et al. 2010;

Neyroud et al. 2013; Williams et al. 2014), although in these latter the correlation was

not specifically addressed.

The equivalent MVC decline (-30%), induced by the three fatiguing tasks in the

present experiment, was associated with a reduction in VAL (-8%) which was also

equivalent among exercises. This finding suggests that the central nervous system

was similarly affected by the three contraction patterns and was no longer able to

maximally activate muscles at TF. Prior studies, with different protocols, have noted a

voluntary activation decrease following CON (e.g. Place et al. 2007; Neyroud et al.

2013) or INT (e.g. Bigland-Ritchie et al. 1986; Behm and St-Pierre 1998; Mademli

and Arampatzis 2008) exercises performed until TF at 40 – 65 % MVC, but our

recent investigation (Lebesque et al. 2023) stands out as the first to directly compare

voluntary activation after these two modalities of contraction showing an equivalent

impact (-11%) of both on VAL.

In the current research, the decline in MVC is also associated with alterations

occurring at the muscular level, as revealed by the decline in Pd_B (-11%) and M_{MAX} (-

18%). The extent of these impairments was equivalent for the three fatiguing

exercises and demonstrated that alteration in muscle contractility can mainly be

explained by the decline in sarcolemma excitability.

Altogether, INT, CON and VAR contractions, performed until exhaustion,

induced a similar reduction in maximal torque production that may be attributed to

both neural and muscular impairments, with a larger contribution from the latter.

These alterations are quantitatively similar among exercises.

The effect of a 10-min recovery period on the neuromuscular function was also

investigated. We observed that the maximal torque production recovered only

partially at Post-10 since the MVC remained lower, at this time point, compared to the

pre-exercise value. This finding agrees with previous investigations showing that

MVC was still depressed 10 min after CON (Fuglevand et al. 1993; Lebesque et al.

2023) or INT (Mademli and Arampatzis 2008; Lebesque et al. 2023) contractions

performed until exhaustion. Specifically, our results suggest that the incomplete

recovery of maximal torque production was mainly due to muscular factors since,

regardless of the contraction pattern, the VAL fully recovered at Post-10, while the

Pd_B was still lower than the pre-exercise value. Since the sarcolemma excitability

(i.e., M_{MAX}) was restored after the recovery period, it can be inferred that the

alterations occurred, more precisely, within the muscle cell (Allen et al. 2008; Cheng

et al. 2018). These findings suggest that intracellular mechanisms including calcic

phenomena, might be involved in the alteration of muscle contractility (i.e., Pd_B) after

the fatiguing exercises (Place et al. 2007; Allen et al. 2008; Cheng et al. 2018).

Maximal torque sustainability

According to our hypothesis, the INT, CON and VAR exercises similarly

altered the $\Delta MVC_{1\text{-}MIN}$. This finding means that exhausting exercises performed with

an identical MET lead to an equivalent impairment of the ability to sustain a maximal

effort, regardless of the ED or the TTI. This outcome appears to support the

significant correlation between the MET and the alteration in maximal torque

sustainability (Lebesque et al. 2023). Indeed, although the contraction pattern was

different among exercises, the MET was equivalent and it led to a similar decline in

maximal torque sustainability. If such a relationship did not exist, we should have detected a different alteration in maximal torque sustainability among the exercises.

The similar alteration of maximal torque sustainability, after the three exercises, was accompanied by a similar increase in $\Delta VAL_{1\text{-MIN}}$. The accumulation of contraction by-products in the intracellular milieu is known to stimulate group III/IV muscle afferents through receptors sensitive to the mechanic and metabolic changes (Garland and Kaufman 1995), leading to the increase of motor cortex inhibition that limits the efferent motor command (Gandevia et al. 1996; Gandevia 2001; Taylor et al. 2016).

The absence of change in $\Delta Pd_{1\text{-MIN}}$ stresses that the decline in maximal torque sustainability may be mainly explained by the reduction in the capacity to maintain a high level of voluntary activation, confirming our previous finding following CON and INT exhausting exercises (Lebesque et al. 2023). This observation has also be made after more ecological exercises such as cycling (Girard et al. 2013). The link between the capacities to maintain the maximal torque and the maximal voluntary activation is also hinted by the fact that the 10 min recovery period was sufficient to restore both the $\Delta VAL_{1\text{-MIN}}$ and the $\Delta MVC_{1\text{-MIN}}$.

Regardless of the contraction pattern, the neuromuscular fatigability is similar when fatiguing exercises are performed until TF. Indeed, the ability to produce and sustain the maximal torque was altered to the same extent at the end of the different exercises. However, the behaviour of MVC and $\Delta MVC_{1\text{-MIN}}$ was different after the rest of 10 min. Contrary to the MVC that did not completely recover after the rest period, the $\Delta MVC_{1\text{-MIN}}$ fully recovered at Post-10. These results are in accordance with previous observations done 10 min after isometric exercises of PF performed to exhaustion (Lebesque et al. 2023) or not (Lebesque et al. 2022). The present data

therefore confirm that maximal torque production and sustainability represent distinct

neuromuscular capacities. Accordingly, it is not possible to extrapolate the behaviour

of one of these from the other, as experimentally shown by the absence of correlation

between MVC and $\Delta MVC_{1\text{-}MIN}$ changes after the fatiguing exercises and following the

recovery period (see Supplemental Figure S1). Therefore, it is important to evaluate

both simultaneously when aiming to explore more deeply the neuromuscular

fatigability of a muscle task or the effects of a training program on neuromuscular

function.

Conclusion

In the present investigation, we demonstrated that INT, CON and VAR

submaximal exercises, performed until exhaustion with an equivalent MET, lead to a

similar reduction in both maximal torque production and sustainability. Therefore, by

evaluating both capacities (Lebesque et al. 2022, 2023), it can be concluded that

these exercises induce the same neuromuscular fatigability. In addition, the distinct

recovery kinetic of maximal torque production and sustainability confirms the

importance of assessing both capacities simultaneously to get a more

comprehensive description of neuromuscular fatigability. Finally, our results show a

correlation between MET and change in $\Delta MVC_{1\text{-}MIN}$, namely a sensitivity of maximal

torque sustainability to MET. This finding is crucial to consider for the management of

neuromuscular fatigability during exercise training and reconditioning in both athletes

and pathological persons.

ACKNOWLEDGMENTS

The authors would like to thank all volunteers that participated in the study.

REFERENCES

Allen DG, Lamb GD, Westerblad H (2008) Skeletal muscle fatigue: cellular mechanisms. Physiol Rev 88:287–332. https://doi.org/10.1152/physrev.00015.2007

Allen GM, Gandevia SC, McKenzie DK (1995) Reliability of measurements of muscle strength and voluntary activation using twitch interpolation. Muscle Nerve 18:593–600. https://doi.org/10.1002/mus.880180605

Ament W, Verkerke GJ (2009) Exercise and Fatigue. Sports Med 39:389–422. https://doi.org/10.2165/00007256-200939050-00005

Bakdash JZ, Marusich LR (2017) Repeated Measures Correlation. Front Psychol 0: https://doi.org/10.3389/fpsyg.2017.00456

Baudry S, Maerz AH, Gould JR, Enoka RM (2011) Task- and time-dependent modulation of Ia presynaptic inhibition during fatiguing contractions performed by humans. J Neurophysiol 106:265–273. https://doi.org/10.1152/jn.00954.2010

Baudry S, Rudroff T, Pierpoint LA, Enoka RM (2009) Load type influences motor unit recruitment in biceps brachii during a sustained contraction. J Neurophysiol 102:1725–1735. https://doi.org/10.1152/jn.00382.2009

Behm DG, St-Pierre DMM (1998) Fatigue Mechanisms in Trained and Untrained Plantar Flexors. The Journal of Strength & Conditioning Research 12:166

Bigland-Ritchie B, Furbush F, Woods JJ (1986) Fatigue of intermittent submaximal voluntary
contractions: central and peripheral factors. Journal of Applied Physiology 61:421–429.
https://doi.org/10.1152/jappl.1986.61.2.421

Borg E (2007) On perceived exertion and its measurement: (529832013-001)

Carpentier A, Duchateau J, Hainaut K (2001) Motor unit behaviour and contractile changes during
fatigue in the human first dorsal interosseus. J Physiol 534:903–912.
https://doi.org/10.1111/j.1469-7793.2001.00903.x

Cheng AJ, Place N, Westerblad H (2018) Molecular Basis for Exercise-Induced Fatigue: The
Importance of Strictly Controlled Cellular Ca $^{2+}$ Handling. Cold Spring Harbor Perspectives in
Medicine 8:a029710. https://doi.org/10.1101/cshperspect.a029710

Degens H, Salmons S, Jarvis JC (1998) Intramuscular pressure, force and blood flow in rabbit tibialis
anterior muscles during single and repetitive contractions. Eur J Appl Physiol Occup Physiol
78:13–19. https://doi.org/10.1007/s004210050381

Douglas CM, Hesketh SJ, Esser KA (2021) Time of Day and Muscle Strength: A Circadian Output?
Physiology (Bethesda) 36:44–51. https://doi.org/10.1152/physiol.00030.2020

Enoka RM, Duchateau J (2008) Muscle fatigue: what, why and how it influences muscle function. J
Physiol 586:11–23. https://doi.org/10.1113/jphysiol.2007.139477

Fuglevand AJ, Zackowski KM, Huey KA, Enoka RM (1993) Impairment of neuromuscular propagation
during human fatiguing contractions at submaximal forces. J Physiol (Lond) 460:549–572.
https://doi.org/10.1113/jphysiol.1993.sp019486

Gandevia SC (2001) Spinal and supraspinal factors in human muscle fatigue. Physiol Rev 81:1725–
1789. https://doi.org/10.1152/physrev.2001.81.4.1725

Gandevia SC, Allen GM, Butler JE, Taylor JL (1996) Supraspinal factors in human muscle fatigue:
evidence for suboptimal output from the motor cortex. J Physiol 490:529–536

Garland SJ, Enoka RM, Serrano LP, Robinson GA (1994) Behavior of motor units in human biceps
brachii during a submaximal fatiguing contraction. J Appl Physiol (1985) 76:2411–2419.
https://doi.org/10.1152/jappl.1994.76.6.2411

Garland SJ, Kaufman MP (1995) Role of muscle afferents in the inhibition of motoneurons during
fatigue. Adv Exp Med Biol 384:271–278. https://doi.org/10.1007/978-1-4899-1016-5_21

Girard O, Bishop DJ, Racinais S (2013) Neuromuscular adjustments of the quadriceps muscle after
repeated cycling sprints. PLoS One 8:e61793. https://doi.org/10.1371/journal.pone.0061793

Guette M, Gondin J, Martin A (2005) Time-of-day effect on the torque and neuromuscular properties
of dominant and non-dominant quadriceps femoris. Chronobiol Int 22:541–558.
https://doi.org/10.1081/CBI-200062407

Halperin I, Aboodarda SJ, Basset FA, et al (2014) Pacing strategies during repeated maximal voluntary
contractions. Eur J Appl Physiol 114:1413–1420. https://doi.org/10.1007/s00421-014-2872-3

Hargreaves M (2005) Metabolic factors in fatigue. Sports Science Exchange 98 18:9

Hunter SK, Rochette L, Critchlow A, Enoka RM (2005) Time to task failure differs with load type when
old adults perform a submaximal fatiguing contraction. Muscle Nerve 31:730–740.
https://doi.org/10.1002/mus.20325

Hunter SK, Ryan DL, Ortega JD, Enoka RM (2002) Task Differences With the Same Load Torque Alter
the Endurance Time of Submaximal Fatiguing Contractions in Humans. Journal of
Neurophysiology 88:3087–3096. https://doi.org/10.1152/jn.00232.2002

Hunter SK, Yoon T, Farinella J, et al (2008) Time to task failure and muscle activation vary with load
type for a submaximal fatiguing contraction with the lower leg. J Appl Physiol 105:463–472.
https://doi.org/10.1152/japplphysiol.90398.2008

Kallenberg LAC, Hermens HJ (2006) Behaviour of motor unit action potential rate, estimated from
surface EMG, as a measure of muscle activation level. J Neuroeng Rehabil 3:15.
https://doi.org/10.1186/1743-0003-3-15

Kawakami Y, Amemiya K, Kanehisa H, et al (2000) Fatigue responses of human triceps surae muscles
during repetitive maximal isometric contractions. Journal of Applied Physiology 88:1969–
1975. https://doi.org/10.1152/jappl.2000.88.6.1969

Klass M, Lévénez M, Enoka RM, Duchateau J (2008) Spinal mechanisms contribute to differences in
the time to failure of submaximal fatiguing contractions performed with different loads. J
Neurophysiol 99:1096–1104. https://doi.org/10.1152/jn.01252.2007

Lebesque L, Scaglioni G, Manckoundia P, Martin A (2023) Neuromuscular fatigability of plantar
flexors following continuous and intermittent contractions. Journal of Applied Physiology
134:1093–1104. https://doi.org/10.1152/japplphysiol.00601.2022

Lebesque L, Scaglioni G, Martin A (2022) The impact of submaximal fatiguing exercises on the ability
to generate and sustain the maximal voluntary contraction. Frontiers in Physiology 13:

Mademli L, Arampatzis A (2008) Effect of voluntary activation on age-related muscle fatigue
resistance. J Biomech 41:1229–1235. https://doi.org/10.1016/j.jbiomech.2008.01.019

Maluf KS, Shinohara M, Stephenson JL, Enoka RM (2005) Muscle activation and time to task failure
differ with load type and contraction intensity for a human hand muscle. Exp Brain Res
167:165–177. https://doi.org/10.1007/s00221-005-0017-y

Marcora S (2010) Effort: Perception of. In: Goldstein E (ed) Encyclopedia of Perception, Encyclopedia
of Perception. SAGE Publications, Inc., Los Angeles, pp 380–383

Matkowski B, Martin A, Lepers R (2011) Comparison of maximal unilateral versus bilateral voluntary
contraction force. Eur J Appl Physiol 111:1571–1578. https://doi.org/10.1007/s00421-010-
1775-1

Matta P, Glories D, Alamia A, et al (2023) Mind over muscle? Time manipulation improves physical
performance by slowing down the neuromuscular fatigue accumulation. Psychophysiology
e14487. https://doi.org/10.1111/psyp.14487

Mira J, Lapole T, Souron R, et al (2017) Cortical voluntary activation testing methodology impacts
central fatigue. European Journal of Applied Physiology 117:1845–1857.
https://doi.org/10.1007/s00421-017-3678-x

Morey RD, Rouder JN (2011) Bayes factor approaches for testing interval null hypotheses. Psychol
774   Methods 16:406–419. https://doi.org/10.1037/a0024377

775 Neyroud D, Rüttimann J, Mannion AF, et al (2013) Comparison of neuromuscular adjustments
776   associated with sustained isometric contractions of four different muscle groups. Journal of
777   Applied Physiology 114:1426–1434. https://doi.org/10.1152/japplphysiol.01539.2012

778 Pageaux B (2016) Perception of effort in Exercise Science: Definition, measurement and perspectives.
779   European Journal of Sport Science 16:885–894.
780   https://doi.org/10.1080/17461391.2016.1188992

781 Place N, Maffiuletti NA, Ballay Y, Lepers R (2005) Twitch potentiation is greater after a fatiguing
782   submaximal isometric contraction performed at short vs. long quadriceps muscle length. J
783   Appl Physiol (1985) 98:429–436. https://doi.org/10.1152/japplphysiol.00664.2004

784 Place N, Martin A, Ballay Y, Lepers R (2007) Neuromuscular fatigue differs with biofeedback type
785   when performing a submaximal contraction. J Electromyogr Kinesiol 17:253–263.
786   https://doi.org/10.1016/j.jelekin.2006.04.001

787 Place N, Martin A, Lepers R (2006a) Changes in neuromuscular function after tasks involving control
788   of EMG versus torque feedback of the same duration. Neurosci Lett 394:37–41.
789   https://doi.org/10.1016/j.neulet.2005.10.013

790 Place N, Matkowski B, Martin A, Lepers R (2006b) Synergists activation pattern of the quadriceps
791   muscle differs when performing sustained isometric contractions with different EMG
792   biofeedback. Exp Brain Res 174:595–603. https://doi.org/10.1007/s00221-006-0504-9

793 Place N, Millet GY (2019) Quantification of Neuromuscular Fatigue: What Do We Do Wrong and
794   Why? Sports Med 50:439–447. https://doi.org/10.1007/s40279-019-01203-9

795 Rohmert W (1960) Determination of the recovery pause for static work of man. Int Z Angew Physiol
796   18:123–164

797 Rudroff T, Christou EA, Poston B, et al (2007) Time to failure of a sustained contraction is predicted
798   by target torque and initial electromyographic bursts in elbow flexor muscles. Muscle &
799   Nerve 35:657–666. https://doi.org/10.1002/mus.20752

800 Rudroff T, Justice JN, Holmes MR, et al (2011) Muscle activity and time to task failure differ with load
801   compliance and target force for elbow flexor muscles. J Appl Physiol 110:125–136.
802   https://doi.org/10.1152/japplphysiol.00605.2010

803 Rudroff T, Justice JN, Matthews S, et al (2010) Muscle activity differs with load compliance during
804   fatiguing contractions with the knee extensor muscles. Exp Brain Res 203:307–316.
805   https://doi.org/10.1007/s00221-010-2233-3

806 Sahlin K (1992) Metabolic Factors in Fatigue. Sports Medicine 13:99–107.
807   https://doi.org/10.2165/00007256-199213020-00005

808 Shahidi AV, Mathieu PA (1995) Endurance time characteristics of human ankle dorsiflexors and
809   plantarflexors. Europ J Appl Physiol 71:124–130. https://doi.org/10.1007/BF00854968

810 Skau S, Sundberg K, Kuhn H-G (2021) A Proposal for a Unifying Set of Definitions of Fatigue. Frontiers
811   in Psychology 12:

Smith CM, Housh TJ, Hill EC, et al (2016) Effects of fatiguing constant versus alternating intensity
intermittent isometric muscle actions on maximal torque and neuromuscular responses. J
Musculoskelet Neuronal Interact 16:318–326

Søgaard K, Gandevia SC, Todd G, et al (2006) The effect of sustained low-intensity contractions on
supraspinal fatigue in human elbow flexor muscles. J Physiol (Lond) 573:511–523.
https://doi.org/10.1113/jphysiol.2005.103598

Strojnik V, Komi PV (1998) Neuromuscular fatigue after maximal stretch-shortening cycle exercise. J
Appl Physiol 84:344–350. https://doi.org/10.1152/jappl.1998.84.1.344

Tamm AS, Lagerquist O, Ley AL, Collins DF (2009) Chronotype Influences Diurnal Variations in the
Excitability of the Human Motor Cortex and the Ability to Generate Torque during a
Maximum Voluntary Contraction. J Biol Rhythms 24:211–224.
https://doi.org/10.1177/0748730409334135

Taylor JL, Amann M, Duchateau J, et al (2016) Neural Contributions to Muscle Fatigue: From the
Brain to the Muscle and Back Again. Med Sci Sports Exerc 48:2294–2306.
https://doi.org/10.1249/MSS.0000000000000923

van Melick N, Meddeler BM, Hoogeboom TJ, et al (2017) How to determine leg dominance: The
agreement between self-reported and observed performance in healthy adults. PLoS One
12:e0189876. https://doi.org/10.1371/journal.pone.0189876

Williams PS, Hoffman RL, Clark BC (2014) Cortical and spinal mechanisms of task failure of sustained
submaximal fatiguing contractions. PLoS One 9:e93284.
https://doi.org/10.1371/journal.pone.0093284

Yoon T, Hawe R, Hunter SK (2009) Variation in limb support influences the time to task failure for a
postural contraction. J Mot Behav 41:393–395. https://doi.org/10.3200/35-09-003

FIGURE CAPTIONS

Figure 1 Sum up of the experimental design of the study. A. The protocol consisted

in a $MVC_{1\text{-}MIN}$ performed before (Pre), after (Post) the fatiguing exercise and after a

10-min recovery period (Post-10). The fatiguing exercise consisted of INT, CON or

VAR contractions performed with an equivalent MET (30% MVC) until TF. Note that

brief MVCs were performed only at Pre and Post-10. Arrows represent electrical

doublets (100 Hz) delivered during the protocol. B. Experimental data of torque

during the $MVC_{1\text{-}MIN}$ from one typical subject. Main neuromuscular measurements are

reported (MVC and ΔMVC_{1-MIN}, VAL and VAL end, Pd_B and Pd_A). Note that Pd_B was

obtained from doublet delivered before the MVC_{1-MIN}, only at Post. Pd_B at Pre and

Post-10 were obtained after the brief MVC. Arrows represent electrical doublets (100

Hz) delivered at rest and superimposed to the contraction. CON = continuous

exercise; INT = intermittent exercise; MET = mean exercise torque; MVC = maximal

voluntary contraction; MVC_{1-MIN} = 1-min sustained MVC; Pd_A = potentiated doublet

after the MVC_{1-MIN}; Pd_B = potentiated doublet before the MVC_{1-MIN}; TF = task failure;

VAL = voluntary activation level; VAR = variable exercise; ΔMVC_{1-MIN} = torque loss

during the MVC_{1-MIN}

Figure 2 Individual repeated-measures correlation between effective ED and change

in electromyographic activities during exercise. The lines correspond to the repeated-

measures correlation fit for each participant (n = 13). For each fatiguing exercise,

data from the same participant are shown in the same colour. $\Delta RMS/M_{MAX}$

represents the change in muscle activities, normalized by M_{MAX}, between the

beginning and the end of the fatiguing exercise. ED = exercise duration; M_{MAX} = sum

of maximal M-wave of triceps surae muscles; RMS = sum of electromyographic root

mean square of triceps surae muscles. $p < 0.050$ expresses a significant correlation

Figure 3 Experimental torque and electromyographic traces from a typical subject

during each fatiguing exercise performed until task failure. INT = intermittent exercise

(15 s on, 5 s off) at 40% MVC; CON = continuous exercise at 30% MVC; VAR =

exercise with a variable contraction intensity (alternation of 10 s at 40% MVC and 10

s at 20% MVC); LG = lateral gastrocnemius; MG = medial gastrocnemius; MVC =

maximal voluntary contraction; SOL = soleus

Figure 4 Maximal torque production, voluntary activation and potentiated doublet measured before (Pre, white violin plots) and after (Post, dark grey violin plots) the fatiguing exercise, as well as after a 10-min recovery period (Post-10, light grey violin plots). Pooled data from the three exercises are violin plotted at each time point and the horizontal black lines represent mean values. Individual data from INT (circles), CON (squares) and VAR (triangles) exercises are also presented ($n = 13$). **A**. Changes in MVC. **B**. Changes in VAL computed at the beginning of the MVC_{1-MIN}. **C**. Changes in Pd_B. CON = continuous exercise; INT = intermittent exercise; MVC = maximal voluntary contraction; MVC_{1-MIN} = sustained MVC for 1 min; Pd_B = potentiated doublet before the MVC_{1-MIN}; VAL = voluntary activation level; VAR = variable exercise. $^*p < 0.05$, $^{***}p < 0.001$ express a significant difference

Figure 5 Maximal torque sustainability and associated variables measured before (Pre, white violin plots) and after (Post, dark violin plots) the fatiguing exercises, and following a 10-min recovery period (Post-10, light grey violin plots). Pooled data from the three exercises are violin plotted at each time point and the horizontal black lines represent mean values. Individual data from INT (circles), CON (squares) and VAR (triangles) exercises are also presented ($n = 13$). **A.** ΔMVC_{1-MIN} represents the percentage of torque loss during each MVC_{1-MIN}. **B**. ΔVAL_{1-MIN} represents the percentage of VAL change during each MVC_{1-MIN}. **C**. ΔPd_{1-MIN} represents the percentage of change in Pd during each MVC_{1-MIN}. CON = continuous exercise; INT = intermittent exercise; MVC_{1-MIN} = 1-minute maximal voluntary contraction; Pd = potentiated doublet; VAL = voluntary activation level; VAR = variable exercise. $^{**}p < 0.01$, $^{***}p < 0.001$ express a significant difference

INT **CON** **VAR**

Torque 25 Nm

SOL 2 mV

LG 2 mV

MG 2 mV

30 s

Table 1 Fatiguing exercise parameters.

	INT	CON	VAR
Total ED (s)	1042.9 ± 546.0	506.2 ± 200.7***	538.4 ± 270.1***
Effective ED (s)	783.4 ± 408.8	506.2 ± 200.7***	538.4 ± 270.1**
Exercise TTI (Nm·s)	35836 ± 14117	17492 ± 5432***	19187 ± 8542***
MET (Nm)	36.2 ± 5.7	35.8 ± 5.0	37.2 ± 6.5
MET (% MVC)	28.9 ± 0.8	28.4 ± 1.5	28.7 ± 1.1
RMS/M_{MAX} beginning (x10^{-3})	9.05 ± 2.98	5.84 ± 1.90**	6.45 ± 1.70*
RMS/M_{MAX} end (x10^{-3})	13.03 ± 4.36	12.88 ± 2.96	12.07 ± 3.58
RPE	97.8 ± 7.4	100.1 ± 7.2	95.5 ± 4.3

CON = continuous exercise; ED = exercise duration; INT = intermittent exercise; MET = mean exercise torque; M_{MAX} = sum of triceps surae maximal M-wave; RMS = sum of triceps surae electromyographic root mean square; RPE = rate of perceived exertion; VAR = variable exercise. The RMS/M_{MAX} ratio was calculated at both the beginning and the end of the fatiguing exercise. $n = 13$. *$p < 0.05$, **$p < 0.01$ and ***$p < 0.001$ represent significant difference from INT value