The effect of altitude on the expression of immune-related genes in Peruvian rural

indigenous

Authors: Luis Jaramillo-Valverde[1,2,3+], Gilderlanio Santana de Araújo[1,4+], Julio A.

Poterico[1,2], Catalina Martinez-Jaramillo[1], Vicky Roa-Linares[1], Sandra Alvites-Arrieta[1], Nelis

Pablo-Ramirez[1], Milward Ubillus[1], Diana Palma-Lozano[1], Rafael Tou[5], Carolina Silva-

Carvalho[5], Luca Vasconcelos-da-Gama[5], Lucas F Costa[5], Manuel Corpas[6,7,8], Eduardo

Tarazona-Santos[5], Soumya Raychaudhuri[9], Heinner Guio[1,2*]

[1]Universidad de Huánuco, Huánuco, Peru

[2]INBIOMEDIC, Research and Technological Center, Lima, Peru

[3]School of Medicine, Universidad Continental, Huancayo, Perú

[4]Instituto de Ciências Biológicas, Universidade Federal do Pará, PA, Brazil

[5]Departamento de Genética, Ecologia e Evolução, Instituto de Ciências Biológicas, Belo

Horizonte, Universidade Federal de Minas Gerais, MG, Brazil

[6]Life Sciences, University of Westminster, London, UK

[7]The Alan Turing Institute, London, UK

[8]Cambridge Precision Medicine Limited, ideaSpace, University of Cambridge Biomedical

Innovation Hub, Cambridge, UK

[9]Department of Biomedical Informatics, Harvard Medical School, Boston, MA, USA

[+]Equal contribution

*Corresponding Author:

E-mail: heinnerguio@gmail.com

AUTHOR CONTRIBUTIONS

Study design: LJ-V, SR and HG. Performed the experiments: LJ-V, GSA, JAP, CMJ, VRL,

SAA, NPR, MU, DPL. Analyzed the data: LJ-V, GSA, RT, CSC, LVG, LFC, MC, ETS, SR

and HG All authors have read and approved the final manuscript.

ACKNOWLEDGEMENTS

We thank Lluis Quintana-Murci and David Comas for their constructive comments and

perspectives. We thank all those who facilitated the recruitment of participants, including the

Direccion Regional de Salud from Huanuco; and all participants in this study.

FUNDING INFORMATION

The Programa Nacional de Investigación Científica y Estudios Avanzados – Prociencia with contract number PE501084276-2023 supported this study.

CONFLICT OF INTEREST STATEMENT

The authors declare that they have no conflicts of interest.

DATA AVAILABILITY STATEMENT

Data have been deposited in the European Genome-phenome Archive (EGA), https://www.ebi.ac.uk/ega/home (accession nos. PRJEB79378). Access to the data is available upon request from the authors. Please contact the corresponding authors for the access directions.

The numerical data, graphs, scripts, and summary are thoroughly referenced in the main text and are readily accessible online.

ORCID

Luis Jaramillo-Valverde https://orcid.org/0000-0001-5093-6067

Gilderlanio Santana de Araújo https://orcid.org/0000-0001-9199-9419

Julio A. Poterico https://orcid.org/0000-0001-7838-3505

Catalina Martinez-Jaramillo https://orcid.org/0000-0001-6846-0140

Vicky Roa-Linares https://orcid.org/0000-0001-9253-615X

Sandra Alvites-Arrieta -

Nelis Pablo-Ramirez -

Milward Ubillus https://orcid.org/0000-0002-3684-9394

Diana Palma-Lozano https://orcid.org/0000-0003-4520-7374

Rafael Tou - https://orcid.org/0000-0003-0907-1510

Carolina Silva-Carvalho https://orcid.org/0000-0002-6356-6779

Luca Vasconcelos-da-Gama -

Lucas F Costa https://orcid.org/0000-0001-9363-0121

Eduardo Tarazona-Santos https://orcid.org/0000-0003-3508-3160

Soumya Raychaudhuri https://orcid.org/0000-0002-1901-8265

Heinner Guio https://orcid.org/0000-0003-0078-1188

ABSTRACT

Background
Genetic factors influencing immune response pathways in Andean populations may underlie adaptations to high-altitude environments. To investigate transcriptomic signatures associated with altitude, we analyzed immune-related gene expression across individuals residing at different elevations.

Methods
We recruited 62 Peruvian volunteers, predominantly from rural regions with high proportions of indigenous ancestry, living at low and high altitudes. Peripheral blood mononuclear cells (PBMCs) were stimulated with bacterial lipopolysaccharide (LPS), Pam3CSK4 (a synthetic triacylated lipopeptide), and R848 (an imidazoquinoline analog of viral nucleic acids). Population structure and ancestry were characterized, and transcriptome-wide differential expression analyses were performed.

Results
We identified 30 genes with significant altitude-associated expression differences, including 22 downregulated (e.g., *FN1*, *CD36*, *FOS*) and nine upregulated genes. Functional enrichment indicated roles in acute inflammatory response, leukocyte migration, and positive regulation of myeloid leukocyte differentiation.

Conclusions

High- and low-altitude Andean individuals exhibit distinct immune gene expression profiles, defining a population-specific transcriptomic signature that may reflect altitude-related immune adaptation.

Keywords: altitude, expression, immune-related genes, rural indigenous

AUTHOR SUMMARY

Andean populations exhibit genetic adaptations in immune response pathways, potentially linked to high-altitude living. To investigate this, we conducted genome-wide and transcriptome-wide analyses to identify immune-related gene expression differences between high- and low-altitude residents. We analyzed the genetic structure and ancestry of Peruvian individuals (primarily rural, with strong indigenous ancestry) living at different altitudes. Peripheral blood mononuclear cells (PBMCs) from 62 volunteers were stimulated with bacterial (LPS, Pam3CSK4) and viral (R848) mimics to assess immune responses. Differential expression analysis identified 22 down-regulated genes (e.g., *FN1*, *CD36*, *FOS*) and nine up-regulated genes, enriched in acute inflammatory response, leukocyte migration, and myeloid leukocyte differentiation. Our findings reveal a unique immune gene expression profile in Andean highlanders, distinct from other populations, suggesting adaptive modulation of both innate and adaptive immunity in response to high-altitude challenges.

INTRODUCTION

Approximately 1% of the world's population resides in high-altitude areas, specifically at elevations over 2500 meters above sea level (m.a.s.l.). Countries such as Peru and Bolivia have a significant portion (30 percent) of their population living at high altitudes in the Andean mountains (1). Living in high-altitude regions poses several environmental challenges for humans, including reduced oxygen levels, increased exposure to ultraviolet radiation, cold temperatures, and altered responses to pathogens, among other factors (2). Pioneering researchers in the field of population genomics at high altitude have suggested that genes of the Hypoxia Inducible Factor (HIF) pathway have evolved under positive natural selection (3,4). In contrast, the genes *ENDRA*, *PRKAA1*, and *NOS2A* have been suggested as being associated with blood pressure regulation and reproductive success in Andeans (5). Additionally, genetic studies have demonstrated that native Andean highlanders depict positive selection of cardiovascular genes, such as *BRINP3*, *NOS2*, *TBX5 (6)*, and the non-coding region *HAND2-AS1* (7).

As previously mentioned, hypoxia impairs oxidative metabolism, which forces cells to rely on oxygen-independent processes like glycolysis to produce ATP. Hypoxia plays a crucial role in immune cell metabolism regulation, which serves as a major regulator of immunometabolism, in addition to controlling immune cell phenotypic and function (8,9). Numerous studies have shown that hypoxia regulates cell types involved in the innate and adaptive immune response, including epithelial cells, neutrophils, macrophages, dendritic cells, T cells, B cells, natural killer cells, and innate lymphoid cells, since the first report of a role for HIF in the control of macrophage function in 2003 (10). A recent study confirms these findings and provides further insights. Analyses of a genome-wide study involving 286 individuals across Peru and over-representation functional analysis identified ten positive natural selection signals among highlander populations related to cardiovascular function (*TLL1*, *DUSP27*, *TBX5*, *PLXNA4*, *SGCD*), the hypoxia-inducible factor pathway (*TGFA*, *APIP*), skin pigmentation (*MITF*), glucose (*GLIS3*), and glycogen metabolism (*PPP1R3C*, *GANC*) (11), as well as genes involved in the *TP53* pathway, such as *SP100*, *DUOX2,* and *CLC (12)*.

These studies indicate that Peruvian Andean populations have exhibited distinct adaptations compared to other groups like Tibetans, Sherpas, or Ethiopians in response to high-altitude

living (11,12). Moreover, specific genes related to immune response have been identified in

Andean populations as being positively selected. These genes might provide insights into the

adaptation of the immune system and responses to pathogens among those residing at high

altitudes. Examples of robust adaptation signals related to altitude in Peruvian and Bolivian

populations can be found in the studies conducted by Borda et al. (7) and Jacovas et al. (12).

Specifically, these studies identified a significant association with the *DUOX2* gene, which

plays a crucial role in maintaining homeostasis in thyroid hormone production, as well as

influencing the innate immune system and inflammatory response.

Likewise, Caro-Consuegra et al. found positive selection in Peruvian Andean populations in

CD40, *TNIP1*, and *CHIA* genes, which are involved in the innate immune system (11).

Furthermore, genes such as *ANXA1*, *CD44*, *ADGRE1*, *KCNA3,* and *FCRL4,* among others,

seem to have been positively selected due to *Mycobacterium tuberculosis* in Andean

Ecuadorian populations, and this event could have emerged before the European contact with

aboriginal communities (13). However, the aforementioned genomic studies have not

included functional analysis to correlate genomic findings with other biomarkers and

phenotypes.

To our knowledge, there is currently no investigation assessing the expression of immune

genes in native participants (*e.g.*, Andean Peruvians) with long-term exposure to high

altitude–*i.e.,* residence since birth or childhood. For that reason, after characterizing the

genetic structure and ancestry of the studied population, we aim to conduct transcriptome

analyses of blood cells in Peruvian individuals, primarily from rural areas, from indigenous

populations. These individuals are exposed to varying living altitudes, comparing individuals

from high versus low altitudes. We collected peripheral blood mononuclear cells (PBMC)

from 62 volunteers, who were exposed to bacterial lipopolysaccharide (LPS), Pam3CSK4 (a

synthetic triacylated lipopeptide responsible for bacterial components), and R848 (an

imidazoquinoline compound related to viral nucleic acids) (14). Subsequently, we discerned

variations in gene expression profiles among populations residing at low and high altitudes

(hypoxia). Our analysis primarily concentrated on genes that exhibited differential expression

regardless of stimuli, with a special focus on those significantly involved in immune

responses.

MATERIAL AND METHODS

Population and samples

We studied the population from Huanuco, a department in Central Peru, focusing on a region in the eastern slope of the Andes that includes the Andean Mountains and the Amazonian Tropical Forest (in particular the so-called Amazon Yunga or Higher Amazonia). In addition to biological samples, we collected demographic data such as sex, age, body mass index (BMI), and altitude of residence. In the first part of the study, 35 individuals from the studied population were analyzed for genomic diversity. In the second part of the study, we studied 62 individuals for transcriptome analyses, with an overlap of 19 individuals between the two parts of the study. The study was approved by the Inmensa Ethics Committee of Peru (number 0011-2021). Written informed consent was obtained from all participants involved in this study.

Genetic structure and ancestry of the studied Huanuco population

Genotyping and Data Quality Control (QC). Thirty-five individuals were genotyped with the Illumina GSA MG v3 array by Macrogen (Seul, Korea). After Illumina GenomeStudio processing, we obtained 737,534 SNVs. After further QC with MosaiQC software (https://github.com/ldgh/Smart-cleaning-public), a tool that automates the Laboratory of Human Genetic Diversity (LDGH) quality control pipeline using PLINK, 703,515 SNVs remained.

Individual ancestry analyses. For the genetic clustering analysis, we ran ADMIXTURE (15), which estimates the fraction of each K parental population that contributes to an individual by fitting the Hardy-Weinberg equilibrium in each of the K populations/clusters (7), using 189,837 unlinked SNVs ($r^2<0.4$). To cover the admixture history and the structure inside Peruvian Native populations, we first ran 14 replicates in the unsupervised mode for K=3 to infer African, European and Native American ancestry using the entire dataset in Supplementary Table S1. We presented the replicate with the higher log-likelihood value.

To avoid the bias inserted by related individuals in ADMIXTURE inferences, we used K=3 to estimate kinship coefficients between Huánuco individuals (Φij) with the software REAP (Relatedness Estimation in Admixed Populations), which is more appropriate for admixed

individuals (16). The estimates of Φij by REAP are conditioned on the estimated individual

ancestry proportions (estimated with ADMIXTURE) for K=3 parental populations. To

identify related individuals to be pruned for further ancestry analyses (K>3), we used

NAToRA, a relatedness-pruning method to minimize the loss of dataset in genetic and omics

analyses (17), using the relatedness cutoff of $\Phi ij > 0.06$.

Then, we pruned our dataset for the few related individuals and ran again 14 replicates of

ADMIXTURE for K from 4 to 12, considering only unrelated individuals and 162,957

unlinked SNVs, presenting the results of the most likely run for each K, and assessing CV

errors for different K. Because the ADMIXTURE results with K=3 informed us that our

dataset had negligible African ancestry, we removed African individuals from the

ADMIXTURE analyses for K>3.

Transcriptome analyses in the Huanuco population

<u>Preparation of biological samples.</u> In 62 individuals, PBMCs were isolated from whole

blood by Ficoll-Paque centrifugation within 4 hours of collection and cryopreserved. PBMCs

were cultured in RPMI-1640 (Fisher) supplemented with 10% heat-inactivated FBS (FBS

premium, US origin, Wisent) and 1% L-glutamine (Fisher). All samples were stimulated by

pathogen molecules (virus and bacteria) in independent processes: for each of the tested

individuals, and one process without stimulation.

Bacterial and Virus Stimulation Assays PBMCs (2 million per condition) were stimulated for

4 hours at 37° C with 5% CO_2 with bacterial lipopolysaccharide (LPS), Pam3CSK4

(Invitrogen – tlrl-pms) a synthetic triacylated lipopeptide responsible for bacterial

components, and R848 (Invitrogen – tlrl-r848-5) an imidazoquinoline compound, related to

viral nucleic acids, according to the manufacturer's instructions.

RNA Extraction Total RNA was extracted with the PureLink RNA Mini Kit from Invitrogen

Inc., including the enzymatic digestion of genomic DNA. Extractions were performed in

batches of 10 samples, and RNA quality and quantity were assessed with a Nanodrop

spectrometer and the Agilent Bioanalyzer RNA 6000 nano kit. We generated a final set of 79

samples from the 62 donors for high-throughput RNA-sequencing, including 17, 16, 24, and

22 samples for the non-simulated, LPS, Pam3CSK4, and R848 conditions, respectively.

<u>RNA Library Construction, Quality Control, and Sequencing.</u> The process of library

construction involved the purification of messenger RNA from total RNA using magnetic

beads attached to poly-T oligos (TruSeq RNA Sample Prep Kit v2, Illumina Inc.).
Subsequently, the mRNA was fragmented, and the first strand of cDNA was synthesized
using random hexamer primers. This was followed by the synthesis of the second strand
cDNA. The library was considered complete after undergoing end repair, A-tailing, adapter
ligation, size selection, amplification, and purification. Quantification and size distribution
analysis of the library was performed using Qubit and real-time PCR, as well as a
Bioanalyzer (Agilent, CA, USA). Finally, the pooled libraries were sequenced using Illumina
platforms by the NOVAGENE facility (https://www.novogene.com/). The total data output
sequencing product was 514.1 GB. The Data Quality report and summary are available in the
Supporting Information and Data.

Transcriptome data pre-processing. Total RNA sequencing was performed by a pair-end
strategy for all samples using Novaseq PE150 (Illumina Inc). Raw data is stored in .fastq
format (.fq.gz compressed) for analysis. The NOVOGENE facility provides raw RNA
sequencing data by demultiplexing (separating multiple samples pooled into a single library),
removing adapters, and low-quality reads regarding bases that cannot be determined (N >
10%) and those with QScore <= 5. For our quality control analysis, each sample in .fastq was
resubmitted for the quality control pipeline using fastQC and multiQC to generate sequencing
quality reports to verify quality per bases, GC content, the proportion of undefined nucleotide
bases, and the presence of adapter sequences. The Phred score Q30 was the base threshold for
filtering high-quality samples. Read alignment was performed with STAR with the Release
41 (GRCh38.p13) genome reference. The software STAR was set to consider that a read may
overlap on the same or opposite strand and to follow the Encyclopedia of DNA Elements
(ENCODE) best practices. The STAR outputs alignments in a sorted BAM format. After
genome alignment, read quantification was performed with HTSeq with the GRCh38 human
genome annotation file (.gff). The read-counting process considered the large set of
transcripts, in total 61801 coding and non-coding elements cataloged in ENCODE. To avoid
non-expressed transcripts, we used the *filterByExpr()* function to filter for genes that meet a
minimum expression threshold. In this analysis, only genes with at least 10 reads across all
samples are retained.

RNA-seq variant calling and global ancestry analysis. Some individuals were sequenced
for RNA-seq multiple times under different stimuli; thus, we merged the corresponding
FASTQ files for each individual. We then used GATK's Best Practices pipeline

(https://github.com/gatk-workflows/gatk4-rnaseq-germline-snps-indels), incorporating steps

for joint variant calling. Variants were filtered using the recommended parameters: a

minimum depth of 5 and up to 10% missing data, resulting in 85,557 variants. For ancestry

inference, we selected 11,861 unlinked SNPs ($r^2 < 0.4$) and performed ADMIXTURE

analysis (Alexander, Novembre, and Lange, 2009) in unsupervised mode, testing K values

from 3 to 12 with three replicates each. African individuals were excluded from panels for

subsequent ADMIXTURE analyses, due to African ancestry showing negligible

contributions. Using the resulting set of variants and quality control filters, we conducted

principal component analysis (PCA) using EIGENSOFT (18,19) to compute 12 principal

components.

Differential gene expression analysis. The 17 control samples (i.e. not stimulated) and 62

treated PBMC samples LPS (n = 16), Pam3CSK4 (n = 24), R848 (n = 22)) were also

organized in two groups: low altitude (less than 1000 m.a.s.l.) and high altitude (>3000

m.a.s.l.) to perform differential expression analysis (DEA) with the edgeR package (20)

Gene expression variance was estimated by variance partition using a linear mixed model

(21). Furthermore, the dispersion of gene expression was assessed to fit a generalized linear

model including covariate correction for individual (categorical variable), sex (categorical

variable), BMI (numerical variable), age (numerical variable), stimulus by pathogens (one

virus and two bacteria, categorical variable), and Native ancestry proportion. Those genes

with a false-discovery rate (FDR < 0,05) and fold-change (|FC| >= 2) were considered

differentially expressed genes (DEG). The distribution of DEG genes (down- and up-

regulated) regarding statistical significance and genes from the immune system, before and

after stimulation, was visualized using volcano plots and heatmaps, created with the

EnhancedVolcano (https://github.com/kevinblighe/EnhancedVolcano) and pheatmap

packages in R (https://cran.r-project.org/package=pheatmap). Hierarchical clustering of genes

and samples was performed and visualized on heat maps using Pearson correlation and the

complete linkage method (19, 20).

Functional Enrichment Analysis. We conducted a thorough functional enrichment analysis

of gene sets, encompassing both up-regulated and down-regulated genes, employing Gene

Ontology (GO) and utilizing the ClusterProfile (22) package in the R language. To enhance

statistical rigor, multiple tests were corrected using FDR. The outcomes, comprising the top

biological processes, were visually represented through bar plots and networks depicting the

interrelation of genes and processes.

Immune system genes dataset. DEA results were merged with biological pathway data,

focusing exclusively on genes related to the Immune System. In the Reactome database, the

adaptive *immune system* catalogs the interaction of 868 genes, the *innate immune system*

shows 724 transcripts, and 1160 genes comprise the *cytokine signaling pathway*. The

Reactome data is available at reacomte.org, and we used version 81 released on June 16,

2024.

RESULTS

Genomic diversity and ancestry of the Huanuco population

ADMIXTURE analyses (Figure 1, K=3 and 4) of 35 individuals reveal that Huanuco

individuals are predominantly Native American (87.3% on average) with a low European

ancestry (average: 11.82%) and negligible African ancestry (<1%) (Supplementary Table

S2). Given the negligible African ancestry, which is consistent with the history of the

Huanuco region, we present ADMIXTURE results excluding African individuals (results in

Supplementary Table S3).

Analyses assuming K=5 ancestry clusters (K=4 and 5) reveal that Native American ancestry

of Huánuco individuals may be dissected in a predominant Andean-associated cluster (84.5%

[SD:5.6%] of the total ancestry) and a minor Amazonian-associated cluster (3.7% [SD: 1.1%]

of the total ancestry). Interestingly, the Andean-associated ancestry cluster is higher in

highlanders (average 86%) than in lowlanders (average 82%, <600 m vs. >3000m, Kruskal-

Wallis H=4.057, $p<0.05$, N=35).

A higher-resolution clustering (K = 6 and 7) reveals that Huanuco individuals are similar to

Tallanes and Moche (indigenous groups from the northern Peruvian Coast) and Chacapoyas

(an indigenous group from the north of the Andes/Amazonian geographic interface).

Altogether, Huanuco individuals are similar to Tallanes, Moche, and Chachapoyas from

Northern Peru, and all of them present different levels of admixture between Andean and

Amazonian ancestries.

Figure 1. Genetic and geographic landscapes for Peruvian populations. (A): Genetic structure of seventeen native populations from the coast (n. 1 and 2), Amazon (n. 3-11), Andes (n. 12-17), and Huanuco samples (18A and 18B), inferred by ADMIXTURE (K=6). Donut plots show the average percentages of individuals for six ancestry clusters in each population. Four clusters were associated with distinct Native American groups: one Andean (brown), one North Amazonian (light green), one South Amazonian (dark green), and one coast (purple). Two clusters were associated with non-Native continental ancestries, both European (red), caused by admixture with these populations in the last 500 years. (B): Information for geographic location and altitude of samples. The color of the population markers goes from white (at sea level) to black (5000 meters above sea level). (https://www.naturalearthdata.com/)

Demographic data overview of the Huanuco transcriptome study

Forty-three (69.4% or total) participants were highland sojourners (3226 – 3539 m.a.s.l.), whereas 19 (30.6%) inhabited lowlands (596 – 634 m.a.s.l.). The samples were obtained at

the residences of participants and comprised 11 males (11.7%) and 51 females (82.3%) with

an age mean of 48 years (SD±15). The samples showed a BMI median of 26.4 Kg/m^2. It is

worth noting that almost all participants (high and lowlanders) had ancestors (e.g., parents,

grandparents) from high Andean areas of the Huánuco department (Table 1).

Table 1. Demographic data of Peruvian individuals from rural areas, categorized by different living

altitudes (high vs low).

Variables	Total		Living altitudes				p-Value
			Low		High		
	N	%	N	%	N	%	
Sex							
Female	51	82.26	15	78.95	36	83.72	
Male	11	17.74	4	21.05	7	16.28	0.650
Age (years)	-	-	42	(34 - 58) †	48	(37 - 64) †	0.392
Cholesterol (mg/dl)	-	-	169	(136 – 185)†	166	(145 – 204)†	0.262
Hemoglobin (g/dl)	-	-	14.4	(13.4 - 15.1)†	14.1	(13.4 - 14.4)†	0.267
Glucose (mg/dl)	-	-	87	(78 - 96)†	83	(76 – 89)†	0.184
BMI (kg/m^2)	-	-	28.3	(24.3 - 30.2)†	25.1	(22.3 - 28.5)†	0.031

*Median. †(Q1 – Q3)
BMI (Body mass index)
Statistically significant (p < 0.05, Chi-squared).

Differentially expressed immune genes in lowlanders versus highlanders

After applying the count filtering criterion, the number of transcripts is reduced from 61,806

to 14,832, effectively avoiding transcripts inflated with low expression (counts < 10) per

sample. After quality control including contamination check, the distribution of samples

stratified for stimulation and altitude is detailed in Supplementary Table S4.

Gene expression variance is primarily driven by individual-specific differences (21.80%),

followed closely by stimulus group effects (16.99%). Moderate contributions come

from altitude (5.45%) and age (2.49%). Smaller but significant influences include population-

specific variation (1.71%), sex differences (1.54%), and body mass index (1.47%). Together,

these factors highlight the complex interplay of genetic, environmental, and demographic

factors shaping gene expression variability (Supplementary Figure S2).

Comparative analysis between individuals from low and high altitudes revealed 31

differentially expressed genes (DEGs), accounting for covariates such as individual differences, sex, BMI, age, Native American ancestry proportions, and pathogen exposure (both viral and bacterial). Among these DEGs, 22 were up-regulated and nine were down-regulated in high-altitude individuals compared to those at low altitude. Notably, three immune-related genes (*FN1, CD36, FOS*) were significantly down-regulated in high-altitude individuals (Figure 2A, Table 2). All DEGs are listed in Supplementary Table S5.

Table 2 – Differential gene expression analysis results.

ENSEMBLE / ID	LogFC	FDR	SYMBOL	EXPRESSION
ENSG00000115414.21	-5.01	3.609194e-15	*FN1*	Down in low altitude
ENSG00000135218.19	-4.09	6.035271e-04	*CD36*	Down in low altitude
ENSG00000170345.10	-3.37	2.291119e-02	*FOS*	Down in low altitude

We performed hierarchical clustering on DEGs identified between lowlanders versus highlanders samples (Figure 2B). The resulting dendrogram revealed two distinct clusters of genes. Furthermore, the sample dendrogram showed a separation between lowlander and highlander groups, indicating distinct gene expression profiles. The heatmap visualization, with dendrograms overlaid, provided a detailed view of the expression patterns, highlighting key clusters of DEGs. Despite accounting for covariates like sex, BMI, age, Native American ancestry proportions, and pathogen exposure, there might be additional unmeasured factors influencing gene expression, causing some samples to cluster differently.

Functional enrichment by gene ontology analysis sheds light on the molecular landscape underlying cellular responses, particularly in the context of immune and inflammatory processes (see Figure 2C-H). The complete functional enrichment analysis results are listed in Supplementary Table S6-S8.

The downregulated genes show a strong enrichment for immune and inflammatory processes, including acute-phase and acute inflammatory responses, leukocyte migration, and myeloid lineage differentiation. Several terms also relate to cell adhesion, suggesting a potential decrease in cell–matrix or cell–substrate interactions. Together, these patterns indicate a suppression of inflammatory signaling and myeloid cell activity, which could reflect reduced immune cell infiltration and altered immune responsiveness.

412 In contrast, the upregulated genes are enriched for reproductive and developmental processes,
413 particularly those associated with gonad development, ovulation cycle regulation, and sexual
414 differentiation. Many terms are specific to female reproductive biology, indicating activation
415 of hormone-driven pathways such as estrogen or progesterone signaling. This transcriptional
416 shift from immune-related programs toward reproductive development could be driven by
417 physiological states, hormonal stimuli, or experimental conditions that favor reproductive
418 tissue function over immune activity.

419 Cell-type-specific Enrichment Analysis (CSEA) for differentially expressed genes was
420 conducted using WebCSEA, a web-based tool designed for Cell-type Specific Enrichment
421 Analysis of Genes. By leveraging WebCSEA, we found cell-type-specific gene expression
422 profiles for macrophages and dendritic cells (Supplementary Figure S3).

Figure 2. Differentially Expressed Genes Analysis: A) The volcano plot illustrates the most
significantly expressed immune system genes between low and high altitude residents. Genes are
plotted based on their expression levels and statistical significance (P ≤ 0.05, FC ≥ 2), highlighting
three down-regulated *FN1, CD36, and FOS*. B) Hierarchical clustering of differentially expressed
genes (DEGs) and samples is shown, with gene expression values normalized by Z-score. Functional
enrichment analysis of Gene Ontology was performed using ClusterProfile's enrichGO function,
focusing on biological processes for the entire set of DEGs (C, F), down-regulated genes (D, G), and
up-regulated genes (E, H). The x-axis represents the number of genes associated with each biological
process, and the false discovery rate (adjusted p-values) is indicated by a red-blue gradient, showing
statistical significance across different methods. P: False Discovery Rate (adjusted p-values); FC:

434   Fold change.

DISCUSSION

Our study is the first to investigate the gene expression of indigenous populations from Peru
(Figure 1) concerning their immune response, considering the altitude of their residence,
which generally exposes them to low oxygen concentration, cold, dryness, ultraviolet
radiation, and a differentiated immune response. In this sense, Quach et al. recently reported
that the effect of immunity may be related to ancestry, which an individual or population
belongs to; specifically, Africans tend to have a stronger immune response than Europeans
(14).

This is the first study to evaluate differential immune response in a high-altitude population
with high native ancestry, where we have identified DEGs. The downregulation of *FN1,*
CD36, and FOS, involved in processes related to extracellular matrix, cell adhesion, and
immune response, could improve our understanding of the modulation of the innate and
adaptive immune systems. Our results are congruent with what has been observed in previous
research conducted on Andean populations, which have identified genes involved in the
immune system, such as *DUOX2*, *CD40*, *TNIP1*, and *CHIA* (7,11).

A recent study published by Li *et al.* presents a meta-analysis of immune genes that
investigated how they varied in expression when exposed to altitude conditions, concerning
studies of cells exposed to normoxia or hypoxia. In the bioinformatics analysis, the cells
came from European descent subjects exposed to altitude but not native residents of a high-
altitude area (23). Li *et al.* demonstrated that the genes *ERH*, *VBP1*, *PSMA4*, *TOMM5*, and
POLR2K were downregulated in association with hypoxia; however, our results show that
these genes are not statistically differentiated. Additionally, these authors show that altitude
promotes the increase of some immune cells (T cells CD4 memory resting, T cells gamma
delta, Dendritic cells activated, Mast cells resting) at the expense of others relevant to the
innate or adaptive immune system (*e.g.,* T cells CD8, NK cells resting, Neutrophils).

In contrast, Manella *et al.* discovered that genes associated with immune response can exhibit
differential expression patterns based on both the duration of acute exposure to high altitude
and the specific altitude level (e.g., 3,800 meters vs. 5,100 meters). Additionally, the timing

of the measurements during the day also plays a role in these gene expression variations (24).

The investigation revealed that time-associated gene alterations were linked to innate

immunity, specifically about the activation of neutrophils. Additionally, distinct

transcriptional patterns were identified concerning the factors *RELA* and *IRF8*, recognized as

mediators of inflammation and immune responses. Furthermore, the study conducted by

Manella et al. unveiled an elevation in M2 macrophages and monocytes, influenced by both

altitude and time. Conversely, a decrease contingent on altitude was observed in NK cells and

regulatory T cells. Moreover, the researchers demonstrated the temporal variation in the

expression of *NR1D1*, with its overall expression being notably reduced in response to

altitude (25).

The Manella team's investigation is particularly intriguing due to its unique focus on

individuals of European descent in one of the world's most densely inhabited cities, located in

Peru. However, it focused on European descent men subjected to a relatively short-term

exposure, spanning several weeks, which cannot be directly compared to the chronic

exposure experienced by native inhabitants at such altitudes. Moreover, our study population

in Huánuco regularly experiences altitude fluctuations due to the necessity for individuals

residing in the high Andean regions to descend to the city of Huánuco for various activities,

such as shopping, making payments, receiving government assistance, and engaging in

recreational pursuits, among others. As a result, a prospective study conducted within the

Huánuco population, using a sample resembling ours, could investigate how genes are

regulated, either up-regulated or down-regulated, with the duration of absence from high-

altitude environments followed by return or conversely, the effects of prolonged high-altitude

exposure and subsequent return to lower altitudes.

Thus, we can assert that our study has demonstrated a distinct genetic expression signature in

high-altitude residents with a native Andean genetic background, as compared to their

counterparts living at lower altitudes. Furthermore, we have identified several genes

exhibiting both up-regulation and down-regulation, with a distribution observed within genes

related to the innate and adaptive immune systems.

It's worth emphasizing that most of our study participants were female, and we did not

implement age-matching procedures during the sample selection process. This absence of age

matching could potentially introduce selection bias, as gene expression can vary based on

both the participant's sex and age. To mitigate potential sex-related bias, we incorporated sex as a covariate variable in our differential expression analysis. However, we acknowledge that our study was limited in its ability to consider all environmental conditions and travel habits, factors that may have exerted an influence on our research findings.

A significant limitation of this study on the genomics and transcriptomics of lowlander and highlander populations is the small sample size. With few samples, the ability to generalize the results to the entire population is limited. Additionally, the small sample size increases susceptibility to statistical biases and reduces the statistical power to detect subtle differences or significant biological effects. However, our study stands as the pioneering effort demonstrating how immune response gene expression varies in individuals chronically exposed to high altitude (natives). Further investigations could validate and expand upon our findings through genomics, including transcription factors, transcriptomic, proteomic, or even epigenetic analyses.

CONCLUSIONS

The deregulated genes *FN1, CD36,* and *FOS* in native inhabitants with a prominent Andean genetic background highlight the active involvement of both the innate and adaptive immune systems. Remarkably, our study suggests that the immune response, even if induced by the same stimulus, varies with altitude. A comprehensive multi-omic approach is indispensable to fully understand the impact of altitude on immune system gene expression and genes related to high-altitude adaptation.

REFERENCES

1. Tremblay JC, Ainslie PN. Global and country-level estimates of human population at high altitude. Proc Natl Acad Sci USA. 2021 May 4;118(18):e2102463118.

2. Moore LG. Measuring high-altitude adaptation. Journal of Applied Physiology. 2017 Nov 1;123(5):1371–85.

3. Bigham A, Bauchet M, Pinto D, Mao X, Akey JM, Mei R, et al. Identifying Signatures of Natural Selection in Tibetan and Andean Populations Using Dense Genome Scan Data. Begun DJ, editor. PLoS Genet. 2010 Sept 9;6(9):e1001116.

4. Bigham AW. Genetics of human origin and evolution: high-altitude adaptations. Current Opinion in Genetics & Development. 2016 Dec;41:8–13.

5. Bigham AW, Mao X, Mei R, Brutsaert T, Wilson MJ, Julian CG, et al. Identifying positive selection candidate loci for high-altitude adaptation in Andean populations. Hum Genomics. 2009 Dec;4(2):79.

6. Crawford JE, Amaru R, Song J, Julian CG, Racimo F, Cheng JY, et al. Natural Selection on Genes Related to Cardiovascular Health in High-Altitude Adapted Andeans. The American Journal of Human Genetics. 2017 Nov;101(5):752–67.

7. Borda V, Alvim I, Mendes M, Silva-Carvalho C, Soares-Souza GB, Leal TP, et al. The genetic structure and adaptation of Andean highlanders and Amazonians are influenced by the interplay between geography and culture. Proceedings of the National Academy of Sciences. 2020;117(51):32557–65.

8. Taylor CT, Colgan SP. Regulation of immunity and inflammation by hypoxia in immunological niches. Nat Rev Immunol. 2017 Dec;17(12):774–85.

9. McGettrick AF, O'Neill LAJ. The Role of HIF in Immunity and Inflammation. Cell Metabolism. 2020 Oct;32(4):524–36.

10. Cramer T, Yamanishi Y, Clausen BE, Förster I, Pawlinski R, Mackman N, et al. HIF-1α Is Essential for Myeloid Cell-Mediated Inflammation. Cell. 2003 Mar;112(5):645–57.

11. Caro-Consuegra R, Nieves-Colón MA, Rawls E, Rubin-de-Celis V,

Lizárraga B, Vidaurre T, et al. Uncovering Signals of Positive Selection in Peruvian Populations from Three Ecological Regions. Mulligan C, editor. Molecular Biology and Evolution. 2022 Aug 3;39(8):msac158.

12.   Jacovas VC, Couto-Silva CM, Nunes K, Lemes RB, De Oliveira MZ, Salzano FM, et al. Selection scan reveals three new loci related to high altitude adaptation in Native Andeans. Sci Rep. 2018 Aug 24;8(1):12733.

13.   Joseph SK, Migliore NR, Olivieri A, Torroni A, Owings AC, DeGiorgio M, et al. Genomic evidence for adaptation to tuberculosis in the Andes before European contact. iScience. 2023 Feb;26(2):106034.

14.   Quach H, Rotival M, Pothlichet J, Loh YHE, Dannemann M, Zidane N, et al. Genetic Adaptation and Neandertal Admixture Shaped the Immune System of Human Populations. Cell. 2016 Oct;167(3):643-656.e17.

15.   Alexander DH, Novembre J, Lange K. Fast model-based estimation of ancestry in unrelated individuals. Genome Res. 2009 Sept;19(9):1655–64.

16.   Thornton T, Tang H, Hoffmann TJ, Ochs-Balcom HM, Caan BJ, Risch N. Estimating Kinship in Admixed Populations. The American Journal of Human Genetics. 2012 July;91(1):122–38.

17.   Leal TP, Furlan VC, Gouveia MH, Duarte JMS, Fonseca PA, Tou R, et al. NAToRA, a relatedness-pruning method to minimize the loss of dataset size in genetic and omics analyses. Computational and structural biotechnology journal. 2022;20:1821–8.

18.   Patterson N, Price AL, Reich D. Population Structure and Eigenanalysis. PLoS Genet. 2006;2(12):e190.

19.   Price AL, Patterson NJ, Plenge RM, Weinblatt ME, Shadick NA, Reich D. Principal components analysis corrects for stratification in genome-wide association studies. Nat Genet. 2006 Aug;38(8):904–9.

20.   Chen Y, Chen L, Lun ATL, Baldoni PL, Smyth GK. edgeR 4.0: powerful differential analysis of sequencing data with expanded functionality and improved support for small counts and larger datasets. 2024;

21.   Hoffman GE, Schadt EE. variancePartition: interpreting drivers of variation in complex gene expression studies. BMC Bioinformatics. 2016 Nov 25;17(1):483.

22.   Wu T, Hu E, Xu S, Chen M, Guo P, Dai Z, et al. clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. The Innovation. 2021 Aug;2(3):100141.

23.   Li Q, Xu Z, Fang F, Shen Y, Lei H, Shen X. Identification of key pathways, genes and immune cell infiltration in hypoxia of high-altitude acclimatization via meta-analysis and integrated bioinformatics analysis. Front Genet. 2023 Mar 24;14:1055372.

24.   Manella G, Ezagouri S, Champigneulle B, Gaucher J, Mendelson M, Lemarie E, et al. The human blood transcriptome exhibits time-of-day-dependent response to hypoxia: Lessons from the highest city in the world. Cell Reports. 2022 Aug;40(7):111213.

25.   Scheiermann C, Kunisaki Y, Frenette PS. Circadian control of the immune system. Nat Rev Immunol. 2013 Mar;13(3):190–8.

SUPPORTING INFORMATION

Figure S1. Huanuco population structure based on the subset of samples with intersecting genomic and transcriptomic data.

Figure S2. Variance Partition Analysis.

Figure S3. Cell-type-specific Enrichment Analysis (CSEA) for differentially expressed genes was obtained using WebCSEA (Web-based Cell-type Specific Enrichment Analysis of Genes).

Table S1 - Population data
Table S2 - Population ancestry
Table S3 - Huanuco ancestry data
Table S4 - Distribution of samples per group and altitude
Table S5 - All DEGs
Table S6 - Functional enrichment analysis for down-regulated genes
Table S7 - Functional enrichment analysis for up-regulated genes
Table S8 - Functional enrichment for all deregulated genes
File: Analysis scripts
File: Quality Control