Prediction of Aggregation Prone Regions in Proteins Using Deep Neural Networks and Their Suppression by Computational Design

Vojtech Cima[1,#], Antonin Kunka[2,3,#,§], Ekaterina Grakova[1], Joan Planas-Iglesias[2,3], Martin Havlasek[2,3], Madhumalar Subramanian[2], Michal Beloch[1], Martin Marek[2,3], Katerina Slaninova[1], Jiri Damborsky[2,3], Zbynek Prokop[2,3,*], David Bednar[2,3,*], Jan Martinovic[1,*]

[1] IT4Innovations, VSB – Technical University of Ostrava, 17. Listopadu 2172/15, 70800, Ostrava-Poruba, Czech Republic

[2] Loschmidt Laboratories, Department of Experimental Biology and RECETOX, Faculty of Science, Masaryk University, Kotlarska 2, Brno, Czech Republic

[3] International Clinical Research Centre, St. Anne's University Hospital, Pekarska 53, Brno, Czech Republic

§The current affiliation of the Author is: Protein Biophysics Group, Department of Biotechnology and Biomedicine, Technical University of Denmark, Søltofts Plads, Building 227, 2800, Kgs. Lyngby, Denmark

#joined first authors; * Corresponding authors: Zbynek Prokop (zbynek@chemi.muni.cz), David Bednar (222755@mail.muni.cz), Jan Martinovic (jan.martinovic@vsb.cz)

Abstract

Protein aggregation is a hallmark of multiple neurodegenerative diseases and a great hindrance in recombinant protein production, handling, and storage. Identification of aggregation prone residues or regions (APRs) in proteins and their suppression by mutations is a powerful and straightforward strategy for improving protein solubility and yield, which significantly increases their application potential. Towards this, we developed a deep neural network based predictor that generates residue level aggregation profile for one or several input protein sequences. The model was trained on a set of hexapeptides with experimentally characterised aggregation propensities and validated on two independent sets of data including hexapeptides and full-length proteins with annotated APRs. In both cases, the model matched, or outperformed the state-of-the-art algorithms. Its performance was further verified using a set of 34 hexapeptides identified in model haloalkane dehalogenase LinB and seven proteins from AmyPro database. The experimental data from Thioflavin T fluorescence and transmission electron microscopy matched the predictions in 79% of the cases, and revealed inaccuracies in the database annotations. Finally, the utility of the algorithm was demonstrated by identifying APRs in a model enzyme (LinB) and designing aggregation-suppressing mutations in the exposed regions. The designed variants showed reduced aggregation propensity, increased solubility and improved yield, with up to a 100% enhancement compared to the wild type for the best one.

Keywords
Amyloid, Aggregation, Deep Neural Network, Computational design, APR

1. Introduction

Recombinant proteins are incredibly versatile with applications spanning from industrial biocatalysts, waste processing, food industry, to highly efficient therapeutics or vaccines. The extraordinary advancements in computational structure prediction[1] greatly accelerated their

transformation towards one of the most promising technologies for a sustainable future and improved quality of life. Although protein structures can now be predicted with high accuracy almost instantaneously[2–4], the underlying computational algorithms are designed assuming a single folded state corresponding to the global minimum of the free energy landscape (FEL)[5]. In reality, folding of protein chain into its functional - native state, driven by formation of intramolecular interactions and burial of hydrophobic residues within protein core, competes with formation of non-native, intermolecular interactions leading to higher-order misfolded assemblies, i.e. protein aggregates. Misfolding pathways are inherent to any protein sequence and depend on both extrinsic (e.g., temperature, pH, ionic strength), and intrinsic (e.g., amino acid sequence, concentration) conditions[6]. Misfolding and aggregation can be triggered *in vivo* by numerous factors, such as oxidative stress, proteasome malfunctioning, posttranslational modifications, or mutations, and is a hallmark of severe pathologies such as Alzheimer's disease, Parkinson's disease, diabetes, prion disease, and many others[7]. Although the underlying mechanisms of these diseases are not yet fully clear, the cellular toxicity of the aggregates by both gain-of-function (generation of toxic species), and loss-of-function (i.e., reduction of proteins native activity due to its aggregation) have been described[8]. Moreover, aggregation and deposition of misfolded, insoluble proteins into inclusion bodies is one of the major hindrances in production of recombinant proteins in the bioindustry. Identification of sequence stretches driving the aggregation, i.e., aggregation prone regions (APRs), is therefore of utmost importance.

Aggregation is inversely correlated with solubility, defined as the concentration of soluble protein in equilibrium with the solid aggregates. These can be classified according to their morphology to amorphous assemblies and amyloid fibrils or crystals. Amorphous aggregates lack well defined high-order structure and their formation is driven by non-specific, mostly hydrophobic interactions[9,10]. Although analytical frameworks that quantitatively describe kinetics of amorphous aggregation have been derived and successfully applied[11–20], detailed characterization of amorphous aggregate assembly attracts little attention. Instead, amorphous aggregation is perceived as an omnipresent problem during protein production and processing[10]. In contrast, amyloids gained significant attention due to their link to over a number of diseases[7], and analysis of their formation has been extensively studied[21,22]. Amyloids are ordered two-dimensional assemblies formed by repeating units of soluble protein precursors (monomers) that are stabilised by an intermolecular network of hydrogen bonds within the cross-β-sheet architecture[23]. The interface within the amyloid subunit is formed by interdigitated complementary amino acid side-chains stabilised by van der Waals interactions, hydrophobic forces, and hydrogen bonds[24,25]. This structural and energetic complementarity in the eight possible arrangements (classes) of these so-called "steric zippers"[26] governs the available amyloid sequence space[27]. Emergence of different fibril polymorphs, i.e., fibril structures with distinct morphologies formed by the same protein sequence, in the increasing number of experimentally determined fibril structures proves that the amyloid landscape is highly frustrated[28,29]. Despite the differences between chain arrangements within their cores, the polymorphs often share a common structural kernel which is believed to be an important driver of amyloid formation and essential for its stability[30–32]. These regions (APRs) are therefore perfect targets for designing mutations that decrease aggregation propensity and, consequently, increase protein solubility.

Existing algorithms for identification of APRs can be broadly categorised into (i) linear and (ii) structural predictors based on their underlying principle and input data. The first class derive the aggregation propensity from the protein sequence using the physio-chemical properties of the amino acids (e.g., hydrophobicity, net charge, patterning,), experimentally derived aggregation scales[33,34], β-sheet propensity[35–39], or their combination[40–44]. The second class of algorithms utilise protein structures to identify aggregation prone patches in globular proteins that are formed by sequentially distant residues that are impossible to identify using the linear predictors[45–48]. Emergence of databases compiling protein and peptide sequences with experimentally validated APRs or aggregation propensities[49–51] boosted the development of machine learning-based predictors[27,52–59]. Their main advantage is that they are fast which (in some cases) enables their application on a proteome-wide scale[52]. However, most of them are only validated on existing datasets and their accuracy and usefulness in real-life applications are yet to be determined. The long-term evolution of aggregation predictors with distinct underlying principles greatly improved our understanding of protein aggregation on molecular level, and aided in solubilization and design of many important targets[60]. However, their performance remains suboptimal and there is still a great potential in improving their accuracy so that they can be efficiently incorporated into protein design pipelines to minimise the experimental screening effort.

Towards this, we have developed a novel protein aggregation predictor based on deep neural networks (DNN) called AggreProt. AggreProt consists of (i) sequential, and (ii) static models that were both trained on the hexapeptides from WaltzDB 2.0[51] using atomic related description of each amino acid (36 features[52]) of the hexapeptides or 18 features found in the database, respectively. The accuracy of AggreProt was higher or comparable to the state-of-the-art predictors Tango, Waltz, Aggrescan, PASTA 2.0, GAP, AnuPP, and FishAmyloid using both hexapeptide- and protein-based validation datasets. Furthermore, we carried out extensive experimental validation of AggreProt to demonstrate its application potential. In contrast to other predictors, AggreProt correctly identified APRs in one of our workhouse proteins - haloalkane dehalogenase (HLD) LinB (UniProt ID:D4Z2G1). This enabled us to fulfil our long-term effort and one of the motivations behind this study, i.e., to design mutations that improve the solubility and double the enzyme yield. We went beyond and experimentally validated AggreProt using hexapeptides derived from the LinB and proteins in the AmyPro database[50]. Based on our close investigation of the AmyPro entries using AggreProt, we have selected hexapeptides that were potentially missanotated in the database, and subsequently confirmed this by the experiment. This not only demonstrates the great accuracy and usefulness of the AggreProt, but implies that the benchmarking of aggregation predictors using AmyPro, CPAD 2.0 or similar datasets may result in biassed statistical evaluation of the tools.

2. Materials and methods
2.1. Design and development of the aggregation prediction models
2.1.1. Training and validation datasets
Training of the AggreProt DNN models was carried out using hexapeptides from the WaltzDB2.0 database[51]. WaltzDB contains 1416 sequences of hexapeptides with experimentally characterised and annotated amyloid-forming properties (515 amyloid, 901 non-amyloid). Additionally, each peptide entry contains features from three different in-silico predictors (TANGO[42], PASTA 2.0[36], and Waltz[40]) together with precalculated hydrophobicity and secondary structure propensities, and local energetic

assessment from FoldX[61]. The hexapeptide dataset was randomly split into training (90%, WaltzDB-90) and validation (10%, WaltzDB-10) subsets. The training subset was used in a train-validation fashion (80:20 leave-one-out) during the hyperparameter optimization of the models, whilst the later subset of peptides was used for model evaluation.

The second validation dataset was selected from proteins in the AmyPro database[50]. AmyPro contains 162 protein sequences (AmyPro162) with manually annotated amyloidogenic regions collected from literature. To ensure there is no overlap between training and validation, proteins that contain sequences of the hexapeptides from WaltzDB2.0 were filtered out. In total, 37 proteins (Amy37) with annotated APRs remained and were used for validation. We further split the Amy37 into two subsets of proteins with APRs longer or shorter than 50 amino acids, and used the latter (n = 27 proteins, AmyPro27) as a high-confidence validation dataset. The number of sequences with short APRs in the complete dataset is 140 (AmyPro140). The main rationale behind such a split is that the annotations are often made based on indirect experimental evidence (as discussed in the results section) and true APRs are generally short segments.

2.1.2. Metrics

The evaluation of DNN performance employed standard metrics derived from the confusion matrix, using both WaltzDB-90 in the training procedure and WaltzDB-10 in the testing (see below). True Positives (TP) are instances where ground truth positives (amyloid prone) are correctly predicted as positives, while False Positives (FP) refer to ground truth negatives (non-amyloid prone) erroneously predicted as positives. Conversely, False Negatives (FN) represent instances where ground truth positives are predicted as negatives, and True Negatives (TN) are instances where ground truth negatives are correctly predicted as negatives. Based on these definitions, Precision (Prec) was calculated as TP/(TP+FP), Recall (Rec) was calculated as TP/(TP+FN), and Specificity (Spec) was calculated as TN / (FP + TN). The performance assessment of the networks involved determining the AuROCC and Precision/Recall Curves (PRC). Youden's J statistic[62] was exploited on Receiver Operating Characteristic Curves (ROCCs) to identify the optimal threshold for prediction.

The evaluation of the final models on whole protein sequence data (AmyPro) was done using the aforementioned metric and the Segment OVerlap score (SOV). SOV was defined by Rost and co-workers[63], and we used the metric to evaluate the correctness of the predictions in a whole sequence context as it was used in used ANuPP(Prabakaran 2021). The SOV score can be calculated both for APR and non-APR segments.

2.1.3. Exploring network attributes and hyperparameters. Architecture of the predicted models

AggreProt DNN models assess the aggregation tendency of an input hexapeptide by analysing either the atomic features associated with each amino acid in the hexapeptide (linear model), following the approach employed in ANuPP[52], or the structural related features obtained from WaltzDB[51]. Consequently, the initial input layer dimension (number of neurons) aligns with the count of initial features, which for the linear predictor is 36 per reside in the input hexapeptide, and for the static one is 18. Throughout the training process, we fine-tuned the architecture and internal parameters of the networks using a 5-fold cross-validation with WaltzDB-90. We optimised the following parameters: i) number of bisectional layers (one or two), ii) number of dense layers (one, two or three), iii) dropout values (from 0 to 0.8, in 0.2-span steps), iv) number of neurons per layer (from 8 to 512, in powers of two-span steps), v) batch size (16 or 32), and vi) learning rate (from 0.0001 to 0.1, in powers of ten-span steps). The models were trained exploiting the TensorFlow Framework[64] in our custom codebase.

A total of 840 and 1680 hyperparameter combinations were examined for the linear and static model architectures respectively. Each of such combinations generated a unique network architecture that was trained in a 5-fold cross-validation manner. Thus, we generated 5 network instances for each architecture assessed. The performance of each architecture was assessed in terms of Area under the Receiver Operating Characteristic Curve (AuROCC). To avoid overfitting, the training was stopped at its optimum, when the training and validation ROC curves diverged (Supplementary Figure 1). As a final model we selected an average ensemble of the different network instances of the top performing architecture for each of the model types (linear and static).

We have designed two different neural network architectures (model types) reflecting the hexapeptide properties (input features dimensions) used for their parameterization, i.e., (i) sequential and (ii) static. In the sequential model, each hexapeptide is parameterized according to the atomic composition as previously described[52]. In the static model, each hexapeptide is parameterized by 18 features from WaltzDB 2.0 database. These include scores from Waltz, TANGO, PASTA2.0 (parallel and antiparallel beta-sheet propensity), hydrophobicity (filter on the percentage of hydrophobic residues, including A, C, F, I, L, M, P, V, W, and Y), Chou-Fasman helix and strand propensities, and 11 FoldX force-field energy parameters calculated using Cordax[27].

Our models were trained on hexapeptide data. In order to make them amenable to process whole protein sequences, the input protein sequence of n amino-acids is divided into n hexapeptides using a sliding window approach with a step of one residue. The aggregation propensity prediction is generated for each hexapeptide. And all the information on a single amino-acid (contributed by up to 6 different hexapeptides) was aggregated using three different metrics: minimum, average and maximum.

2.1.4. Model validation and benchmarking

The final models were evaluated using two different testing sets. First, the remaining 10% of WatzDB not used during the training procedure (WaltzDB10). Here, we calculated AuROCC as we did during training and compared to the training value.

Second, we validated on complete sequence proteins from AmyPro. To identify APRs on input protein sequences, these were fragmented into overlapping hexapeptides using a six-amino-acid-windows shifted one amino acid at a time. Each hexapeptide was evaluated by each of our two models (linear and static) and the results were averaged per residue mirroring the sliding window procedure used before. The performance of each network was assessed in terms of AuROCC, AuPR, and SOV.

2.2. Predictor testing and computational design of mutations suppressing the aggregation
2.2.1. Detection of aggregation prone regions in model protein and computational design of aggregation-suppressing mutations.

Sphingobium japonicum haloalkane dehalogenase LinB (UniProt ID: D4Z2G1) was analysed using the sequential model of AggreProt. The surface accessible area (SASA) was calculated using the DSSP algorithm[65] on the publicly available LinB structure (PDB IDs: 1mj5). Residue accessibility values were normalized by the amino acid total volume to obtain SASA value per residue. Altogether, seven sequence stretches with highest AggreProt values were identified as potential APRs. The most solvent exposed residue with the highest AggreProt score in the middle of each APR, and two exposed residues that defined the APR borders were considered for mutagenesis. We iteratively mutated each selected position (APRs centres) or pairs of positions (APRs borders) to all possible amino acids and calculated the resulting AggreProt scores. Next, we evaluated the effects of all possible mutations on the structural integrity of the protein using HotSpot Wizard (HS)[66] and Rossetta ddg_monomer[67]. The mutations that had low AggreProt score, high mutability according to HS, reasonable occurrence (i.e., not rare) in homologous sequences based on multiple sequence alignment, and did not significantly compromise the conformational stability of the protein, were selected for experimental characterization.

2.2.3. Identification of hexapeptides for experimental validation of their aggregation propensity.

Based on the manual inspection of our AmyPro37 validation dataset, we selected seven relatively short proteins (lysozyme c, apomyoglobin, serum amyloid A, envelope small membrane protein, cell division topological specificity factor, gallectin-7, and gamma-crystallin D) with clear deviations between our predictions and ground truth based on AmyPro. Within these proteins, we identified 17 hexapeptides which might be incorrectly labelled in the AmyPro as amyloids (i.e., false positives - FP, n = 11) or non-amyloids (i.e., false negatives - FN, n = 6). Our selection criteria included inconsistency between labels and primary literature, and low or high AggreProt scores for FP and FN, respectively. As a second dataset, we chose 7 hexapeptides corresponding to the putative APRs found in LinB and their 10 counterparts containing aggregation-suppressing mutations described above. Altogether, we selected 34 hexapeptides for experimental validation of their aggregation propensity (**Supplementary Table 1**).

2.3. Experimental validation

2.3.1. Material and chemicals

The genes encoding LinB variants with C-terminal His-tag were cloned in the pET21b plasmids and expressed in *E. coli* BL21(DE3) (New England Biolabs). Target mutations were introduced using megaprimer PCR mutagenesis using pairs of mutagenic and non-mutagenic primers (**Table 1**). All chemicals used in this study, classified as the quality of *pro analysis,* were purchased from Sigma unless specified otherwise. The hexapeptides selected for the experimental validation were synthesised by ABI Scientific Inc. (USA) as the trifluoroacetate salts with acetylated and aminated N- and C-termini, respectively. Their purity was 98 % according to the HPLC analysis carried out by the manufacturer.

2.3.2. Site-directed mutagenesis

The mutagenic primers used for the introduction of the designed mutation were ordered from Merck (Merck kGaA, Germany). The results of the megaprimer mutagenesis, consisting of two-round PCR, were verified using sequencing (Eurofins Genomics, Germany).

Table 1: List of mutagenic primers. LinB175 (denoted with an asterisk) was prepared using the DNA sequence of LinB165 as a template. All other mutants were prepared using wild-type as a template.

LinB variant	Forward primer	Reverse primer
161	5'-ctggtgttgcaggacaatgtctttaaggagcaggtcttgccaggcctgatc-3'	5'-gctagttattgctcagcgg-3'
162	5'-ttgcaggacaatgtctttgttgagcgcgtcttgccaggcctgatcctgcgt-3'	5'-gctagttattgctcagcgg-3'
163	5'-agcaggcgaagaactggtcgcgcaggacaatgtctttgttgagcaggtcttgccacgcctgatcctgcgtccgctg-3'	5'-gctagttattgctcagcgg-3'
164	5'-taatacgactcactataggg-3'	5'-ccagcctgcgtagtcacgcgcaatttcgaccacatcagcaggcgtgccggc-3'
165	5'-taatacgactcactataggg-3'	5'-cagccagcctgcgtagtcttccgcaatcgcgaccacatccggaggcgtgccggcaatcgg-3'
166	5'-taatacgactcactataggg-3'	5'-gcctgcgtagtcacgcgcaatcgcatgcacatcagcaggcgtgccggcaat-3'
167	5'-taatacgactcactataggg-3'	5'-cggcgaatcttcctggataaagtggccacccgccacggtaatctcggtttg-3'

168	5'-taatacgactcactataggg-3'	5'-cggcgaatcttcctggataaagtggcgacccgccacggtaatctcggtttg-3'
169	5'-taatacgactcactataggg-3'	5'-atcttcctggataaagtgagcacccggcacggtaatctcggtttggttcgg-3'
170	5'-taatacgactcactataggg-3'	5'-ctggataaagtgagcacccgccacgcgaatctcggtttggttcggccacgt-3'
171	5'-taatacgactcactataggg-3'	5'-acgcagacgacgaacaaatgctgcatcggccgcaccaatctcgtccggcga-3'
172	5'-taatacgactcactataggg-3'	5'-acgacgaacaaatgctgcgatggcttcaccaatctcgtccggcgaatcttc-3'
173	5'-taatacgactcactataggg-3'	5'-tgcaggacgcagacgacgaacaaattctgcgatggcttcaccaatctcgtccggcgaatcttc-3'
174	5'-taatacgactcactataggg-3'	5'-gcaggcctgggtcgcctgattgcgccggatctgatcggtatgggtgacagc-3'
175*	5'-taatacgactcactataggg-3'	5'-tgcaggacgcagacgacgaacaaattctgcgatggcttcaccaatctcgtccggcgaatcttc-3'

2.3.3. Protein expression, purification, and quality control

LinB variants were expressed in *E. coli* BL21 (DE3) transformed by the mutagenic plasmids (pET21b) encoding respective LinB mutants. The protein expression was induced by the addition of IPTG (0.4 mM final concentration) and carried out at 20°C for 16 hours . The cells were harvested by centrifugation (4,000 x g, 4 °C, 30 min) and frozen at -70 °C. After defrosting, the resuspended cells were disrupted by the addition of DNAse I and 3 rounds of sonication. The disrupted cells were centrifuged (21 000g, 4 °C, 60 min) and the proteins were purified from the cell lysates by metallo-affinity chromatography with HisTrap HP column (Cytiva, USA) charged with N^{i2+} ions using stepwise increase of imidazole in elution buffer. The pure monomeric fraction for the following aggregation experiments was isolated by the size-exclusion chromatography using the Superdex 200 pg 10/300 GL column (Cytiva, USA) with the 50mM Potassium phosphate buffer (pH 7.5) used as the mobile phase. Protein purity was verified by SDS-PAGE and the concentration determined by UV-absorbance measurement (DeNovix DS-11 Spectrophotometer, DeNo-vix Inc.,USA).

2.3.4. Temperature scanning experiments

The aggregation propensity of all tested LinB variants was screened using UNcle (Unchained labs, USA) in a 50 mM potassium phosphate buffer pH 7.5. The protein samples were continuously heated

from 20 to 95 °C at 1°C/min scan rate and the static light scattering (SLS) at 266 nm and fluorescence emission spectra were recorded. The aggregation propensity was estimated from the maximum amplitude of the SLS curve while the apparent melting temperature (T_m^{app}) was evaluated from the midpoint of the average emission wavelength (i.e., Barycentric mean - BCM) curve. Each sample was analysed in a duplicate at protein concentration of 0.275 mg/ml to ensure comparability between individual variants.

2.3.5. Aggregation kinetics

Selected LinB variants were selected based on the previous screening and their aggregation kinetics at 37 °C was monitored using SLS and Thioflavin T assay (ThT) in 50 mM Potassium phosphate buffer (pH 7.5). The scattering intensity at 266 nm was measured using UNcle (Unchained labs, USA) and the amount of aggregates was estimated from the curve amplitude. Each sample was analysed in a duplicate at two different protein concentrations of 0.5 and 1.0 mg/ml.

The ThT assay was measured in 384-well plates (Corning® 384-well black, clear flat bottom polystyrene microplate) sealed with aluminium tape (Corning® 96 Well Microplate Aluminum Sealing Tape in duplicates using Synergy H4 reader (BioTek, USA). The ThT fluorescence (30 µM) added to samples was excited at 440 nm and emission measured at 485 nm using bottom read over the course of 24 hours in 5-minute intervals at 37°C. Each variant was measured at two protein concentrations of 0.5 and 1.0 mg/ml in duplicates.

2.3.6. Flow-induced dispersion analysis

The amount of soluble protein in the samples after aggregation was analysed for the two least aggregation-prone LinB variants and wild-type using flow-induced dispersion analysis (FIDA) using FIDA 1 (Fidabio, Denmark) equipped with the protein intrinsic fluorescence detection (280 nm excitation).. The three proteins (1 mg/ml) were incubated at 37 °C for 24 hours in low binding tubes. Aliquots withdrawn from the aggregation reaction at the start of incubation (0 h), and after 1 and 24 hours were centrifuged (18000g, 4 °C, 20 min) and the supernatant analysed using FIDA with the following method:
Step 1: NaOH wash (60 s, 3500 mbar)
Step 2: H_2O wash (60 sec, 3500 mbar)
Step 3: Buffer wash (40 sec, 1500 mbar)
Step 4: Sample application (20 s, 75 mbar)
Step 5: Sample analysis (buffer, 75 s, 1500 mbar)
The amount of soluble protein at each time-point was determined from the integral of the resulting Gaussian peak and expressed relative to the amount at the beginning of the reaction.

2.3.6. Dehalogenase activity measurement

Effects of the introduced mutations to the protein functionality were verified by determination of their specific activity towards 1,2-dibromoethane using the colorimetric assay according to Iwasaki[68]. The dehalogenation reactions were carried out at 37 °C in 25-ml ReactiFlasks closed by Mininert Valves. The reaction mixture was composed of 10 mL of 100mM glycine buffer (pH 8.6) and 10 µl of the substrate (1,2-dibromoethane, purity ≥ 98%). The reaction was initiated by the addition of the enzyme (final concentration ~ 2 µg/ml). The progress of the reaction was monitored over 30 minutes by withdrawing 1 ml aliquots from the reaction mixture in 4-minute intervals. At each time point, the reaction was terminated by the addition of 0.1 ml of 35% nitric acid to the mixture. The bromides formed during the reaction were detected spectrophotometrically upon the reaction with mercuric thiocyanate and ferric ammonium sulphate using absorbance at 460 nm (Eon plate spectrophotometer, BioTek, USA). The absorbance read-out was recalculated to the concentration of the product using the calibration curve with known concentration of bromides. The activity was determined from the slope of the product concentration over time.

2.3.6. Circular dichroism spectroscopy

The secondary structure of the two least aggregation-prone LinB variants and wild-type and their thermal stability was analysed using circular dichroism (CD) spectroscopy. The CD spectra of 0.175

mg/ml proteins in a 50 mM potassium phosphate buffer were measured from 185 to 260 nm (0.25 s integration time, 1 nm bandwidth) at 20 °C using Chirascan spectropolarimeter (Applied Photophysics, UK). Thermal stability was probed by monitoring the ellipticity changes at 227 nm while the temperature was gradually increased at 1°C/min scan rate from 20 to 75 °C. The apparent melting temperatures were assessed from the midpoints of the denaturation curves.

2.3.7. Thioflavin T assay and transmission electron microscopy of the hexapeptide aggregates

The peptide lyophilisates were first dissolved in a 50 mM sodium phosphate buffer (pH 7.4) to a final concentration of 5 mM. The insoluble ones (based on turbidity and physical appearance of the buffer stocks) were additionally dissolved in either DMSO or HFIP (for peptides containing cysteine to avoid their oxidation) to a final concentration of ~100 mM. For the aggregation assays, all peptides were diluted to the final working concentration of 1 mM using 50 mM sodium phosphate buffer pH 7.4 containing 0.05% sodium azide to prevent bacterial contamination. Samples were incubated at 37 °C for 4 weeks and the progress of aggregation was monitored using ThT assay as described above. At the end of the incubation, full ThT emission spectrum was recorded (ex. 440, em. 460-560 nm) for each sample to determine whether the hexapeptide aggregate is positive for ThT. Finally, 4 µl of the 1 mM peptide solution was spotted onto a carbon-coated 300 mesh copper grid and incubated for 2 minutes, followed by 2 minute negative staining using 2% uranyl acetate solution. Imaging was performed using Talon™ F200C microscope (Thermo Scientific™). For reliable comparison, three independent squares were imaged for all peptides and the obtained structures were characterised into five different categories: fibrils, short fibrils, amorphous aggregates, hydrogels/unspecific structures and also grids with no aggregation. Classification of the hexapeptide states to amyloids and non-amyloids for AggreProt benchmarking was done primarily based on the TEM images by the experimentalist without knowledge of the predicted aggregation scores of the hexapeptides to avoid biassed assessment.

3. Results and Discussion

3.1. Derivation of the input features for the training dataset and architecture of the deep neural networks

One of the major hindrances in the development of reliable protein aggregation predictors is the scarcity of reliable experimental data for the training. We have identified AmyPro[50] and WaltzDB2.0[51] as the most suitable data repositories that could serve our purpose. AmyPro contains 162 protein sequences with manually annotated amyloid forming sequence stretches (APRs) whereas WaltzDB2.0 involves 1416 hexapeptides with experimentally verified amyloid-forming propensities. Considering the unambiguous data labels in the later, and the fact that the hexapeptide is widely considered as the minimal structural unit capable of forming the steric zipper of the amyloid core[23], we have decided to utilise the set of hexapeptides from WaltzDB for the training.

In the next step, we have selected three classes of descriptors for the hexapeptides. For the first model class (termed "sequential"), input features for every amino-acid in each hexapeptide were generated using the amino-acid atomic composition (36 descriptors[52]). For the second class (termed "static"), we have used the parameters from the WaltzDB itself. These include aggregation propensity scores calculated using Waltz, TANGO, and PASTA software, Chou-Fasman alpha-helix and beta-sheet propensity[69], hydrophobicity scale (as described in WaltzDB[51]), and eleven energy parameters caluclated using FoldX[61] and Cordax[27].

The two models we present here were trained on a 5-fold cross-validation procedure (see Methods), diverged on the type of input features (see above), and both took a hexapeptide as input. From the training procedure, we chose the top performing architectures (summarized in **Figure 1**) and we built each model as an ensemble of the 5 variations of each top-ranking architecture produced by the 5-fold cross-validation procedure. The total number of parameters incorporated to the networks was 46113 and 2049 for the linear and static model, respectively.

Figure 1. Model architectures: **A** In the linear model, an input layer processes sequential inputs into a cascade of bidirectional long short-term memory units, followed by two dense (fully-connected) layers, each connected to their corresponding dropout layer, finishing in a single-unit dense output layer. **B** The static model is composed of an input layer that processes normalized static features (hexapeptide-level properties) into a dense (fully connected) layer, followed by a dropout layer and a single-unit dense output layer. All model architectures are supported with by dropout layers aiming to reduce potential model overfitting (**Supplementary Figure S1**).

3.2. Model validation, comparison, and benchmarking

We validated our models against two different datasets. First, the remaining 10% of WaltzDB (WlatzDB-10) that was not used during the training-validation procedure. Here, we expected that our networks will perform similarly than with the validation set and worse than with the training one (so that overfitting and data leakage were avoided). Indeed, our models achieved an AuROCC of 0.89 and 0.85 (linear and static models, respectively) when they achieved 0.88 and 0.89 on the validation, respectively. On the training set both models reached an AuROCC of 1 (**Supplementary Figure 1**).

Second, we enquired how or models performed on complete protein sequences. Here, we implemented a two-step sliding window procedure, to first decompose the input protein into hexapeptides and then map and aggregate each hexapeptide prediction (value) to every amino-acid in the protein sequence (see Methods). We used three different aggregation metrics (minimum, average, and maximum) to map the hexapeptide prediction onto individual amino-acids. On a dataset completely oblivious to any hexapeptide used during the training validation procedure (AmyPro37), the linear predictor achieved an AuROCC of 0.53 for all three aggregating metrics, and an AuPRC of 0.23. The static predictor reached an AuROCC ranging

between 0.49 (minimum) and 0.50 (average and maximum); all three metrics resulted in an AuPRC of 0.23 (**Supplementary Figure 2 A, B**).

We were struck with these big differences in between the validation on the whole-sequence data and the hexapeptide data. We explored the nature of the data stored in AmyPro and we realised that there were very long APRs annotated (more than 50 consecutive residues). It is highly unlikely that all residues in these stretches contribute equally to the aggregation propensity of the protein which is not reflected by the binary annotation of the database (i.e., amyloid, non-amyloid) and can thus introduce bias during validation. Removing proteins containing such long APRs from the previous dataset, we obtained AmyPro27 and we proceeded again to validate our models. On this benchmark better representing the biological nature of aggregation (and still oblivious to any hexapeptide used during training), the linear model reached AuROCCs of 0.59, 0.64 and 0.65 for each of minimum, average, and maximum aggregation metrics, respectively. The corresponding AuPRCs were 0.28, 0.32 and 0.33. The static predictor, achieved AuROCCs of 0.61, 0.66, and 0.65 for the three aggregation matrices, and their corresponding AuPRCs were 0.32, 0.35 and 0.34. (**Figure 2 A, B**) Thus, the predictor cannot learn about long APRs from hexapeptide data, but those long APRs are likely to be artifacts of the method of detection.

We also computed the SOV metric, which represents how well (in a range between 0 and 1) our predictions cover the APRs and non-APRs of the annotated data in AmyPro. The different aggregation metrics on the linear predictor ranged between 0.51 and 0.61 in APRs, and between 0.29 and 0.42 in non-APRs when validated on AmyPro27 (**Figure 2 C, D**). On the other hand, on AmyPro32 these values dropped to a range between 0.40 and 0.47 for APRs and a range between 0.23 and 0.34 for non-APRs, in the case of the linear predictor (**Supplementary Figure 2 C, D**). In general, we didn't observe big differences among the linear and static predictor. The second, under some aggregation metrics, performs better, but this improvement comes at a price of much larger computation time. On the rest of the manuscript (section 3.3 and below), we use predictions by the linear model with an average aggregation procedure (AggreProt_linear_mean), with an optimal threshold of 0.25 calculated by Youden's J[62] on AmyPro140 (the larger validation dataset available without long APRs). This same predictor has been implemented in AggreProt web server[70], where details of its validation are also shown.

Finally, we compared the results of our models to those of other sequence-based state-of-the-art methods, namely Waltz[40], Tango[42], ANuPP[52], and AGGRESCAN[33]. Except in the case of SOV for non-APRs, our models performed equal or better than the others (**Figure 2, Supplementary Figure 2**). The underperformance on the SOV metric for non-APRs, indicates that our networs estimate aggregation-prone regions that are not annotated as such. We wanted to understand whether this was an undesired feature of our predictors or, on the other hand, was a problem with the annotated data. In order to address this conundrum, we devised a experimental procedure that is detailed in section 3.4.

Figure 2. **Performance of the trained networks using different aggregating metrics and other state-of the art predictors on AmyPro27.** Different evaluation metrics are shown for the performance of the linear (blue hues) and the static (red hues) predictors on AmyPro27, along with other state-of-the-art sequence-based methods (Waltz, pink; Tango, green; Aggrescan, orange; ANuPP, grey). **A**: Area under the Receiver Operating Characteristic Curve (AuROCC). **B**: Area under the Precision-Recall Curve (AuPRC). **C**: Segment OVerlap score for Aggregation Prone Regions (SOV_APR). **D**: Segment OVerlap score for Non-Aggregation Prone Regions (SOV_nonAPR). The magnitude of each metric is shown in the y axis. Data was not calculated for AggreProt_static predictors in panels **C** and **D**

3.3. Introduction of mutations to the APRs identified by AggreProt reduces aggregation and increases solubility of LinB

Next, we tested our newly trained predictor on a model haloalkane dehalogenase LinB. HLDs (EC3.8.1.5) are well-characterised model enzymes belonging to a larger superfamily of α/β hydrolases that hydrolytically cleave carbon-halogen bonds in a broad range of aliphatic halogenated compounds. LinB has been isolated from bacterium *Sphingobium japonicum UT26* and its ability to convert persistent manmade pollutants and xenobiotics, most notably hexachlorocyclohexane, makes it a valuable target for industrial biodegradation or bioremediation[71–73]. Its broader and efficient application is, however, hindered by its poor solubility and low expression yield compared to the rest of structurally similar HLDs. The enzyme was previously stabilised by approx. 23 °C in terms of apparent melting temperature using a combination of energy calculations, disulfide bridge design, and molecular dynamics simulations[74]. However, the remarkable increase in its thermostability has not been accompanied by improvement in yield and solubility suggesting distinct determinants of LinB's

conformational and colloidal stability. Our previous efforts to identify APRs in LinB by the state-of-the-art predictors and design solubilizing mutations have been futile.

Using the sequential model of AggreProt we identified 7 regions within the LinB sequence whose score was higher compared to the rest of the sequence and could thus represent potential APRs (**Figure 3A**). Two of these APRs were also consensually detected using other aggregation prediction algorithms, whereas the others were identified only by a few or none (**Supplementary figure 3**). We immediately excluded the three stretches towards the N-terminus since they are buried within the central beta-sheet of the hydrolase core (**Figure 3A**, grey APRs), and focused on the remaining four. We defined each APR empirically as the residues with the highest aggregation score within the local sequence region around the maximum. Next, we selected residues that were near the centre or the edges of the APR and evaluated changes in aggregation score upon their *in silico* saturation mutagenesis. We purposely selected APR residues with different degrees of solvent exposure to evaluate the importance of SASA for the design of aggregation suppressing mutations. The five residues that lowered the aggregation the most regardless of the APR according to AggreProt were R, K, P, E, and D, which are known to be the "aggregation gatekeepers"[75]. Conversely, hydrophobic residues such as V, I, or C were among the worst, as expected. We further analysed the proposed mutations by the HotSpot Wizard[66] and Rosetta[67] to assess their mutability and effect on conformational stability, respectively. Finally, we selected 3-4 mutations from each APR, yielding 14 LinB variants for experimental validation.

We expressed all proteins in two batches of seven mutants + wild type each, and purified them from cell free extract using a combination of metallo-affinity and size-exclusion (SEC) chromatography. We manage to obtain all but one (I286D) variant to excellent purity (>95% based on SDS-PAGE) and homogeneity (>99% based on SEC). We evaluated the solubility of each variant from (i) the SDS-PAGE analysis of the total, soluble, and insoluble fractions during expression, and (ii) the final yield of monomeric protein. Unfortunately, the former strategy proved to be highly irreproducible with results of two large-scale and two small-scale expressions of wild-type alone differing by 80% (**Supplementary Figure 4**). In contrast, the deviation between yields from two large-scale expression and purification batches of wild type differed by only 11% so we used this parameter for comparing soluble expression of individual variants (Figure XXX c, grey). Notably, the expression and purification of all variants has been carried out over the course of two consecutive weeks by a single operator, using the same batches of media, incubation times, and purification protocol to ensure maximum comparability.

In total, eight out of fourteen variants had significantly improved yield, two mutations had neutral effect, and four variants had lower yield compared to the wild type (**Figure 3c**, grey). The latter are mutations of the buried residues, confirming our hypothesis regarding the importance of SASA during design (**Table 1**). The biggest improvements of yield were achieved with mutations in the APR2 (residues 218-225), APR3 (267-274) and APR4 (284-290) confirming their correct identification by AggreProt. Arguably, APR1 has been identified correctly (judged by the Q172R, Table 1) but more efficient suppression of its effect will require further designing effort. To assess the effect of the mutations on the aggregation propensity of LinB, we carried out a temperature melting scan and used static light scattering (SLS) as a measure of the aggregate formation (**Figure 3c**, black; **Supplementary Figure 5**). Similarly to yield, the mutations of buried residues increased aggregation (I286D which introduces buried negative charge could not even be obtained in monomeric form). The rest of mutations suppressed the aggregation or had a neutral effect (**Figure 3c**, black). We found weak negative correlation ($R^2 = 0.36$, PCC = -0.60) between changes in aggregation and soluble expression (**Figure 3d**), suggesting that these two properties are weakly linked. On average, introduction of negative charge (mutations to E or D) has a better effect on aggregation and yield compared

to the positive charges (R or K). However, a systematic screening of more mutants is needed before broader generalisation can be drawn.

Finally, we picked the best mutant from each APR for subsequent validation (highlighted by a coloured box in **Figure 3c**). We monitored their aggregation kinetics at 37 °C for 24 hours using SLS and Thioflavin T (ThT), which is an established amyloid fluorescence probe (**Supplementary Figure 6A, B**). The aggregation kinetics are characterised by fast increase of SLS signal followed by its slower attenuation and subsequent slow increase into plateau. The ThT signal shows a short lag phase on a time scale corresponding to the fast phase of SLS signal, and is followed by an exponential increase to a final plateau. The multiphasic nature of both signals suggests a complex aggregation mechanism of LinB whose description is beyond the scope of this work. However, the final SLS and ThT signals report well on the amount of aggregates in the samples (**Supplementary Figure 6C**). The SLS signals of the single point mutants in APR1 and 3 are ca. 50% higher compared to the wild type, suggesting that their aggregation propensities have not been significantly improved. In contrast, double-point mutants in both APR2 and APR4 show remarkably reduced aggregation based on both SLS and ThT. This is further supported by additional analysis of their soluble fraction upon centrifugation using flow-induced dispersion analysis (FIDA), which detects only 40% of residual monomer in the wild type sample, but 76 and 78% in case of LinB165 (A218P + R225E, APR2) and LinB173 (A284E + A288E, APR4), respectively (**Supplementary Figure 7**). This validates the results of our initial screening and confirms that the aggregation propensity of the designed LinB variants has been significantly decreased. Moreover, we further verified that the activity of both LinB variants towards 1,2-Dibromoethane has not been compromised by the mutations which makes them perfect targets for future research and biotechnological applications.

In summary, AggreProt identified four APRs in haloalkane dehalogenase LinB which were successfully suppressed by mutagenesis leading to LinB variants with increased solubility and yield, and decreased aggregation propensity.

Figure 3: Identification of APRs in the model haloalkane dehalogenase LinB and their suppression by computational design of mutations. a) Aggregation profile of LinB sequence.The AggreProt score for each LinB residue (black line

and symbols) is overlaid with the relative SASA score (grey areas). Residues selected for mutagenesis are marked by red. The zoom-ins of the seven regions with the highest aggregation score are shown on top. The APRs1, 2, 3, and 4 are coloured by red, green, blue, and violet, respectively. **b) Structural context of the APRs** identified by AggreProt. PDB-ID of the LinBwt structure - 1cv2 . **c) Change in aggregation propensity and yield by the designed mutations.** Total yield is calculated from the amount of soluble monomer isolated using SEC. Aggregation score is taken as a maximum value of the SLS signal during melting scans of the variants at 0.275 mg/ml and 1°C/min scan rate. The relative scores are calculated as wt - wt/mut. The mutations subjected to subsequent characterization are color coded according to the respective APR from a and highlighted with the box. **D) Correlation between solubility and aggregation.** Relative scores from c are plotted and their correlation depicted by the red dashed line. The variants selected for further characterization are highlighted.

Table 1: Design of mutations to the APRs that suppress aggregation. The SASA (solvent-accessible surface area) has been calculated from the structure (PDB-ID:1MJ5) using DSSP[65,76] and ranges from 0 (completely buried - B) to 100 (completely exposed - E). HotSpot Wizards (HSW) score provides mutability of a given residue based on the multiple sequence alignment of homologous sequences and ranges from 0 (not mutable) to 10 (mutable). Yield of the soluble monomeric protein from 1 L of bacterial culture. The value given for wild type is a mean and standard deviation between two individual batches. Maximum SLS corresponds to the highest values experimentally measured during melting scan from 15 to 95 °C at 1 °C/min scan rate at 1 mg.ml^{-1} of protein.

LinB variant	APR residues	Mutations	SASA (%)	Buried/Exposed	AggreProt score	HSW mutability	yield (mg.L^{-1})	Maximum SLS (AU)
wt	-	-	-	-			13.5 ± 1.5	161632 ± 2059
161		V170K	0.0	B	0.58	5	6	161480 ± 1571
162	167-175	**Q172R**	**50**	**E**	**0.52**	**8**	**21**	**159090 ± 1688**
163		L164R + G176R	5.2, 57.0	B,E	0.18, 0.26	6, 8	10	197672 ± 1264
164		A222E	43.5	E	0.42	6	28	120314 ± 8404
165	218-222	**A218P + R225E**	**64.3, 65.0**	**E,E**	**0.18, 0.15**	**6, 7**	**28**	**94883 ± 329**
166		V221H	17.4	B	0.43	6	30	149092 ± 3376
167		**A271G**	**7.8**	**B**	**0.39**	**7**	**28**	**139457 ± 13647**
168	267-275	A271R	7.8	B	0.39	7	18	140772 ± 295
169		A269P	47.0	E	0.33	8	15	143063 ± 5251
170		T267R	43.3	E	0.30	9	15	173294 ± 521
171		I286D	0.0	B	0.54	4	3	n.d.
172	283-289	A284E	40.9	E	0.29	8	19	111726 ± 4623
173		**A284E + A288E**	**40.9, 24.3**	**E,E**	**0.29, 0.42**	**8, 6**	**22**	**49941 ± 369**
174	55-65	C61P	0.0	B	0.58	4.0	9	231116 ± 5261

3.4. Experimental validation on hexapeptides reveals inaccuracies in the annotations of aggregation databases.

Finally, we experimentally validated the AggreProt predictions using a set of 34 hexapeptides. These included (i) 7 peptides corresponding to the APRs identified by AggreProt in LinBwt, (ii) 10 LinB-derived peptides carrying mutations aiming to suppress the aggregation described above, and (iii) 17 peptides selected from proteins in AmyPro database (**Supplementary Table 1**). The first two subsets were selected to test the performance of AggreProt on the identified APRs outside their structural context in LinB. Amyloid fibrils were found in all 7 samples after 1-month incubation, confirming their accurate identification.

Interestingly, aggregates were also observed in the samples of peptides carrying putative aggregation-suppressing mutations (**Figure 4b**). However, the amount of aggregates was generally lower and their morphology varied from amorphous clumps and hydrogels to short fibrils, indicating that the aggregation propensity of the peptides was reduced, but not completely suppressed by the mutations. The only peptides that showed absolutely no signs of aggregation were the two derived from LinB APR4 (corresponding to LinB167, and 168). The results obtained with these two subsets of hexapeptides together with the model case above illustrate that although AggreProt was successful in identification of APRs, the design of the suppressing mutations requires further optimization.

Next, we experimentally validated the aggregation propensity of 17 hexapeptides selected from a subset of proteins from AmyPro database[50]. AmyPro is one of the several databases that compiles sequences of aggregation-prone proteins together with manually curated amyloidogenic sequence regions (APRs) found in literature. Noticeably, a lot of the annotated APRs are unusually long as discussed in the *Model validation, comparison, and benchmarking* section above (> 50 amino acids). We therefore revisited the primary literature of several proteins from the database to investigate how the annotation has been made. One of the examples is γ-crystallin D (P07320) which is a protein found in human eye lenses and its aggregation is a hallmark of cataract[77,78]. Based on the AmyPro database, residues 1-79 are non-amyloidogenic whereas residues 80-163 are annotated as APR (**Figure 4a**, green and red background, respectively). The annotation has been made based on the study which used pepsin digestion of the aggregates followed by MS/MS identification of the amyloid core spanning the residues 81-163 as annotated in the AmyPro[79]. The fragment was further digested in-gel using trypsin and subsequent MALDI-MS analysis enabled identification of 6 peptide fragments (80-88, 100-115, 118-140, 143-152, 154-163) suggesting that the cleavage sites are accessible by the protease. Interestingly, these fragments nicely overlap with sequence stretches with high AggreProt score (**Figure 4a**) which might indicate that they play an important role in stabilisation of the amyloid core via steric-zipper interactions. To test this, we selected two hexapeptides corresponding to the residues found in the trypsin fragments (peptide 13: 129-135, peptide 14: 121-126) with the highest AggreProt score, and two hexapeptides (peptide 16: 94-99, peptide 17: 150-155) which were not identified by the trypsin digestion and their AggreProt score is close to zero (**Figure 4a**). Additionally, we included one more hexapeptide with high AggreProt score (peptide 15: 41-46) which is annotated as non-amyloid in the AmyPro despite experimentally characterised to form amyloid fibrils[80]. The TEM analysis of the hexapeptides after 4 weeks of incubation reveals that the AggreProt predictions were correct (**Figure 4a**) and demonstrating that the annotations in both AmyPro and CPAD2.0[49] databases are indeed not accurate. We identified further missannotations in six additional proteins from the AmyPro database (**Supplementary Figure 8**) which included 8 false positives and 6 false negatives (**Supplementary Table 1**). This is an exceedingly important finding (though not the first one of this kind[27]) since there are numerous new aggregation prediction algorithms that use these databases for benchmarking without any further scrutiny or experimental validation[52,81–83].

Altogether, we demonstrate successful utilisation of AggreProt in two experimental case studies. Using our newly developed algorithm, we identified seven APRs in a model enzyme and correctly accessed the biological relevance of four of them based on their solvent accessibility within the structure. Moreover, we were able to design mutations into these APRs that increase solubility and decrease aggregation propensity of the protein. We firmly believe that we can improve these properties even further in the future studies by systematically optimising the design strategy. Finally, AggreProt showed great accuracy (80 %) in identifying aggregation propensity of isolated hexapeptides which were wrongly annotated in the databases of protein aggregation. The newly characterised 34 hexapeptide will be deposited to the

WaltzDB2.0 database to expand its content and increase its impact for the further development of *in-silico* aggregation prediction tools in the future.

Figure 4: Experimental validation of AggreProt using hexapeptides identified in proteins from AmyPro database and LinB. a) Example of hexapeptide selection using gamma-crystallin D entry from AmyPro (CPAD 2.0) database. AggreProt score (black line and points) is not in agreement with the database annotation of amyloid (red shading) and non-amyloid (green shading) regions. Five hexapeptides were selected to represent potentially false negative (peptide 15; red), false positive (peptides 16, 17; green), and true positive (peptides 13, 14) amyloid annotation. Experimental validation by transmission electron microscopy (TEM, insets; red) and Thioflavin T assay confirmed that the AggreProt predictions are correct and three of the respective segments are missanotated in the databases (sequences corresponding to peptides 15-17). Similar cases of missanotations were found in other proteins in the databases (**Supplementary Table 1**). * experimentally confirmed by TEM and ThT, ** annotation according to AmyPro (http://amypro.net/) and CPAD2.0 (https://web.iitm.ac.in/bioinfo2/cpad2). **b) Analysis of peptides derived from predicted APRs in LinB.** Seven hexapeptides derived from the APRs identified by LinB (top) were synthesised together with ten variants carrying mutations designed to suppress their aggregation (marked by red and green letters ,respectively). All seven hexapeptides derived from LinBwt were experimentally verified to form fibrils (representative micrographs shown). The presence or absence of fibrils in the samples of peptides carrying designed mutations is indicated by tick mark or cross, respectively. The correspondence between the AggreProt score (aggregation threshold > 0.25) and experimental results is encoded by green colour, failed prediction of the outcome by red. **c) Results of experimental validation of AggreProt prediction using a set of 34 hexapeptides derived from AmyPro database and model protein LinB.**

Conclusions

In this study, we trained two novel protein aggregation predictors based on deep neural networks using two different feature representations of hexapeptides from WaltzDB. The predictors achieved great performance on the hexapeptide validation dataset (AuROCC scores of 0.85 and 0.89 for linear and static model, respectively). Using whole sequence validation datasets based on the AmyPro database, the performance was lower, yet still equivalent or better compared to the state-of-the-art algorithms. The lower scores can be attributed to the discrepancy between sequence length of the training (i.e., six amino acids) and validation (full sequences), small size of the training dataset (1406 entries), binary labels in the AmyPro database, and most importantly, to the incorrect annotations in the database discovered during this work.

To truly test the performance of our predictors and to demonstrate its applicability, we carried out two experimental validations. First, we selected 34 hexapeptides identified by AggreProt in model haloalkane dehalogenase LinB and seven other proteins from AmyPro, and experimentally determined their aggregation. The experimental results matched the predictions in 79% of cases and, most importantly, confirmed our initial scepticism regarding missannotations of long (>50 amino acid) APRs in the validation dataset. Based on this experience, we advise caution when relying on purely in silico validation of computational aggregation predictors. The newly characterised 34 hexapeptide will be deposited to the WaltzDB2.0 database to extend its content and increase its impact for further development of *in-silico* aggregation prediction tools in the future.

Finally, we showcase an example of the AggreProt use for protein design. Specifically, we correctly identified seven APRs in a model haloalkane dehalogenase LinB and correctly accessed the biological relevance of four of them based on their solvent accessibility within the structure. Moreover, we were able to design mutations into these APRs that increase solubility and decrease aggregation propensity of the protein, resulting in up to doubled soluble protein expression compared to the wild type without compromising its functionality. We firmly believe that we can improve these properties even further in the future studies by systematically optimising the design strategy, making AggreProt a valuable tool for solubilizing proteins.

Acknowledgements

Computational resources were provided by the e-INFRA CZ and ELIXIR-CZ projects (LM2023055 and LM2018140), supported by the Ministry of Education, Youth and Sports of the Czech Republic. Authors thank the RECETOX Research Infrastructure (No. LM2023069) and Operational Programme Research, Development and Education (the CETOCOEN EXCELLENCE project No. CZ.02.1.01/0.0/0.0/17_043/0009632). This research was also supported by the European Union's Horizon 2020 Research and Innovation Programme under grant agreement No. 857560 and by the project National Institute for Neurology Research (nr. LX22NPO5107 MEYS): Financed by European Union – Next Generation EU.

References

1.   Wodak, S. J., Vajda, S., Lensink, M. F., Kozakov, D. & Bates, P. A. Critical Assessment of Methods for Predicting the 3D Structure of Proteins and Protein Complexes. *Annu Rev Biophys* **52**, 183–206 (2023).

2.   Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583–589 (2021).

3.   Baek, M. *et al.* Accurate prediction of protein structures and interactions using a three-track neural network. *Science (1979)* **373**, 871–876 (2021).

4.   Mirdita, M. *et al.* ColabFold: making protein folding accessible to all. *Nat Methods* **19**, 679–682 (2022).

5.   Pinheiro, F., Santos, J. & Ventura, S. AlphaFold and the amyloid landscape. *J Mol Biol* **433**, 167059 (2021).

6.   Knowles, T. P. J., Vendruscolo, M. & Dobson, C. M. The amyloid state and its association with protein misfolding diseases. *Nat Rev Mol Cell Biol* **15**, 384–396 (2014).

7.   Chiti, F. & Dobson, C. M. Protein Misfolding, Amyloid Formation, and Human Disease: A Summary of Progress Over the Last Decade. *Annu Rev Biochem* **86**, 27–68 (2017).

8.   Gerasimavicius, L., Livesey, B. J. & Marsh, J. A. Loss-of-function, gain-of-function and dominant-negative mutations have profoundly different effects on protein structure. *Nat Commun* **13**, 3895 (2022).

9.   Kuroda, Y. Biophysical studies of protein solubility and amorphous aggregation by systematic mutational analysis and a helical polymerization model. *Biophys Rev* **10**, 473–480 (2018).

10.   Kuroda, Y. Biophysical studies of amorphous protein aggregation and in vivo immunogenicity. *Biophys Rev* **14**, 1495–1501 (2022).

11.   Smoluchowski, M. V. Drei Vortrage uber Diffusion, Brownsche Bewegung und Koagulation von Kolloidteilchen. *Physik. Zeit.* **17**, 557–585 (1916).

12.   Roberts, C. J. Non-native protein aggregation kinetics. *Biotechnol Bioeng* **98**, 927–938 (2007).

13.   Roberts, C. J., Darrington, R. T. & Whitley, M. B. Irreversible aggregation of recombinant bovine granulocyte-colony stimulating factor (bG-CSF) and implications for predicting protein shelf life. *J Pharm Sci* **92**, 1095–1111 (2003).

14.   Andrews, J. M. & Roberts, C. J. A Lumry−Eyring Nucleated Polymerization Model of Protein Aggregation Kinetics: 1. Aggregation with Pre-Equilibrated Unfolding. *J Phys Chem B* **111**, 7897–7913 (2007).

15.   Li, Y., Weiss, W. F. & Roberts, C. J. Characterization of High-Molecular-Weight Nonnative Aggregates and Aggregation Kinetics by Size Exclusion Chromatography With Inline Multi-Angle Laser Light Scattering. *J Pharm Sci* **98**, 3997–4016 (2009).

16.   Brummitt, R. K. *et al.* Nonnative Aggregation of an IgG1 Antibody in Acidic Conditions: Part 1. Unfolding, Colloidal Interactions, and Formation of High-Molecular-Weight Aggregates. *J Pharm Sci* **100**, 2087–2103 (2011).

17.   Arosio, P., Rima, S., Lattuada, M. & Morbidelli, M. Population Balance Modeling of Antibodies Aggregation Kinetics. *J Phys Chem B* **116**, 7066–7075 (2012).

18.   Zidar, M., Kuzman, D. & Ravnik, M. Characterisation of protein aggregation with the Smoluchowski coagulation approach for use in biopharmaceuticals. *Soft Matter* **14**, 6001–6012 (2018).

19.   Hirota, N., Edskes, H. & Hall, D. Unified theoretical description of the kinetics of protein aggregation. *Biophys Rev* **11**, 191–208 (2019).

20.   Szała-Mendyk, B., Drajkowska, A. & Molski, A. Modified Smoluchowski Rate Equations for Aggregation and Fragmentation in Finite Systems. *J Phys Chem B* **127**, 6154–6162 (2023).

21.   Rambaran, R. N. & Serpell, L. C. Amyloid fibrils. *Prion* **2**, 112–117 (2008).

22.   Michaels, T. C. T. *et al.* Amyloid formation as a protein phase transition. *Nature Reviews Physics* **5**, 379–397 (2023).

23.   Sawaya, M. R. *et al.* Atomic structures of amyloid cross-β spines reveal varied steric zippers. *Nature* **447**, 453–457 (2007).

24. Nelson, R. *et al.* Structure of the cross-β spine of amyloid-like fibrils. *Nature* **435**, 773–778 (2005).

25. Riek, R. The Three-Dimensional Structures of Amyloids. *Cold Spring Harb Perspect Biol* **9**, a023572 (2017).

26. Thompson, M. J. *et al.* The 3D profile method for identifying fibril-forming segments of proteins. *Proceedings of the National Academy of Sciences* **103**, 4074–4078 (2006).

27. Louros, N., Orlando, G., De Vleeschouwer, M., Rousseau, F. & Schymkowitz, J. Structure-based machine-guided mapping of amyloid sequence space reveals uncharted sequence clusters with higher solubilities. *Nat Commun* **11**, 3314 (2020).

28. Fändrich, M. *et al.* Amyloid fibril polymorphism: a challenge for molecular imaging and therapy. *J Intern Med* **283**, 218–237 (2018).

29. Lövestam, S. *et al.* Disease-specific tau filaments assemble via polymorphic intermediates. *Nature* **625**, 119–125 (2024).

30. Wang, H. *et al.* Polymorphic Aβ42 fibrils adopt similar secondary structure but differ in cross-strand side chain stacking interactions within the same β-sheet. *Sci Rep* **10**, 5720 (2020).

31. Sawaya, M. R., Hughes, M. P., Rodriguez, J. A., Riek, R. & Eisenberg, D. S. The expanding amyloid family: Structure, stability, function, and pathogenesis. *Cell* **184**, 4857–4873 (2021).

32. van der Kant, R., Louros, N., Schymkowitz, J. & Rousseau, F. Thermodynamic analysis of amyloid fibril structures reveals a common framework for stability in amyloid polymorphs. *Structure* **30**, 1178-1189.e3 (2022).

33. Conchillo-Solé, O. *et al.* AGGRESCAN: a server for the prediction and evaluation of 'hot spots' of aggregation in polypeptides. *BMC Bioinformatics* **8**, 65 (2007).

34. Tartaglia, G. G. & Vendruscolo, M. The Zyggregator method for predicting protein aggregation propensities. *Chem Soc Rev* **37**, 1395 (2008).

35. Zibaee, S., Makin, O. S., Goedert, M. & Serpell, L. C. A simple algorithm locates β-strands in the amyloid fibril core of α-synuclein, Aβ, and tau using the amino acid sequence alone. *Protein Science* **16**, 906–918 (2007).

36. Walsh, I., Seno, F., Tosatto, S. C. E. & Trovato, A. PASTA 2.0: an improved server for protein aggregation prediction. *Nucleic Acids Res* **42**, W301–7 (2014).

37. Ahmed, A. B., Znassi, N., Château, M. & Kajava, A. V. A structure-based approach to predict predisposition to amyloidosis. *Alzheimer's & Dementia* **11**, 681–690 (2015).

38. Hamodrakas, S. J., Liappa, C. & Iconomidou, V. A. Consensus prediction of amyloidogenic determinants in amyloid fibril-forming proteins. *Int J Biol Macromol* **41**, 295–300 (2007).

39. Bryan, A. W. Prediction of parallel in-register amyloidogenic beta-structures In highly beta-rich protein sequences by pairwise propensity analysis. (Harvard University, 2009).

40. Maurer-Stroh, S. *et al.* Exploring the sequence determinants of amyloid structure using position-specific scoring matrices. *Nat Methods* **7**, 237–242 (2010).

41. Tartaglia, G. G., Cavalli, A., Pellarin, R. & Caflisch, A. Prediction of aggregation rate and aggregation-prone segments in polypeptide sequences. *Protein Science* **14**, 2723–2734 (2005).

42. Fernandez-Escamilla, A.-M., Rousseau, F., Schymkowitz, J. & Serrano, L. Prediction of sequence-dependent and mutational effects on the aggregation of peptides and proteins. *Nat Biotechnol* **22**, 1302–1306 (2004).

43. Tsolis, A. C., Papandreou, N. C., Iconomidou, V. A. & Hamodrakas, S. J. A Consensus Method for the Prediction of 'Aggregation-Prone' Peptides in Globular Proteins. *PLoS One* **8**, e54175 (2013).

44. Emily, M., Talvas, A. & Delamarche, C. MetAmyl: A META-Predictor for AMYLoid Proteins. *PLoS One* **8**, e79722 (2013).

45. Chennamsetty, N., Voynov, V., Kayser, V., Helk, B. & Trout, B. L. Prediction of Aggregation Prone Regions of Therapeutic Proteins. *J Phys Chem B* **114**, 6614–6624 (2010).

46. Zambrano, R. *et al.* AGGRESCAN3D (A3D): server for prediction of aggregation properties of protein structures. *Nucleic Acids Res* **43**, W306–13 (2015).

47. Kuriata, A. *et al.* Aggrescan3D (A3D) 2.0: prediction and engineering of protein solubility. *Nucleic Acids Res* **47**, W300–W307 (2019).

48. Sormanni, P., Aprile, F. A. & Vendruscolo, M. The CamSol method of rational design of protein mutants with enhanced solubility. *J Mol Biol* **427**, 478–490 (2015).

49. Rawat, P. *et al.* CPAD 2.0: a repository of curated experimental data on aggregating proteins and peptides. *Amyloid* **27**, 128–133 (2020).

50. Varadi, M., De Baets, G., Vranken, W. F., Tompa, P. & Pancsa, R. AmyPro: a database of proteins with validated amyloidogenic regions. *Nucleic Acids Res* **46**, D387–D392 (2018).

51. Louros, N. *et al.* WALTZ-DB 2.0: an updated database containing structural information of experimentally determined amyloid-forming peptides. *Nucleic Acids Res* **48**, D389–D393 (2020).

52. Prabakaran, R., Rawat, P., Kumar, S. & Michael Gromiha, M. ANuPP: A Versatile Tool to Predict Aggregation Nucleating Regions in Peptides and Proteins. *J Mol Biol* **433**, 166707 (2021).

53. Wojciechowski, J. W. & Kotulska, M. PATH - Prediction of Amyloidogenicity by Threading and Machine Learning. *Sci Rep* **10**, 7721 (2020).

54. Niu, M., Li, Y., Wang, C. & Han, K. RFAmyloid: A Web Server for Predicting Amyloid Proteins. *Int J Mol Sci* **19**, 2071 (2018).

55. Keresztes, L. *et al.* The Budapest Amyloid Predictor and Its Applications. *Biomolecules* **11**, 500 (2021).

56. Orlando, G., Silva, A., Macedo-Ribeiro, S., Raimondi, D. & Vranken, W. Accurate prediction of protein beta-aggregation with generalized statistical potentials. *Bioinformatics* **36**, 2076–2081 (2020).

57. Li, Y., Zhang, Z., Teng, Z. & Liu, X. PredAmyl-MLP: Prediction of Amyloid Proteins Using Multilayer Perceptron. *Comput Math Methods Med* **2020**, 1–12 (2020).

58. Tian, J., Wu, N., Guo, J. & Fan, Y. Prediction of amyloid fibril-forming segments based on a support vector machine. *BMC Bioinformatics* **10**, S45 (2009).

59. Burdukiewicz, M. *et al.* Amyloidogenic motifs revealed by n-gram analysis. *Sci Rep* **7**, 12961 (2017).

60. Navarro, S. & Ventura, S. Computational methods to predict protein aggregation. *Curr Opin Struct Biol* **73**, 102343 (2022).

61. Schymkowitz, J. *et al.* The FoldX web server: an online force field. *Nucleic Acids Res* **33**, W382–W388 (2005).

62. Ruopp, M. D., Perkins, N. J., Whitcomb, B. W. & Schisterman, E. F. Youden Index and Optimal Cut-Point Estimated from Observations Affected by a Lower Limit of Detection. *Biometrical Journal* **50**, 419–430 (2008).

63. Zemla, A., Venclovas, Č., Fidelis, K. & Rost, B. A modified definition of Sov, a segment-based measure for protein secondary structure prediction assessment. *Proteins: Structure, Function, and Genetics* **34**, 220–223 (1999).

64. Abadi, M. *et al.* TensorFlow: Large-Scale Machine Learning on Heterogeneous Systems. Preprint at https://doi.org/10.5281/zenodo.4724125 (2015).

65. Kabsch, W. & Sander, C. Dictionary of protein secondary structure: Pattern recognition of hydrogen-bonded and geometrical features. *Biopolymers* **22**, 2577–2637 (1983).

66. Sumbalova, L., Stourac, J., Martinek, T., Bednar, D. & Damborsky, J. HotSpot Wizard 3.0: web server for automated design of mutations and smart libraries based on sequence input information. *Nucleic Acids Res* **46**, W356–W362 (2018).

67. Kellogg, E. H., Leaver-Fay, A. & Baker, D. Role of conformational sampling in computing mutation-induced changes in protein structure and stability. *Proteins* **79**, 830–838 (2011).

68. Iwasaki, I., Utsumi, S. & Ozawa, T. New colorimetric determination of chloride using mercuric thiocyanate and ferric ion. *Bull Chem Soc Jpn* **25**, 226 (1952).

69. Chou, P. Y. & Fasman, G. D. Prediction of protein conformation. *Biochemistry* **13**, 222–245 (1974).

70. Planas-Iglesias, J. *et al.* AggreProt: A Web Server for Predicting and Engineering Aggregation Prone Regions in Proteins. *Nucleic Acids Res* (2024).

71. Moriuchi, R. *et al.* Stepwise enhancement of catalytic performance of haloalkane dehalogenase LinB towards β-hexachlorocyclohexane. *AMB Express* **4**, 72 (2014).

72. Bidmanova, S., Damborsky, J. & Prokop, Z. Immobilization of Haloalkane dehalogenase LinB from Sphingobium japonicum UT26 for Biotechnological Applications. *Journal of Biocatalysis & Biotransformation* **02**, (2013).

73. Ito, M. *et al.* Degradation of β-hexachlorocyclohexane by haloalkane dehalogenase LinB from γ-hexachlorocyclohexane-utilizing bacterium Sphingobium sp. MI1205. *Arch Microbiol* **188**, 313–325 (2007).

74. Floor, R. J. *et al.* Computational Library Design for Increasing Haloalkane Dehalogenase Stability. *ChemBioChem* **15**, 1660–1672 (2014).

75. Beerten, J. *et al.* Aggregation gatekeepers modulate protein homeostasis of aggregating sequences and affect bacterial fitness. *Protein Engineering Design and Selection* **25**, 357–366 (2012).

76. Joosten, R. P. *et al.* A series of PDB related databases for everyday needs. *Nucleic Acids Res* **39**, D411–D419 (2011).

77. Surguchev, A. & Surguchov, A. Conformational diseases: Looking into the eyes. *Brain Res Bull* **81**, 12–24 (2010).

78. Wang, Y., Lomakin, A., McManus, J. J., Ogun, O. & Benedek, G. B. Phase behavior of mixtures of human lens proteins Gamma D and Beta B1. *Proceedings of the National Academy of Sciences* **107**, 13282–13287 (2010).

79. Moran, S. D., Decatur, S. M. & Zanni, M. T. Structural and Sequence Analysis of the Human γD-Crystallin Amyloid Fibril Core Using 2D IR Spectroscopy, Segmental ^{13}C Labeling, and Mass Spectrometry. *J Am Chem Soc* **134**, 18410–18416 (2012).

80. Abu-Hussien, M. *et al.* Inhibition of amyloid fibrillation of γD-crystallin model peptide by the cochineal Carmine. *Int J Biol Macromol* **169**, 342–351 (2021).

81. Charoenkwan, P., Kanthawong, S., Nantasenamat, C., Hasan, Md. M. & Shoombuatong, W. iAMY-SCM: Improved prediction and analysis of amyloid proteins using a scoring card method with propensity scores of dipeptides. *Genomics* **113**, 689–698 (2021).

82. Charoenkwan, P. *et al.* AMYPred-FRL is a novel approach for accurate prediction of amyloid proteins by using feature representation learning. *Sci Rep* **12**, 7697 (2022).

83. Kang, S., Kim, M., Sun, J., Lee, M. & Min, K. Prediction of Protein Aggregation Propensity via Data-driven Approaches. (2023).