# HSV-1 infection induces a downstream-shift of the +1 nucleosome

Elena Weiß[1], Adam W. Whisnant[2,3], Thomas Hennig[2,3], Lara Djakovic[2], Lars Dölken[2,3,4#], Caroline C. Friedel[1#]

[1] Institute of Informatics, Ludwig-Maximilians-Universität München, Amalienstr. 17, 80333 Munich, Germany

[2] Institute for Virology and Immunobiology, Julius-Maximilians-University Würzburg, Versbacher Straße 7, 97078 Würzburg, Germany

[3] Institute for Virology, Medizinische Hochschule Hannover, Carl-Neuberg-Str. 1, 30625 Hannover, Germany

[4] Helmholtz Institute for RNA-based Infection Research (HIRI), Helmholtz-Center for Infection Research (HZI), 97080 Würzburg, Germany

[#] Corresponding authors

E-mail: lars.doelken@uni-wuerzburg.de; Caroline.Friedel@bio.ifi.lmu.de

**Abstract**

Herpes simplex virus 1 (HSV-1) infection induces a genome-wide loss of host transcriptional

activity and widespread disruption of host transcription termination. The latter leads to read-

through transcription for thousands of nucleotides beyond poly(A) sites and is associated with

induction of open chromatin downstream of genes. In this study, we show that lytic HSV-1

infection alters chromatin accessibility around host promoters resulting in an extension of

nucleosome-free regions at transcription start sites into downstream regions. This was most

prominent for highly expressed genes and still observed upon knockout of the viral immediate-

early proteins ICP0, ICP22, and ICP27 and the virion host shutoff protein *vhs*. ChIPmentation

analysis of the noncanonical histone variant H2A.Z, which is strongly enriched at +1 and -1

nucleosomes, indicated that the downstream extension of accessible chromatin at promoters

was linked to a downstream shift of +1 nucleosomes. In yeast, downstream shifts of +1

nucleosomes can be induced by RNA Polymerase II (Pol II) degradation. Accordingly,

irreversible depletion of Pol II from genes in human cells using α-amanitin altered H2A.Z

occupancy and +1 nucleosome positioning similar to what was observed in HSV-1 infection.

Consequently, treatment with phosphonoacetic acid (PAA) and knockout of ICP4, which both

prevent viral DNA replication and alleviate the loss of Pol II from host genes, largely abolished

the downstream extension of accessible chromatin in HSV-1 infection. In absence of viral DNA

replication, doxycycline-induced expression of ICP27, which redirects Pol II from gene bodies

into intergenic regions by disrupting transcription termination, can induce an attenuated effect

that is further enhanced by co-expression with ICP22. In summary, our study provides strong

evidence that HSV-1-induced depletion of Pol II from the host genome leads to a widespread

downstream shift of +1 nucleosomes at host gene promoters.

## Author summary

Herpes simplex virus 1 (HSV-1) infection leads to a profound shutoff of host transcription during lytic infection. Loss of RNA Polymerase II (Pol II) in yeast has previously been shown to relax +1 nucleosome positioning to more thermodynamically favorable sites downstream of transcription start sites. In this study, we show that a similar phenomenon is likely also at play in lytic HSV-1 infection. Sequencing of accessible chromatin revealed a widening of nucleosome-free regions at host gene promoters into downstream regions. By mapping genome-wide positions of the noncanonical histone variant H2A.Z enriched at +1 and -1 nucleosomes, we demonstrate a downstream shift of +1 nucleosomes for the majority of cellular genes in lytic HSV-1 infection. We confirm that chemical depletion of Pol II from genes leads to a downstream shift of +1 nucleosomes in human cells. Thus, changes in chromatin architecture at promoters in HSV-1 infection are likely a consequence of HSV-1-induced loss of Pol II activity from the host genome.

## Introduction

Herpes simplex virus 1 (HSV-1) is one of nine human herpesviruses and more than half of the global population are latently infected with HSV-1 (1-4). HSV-1 does not only cause the common cold sores but is also responsible for life-threatening diseases particularly in the immunocompromised (5). The HSV-1 genome consists of 152 kilobases (kb) of double-stranded DNA and encodes for at least 121 large open reading frames (ORFs) and >100 small ORFs (6). Upon lytic infection, four of the five viral immediate early (IE) genes, i.e., ICP0, ICP4, ICP22 and ICP27, hijack the host gene expression machinery to ensure viral replication during early stages of infection (7-11). This results in recruitment of RNA polymerase II (Pol II) from the host chromatin to viral genomes and a general shutoff of host transcription (7, 9, 11-13). Pol II recruitment to viral genomes is predominantly facilitated by the viral IE protein ICP4 (13), which effectively sequesters Pol II into the viral replication compartments (14). However, ICP4 already depletes Pol II from cellular promoters as early as 2 h post infection (p.i.), i.e. before the onset of viral genome replication, and independent of viral genome copy number (13). The global host transcription shutoff is further exacerbated by ICP22-mediated inhibition of host transcription elongation (15) and the viral *vhs* (virion host shutoff) protein. *vhs* is a nuclease delivered by the tegument of the incoming viral particles that cleaves both viral and host mRNAs (16-18). We recently demonstrated that *vhs* continuously degrades about 30 % of host mRNAs per hour during the first 8 h of high multiplicity lytic infection of primary human fibroblasts (19).

HSV-1 further disturbs host transcription by inducing a widespread disruption of host transcription termination (DoTT) resulting in read-through transcription beyond polyadenylation (pA) sites of host genes for tens of thousands of nucleotides (nt) (20). Read-through transcription commonly extends into downstream genes, leading to a seeming

transcriptional induction of these genes. DoTT negatively affects host gene expression in two

ways: First, as read-through transcripts are not exported (21), it effectively prevents translation

of newly transcribed mRNAs and thus dampens the host transcriptional response to infection.

Second, it redirects a large fraction of ongoing Pol II transcription from host gene bodies into

downstream intergenic regions and, thus, further exacerbates the depletion of Pol II from host

genes. We recently showed that ICP27 is sufficient for inducing DoTT (22). Interestingly,

transcription downstream of genes (DoG) is also induced in cellular stress responses (21, 23-

25). Accordingly, infection with an ICP27-null mutant still induces read-through transcription

albeit at much lower levels, presumably due to a virus-induced cellular stress response (26). In

contrast to stress-induced DoGs, HSV-1-induced DoTT is associated with a massive increase

of chromatin accessibility downstream of genes (denoted as downstream open chromatin

regions (dOCRs)) detectable by ATAC-seq (Assay for Transposase-Accessible Chromatin

using sequencing) (21). dOCR induction requires strong absolute levels of transcription

downstream of genes, thus it is predominantly observed for highly transcribed genes with strong

read-through. Treatment with phosphonoacetic acid (PAA) during HSV-1 infection, which

inhibits viral DNA replication (27, 28), results in increased dOCR induction (29). As viral DNA

redirects Pol II from the host chromatin, PAA treatment mitigates the loss of Pol II from the

host genome and thus increases absolute levels of transcription on host genes and – importantly

for dOCR induction – read-through regions. Although dOCRs can extend into downstream

genes following read-through transcription, this occurs only in cases of very strong read-

through transcription into these downstream genes. We recently showed that ICP22 is both

required and sufficient for inducing dOCRs in presence of ICP27-induced read-through (29).

As we previously only investigated chromatin accessibility *downstream* of genes, we now

further explore our previously published ATAC-seq data to investigate changes in chromatin

accessibility around host gene promoters. This revealed a broadening of open chromatin regions

at the promoter into regions downstream of transcription start sites (TSS) for the majority of

host genes. While it was consistently observed in absence of ICP0, ICP22, ICP27, and *vhs*,

PAA treatment and ICP4 knockout, both of which inhibit viral DNA replication and alleviate

the HSV-1-induced loss of host transcription, substantially reduced the extension of open

chromatin at the TSS. Weiner *et al.* previously showed that Pol II depletion in yeast induces

downstream shifts of +1 nucleosomes, in particular for highly expressed genes (30).

Transcribing Pol II disturbs histone-DNA interactions and at high transcription rates leads to

eviction of nucleosomes from the chromatin (31). Weiner *et al.* proposed that Pol II depletion

relaxes chromatin allowing nucleosomes to shift to more thermodynamically favorable sites.

The noncanonical histone H2A variant H2A.Z is strongly enriched at gene promoters at +1 and

-1 nucleosome positions (32, 33), likely as H2A.Z deposition increases nucleosome mobility

and makes DNA more accessible to the transcriptional machinery (34). We thus performed

ChIPmentation for H2A.Z during lytic HSV-1 infection to map +1 nucleosome positions. This

showed that changes in chromatin accessibility in HSV-1 infection reflected a downstream shift

of +1 nucleosomes. We observed similar changes in H2A.Z occupancy upon treatment with α-

amanitin, a deadly toxin found in Amanita mushrooms, which inhibits transcription by

preventing translocation of Pol II and triggering degradation of Rpb1, the largest subunit of Pol

II (35). Our findings thus provide strong evidence that depletion of Pol II from host genes during

HSV-1 infection leads to the downstream shift of +1 nucleosomes to more thermodynamically

favorable sites.

## Results

**Widespread extension of chromatin-free regions downstream of transcription start sites**

To investigate changes in chromatin accessibility within promoter regions during HSV-1 infection, we re-analyzed our recently published ATAC-seq data (29) for mock and wild-type (WT) HSV-1 infection of human fetal foreskin fibroblasts (HFF) as well as infection with null mutant viruses for the HSV-1 IE proteins ICP0, ICP22, and ICP27 as well as the tegument-delivered late protein *vhs*. Samples for mock, WT, Δ*vhs* and ΔICP27 infection were collected at 8 h p.i., while samples for ΔICP0 and ΔICP22 infection were collected at 12 h p.i. To assess chromatin accessibility changes in promoter regions, we identified genomic regions with differential ATAC-seq read density using our recently published RegCFinder method (36). RegCFinder searches for genomic subregions for which the read distribution differs between two conditions (e.g., mock and WT infection) and is targeted towards regions of interest by specifying genomic windows as input (**Sup. Fig 1a** in **S1 File**). For each input window, subregions are identified that show differences in the distribution of reads within this window. Statistical significance and log2 fold-changes for identified subregions are determined by DEXSeq (37). For our purposes, we defined the windows of interest as the ± 3kb promoter region around the TSS of 7,649 human genes. These TSSs were previously defined in our analysis of promoter-proximal Pol II pausing in HSV-1 WT infection (38) based on a re-analysis of published precision nuclear run-on sequencing (PRO-seq) and PROcap-seq (a variation of PRO-seq) data of flavopiridol-treated uninfected HFF. CDK9 inhibition by flavopiridol arrests Pol II in a paused state at the TSS, allowing the mapping of Pol II initiation sites with both PRO- and PROcap-seq. Filtering for high-confidence TSSs that were confirmed by both data sets and within a maximum distance of 500 bp to the nearest annotated gene on the same strand identified the major TSS for 7,649 genes. Using these 6 kb promoter windows as input, we applied RegCFinder to our ATAC-seq data for WT and all null-mutant infections

in comparison to mock to identify promoter subregions that show differences in the distribution

of open chromatin between infection and mock. We did not filter these promoter windows

beforehand to exclude genes for which dOCRs extended from an upstream gene into their

promoter window for two reasons: (i) Since dOCRs represent changes in chromatin

accessibility, RegCFinder should by design detect them; (ii) The farther the distance from the

upstream gene, the less pronounced are changes in chromatin accessibility in dOCR regions,

making it difficult to distinguish them from background. Any approach we tested to filter

promoter windows overlapping with dOCRs from upstream genes before the RegCFinder

analysis either excluded too many or too few windows. Thus, we decided to first run

RegCFinder and then investigate which of the detected changes were due to induction of dOCRs

from upstream genes.

RegCFinder identified 23,000-24,000 differential subregions (in the following also denoted as

RegC, short for regions of change) for each virus infection compared to mock (WT: 23,357

RegC, ΔICP0: 23,532, ΔICP22: 24,640, ΔICP27: 23,523, Δ*vhs*: 23,716, **Sup. Fig 1b** in **S1 File**).

Between 29 and 37 % of these showed a statistically significant difference in read density within

the corresponding 6 kb promoter windows upon infection (multiple testing adjusted p-value

(adj. p.) $\leq$ 0.01, WT: 8,552 RegC, ΔICP0: 6,740, ΔICP22: 5,710, ΔICP27: 7,188, Δ*vhs*: 6,860).

These represented between 3,129 (ΔICP22 infection) and 4,391 (WT) genes (**Sup. Fig 1c** in **S1**

**File**). The lower fraction of statistically significant differential regions in ΔICP22 infection is

likely due to the relatively low number of ATAC-seq reads mapping to the host genome

compared to the other infections (~3.8-fold fewer, see **Sup. Table 1** in **S2 File**). The main

reason for this was the high fraction of reads mapping to the HSV-1 genome in ΔICP22

infection (70-94%) compared to WT infection (~52%). We confirmed this observation in an

independent experiment for 8 and 12 h infection with the ΔICP22 mutant and its parental WT

strain F (WT-F, **Sup. Table 1** in **S2 File**).

**Fig 1a** visualizes the positions of identified RegC for the 4,981 promoter windows containing

at least one statistically significant (adj. p. ≤ 0.01) differential region for at least one virus

infection compared to mock. Log2 fold-changes for statistically significant regions are

illustrated in **Sup. Fig 1d** in **S1 File**.  Here, blue indicates that relative read density in that

subregion is increased during virus infection compared to mock (infection>mock, denoted as i-

RegC in the following) and red that relative read density in that subregion is higher in mock

compared to infection (mock>infection, denoted as m-RegC). Strikingly, our results showed a

highly homogeneous picture for WT and null-mutant infections compared to mock, with

changes in the same direction observed at approximately the same locations. To identify distinct

patterns of changes in chromatin accessibility, we performed hierarchical clustering on the

RegC location heatmap in **Fig 1a** and selected a cutoff on the dendrogram to obtain 14 clusters

identified by visual inspection. To visualize average read density for each cluster, we performed

metagene analyses of promoter windows combined with statistics on RegC locations for each

cluster (**Fig 1b,d,f,g, Sup. Fig 2** in **S1 File**). For metagene analyses, each 6 kb input window

was divided into 101 bins of ~59 bp. For each bin, ATAC-seq read counts were determined,

normalized to sequencing depth, normalized to sum up to 1 for each input window, averaged

across all windows in each cluster and then averaged across replicates. In addition, we show

which fraction of windows have an m- (red) or i-RegC (blue) for each bin.  Based on the location

heatmap and the metagene analyses, we identified three major patterns of changes in chromatin

accessibility around promoters during WT infection: (i) ATAC-seq peaks at the TSS that shift

and/or broaden into regions downstream of the TSS during HSV-1 infection (clusters 1, 2, 4, 5,

9, 10, 12, 14, a total of 3,472 genes, **Fig 1b, Sup. Fig 2a-g** in **S1 File,** examples in **Fig 1c, Sup.**

**Fig 3a-h** in **S1 File**), (ii) ATAC-seq peaks at the TSS that shift and/or broaden into regions

upstream of the TSS (clusters 6, 7, 11, a total of 919 genes, **Fig 1d, Sup. Fig 2h,i** in **S1 File,**

examples in **Fig 1e, Sup. Fig 3i-k** in **S1 File**), and (iii) an increase in chromatin accessibility

both up- and downstream of the TSS peak in infection compared to mock (cluster 13, 126 genes,

**Fig 1f,** example in **Sup. Fig 3l** in **S1 File**). Two clusters (3 and 8, a total of 464 genes) exhibited

a combination of patterns (i) and (ii) with an extension of the TSS peak in both up- and

downstream direction (**Fig 1g, Sup. Fig 2j, Sup. Fig 3m,n** in **S1 File**).

**Most chromatin accessibility changes at promoters are independent of dOCR induction**

First, we evaluated which of the observed changes were due to dOCRs extending into promoter

regions. For this purpose, we determined the fraction of promoter windows in each cluster that

overlapped with dOCRs of the 1,296 genes for which we previously showed consistent dOCR

induction across different HSV-1 strains (29) (**Fig 2a**). dOCR regions in mock, WT and all null

mutant infections (average across two replicates) were identified as previously described (21,

29) (see also Materials and Methods). It is important to note here that some relatively short

dOCRs are already observed in mock infection but these substantially extend during HSV-1

infection for read-through regions. In this analysis, we also included ATAC-seq data obtained

in the same experiment for 8 h p.i. WT infection combined with PAA treatment (WT+PAA) to

identify dOCR regions that are not as clearly detectable without PAA treatment but may still

bias promoter analyses. WT (± PAA) and null mutant infections are ordered according to the

overall extent of dOCR induction in **Fig 2a**. While no differences to mock infection were

observed for ΔICP22 infection as expected, some clusters showed an enrichment for dOCR

overlaps in other virus infections. This increased with the overall extent of dOCR induction in

these viruses. However, no cluster had more than 20% of promoter windows overlapping with

dOCRs even with PAA treatment and most had <5% overlap, indicating that dOCRs represent

only a very small fraction of observed changes in chromatin accessibility around promoters.

Most importantly, the extension of accessible chromatin regions at promoters was also observed

upon infection with an ICP22-null mutant but strongly reduced by PAA treatment indicating

that the extension is not linked to dOCR formation (**Fig 2b,c**, **Sup. Fig 4-17** in **S1 File**). Even

for clusters 12-14, which exhibited some enrichment for dOCR overlaps, most differential

regions were also detected in ΔICP22 infection (**Sup. Fig 15-17** in **S1 File**) and thus not linked

to dOCR induction. This was also confirmed in the independent ATAC-seq experiment for 8

and 12 h infection with WT-F and its ΔICP22 mutant (**Sup. Fig 18**). As absence of ICP0, ICP27

or *vhs* also had little impact on changes in chromatin accessibility around promoters (**Sup. Fig**

**4-17** in **S1 File**), we conclude that neither dOCR induction nor activity of ICP0, ICP22, ICP27

or *vhs* alone explain the majority of observed changes in chromatin accessibility at promoters

in HSV-1 infection.

**Extension of accessible chromatin regions around promoters is linked to transcription**

To correlate observed changes to potential transcription factor binding, we next performed

motif discovery for novel and known motifs in m- and i-RegC grouped either by cluster or

pattern compared to the background of all promoter windows using HOMER (39), but found

no enriched motifs. Functional enrichment analysis for Gene Ontology (GO) terms for each

cluster also yielded no significant results, with the exception of cluster 13 (pattern (iii)). Cluster

13 was enriched for "mRNA splicing via the spliceosome" and related terms, however, this was

mainly due to several snRNA genes included in this cluster. Accordingly, cluster 13 was

significantly enriched for snRNA genes (**Fig 2d**, adj. p. $= 3 \times 10^{-16}$, see Materials and

Methods). In contrast, pattern (i) clusters tended to contain high fractions of protein-coding

genes, but no or very few snRNAs. Manual investigation of these few snRNAs either showed

no shift or an overlap with other protein-coding genes. Thus, most snRNAs either showed no

significant changes or a relative increase in chromatin accessibility on both sides of the TSS (=

pattern (iii) in cluster 13), but no down- or upstream shifts in chromatin accessibility.  Notably,

snRNAs are transcribed by RNA Polymerase III (Pol III) and not Pol II.  However, snRNA loci

as well as other genes not transcribed by Pol II (rRNAs, tRNAs) are repeated several times in

the human genome, thus quality of read mappings to these loci is not sufficient to draw a

definitive conclusion. Consequently, few snRNA (55), almost no rRNA (12) and no tRNA loci

were included in our analysis. In summary, this analysis provides clear evidence for an

extension of accessible chromatin at promoters of a large number of genes transcribed by Pol II,

in particular protein-coding genes.

Strikingly, cluster 7 (= the most pronounced case of pattern (ii)) was significantly enriched for

antisense transcripts (adj. p. $= 5 \times 10^{-15}$). The other pattern (ii) clusters 6 and 11 as well as

cluster 8 (combined pattern (i) + (ii)) also contained a relatively high fraction of antisense

transcripts, although this was not statistically significant. Consistently, we found that cluster 7

(adj. p. $= 3.8 \times 10^{-14}$) and to a lesser degree clusters 6 and 11 (adj. p. $= 0.006$ and $0.041$,

respectively) were enriched for bidirectional promoters, i.e. promoters containing an annotated

gene start within 1 kb upstream of the TSS of the target gene (= the gene around whose TSS

the promoter window was originally defined, **Fig 2e,** see **Fig 1e** for an example). We thus

hypothesized that pattern (ii) essentially represented the mirror image of pattern (i) (mirrored

at a vertical axis through the TSS) with the "downstream" broadening/shift of chromatin

accessibility occurring in antisense direction for bidirectional promoters. To confirm this, we

calculated the ratio of sense to antisense transcriptional activity in promoter windows for all

genes. For this purpose, we analyzed RNA-seq data of chromatin-associated RNA, which

depicts nascent transcription, for mock and 8 h p.i. WT HSV-1 infection from our recent study

(19). We also included in this analysis the 2,668 genes (denoted as NA group) without

significant chromatin accessibility changes that were excluded from **Fig 1a**. Interestingly,

metagene analyses for these genes also revealed a slight broadening of the TSS peak into

downstream regions, although this was much less pronounced than for pattern (i) clusters and

therefore not detected by RegCFinder (**Sup. Fig 19a** in **S1 File**). To calculate the ratio of sense

to antisense transcription, promoter windows were divided into the regions down- and upstream

of the TSS and expression in chromatin-associated RNA was determined in sense direction for

the downstream region (=DSR) and in antisense direction for the upstream region (=UAR). We

then compared log2 ratios of DSR to UAR expression between clusters (**Fig 3a**). Here, positive

log2(DSR:UAR) ratios indicate that sense transcription downstream of the TSS is stronger than

antisense transcription upstream of the TSS and negative values indicate the opposite.

Strikingly, all pattern (ii) clusters had significantly lower DSR to UAR ratios than the remaining

genes, while pattern (i) tended to have significantly higher ratios. While for pattern (ii) clusters

6 and 11 median values were still positive, values for cluster 7 with the most pronounced pattern

(ii) were commonly negative. Thus, transcription upstream of the TSS in antisense direction

was the dominant mode of transcription for cluster 7 promoters, which means pattern (ii)

essentially just represents pattern (i) for the genes on the antisense strand.

Next, we investigated differences in gene expression changes in WT compared to mock

infection between patterns (**Sup. Fig 19b** in **S1 File**). This showed a significant difference (adj.

$p < 0.001$) for cluster 1, for which gene expression was more strongly down-regulated compared

to all other genes, but otherwise no consistent trends between patterns. In contrast, gene

expression (quantified as FPKM = fragments per kilobase million mapped reads, **Fig 3b**)

differed considerably between patterns. Most pattern (i) clusters as well as cluster 3 (combined

pattern (i)+(ii)) exhibited median expression levels above average in mock (**Fig 3b**), which was

statistically significant for 5 clusters. In contrast, clusters 6, 7 (pattern (ii)) and 8 (combined

pattern (i)+(ii)) exhibited relatively low expression values, with cluster 7 genes having

significantly lower expression. The latter is consistent with these genes being less expressed

than their antisense counterpart in these bidirectional promoters. Genes without significant

chromatin accessibility changes (NA group) also showed significantly lower expression, albeit

not as low as cluster 7. These differences between clusters/patterns were generally maintained

in HSV-1 infection at lower overall expression levels, consistent with the absence of differences

in gene expression fold-changes between most clusters. When stratifying analyzed genes into

five equal-sized groups according to FPKM in chromatin-associated RNA in uninfected cells,

we observed that the fraction of genes with at least one significant RegC increased with gene

expression (**Fig. 3c**). Consistently, pattern (ii) was significantly enriched among the lowliest

expressed genes and depleted among genes with medium to high expression (**Fig. 3d**). In

contrast, cluster 5, 9 and 10 with pattern (i) were enriched among the most highly expressed

genes. Metagene analyses on the five groups of gene expression confirm this trend, with the

downstream extension of chromatin accessibility increasing with gene expression (**Fig. 3e-g**).

Interestingly, pattern (i) cluster 1, which showed a significantly higher reduction in gene

expression during HSV-1 infection than the other analyzed genes, was most frequent among

genes with low to high expression, but weakly significantly depleted among the most highly

expressed genes. In summary, these results indicate that HSV-1 infection leads to an extension

of the accessible chromatin around promoters in the dominant direction of transcription for the

majority of host genes. Here, highly expressed genes and moderately expressed genes with

stronger transcription reduction in HSV-1 infection are most strongly affected. Notably,

although highly expressed genes do not generally exhibit a stronger reduction of transcription

*relative* to their original expression levels in mock, the *absolute* drop in Pol II occupancy

between mock and HSV-1 infection is more pronounced than for more lowly expressed genes.

Thus, our results show parallels to observations for yeast that Pol II depletion induces

downstream shifts of +1 nucleosomes, which extends nucleosome-free regions and accordingly

accessible chromatin at promoters, most prominently for highly expressed genes (30). We thus

hypothesized that the loss of Pol II during HSV-1 infection causes the extension of accessible

chromatin at promoters during HSV-1 infection.

**Changes in chromatin accessibility manifest between 4 and 6 h of HSV-1 infection**

To investigate how early in infection changes in chromatin accessibility around promoters can be detected, we ran RegCFinder on an ATAC-seq time-course for 1, 2, 4, 6, and 8 h p.i. WT infection from our previous publication (21) (**Fig 4a,b**). While barely any significant differential regions were identified at 1 and 2 h p.i., 1,963 significant regions in 1,249 promoter windows were identified by 4 h p.i. and 7,558 significant regions in 3,989 promoter windows by 8 h p.i. (**Fig 4b**). Log2 fold-changes for significant differential regions generally reflected the patterns observed in the analysis for WT and null mutant infections (**Fig 4a**). The same applied when we separately determined fold-changes and significance on the time-course data for the differential regions determined for WT vs. mock from our initial analysis (**Sup. Fig 20 in S1 File**) and when we performed metagene analyses for the clusters from **Fig 1a (Fig 4d-f**, **Sup. Fig 21 in S1 File**, example genes in **Sup. Fig 22 in S1 File)**. Although the changes in chromatin accessibility at 8 h p.i. of the time-course were less pronounced than for the original WT vs. mock comparison at 8 h p.i., this nevertheless confirms the changes in chromatin accessibility around promoters in HSV-1 infection in an independent experiment. The lower fraction of viral reads at 8 h p.i. in the time-course experiment (~25%, **Sup. Table 1 in S2 File**) than for the first WT experiment (~52%) suggests a slower progression of infection in the time-course experiment, which likely explains why the effect was less pronounced.

The time-course analysis also reveals that these changes begin to manifest by 4 h p.i., with 1,249 genes already showing a significant change. Moreover, genes that showed an early effect (by 4 h p.i. or earlier) were significantly more highly expressed than genes that showed an effect by 6 h p.i. or later (**Fig 4c**). The lowest expression was observed for genes with a significant effect in the first WT experiment, but not in the time-course. This again confirms the link between expression levels of a gene and the change in chromatin accessibility. Notably, we previously showed that transcriptional activity on host genes drops substantially in the first 4 h

of infection to only 40% of transcription in uninfected cells, which was further halved until 8 h

p.i. (19). Thus, the onset of the extension of accessible chromatin regions at promoters into

downstream (for pattern (i)) and upstream (for pattern (ii)) regions follows the onset of the drop

in host transcription during infection. This provides further evidence for the hypothesis that loss

of Pol II from host genes drives changes in chromatin accessibility at promoters. Since slower

progression of virus infection in the time-course experiment likely leads to a less pronounced

loss of Pol II from host genes, this would explain the differences in effect between the time-

course and the null mutant experiment.

**ICP4 knockout reduces and combined expression of ICP22 and ICP27 induces the**

**extension of accessible chromatin**

We already showed that PAA treatment substantially reduced the broadening and/or

downstream shift of accessible chromatin regions at promoters (**Fig 2b,c, Sup. Fig 4-17** in **S1**

**File**). As PAA alleviates the depletion of Pol II from host genomes by inhibiting viral DNA

replication, this further corroborates that Pol II depletion leads to the observed shifts in

chromatin accessibility at promoters. Exceptions were cluster 13, and parts of cluster 12 and

14, for which chromatin accessibility increased up- (for clusters 12-14) and downstream (for

cluster 13) of the TSS peak (**Sup. Fig 15-17** in **S1 File**). Since these clusters were enriched for

overlaps with dOCRs from an upstream gene, these changes reflect increased dOCR induction

upon PAA treatment. Notably, however, the broadening of the TSS peak into downstream

regions in clusters 12 and 14 was reduced by PAA treatment.

The original ATAC-seq experiment with null mutant infections also included infection with an

ICP4 null mutant at 8 h p.i., which had not been previously published. Since ICP4 facilitates

recruitment of Pol II to viral replication compartments (13, 14), we now also investigated

changes in chromatin accessibility in ΔICP4 infection. Strikingly, while we still observed the

extension of accessible chromatin in down- (**Fig 5a**, **Sup. Figs 23a-g,k,l** in **S1 File**) or upstream

direction (**Fig 5b**, **Sup Fig. 23h,i** in **S1 File**), it was similarly reduced in ΔICP4 infection

compared to WT infection as upon PAA treatment. Interestingly, the fraction of viral reads in

ATAC-seq experiments were comparably low at 3-4% in both ΔICP4 and WT+PAA infection

(**Sup. Table 1** in **S2 File**), reflecting little viral DNA replication in absence of ICP4. We next

tested for enrichment in promoter regions of ICP4 binding sites previously identified by Dremel

*et al.* for 2 h p.i. HSV-1 infection (40). This showed a significant enrichment (adj. p. =

$1.1 \times 10^{-5}$) of ICP4 binding sites for genes with significant RegC (compared to the NA group

of genes without any significant changes) and among these, a weakly significant enrichment in

clusters 9 (pattern (i), adj. p = 0.0046) and 11 (pattern (ii), adj. p.=0.0018). Notably, the only

cluster with less frequent ICP4 binding than the NA group was cluster 13 (pattern (iii)), which

did not show shifts in chromatin accessibility. Thus, genes with significant shifts in chromatin

accessibility are more frequently bound by ICP4 than genes without shifts, providing evidence

that Pol II depletion from host genomes by ICP4 contributes to this phenomenon.

ICP27 and ICP22 are still expressed and active in absence of ICP4, albeit at lower levels, as

both read-through transcription and dOCRs are observed in ICP4 null mutants (22). Thus,

ICP22 and ICP27 activities still lead to some depletion of Pol II from host gene bodies, which

likely explains the residual changes in chromatin accessibility observed in absence of ICP4. To

confirm that ICP27 and/or ICP22 expression alone can induce such changes, we re-investigated

our recently published Omni-ATAC-seq (a recent improvement of ATAC-seq (41)) data for

telomerase-immortalized human foreskin fibroblasts (T-HFs) that express either ICP22 (T-HF-

ICP22 cells) or ICP27 (T-HF-ICP27 cells) in isolation or in combination (T-HF-ICP22/ICP27

cells) upon doxycycline (dox) exposure. While we did not identify any significant RegC upon

ICP22 expression alone, ICP27 expression alone lead to significant changes for >2,000 genes

(**Fig 5c**, **Sup. Fig 24a,b** in **S1 File**). Combined ICP22 and ICP27 expression led to more

pronounced results, with the same patterns observed as for WT infection but in an attenuated

manner (**Fig 5c-e**, **Fig 24** in **S1 File**). This shows that combined activity of ICP22 and ICP27

is sufficient to induce a moderate extension of chromatin accessibility at promoters. However,

the pronounced changes observed in WT infection require viral DNA replication but neither

ICP22 nor ICP27 expression. Notably, although ICP27 is required for optimal viral DNA

replication (42), knockout of ICP27 does not reduce viral DNA replication as effectively as

ICP4 knockout or PAA treatment as ~23% of ATAC-seq reads have a viral origin in ΔICP27

infection (**Sup. Table 1** in **S2 File**). This is comparable to the 8 h p.i. time-point in the time-

course experiment. We conclude that a substantial depletion of Pol II from host genomes due

to recruitment to viral genomes is required to fully manifest the changes in chromatin

architecture at promoters during HSV-1 infection.

**HSV-1 infection induces a downstream shift of +1 nucleosomes**

The HSV-1-induced changes in chromatin accessibility at promoters indicate a broadening of

the nucleosome-free region around promoters in either sense (pattern (i)) or antisense (pattern

(ii)) direction. The extension of nucleosome-free regions requires shifts in +1 or -1 nucleosome

positions and depletion of Pol II has been shown to induce a downstream shift of +1

nucleosomes in yeast (30). We, thus, performed ChIPmentation in HFF for mock and WT

infection at 8 h p.i. with an antibody recognizing the C-terminal part of the non-canonical

histone H2A.Z (n=3 replicates). H2A.Z is highly enriched at gene promoters at -1 and +1

nucleosome positions (32, 33) and is encoded by two genes, whose protein products H2A.Z.1

and H2A.Z.2 differ by only three amino acids. While their genomic occupancy patterns are

similar, there are quantitative differences with H2A.Z.1 being more abundant at active

promoters than H2A.Z.2 (43). Since the C-terminal regions of H2A.Z.1 and H2A.Z.2 differ

only by the last amino acid, the antibody used for ChIPmentation recognizes both isoforms. A

metagene analysis of all analyzed promoter windows showed the expected distribution of

H2A.Z occupancy in mock infection with two peaks on both sides of the TSS, corresponding

to -1 and +1 nucleosome positions (**Sup. Fig 25a** in **S1 File**).

Application of RegCFinder to H2A.Z ChIPmentation data generally identified a relative

increase in H2A.Z downstream of the TSS during infection and a relative decrease upstream of

the TSS for pattern (i) and combined pattern (i)+(ii) clusters (**Fig 6a**, log2 fold-changes for

differential regions shown in **Sup. Fig 25b** in **S1 File**, examples in **Fig 6b**, **Sup. Fig 26a-g,k,l**

in **S1 File**). Metagene analyses showed that this reflected a downstream shift and broadening

of +1 nucleosome peaks as well as a relative increase of +1 nucleosome peaks compared to -1

nucleosome peaks (**Fig 6d**, **Sup. Fig 25c-i,m,n** in **S1 File**). In contrast, cluster 7 (most

pronounced pattern (ii) cluster) showed the opposite trend with relative increases in H2A.Z

upstream of the TSS and decreases downstream of the TSS (example in **Fig 6c**). Consistent

with pattern (ii) representing pattern (i) in antisense direction, the metagene analysis showed an

upstream shift and broadening of the -1 nucleosome peak and a relative increase of the -1

nucleosome peak compared to the +1 nucleosome peak (**Fig 6e**). However, this was only

observed for few genes in cluster 6 and cluster 11 (**Fig 6a**, **Sup. Fig 25j,k** in **S1 File**, examples

in **Sup. Fig 26h,i** in **S1 File**), which fits with the observation that pattern (ii) in the ATAC-seq

data was also much less pronounced for these clusters. In contrast to pattern (i) and (ii), pattern

(iii) in the ATAC-seq data (cluster 13) was not associated with distribution changes in H2A.Z

(**Sup. Fig 25l**, example in **Sup. Fig 26j** in **S1 File**), thus it is not shaped by shifts in +1 and/or

-1 nucleosome positioning. This is consistent with pattern (iii) being at least partly associated

with dOCR induction for upstream genes, which we previously linked to impaired nucleosome

repositioning following Pol II transcription downstream of genes (29).

In summary, our data confirm that the broadening of chromatin accessibility around promoters

indeed results from down- and upstream shifts of +1 or -1 nucleosomes, respectively. Here, the

direction of shift depends on whether sense or antisense transcription represents the dominant

direction of transcription at this promoter. Notably, a recent study in yeast proposed that -1

nucleosomes should actually be considered +1 nucleosomes for antisense transcription (44).

This strongly suggests that the loss of Pol II from host promoters during HSV-1 infection

induces a shift of +1 and -1 nucleosome positions similar to what was previously observed for

yeast. To confirm that inhibition of transcription alone can lead to shifts of +1 and -1

nucleosomes in human cells, we re-analyzed published H2A.Z ChIP-seq data of HCT116 cells

with and without α-amanitin treatment from the recent study by Lashgari *et al*. (45). They

showed that α-amanitin treatment leads to reduced Pol II levels at gene TSSs and increased

incorporation of H2A.Z at TSSs of transcribed genes. Here, Pol II ChIP and RT-qPCR of

control genes showed that α-amanitin treatment reduced Pol II signal at the TSS to 10-20% of

untreated cells and transcriptional activity to less than 40%. This is not much lower than what

has previously been reported for HSV-1 infection: Abrisch *et al.* found that Pol II occupancy

on gene bodies upon 4 h HSV-1 infection was reduced on average to around 20% (9) and we

previously estimated that transcriptional activity was reduced to 20-40% between 4 and 8 h p.i.

(19). Notably, in contrast to DRB and flavopiridol, which inhibit transcription via arresting Pol

II in a paused state at the promoter,  α-amanitin actually induces a loss of Pol II from the host

genome, thus making it a better model of HSV-1-induced Pol II depletion.

Lashgari *et al*. previously only analyzed changes in absolute H2A.Z levels at promoters and did

not focus on changes in +1 and -1 nucleosome positioning. Metagene and RegCFinder analyses

of the H2A.Z ChIP-seq data indeed showed a similar trend upon α-amanitin treatment as during

HSV-1 infection (**Fig 6f,g, Sup. Fig 27,** examples are shown in **Sup. Fig 28** in **S1 File**). Pattern

(i) and pattern (i)+(ii) clusters showed a massive broadening of the +1 nucleosome peak into

downstream regions, even more pronounced than what is observed upon HSV-1 infection. In

addition, they also exhibited a (less pronounced) broadening of -1 nucleosome peaks. Similarly,

pattern (ii) clusters showed a strong broadening of both -1 and +1 nucleosome peaks into

upstream regions. In case of cluster 7 (strongest pattern (ii)), the broadening was much more

pronounced for the -1 nucleosome (i.e. the +1 nucleosome in antisense direction) than for the

+1 nucleosome (i.e. the -1 nucleosome in antisense direction). This analysis confirms that

depletion of Pol II from promoters leads to a downstream shift of +1 nucleosomes in human

cells. In summary, our results provide strong evidence that the shift of +1 and -1 nucleosomes

in the direction of transcription in HSV-1 infection is a consequence of Poll II depletion from

host chromatin.

## Discussion

Previously, we showed that HSV-1 infection disrupts chromatin architecture downstream of

genes with strong read-through transcription. In this study, we reveal that chromatin

architecture is also substantially altered at host gene promoters during HSV-1 infection. Here,

57% of genes showed a statistically significant change in the distribution of chromatin

accessibility at promoters in WT HSV-1 infection and metagene analyses indicated similar but

less pronounced changes even for genes without statistically significant changes. In essence,

we identified three types of changes: Most genes showed a shift and/or broadening of accessible

chromatin into regions downstream of the TSS (pattern (i)), while ~900 genes showed the

opposite trend with a shift and/or broadening of accessible chromatin into regions upstream of

the TSS (pattern (ii)). A further ~460 genes exhibited a combination of pattern (i) and (ii), with

the broadening of accessible chromatin in downstream direction being more pronounced than

in upstream direction.  Only a small set of genes (126 genes) showed relative increases in

chromatin accessibility up- and downstream of the TSS, which could partly be attributed to

overlaps with dOCRs from upstream genes and partly to an enrichment for short non-coding

RNAs not transcribed by Pol II, for which promoter windows were longer than the actual gene.

We thus did not further investigate this pattern.

As patterns (i) and (ii) were still observed in absence of ICP22, which is necessary for dOCR

induction, these are not artefacts of dOCRs extending into downstream genes. On the contrary,

PAA treatment, which increases dOCR induction, substantially reduced – though not

completely abolished – pattern (i) and (ii). Furthermore, knockout of neither ICP0, ICP22,

ICP27 nor *vhs* substantially affected pattern (i) and (ii). In contrast, ICP4 knockout, similar to

PAA treatment, also alleviated the down- and upstream broadening of chromatin accessibility.

Both PAA treatment and ICP4 knockout largely abolish viral DNA replication and

consequently alleviate the depletion of Pol II from host genes, leading us to hypothesize that

this HSV-1-induced loss of Pol II causes the widespread extension of open chromatin at host

gene promoters. This is further supported by the observation that combined dox-induced

expression of ICP22 and ICP27 (and to a lesser degree dox-induced ICP27 expression alone)

led to some broadening of chromatin accessibility at promoters despite knockdown of neither

of these proteins having an effect on the HSV-1-induced changes. Knockout of ICP22 or even

ICP27 does not sufficiently abolish viral DNA replication and thus has only a minor effect on

the depletion of Pol II from host genomes and consequently on chromatin accessibility changes.

In contrast, in absence of viral DNA replication, activities of ICP22 and ICP27 (and potentially

other viral factors) sufficiently reduce Pol II levels on gene bodies to induce an alleviated form

of those chromatin accessibility changes. Moreover, both the relatively late onset of chromatin

accessibility changes between 4 and 8 h p.i. in our time-course ATAC-seq analysis and the

generally reduced levels of changes in the time-course, which had a lower fraction of viral reads

at 8 h p.i., confirm that strong depletion of Pol II is necessary to observe substantial effects on

chromatin accessibility.  This also highlights that early (at ~2 h p.i.) ICP4-mediated depletion

of Pol II from host promoters before the onset of genome replication does not have a detectable

effect.

Analysis of transcriptional activity using RNA-seq of chromatin-associated RNA revealed that

pattern (i) and (ii) are indeed linked to transcription, with pattern (i) associated with more highly

expressed genes and pattern (ii) associated with bidirectional promoters with strong antisense

transcription on the opposite strand. In case of cluster 7, for which pattern (ii) was most

pronounced, antisense transcription was indeed much higher than sense transcription. This was

not due to the widespread induction of antisense transcription in HSV-1 infection, which we

previously reported on (46), but these genes already exhibited strong transcription in antisense

direction in uninfected cells originating from bidirectional promoters. Consistently, cluster 7

was enriched for genes annotated as antisense and more than a third of these promoters

contained an annotated gene starting on the opposite strand within 1 kb upstream of the TSS of

the target gene. Thus, pattern (ii) is essentially just pattern (i) for the genes on the opposite

strand in bidirectional promoters.

Since the broadening of chromatin accessibility in promoter regions suggested an extension of

nucleosome-free regions at promoters, we mapped +1 and -1 nucleosome positions by

ChIPmentation of the H2A.Z histone variant enriched at these nucleosomes. Analysis of H2A.Z

occupancy indeed showed a downstream shift of +1 nucleosomes for pattern (i) genes in HSV-1

infection and an upstream shift of -1 nucleosomes for genes with most pronounced pattern (ii)

(cluster 7). This further confirmed that pattern (ii) just represents pattern (i) for genes on the

antisense strand. It is also consistent with recent results from yeast by Bagchi *et al.* that -1

H2A.Z-containing nucleosomes should actually be considered as +1 nucleosomes for antisense

transcription (44). Downstream shifts of +1 nucleosomes have previously been reported upon

Pol II degradation in yeast by Weiner *et al.* (30), in particular for highly expressed genes. Re-

analysis of published H2A.Z ChIP-seq data with and without α-amanitin-induced depletion of

Pol II from gene promoters confirmed that depletion of Pol II alone is sufficient to induce

downstream shifts of +1 nucleosomes in human cells. Moreover, for genes with dominant

antisense transcription (cluster 7), it resulted in upstream shifts of -1 nucleosomes. To our

knowledge, this is the first study that confirms the effect of Pol II depletion on nucleosome

positioning for human cells. The broadening of H2A.Z peaks in both HSV-1 infection and upon

α-amanitin treatment, respectively, indicate a less precise positioning of corresponding

nucleosomes. This was more pronounced upon α-amanitin treatment and also observed for -1

nucleosomes, suggesting that loss of Pol II upon α-amanitin treatment is more pronounced than

in HSV-1 infection.

One open question remaining concerns the biological significance of the observed changes in

nucleosome positioning. Our analysis of gene expression changes did not suggest an effect on

differential gene expression, rather the opposite with strong down-regulation leading to the

same effect for less strongly expressed genes (cluster 1) as for highly expressed genes that are

not more down-regulated than other genes. However, the downstream shifts in +1 nucleosomes

may provide an explanation for the downstream shift of Pol II pausing we recently reported for

HSV-1 infection (38). The +1 nucleosome was shown to play a role in promoter-proximal

Pol II pausing between the promoter and the +1 nucleosome (47). In particular, the +1

nucleosome represents a $2^{nd}$ barrier to Pol II pause release independent of the main pausing

factor negative elongation factor (NELF). Upon NELF depletion, Pol II stops at a $2^{nd}$ pausing

region around the +1 nucleosomal dyad-associated region (48). We previously observed that

Pol II pausing in HSV-1 infection is shifted to more downstream and less well-positioned sites

for the majority of genes (38), consistent with +1 nucleosome positioning also appearing less

well-positioned upon HSV-1 infection in our H2A.Z ChIPmentation data. Alternatively, the

shifts in +1 nucleosomes might also be linked to the mobilization of H1, H2 (including H2A.Z)

and H4 histones during HSV-1 infection (49-51), which has been proposed to serve as a source

of histones for viral chromatin assembly. Nevertheless, even if the changes in nucleosome

positioning upon Pol II depletion are of no further functional consequence for HSV-1 infection,

they are highly relevant for any functional genomics studies on chromatin architecture in

HSV-1 infection or other conditions that deplete Pol II from the genome. If not properly taken

into account, the broadening of accessible chromatin at promoters may be mistaken e.g. for

differential transcription factor binding, which could lead to wrong conclusions. Together with

read-through transcription and dOCRs extending into downstream genes, this represents one

more example of how HSV-1 infection confounds standard functional genomics analyses. In

any case, our study highlights that HSV-1 infection impacts chromatin architecture at promoters

independently of the widespread changes downstream of genes mediated by ICP22.

**Materials and Methods**

*Previously published sequencing data analyzed in this study*

ATAC-seq for mock and WT HSV-1 infection (strain 17, 8 h p.i.), WT infection with 8 h PAA

treatment, infection with ICP0-null mutant (ΔICP0, strain 17, (52), 12 h p.i.), ICP22-null mutant

(ΔICP22, R325, strain F, (53), 12 h p.i.), ICP27-null mutant (ΔICP27, strain KOS, (54), 8 h

p.i.) and *vhs*-null mutant (Δvhs, strain 17, (55)) in of HFF were taken from our recent

publication (29) (n=2 apart from ΔICP22 infection with n=4, GEO accession: GSE185234).

This experiment also included infection with an ICP4-null mutant (ΔICP4, n12, strain KOS,

(56), 8 h p.i.) which had not previously been published (GEO accession: GSE212820). ATAC-

seq data for mock and WT HSV-1 infection of HFF at 1, 2, 4, 6 and 8 h p.i. WT infection (n=2

replicates, GEO accession: GSE100611) and chromatin-associated RNA-seq data for mock and

8 h p.i. WT infection (GEO accession: GSE100576) were taken from our previous publication

(21). H2A.Z ChIP-seq data of untreated and α-amanitin-treated HCT116 cells were taken from

the study by Lashgari *et al*. (45) (n=2, GEO accession: GSE101427).

*ChIPmentation, library preparation and sequencing*

HFF were purchased from ECACC and cultured in Dulbecco's Modified Eagle Medium

(DMEM, ThermoFisher #41966052) supplemented with 10% (v/v) Fetal Bovine Serum (FBS,

Biochrom #S0115), 1× MEM Non-Essential Amino Acids (ThermoFisher #11140050) and 1%

penicillin/streptomycin. Two days prior to infection, two million HFF cells were seeded in 15

cm dishes. At the day of infection, cells had expanded to ~80% confluency. Cells were infected

with the respective viruses as described in results section (n=3). At 8 p.i., cells were fixed for

10 minutes at room temperature by adding 1% formaldehyde (final) directly to the medium.

Cells were scraped in 1mL of ice-cold 1x PBS containing protease inhibitor cocktail (1x)

(Roche #11836153001) with an additional 1mM phenylmethylsulfonyl fluoride (PMSF). Cells

were pelleted at 1500 rpm for 20 min at 4 °C. Supernatant was aspirated and cell pellets were

frozen in liquid $N_2$.

Cell pellets were resuspended in 1.5 mL 0.25% [w/v] SDS sonication buffer (10 mM Tris

pH=8.0, 0.25% [w/v] SDS, 2 mM EDTA) with 1x protease inhibitors and 1 mM additional

PMSF and incubated on ice for 10 minutes. Cells were sonicated in fifteen 1 minute intervals,

25% amplitude, with Branson Ultrasonics SonifierTM S-450 until most fragments were in the

range of 200-700 bp as determined by agarose gel electrophoresis. Two million cells used for

the preparation of the ChIPmentation libraries were diluted 1:1.5 with equilibration buffer (10

mM Tris, 233 mM NaCl, 1.66% [v/v] Triton X-100, 0.166% [w/v] sodium deoxycholate, 1 mM

EDTA, protease inhibitors) and spun at 14,000x g for 10 minutes at 4 °C to pellet insoluble

material. Supernatant was transferred to a new 1.5 mL screw-cap tube and topped up with

RIPA-LS (10 mM Tris-HCl pH 8.0, 140 mM NaCl, 1 mM EDTA pH 8.0, 0.1% [w/v] SDS,

0.1% [w/v] sodium deoxycholate, 1% [v/v] Triton X-100, protease inhibitors) to 200 μL. Input

and gel samples were preserved. Lysates were incubated with 1μg/IP of anti-H2A.Z antibody

(Diagenode, #C15410201) on a rotator overnight at 4 °C.

Dependent on the added amount of antibody, the amount of Protein A magnetic beads

(ThermoFisher Scientific #10001D) was adjusted (e.g. for 1-2 μg of antibody/IP = 15 μL of

beads) and blocked overnight with 0.1% [w/v] bovine serum albumin in RIPA buffer. On the

following day, beads were added to the IP samples for 2 h on a rotator at 4 °C to capture the

antibody-bound fragments. The immunoprecipitated chromatin was subsequently washed twice

with 150 μL each of ice-cold buffers RIPA-LS, RIPA-HS (10 mM Tris-HCl pH 8.0, 50 0mM

NaCl, 1 mM EDTA pH 8.0, 0.1% [w/v] SDS, 0.1% [v/v] sodium deoxycholate, 1% [v/v] Triton

X-100), RIPA-LiCl (10 mM Tris-HCl pH 8.0, 250 mM LiCl, 1 mM EDTA pH 8.0, 0.5% [w/v]

sodium deoxycholate, 0.5% [v/v] Nonidet P-40) and 10 mM Tris pH 8.0 containing protease

inhibitors. Beads were washed once more with ice-cold 10 mM Tris pH 8.0 lacking inhibitors

and transferred into new tubes.

Beads were resuspended in 25 μL of the tagmentation reaction mix (Nextera DNA Sample Prep

Kit, Illumina) containing 5 μL of 5x Tagmentation buffer, 1 μL of Tagment DNA enzyme,

topped up with $H_2O$ to the final volume and incubated at 37 °C for 10 minutes in a thermocycler.

Beads were mixed after 5 minutes by gentle pipetting. To inactivate the Tn5 enzyme, 150 μL

of ice-cold RIPA-LS was added to the tagmentation reaction. Beads were washed twice with

150 μL of RIPA-LS and 1x Tris-EDTA and subjected to de-crosslinking by adding 100 μL

ChIPmentation elution buffer (160 mM NaCl, 40 μg/mL Rnase A (Sigma-Aldrich #R4642), 1x

Tris-EDTA (Sigma #T9285) and incubating for 1h at 37 °C followed by overnight shaking at

65 °C. The next day, 4 mM EDTA and 200 μg/mL Proteinase K (Roche, #03115828001) were

added, and samples incubated for another 2h at 45 °C with 1000 rpm shaking. Supernatant was

transferred into a new tube and another 100 μL of ChIPmentation elution buffer was added for

another hour at 45 °C with 1000 rpm shaking. DNA was isolated with MinElute PCR

Purification Kit (Qiagen #28004) and eluted in 21 μL of $H_2O$.

DNA for the final library was prepared with 25 μL NEBNext Ultra II Q5 Master Mix, 3.75 μL

IDT custom primer i5_n_x (10 μM); 3.75 μL IDT custom primer i7_n_x (10 μM); 3.75 μL $H_2O$

and 13.75 μL ChIPmentation DNA. The Cq value obtained from the library quantification,

rounded up to the nearest integer plus one additional cycle, was used to amplify the rest of the

ChIPmentation DNA. Library qualities were verified by High Sensitivity DNA Analysis on the

Bioanalyzer 2100 (Agilent) before performing sequencing on NextSeq 500 (paired-end 35bp

reads) at the Core Unit Systemmedizin, Würzburg, Germany. All samples were sequenced at

equimolar ratios.

*Read alignment*

The read alignment pipeline was implemented and run in the workflow management system

Watchdog (57, 58) as already previously described (38). Public sequencing data were

downloaded from SRA using the sratoolkit version 2.10.8. Sequencing reads were aligned

against the human genome (GRCh37/hg19), the HSV-1 genome (Human herpesvirus 1 strain

17, GenBank accession code: JN555585) and human rRNA sequences using ContextMap2

version 2.7.9 (59) (using BWA as short read aligner (60) and allowing a maximum indel size

of 3 and at most 5 mismatches). For the two repeat regions in the HSV-1 genome, only one

copy was retained each, excluding nucleotides 1–9,213 and 145,590–152,222 from the

alignment. SAM output files of ContextMap2 were converted to BAM files using samtools

(61). Read coverage in bedGraph format was calculated from BAM files using BEDTools (62).

Subregions of promoter windows (TSS ± 3 kb) with differential read coverage in ATAC-seq

and H2A.Z ChIPmentation/-seq data were determined with RegCFinder (36). For this purpose,

RegCFinder was applied to ATAC-seq data for all pairwise comparisons of mock to WT and

null mutant infections as well as mock to each timepoint of infection for the time-course data

and to H2A.Z ChIPmentation and ChIP-seq data for the comparison of mock and WT infection

as well as untreated and α-amanitin-treated HCT116 cells.

*Quality control*

Statistics on numbers of mapped reads and reads mapped to human and HSV-1 genomes were

determined with samtools (65). Promoter/Transcript body (PT) scores were determined with

ATACseqQC (66). For peak calling, BAM files with mapped reads were converted to BED

format using BEDvTools (67) and peaks were determined from these BED files using F-Seq

with default parameters (68). Fraction of reads in peaks (FRiP) was calculated with

featureCounts (69) using identified peaks as annotation. Annotation of peaks relative to genes

was performed using ChIPseeker (70).

*Data plotting and statistical analysis*

All figures were created in R and all statistical analyses were performed in R (63). Read

coverage plots were created using the R Bioconductor package Gviz (64).

*Metagene and clustering analysis*

Metagene analyses were performed as previously described (65) using the R program developed

for this previous publication (available with the Watchdog *binGenome* module in the Watchdog

module repository (https://github.com/watchdog-wms/watchdog-wms-modules/)). For

promoter region analyses, the regions −3 kb to +3 kb of the TSS were divided into 101 equal-

sized bins for each gene. For each bin, the average coverage per genome position was calculated

and bin read coverages were then normalized by dividing by the total sum of all bins. Metagene

curves for each replicate were created by averaging results for corresponding bins across all

genes in a cluster/group and then averaged across replicates. For hierarchical clustering

analysis, RegCFinder profiles of differential regions were calculated for each promoter window

and comparison by setting each position in an m-RegC to 1, in an i-RegC to -1 and all other

positions to 0.  RegCFinder profiles of each promoter window for each comparison were

concatenated into one row in the matrix. Hierarchical clustering of the resulting matrix was then

performed using the hclust function in R according to Euclidean distances and Ward's

clustering criterion.

*Analysis of downstream open chromatin regions (dOCRs)*

Open chromatin regions (OCRs) were determined from ATAC-seq data by first converting

BAM files with mapped reads to BED format using BEDTools (66) and then determining

enriched regions from these BED files using F-Seq with default parameters (67). dOCRs for

individual genes were calculated from OCRs as previously described (21, 29). In brief, dOCRs

are determined for genes by first assigning all OCRs overlapping with the 10 kb downstream

of a gene to this gene. Second, OCRs starting at most 5 kb downstream of the so far most

downstream OCR of a gene are also assigned to this gene. In both steps, individual OCRs can

be assigned to multiple genes. The second step is iterated until no more OCRs can be assigned.

The dOCR of a gene is then defined as the region from the gene 3'end to the end of the most

downstream OCR assigned to this gene.

*Gene expression analysis*

Number of fragments (=read pairs) per gene were determined from mapped paired-end RNA-

seq reads in a strand-specific manner using featureCounts (68) and gene annotations from

Ensembl (version 87 for GRCh37). For genes, all fragments overlapping exonic regions on the

corresponding strand by $\geq$ 25bp were counted for the corresponding gene. Fold-changes in gene

expression and statistical significance of changes were determined using DESeq2 (69) and p-

values were adjusted for multiple testing using the method by Benjamini and Hochberg (70).

Gene expression was quantified in terms of fragments per kilobase of exons per million mapped

reads (FPKM). Only reads mapped to the human genome were counted for the total number of

mapped reads for FPKM calculation.

*Motif discovery and enrichment analysis*

Motif discovery was performed for each cluster separately for the i- and m-RegC regions using

findMotifsGenome.pl script of the Homer suite (39), with our 7,649 input promoter windows

as background. For this purpose, we used the hg19 annotation provided by Homer and

automated trimming of input windows was disabled. Significant motifs were identified at a

q-value (=False Discovery Rate (FDR) calculated with the Benjamini-Hochberg method (70))

cutoff of 0.01. Over-representation of Gene Ontology (GO) terms was performed separately for

each cluster using the g:Profiler webserver (71) and the R package gprofiler2 (72), which

provides an R interface to the webserver. As background gene list, the genes corresponding to

our 7,649 input promoter windows were provided. P-values were corrected for multiple testing

using the Benjamini-Hochberg method (70) and significantly over-represented GO terms were

identified at an multiple testing adjusted p-value cutoff of 0.001. Enrichment of gene types

(obtained from the Ensembl annotation (version 87 for GRCh37)) within clusters was

determined using one-sided Fisher's exact tests (with alternative = greater). Enrichment (odds-

ratio >1) or depletion (odds-ratio < 1) of ICP4 binding sites from the study of Dremel *et al.* (40)

within clusters as well as enrichment and depletion of clusters within gene groups with different

expression levels were determined with two-sided Fisher's exact tests. P-values were always

corrected for multiple testing using the Benjamini-Hochberg method.

**References**

1. Roizman B KD, RJ W. Herpes Simplex Viruses. In: Knipe DM HP, editor. Fields

Virology. 5th. Philadelphia: Lippincott Williams & Wilkins; 2007. p. 2501-601.

2. Pellet PE RB. Herpesviridae. In: Knipe DM HP, editor. Fields Virology. 6th.

Philadelphia: Wolters Kluwer Health/Lippincott Williams & Wilkins; 2013. p. 1802-2128.

3. Ablashi D, Agut H, Alvarez-Lafuente R, Clark DA, Dewhurst S, DiLuca D, et al.

Classification of HHV-6A and HHV-6B as distinct viruses. Arch Virol. 2014;159(5):863-70.

4. Korr G, Thamm M, Czogiel I, Poethko-Mueller C, Bremer V, Jansen K. Decreasing

seroprevalence of herpes simplex virus type 1 and type 2 in Germany leaves many people

susceptible to genital infection: time to raise awareness and enhance control. BMC Infect Dis.

2017;17(1):471.

5. Whitley RJ, Roizman B. Herpes simplex virus infections. Lancet.

2001;357(9267):1513-8.

6. Whisnant AW, Jürges CS, Hennig T, Wyler E, Prusty B, Rutkowski AJ, et al. Integrative

functional genomics decodes herpes simplex virus 1. Nat Commun. 2020;11(1):2038.

7. Spencer CA, Dahmus ME, Rice SA. Repression of host RNA polymerase II

transcription by herpes simplex virus type 1. J Virol. 1997;71(3):2031-40.

8. Rice SA, Davido DJ. HSV-1 ICP22: hijacking host nuclear functions to enhance viral

infection. Future Microbiol. 2013;8(3):311-21.

9. Abrisch RG, Eidem TM, Yakovchuk P, Kugel JF, Goodrich JA. Infection by Herpes

Simplex Virus 1 Causes Near-Complete Loss of RNA Polymerase II Occupancy on the Host

Cell Genome. J Virol. 2015;90(5):2503-13.

10. Dembowski JA, DeLuca NA. Selective recruitment of nuclear factors to productively

replicating herpes simplex virus genomes. PLoS Pathog. 2015;11(5):e1004939.

11.   Birkenheuer CH, Danko CG, Baines JD. Herpes Simplex Virus 1 Dramatically Alters

Loading and Positioning of RNA Polymerase II on Host Genes Early in Infection. J Virol.

2018;92(8).

12.   Rivas HG, Schmaling SK, Gaglia MM. Shutoff of Host Gene Expression in Influenza

A Virus and Herpesviruses: Similar Mechanisms and Common Themes. Viruses.

2016;8(4):102.

13.   Dremel SE, DeLuca NA. Herpes simplex viral nucleoprotein creates a competitive

transcriptional environment facilitating robust viral transcription and host shut off. Elife.

2019;8.

14.   McSwiggen DT, Hansen AS, Teves SS, Marie-Nelly H, Hao Y, Heckert AB, et al.

Evidence for DNA-mediated nuclear compartmentalization distinct from phase separation.

eLife. 2019;8:e47098.

15.   Isa NF, Bensaude O, Aziz NC, Murphy S. HSV-1 ICP22 Is a Selective Viral Repressor

of Cellular RNA Polymerase II-Mediated Transcription Elongation. Vaccines.

2021;9(10):1054.

16.   Kwong AD, Frenkel N. Herpes simplex virus-infected cells contain a function(s) that

destabilizes both host and viral mRNAs. Proc Natl Acad Sci U S A. 1987;84(7):1926-30.

17.   Oroskar AA, Read GS. Control of mRNA stability by the virion host shutoff function

of herpes simplex virus. J Virol. 1989;63(5):1897-906.

18.   Feng P, Everly DN, Jr., Read GS. mRNA decay during herpesvirus infections:

interaction between a putative viral nuclease and a cellular translation factor. J Virol.

2001;75(21):10272-80.

19.   Friedel CC, Whisnant AW, Djakovic L, Rutkowski AJ, Friedl MS, Kluge M, et al.

Dissecting Herpes Simplex Virus 1-Induced Host Shutoff at the RNA Level. J Virol.

2021;95(3).

20.    Rutkowski AJ, Erhard F, L'Hernault A, Bonfert T, Schilhabel M, Crump C, et al.

Widespread disruption of host transcription termination in HSV-1 infection. Nat Commun.

2015;6:7126.

21.    Hennig T, Michalski M, Rutkowski AJ, Djakovic L, Whisnant AW, Friedl MS, et al.

HSV-1-induced disruption of transcription termination resembles a cellular stress response but

selectively    increases    chromatin    accessibility    downstream    of    genes.    PLoS    Pathog.

2018;14(3):e1006954.

22.    Wang X, Hennig T, Whisnant AW, Erhard F, Prusty BK, Friedel CC, et al. Herpes

simplex virus blocks host transcription termination via the bimodal activities of ICP27. Nat

Commun. 2020;11(1):293.

23.    Vilborg A, Passarelli MC, Yario TA, Tycowski KT, Steitz JA. Widespread Inducible

Transcription Downstream of Human Genes. Mol Cell. 2015;59(3):449-61.

24.    Rosa-Mercado NA, Zimmer JT, Apostolidi M, Rinehart J, Simon MD, Steitz JA.

Hyperosmotic stress alters the RNA polymerase II interactome and induces readthrough

transcription despite widespread transcriptional repression. Mol Cell. 2021;81(3):502-13 e4.

25.    Hadar S, Meller A, Saida N, Shalgi R. Stress-induced transcriptional readthrough into

neighboring genes is linked to intron retention. iScience. 2022;25(12):105543.

26.    Wu N, Watkins SC, Schaffer PA, DeLuca NA. Prolonged gene expression and cell

survival after infection by a herpes simplex virus mutant defective in the immediate-early genes

encoding ICP4, ICP27, and ICP22. Journal of Virology. 1996;70(9):6358-69.

27.    Honess RW, Watson DH. Herpes simplex virus resistance and sensitivity to

phosphonoacetic acid. Journal of Virology. 1977;21(2):584-600.

28.    Becker Y, Asher Y, Cohen Y, Weinberg-Zahlering E, Shlomai J. Phosphonoacetic

Acid-Resistant Mutants of Herpes Simplex Virus: Effect of Phosphonoacetic Acid on Virus

Replication and In Vitro Deoxyribonucleic Acid Synthesis in Isolated Nuclei. Antimicrobial

Agents and Chemotherapy. 1977;11(5):919-22.

29.    Djakovic L, Hennig T, Reinisch K, Milic A, Whisnant AW, Wolf K, et al. The HSV-1

ICP22 protein selectively impairs histone repositioning upon Pol II transcription downstream

of genes. Nat Commun. 2023;14(1):4591.

30.    Weiner A, Hughes A, Yassour M, Rando OJ, Friedman N. High-resolution nucleosome

mapping reveals transcription-dependent promoter packaging. Genome Res. 2010;20(1):90-

100.

31.    Schwabish MA SK. Evidence for eviction and rapid deposition of histones upon

transcriptional elongation by RNA polymerase II. Mol Cell Biol. 2004;24(23):10111-7.

32.    Barski A, Cuddapah S, Cui K, Roh TY, Schones DE, Wang Z, et al. High-resolution

profiling of histone methylations in the human genome. Cell. 2007;129(4):823-37.

33.    Raisner RM, Hartley PD, Meneghini MD, Bao MZ, Liu CL, Schreiber SL, et al. Histone

variant H2A.Z marks the 5' ends of both active and inactive genes in euchromatin. Cell.

2005;123(2):233-48.

34.    Li S, Wei T, Panchenko AR. Histone variant H2A.Z modulates nucleosome dynamics

to promote DNA accessibility. Nature Communications. 2023;14(1):769.

35.    Bensaude O. Inhibiting eukaryotic transcription. Which compound to choose? How to

evaluate its activity? Transcription. 2011;2(3):103-8.

36.    Weiss E, Friedel CC. RegCFinder: targeted discovery of genomic subregions with

differential read density. Bioinform Adv. 2023;3(1):vbad085.

37.    Anders S RA, Huber W. Detecting differential usage of exons from RNA-seq data.

Genome Research. 2012;22(10):2008-17.

38.    Weiss E, Hennig T, Grassl P, Djakovic L, Whisnant AW, Jurges CS, et al. HSV-1

Infection Induces a Downstream Shift of Promoter-Proximal Pausing for Host Genes. J Virol.

2023;97(5):e0038123.

39. Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, et al. Simple combinations

of lineage-determining transcription factors prime cis-regulatory elements required for

macrophage and B cell identities. Mol Cell. 2010;38(4):576-89.

40. Dremel SE, Sivrich FL, Tucker JM, Glaunsinger BA, DeLuca NA. Manipulation of

RNA polymerase III by Herpes Simplex Virus-1. Nature Communications. 2022;13(1):623.

41. Corces MR, Trevino AE, Hamilton EG, Greenside PG, Sinnott-Armstrong NA, Vesuna

S, et al. An improved ATAC-seq protocol reduces background and enables interrogation of

frozen tissues. Nature Methods. 2017;14(10):959-62.

42. Uprichard SL, Knipe DM. Herpes simplex ICP27 mutant viruses exhibit reduced

expression of specific DNA replication genes. Journal of Virology. 1996;70(3):1969-80.

43. Greenberg RS, Long HK, Swigut T, Wysocka J. Single Amino Acid Change Underlies

Distinct Roles of H2A.Z Subtypes in Human Syndrome. Cell. 2019;178(6):1421-36 e24.

44. Bagchi DN, Battenhouse AM, Park D, Iyer VR. The histone variant H2A.Z in yeast is

almost exclusively incorporated into the +1 nucleosome in the direction of transcription.

Nucleic Acids Research. 2020;48(1):157-70.

45. Lashgari A, Millau J-F, Jacques P-É, Gaudreau L. Global inhibition of transcription

causes an increase in histone H2A.Z incorporation within gene bodies. Nucleic Acids Research.

2017;45(22):12715-22.

46. Wyler E, Menegatti J, Franke V, Kocks C, Boltengagen A, Hennig T, et al. Widespread

activation of antisense transcription of the host genome during herpes simplex virus 1 infection.

Genome Biology. 2017;18(1):209.

47. Jimeno-González S, Ceballos-Chávez M, Reyes JC. A positioned +1 nucleosome

enhances promoter-proximal pausing. Nucleic Acids Research. 2015;43(6):3068-78.

48. Aoi Y, Smith ER, Shah AP, Rendleman EJ, Marshall SA, Woodfin AR, et al. NELF

Regulates a Promoter-Proximal Step Distinct from RNA Pol II Pause-Release. Molecular Cell.

2020;78(2):261-74.e5.

49.    Conn KL HM, Schang LM. Linker histones are mobilized during infection with herpes

simplex virus type 1. J Virol. 2008;82:8629–46.

50.    Conn KL HM, Schang LM. Core histones H2B and H4 are mobilized during infection

with herpes simplex virus 1. J Virol. 2011;85:13234–52.

51.    Conn KL, Schang LM. Chromatin Dynamics during Lytic Infection with Herpes

Simplex Virus 1. Viruses. 2013;5(7):1758-86.

52.    Stow ND, Stow EC. Isolation and Characterization of a Herpes Simplex Virus Type 1

Mutant Containing a Deletion within the Gene Encoding the Immediate Early Polypeptide

Vmw110. Journal of General Virology. 1986;67(12):2571-85.

53.    Post LE, Roizman B. A generalized technique for deletion of specific genes in large

genomes: a gene 22 of herpes simplex virus 1 is not essential for growth. Cell. 1981;25(1):227-

32.

54.    Smith IL, Hardwicke MA, Sandri-Goldin RM. Evidence that the herpes simplex virus

immediate early protein ICP27 acts post-transcriptionally during infection to regulate gene

expression. Virology. 1992;186(1):74-86.

55.    Fenwick ML, Everett RD. Inactivation of the Shutoff Gene (UL41) of Herpes Simplex

Virus Types 1 and 2. Journal of General Virology. 1990;71(12):2961-7.

56.    DeLuca NA, Schaffer PA. Physical and functional domains of the herpes simplex virus

transcriptional regulatory protein ICP4. Journal of Virology. 1988;62(3):732-43.

57.    Kluge M, Friedel CC. Watchdog - a workflow management system for the distributed

analysis of large-scale experimental data. BMC Bioinformatics. 2018;19(1):97.

58.    Kluge M, Friedl MS, Menzel AL, Friedel CC. Watchdog 2.0: New developments for

reusability, reproducibility, and workflow execution. Gigascience. 2020;9(6).

59.    Bonfert T, Kirner E, Csaba G, Zimmer R, Friedel CC. ContextMap 2: fast and accurate

context-based RNA-seq mapping. BMC Bioinformatics. 2015;16:122.

60.    Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler

transform. Bioinformatics. 2009;25(14):1754-60.

61.    Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, et al. Twelve years

of SAMtools and BCFtools. Gigascience. 2021;10(2).

62.    Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic

features. Bioinformatics. 2010;26(6):841-2.

63.    R Core Team. R: A Language and Environment for Statistical Computing. Vienna,

Austria2022.

64.    Hahne F, Ivanek R. Visualizing Genomic Data Using Gviz and Bioconductor. Methods

Mol Biol. 2016;1418:335-51.

65.    Chirackal Manavalan AP, Pilarova K, Kluge M, Bartholomeeusen K, Rajecky M,

Oppelt J, et al. CDK12 controls G1/S progression by regulating RNAPII processivity at core

DNA replication genes. EMBO Rep. 2019;20(9):e47592.

66.    Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic

features. Bioinformatics. 2010;26(6):841-2.

67.    Boyle AP, Guinney J, Crawford GE, Furey TS. F-Seq: a feature density estimator for

high-throughput sequence tags. Bioinformatics. 2008;24(21):2537-8.

68.    Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for

assigning sequence reads to genomic features. Bioinformatics. 2014;30(7):923-30.

69.    Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. Genome Biol. 2014;15(12):550.

70.    Benjamini Y HY. Controlling the False Discovery Rate: A Practical and Powerful

Approach to Multiple Testing. Journal of the Royal Statistical Society Series B

(Methodological). 1995;57(1):289-300.

71.    Raudvere U, Kolberg L, Kuzmin I, Arak T, Adler P, Peterson H, et al. g:Profiler: a web

server for functional enrichment analysis and conversions of gene lists (2019 update). Nucleic

Acids Res. 2019;47(W1):W191-W8.

72.    Kolberg L, Raudvere U, Kuzmin I, Vilo J, Peterson H. gprofiler2 -- an R package for

gene list functional enrichment analysis and namespace conversion toolset g:Profiler.

F1000Res. 2020;9.

**Supporting information**

**S1 File. Supplementary Figures.**

Contains Supplementary Figures 1-28 and legends (PDF).

**S2 File. Supplementary Tables.**

Contains Supplementary Table 1 with statistics on all ATAC-seq samples analyzed for this

study (XLSX).

**Figures**

**Fig 1: HSV-1 infection impacts chromatin accessibility around promoters for the majority**

**of host genes.**

**(a)** Heatmap illustrating the location of differential regions (m- and i-RegC) identified by

RegCFinder on the ATAC-seq data for WT, ΔICP0, ΔICP22, ΔICP27, and Δ*vhs* infection in

comparison to mock infection. Results are shown for 4,981 (out of 7,649) promoter windows

containing at least one statistically significant (adj. p. ≤ 0.01) differential region for at least one

virus infection compared to mock. Red and blue colors represent m-RegC (mock>infection)

and i-RegC (infection>mock) locations, respectively, within promoter windows. Here, each

row of the heatmap shows results for the same input window in different comparisons of virus

infection to mock. Colored rectangles on top of the heatmap indicate which virus infection was

compared to mock. Black vertical lines in the center of each part of the heatmap indicate the

position of the TSS. Hierarchical clustering was performed according to Euclidean distances

and Ward's clustering criterion and the cutoff on the hierarchical clustering dendrogram was

selected to obtain 14 clusters (marked by colored rectangles between dendrogram and heatmap

and numbered from top to bottom as indicated). Log2 fold-changes for differential regions are

shown in **Sup. Fig 1d** in **S1 File. (b)** Metagene plot showing the average ATAC-seq profile

around the TSS ± 3 kb in mock (red) and WT (blue) infection for cluster 5, an example for

pattern (i). For a description of metagene plots, see Materials and Methods. The colored bands

below the metagene curves in each panel indicate the percentage of genes having an m- or i-

RegC (red or blue, respectively) at that position. **(c)** Read coverage in a ±3.6 kb window around

the TSS in ATAC-seq data of mock (red) and WT (blue) infection for example gene FBXO28

with pattern (i). Read coverage was normalized to total number of mapped reads for each

sample and averaged between replicates. The TSS is indicated by a short vertical line below the

read coverage track for mock infection. Gene annotation is indicated at the top. Boxes represent

exons, lines represent introns, and direction of transcription is indicated by arrowheads. Below

the read coverage track for WT infection, m-RegC (red bars) and i-RegC (blue bars) are shown

for the comparison of WT vs. mock infection. **(d)** Metagene plot as in **(b)** for cluster 7, the most

pronounced case of pattern (ii). **(e)** Read coverage plot as in **(c)** for example gene ARHGAP1

with pattern (ii). Here, the promoter window also contains the TSS of the ZNF408 gene on the

opposite strand and the accessible chromatin region is extended upstream of the ARHGAP1

TSS, i.e. downstream of the ZNF408 TSS. **(f,g)** Metagene plots as in **(b)** for **(f)** cluster 13,

which exhibits pattern (iii), and **(g)** cluster 8, one of two clusters exhibiting the combined (i)+(ii)

pattern.

**Fig 2: Changes in chromatin accessibility at promoters are mostly independent of dOCR**

**induction.**

**(a)** Bar plots showing the percentage of promoter regions for each cluster overlapping with

dOCRs downstream of the 1,296 genes for which we previously showed consistent dOCR

induction across different HSV-1 strains (29). dOCR regions in mock, WT ($\pm$ PAA treatment)

and null mutant infection were calculated as previous described (29) (see Materials and

Methods for further details) and the overlaps of promoter windows to dOCR regions originating

from an upstream gene were determined for each cluster. WT ($\pm$ PAA treatment) and null

mutant infection are ordered according to the overall extent of dOCR induction. Only few

clusters (12-14) showed some enrichment for dOCRs from upstream genes, which was

abolished in $\Delta$ICP22 infection. **(b-c)** Metagene curves showing average ATAC-seq profiles in

promoter windows for mock (red), WT (blue), WT+PAA (green) and $\Delta$ICP22 (orange)

infection for clusters 5 **(b)** and 7 **(c)**. (**d**) Barplot showing the percentage of genes in each cluster

annotated as either protein-coding, long intervening/intergenic noncoding RNAs (lincRNA),

snRNA, antisense, or other in Ensembl. Significance of enrichment of each gene type in each

cluster was determined using a one-sided Fisher's exact test (with alternative = greater) and p-

values were corrected for multiple testing using the Benjamini-Hochberg method. Adj. p. <

0.05 are indicated in the corresponding field of the barplot. (**e**) Percentage of bidirectional

promoters in each cluster was calculated as the percentage of promoter windows containing a

protein-coding, lincRNA or antisense gene (according to Ensembl annotation) on the opposite

strand to the target gene starting within 1 kb upstream of the TSS of the target gene. Enrichment

and significance analysis and multiple testing correction was performed as for (**d**) and adj. p. <

0.05 are indicated on top of bars.

1012

1013

**Fig 3: Changes in chromatin accessibility at promoters are linked to transcription.**

**(a,b)** Boxplots showing the distribution of **(a)** log2(DSR:UAR) ratios and **(b)** gene expression (FPKM) values in chromatin-associated RNA in mock (red) and 8 h p.i. WT (blue) HSV-1 infection for all clusters (grouped by pattern, cluster numbers are shown below each boxplot) and remaining genes without significant chromatin accessibility changes (NA group). DSR was calculated as the expression in chromatin-associated RNA for the region downstream of the TSS in sense direction of the target gene and UAR as the expression in the upstream region in antisense direction. P-values for Wilcoxon rank sum tests comparing values in mock infection for each cluster against all other analyzed genes are indicated below cluster numbers and were corrected for multiple testing using the Bonferroni method. **(c)** Barplot showing the percentage of genes with very low, low, medium, high and very high gene expression with at least one significant differential region (RegC). Here, gene expression cutoffs for the five groups were determined such that each group contains the same number of genes and are indicated below labels on the x-axis. Multiple testing corrected p-values for two-sided Fisher's exact tests comparing the fraction of genes with RegC between subsequent gene expression groups are indicated on top of bars. **(d)** Percentage of genes within each cluster among those genes with at least one significant RegC for the five gene expression groups from **(c)**. Significance of enrichment or depletion for each cluster in each gene group was determined with two-sided Fisher's exact tests and multiple testing correction was performed with the method by Benjamini and Hochberg. Adj. p. < 0.05 are shown and are underlined in case the cluster is enriched and not underlined if it is depleted. **(e-g)** Metagene curves showing average ATAC-seq profiles in promoter windows for mock (red) and WT infection (blue) for genes with very low, medium and very high expression from **(b)**.

 **Fig. 4: Changes in chromatin accessibility begin to manifest between 4 and 8 h p.i.**

 **depending on gene expression**

**(a)** Heatmap showing the log2 fold-changes determined with DEXSeq on the ATAC-seq time-

course data (1, 2, 4, 6 and 8 h p.i. WT infection compared to mock infection) for statistically

significant differential regions (m- and i-RegC) identified by RegCFinder. For comparison,

log2 fold-changes for the WT vs. mock comparison from **Fig 1a** are also shown. Each row

represents the results for one of the input windows included in **Fig 1a**. Colored rectangles on

top indicate the time-point of infection or whether the original WT vs. mock comparison is

shown. Statistically significant differential regions are colored according to the log2 fold-

change determined by DEXSeq. Here, the color scale is continuous between -1 and 1 and all

log2 fold-changes >1 are colored the same red and all log2 fold-changes <1 the same blue.

Promoter windows are ordered as in **Fig 1a** and clusters are annotated by colored and numbered

rectangles on the left. **(b)** Number of statistically significant differential regions (m- and i-

RegC) and number of genes with at least one statistically significant differential region

identified for each time-point of infection in the ATAC-seq time-course data. **(c)** Boxplot

showing the distribution of gene expression (FPKM) values in chromatin-associated RNA for

mock and 8 h p.i. WT infection for genes with significant changes in chromatin accessibility in

promoter windows (i) at 4 h p.i. or earlier, (ii) at 6 h p.i. but not yet at 4 h p.i., (iii) at 8 h p.i.

but not yet at 6 h p.i. and (iv) in the analysis shown in **Fig 1a**, but not yet at 8 h p.i. in the time-

course ATAC-seq experiment. P-values for Wilcoxon rank sum tests comparing FPKM values

in mock infection between subsequent groups are also indicated. **(d-f)** Metagene curves of

ATAC-seq profiles in mock infection (red) and all time-points of infection (blue shades) from

the time-course experiment for example clusters 5 (pattern (i), **d**), 7 (pattern (ii), **e**) and 13

(pattern (iii), **f**). The colored bands below the metagene curves indicate the percentage of genes

having an i- or m-RegC (blue or red, respectively) at that position in the comparison of 8 h p.i.

to mock infection from the time-course experiment. Metagene plots for all clusters are shown

in **Sup. Fig 21** in **S1 File**.

**Fig 5: Chromatin accessibility changes upon ICP4 knockout and dox-induced combined**

**ICP22 and ICP27 expression**

**(a-b)** Metagene curves showing average ATAC-seq profiles in promoter windows for mock

(red), WT (blue), WT+PAA (green) and ΔICP4 (violet) infection for clusters 5 **(a)** and 7 **(b)**.

Metagene plots for all other clusters are shown in **Sup. Fig 23** in **S1 File**. **(c)** Heatmap showing

the location of differential regions (m- and i-RegC) identified by RegCFinder for T-HF-ICP27

cells and T-HF-ICP22/ICP27 cells upon dox exposure. Each row represents the results for one

of the input windows included in **Fig 1a**. Promoter windows are ordered as in **Fig 1a** and

clusters are annotated as colored and numbered rectangles on the left. Log2 fold-changes for

differential regions are shown in **Sup. Fig 24b** in **S1 File**. **(d-e)** Metagene curves showing

average ATAC-seq profiles in promoter windows for T-HF-ICP22/ICP27 cells with (blue) and

without (red) dox exposure for clusters 5 **(d)** and 7 **(e)**. The colored bands below the metagene

curves indicate the percentage of genes having an m-RegC (red, decreased upon dox exposure)

or i-RegC (blue, increased upon dox exposured) or at that position. Metagene plots for all

clusters are shown in **Sup. Fig 24** in **S1 File**.

1080

**Fig 6: HSV-1 infection and depletion of Pol II from human genes lead to a downstream**

**shift of H2A.Z-containing +1 nucleosomes.**

**(a)** Heatmap showing the location for differential regions (m- and i-RegC) identified in the WT

vs. mock comparison on the ATAC-seq (left half, identical to left-most part of **Fig 1a**) and

H2A.Z ChIPmentation data (right half) for the promoter windows included in **Fig 1a.**

Differential regions with mock>infection (m-RegC) are marked in red, differential regions with

infection>mock (i-RegC) in blue. The order of promoter windows is the same as in **Fig 1a** and

clusters are indicated by colored and numbered rectangles on the left. Log2 fold-changes for

differential regions are shown in **Sup. Fig 25b** in **S1 File**. **(b,c)** Read coverage in a ±3.6 kb

window around the TSS in H2A.Z ChIPmentation data for mock (red) and WT (blue) infection

for example genes with **(b)** pattern (i) (FBXO28) and **(c)** pattern (ii) (ARHGAP1). For

descriptions of read coverage plots see caption to **Fig 1**. **(d-g)** Metagene plots of H2A.Z profiles

for mock and WT infection **(d,e)** and untreated and α-amanitin-treated HCT116 cells **(f,g)** for

clusters 5 (pattern (i), **d,f**) and 7 (pattern (ii), **e,g**). Metagene plots for other clusters can be

found in **Sup. Fig 25** and **27** in **S1 File**, respectively. The colored bands below the metagene

curves in each panel indicate the percentage of genes having m- or i-RegC at that position. For

**(f,g)**, m-RegC (green) are differential regions with relative read coverage higher in untreated

cells and i-RegC (violet) differential regions with relative read coverage higher upon α-

amanitin-treatment.