HSV-1 infection induces a downstream-shift of +1 nucleosomes

Elena Weiß[1], Adam W. Whisnant[2,3], Thomas Hennig[2,3], Lara Djakovic[2], Lars Dölken[2,3,4#],

Caroline C. Friedel[1#]

[1] Institute of Informatics, Ludwig-Maximilians-Universität München, Amalienstr. 17, 80333

Munich, Germany

[2] Institute for Virology and Immunobiology, Julius-Maximilians-University Würzburg,

Versbacher Straße 7, 97078 Würzburg, Germany

[3] Institute for Virology, Medizinische Hochschule Hannover, Carl-Neuberg-Str. 1, 30625

Hannover, Germany

[4] Helmholtz Institute for RNA-based Infection Research (HIRI), Helmholtz-Center for

Infection Research (HZI), 97080 Würzburg, Germany

[*] Corresponding authors

E-mail: lars.doelken@uni-wuerzburg.de; Caroline.Friedel@bio.ifi.lmu.de

Abstract

Herpes simplex virus 1 (HSV-1) infection induces a genome-wide loss of host transcriptional activity and widespread disruption of host transcription termination. The latter leads to read-through transcription for thousands of nucleotides beyond poly(A) sites and is associated with induction of open chromatin downstream of genes. In this study, we show that chromatin accessibility around promoters is also massively altered by HSV-1 infection as nucleosome-free regions around promoters are extended into regions downstream of transcription start sites, in particular for highly expressed genes. This was independent of HSV-1 immediate-early genes ICP0, ICP22, and ICP27 and the virion host shutoff protein *vhs*, but was reduced by treatment with phosphonoacetic acid (PAA), which inhibits viral DNA replication and alleviates the loss of RNA Polymerase II (Pol II) from host genes in HSV-1 infection. ChIPmentation analysis of the noncanonical histone variant H2A.Z, which is strongly enriched at +1 and -1 nucleosomes, indicated that the downstream extension of accessible chromatin at promoters was linked to a downstream shift of +1 nucleosomes. In yeast, downstream shifts of +1 nucleosomes can be induced by Pol II degradation. We confirmed that Pol II depletion is sufficient to similarly alter +1 nucleosome positioning in human cells by analyzing H2A.Z occupancy upon treatment with α-amanitin, which triggers degradation of the largest Pol II subunit. In summary, our study provides strong evidence that HSV-1-induced depletion of Pol II from the host genome leads to a widespread downstream shift of +1 nucleosomes at host gene promoters.

Author summary

Herpes simplex virus 1 (HSV-1) infection leads to a massive shutoff of host transcription. Loss of RNA Polymerase II (Pol II) in yeast has previously been shown to lead to a relaxation of +1 nucleosome positioning to more thermodynamically favorable sites downstream of transcription start sites. In this study, we show that a similar phenomenon is likely also at play in HSV-1 infection as sequencing of accessible chromatin reveals a widening of nucleosome-free regions at promoters into downstream regions. By mapping genome-wide positions of the noncanonical histone variant H2A.Z enriched at +1 and -1 nucleosomes, we reveal a downstream shift of +1 nucleosomes for the majority of genes in HSV-1 infection. We furthermore confirm that depletion of Pol II alone leads to downstream shifts of +1 nucleosomes in human cells. Thus, changes in chromatin architecture at promoters in HSV-1 infection are likely a consequence of HSV-1-induced depletion of Pol II from the host genome.

Introduction

Herpes simplex virus 1 (HSV-1) is one of nine human herpesviruses, with more than half of the global population being latently infected with HSV-1 (1-4). HSV-1 is known for causing the common cold sores but is also responsible for life-threatening diseases particularly in the immunocompromised (5). The HSV-1 genome consists of 152 kilobases (kb) of double-stranded DNA and encodes for at least 121 large open reading frames (ORFs) and >100 small ORFs (6). Upon lytic infection, the five immediate early (IE) genes ICP0, ICP4, ICP22, ICP27, and ICP47 are expressed and hijack the host gene expression machinery to ensure viral replication during early stages of infection (7-11). This results in recruitment of RNA polymerase II (Pol II) from the host chromatin to viral genomes and a general shutoff of host transcription (7, 9, 11-13). Pol II recruitment to viral genomes is predominantly facilitated by the viral IE protein ICP4 (13), which effectively sequesters Pol II into the viral replication compartment (14). However, ICP4 already depletes Pol II from cellular promoters as early as 2 h post infection (p.i.), i.e. before the onset of viral genome replication, and independent of viral genome copy number (15). The global host transcription shutoff is further exacerbated by ICP22-mediated inhibition of host transcription elongation (16) and the viral *vhs* (virion host shutoff) protein. *vhs* is a nuclease delivered by the tegument of the incoming viral particles and cleaves both viral and host mRNAs (17-19). We recently demonstrated that *vhs* continuously degrades about 30 % of host mRNAs per hour during the first 8 h of high multiplicity lytic infection of primary human fibroblasts (20).

HSV-1 further disturbs host transcription by inducing a widespread disruption of host transcription termination (DoTT) resulting in read-through transcription beyond polyadenylation (pA) sites of host genes for tens of thousands of nucleotides (nt) (21). Read-through transcription can even extend into downstream genes, leading to a seeming

transcriptional induction of these genes. DoTT negatively affects host gene expression in two

ways: First, as read-through transcripts are not exported (22), it effectively prevents

translation of newly transcribed mRNAs and thus dampens the host transcriptional response

to infection. Second, it redirects a large fraction of ongoing Pol II transcription from host gene

bodies into downstream intergenic regions and, thus, further exacerbates the depletion of Pol

II from host genes. We recently showed that ICP27 is sufficient for inducing DoTT, but DoTT

is also observed in absence of ICP27, albeit at substantially reduced levels (23). As cellular

stresses also induce transcription downstream of genes (DoG) (22, 24-26), HSV-1-induced

read-through transcription in absence of ICP27 presumably results from a virus-induced

cellular stress response (27). In contrast to stress-induced DoGs, HSV-1-induced DoTT is

associated with a massive increase of chromatin accessibility downstream of genes (denoted

as downstream open chromatin regions (dOCRs)) (22). dOCR induction requires strong

absolute levels of transcription downstream of genes, thus it is predominantly observed for

highly transcribed genes with strong read-through. Treatment with phosphonoacetic acid

(PAA) during HSV-1 infection, which inhibits viral DNA replication (28, 29), results in

increased dOCR induction (30). As viral DNA redirects Pol II from the host chromatin, PAA

treatment mitigates the loss of Pol II from the host genome and thus increases absolute levels

of transcription on host genes and – importantly for dOCR induction – read-through regions.

Although dOCRs can extend into downstream genes following read-through transcription, this

occurs only in cases of very strong read-through transcription into these downstream genes.

We recently showed that ICP22 is both required and sufficient for inducing dOCRs in

presence of ICP27-induced read-through (30). This discovery was based on extensive ATAC-

seq (Assay for Transposase-Accessible Chromatin using sequencing) analysis of both wild-

type (WT) HSV-1 infection as well as infection with null mutants for ICP0, ICP22, ICP27,

and *vhs*.

As we previously only investigated chromatin accessibility *downstream* of genes, we now

further explore these data to investigate changes in chromatin accessibility around host gene

promoters. This revealed a broadening of open chromatin regions at the promoter into regions

downstream of transcription start sites (TSS) for the majority of host genes. While it was

consistently observed in absence of ICP0, ICP22, ICP27, and *vhs*, PAA treatment, which

alleviates the HSV-1-induced loss of transcription, substantially reduced the extension of

open chromatin at the TSS. Analysis of *de novo* transcription using RNA-seq of chromatin-

associated RNA revealed a link of this phenomenon to transcription as (i) more highly

expressed genes were more strongly affected and (ii) genes with strong antisense transcription

exhibited the opposite effect with a broadening of open chromatin into regions upstream of

the TSS. ChIPmentation for the noncanonical histone H2A variant H2A.Z enriched at +1 and

-1 nucleosomes (31, 32) indicated that changes in chromatin accessibility reflected a

downstream shift of +1 nucleosomes in HSV-1 infection. We observed similar changes in

H2A.Z occupancy upon α-amanitin treatment, a deadly toxin found in Amanita mushrooms,

which inhibits transcription by preventing translocation of Pol II and triggering degradation of

Rpb1, the largest subunit of Pol II (33). Furthermore, Weiner *et al.* previously showed that Pol

II depletion in yeast induces downstream shifts of +1 nucleosomes, in particular for highly

expressed genes (34). Transcribing Pol II disturbs histone-DNA interactions and at high

transcription rates leads to eviction of nucleosomes from the chromatin (35). Weiner *et al.*

proposed that Pol II depletion relaxes chromatin allowing nucleosomes to shift to more

thermodynamically favorable sites. As nucleosome positioning of highly expressed genes

deviates more strongly from *in vitro* nucleosome preferences (36), this explains why these

genes showed more pronounced shifts. These results provide strong evidence that depletion of

Pol II from host genes during HSV-1 infection leads to the downstream shifts of +1

nucleosomes to more thermodynamically favorable sites.

Results

Widespread extension of chromatin-free regions downstream of transcription start sites

To investigate changes in chromatin accessibility within promoter regions during HSV-1 infection, we re-analyzed our recently published ATAC-seq data (30) for mock and wild-type (WT) HSV-1 infection of human foreskin fibroblasts (HFF) as well as infection with null mutant viruses for the HSV-1 IE proteins ICP0, ICP22, and ICP27 as well as *vhs*. Samples for mock, WT, and ΔICP27 infection were collected at 8 h p.i., while samples for ΔICP0 and ΔICP22 infection were collected at 12 h p.i. as these null-mutant infections progress more slowly. To assess chromatin accessibility changes in promoter regions, we identified genomic regions with differential ATAC-seq read density using our newly published RegCFinder method (37). RegCFinder searches for genomic regions for which the read distribution differs between two conditions (e.g., mock and WT infection) and is targeted towards regions of interest by specifying genomic windows as input (**Sup. Fig. 1a**). For each input window, subregions are identified that show differences in the distribution of reads within this window. Statistical significance for identified subregions is determined by DEXSeq (38). For our purposes, we defined the windows of interest as the ± 3kb promoter region around the TSS of 7,649 human genes. These TSSs were previously defined in our analysis of promoter-proximal Pol II pausing in HSV-1 WT infection (39) based on a re-analysis of published precision nuclear run-on sequencing (PRO-seq) and PROcap-seq (a variation of PRO-seq) data of flavopiridol-treated uninfected HFF. CDK9 inhibition by flavopiridol arrests Pol II in a paused state at the TSS, allowing the mapping of Pol II initiation sites with both PRO-seq and PROcap-seq. Filtering for high-confidence TSSs that were confirmed by both data sets and within a maximum distance of 500 bp to the nearest annotated gene on the same strand identified the major TSS for 7,649 genes. Using these 6 kb promoter windows as input, we applied RegCFinder to our ATAC-seq data for WT and all null-mutant infections in

comparison to mock to identify promoter subregions for each gene that show differences in

the distribution of open chromatin between infection and mock. It should be noted that we did

not filter these windows beforehand to exclude genes for which dOCRs extended from an

upstream gene into their promoter window for two reasons: (i) Since dOCRs represent

changes in chromatin accessibility, RegCFinder should also detect them; (ii) The further the

distance from the upstream gene, the less pronounced are changes in chromatin accessibility

in dOCR regions, making it difficult to distinguish them from background. Any approach we

tested to filter promoter windows overlapping with dOCRs from upstream genes before the

RegCFinder analysis either excluded too many or too few windows. Thus, we decided to first

run RegCFinder and then investigate which of the detected changes were due to induction of

dOCRs for upstream genes.

RegCFinder identified 23-24,000 differential subregions (in the following also denoted as

RegC, short for regions of change) for each virus infection (WT: 23,357 RegC, ΔICP0:

23,532, ΔICP22: 24,640, ΔICP27: 23,523, Δ*vhs*: 23,716, **Sup. Fig. 1b**). Between 29 and 37 %

of these showed a statistically significant difference in read density within the corresponding

6 kb promoter windows in infection compared to mock (multiple testing adjusted p-value (adj.

p.) ≤ 0.01, WT: 8,552 RegC, ΔICP0: 6,740, ΔICP22: 5,710, ΔICP27: 7,188, Δ*vhs*: 6,860).

These represented between 3,129 (ΔICP22 infection) and 4,391 (WT) genes (**Sup. Fig. 1c**).

The lower fraction of statistically significant differential regions in ΔICP22 infection is likely

due to the relatively low number of ATAC-seq reads mapped to the host genome compared to

the other infections (~3.8-fold fewer, see Supplementary Data 1 in (30)). **Fig. 1a** visualizes

the positions of identified RegC for the 4,981 promoter windows containing at least one

statistically significant (adj. p. ≤ 0.01) differential region for at least one virus infection

compared to mock and **Sup. Fig. 1d** illustrates log2 fold-changes for statistically significant

regions. Here, blue indicates that relative read density in that subregion is increased during

virus infection compared to mock (infection>mock, denoted as i-RegC in the following) and

red that relative read density in that subregion is higher in mock compared to infection

(mock>infection, denoted as m-RegC).

Strikingly, our results showed a highly homogeneous picture for WT and null-mutant

infections compared to mock, with changes in the same direction observed at approximately

the same locations. To identify distinct patterns of changes in chromatin accessibility, we

performed hierarchical clustering on the RegC location heatmap in **Fig. 1a** and selected a

cutoff on the dendrogram to obtain 14 clusters identified by visual inspection. To visualize

average read density for each cluster, we performed metagene analyses of promoter windows

combined with statistics on RegC locations for each cluster (**Fig. 1b,d,f,g, Sup. Fig. 2**). For

metagene analyses, each 6 kb input window was divided into 101 bins of ~59 bp. For each

bin, ATAC-seq read counts were determined, normalized to sequencing depth, normalized to

sum up to 1 for each input window, averaged across all windows in each cluster and then

averaged across replicates. For each bin in the promoter window, metagene curves also show

which fraction of windows have an i- (blue) or m-RegC (red) at that position in WT infection.

Based on the location heatmap and the metagene analyses, we identified three major patterns

in changes in chromatin accessibility around promoters during WT infection: (i) ATAC-seq

peaks at the TSS that shift and/or broaden into regions downstream of the TSS during HSV-1

infection (clusters 1, 2, 4, 5, 9, 10, 12, 14, a total of 3,472 genes, **Fig. 1b, Sup. Fig. 2a-g,**

examples in **Fig. 1c, Sup. Fig. 3a-h**), (ii) ATAC-seq peaks at the TSS that shift and/or

broaden into regions upstream of the TSS (clusters 6, 7, 11, a total of 919 genes, **Fig. 1d, Sup.**

Fig. 2h,i, examples in **Fig. 1e, Sup. Fig. 3i-k**), and (iii) an increase in chromatin-accessibility

both up- and downstream of the TSS peak in infection compared to mock (cluster 13, 126

genes, **Fig. 1f,** example in **Sup. Fig. 3l**). Two clusters (3 and 8, a total of 464 genes) exhibited

a combination of patterns (i) and (ii) with an extension of the TSS peak in both up- and

downstream direction (**Fig. 1g, Sup. Fig. 2j, Sup. Fig. 3m,n**).

As a first analysis, we performed motif discovery both for novel and known motifs in m- and

i-RegC grouped either by cluster or pattern compared to the background of all promoter

windows using HOMER (40), but found no enriched motifs. Functional enrichment analysis

for Gene Ontology (GO) terms also yielded no significant results, with the exception of

cluster 13 (pattern (iii)). Cluster 13 was enriched for "mRNA splicing via the spliceosome"

and related terms, however, this was mainly due to several snRNA genes included in this

cluster. Accordingly, when investigating enrichment of clusters for gene types (**Sup. Fig. 1e**),

we found that cluster 13 was significantly enriched for snRNA genes (adj. p. $= 3 \times 10^{-16}$, see

Materials and Methods). In contrast, pattern (i) clusters tended to have relatively high

fractions of protein-coding genes, with clusters 1, 2, 5, and 9 being statistically significantly

enriched. Strikingly, cluster 7 (= most pronounced case of pattern (ii)) was significantly

enriched for antisense transcripts (adj. p. $= 5 \times 10^{-15}$). The two other pattern (ii) clusters 6

and 11 as well as cluster 8 (combined pattern (i) + (ii)) also contained a relatively high

fraction of antisense transcripts, although this was not statistically significant.   The

association of pattern (ii) with antisense transcription is further investigated below.

Most chromatin accessibility changes at promoters are independent of dOCR induction

Next, we evaluated which of the observed changes were due to dOCRs extending into

promoter regions. For this purpose, we determined the fraction of promoter windows in each

cluster that overlapped with dOCRs of the 1,296 genes for which we previously showed

consistent dOCR induction across different HSV-1 strains (30) (**Fig. 2a**). dOCR regions in

mock, WT and all null mutant infections (average across two replicates) were identified as

previously described (22, 30) (see also Materials and Methods). It is important to note here

that some relatively short dOCRs are also observed in mock infection but these substantially

extend during HSV-1 infection. In this analysis, we also included ATAC-seq data obtained in

the same experiment for 8 h p.i. WT infection combined with PAA treatment (WT+PAA) to

identify dOCR regions that are not as clearly detectable without PAA treatment but may still

bias promoter analyses. As ICP22 is required for dOCR induction, overlaps of promoter

windows to dOCR regions in ΔICP22 infection were as low as in mock infection. In other

virus infections, some clusters showed enrichment for dOCR overlaps that increased with the

overall extent of dOCR induction in these viruses. To highlight this trend, WT (± PAA) and

null mutant infections are ordered according to the extent of dOCR induction in **Fig. 2a**. No

cluster had more than 20% of promoter windows overlapping with dOCRs even with PAA

treatment and most had < 5% overlap, indicating that dOCRs represent only a very small

fraction of observed changes in chromatin accessibility around promoters.

Interestingly, cluster 13, which exhibited increased chromatin accessibility surrounding the

TSS peak (pattern (iii)) also exhibited the strongest enrichment for dOCRs during infection.

An example for a gene with dOCRs extending into the promoter is shown in **Sup. Fig. 3l**.

Weaker enrichment for dOCRs was also observed for clusters 12 and 14, for which we

detected regions with increased chromatin accessibility in infection also upstream of the TSS,

in contrast to other pattern (i) clusters (**Sup. Fig. 2f,g,** example with dOCR overlap in **Sup.**

Fig. 3h). In contrast, clusters 3 and 8, which showed a broadening of the TSS peak in both up-

and downstream direction (combined pattern (i)+(ii), **Fig. 1g**, **Sup. Fig. 2j**) were not

associated with dOCR induction. Importantly, in absence of ICP22, which is required for

dOCR induction, we generally observed the same trends in chromatin accessibility changes at

promoters as for WT and the other null-mutant infections (**Sup. Fig. 4-17**). This largely also

applied to clusters 12-14, though a subset of differential regions was no longer observed in

ΔICP22 infection. This applied to some (but not all) regions with increased chromatin

accessibility upstream of genes in clusters 12 and 14 and around TSS peaks in cluster 13 (**Fig.**

1a, Sup. Fig. 1d, Sup. Fig. 15-17). Thus, only some chromatin accessibility changes in

clusters 12-14 were linked to dOCR induction. As absence of ICP0, ICP27 or *vhs* also had

little impact on changes in chromatin accessibility around promoters, we conclude that neither

dOCR induction nor activity of ICP0, ICP22, ICP27 or *vhs* alone explain the majority of

observed changes in chromatin accessibility at promoters in HSV-1 infection.

Extension of accessible chromatin regions around promoters is linked to transcription

In contrast to the single-gene knockouts, PAA treatment substantially reduced, though not

completely abolished, the downstream broadening and/or shift of TSS peaks in pattern (i)

clusters and the upstream broadening and/or shift in pattern (ii) clusters (**Fig. 2b,c, Sup. Fig.**

4-17). Furthermore, fold-changes for differential regions were generally less pronounced

(**Sup. Fig. 18a**). Exceptions were cluster 13, and parts of cluster 12 and 14, for which

chromatin accessibility increased up- (for clusters 12-14) and downstream (for cluster 13) of

the TSS peak (**Fig. 2d, Sup. Fig. 15-17**). This is consistent with these clusters being enriched

for dOCR induction originating from an upstream gene, which is increased upon PAA

treatment. Notably, however, the broadening of the TSS peak into downstream regions in

clusters 12 and 14 was greatly reduced by PAA treatment.

Since PAA treatment alleviates the depletion of Pol II from host genomes, we hypothesized

that loss of Pol II contributes to the observed changes in chromatin accessibility around

promoters in HSV-1 infection. Notably, PAA only reduces expression of late, but not IE and

early HSV-1 proteins (28). Thus, activities of IE, early and tegument-delivered proteins can

still lead to some loss of host gene transcription in presence of PAA. These activities include

depletion of Pol II by ICP4 prior to genome replication (15), loss of Pol II phosphorylation

and inhibition of elongation mediated by ICP22 (16), as well as ICP27-induced inhibition of

transcription termination, which redirects Pol II from gene bodies to intergenic regions (21).

To correlate transcriptional activity of genes to the observed patterns in chromatin

accessibility changes, we analyzed RNA-seq data of chromatin-associated RNA for mock and

8 h p.i. WT HSV-1 infection from our recent study (20). Chromatin-associated RNA depicts

ongoing nascent transcription on the chromatin. Furthermore, changes in chromatin-

associated RNA better reflect transcriptional changes in HSV-1 infection than total RNA

measurements as the latter are confounded by *vhs*-mediated mRNA degradation (20). Our

analysis also included the 2,668 genes (denoted as NA group) without significant chromatin

accessibility changes that were excluded from **Fig. 1a**. Interestingly, metagene analyses for

these genes also revealed a slight broadening of the TSS peak into downstream regions,

although this was much less pronounced than for pattern (i) clusters and therefore not detected

by RegCFinder (**Sup. Fig. 18b**). We analyzed gene expression values in mock and WT

infection (quantified as FPKM = fragments per kilobase million mapped reads, **Fig. 2e**) as

well as log2 fold-changes in gene expression between WT and mock infection (**Sup. Fig.**

18c). Furthermore, we hypothesized that pattern (ii) was shaped by antisense transcription

since pattern (ii) essentially represented the mirror image of pattern (i) (mirrored at a vertical

axis through the TSS), cluster 7 (most pronounced pattern (ii)) was enriched for antisense

transcripts (**Sup. Fig. 1e**), and several pattern (ii) examples showed overlapping or nearby

genes on the opposite strand (e.g. **Fig. 1e**). We thus also calculated the ratio of sense to

antisense transcription in promoter windows for all genes. For this purpose, promoter

windows were divided into the regions down- and upstream of the TSS and expression was

determined in sense direction for the downstream region (=DSR) and in antisense direction

for the upstream region (=UAR). Log2 ratios of DSR to UAR expression are shown in **Fig.**

2f.

We observed no consistent differences in log2 gene expression fold-changes associated with

any of the patterns in chromatin accessibility changes (**Sup. Fig. 18c**). While cluster 1 (pattern

(i)) showed a significant decrease in log2 fold-changes compared to other genes, the

remaining pattern (i) clusters showed either no significant difference or a significant increase

in log2 fold-changes (cluster 9). Similarly, although one pattern (ii) cluster showed a

significant decrease in log2 fold-changes (cluster 6), the other two clusters did not. It should

be noted here that log2 fold-changes represent relative changes in gene expression compared

to all other genes. Thus, genes with positive log2 fold-changes are either increased in

expression or less decreased compared to all other genes, while genes with negative log2 fold-

changes are more decreased in expression compared to all other genes. In contrast, gene

expression levels differed significantly between patterns in both mock and WT infection.

Most pattern (i) clusters as well as cluster 3 (combined pattern (i)+(ii)) exhibited median

expression levels above average in mock (**Fig. 2e**), which was statistically significant for 5

clusters. In contrast, clusters 6, 7 (pattern (ii)) and 8 (combined pattern (i)+(ii)) exhibited

relatively low expression values, with cluster 7 genes having significantly lower expression.

Genes without significant chromatin accessibility changes (NA group) also showed

significantly lower expression, albeit not as low as cluster 7. These differences between

clusters/patterns were generally maintained in HSV-1 infection at lower overall expression

levels, consistent with the absence of differences in gene expression fold-changes between

most clusters.

Strikingly, when investigating DSR to UAR ratios (**Fig. 2f**), we found that all pattern (ii)

clusters had significantly lower DSR to UAR ratios than remaining genes, in particular than

those with pattern (i). In part this was due to their lower expression as the ratio to background

transcription on the antisense strand is lower. Lower expression in WT infection also explains

why log2 DSR:UAR ratios were closer to zero in WT infection for all clusters. However, for

cluster 7, which represented the most pronounced pattern (ii) cluster, the log2 DSR:UAR

ratios were commonly negative, i.e. antisense transcription upstream of the TSS was higher

than sense transcription downstream of the TSS. This suggests that pattern (ii) effectively

represents pattern (i) for the genes on the antisense strand. Notably, for clusters 3 and 8,

which we classified as showing both pattern (i) and (ii) (or pattern (i) in both sense and

antisense direction), DSR:UAR ratios were generally lower than for pattern (i) clusters, albeit

still higher than for pattern (ii) clusters and genes without significant chromatin changes (NA

group). For these genes the extension of accessible chromatin at the promoter was also more

pronounced in sense than in antisense direction. In summary, these results indicate that HSV-

1 infection leads to an extension of the accessible chromatin around promoters for the

majority of genes in the dominant direction of transcription. Here, highly expressed genes

appear to be most strongly affected.

To investigate how early in infection changes in chromatin accessibility around promoters can

be detected, we ran RegCFinder on an ATAC-seq time-course for 1, 2, 4, 6, and 8 h p.i. WT

infection from our previous publication (22) (**Fig. 3a,b**). While barely any significant

differential regions were identified at 1 and 2 h p.i., 1,963 significant regions were identified

in 1,249 promoter windows by 4 h p.i. and 7,558 significant regions in 3,989 promoter

windows by 8 h p.i. (**Fig. 3b**). Log2 fold-changes for significant differential regions generally

reflected the patterns observed in the analysis for WT and null mutant infections (**Fig. 3a**).

The same applied when we separately determined fold-changes and significance on the time-

course ATAC-seq data for the differential regions determined for WT vs. mock from our

initial analysis (**Sup. Fig. 19**) and when we performed metagene analyses for the clusters

from **Fig. 1a** (**Sup. Fig. 20**, example genes in **Sup. Fig. 21**). Although the changes in

chromatin accessibility at 8 h p.i. of the time-course were less pronounced than for the

original WT vs. mock comparison at 8 h p.i., this nevertheless confirms the changes in

chromatin accessibility around promoters in HSV-1 infection in an independent experiment.

Furthermore, it reveals that these changes begin to manifest between 2 and 4 h p.i. We

previously showed that transcriptional activity on host genes drops substantially in the first 4

h of infection to only 40% of transcription in uninfected cells, which was further halved until

8 h p.i. (20). Thus, the onset of the extension of accessible chromatin regions at promoters

into downstream (for pattern (i)) and upstream (for pattern (ii) regions coincides with the

onset of the drop in host transcription during infection.

HSV-1 infection induces a downstream shift of +1 nucleosomes

The HSV-1-induced changes in chromatin accessibility at promoters suggest a broadening of

the nucleosome-free region around promoters in either sense (pattern (i)) or antisense (pattern

(ii)) direction. Consistent with this, we observed downstream and upstream shifts for pattern

(i) and (ii) clusters, respectively, in the nucleosome-depleted regions at promoters in

H3K36me3 ChIPmentation data for mock and 8 h WT HSV-1 infection from our previous

study (30) (**Sup. Fig. 22**). H3K36me3 marks H3 histones on actively transcribed genes (41).

These changes in nucleosome-free regions require a shift in nucleosome positioning, in

particular for +1 or -1 nucleosomes. Several studies performed in yeast previously

investigated the role of transcriptional activity on nucleosome positioning (42-44). In

particular, Weiner *et al.* reported a downstream shift of +1 nucleosomes upon Pol II depletion

in yeast (34). Here, the strongest downstream shifts were observed for very highly expressed

genes, consistent with our observation that pattern (i) was associated with higher gene

expression (**Fig. 2e**). To investigate whether changes in chromatin accessibility at promoters

during HSV-1 infection were linked to shifts in +1 and -1 nucleosome positions, we

performed ChIPmentation in HFF for mock and 8 h p.i. WT infection with an antibody

recognizing the C-terminal part of the non-canonical histone H2A.Z (n=3 replicates). H2A.Z

is highly enriched at gene promoters at -1 and +1 nucleosome positions (31, 32) and is

encoded by two genes, whose protein products H2A.Z.1 and H2A.Z.2 differ by only three

amino acids. While their genomic occupancy patterns are similar, there are quantitative

differences with H2A.Z.1 being more abundant at active promoters than H2A.Z.2 (45). Since

the C-terminal regions of H2A.Z.1 and H2A.Z.2 differ only by the last amino acid, the

antibody used for ChIPmentation recognizes both isoforms. A metagene analysis of all

analyzed promoter windows showed the expected distribution of H2A.Z occupancy in mock

infection with two peaks on both sides of the TSS, corresponding to -1 and +1 nucleosome

positions (**Sup. Fig. 23a**).

Application of RegCFinder to H2A.Z ChIPmentation data generally identified a relative

increase in H2A.Z downstream of the TSS during infection and a relative decrease upstream

of the TSS for pattern (i) and combined pattern (i)+(ii) clusters (**Fig. 4a**, log2 fold-changes for

differential regions shown in **Sup. Fig. 23b**, examples in **Fig 4b**, **Sup. Fig. 24a-g,k,l**).

Metagene analyses showed that this reflected a downstream shift and broadening of +1

nucleosome peaks as well as a relative increase of +1 nucleosome peaks compared to -1

nucleosome peaks (**Fig. 4d**, **Sup. Fig. 23c-i,m,n**). In contrast, cluster 7 (most pronounced

pattern (ii) cluster) showed the opposite trend with relative increases in H2A.Z upstream of

the TSS and decreases downstream of the TSS (example in **Fig 4c**). Consistent with pattern

(ii) representing pattern (i) in antisense direction, the metagene analysis showed an upstream

shift and broadening of the -1 nucleosome peak and a relative increase of the -1 nucleosome

peak compared to the +1 nucleosome peak (**Fig. 4e**). However, this was only observed for

few genes in cluster 6 and cluster 11 (**Fig. 4a**, **Sup. Fig. 23j,k**, examples in **Fig. 4c, Sup. Fig.**

24h,i), which fits with the observation that pattern (ii) in the ATAC-seq data was also much

less pronounced for these clusters. In contrast to pattern (i) and (ii), pattern (iii) in the ATAC-

seq data (cluster 13) was not associated with notable changes in H2A.Z (**Sup. Fig. 23l**,

example in **Sup. Fig. 24j**), indicating that it is not shaped by shifts in +1 and/or -1

nucleosome positioning. This is consistent with pattern (iii) being at least partly associated

with dOCR induction for upstream genes, which we previously linked to impaired

nucleosome repositioning following Pol II transcription downstream of genes (30).

In summary, our data suggest that the alterations in chromatin accessibility around promoters

result from an extension of the nucleosome-free region around promoters due to downstream

and upstream shifts of either +1 or -1 nucleosomes. Here, the direction of shift depends on

whether sense or antisense transcription represents the dominant direction of transcription at

this genome position. Notably, a recent study in yeast proposed that -1 nucleosomes should

actually be considered +1 nucleosomes for antisense transcription (46). We hypothesized that

– similar to what was previously observed for yeast – the loss of Pol II from host promoters

during HSV-1 infection induced the shift of +1 and -1 nucleosome positions. To confirm that

inhibition of transcription alone can lead to similar shifts of +1 and -1 nucleosomes in human

cells, we re-analyzed published H2A.Z ChIP-seq data of HCT116 cells with and without α-

amanitin treatment from the study by Lashgari *et al.* (47). They previously showed that α-

amanitin treatment leads to reduced Pol II levels at gene TSSs and increased incorporation of

H2A.Z at TSSs of transcribed genes. However, they only analyzed changes in absolute H2A.Z

levels at TSS promoters and did not focus on changes in +1 and -1 nucleosome positioning.

Metagene and RegCFinder analyses of the H2A.Z ChIP-seq data for our clusters indeed

showed a similar trend upon α-amanitin treatment as during HSV-1 infection (**Fig. 4f,g, Sup.**

Fig. 25, examples are shown in **Sup. Fig. 26**). Pattern (i) and pattern (i)+(ii) clusters showed a

massive broadening of the +1 nucleosome peak into downstream regions, even more

pronounced than what is observed upon HSV-1 infection. In addition, they also showed a less

pronounced broadening of -1 nucleosome peaks. Similarly, pattern (ii) clusters showed a

strong broadening of both -1 and +1 nucleosome peaks into upstream regions. In case of

cluster 7 (strongest pattern (ii)), the broadening was much more pronounced for the -1

nucleosome (i.e. the +1 nucleosome in antisense direction) than for the +1 nucleosome (i.e.

the -1 nucleosome in antisense direction). This analysis confirms that depletion of Pol II from

promoters leads to a downstream shift of +1 nucleosomes in human cells. Together with the

mitigating effect of PAA treatment on the chromatin accessibility changes in HSV-1 infection

and the timing of onset, our results thus provide strong evidence that the shift of +1 and -1

nucleosomes in the direction of transcription in HSV-1 infection is a consequence of Poll II

depletion from host chromatin.

Discussion

Previously, we showed that HSV-1 infection disrupts chromatin architecture downstream of

genes with strong read-through transcription. In this study, we reveal that chromatin

architecture is also substantially altered at host gene promoters during HSV-1 infection. Here,

57% of genes showed a statistically significant change in the distribution of chromatin

accessibility at promoters in WT HSV-1 infection and metagene analyses indicated similar

but less pronounced changes even for genes without statistically significant changes. In

essence, we identified three types of changes: Most genes showed a shift of the TSS peak

and/or broadening of chromatin accessibility into regions downstream of the TSS (pattern (i)),

while ~900 genes showed the opposite trend with a shift of the TSS peak and/or broadening

of chromatin accessibility into regions upstream of the TSS (pattern (ii)). A further ~460

genes exhibited a combination of pattern (i) and (ii), with the broadening of accessible

chromatin in downstream direction being more pronounced than in upstream direction. Only

a small set of genes (126 genes) showed relative increases in chromatin accessibility up- and

downstream of the TSS, which could partly be attributed to overlaps with dOCRs from

upstream genes and partly to an enrichment for (short) non-coding RNAs for which promoter

windows were larger than the actual gene length. We thus did not further investigate this

pattern.

As patterns (i) and (ii) were still observed in absence of ICP22, which is necessary for dOCR

induction, these are not artefacts of dOCRs extending into downstream genes. On the

contrary, PAA treatment, which increases dOCR induction, substantially reduced – though

not completely abolished – pattern (i) and (ii). In contrast, knockdown of neither ICP0,

ICP22, ICP27 nor *vhs* substantially affected pattern (i) and (ii). This led us to hypothesize that

the loss of Pol II from host genes during HSV-1 infection, which is alleviated by PAA, causes

the widespread extension of open chromatin at host gene promoters. Analysis of

transcriptional activity using RNA-seq of chromatin-associated RNA revealed that pattern (i)

and (ii) are indeed linked to transcription, with pattern (i) associated with more highly

expressed genes and pattern (ii) associated with genes with strong antisense transcription on

the opposite strand. In case of cluster 7, for which pattern (ii) was most pronounced, antisense

transcription was indeed much higher than sense transcription. This was not due to the

widespread induction of antisense transcription we previously reported in HSV-1 infection

(48), but these genes already exhibited strong transcription in antisense direction in uninfected

cells originating from overlapping genes on opposite strands or bidirectional promoters. Thus,

pattern (ii) is essentially just pattern (i) for the genes on the antisense strand.

Since the broadening of chromatin accessibility in promoter regions suggested an extension of

nucleosome-free regions at promoters, we mapped +1 and -1 nucleosome positions by

ChIPmentation of the H2A.Z histone variant enriched at these nucleosomes. Analysis of

H2A.Z occupancy indeed showed a downstream shift of +1 nucleosomes for pattern (i) genes

in HSV-1 infection and an upstream shift of -1 nucleosomes for genes with most pronounced

pattern (ii) (cluster 7). This further confirmed that pattern (ii) just represents pattern (i) for

genes on the antisense strand. It is also consistent with recent results from yeast by Bagchi *et*

al. that -1 H2A.Z-containing nucleosomes should actually be considered as +1 nucleosomes

for antisense transcription (46). Downstream shifts of +1 nucleosomes have previously been

reported upon Pol II degradation in yeast by Weiner *et al.* (34), in particular for highly

expressed genes. Re-analysis of published H2A.Z ChIP-seq data with and without α-amanitin-

induced depletion of Pol II from gene promoters confirmed that loss of Pol II alone is

sufficient to induce downstream shifts of +1 nucleosomes in human cells. Moreover, for

genes with dominant antisense transcription (cluster 7), it resulted in upstream shifts of -1

nucleosomes. Here, the broadening of H2A.Z peaks in both HSV-1 infection and upon α-

amanitin treatment, respectively, indicate a less precise positioning of corresponding

nucleosomes. This was more pronounced upon α-amanitin treatment and also observed for -1

nucleosomes, suggesting that loss of Pol II upon α-amanitin treatment is more complete than

in HSV-1 infection.

The +1 nucleosome has been shown to play a role in promoter-proximal Pol II pausing,

which occurs between the promoter and the +1 nucleosome (49). It is thus tempting to

speculate that the downstream shifts in +1 nucleosomes we observed in this study may

provide an explanation for the downstream shift in Pol II pausing in HSV-1 infection we

recently reported (39). Notably, there are a few parallels between the downstream shifts of +1

nucleosomes and changes in Pol II pausing during HSV-1 infection. Both appear to be

established relatively early in infection, with the extension of the nucleosome-free region

detectable at 4 h p.i. but not at 2 h p.i. in our ATAC-seq data and the downstream shift of Pol

II pausing clearly detectable in PRO-seq data at 3 h p.i. but not yet at 1.5 h p.i. (49).

Furthermore, both +1 nucleosomes and Pol II pausing sites, which are well-positioned in

uninfected cells, appear much less well-positioned in HSV-1 infection. In this way, both

phenomena would largely be a bystander effect of the loss of host transcriptional activity and

not necessarily play a functional role for HSV-1 infection. Alternatively, the shifts in +1

nucleosomes might also be linked to the mobilization of H1, H2 (including H2A.Z) and H4

histones during HSV-1 infection (50-52), which has been proposed to serve as a source of

histones for viral chromatin assembly. In any case, our study highlights that HSV-1 infection

impacts chromatin architecture at promoters independently of the widespread changes

downstream of genes mediated by ICP22.

Materials and Methods

Previously published sequencing data analyzed in this study

ATAC-seq for mock and WT HSV-1 infection (strain 17, 8 h p.i.), WT infection with 8 h

PAA treatment, infection with ICP0-null mutant (ΔICP0, strain 17, (53), 12 h p.i.), ICP22-

null mutant (ΔICP22, R325, strain F, (54), 12 h p.i.), ICP27-null mutant (ΔICP27, strain

KOS, (55), 8 h p.i.) and *vhs*-null mutant (Δvhs, strain 17, (56)) in of HFF were taken from our

recent publication (30) (n=2 apart from ΔICP22 infection with n=4, GEO accession:

GSE185234). ATAC-seq data for mock and WT HSV-1 infection of HFF at 1, 2, 4, 6 and 8 h

p.i. WT infection (n=2 replicates, GEO accession: GSE100611) and chromatin-associated

RNA-seq data for mock and 8 h p.i. WT infection (GEO accession: GSE100576) were taken

from our previous publication (22). H3K36me3 ChIPmentation data for mock and 8 h p.i. WT

infection of HFF were also taken from our recent publication (30) (n=3, GEO accession:

GSE185246). H2A.Z ChIP-seq data of untreated and α-amanitin-treated HCT116 cells were

taken from the study by Lashgari *et al.* (47) (n=2, GSE101427).

535
ChIPmentation, library preparation and sequencing

Two days prior to infection, two million HFF cells were seeded in 15 cm dishes. At the day of

infection, cells had expanded to ~80% confluency. Cells were infected with the respective

viruses as described in results section (n=3). At 8 p.i., cells were fixed for 10 minutes at room

temperature by adding 1% formaldehyde (final) directly to the medium. Cells were scraped in

1mL of ice-cold 1x PBS containing protease inhibitor cocktail (1x) (Roche #11836153001)

with an additional 1mM phenylmethylsulfonyl fluoride (PMSF). Cells were pelleted at 1500

rpm for 20 min at 4 °C. Supernatant was aspirated and cell pellets were frozen in liquid N_2.

Cell pellets were resuspended in 1.5 mL 0.25% [w/v] SDS sonication buffer (10 mM Tris

pH=8.0, 0.25% [w/v] SDS, 2 mM EDTA) with 1x protease inhibitors and 1 mM additional

PMSF and incubated on ice for 10 minutes. Cells were sonicated in fifteen 1 minute intervals,

25% amplitude, with Branson Ultrasonics SonifierTM S-450 until most fragments were in the

range of 200-700 bp as determined by agarose gel electrophoresis. Two million cells used for

the preparation of the ChIPmentation libraries were diluted 1:1.5 with equilibration buffer (10

mM Tris, 233 mM NaCl, 1.66% [v/v] Triton X-100, 0.166% [w/v] sodium deoxycholate, 1

mM EDTA, protease inhibitors) and spun at 14,000x g for 10 minutes at 4 °C to pellet

insoluble material. Supernatant was transferred to a new 1.5 mL screw-cap tube and topped

up with RIPA-LS (10 mM Tris-HCl pH 8.0, 140 mM NaCl, 1 mM EDTA pH 8.0, 0.1% [w/v]

SDS, 0.1% [w/v] sodium deoxycholate, 1% [v/v] Triton X-100, protease inhibitors) to 200

μL. Input and gel samples were preserved. Lysates were incubated with 1μg/IP of anti-H2A.Z

antibody (Diagenode, #C15410201) on a rotator overnight at 4 °C.

Dependent on the added amount of antibody, the amount of Protein A magnetic beads

(ThermoFisher Scientific #10001D) was adjusted (e.g. for 1-2 μg of antibody/IP = 15 μL of

beads) and blocked overnight with 0.1% [w/v] bovine serum albumin in RIPA buffer. On the

following day, beads were added to the IP samples for 2 h on a rotator at 4 °C to capture the

antibody-bound fragments. The immunoprecipitated chromatin was subsequently washed

twice with 150 μL each of ice-cold buffers RIPA-LS, RIPA-HS (10 mM Tris-HCl pH 8.0, 50

0mM NaCl, 1 mM EDTA pH 8.0, 0.1% [w/v] SDS, 0.1% [v/v] sodium deoxycholate, 1%

[v/v] Triton X-100), RIPA-LiCl (10 mM Tris-HCl pH 8.0, 250 mM LiCl, 1 mM EDTA pH

8.0, 0.5% [w/v] sodium deoxycholate, 0.5% [v/v] Nonidet P-40) and 10 mM Tris pH 8.0

containing protease inhibitors. Beads were washed once more with ice-cold 10 mM Tris pH

8.0 lacking inhibitors and transferred into new tubes.

Beads were resuspended in 25 μL of the tagmentation reaction mix (Nextera DNA Sample

Prep Kit, Illumina) containing 5 μL of 5x Tagmentation buffer, 1 μL of Tagment DNA

enzyme, topped up with H_2O to the final volume and incubated at 37 °C for 10 minutes in a

thermocycler. Beads were mixed after 5 minutes by gentle pipetting. To inactivate the Tn5

enzyme, 150 μL of ice-cold RIPA-LS was added to the tagmentation reaction. Beads were

washed twice with 150 μL of RIPA-LS and 1x Tris-EDTA and subjected to de-crosslinking

by adding 100 μL ChIPmentation elution buffer (160 mM NaCl, 40 μg/mL RNase A (Sigma-

Aldrich #R4642), 1x Tris-EDTA (Sigma #T9285) and incubating for 1h at 37 °C followed by

overnight shaking at 65 °C. The next day, 4 mM EDTA and 200 μg/mL Proteinase K (Roche,

#03115828001) were added, and samples incubated for another 2h at 45 °C with 1000 rpm

shaking. Supernatant was transferred into a new tube and another 100 μL of ChIPmentation

elution buffer was added for another hour at 45 °C with 1000 rpm shaking. DNA was isolated

with MinElute PCR Purification Kit (Qiagen #28004) and eluted in 21 μL of H_2O.

DNA for the final library was prepared with 25 μL NEBNext Ultra II Q5 Master Mix, 3.75 μL

IDT custom primer i5_n_x (10 μM); 3.75 μL IDT custom primer i7_n_x (10 μM); 3.75 μL

H_2O and 13.75 μL ChIPmentation DNA. The Cq value obtained from the library

quantification, rounded up to the nearest integer plus one additional cycle, was used to

amplify the rest of the ChIPmentation DNA. Library qualities were verified by High

Sensitivity DNA Analysis on the Bioanalyzer 2100 (Agilent) before performing sequencing

on NextSeq 500 (paired-end 35bp reads) at the Core Unit Systemmedizin, Würzburg,

Germany. All samples were sequenced at equimolar ratios.

Read alignment

The read alignment pipeline was implemented and run in the workflow management system

Watchdog (57, 58) as already previously described (39). Public sequencing data were

downloaded from SRA using the sratoolkit version 2.10.8. Sequencing reads were aligned

against the human genome (GRCh37/hg19), the HSV-1 genome (Human herpesvirus 1 strain

17, GenBank accession code: JN555585) and human rRNA sequences using ContextMap2

version 2.7.9 (59) (using BWA as short read aligner (60) and allowing a maximum indel size

of 3 and at most 5 mismatches). For the two repeat regions in the HSV-1 genome, only one

copy was retained each, excluding nucleotides 1–9,213 and 145,590–152,222 from the

alignment. SAM output files of ContextMap2 were converted to BAM files using samtools

(61). Read coverage in bedGraph format was calculated from BAM files using BEDTools

(62). Subregions of promoter windows (TSS ± 3 kb) with differential read coverage in

ATAC-seq and H2A.Z ChIPmentation/-seq data were determined with RegCFinder (37). For

this purpose, RegCFinder was applied to ATAC-seq data for all pairwise comparisons of

mock to WT and null mutant infections as well as mock to each timepoint of infection for the

time-course data and to H2A.Z ChIPmentation and ChIP-seq data for the comparison of mock

and WT infection as well as untreated and α-amanitin-treated HCT116 cells.

Data plotting and statistical analysis

All figures were created in R and all statistical analyses were performed in R (63). Read

coverage plots were created using the R Bioconductor package Gviz (64).

Metagene and clustering analysis

Metagene analyses were performed as previously described (65) using the R program

developed for this previous publication (available with the Watchdog *binGenome* module in

the   Watchdog   module   repository   (https://github.com/watchdog-wms/watchdog-wms-

modules/). For promoter region analyses, the regions −3 kb to +3 kb of the TSS were divided

into 101 equal-sized bins for each gene. For each bin, the average coverage per genome

position was calculated and bin read coverages were then normalized by dividing by the total

sum of all bins. Metagene curves for each replicate were created by averaging results for

corresponding bins across all genes in a cluster/group and then averaged across replicates. For

hierarchical clustering analysis, RegCFinder profiles of differential regions were calculated

for each promoter window and comparison by setting each position in an m-RegC to 1, in an

i-RegC to -1 and all other positions to 0. RegCFinder profiles of each promoter window for

each comparison were concatenated into one row in the matrix. Hierarchical clustering of the

resulting matrix was then performed using the hclust function in R according to Euclidean

distances and Ward's clustering criterion.

Motif discovery and enrichment analysis

Motif discovery was performed for each cluster separately for the i- and m-RegC regions

using findMotifsGenome.pl script of the Homer suite (40), with our 7,649 input promoter

windows as background. For this purpose, we used the hg19 annotation provided by Homer

and automated trimming of input windows was disabled. Significant motifs were identified at

a q-value (=False Discovery Rate (FDR) calculated with the Benjamini-Hochberg method

(66)) cutoff of 0.01. Over-representation of Gene Ontology (GO) terms was performed

separately for each cluster using the g:Profiler webserver (67) and the R package gprofiler2

(68), which provides an R interface to the webserver. As background gene list, the genes

corresponding to our 7,649 input promoter windows were provided. P-values were corrected

for multiple testing using the Benjamini-Hochberg method (66) and significantly over-

represented GO terms were identified at an multiple testing adjusted p-value cutoff of 0.001.

Enrichment of gene types (obtained from the Ensembl annotation (version 87 for GRCh37))

within clusters was determined using one-sided Fisher's exact tests (with alternative =

greater). P-values were again corrected for multiple testing using the Benjamini-Hochberg

method and an adjusted p-value cutoff of 0.001 was used.

Analysis of downstream open chromatin regions (dOCRs)

Open chromatin regions (OCRs) were determined from ATAC-seq data by first converting

BAM files with mapped reads to BED format using BEDTools (69) and then determining

enriched regions from these BED files using F-Seq with default parameters (70). dOCRs for

individual genes were calculated from OCRs as previously described (22, 30). In brief,

dOCRs are determined for genes by first assigning all OCRs overlapping with the 10 kb

downstream of a gene to this gene. Second, OCRs starting at most 5 kb downstream of the so

far most downstream OCR of a gene are also assigned to this gene. In both steps, individual

OCRs can be assigned to multiple genes. The second step is iterated until no more OCRs can

be assigned. The dOCR of a gene is then defined as the region from the gene 3'end to the end

of the most downstream OCR assigned to this gene.

Gene expression analysis

Number of fragments (=read pairs) per gene were determined from mapped paired-end RNA-

seq reads in a strand-specific manner using featureCounts (71) and gene annotations from

Ensembl (version 87 for GRCh37). For genes, all fragments overlapping exonic regions on

the corresponding strand by $\geq$ 25bp were counted for the corresponding gene. Fold-changes in

gene expression and statistical significance of changes were determined using DESeq2 (72)

and p-values were adjusted for multiple testing using the method by Benjamini and Hochberg

(66). Gene expression was quantified in terms of fragments per kilobase of exons per million

mapped reads (FPKM). Only reads mapped to the human genome were counted for the total

number of mapped reads for FPKM calculation.

672
Acknowledgements

This work was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research

Foundation) in the framework of the Research Unit FOR5200 DEEP-DV (443644894)

projects FR 2938/11-1 to C.C.F. and DO 1275/12-1 to L.D.

Competing interests

The authors declare no competing interests.

References

1.   Roizman B KD, RJ W. Herpes Simplex Viruses. In: Knipe DM HP, editor. Fields

Virology. 5th. Philadelphia: Lippincott Williams & Wilkins; 2007. p. 2501-601.

2.   Pellet PE RB. Herpesviridae. In: Knipe DM HP, editor. Fields Virology. 6th.

Philadelphia: Wolters Kluwer Health/Lippincott Williams & Wilkins; 2013. p. 1802-2128.

3.   Ablashi D, Agut H, Alvarez-Lafuente R, Clark DA, Dewhurst S, DiLuca D, et al.

Classification of HHV-6A and HHV-6B as distinct viruses. Arch Virol. 2014;159(5):863-70.

4.   Korr G, Thamm M, Czogiel I, Poethko-Mueller C, Bremer V, Jansen K. Decreasing

seroprevalence of herpes simplex virus type 1 and type 2 in Germany leaves many people

susceptible to genital infection: time to raise awareness and enhance control. BMC Infect Dis.

2017;17(1):471.

5.   Whitley   RJ,   Roizman   B.   Herpes   simplex   virus   infections.   Lancet.

2001;357(9267):1513-8.

694 6. Whisnant AW, Jürges CS, Hennig T, Wyler E, Prusty B, Rutkowski AJ, et al.

695 Integrative functional genomics decodes herpes simplex virus 1. Nat Commun.

696 2020;11(1):2038.

697 7. Spencer CA, Dahmus ME, Rice SA. Repression of host RNA polymerase II

698 transcription by herpes simplex virus type 1. J Virol. 1997;71(3):2031-40.

699 8. Rice SA, Davido DJ. HSV-1 ICP22: hijacking host nuclear functions to enhance viral

700 infection. Future Microbiol. 2013;8(3):311-21.

701 9. Abrisch RG, Eidem TM, Yakovchuk P, Kugel JF, Goodrich JA. Infection by Herpes

702 Simplex Virus 1 Causes Near-Complete Loss of RNA Polymerase II Occupancy on the Host

703 Cell Genome. J Virol. 2015;90(5):2503-13.

704 10. Dembowski JA, DeLuca NA. Selective recruitment of nuclear factors to productively

705 replicating herpes simplex virus genomes. PLoS Pathog. 2015;11(5):e1004939.

706 11. Birkenheuer CH, Danko CG, Baines JD. Herpes Simplex Virus 1 Dramatically Alters

707 Loading and Positioning of RNA Polymerase II on Host Genes Early in Infection. J Virol.

708 2018;92(8).

709 12. Rivas HG, Schmaling SK, Gaglia MM. Shutoff of Host Gene Expression in Influenza

710 A Virus and Herpesviruses: Similar Mechanisms and Common Themes. Viruses.

711 2016;8(4):102.

712 13. Dremel SE, DeLuca NA. Herpes simplex viral nucleoprotein creates a competitive

713 transcriptional environment facilitating robust viral transcription and host shut off. Elife.

714 2019;8.

715 14. McSwiggen DT, Hansen AS, Teves SS, Marie-Nelly H, Hao Y, Heckert AB, et al.

716 Evidence for DNA-mediated nuclear compartmentalization distinct from phase separation.

717 eLife. 2019;8:e47098.

15.   Dremel SE, DeLuca NA. Herpes simplex viral nucleoprotein creates a competitive

transcriptional environment facilitating robust viral transcription and host shut off. eLife.

2019;8:e51109.

16.   Isa NF, Bensaude O, Aziz NC, Murphy S. HSV-1 ICP22 Is a Selective Viral

Repressor of Cellular RNA Polymerase II-Mediated Transcription Elongation. Vaccines.

2021;9(10):1054.

17.   Kwong AD, Frenkel N. Herpes simplex virus-infected cells contain a function(s) that

destabilizes both host and viral mRNAs. Proc Natl Acad Sci U S A. 1987;84(7):1926-30.

18.   Oroskar AA, Read GS. Control of mRNA stability by the virion host shutoff function

of herpes simplex virus. J Virol. 1989;63(5):1897-906.

19.   Feng P, Everly DN, Jr., Read GS. mRNA decay during herpesvirus infections:

interaction between a putative viral nuclease and a cellular translation factor. J Virol.

2001;75(21):10272-80.

20.   Friedel CC, Whisnant AW, Djakovic L, Rutkowski AJ, Friedl MS, Kluge M, et al.

Dissecting Herpes Simplex Virus 1-Induced Host Shutoff at the RNA Level. J Virol.

2021;95(3).

21.   Rutkowski AJ, Erhard F, L'Hernault A, Bonfert T, Schilhabel M, Crump C, et al.

Widespread disruption of host transcription termination in HSV-1 infection. Nat Commun.

2015;6:7126.

22.   Hennig T, Michalski M, Rutkowski AJ, Djakovic L, Whisnant AW, Friedl MS, et al.

HSV-1-induced disruption of transcription termination resembles a cellular stress response

but selectively increases chromatin accessibility downstream of genes. PLoS Pathog.

2018;14(3):e1006954.

23.   Wang X, Hennig T, Whisnant AW, Erhard F, Prusty BK, Friedel CC, et al. Herpes

simplex virus blocks host transcription termination via the bimodal activities of ICP27. Nat

Commun. 2020;11(1):293.

24.   Vilborg A, Passarelli MC, Yario TA, Tycowski KT, Steitz JA. Widespread Inducible

Transcription Downstream of Human Genes. Mol Cell. 2015;59(3):449-61.

25.   Rosa-Mercado NA, Zimmer JT, Apostolidi M, Rinehart J, Simon MD, Steitz JA.

Hyperosmotic stress alters the RNA polymerase II interactome and induces readthrough

transcription despite widespread transcriptional repression. Mol Cell. 2021;81(3):502-13 e4.

26.   Hadar S, Meller A, Saida N, Shalgi R. Stress-induced transcriptional readthrough into

neighboring genes is linked to intron retention. iScience. 2022;25(12):105543.

27.   Wu N, Watkins SC, Schaffer PA, DeLuca NA. Prolonged gene expression and cell

survival after infection by a herpes simplex virus mutant defective in the immediate-early

genes encoding ICP4, ICP27, and ICP22. Journal of Virology. 1996;70(9):6358-69.

28.   Honess RW, Watson DH. Herpes simplex virus resistance and sensitivity to

phosphonoacetic acid. Journal of Virology. 1977;21(2):584-600.

29.   Becker Y, Asher Y, Cohen Y, Weinberg-Zahlering E, Shlomai J. Phosphonoacetic

Acid-Resistant Mutants of Herpes Simplex Virus: Effect of Phosphonoacetic Acid on Virus

Replication and In Vitro Deoxyribonucleic Acid Synthesis in Isolated Nuclei. Antimicrobial

Agents and Chemotherapy. 1977;11(5):919-22.

30.   Djakovic L, Hennig T, Reinisch K, Milic A, Whisnant AW, Wolf K, et al. The HSV-1

ICP22 protein selectively impairs histone repositioning upon Pol II transcription downstream

of genes. Nat Commun. 2023;14(1):4591.

31.   Barski A, Cuddapah S, Cui K, Roh TY, Schones DE, Wang Z, et al. High-resolution

profiling of histone methylations in the human genome. Cell. 2007;129(4):823-37.

32.   Raisner RM, Hartley PD, Meneghini MD, Bao MZ, Liu CL, Schreiber SL, et al.

Histone variant H2A.Z marks the 5' ends of both active and inactive genes in euchromatin.

Cell. 2005;123(2):233-48.

33.   Bensaude O. Inhibiting eukaryotic transcription. Which compound to choose? How to

evaluate its activity? Transcription. 2011;2(3):103-8.

34.   Weiner A, Hughes A, Yassour M, Rando OJ, Friedman N. High-resolution

nucleosome mapping reveals transcription-dependent promoter packaging. Genome Res.

2010;20(1):90-100.

35.   Schwabish MA SK. Evidence for eviction and rapid deposition of histones upon

transcriptional elongation by RNA polymerase II. Mol Cell Biol. 2004;24(23):10111-7.

36.   Kaplan N MI, Fondufe-Mittendorf Y, Gossett AJ, Tillo D, Field Y, LeProust EM,

Hughes TR, Lieb JD, Widom J, Segal E. The DNA-encoded nucleosome organization of a

eukaryotic genome. Nature. 2008;458(7236):362-6.

37.   Weiss E, Friedel CC. RegCFinder: targeted discovery of genomic subregions with

differential read density. Bioinform Adv. 2023;3(1):vbad085.

38.   Anders S RA, Huber W. Detecting differential usage of exons from RNA-seq data.

Genome Research. 2012;22(10):2008-17.

39.   Weiss E, Hennig T, Grassl P, Djakovic L, Whisnant AW, Jurges CS, et al. HSV-1

Infection Induces a Downstream Shift of Promoter-Proximal Pausing for Host Genes. J Virol.

2023;97(5):e0038123.

40.   Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, et al. Simple combinations

of lineage-determining transcription factors prime cis-regulatory elements required for

macrophage and B cell identities. Mol Cell. 2010;38(4):576-89.

41.   Barski A, Cuddapah S, Cui K, Roh T-Y, Schones DE, Wang Z, et al. High-Resolution

Profiling of Histone Methylations in the Human Genome. Cell. 2007;129(4):823-37.

42.   Shivaswamy S, Bhinge A, Zhao Y, Jones S, Hirst M, Iyer VR. Dynamic remodeling of

individual nucleosomes across a eukaryotic genome in response to transcriptional

perturbation. PLoS Biol. 2008;6(3):e65.

43.   Tirosh I, Barkai N. Two strategies for gene regulation by promoter nucleosomes.

Genome Res. 2008;18(7):1084-91.

44.   Hartley PD, Madhani HD. Mechanisms that specify promoter nucleosome location and

identity. Cell. 2009;137(3):445-58.

45.   Greenberg RS, Long HK, Swigut T, Wysocka J. Single Amino Acid Change Underlies

Distinct Roles of H2A.Z Subtypes in Human Syndrome. Cell. 2019;178(6):1421-36 e24.

46.   Bagchi DN, Battenhouse AM, Park D, Iyer VR. The histone variant H2A.Z in yeast is

almost exclusively incorporated into the +1 nucleosome in the direction of transcription.

Nucleic Acids Research. 2020;48(1):157-70.

47.   Lashgari A, Millau J-F, Jacques P-É, Gaudreau L. Global inhibition of transcription

causes an increase in histone H2A.Z incorporation within gene bodies. Nucleic Acids

Research. 2017;45(22):12715-22.

48.   Wyler E, Menegatti J, Franke V, Kocks C, Boltengagen A, Hennig T, et al.

Widespread activation of antisense transcription of the host genome during herpes simplex

virus 1 infection. Genome Biology. 2017;18(1):209.

49.   Jimeno-González S, Ceballos-Chávez M, Reyes JC. A positioned +1 nucleosome

enhances promoter-proximal pausing. Nucleic Acids Research. 2015;43(6):3068-78.

50.   Conn KL HM, Schang LM. Linker histones are mobilized during infection with herpes

simplex virus type 1. J Virol. 2008;82:8629–46.

51.   Conn KL HM, Schang LM. Core histones H2B and H4 are mobilized during infection

with herpes simplex virus 1. J Virol. 2011;85:13234–52.

52.   Conn KL, Schang LM. Chromatin Dynamics during Lytic Infection with Herpes

Simplex Virus 1. Viruses. 2013;5(7):1758-86.

53.   Stow ND, Stow EC. Isolation and Characterization of a Herpes Simplex Virus Type 1

Mutant Containing a Deletion within the Gene Encoding the Immediate Early Polypeptide

Vmw110. Journal of General Virology. 1986;67(12):2571-85.

54. Post LE, Roizman B. A generalized technique for deletion of specific genes in large

genomes: a gene 22 of herpes simplex virus 1 is not essential for growth. Cell.

1981;25(1):227-32.

55. Smith IL, Hardwicke MA, Sandri-Goldin RM. Evidence that the herpes simplex virus

immediate early protein ICP27 acts post-transcriptionally during infection to regulate gene

expression. Virology. 1992;186(1):74-86.

56. Fenwick ML, Everett RD. Inactivation of the Shutoff Gene (UL41) of Herpes Simplex

Virus Types 1 and 2. Journal of General Virology. 1990;71(12):2961-7.

57. Kluge M, Friedel CC. Watchdog - a workflow management system for the distributed

analysis of large-scale experimental data. BMC Bioinformatics. 2018;19(1):97.

58. Kluge M, Friedl MS, Menzel AL, Friedel CC. Watchdog 2.0: New developments for

reusability, reproducibility, and workflow execution. Gigascience. 2020;9(6).

59. Bonfert T, Kirner E, Csaba G, Zimmer R, Friedel CC. ContextMap 2: fast and

accurate context-based RNA-seq mapping. BMC Bioinformatics. 2015;16:122.

60. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler

transform. Bioinformatics. 2009;25(14):1754-60.

61. Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, et al. Twelve

years of SAMtools and BCFtools. Gigascience. 2021;10(2).

62. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic

features. Bioinformatics. 2010;26(6):841-2.

63. R Core Team. R: A Language and Environment for Statistical Computing. Vienna,

Austria2022.

64. Hahne F, Ivanek R. Visualizing Genomic Data Using Gviz and Bioconductor.

Methods Mol Biol. 2016;1418:335-51.

65. Chirackal Manavalan AP, Pilarova K, Kluge M, Bartholomeeusen K, Rajecky M,

Oppelt J, et al. CDK12 controls G1/S progression by regulating RNAPII processivity at core

DNA replication genes. EMBO Rep. 2019;20(9):e47592.

66. Benjamini Y HY. Controlling the False Discovery Rate: A Practical and Powerful

Approach to Multiple Testing. Journal of the Royal Statistical Society Series B

(Methodological). 1995;57(1):289-300.

67. Raudvere U, Kolberg L, Kuzmin I, Arak T, Adler P, Peterson H, et al. g:Profiler: a

web server for functional enrichment analysis and conversions of gene lists (2019 update).

Nucleic Acids Res. 2019;47(W1):W191-W8.

68. Kolberg L, Raudvere U, Kuzmin I, Vilo J, Peterson H. gprofiler2 -- an R package for

gene list functional enrichment analysis and namespace conversion toolset g:Profiler.

F1000Res. 2020;9.

69. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic

features. Bioinformatics. 2010;26(6):841-2.

70. Boyle AP, Guinney J, Crawford GE, Furey TS. F-Seq: a feature density estimator for

high-throughput sequence tags. Bioinformatics. 2008;24(21):2537-8.

71. Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for

assigning sequence reads to genomic features. Bioinformatics. 2014;30(7):923-30.

72. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. Genome Biol. 2014;15(12):550.

Figures

Fig. 1: HSV-1 infection impacts chromatin accessibility around promoters for the

majority of host genes.

(a) Heatmap illustrating the location of differential regions (m- and i-RegC) identified by

RegCFinder on the ATAC-seq data for WT, ΔICP0, ΔICP22, ΔICP27, and Δ*vhs* infection in

comparison to mock infection. Results are shown for 4,981 (out of 7,649) promoter windows

containing at least one statistically significant (adj. p. ≤ 0.01) differential region for at least

one virus infection compared to mock. Red and blue colors represent m-RegC

(mock>infection) and i-RegC (infection>mock) locations, respectively, within promoter

windows. Here, each row of the heatmap shows results for the same input window in different

comparisons of virus infection to mock. Colored rectangles on top of the heatmap indicate

which virus infection was compared to mock. Black vertical lines in the center of each part of

the heatmap indicate the position of the TSS. Hierarchical clustering was performed according

to Euclidean distances and Ward's clustering criterion and the cutoff on the hierarchical

clustering dendrogram was selected to obtain 14 clusters (marked by colored rectangles

between dendrogram and heatmap and numbered from top to bottom as indicated). **(b)**

Metagene plot showing the average ATAC-seq profile around the TSS ± 3 kb in mock (red)

and WT (blue) infection for cluster 5, an example for pattern (i). For a description of

metagene plots, see Materials and Methods. The colored bands below the metagene curves in

each panel indicate the percentage of genes having an i- or m-RegC (blue or red, respectively)

at that position. **(c)** Read coverage in a ±3.6 kb window around the TSS in ATAC-seq data of

mock (red) and WT (blue) infection for an example gene (FBXO28) with pattern (i). Read

coverage was normalized to total number of mapped reads for each sample and averaged

between replicates. The TSS is indicated by a short vertical line below the read coverage track

for mock infection. Gene annotation is indicated at the top. Boxes represent exons, lines

represent introns, and direction of transcription is indicated by arrowheads. Below the read

coverage track for WT infection, m-RegC (red bars) and i-RegC (blue bars) are shown for the

comparison of WT vs. mock infection. **(d)** Metagene plot as in **(b)** for cluster 7, the most

pronounced case of pattern (ii). **(e)** Read coverage plot as in **(c)** for an example gene

(ARHGAP1) with pattern (ii). Here, the promoter window also contains the TSS of the

ZNF408 gene on the opposite strand and the accessible chromatin region extends upstream of

the ARHGAP1 TSS, i.e. downstream of the ZNF408 TSS. **(f,g)** Metagene plots as in **(b)** for

(f) cluster 13, which exhibits pattern (iii), and **(g)** cluster 8, one of two clusters exhibiting the

combined (i)+(ii) pattern.

Fig. 2: Changes in chromatin accessibility at promoters are mostly independent of

dOCR induction and linked to transcription.

(a) Bar plots showing the percentage of promoter regions for each cluster overlapping with

dOCRs downstream of the 1,296 genes for which we previously showed consistent dOCR

induction across different HSV-1 strains (30). dOCR regions in mock, WT (± PAA treatment)

and null mutant infection were calculated as previous described (30) (see Materials and

Methods for further details) and the overlaps of promoter windows to dOCR regions

originating from an upstream gene were determined for each cluster. WT (± PAA treatment)

and null mutant infection are ordered according to the overall extent of dOCR induction. Only

few clusters (12-14) showed some enrichment for dOCRs from upstream genes, which was

abolished in ΔICP22 infection. **(b-d)** Metagene curves showing average ATAC-seq profiles in

promoter windows for mock (red), WT (blue) and WT+PAA (green) infection for clusters 5

(b), 7 **(c),** and 13 **(d)**. See Materials and Methods for a detailed description of metagene plots.

PAA treatment reduces the downstream and upstream broadening for patterns (i) (example

cluster in **b**) and (ii) (example cluster in **c**), respectively, but enhances pattern (iii) (in **d**),

which is enriched for dOCR induction from upstream genes. **(e,f)** Boxplots showing the

distribution of **(e)** gene expression (FPKM) values and **(f)** log2 DSR:UAR ratios in

chromatin-associated RNA in mock (red) and 8 h p.i. WT (blue) HSV-1 infection for all

clusters (grouped by pattern, cluster numbers are shown below each boxplot) and remaining

genes without significant chromatin accessibility changes (NA group). DSR was calculated as

the expression (FPKM) in chromatin-associated RNA for the region downstream of the TSS

in sense direction and UAR as the expression (FPKM) in the upstream region in antisense

direction. P-values for Wilcoxon rank sum tests comparing values in mock infection for each

cluster/group against all other analyzed genes are indicated below cluster numbers and were

corrected for multiple testing using the Bonferroni method.

Fig. 3: Changes in chromatin accessibility at promoters are detectable as early as 4 h p.i.

(a) Heatmap showing the log2 fold-changes determined with DEXSeq on the ATAC-seq

time-course data (1, 2, 4, 6 and 8 h p.i. WT infection compared to mock infection) for

statistically significant differential regions (m- and i-RegC) identified by RegCFinder. For

comparison, log2 fold-changes for the WT vs. mock comparison from **Fig 1a** are also shown.

Each row represents the results for one of the input windows included in **Fig. 1a**. Colored

rectangles on top indicate the time-point of infection or whether the original WT vs. mock

comparison is shown. Statistically significant differential regions are colored according to the

log2 fold-change determined by DEXSeq. Here, the color scale is continuous between -1 and

1 and all log2 fold-changes >1 are colored the same red and all log2 fold-changes <1 the same

blue. Promoter windows are ordered as in **Fig. 1a** and clusters are annotated as colored and

numbered rectangles on the left. **(b)** Number (no.) of statistically significant differential

regions (m- and i-RegC) and number of genes with at least one statistically significant

differential region identified for each time-point of infection in the ATAC-seq time-course

data. **(c)** Metagene plot of ATAC-seq profiles in mock infection (red) and all time-points of

infection (blue shades) from the ATAC-seq time-course experiment for example cluster 5

(pattern (i)). The colored bands below the metagene curves in each panel indicate the

percentage of genes having an i- or m-RegC (blue or red, respectively) at that position in the

comparison of 8 h p.i. to mock infection from the time-course experiment. Metagene plots for

all clusters are shown in **Sup. Fig. 20**.

Fig. 4: HSV-1 infection and depletion of Pol II from human genes lead to a downstream

shift of H2A.Z-containing +1 nucleosomes.

(a) Heatmap showing the location for differential regions (m- and i-RegC) identified in the

WT vs. mock comparison on the ATAC-seq (left half, identical to left-most part of **Fig. 1a**)

and H2A.Z ChIPmentation data (right half) for the promoter windows included in **Fig. 1a.**

Differential regions with mock>infection (m-RegC) are marked in red, differential regions

with infection>mock (i-RegC) in blue. The order of promoter windows is the same as in **Fig.**

1a and clusters are indicated by colored and numbered rectangles on the left. **(b,c)** Read

coverage in a ±3.6 kb window around the TSS in H2A.Z ChIPmentation data for mock (red)

and WT (blue) infection for example genes with **(b)** pattern (i) (FBXO28) and **(c)** pattern (ii)

(ARHGAP1). For descriptions of read coverage plots see caption to **Fig. 1**. **(d-g)** Metagene

plots of H2A.Z profiles for mock and WT infection **(d,e)** and untreated and α-amanitin-treated

HCT116 cells **(f,g)** for clusters 5 (pattern (i), **d,f**) and 7 (pattern (ii), **e,g**). Metagene plots for

other clusters can be found in **Sup. Fig. 23** and **25**, respectively. The colored bands below the

metagene curves in each panel indicate the percentage of genes having m- or i-RegC at that

position. For **(f,g)**, m-RegC (green) are differential regions with relative read coverage higher

in untreated cells and i-RegC (violet) differential regions with relative read coverage higher

upon α-amanitin-treatment.

a

log2FC mock vs infection

4981 windows, logFC cutoff 1.0

b

c

cluster 5 (n=771)

a

mock>infection
infection>mock

ATAC-seq   H2A.Z

1 2 3 4 5 6 7 8 9 10 11 12 13 14

b FBXO28

mock [0, 0.22]
WT [0, 0.21]

m-RegC
i-RegC

224.299 mb   224.301 mb   224.303 mb   224.305 ml
224.3 mb   224.302 mb   224.304 mb

c ARHGAP1   ZNF408

mock [0, 0.23]
WT [0, 0.21]

m-RegC
i-RegC

46.719 mb   46.721 mb   46.723 mb   46.725 mb
46.72 mb   46.722 mb   46.724 mb

cluster 5 (n=771)

d

m-RegC   i-RegC
<25%   <25%
25−50%   25−50%
50−75%   50−75%
>75%   >75%

mock
WT

H2A.Z profile

−3000   −1500   TSS   1500   3000

cluster 7 (n=245)

e

m-RegC   i-RegC
<25%   <25%
25−50%   25−50%
50−75%   50−75%
>75%   >75%

mock
WT

H2A.Z profile

−3000   −1500   TSS   1500   3000

cluster 5 (n=771)

f

m-RegC   i-RegC
<25%   <25%
25−50%   25−50%
50−75%   50−75%
>75%   >75%

non-treated
α-amanitin

H2A.Z profile

−3000   −1500   TSS   1500   3000

cluster 7 (n=245)

g

m-RegC   i-RegC
<25%   <25%
25−50%   25−50%
50−75%   50−75%
>75%   >75%

non-treated
α-amanitin

H2A.Z profile

−3000   −1500   TSS   1500   3000