Human immunodeficiency virus-1 induces host genomic R-loop and preferentially integrates its genome near the R-loop regions

Kiwon Park[1,2¶], Dohoon Lee[3,4¶], Jiseok Jeong[1,2], Sungwon Lee[1,2], Sun Kim[5], and Kwangseog Ahn[1,2,6*]

[1]Center for RNA Research, Institute for Basic Science, Seoul 08826, Republic of Korea

[2]School of Biological Sciences, Seoul National University, Seoul 08826, Republic of Korea

[3]Bioinformatics Institute, Seoul National University, Seoul 08826, Republic of Korea

[4]BK21 FOUR Intelligence Computing, Seoul National University, Seoul 08826, Republic of Korea

[5]Department of Computer Science and Engineering, Seoul National University, Seoul 08826, Republic of Korea

[6]SNU Institute for Virus Research, Seoul National University, Seoul 08826, Republic of Korea

*Corresponding author

Email: ksahn@snu.ac.kr (KA)

[¶]These authors contributed equally to this work.

Abstract

Although HIV-1 integration sites favor active transcription units in the human genome, high-resolution analysis of individual HIV-1 integration sites has shown that the virus can integrate into a variety of host genomic locations, including non-genic regions. The invisible infection by HIV-1 integrating into non-genic regions, challenging the traditional understanding of HIV-1 integration site selection, is more problematic because they are selected for preservation in the host genome during prolonged antiretroviral therapies. Here, we showed that HIV-1 integrates its viral genome into the vicinity of R-loops, a genomic structure composed of DNA–RNA hybrids. VSV-G-pseudotyped HIV-1 infection initiates the formation of R-loops in both genic and non-genic regions of the host genome and preferentially integrates into R-loop-rich regions. Using a HeLa cell model that can independently control transcriptional activity and R-loop formation, we demonstrated that the exogenous formation of R-loops directs HIV-1 integration-targeting sites. We also found that HIV-1 integrase proteins physically bind to the host genomic R-loops. These findings provide novel insights into the mechanisms underlying retroviral integration and the new strategies for antiretroviral therapy against HIV-1 latent infection.

Introduction

Retroviruses cause permanent host infections by integrating their reverse-transcribed viral genomes into the host genome. Retroviral integration considerably affects a wide range of biological phenomena, including the persistence of fatal human diseases and the shaping of metazoan evolution (*1*). Human immunodeficiency virus (HIV)-1 is a representative retrovirus underlying the global burden of acquired immune deficiency syndrome (AIDS) (*2*). The chromosomal landscape of HIV-1 integration has been shown to influence proviral gene expression, the persistence of integrated proviruses, and prognosis in antiretroviral therapy (*3-5*). Integration into the host genome is not random and displays distinct preferences for gene-dense regions, where active transcription occurs (*6*), by interacting with host factors, such as transcription activators, epigenetic marker binding proteins, and super enhancers (*7-13*). However, transcriptional activity is not the sole determinant of the HIV-1 integration site landscape (*10*). For instance, some highly expressed genes do not necessarily correspond to a high level of integration in PBMCs infected in culture with HIV (*14*). Despite their lower probability of integration, HIV-1 proviruses are observed in non-genic regions of the genomes of infected individuals (*4*). This indicates the possibility of there being an undiscovered mechanism or determinant that constitutes the correct genomic environment for HIV-1 integration.

An R-loop is a three-stranded nucleic acid structure that comprises a DNA–RNA hybrid and a displaced strand of DNA, and has long been considered a transcription byproduct (*15, 16*). R-loops in cellular genomes are enriched in actively transcribed genes as they occur naturally during transcription (*15, 17*). However, R-loop formation is not limited to gene body regions and is widespread in the genome (*15*). As a result of *in trans* R-loop formation,

R-loops are also abundant in non-genic regions, such as intergenic regions and repetitive

sequences, including transposable elements, centromeres, and telomeres (*15, 18-20*),

independent of the transcriptional activity of the genes harboring the R-loops. Although R-

loops have been identified as critical intermediates and regulators of a number of biological

processes (*15, 16, 21*), the dynamics and the role played by cellular R-loops in pathological

contexts remain unclear.

R-loops are important contributors to molding the genomic environment and spatial

organization of the cellular genome, and can potentially play a novel role in host-pathogen

interaction. In the cellular genome, R-loops relieve superhelical stresses and are often

associated with open chromatin marks and active enhancers (*22, 23*), which are also distributed

over HIV-1 integration sites (*6, 9, 10*). In transcription-induced R-loop formation, a guanine-

quadruplex (G4) structure can be generated in the non-template DNA strand of the R-loop (*24*).

A recent study showed that G4 DNA can influence both productive and latent HIV-1 integration

(*25*). In addition, R-loops are prevalent non-canonical B-form DNA structures (*26*) and

intermediates between the B-form DNA and A-form RNA conformations (*27*). It has recently

been reported that the B-to-A transition in the target DNA occur during retroviral integration

(*27, 28*). Accumulating evidence suggests that host genomic R-loops are undiscovered host

factors in the HIV-1 integration site selection mechanism that dynamically interact with the

host genomic environment.

Here, we showed a notable role for R-loops in the interaction between HIV-1 and its

host, specifically in HIV-1 integration. VSV-G-pseudotyped HIV-1-infection induces host

cellular R-loop formation, and the R-loop rich regions of the host genome are preferred for

HIV-1 integration in diverse cell types. HIV-1 integrase proteins showed considerable binding

affinity for nucleic acid substrates containing R-loop structures. Our results suggest that R-

loops are important components of the host genomic environment for HIV-1 integration site

determination.

Results

Host genomic R-loops accumulate by VSV-G-pseudotyped HIV-infection in diverse cell

types

To investigate the relationship between HIV-1 infection and host cellular R-loops, we

first analyzed R-loop dynamics in different types of cells infected with HIV-1 at early post-

infection time points using DNA–RNA immunoprecipitation followed by cDNA conversion

coupled with high-throughput sequencing (DRIPc-seq) using a DNA–RNA hybrid-specific

binding antibody, anti-S9.6 (*29*). HeLa cells, primary CD4$^+$ T cells isolated from two individual

donors, and the CD4$^+$/CD8$^-$ T cell lymphoma Jurkat cell line were infected with the VSV-G-

pseudotyped HIV-1-EGFP in a time-course manner. The infected cells were harvested at the

same post-seeding time point but at 0, 3, 6, and 12 h post infection (hpi) for DRIPc-seq library

construction (Fig. 1A and S1A-C Fig.). Our DRIPc-seq analysis yielded loci-specific R-loop

signals at the referenced R-loop-positive loci (RPL13A and CALM3) and an R-loop-negative

locus (SNRPN) (*29*), which were both strand-specific and highly sensitive to pre-

immunoprecipitation *in vitro* RNase H treatment in HeLa cells, CD4$^+$, and Jurkat T cells (Table

S1-3). Notably, the number of DRIPc-seq peaks mapped to the human reference genome

increased markedly during the early post infection with HIV-1 (3 and 6 hpi for HeLa cells and

6 and 12 hpi for CD4$^+$ and Jurkat T cells; Fig. 1B). Most of the peaks that mapped in cells

harvested at 0 hpi were commonly found in all other samples, but a significant numbers of

unique peaks were observed after infection (Fig. 1C).

In addition to the DRIPc-seq data analysis, we used different biochemical approaches

to examine R-loop accumulation after HIV-1 infection in HeLa cells. R-loop accumulation in

HIV-1-infected cells was observed using DNA–RNA hybrid dot blots with the anti-S9.6

antibodies (Fig. 1D). Dot intensity significantly increased upon HIV-1 infection at 6 hpi and

the enhanced R-loop signals on the dot blots of HIV-1-infected cells were highly sensitive to

in vitro treatment with RNase H (Fig. 1D). This result is highly consistent with our DRIPc-seq

data analysis of HIV-1-infected HeLa cells. Subsequently, we observed HIV-1-induced R-loops

using an immunofluorescence assay by probing HIV-1-infected or non-infected control cells

with S9.6 antibody at 6 hpi (Fig. 1E). After subtracting the nucleolar signal, the nuclear

fluorescence signal associated with R-loops was significantly enhanced in cells infected with

HIV-1 (Fig. 1E). We validated and quantified HIV-1-infection induced R-loop formation on the

host genome in a genome-site-specific manner by using DRIP followed by real-time

polymerase chain reaction (DRIP-qPCR). In this experiment, the S9.6 signal was determined

for three and two HIV-1-induced-R-loop-positive (P1, P2, and P3) and -negative regions (N1

and N2), respectively, which were defined by DRIPc-seq data analysis (S2A-E Fig.). We

detected significantly increased R-loop signals that were highly sensitive to RNase H treatment

of pre-immunoprecipitates in the P1, P2, and P3 regions of HIV-1-infected cells at 6 hpi

compared to those in the cells harvested at 0 hpi (S3A Fig.). However, the HIV-1-induced R-

loop-negative regions N1 and N2, did not show significant R-loop accumulations (S3A Fig.).

Importantly, the R-loop signal was enriched even in cells infected with HIV-1 when

the reverse transcription or integration of HIV-1 was blocked by enzyme inhibitors, such as

nevirapine (NVP) or raltegravir (RAL) (S3B and S3C Fig.). This result indicates that the

enrichment of R-loop signals in cells originates from the host genome, but not from DNA-RNA

hybrid formation during the viral life cycle or transcriptional burst from integrated HIV-1

proviruses. In addition, we confirmed that nearly 100% of DRIPc-seq reads from HIV-1-

infected HeLa, CD4[+], and Jurkat T cells were aligned to the host cellular genome, but not to

that of HIV-1 (S3D Fig.). Together, these data demonstrated that HIV-1 infection induce host

genomic R-loop enrichment at early post-infection. This also suggests that HIV-1 infection

regulates R-loop dynamics in host cells, possibly by inducing either new R-loop formation or

stabilizing existing R-loops in the host cellular genome.

R-loops accumulation after HIV-1 infection are widely distributed in both genic and non-

genic regions

To investigate the distribution of cellular genomic R-loops during HIV-1 infection, we

conducted a genome-wide analysis of the DRIPc-seq data. Unique DRIPc-seq peaks observed

after HIV-1 infection were numerous and relatively long (Fig. 2A). This suggests that the R-

loops induced by HIV-1 infection occupy a genomic region larger than that of the R-loops

without HIV-1 infection. We observed a significant accumulation of R-loops over diverse

genomic compartments at the hpi of HIV-1-infection induced R-loop formation (Fig. 2B). The

presence of R-loops is often correlated with high transcriptional activity, and we found a

significantly high proportion of DRIPc-seq peaks enrichment upon HIV-1 infection in the gene

body regions (Fig. 2B). However, we also observed enrichment of HIV-1-infection induced

DRIPc-seq peaks proportions mapped to intergenic or repeat regions, including short

interspersed nuclear elements (SINEs), long interspersed nuclear elements (LINEs), and long

terminal repeat (LTR) retrotransposons, where transcription is typically repressed (Fig. 2B). In

addition, the proportion of DRIPc-seq peaks mapped to various genomic compartments

remained consistent over the hours following HIV-1 infection (Fig. 2C). This suggests that

HIV-1 infection does not induce R-loop enrichment at specific genomic features, but that R-

loops accumulation after HIV-1 infection is widely distributed. Although the expression of

repetitive elements is mostly repressed during normal cellular activities, HIV-1 infection could

activate endogenous retroviral promoters (*30, 31*). To investigate whether R-loop induction in

gene-silent regions is associated with transcriptome changes during HIV-1 infection, we

performed RNA sequencing (RNA-seq) of HIV-1-infected HeLa cells at 0, 3, 6, and 12 hpi.

Consistent with previous reports, we observed an increase in the expression levels of repetitive

elements at later time points post-infection (S4A Fig. 12 hpi). In contrast, we found that the

expression levels of SINEs, LINEs, and LTRs were even lower at both 3 and 6 hpi compared

with those at 0 hpi whereas HIV-1-induced R-loops were significantly accumulated compared

with those at 0 hpi (S4A Fig.). We further examined the expression profiles of genes containing

the R-loop in HeLa cells. The expression profile of genes harboring HIV-1-induced R-loops in

their gene bodies showed very weak correlations with the signals of DRIPc-seq peaks at 3 hpi

(Pearson's r = 0.21, P = 1.08×10^{-84}; Fig. 2D) and at 6 hpi (Pearson's r = -0.34, P = 2.40×10^{-228};

^{228}; Fig. 2D), which implies that the unique R-loop peaks upon HIV-1 infection do not engage

in a transcriptional burst. In agreement with our DRIPc-seq and global RNA-seq data analysis,

the expression level of the genes harboring HIV-1-infection induced R-loops, which were

quantified by DRIP-qPCR (S3A Fig.), were not significantly affected by HIV-1 infection (S4B

Fig. and Table S4). Together, our data demonstrate that host cellular R-loop accumulation upon

HIV-1 infection is widely distributed in both genic and non-genic regions and is not necessarily

correlate with the expression levels of the genes harboring the R-loops.

HIV-1 integration sites are enriched at systemically induced sequence-specific R-loop regions in cell model

HIV-1 infection is completed by integrating the viral genome into the host's through dynamic interactions with the host genome (*32*). In addition, as HIV-1 infection induces R-loop accumulation at early post infection hours, when the HIV-1 genome are imported into the nucleus and integration may initiate (*33-35*), we hypothesized that host genomic R-loops play a role in HIV-1 integration and possibly in integration site selection. To systemically and directly assess the relationship between host genomic R-loops and HIV-1 integration in a genome site-specific manner, we adapted and modified an elegantly designed episomal system that induces sequence-specific R-loops through DOX-inducible promoters (*17*). To most closely mimic the presence of the R-loop in the host cellular genome, we subcloned the R-loop-forming portion of the mouse gene encoding AIRN (mAIRN) (*18*) or the non-R-loop-forming ECFP sequence (*17*) with a DOX-inducible promoter into the piggyBac transposon vector and co-expressed piggyBac transposase in HeLa cells. The mAIRN gene sequence possesses a high GC skew that causes R-loop structure formation upon transcription by hybridizing newly synthesized RNA back to the template DNA strand, and the non-template DNA strand to remains looped out in a single-stranded form (*18, 36*). In contrast, the ECFP sequence possesses a low GC skew that does not form a stable R-loop at the site of transcription. Thus, it was used as an R-loop-forming negative sequence. These R-loop forming (mAIRN) or non-R-loop forming (ECFP) sequences are nonhuman sequences. Therefore, our cell model allowed us to induce and quantify R-loop formation at designated genomic regions and distinguish R-loop formation from endogenous R-loops on the cellular genome, which are not sequence-specific and impossible to control for induction. Moreover, using this system we can quantify R-loop-associated site-specific HIV-1 integration events at designated regions, which

can also be distinguished from HIV-1 integration events at endogenous host genomic loci. We

designated the pool of cells with the R-loop forming sequence (mAIRN) inserted into its

genome as "pgR-rich (piggyBac R-loop rich)" cell line and the pool of cells with the non-R-

loop forming sequence (ECFP) inserted into its genome as "pgR-poor (piggyBac R-loop poor)"

cell line (Fig. 3A).

A similar number of the copies of the piggyBac transposon were successfully delivered

to the genome of each cell line (S5A Fig.) and DOX treatment strongly induced the

transcriptional activity of mAIRN or ECFP without affecting the transcription of endogenous

loci in either cell line (S5B and S5C Fig.). Although the transcription of mAIRN or ECFP was

strongly induced by DOX treatment, the activity did not exceed that of the endogenous loci in

either cell line (S5D and S5E Fig.). Although the two cell lines showed comparable levels of

DOX-inducible transcriptional activity at the designated sequences (Fig. 3B), only the pgR-

rich cells exhibited robust RNase H-sensitive stable R-loop formation upon DOX treatment

(Fig. 3C, mAIRN). In contrast, R-loops were weakly formed in the pgR-poor cells where a

non-R-loop-forming sequence (ECFP) was inserted into the genome (Fig. 3C, ECFP).

To examine whether the formation of 'extra' R-loops in the host genome influence

HIV-1-infection in host cells, we infected both cell lines with VSV-G-pseudotyped HIV-1-

luciferase viruses and examined the luciferase activity. Notably, we found that pgR-rich cells

showed significantly higher luciferase activity only when R-loops were induced by DOX

treatment, whereas pgR-poor cells showed comparable luciferase activity regardless of

transcriptional activation by DOX treatment (Fig. 3D). We sequenced HIV-1 integration sites

in HIV-1-infected pgR-poor and pgR-rich cells to directly quantify site-specific integration

events in sequence-specific R-loop regions. Remarkably, integration events were significantly

higher in pgR-rich cells only when R-loops were induced by DOX treatment (Fig. 3E).

However, the number of HIV-1 integration-seq mapped reads within the non-R-loop forming

sequence in pgR-poor cells remained very low, even after transcriptional activation by DOX

treatment (Fig. 3E). HIV-1 integration was enriched in the vicinity of R-loop forming regions

in the pgR-rich cell line upon DOX treatment, but the enrichment was not observed in pgR-

poor cells that did not form stable R-loops, even after transcriptional activation by DOX

treatment (Figs. 3F and 3G). This cell-based R-loop-inducing system with independent control

over transcription and R-loop formation enabled the direct measurement of HIV-1 integration

events at defined R-loop regions. The results indicated that host genomic R-loops are preferred

by HIV-1 integration. Moreover, our data suggest that transcriptional activity itself is not

sufficient for HIV-1 integration site determination, but that the formation of R-loops accounts

for HIV-1 integration site selection.

Host genomic R-loop regions are frequently targeted by HIV-1 integration

We attempted to further validate the relationship between R-loops and HIV-1

integration site selection by global analysis of HIV-1 integration sites in the endogenous

genomic regions of HIV-1 infected host cells. We performed HIV-1 integration site sequencing

in HIV-1 infected HeLa cells, CD4$^+$, and Jurkat T cells, and analyzed the sequencing data

combined with our DRIPc-seq data. We counted and compared the number of successfully

integrated proviruses in the R-loop regions (the combined genomic regions within 30-kb

windows centered on DRIPc-seq peaks from 0, 3, 6, and 12 hpi) to those in non-R-loop-forming

regions (the total genomic regions outside of the 30-kb windows centered on DRIPc-seq peaks).

Notably, we detected approximately three to four times more integration in the R-loop regions

than in other genomic regions without R-loops in HeLa cells, CD4[+], and Jurkat T cells (Fig.

4A). Notably, HIV-1 integration sites preferred the central and nearby areas of the R-loops

regions (Fig. 4B). We observed biases in HIV-1 integration in HIV-1-induced R-loop-positive

regions, P1-P3, where showed highly induced R-loop signals upon HIV-1 infection in DRIPc-

seq analysis and DRIP-qPCR (Fig. 4C). In contrast, HIV-1 integration sites were not detected

in the R-loop-negative regions N1 and N2 (Fig. 4D). Overall, our results from bioinformatics

analysis using different types of naïve host cells infected with HIV-1 are consistent with the

idea that the virus has a preference for targeting R-loop-forming regions for integration (Fig.

3), suggesting that R-loops are an important component of the host genomic environment for

HIV-1 integration site determination.

HIV-1 integrase proteins physically interacts with R-loops on the host genome

The HIV-1 intasome is tethered to the host genome for viral cDNA integration.

Intasomes consist of HIV-1 viral cDNA and the HIV-1 integrase proteins. We observed that

HIV-1 was preferentially integrated into the R-loops regions in the host genome; thus, we

hypothesized that the HIV-1 integrase protein could directly bind and be recruited to the

genomic R-loops. To test this hypothesis, we investigated whether HIV-1 integrase proteins

have a physical binding affinity for nucleic acid substrates with an R-loop structure. Although

HIV-1 integrases are DNA and RNA binding proteins (*37, 38*), their ability to bind to a three-

stranded nucleic acid structure that is composed of a DNA-RNA hybrid like R-loop has not

been investigated. We carried out an *in vitro* protein-nucleic acid binding assay by

electrophoretic mobility shift assay (EMSA) with Sso7d-tagged HIV-1 integrase (E152Q)

recombinant proteins and diverse structures of nucleic acid substrates including R-loops and

simple dsDNA duplexes. In this experiment, we used HIV-1 integrase protein with an active

site amino acid substitution, E152Q, to prevent any undesirable alternation in nucleic acid

substrates by enzymatic activities of integrase proteins, such as 3'-processing. Notably, nucleic

acid substrate consisted of an R-loop structure bound to HIV-1 integrase proteins with higher

binding affinity than that of the simple dsDNA duplex (Fig. 5A). Additionally, the R-loop

composing forms of nucleic acid structures, such as RNA-DNA hybrid with exposed ssDNA

(R:D+ssDNA) and RNA-DNA hybrid (hybrid), also showed high binding affinity to integrases

(S6A Fig. and Fig. 5A).

We validated the interaction between cellular genomic R-loops and HIV-1 integrase

proteins by DNA–RNA hybrid immunoprecipitation using S9.6 antibodies against FLAG-

tagged HIV-1 integrase-expressing HeLa cells (Fig. 5B). Under our experimental conditions,

R-loops were reproducibly immunoprecipitated (S6B Fig.) and HIV-1 integrase proteins co-

immunoprecipitated with the R-loops (Fig. 5C). DNA–RNA hybrids also co-

immunoprecipitated with the positive control H3 (*39*) but not with the negative controls

LaminA/C and Actin (*39*) (Fig. 5C). To verify the specificity of our co-immunoprecipitation

results for R-loops and HIV-1 integrases, we performed DNA–RNA hybrid

immunoprecipitation with RNase H treatment (S6C Fig.). The S9.6 signal of the

immunoprecipitated nucleic acids was highly sensitive to RNase H treatment of the pre-

immunoprecipitates (Fig. 5D). Accordingly, the blotting signal of the co-immunoprecipitated

HIV-1 integrase and H3 proteins was significantly reduced upon RNase H treatment (Fig. 5E).

We performed reciprocal immunoprecipitation using an anti-FLAG monoclonal antibody and

detected the immunoprecipitated R-loops using dot blot analysis with anti-S9.6. R-loops, which

were immunoprecipitated using HIV-1 integrase, and the S9.6 signal of immunoprecipitated

nucleic acids was highly sensitive to RNase H treatment (Fig. 5F and S6D Fig.). Subsequently,

we examined the interaction between the R-loops and HIV-1 integrase using a proximity-

ligation assay (PLA)   in HIV-1-infected cells. We used two antibodies: one that binds to the

R-loops (anti-S9.6) and another that binds to GFP-tagged HIV-1 integrase. We detected PLA

signals in cells infected with HIV-IN-EGFP virions (*34*) and non-infected control cells. PLA

signals in non-infected cells were comparable to those in S9.6-alone and GFP-alone single

antibody-negative controls; however, PLA signals significantly increased upon HIV-1 infection

(Fig. 5G and S6E Fig.). Our data suggest that the HIV-1 frequently targets R-loop-rich regions

for viral genome integration by physically binding of HIV-1 integrase proteins to R-loop

structures on the host genome.

Discussion

301   In this study, we found that HIV-1 preferentially integrated into regions rich in R-loops, suggesting that R-loops are novel host factors that contribute to HIV-1 integration site selection. In our bioinformatics analysis, host cellular R-loops were induced by VSV-G-pseudotyped HIV-1 infection and widespread in host genomic regions. Using our R-loop-inducible cell models, R-loop formation, and not necessarily the transcriptional activity itself, was found to be important for HIV-1 integration site determination. In addition, HIV-1 integrase proteins favor physical binding with R-loops *in vitro*, and interact with host genomic R-loops in HIV-1-infected cells. These results demonstrate that HIV-1 exploits and frequently targets the host genomic R-loop regions for successful integration and infection.

311   One possible explanation for why HIV-1 integration shows a preference for host genomic R-loop regions is that the R-loop structure may drive dynamics in the genomic environment and the spatial organization of the genome, resulting in increased accessibility of HIV-1 intasome binding to the target host genomic region. The R-loops display enhancer and insulator chromatin states, that can act as distal regulatory elements by recruiting diverse chromatin binding factors (*22*). This not only allows R-loops to drive dynamics in the genome, but also possibly drives R-loop-mediated integration over long-range genomic regions. R-loop regions exhibit increased chromatin accessibility. In the cellular genome, these structures relieve superhelical stresses and are often associated with open chromatin marks and active enhancers (*22, 23*), which are also distributed over HIV-1 integration sites (*6*). In the case of transcription-induced R-loop formation, a guanine-quadruplex (G4) structure can be generated in the non-template DNA strand of the R-loop, which is another contributor to genome architecture (*24*). A recent study showed that G4 DNA can influence both productive and latent HIV-1 integration, as well as the potential for reactivation of latent proviruses (*25*).

Another possible explanation for the preference of HIV-1 integration for host genomic

R-loops is that R-loops may play a collaborative role with host factors governing the HIV-1

integration site selection. Cellular R-loops are recognized and regulated by numerous cellular

proteins (*39, 40*). LEDGF/p75 (*9, 13, 41*) and CPSF6 (*7, 9*) are two decisive host factors that

direct HIV-1 integration by interacting with integrase or trafficking the viral preintegration

complex towards the nuclear interior (*7, 9*). In fact, these host factors have recently been

identified as potential R-loop-binding proteins in DNA–RNA interactome analysis (*39*) and R-

loop proximity proteomics (*40*), respectively. R-loops are tightly regulated by DNA damage

response proteins (*42*) and the DNA repair machinery plays an important role in HIV-1

integration process (*32*). For example, the Fanconi anemia pathway (*43, 44*), a well-known R-

loop regulatory pathway, has been recently proposed as an HIV-1 integration regulatory factor

exploited by HIV-1 (*45*). Considering theses previous studies and our current findings, we

propose R-loops as another putative host factor driving HIV-1 integration site determination,

and a possible intermediate regulator of HIV-1 integration site selection by such host proteins.

Our data showed that HIV-1 integrase proteins physically interacted with genomic R-

loops *in vitro* and in cells. Recent advancements in cryogenic electron microscopy (cryo-EM)

technology have revealed the conformational characteristics of the target DNA during retroviral

integration (*27, 28*). During retroviral integration, the target DNA undergoes a transition of its

conformation from B-form to A-form. R-loops, which represent intermediates between the B-

form DNA and A-form RNA conformations (*27*), may have an intrinsic preferential binding

ability to retroviral intasomes over other nucleic acid structures.

Viruses often take advantage of various host factors, and targeting the viral

components that manipulate the host cellular environment can be an effective strategy for

antiviral therapy. Our study showed the host genomic R-loops significantly accumulate shortly

after HIV-1 infection. Thus, it is possible that virion-associated HIV-1 proteins are responsible

for inducing these R-loops. For instance, the HIV-1 accessory protein Vpr causes genomic

damage (*46*) and transcriptomic changes during the early stages post infection(*47*), both of

which can lead to *in cis* and *in trans* R-loop formation (*16*). Another HIV-1 accessory protein,

Vif, counteracts the host antiviral factor, APOBEC3 (*48, 49*), which was recently found to

regulates cellular R-loop levels (*50*). Identifying the HIV-1 components responsible for

inducing host cellular R-loops, elucidating the mechanism by which they induce genome-wide

R-loop formation, and contributing to successful viral integration into selective genomic

regions, are areas for further research.

Although most HIV-1 integration occurs in genic regions (*4, 6*), HIV-1 proviruses are

also found in non-genic regions (*51*) and understanding these "transcriptionally silent"

proviruses is critical for developing strategies to completely eliminate HIV-1. In HIV-1 elite

controllers, who suppress viral gene expression to undetectable levels, HIV-1 proviruses in

heterochromatic regions are not eliminated but selected by the immune system (*5*). Moreover,

proviruses with lower expression level can persist in the host genome even during antiretroviral

therapy (*4*). However, the mechanism by which HIV-1 targets gene-silent regions for

"invisible" integration remains unclear. Our study has revealed that R-loops are enriched in

both genic and non-genic regions during HIV-1 infection, and that the virus preferentially

targets these R-loops for integration. We propose that R-loops, particularly those enriched in

non-genic regions, may represent the mechanism by which the virus achieves "invisible" and

permanent infection.

Materials and methods

Cell culture

HeLa and HEK293T cells were cultured in Dulbecco's modified Eagle's medium (Gibco) supplemented with 10% (v/v) fetal bovine serum (FBS, Cytiva), antibiotic mixture (100 units/ml penicillin–streptomycin, Gibco), and 1% (v/v) GlutaMAX-I (Gibco). Jurkat cells were cultured in Roswell Park Memorial Institute (RPMI) 1640 medium (ATCC) supplemented with 10% (v/v) FBS (Cytiva). Cells were incubated at 37°C and 5% CO_2.

Virus production and infection

VSV-G-pseudotyped HIV-1 virus stocks were prepared by performing standard polyethylenimine-mediated transfection of HEK293T monolayers with pNL4-3 ΔEnv EGFP (NIH AIDS Reagent Program 11100) or pNL4-3. Luc.R-E (NIH AIDS Reagent Program, 3418) along with pVSV-G at a ratio of 5:1. HIV-IN-EGFP virions were produced as previously described (34) by performing polyethylenimine-mediated transfection of HEK293T cells with 6 μg of pVpr-IN-EGFP, 6 μg of HIV-1 NL4-3 non-infectious molecular clone (pD64E; NIH AIDS Reagent Program 10180), and 1 μg of pVSV-G. The cells were incubated for 4 h before the medium was replaced with fresh complete medium. Virion-containing supernatants were collected after 48 h, filtered through a 0.45-μm syringe filter, and pelleted using the Lenti-X Concentrator (631232; Clontech) according to the manufacturer's instructions. The multiplicity of infection (MOI) of virus stocks was determined by transducing a known number of HeLa cells with a known amount of virus particles and then counting GFP-positive cells using flow cytometry. For luciferase reporter HIV-1 virus, the HIV-1 p24 antigen content in viral stock were quantified using the HIV1 p24 ELISA kit (Abcam, ab218268), according to the manufacturer's instruction. For virus infection, HeLa cells were seeded at a density of 0.5–4 ×

395   10^5 cells/mL on the day before infection. The culture medium was replaced with fresh complete

396   culture medium 2 hpi. The infected cells were washed twice with PBS and harvested at the

397   indicated time points. Jurkat cells were seeded at a density of 1×10^6 cells/mL and inoculated

398   with 300ng/p24 capsid antigen. The plates were centrifuged at 1000 g at 30°C for 1 h. The

399   medium was replaced with fresh RPMI 2 h after infection.

Primary cell isolation, culture, T cell activation, and infection

For CD4$^+$ T cells isolation, human PBMC (ST70025, STEMCELL Technologies) was mixed

and incubated with MACS CD4 MicroBeads (130-045-101, Miltenyi Biotec) and FITC-

conjugated mouse anti-CD4 (561005, BD Bioscience) according to the manufacturer's

instructions. Then the CD4$^+$ T cells were enriched by using LS Columns (130-042-401,

Miltenyi Biotec) and MidiMACS Separator (130-042-302, Miltenyi Biotec). The efficiency of

magnetic separation was analyzed by using Flow-Activated Cell Sorter Canto II (BD

Bioscience) and Flowjo software (Flowjo).

CD4$^+$ T cells were cultured in Roswell Park Memorial Institute (RPMI) 1640 medium (Gibco),

supplemented with 10% (v/v) fetal bovine serum (FBS, Cytiva), antibiotic mixture (100

units/ml penicillin–streptomycin, Gibco), 1% (v/v) GlutaMAX-I (Gibco), and 20 ng/ml of IL-

2 (PHC0026, Gibco), left in resting state or activated with Dynabeads Human T-Activator

CD3/CD28 (1161D, Thermo Fisher Scientific) for 72 h. CD4$^+$ T cells activation efficiency was

assessed by staining cells with FITC-conjugated mouse anti-CD25 (340694, BD Bioscience)

and APC-conjugated mouse anti-CD69 (130-114-046, Miltenyi Biotec) and using Flow-

Activated Cell Sorter Canto II (BD Bioscience) and Flowjo software (Flowjo).

Purified and activated CD4$^+$ T cells were seeded at a density of 1×10^6 cells/mL and inoculated

with 600ng/p24 capsid antigen in presence of polybrene. The plates were centrifuged at 1000

g at 30°C for 1 h. The medium was replaced with fresh RPMI 2 h after infection.

DRIP-qPCR

DRIP was performed as described for the construction of the DRIPc-seq library. After the

elution of isolated complexes, nucleic acids were purified using the standard phenol-

chloroform extract method and used for qPCR. S6 Table presents details of the primer

sequences used for DRIP-qPCR analysis.

RNA-seq library construction

For RNA-seq, HeLa cells were infected with VSV-G-pseudotyped HIV-1 NL4-3 ΔEnv EGFP

virus at a MOI of 0.6 and harvested at 0, 3, 6, and 12 hpi. Sequencing was performed with

biological replicates. Total RNA was extracted using TRIzol reagent (Invitrogen), according to

the manufacturer's instructions. An mRNA sequencing library was constructed using Illumina

adaptors harboring p5 and p7 sequences and Rd1 SP and Rd2 SP sequences. Sequencing was

performed using the HiSeq2500 system (Illumina).

Luciferase assay

HeLa cells infected with VSV-G-pseudotyped pNL4-3.Luc.R-E HIV-1 viruses were harvested

at 48 hpi, and luminescence was measured using the Dual-Luciferase Reporter Assay System

(Promega) according to the manufacturer's instructions. Briefly, 250 μl of passive lysis buffer

was used to lyse cells for each sample, 20 μl of the lysate was mixed with 100 μl of the

Luciferase Assay Reagent II, and the luminescence of firefly luciferase was measured using a

microplate luminometer (Berthold). The luminescence signal were normalized with total

protein content, measured by BCA assay.

Quantitative real-time PCR (qPCR)

For RT (reverse transcription)-qPCR, 1 μg of RNA was reverse-transcribed using the ReverTra Ace qPCR RT Kit (TOYOBO) following the manufacturer's instructions. For qPCR, DNA extracts were prepared using a DNA purification kit (Qiagen, 51106) according to the manufacturer's instructions. Equivalent amounts of purified gDNA from each sample were analyzed using qPCR. qPCR was performed using TOPreal qPCR PreMIX (Enzynomics, RT500M). The reactions were performed in duplicate or triplicate for technical replicates. PCR was performed using the iCycler iQ real-time PCR detection system (Bio-Rad). All the primers used for qPCR are listed in S6 Table.

DRIPc-seq library construction

DRIP followed by library preparation, next-generation sequencing, and peak calling were performed as described earlier (*29*). Briefly, the corresponding cells were harvested and their gDNA was extracted. The extracted nucleic acids were fragmented using a restriction enzyme cocktail with BsrB I (NEB, R0102S), HindIII (NEB, R0136L), Xba I (NEB, R0145L), and EcoRI (NEB, R3101L) overnight at 37°C. Half of the fragmented nucleic acids were digested with RNase H (New England Biolabs) overnight at 37°C to serve as a negative control. The digested nucleic acids were cleaned using standard phenol-chloroform extraction and resuspended in DNase/RNase-free water. DNA–RNA hybrids were immunoprecipitated from total nucleic acids using mouse anti-DNA–RNA hybrid S9.6 (Kerafast, ENH001) DRIP binding buffer and incubated overnight at 4°C. Dynabeads Protein A (Invitrogen, 10001D) was used to pull down the DNA-antibody complexes by incubation for 4 h at 4°C. The isolated complexes were washed twice with DRIP binding buffer before elution. For elution, the isolated complexes were incubated in an elution buffer containing proteinase K for 45 min at

55 °C. Subsequently, DNA was purified using the standard phenol-chloroform extract method

and subjected to DNase I (Takara, 2270 B) treatment and reverse transcription for DRIPc-seq

library construction. DRIPc-seq was performed in biological replicates. S5 Table shows details

of the oligonucleotides used for DRIPc-seq library construction. DRIPc-seq libraries were

analyzed using 150 bp paired-end sequencing on a HiSeqX Illumina instrument.

Immunofluorescence microscopy

For immunofluorescence assays of S9.6 nuclear signals, when indicated, the cells were pre-

extracted with cold 0.5% NP-40 for 3 min on ice. Cells were fixed with 100% ice-cold methanol

for 10 min on ice and then incubated with 100% ice-cold acetone for 1 min. The slides were

washed three times with 1× PBS and incubated with or without 60 U/mL RNase H (M0297S,

NEB) at 37°C for 36 h or left untreated. The slides were subsequently briefly rinsed thrice with

2% BSA/0.05% Tween (in PBS) and incubated with mouse anti-DNA–RNA hybrid S9.6

(Kerafast, ENH001; 1:100) and rabbit anti-nucleolin (Abcam, ab22758; 1:300) in 2%

BSA/0.05% Tween (in PBS) for 4 h at 4°C. The slides were then washed three times with 2%

BSA/0.05% Tween (in PBS) and incubated with goat anti-rabbit AlexaFluor-488-conjugated

(Invitrogen, A-11008) and goat anti-mouse AlexaFluor-568-conjugated (Molecular Probes,

A11004) secondary antibodies (1:200) for 2 h at room temperature. The slides were then

washed three times with 2% BSA/0.05% Tween (in PBS) and mounted using the ProLong Gold

AntiFade reagent (Invitrogen). Images were obtained using an inverted microscope Nikon

Eclipse Ti2, equipped with a 1.45 numerical aperture, plan apochromat lambda 100× oil

objective, and an scientific complementary metal–oxide–semiconductor camera (Photometrics

prime 95 B 25 mm). For each field of view, images were obtained with DAPI395, GFP488,

and Alexa594 channels using the NIS-Elements software. For quantification analysis, binary

masks of nuclei and nucleoli were generated using the ROI manager and auto local thresholding

using the ImageJ software. The intensity of nuclear signals for DNA–RNA hybrids and

nucleolin was then quantified. The final DNA–RNA hybrid signals in the nucleus were

calculated by subtracting the nucleolin signals from the DNA–RNA hybrid signals.

pgR-rich and -poor cell line generation with piggyBac transposition

We adapted and modified an elegantly designed episomal system that induces defined R-loops

with controlled transcription levels (*17*) for R-loop-forming or non-R-loop-forming sequence

subcloning into the piggyBac transposon vector. HeLa cells were seeded at a density of 5×10^4

cells/ml in a 6-well plate. The next day, cells were transfected with 0.2 µg of Super PiggyBac

Transposase Expression Vector (System Biosciences, PB210PA-1) and 0.2, 1, or 2 µg of

transposon vectors with appropriate "cargo" sub cloned using Lipofectamine 3000 (Invitrogen)

according to the manufacturer's instructions. After 3 days, the cells were treated with 10 µg/ml

blasticidin S (Gibco, A1113903) for selection. Cells with positive integrants for more than 7

days were validated using immunoblotting or RT-qPCR following treatment with DOX. Jurkat

cells were seeded at a density of 8×10^5 cells/ml in a 6-well plate and transfected with 0.2 µg

of transposase and 1 µg of corresponding transposon vectors with Lipofectamine 3000, like

HeLa cells. After 3 days, the cells were treated with 10 µg/ml blasticidin S (Gibco, A1113903)

for selection. For each passage, cells were cushioned onto Ficoll-Pacque (Cytiva, 17144002)

to separate live cells from dead cell debris. The cells over the cushion were washed with PBS

and incubated in cell culture medium with 10 µg/ml of blasticidin for further selection for at

least 14 days. Cells with positive integrants were validated by immunoblotting after treatment

with DOX. Quantification of successfully integrated piggyBac transposons was performed

using a piggyBac qPCR copy number kit (System Biosciences, PBC100A-1).

HIV-1 integration site sequencing library construction

HIV-1 integration site sequencing library construction was performed as described earlier (*7, 9*). Summarily, HeLa cells and primary CD4$^+$ T cells were infected with VSV-G-pseudotyped HIV-1 NL4-3 ΔEnv EGFP virus at a MOI of 0.6 and harvested 5 days post infection. gDNA was isolated using a DNA purification kit (Qiagen, 51106), according to the manufacturer's instructions. gDNA (10 µg) was digested overnight at 37°C with 100 U each of the restriction endonucleases MseI (NEB, R0525L) and BglII (NEB, R0144L). Linker oligonucleotides, which were compatible for ligation with the MseI-generated DNA ends, were ligated with gDNA overnight at 12°C in reactions containing 1.5 µM ligated linker, 1 µg fragmented DNA, and 800 U T4 DNA ligase (NEB, M0202S). Viral LTR–host DNA junctions were amplified using semi-nested PCR with a unique linker-specific primer and LTR primers. The second round of PCR was carried out with primers binding to the LTR and the linkers for next-generation sequencing. Two PCRs were performed in parallel for the first round of PCR and five PCRs were performed in parallel for the second round of PCR to enhance library diversity. S7 Table presents details of the oligonucleotides used for HIV-1 integration site sequencing library construction. HIV-1 integration site sequencing was performed in biological replicates. Integration site libraries were analyzed using 150 bp paired-end sequencing on a HiSeqX Illumina instrument.

Co-immunoprecipitation of DNA–RNA hybrid

DNA–RNA hybrid immunoprecipitation was performed as described earlier (*39*). Summarily, non-crosslinked HeLa cells transfected with the pFlag-IN codon-optimized plasmid were lysed in 85 mM KCl, 5 mM PIPES (pH 8.0), and 0.5% NP-40 for 10 min on ice, and then, the lysates were centrifuged at 750 *g* for 5 min to pellet the nuclei. The pelleted nuclei were resuspended

in sodium deoxycholate, SDS, and sodium lauroyl sarcosinate in RSB buffer and were

sonicated for 10 min (Diagenode Bioruptor). Extracts were then diluted (1:4 in RSB + T buffer)

and subjected to immunoprecipitation with the S9.6 antibody overnight at 4°C. Antibody-

bound complexes were incubated with Protein A Dynabeads (Invitrogen) for 4 h at 4°C for

immunoprecipitation. Normal mouse IgG antibodies (Santa Cruz, sc-2025) were used as

negative controls. RNase A (Thermo Scientific, EN0531) was added during

immunoprecipitation at 0.1 ng RNase A per µg gDNA. Beads were washed four times with

RSB + T; twice with RSB, and eluted either in 2× LDS (Novex, NP0007), 100 mM DTT for

10 min at 70°C (for western blot), or 1% SDS and 0.1 M NaHCO$_3$ for 30 min at room

temperature (for DNA–RNA hybrid dot blot).

For co-immunoprecipitation of DNA–RNA hybrids with RNase H treatment, gDNA containing

RNA–DNA hybrids was isolated from HeLa cells transfected with a pFlag-IN codon-optimized

plasmid using a QIAmp DNA Mini Kit (Qiagen, 51304). gDNA was sonicated for 10 min

(Diagenode Bioruptor) and then treated with 5.5 U RNase H (NEB, M0297) per µg of DNA

overnight at 37 °C. A fraction of gDNA was stored as "nucleic acid input" for dot blot analysis.

gDNA was cleaned using standard phenol-chloroform extraction, resuspended in

DNase/RNase-free water, enriched for DNA–RNA hybrids using immunoprecipitation with the

S9.6 antibody (overnight at 4°C), isolated with Protein A Dynabeads (Invitrogen; 4 h at 4°C),

washed thrice with RSB+T. The immunoprecipitated complexes were incubated with nuclear

extracts of HeLa cells transfected with the pFlag-IN codon-optimized plasmid for 2 h at 4°C

with diluted HeLa nuclear extracts. The cell lysate containing proteins were pre-treated with

0.1 mg/ml RNase A (Thermo Scientific, EN0531) for 1 h at 37°C to degrade all RNA–DNA

hybrids, and the excess of RNase A was blocked by adding 200 U of SUPERase in RNase

inhibitor (Invitrogen, AM2694) for immunoprecipitation. In addition, 100 µL fraction of

diluted and RNase A pre-treated extracts prior to immunoprecipitation was stored as "protein

input" for western blotting. Beads were washed four times with RSB + T; twice with RSB, and

eluted either in 2× LDS (Novex, NP0007), 100 mM DTT for 10 min at 70°C (for western blot),

or 1% SDS, and 0.1 M NaHCO$_3$ for 30 min at room temperature (for DNA–RNA hybrid dot

blot).

561 Recombinant Sso7d-IN protein purification

Sso7d-integrase active site mutant E152Q was expressed in Escherichia coli BL21-AI and

purified essentially as previously described (*52*). Briefly, Sso7d-IN (E152Q) expressed BL21-

AI cells were lysis in lysis buffer (20 mM HEPES pH 7.5, 2 mM 2-mercaptoethanol, 1 M NaCl,

10% (w/v) glycerol, 20 mM imidazole, 1 mg RNase A, and 1000 U DNase I) and purified by

nickel affinity chromatography (Qiagen, 30210). Protein were first loaded on HeparinHP

column (GE Healthcare) equilibrated with equilibrated with 20 mM Tris, pH 8.0, 0.5 mM TCEP,

200 mM NaCl, 10% glycerol for anion exchange chromatography prior to the size exclusion

chromatography. Proteins were eluted with a linear gradient of NaCl from 200 mM to 1 M.

Eluted fractions were pooled and then separated on a Superdex-200 PC 10/300   GL column

(GE Healthcare) equilibrated with 20 mM Tris pH 8.0, 0.5 mM TCEP, 500 mM NaCl and 6%

(w/v) glycerol. The purified protein was concentrated to 0.6 mg/ml using an Amicon centrifugal

contentrator (EMD Millipore), flash-frozen in liquid nitrogen and stored at -80°C.

574 Electrophoretic mobility shift assay for R-loop binding of Sso7d-IN

To test the binding affinity of Sso7d-tagged HIV-1 IN to different types of nucleic acid

substrates, we prepared R-loop, dsDNA, RNA-DNA hybrid with exposed ssDNA

(R:D+ssDNA), RNA-DNA hybrid (Hybrid), ssDNA, and ssRNA by annealing different

combinations of Cy3, Cy5 or non-labeled oligonucleotides following the previous protocol (*53*).

10 nM of DNA substrate was incubated with Sso7d-IN at different concentrations in assembly

buffer (20 mM HEPES pH 7.5, 5 mM $CaCl_2$, 8 mM 2-mercaptoethanol, 4 uM $ZnCl_2$, 100 mM

NaCl, 25% (w/v) glycerol and 50 mM 3-(Benzyldimethylammonio) propanesulfonate (NDSB-

256)), for 1 h at 30°C then incubated for 15 min on ice. All the reactants were run on 4.5% non-

denaturing PAGE in 1× TBE and then Cy3 or Cy5 fluorescence signal was imaged by

ChemiDoc MP imaging system (Bio-Rad). S8 Table presents details of the oligonucleotide

sequence used for EMSA.

PLA

For PLA, HeLa cells were grown on coverslips and infected with HIV-IN-EGFP virions. At 6

hpi, cells were pre-extracted with cold 0.5% NP-40 for 3 min on ice. The cells were fixed with

4% paraformaldehyde in PBS for 15 min at 4 °C. The cells were then blocked with 1× blocking

solution (Merck, DUO92102) for 1 h at 37°C in a humidity chamber. After blocking, cells were

incubated with the following primary antibodies overnight at 4°C for S9.6-HIV-1-IN_PLA:

mouse anti-DNA–RNA hybrid S9.6 (1:250; Kerafast, ENH001) and rabbit anti-GFP (1:500;

Abcam, ab6556). The following day, after washing with once with buffer A twice (Merck,

DUO92102), cells were incubated with pre-mixed Duolink PLA plus (anti-mouse) and PLA

minus probes (anti-rabbit) antibodies for 1 h at 37°C. The subsequent steps in the proximal

ligation assay were performed using the Duolink PLA Fluorescence kit (Sigma) according to

the manufacturer's instructions. To obtain images, the mounted specimens were visually

scanned and representative images were acquired using a Zeiss LSM 710 laser scanning

confocal microscope (Carl Zeiss). The number of intranuclear PLA puncta was quantified using

the ImageJ software. For each biological replicate and experiment, a PLA with a single

antibody was performed as a negative control under the same conditions.

DRIPc-Seq data processing and peak calling

DRIPc-seq reads were quality-controlled using FastQC v0.11.9 (*54*), and sequencing adapters were trimmed using Trim Galore! v0.6.6 (*55*) based on Cutadapt v2.8 (*56*). Trimmed reads were aligned to the hg38 reference genome using bwa v0.7.17-r1188 (*57*). Read deduplication and peak calling were performed using MACS v2.2.7.1 (*58*). Because R-loops appear as both narrow and broad peaks in DRIPc-seq read alignment owing to its variable length, two independent "MACS2 callpeak" runs were performed for narrow and broad peak calling. The narrow and broad peaks were merged using Bedtools v2.26.0 (*59*). To increase the sensitivity of DRIPc-seq peak identification, peaks were called after pooling the two biological replicates of the DRIPc-seq sequencing data for each condition.

Consensus R-loop peak calling

The R-loop peaks at 0, 3, 6 and 12 hpi were first merged using "bedtools merge" to create a universal set of R-loop peaks across time points (n = 46542). Then, each of the universal R-loop peaks was tested for overlap with the R-loop peaks for 0, 3, 6 and 12 hpi using "bedtools intersect". In all, 9,190, 21,403, 33,544, and 9,941 peaks overlapped with 0, 3, 6, and 12 hpi R-loop peaks, respectively. For CD4 cells, we identified a universal R-loop set consisting of 3,928 R-loops, and among them, 737, 722, 1,796 and 2,766 peaks overlapped with 0, 3, 6 and 12hpi R-loop peaks.

HIV-1 integration site sequencing data processing

Quality control of HIV-1 integration site-sequencing reads was performed using FastQC v0.11.9. To discard primers and linkers specific for integration site-sequencing from reads, we used Cutadapt v2.8 with the following option: "-u 49 -U 38 --minimum-length 36 --pair-filter

any --action trim -q0,0 –a linker -A TGCTAGAGATTTTCCACACTGACTGGGTCTGAGGG

-A GGGTCTGAGGG --no-indels --overlap 12". This allowed the first position of the read

alignment to directly represent the genomic position of HIV-1 integration. Processed reads

were aligned to the hg38 reference genome using bwa v0.7.17-r1188, and integration sites were

identified using an in-house Python script. Genomic positions supported by more than five read

alignments were regarded as HIV-1 integration sites. For Jurkat cells, we adopted integration

site sequencing data of HIV-1 infected wild type Jurkat cells from SRR12322252 (*60*).

Co-localization analysis of R-loops and integration sites

Enrichment of integration sites near the R-loop peaks was tested using a randomized

permutation test. Randomized R-loop peaks were generated using "bedtools shuffle" command,

thus preserving the number and the length distribution of the R-loop peaks during the

randomization process. Similarly, integration sites were randomized using the "bedtools shuffle"

command. Randomization was performed 100 times. ENCODE blacklist regions (*61*) were

excluded while shuffling the R-loops and integration sites to exclude inaccessible genomic

regions from the analysis. For each of the observed (or randomized) integration sites, the

closest observed (or randomized) R-loop peak and the corresponding genomic distance were

identified using the "bedtools closest" command. The distribution of the genomic distances

was displayed to show the local enrichment of integration sites in terms of the increased

proportion of integration sites within the 30-kb window centered on R-loops compared to their

randomized counterparts.

DNA plasmid construction and transfection

R-loop-forming mAIRN and non-R-loop forming ECPF sequences were subcloned from

pSH26 and pSH36 plasmids, which were generously provided by Prof. Karlene A. Cimprich,

into the piggyBac transposon vector, where the tet operator sequences were located upstream

of the minimal CMV promoter. The pFlag-IN codon-optimized plasmid and pVpr-IN-EGFP

were kindly provided by Prof. A. Engelman and Prof. Anna Cereseto, respectively.

Lipofectamine 3000 (Invitrogen) transfection reagent was used for the transfection of all

plasmids into cells, according to the manufacturer's protocol.

DNA–RNA hybrid dot blotting

653 Total gDNA was extracted using the QIAmp DNA Mini Kit (Qiagen, 51304) according to the

654 manufacturer's instructions. gDNA (1.2 µg) was treated with 2 U RNase H (NEB, M2097) per

655 µg of gDNA for 4 h at 37°C, with half of the sample left untreated but denatured. Half of the

656 DNA sample was probed with S9.6 antibody (1:1000), and the other half was probed with an

657 anti-ssDNA antibody (MAB3034, Millipore, 1:10000).

Immunoblotting

Cells were lysed using RIPA buffer (50 mM Tris, 150 mM sodium chloride, 0.5% sodium

deoxycholate, 0.1% SDS, and 1.0% NP-40) supplemented with 10 µM leupeptin (Sigma-

Aldrich) and 1 mM phenylmethanesulfonyl fluoride (Sigma-Aldrich) and boiled at 98°C for

10 min with SDS sample buffer prior to SDS-PAGE. The primary antibodies used were mouse

monoclonal anti-FLAG M2 (Sigma, F3165), monoclonal mouse anti-HSC70 (Abcam, ab2788),

polyclonal rabbit anti-histone H3 (tri methyl K4) antibody (Abcam, ab8580), monoclonal

mouse anti- HIV-1 Integrase (Santa Cruz, sc-69721), rabbit anti-LaminA/C antibody (Cell

Signaling, 2032), and monoclonal mouse anti-Actin (Invitrogen, MA1-744). All primary

antibodies were used at a dilution of 1:1000 for western blotting. Peroxidase-conjugated anti-

mouse IgG (115-035-062) and anti-rabbit IgG (111-035-003; both Jackson Laboratories) were

used as secondary antibodies at 1:5000 dilution. Signals were detected using the SuperSignal

West Pico chemiluminescence kit (Thermo Fisher Scientific).

RNA-seq data processing

RNA-seq reads were quality-controlled and adapter-trimmed as in DRIPc-seq processing. To

quantify the expression levels of protein-coding genes, processed reads were aligned to the

hg38 reference genome with GENCODE v37 gene annotation (*62*) using STAR v2.7.3a (*63*).

Gene expression quantification was performed using RSEM v1.3.1. To quantify the expression

levels of transposable elements (TEs), we used TEtranscripts v2.2.1 (*64*). Processed reads were

first aligned to the hg38 reference genome using GENCODE v37 and RepeatMasker TE

annotation using STAR v2.7.3a. In this case, STAR options were modified as follows to utilize

multimapping reads in downstream analyses: "--outFilterMultimapNmax 100 --

winAnchorMultimapNmax 100 --outMultimapperOrder random --runRNGseed 77 --

outSAMmultNmax 1 --outFilterType BySJout --alignSJoverhangMin 8 --

alignSJDBoverhangMin 1 --alignIntronMin 20 --alignIntronMax 1000000 --

alignMatesGapMax 1000000". Expression levels of TEs were quantified as read counts with

the "TEcount" command.

Genome annotations

All bioinformatic analyses were performed using the hg38 reference genome and GENCODE

v37 gene annotation. Promoters were defined as a 2-kb region centered at the transcription start

sites of the APPRIS principal isoform of protein-coding genes. TTS regions were defined as

the 2-kb region centered at the 3′ terminals of protein-coding transcripts. CpG island

annotations were downloaded from the UCSC table browser. CpG shores were defined as 2-kb

regions flanking CpG islands, excluding the regions overlapping with CpG islands. Similarly,

CpG shelves were defined as 2-kb regions flanking the stretch of CpG islands and shores while

excluding the regions overlapping with CpG islands and shores. Annotations for LINE, SINE,

and LTR were extracted from the RepeatMasker track in the UCSC table browser.

Identification of viral sequencing reads in DRIPc-seq

To identify sequencing reads originating from the viral genome, we aligned DRIPc-seq reads

to a composite reference genome consisting of the human and HIV1 genome (Genbank

accession number: AF324493.2) and computed the proportion of the reads mapped to HIV1

genome.

Code availability

Bioinformatics pipelines and scripts used in this study are accessible from

https://github.com/dohlee/hiv1-rloop.

Acknowledgements

We are grateful to Prof. Karlene A. Cimprich (Standford University) for providing the pSH26 and pSH36 plasmids, Prof. A. Engelman (Harvard Medical School) for providing pFlag-IN codon optimized plasmid and Prof. Anna Cereseto (University of Trento) for providing pVpr-IN-EGFP. The NL4-3 ΔEnv EGFP and pNL4-3.Luc.R-E- viral plasmids were obtained through the NIH HIV Reagent Program, Division of AIDS, NIAID, NIH. We thank Dr. Sungchul Kim (IBS center for RNA Research) and Seongjin An (Korea University) for their technical support in recombinant protein purification.

Author contributions

K.P. and K.A. designed experiments. K.P., J.J and S.L. performed experiments. D.L. performed the bioinformatical and statistical analyses. K.P., D.L., K.A. and S.K. analyzed the data. K.P., D.L., and K.A. wrote the manuscript.

Funding

This work was supported by the Institute for Basic Science of the Ministry of Science Grant (IBS-R008-D1) and the National Research Foundation of Korea (NRF) grant funded by the Korea government (NRF-2020R1A2C3011298) (to K. A.) and (NRF-2020R1A5A1018081) (to K.A.). The funders had no role in the study design, data collection, analysis, decision to publish, or preparation of the manuscript.

Competing interests

The authors have declared that no competing interests exist.

Figures

Fig 1. HIV-1 infection induces genomic R-loop accumulation in cells at early post-infection. (**A**) Summary

of experimental design for DRIPc-seq in HeLa cells, primary CD4$^+$ T cells and Jurkat cells infected with HIV-1.

(**B**) Bar graphs indicating DRIPc-seq peak counts for HIV-1-infected HeLa cells, primary CD4$^+$ T cells and Jurkat

cells harvested at the indicated hours post infection (hpi). Pre-immunoprecipitated samples were untreated (−) or

treated (+) with RNase H, as indicated. Each bar corresponds to pooled datasets from two biologically independent

experiments. (**C**) All genomic loci overlapping a DRIPc-seq peak from HIV-1 infected HeLa cells, primary CD4$^+$

T cells and Jurkat cells in at least one sample are stacked vertically; the position of each peak in a stack is constant

horizontally across samples. Each hpi occupies a vertical bar, as indicated. Each bar corresponds to pooled datasets

from two biologically independent experiments. Common peaks for all samples are represented in black, and in

dark gray for those common for at least two samples. The lack of a DRIP signal over a given peak in any sample

is shown in light gray. The sample-unique peaks are colored blue, yellow, green, and red at 0, 3, 6, and 12 hpi,

respectively. (**D**) Dot blot analysis of the R-loop in gDNA extracts from HIV-1 infected HeLa cells with MOI of

0.6 harvested at the indicated hpi. gDNAs were probed with anti-S9.6. gDNA extracts were incubated with or

without RNase H in vitro before membrane loading (anti-RNA/DNA signal). Fold-induction was normalized to

the value of harvested cells at 0 hpi by quantifying the dot intensity of the blots and calculating the ratios of the

S9.6 signal to the total amount of gDNA (anti-ssDNA signal). (**E**) Representative images of the

immunofluorescence assay of S9.6 nuclear signals in HIV-1 infected HeLa cells with MOI of 0.6 harvested at 6

hpi. The cells were pre-extracted of cytoplasm and co-stained with anti-S9.6 (red), anti-nucleolin antibodies

(green), and DAPI (blue). The cells were incubated with or without RNase H in vitro before staining with anti-

S9.6 antibodies, as indicated. Quantification of S9.6 signal intensity per nucleus after nucleolar signal subtraction

for the immunofluorescence assay. The mean value for each data point is indicated by the red line. Statistical

significance was assessed using one-way ANOVA (n >53).

Fig. 2. HIV-1-induced R-loops are enriched at both transcriptionally active and silent regions. (**A**)

Distribution of DRIPc-seq peak lengths for HIV-1-infected HeLa cells, primary CD4[+] T cells and Jurkat cells

harvested at the indicated time points (blue, 0 hpi; yellow, 3 hpi; green, 6 hpi; red, 12 hpi). (**B**) Stacked bar graphs

indicating the proportion of DRIPc-seq peaks mapped for HIV-1-infected HeLa cells, primary CD4[+] T cells and

Jurkat cells harvested at the indicated hpi over different genomic features. (**C**) Stacked bar graphs indicating the

proportion of DRIPc-seq peaks mapped to indicated genomic compartments for HIV-1-infected HeLa cells,

primary CD4[+] T cells and Jurkat cells harvested at the 0, 3, 6, and 12 hpi . (**D**) Correlation between gene expression

and DRIPc-seq signals of HIV-1-infected HeLa cells with MOI of 0.6 harvested at the indicated hpi. Statistical

significance was assessed using Pearson's r and p-values.

Fig. 3. R-loop inducible cell line model directly addresses R-loop-mediated HIV-1 integration site selection.
(**A**) Summary of the experimental design for R-loop inducible cell lines, pgR-poor and pgR-rich. (**B**) Gene expression of ECFP (gray) and mAIRN (red), as measured using RT-qPCR in pgR-poor or pgR-rich cells. Where indicated, the cells were incubated with 1 μg/ml DOX for 24 h. Gene expression was normalized relative to *β-actin*. Data are presented as the mean ± SEM, n = 3. (**C**) DRIP-qPCR using the anti-S9.6 antibody against ECFP and mAIRN in pgR-poor or pgR-rich cells. Where indicated, the cells were incubated with 1 μg/ml DOX for 24 h. Pre-immunoprecipitated samples were untreated or treated with RNase H as indicated. Values are relative to those of DOX-treated (+) RNase H-untreated (−) pgR-poor cells. Data are presented as the mean ± SEM; statistical significance was assessed using two-way ANOVA (n = 2). (**D**) Bar graphs indicate luciferase activity at 48 hpi in pgR-poor or prR-rich cells infected with 100ng/p24 capsid antigen of luciferase reporter HIV-1 virus per 1×10^5 cells/mL. Data are presented as the mean ± SEM; P values were calculated using one-way ANOVA (n = 6). (**E**) Box graph indicating the quantified HIV-1 integration site sequencing read count across pgR-poor and pgR-rich transposon sequences in untreated (−) or DOX-treated (+) pgR-poor or pgR-rich cell line infected with 100ng/p24

capsid antigen of luciferase reporter HIV-1 virus per 1×10^5 cells/mL. Each bar corresponds to pooled datasets

from three biologically independent experiments (n =3). In each boxplot, the centerline denotes the median, the

upper and lower box limits denote the upper and lower quartiles, and the whiskers denote the $1.5 \times$ interquartile

range. Statistical significance was assessed using a two-sided Mann–Whitney U test. (**F** and **G**) Heat maps

representing the number of HIV-1 integration-seq mapped read across pgR-poor (**F**) or pgR-rich (**G**) transposon

sequence in untreated (-) or DOX-treated (+) pgR-poor (**F**) or pgR-rich (**G**) cell line. Each rectangular box

corresponds to the pooled the number of HIV-1 integration-seq mapped read from three biologically independent

experiments (n =3) at the indicated position within pgR-poor (**F**) or pgR-rich (**G**) transposon vector. Each light

blue box represents actual position of R-loop forming or non-R-loop forming sequence (ECFP or mAIRN) and

the yellows stars indicate TRE promoter position within vector.

Fig. 4. HIV-1 prefers host genomic R-loop regions for its viral cDNA integration. (A) Bar graphs showing quantified number of HIV-1 integration sites per Mb pairs in total regions of 30-kb windows centered on DRIPc-seq peaks from HIV-1 infected HeLa cells, primary CD4+ T cells and Jurkat cells (magenta) or non-R-loop region in the cellular genome (gray). (B) Proportion of integration sites within the 30-kb windows centered on DRIPc-

seq peaks (magenta solid lines) or randomized DRIPc-seq peaks (gray dotted lines). Control comparisons between

randomized integration sites with DRIPc-seq peaks and randomized DRIPc-seq peaks are indicated by black

dotted lines and gray solid lines, respectively. (**C** and **D**) Superimpositions of HIV-1-induced R-loop positive

chromatin regions, P1-P3 (**C**), and HIV-1-induced R-loop negative chromatin regions, N1 and N2 (**D**), on DRIPc-

seq (blue, 0 hpi; yellow, 3 hpi; green, 6 hpi; red, 12 hpi) and number of mapped read of HIV-1 integration-seq

(integration, black). Magenta dotted lines represent primer binding sites in qPCR following DRIP.

Fig. 5. HIV-1 integrase proteins directly bind to host genomic R-loops. (**A**) Representative gel images for EMSA of Sso7d-tagged HIV-1-integrase (E152Q) with R-loop and dsDNA, 10 nM nucleic acid substrate was incubated with Sso7d-tagged HIV-1-integrase (E152Q) at 0 nM, 20 nM, 50 nM, 100 nM, 200 nM, and 400 nM (left). Unbound fraction were quantified for EMSA of Sso7d-tagged HIV-1-integrase (E152Q) with different types substrates (R-loop, dsDNA, R-loop, R:D+ssDNA and Hybrid). Data are presented as the mean ± SEM, n = 3 (right). (**B**) Summary of the experimental design for R-loop immunoprecipitation using S9.6 antibody in FLAG-

tagged HIV-1 integrase protein-expressing HeLa cells. (**C**) Western blotting for HIV-1 integrase protein, H3, and

LaminA/C of DNA–RNA hybrid immunoprecipitation using the S9.6 antibody. (**D**) **and** (**E**) HeLa gDNA input

was either untreated (–) or treated (+) with RNase H before enrichment for DNA–RNA hybrids using the S9.6

antibody. gDNA–RNA hybrids were incubated with nuclear extracts depleted of DNA–RNA hybrids with RNase

A followed by S9.6 immunoprecipitation. DNA–RNA hybrid dot blot (**D**) and western blot of DNA–RNA hybrid

immunoprecipitation, probed with the indicated antibodies (**E**). (**F**) DNA–RNA hybrid dot blot of FLAG antibody-

immunoprecipitated nucleic acid extracts. Where indicated, nucleic acid extracts were untreated (–) or treated (+)

with RNase H before probing with the S9.6 antibodies. (**G**) Representative images of the proximity-ligation assay

(PLA) between GFP and S9.6 antibodies in HIV-IN-EGFP virion-infected HeLa cells at 6 hpi. Cells were

subjected to PLA (orange) and co-stained with DAPI (blue). PLA puncta in the nucleus are indicated by the yellow

arrows. Quantification analysis of number of PLA foci per nucleus (left). GFP_alone and S9.6_alone were used

as single-antibody controls from HIV-IN-EGFP virion-infected HeLa cells (right). The mean value for each data

point is indicated by the red line. P value was calculated using a two-tailed unpaired *t*-test (n > 50).

Supplemental Information

Materials and Methods

S1 Fig. Primary CD4$^+$ T cells sorting strategies and GFP-HIV-1 infection

S2 Fig. Genome browser screenshot over the HIV-1-induced R-loop forming positive or negative genomic regions

S3 Fig. Host cellular R-loop induction by HIV-1 infection is host-genome specific.

S4 Fig. R-loop induction by HIV-1 infection does not follow transcriptome changes in HeLa cells

S5 Fig. PiggyBac transposon-transposase insertion of R-loop forming and non-R-loop forming sequences in HeLa cells

S6 Fig. HIV-1 integrase proteins directly binds to host genomic R-loops

S1 Table. Chromosomal position and DRIPc-seq signal for referenced R-loop-positive and – negative regions in HIV-1 infected HeLa cells

S2 Table. Chromosomal position and DRIPc-seq signal for referenced R-loop-positive and – negative regions in HIV-1 infected primary CD4$^+$ T cells

S3 Table. Chromosomal position and DRIPc-seq signal for referenced R-loop-positive and – negative regions in HIV-1 infected Jurkat cells

S4 Table. RNA-seq analysis of relative gene expression levels of P1-3 and N1,2 R-loop regions

S5 Table. Oligonucleotides used for DRIPc-seq library construction

S6 Table. Primers used for qPCR

S7 Table. Oligonucleotides used for HIV-1 integration site sequencing library construct

S8 Table. Oligonucleotides used for electrophoretic mobility shift assay substrate preparation

S9 Table. Accession numbers and data sources.

Supplementary figures

S1 Fig. Primary CD4$^+$ T cells sorting strategies and GFP-HIV-1 infection. (**A**) Gating strategy used to

determine the efficiency of CD4$^+$ T cells sorting from human PBMC. Pre-sorted PBMCs were staining with FITC-

conjugated anti-CD4 and subjected for positive CD4+ T cell sorting. The percentages of FITC stained cell

population at each step of cell sorting are as indicated. (**B**) Gating strategy used to determine non-activated (Naïve)

and activated cells (αCD3/28) with two markers, CD25 (FITC) and CD69 (APC), for each donor (upper panels,

Donor 1; lower panels, Donor 2). (**C**) Gating strategy used to determine HIV-1-infectivity of CD4+ T cells from

each donor infected with GFP reporter HIV-1 virus at 48 hpi. The percentages of GFP positive cell population at

are as indicated.

S2 Fig. Genome browser screenshot over the HIV-1-induced R-loop forming positive or negative genomic

regions. (A-C), Genome browser screenshot over the P1 (**A**), P2 (**B**), and P3 (**C**) HIV-1 induced R-loop-positive

chromosomal regions showing result from DRIPc-seq in HIV-1-infected HeLa cells (blue, 0 hpi; yellow, 3 hpi;

green, 6 hpi; red, 12 hpi; black, input signals for each indicated sample) on plus (+) or minus (-) DNA strand.

Magenta dotted lines represent primer binding sites in qPCR following DRIP. **(D** and **E),** Genome browser

screenshot over the N1 (**D**), and N2 (**E**) HIV-1 induced R-loop-negative chromosomal regions showing result

from DRIPc-seq in HIV-1-infected HeLa cells (blue, 0 hpi; yellow, 3 hpi; green, 6 hpi; red, 12 hpi; black, input

signals for each indicated sample) on plus (+) or minus (-) DNA strand. Magenta dotted lines represent primer

binding sites in qPCR following DRIP.

S3 Fig. Host cellular R-loop induction by HIV-1 infection is host-genome specific. (A) DRIP-qPCR using the

anti-S9.6 antibody at P1, P2, P3, N1, and N2 in HIV-1-infected cells with MOI of 0.6 harvested at the indicated

hpi (blue, 0 hpi; green, 6 hpi). Pre-immunoprecipitated materials were untreated (−) or treated (+) with RNase H,

as indicated. Data are presented as the mean ± SEM; P-values were calculated using one-way ANOVA (n = 2).

(**B**) Dot blot analysis of the R-loop in gDNA extracts from HIV-1 infected HeLa cells with MOI of 0.6 harvested

at 6hpi. The cells were treated with DMSO, 10uM of Nevirapine (NVP), or 10uM of Raltegravir (RAL) for 24 h

before infection, as indicated. gDNAs were probed with anti-S9.6. gDNA extracts were incubated with or without

RNase H in vitro before membrane loading (anti-RNA/DNA signal). Fold-induction was normalized to the value

of harvested cells at 0 hpi by quantifying the dot intensity of the blots and calculating the ratios of the S9.6 signal

to the total amount of gDNA (anti-ssDNA signal). (**C**) Representative images of the immunofluorescence assay

of S9.6 nuclear signals in HIV-1 infected HeLa cells with MOI of 0.6 at 6 hpi. The cells were pre-extracted of

cytoplasm and co-stained with anti-S9.6 (red), anti-nucleolin antibodies (green), and DAPI (blue). The cells were

treated with DMSO, 10uM of Nevirapine (NVP), or 10uM of Raltegravir (RAL) for 24 h before infection, as

indicated. Quantification of S9.6 signal intensity per nucleus after nucleolar signal subtraction for the

immunofluorescence assay. The mean value for each data point is indicated by the red line. Statistical significance

was assessed using one-way ANOVA (n >51). (**D**) Pie graphs indicating the percentage of DRIPc-seq reads aligned

to host cellular genome (aquamarine) or to HIV-1 viral genome (gray), out of the total consensus DRIPc-seq peaks

from HIV-infected HeLa cells, primary CD4$^+$ T cells and Jurkat cells.

S4 Fig. R-loop induction by HIV-1 infection does not follow transcriptome changes in HeLa cells. (**A**) Line graphs and heat maps representing expression levels of indicated repetitive elements (SINE, right; LINE, middle; LTR, left) at the indicated hpi of HIV-1 in HeLa cells. Data are presented as the mean expression levels of two biologically independent experiments. (**B**) Indicated gene expression as measured by RT-qPCR in 0 or 6 hpi harvested HIV-1-infected HeLa cells. Data represent mean ± SEM, n = 3, P values were calculated according to two-tailed Student's t-test. P > 0.05; n.s, not significant.

S5 Fig. Regulation of cellular R-loops by RNase H1 expression, or by transposon-transposase insertion of

R-loop forming and non-R-loop forming sequences in HeLa cells. (**A**) Copy number of piggyBac transposon

inserts in each cell line constructed by transfecting the transposon vector and transposase-expressing vector. (**B**

and **C**) Fold induction of gene expression for the indicated genes as measured by RT-qPCR. Fold induction were

calculated by dividing the gene expression level of DOX-treated (+) by that of DOX-untreated (-) in pgR-poor

cells (**B**) or pgR-rich cells (**C**). Data represent mean ± SEM, n = 2, P values were calculated according to two-

way ANOVA. P > 0.05; n.s, not significant. (**D** and **E**) Relative gene expression of the indicated genes as measured

by RT-qPCR in DOX-treated (+) or DOX-untreated (-) pgR-poor cells (**D**) or pgR-rich cells (**E**). Data represent

mean ± SEM, n = 2, P values were calculated according to two-way ANOVA. P > 0.05; n.s, not significant.

S6 Fig. HIV-1 integrase proteins directly binds to host genomic R-loops. (A) Representative gel images for

EMSA of Sso7d-tagged HIV-1-integrase (E152Q) with different types of nucleic acids substrates (R:D+ssDNA

and Hybrid). 100 nM nucleic acid substrate was incubated with Sso7d-tagged HIV-1-integrase (E152Q) at 0 nM,

20 nM, 50 nM, 100 nM, 200 nM, and 400 nM (n = 3). (B) Nucleic acid extracts from FLAG-HIV-1-integrase-

expressing cells were immunoprecipitated using S9.6 antibody. gDNA was precipitated from the elutes of

immunoprecipitation and subjected to DNA–RNA hybrid dot blotting. Where indicated, the gDNA extracts were

either untreated (–) or treated (+) with RNase H after elution of immunoprecipitated materials. (C) Summary of

the experimental design for R-loop immunoprecipitation using S9.6 antibody in FLAG-tagged HIV-1 integrase

protein-expressing HeLa cells with pre-immunoprecipitation in vitro RNase H treatment. (**D**) Protein extracts

from FLAG-HIV-1-integrase-expressing cells were immunoprecipitated using anti-FLAG antibody. Western blot

of FLAG immunoprecipitation was probed with anti-FLAG or anti-H3 antibodies. (**E**) Representative images of

the proximity-ligation assay (PLA) using single antibody (anti-GFP or anti-S9.6) in HIV-IN-EGFP virion-infected

HeLa cells at 6 hpi, as PLA signal negative controls. Cells were subjected to PLA (orange) and co-stained with

DAPI (blue) (n > 50).

HeLa				
Gene	**Chromosom**	**Position (hg38)**	**Description**	**Average DRIPc-seq signal**
RPL13A	chr19	49487608-49493057	Input (-)_0hpi	3.59
			Input (-)_3hpi	0.24
			Input (-)_6hpi	2.39
			Input (-)_12hpi	3.51
			Input (+)_0hpi	82.29
			Input (+)_3hpi	51.76
			Input (+)_6hpi	39.14
			Input (+)_12hpi	176.73
			IP_RNase H- (-)_0hpi	2.21
			IP_RNase H- (-)_3hpi	2.73
			IP_RNase H- (-)_6hpi	2.39
			IP_RNase H- (-)_12hpi	4.25
			IP_RNase H- (+)_0hpi	110.32
			IP_RNase H- (+)_3hpi	140.22
			IP_RNase H- (+)_6hpi	58.36
			IP_RNase H- (+)_12hpi	137.37
			IP_RNase H+ (-)_0hpi	0.00
			IP_RNase H+ (-)_3hpi	4.48
			IP_RNase H+ (-)_6hpi	3.74
			IP_RNase H+ (-)_12hpi	0.00
			IP_RNase H+ (+)_0hpi	1.98
			IP_RNase H+ (+)_3hpi	3.36
			IP_RNase H+ (+)_6hpi	1.60
			IP_RNase H+ (+)_12hpi	6.81
Gene	**Chromosom**	**Position (hg38)**	**Description**	**Average DRIPc-seq signal**
CALM3	chr19	46601330-46610782	Input (-)_0hpi	1.47
			Input (-)_3hpi	1.02
			Input (-)_6hpi	2.46
			Input (-)_12hpi	0.74
			Input (+)_0hpi	26.50
			Input (+)_3hpi	19.95
			Input (+)_6hpi	11.61
			Input (+)_12hpi	56.92
			IP_RNase H- (-)_0hpi	0.90
			IP_RNase H- (-)_3hpi	1.54
			IP_RNase H- (-)_6hpi	1.23
			IP_RNase H- (-)_12hpi	1.73
			IP_RNase H- (+)_0hpi	13.97
			IP_RNase H- (+)_3hpi	28.68
			IP_RNase H- (+)_6hpi	10.58
			IP_RNase H- (+)_12hpi	24.70
			IP_RNase H+ (-)_0hpi	0.71
			IP_RNase H+ (-)_3hpi	1.83
			IP_RNase H+ (-)_6hpi	2.78
			IP_RNase H+ (-)_12hpi	1.04
			IP_RNase H+ (+)_0hpi	2.12
			IP_RNase H+ (+)_3hpi	1.64
			IP_RNase H+ (+)_6hpi	2.26
			IP_RNase H+ (+)_12hpi	1.65
Gene	**Chromosom**	**Position (hg38)**	**Description**	**Average DRIPc-seq signal**
SNRPN	chr15	24823647-24978582	Input (-)_0hpi	1.46
			Input (-)_3hpi	1.27
			Input (-)_6hpi	1.34
			Input (-)_12hpi	1.76
			Input (+)_0hpi	1.21
			Input (+)_3hpi	0.81
			Input (+)_6hpi	1.25
			Input (+)_12hpi	0.41
			IP_RNase H- (-)_0hpi	0.45
			IP_RNase H- (-)_3hpi	0.47
			IP_RNase H- (-)_6hpi	0.37
			IP_RNase H- (-)_12hpi	0.05
			IP_RNase H- (+)_0hpi	0.37
			IP_RNase H- (+)_3hpi	0.24
			IP_RNase H- (+)_6hpi	0.54
			IP_RNase H- (+)_12hpi	0.07
			IP_RNase H+ (-)_0hpi	1.40
			IP_RNase H+ (-)_3hpi	0.93
			IP_RNase H+ (-)_6hpi	1.10
			IP_RNase H+ (-)_12hpi	1.31
			IP_RNase H+ (+)_0hpi	1.18
			IP_RNase H+ (+)_3hpi	1.12
			IP_RNase H+ (+)_6hpi	1.26
			IP_RNase H+ (+)_12hpi	1.10

923 **S1 Table. Chromosomal position and DRIPc-seq signal for referenced R-loop-positive and –negative regions**

924 **in HIV-1 infected HeLa cells**

				CD4+	
Gene	Chromosom	Position (hg38)	Description		Average DRIPc-seq signal
RPL13A	chr19	49487608-49493057	Input (-)_0hpi		2.33
			Input (-)_3hpi		1.51
			Input (-)_6hpi		2.56
			Input (-)_12hpi		0.77
			Input (+)_0hpi		2.91
			Input (+)_3hpi		1.94
			Input (+)_6hpi		2.36
			Input (+)_12hpi		2.19
			IP_RNase H- (-)_0hpi		0.00
			IP_RNase H- (-)_3hpi		3.63
			IP_RNase H- (-)_6hpi		0.00
			IP_RNase H- (-)_12hpi		0.00
			IP_RNase H- (+)_0hpi		144.19
			IP_RNase H- (+)_3hpi		77.26
			IP_RNase H- (+)_6hpi		130.86
			IP_RNase H- (+)_12hpi		190.08
			IP_RNase H+ (-)_0hpi		1.42
			IP_RNase H+ (-)_3hpi		0.00
			IP_RNase H+ (-)_6hpi		0.00
			IP_RNase H+ (-)_12hpi		0.00
			IP_RNase H+ (+)_0hpi		0.93
			IP_RNase H+ (+)_3hpi		0.00
			IP_RNase H+ (+)_6hpi		0.00
			IP_RNase H+ (+)_12hpi		2.28
Gene	Chromosom	Position (hg38)	Description		Average DRIPc-seq signal
CALM3	chr19	46601330-46610782	Input (-)_0hpi		4.58
			Input (-)_3hpi		4.64
			Input (-)_6hpi		2.96
			Input (-)_12hpi		4.04
			Input (+)_0hpi		3.62
			Input (+)_3hpi		3.65
			Input (+)_6hpi		3.40
			Input (+)_12hpi		4.11
			IP_RNase H- (-)_0hpi		0.00
			IP_RNase H- (-)_3hpi		0.00
			IP_RNase H- (-)_6hpi		0.00
			IP_RNase H- (-)_12hpi		2.70
			IP_RNase H- (+)_0hpi		108.23
			IP_RNase H- (+)_3hpi		183.80
			IP_RNase H- (+)_6hpi		87.73
			IP_RNase H- (+)_12hpi		181.80
			IP_RNase H+ (-)_0hpi		2.80
			IP_RNase H+ (-)_3hpi		0.00
			IP_RNase H+ (-)_6hpi		0.00
			IP_RNase H+ (-)_12hpi		1.94
			IP_RNase H+ (+)_0hpi		4.11
			IP_RNase H+ (+)_3hpi		1.19
			IP_RNase H+ (+)_6hpi		9.88
			IP_RNase H+ (+)_12hpi		6.17
Gene	Chromosom	Position (hg38)	Description		Average DRIPc-seq signal
SNRPN	chr15	24823647-24978582	Input (-)_0hpi		1.65
			Input (-)_3hpi		1.41
			Input (-)_6hpi		1.74
			Input (-)_12hpi		1.15
			Input (+)_0hpi		1.72
			Input (+)_3hpi		1.46
			Input (+)_6hpi		1.97
			Input (+)_12hpi		1.29
			IP_RNase H- (-)_0hpi		0.31
			IP_RNase H- (-)_3hpi		0.27
			IP_RNase H- (-)_6hpi		0.10
			IP_RNase H- (-)_12hpi		0.27
			IP_RNase H- (+)_0hpi		0.98
			IP_RNase H- (+)_3hpi		1.00
			IP_RNase H- (+)_6hpi		0.53
			IP_RNase H- (+)_12hpi		0.56
			IP_RNase H+ (-)_0hpi		0.94
			IP_RNase H+ (-)_3hpi		1.57
			IP_RNase H+ (-)_6hpi		0.00
			IP_RNase H+ (-)_12hpi		2.17
			IP_RNase H+ (+)_0hpi		1.37
			IP_RNase H+ (+)_3hpi		1.14
			IP_RNase H+ (+)_6hpi		1.42
			IP_RNase H+ (+)_12hpi		1.19

S2 Table. Chromosomal position and DRIPc-seq signal for referenced R-loop-positive and –negative regions

in HIV-1 infected primary CD4+ T cells

Jurkat				
Gene	**Chromosom**	**Position (hg38)**	**Description**	**Average DRIPc-seq signal**
RPL13A	chr19	49487608-49493057	Input (-)_0hpi	1.46
			Input (-)_3hpi	1.92
			Input (-)_6hpi	1.92
			Input (-)_12hpi	1.58
			Input (+)_0hpi	1.40
			Input (+)_3hpi	2.02
			Input (+)_6hpi	1.15
			Input (+)_12hpi	1.54
			IP_RNase H- (-)_0hpi	0.00
			IP_RNase H- (-)_3hpi	10.17
			IP_RNase H- (-)_6hpi	9.60
			IP_RNase H- (-)_12hpi	2.64
			IP_RNase H- (+)_0hpi	404.40
			IP_RNase H- (+)_3hpi	183.88
			IP_RNase H- (+)_6hpi	486.50
			IP_RNase H- (+)_12hpi	526.25
			IP_RNase H+ (-)_0hpi	0.00
			IP_RNase H+ (-)_3hpi	3.53
			IP_RNase H+ (-)_6hpi	0.00
			IP_RNase H+ (-)_12hpi	0.00
			IP_RNase H+ (+)_0hpi	6.13
			IP_RNase H+ (+)_3hpi	0.00
			IP_RNase H+ (+)_6hpi	0.00
			IP_RNase H+ (+)_12hpi	0.00
Gene	**Chromosom**	**Position (hg38)**	**Description**	**Average DRIPc-seq signal**
CALM3	chr19	46601330-46610782	Input (-)_0hpi	2.40
			Input (-)_3hpi	2.18
			Input (-)_6hpi	2.26
			Input (-)_12hpi	2.78
			Input (+)_0hpi	2.08
			Input (+)_3hpi	2.78
			Input (+)_6hpi	1.99
			Input (+)_12hpi	2.38
			IP_RNase H- (-)_0hpi	0.00
			IP_RNase H- (-)_3hpi	11.73
			IP_RNase H- (-)_6hpi	5.58
			IP_RNase H- (-)_12hpi	5.22
			IP_RNase H- (+)_0hpi	208.25
			IP_RNase H- (+)_3hpi	182.67
			IP_RNase H- (+)_6hpi	167.98
			IP_RNase H- (+)_12hpi	220.30
			IP_RNase H+ (-)_0hpi	0.00
			IP_RNase H+ (-)_3hpi	2.04
			IP_RNase H+ (-)_6hpi	0.00
			IP_RNase H+ (-)_12hpi	4.84
			IP_RNase H+ (+)_0hpi	13.84
			IP_RNase H+ (+)_3hpi	1.62
			IP_RNase H+ (+)_6hpi	4.37
			IP_RNase H+ (+)_12hpi	3.29
Gene	**Chromosom**	**Position (hg38)**	**Description**	**Average DRIPc-seq signal**
SNRPN	chr15	24823647-24978582	Input (-)_0hpi	1.75
			Input (-)_3hpi	1.94
			Input (-)_6hpi	1.87
			Input (-)_12hpi	1.84
			Input (+)_0hpi	1.86
			Input (+)_3hpi	1.89
			Input (+)_6hpi	1.81
			Input (+)_12hpi	1.73
			IP_RNase H- (-)_0hpi	0.12
			IP_RNase H- (-)_3hpi	0.00
			IP_RNase H- (-)_6hpi	0.17
			IP_RNase H- (-)_12hpi	0.00
			IP_RNase H- (+)_0hpi	2.43
			IP_RNase H- (+)_3hpi	2.19
			IP_RNase H- (+)_6hpi	2.23
			IP_RNase H- (+)_12hpi	2.36
			IP_RNase H+ (-)_0hpi	2.58
			IP_RNase H+ (-)_3hpi	3.46
			IP_RNase H+ (-)_6hpi	1.62
			IP_RNase H+ (-)_12hpi	1.87
			IP_RNase H+ (+)_0hpi	1.78
			IP_RNase H+ (+)_3hpi	2.38
			IP_RNase H+ (+)_6hpi	1.06
			IP_RNase H+ (+)_12hpi	1.43

929 **S3 Table. Chromosomal position and DRIPc-seq signal for referenced R-loop-positive and –negative regions**

930 **in HIV-1 infected Jurkat cells**

	gene_symbol	0hpi	3hpi	6hpi	12hpi	48hpi
P1	TOR1AIP2	1	1.163832	1.247899	1.024926	0.619497
P2	DVL1	1	0.781593	0.571348	0.901502	0.270459
P3	PKN2	1	1.280974	1.891552	1.31842	1.515107
N1	N/A	N/A	N/A	N/A	N/A	N/A
N2	CDK5RAP1	1	0.73977	0.775	1.143662	0.472377

933 **S4 Table. RNA-seq analysis of relative gene expression levels of P1-3 and N1,2 R-loop regions**

Oligonucleotides	Sequence 5' to 3'	Remark
PCR primer 1.0 P5	AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGA	amplification primer
PCR primer 2.0 P7	CAAGCAGAAGACGGCATACGAGAT	amplification primer
Index Adapter 1	GATCGGAAGAGCACACGTCTGAACTCCAGTCACATCACGATCTCGTATGCCGTCTTCTGCTTG	HeLa 0hpi Input replicate 1
Index Adapter 2	GATCGGAAGAGCACACGTCTGAACTCCAGTCACCGATGTATCTCGTATGCCGTCTTCTGCTTG	HeLa 3hpi Input replicate 1
Index Adapter 3	GATCGGAAGAGCACACGTCTGAACTCCAGTCACTTAGGCATCTCGTATGCCGTCTTCTGCTTG	HeLa 6hpi Input replicate 1
Index Adapter 4	GATCGGAAGAGCACACGTCTGAACTCCAGTCACTGACCAATCTCGTATGCCGTCTTCTGCTTG	HeLa 12hpi Input replicate 1
Index Adapter 5	GATCGGAAGAGCACACGTCTGAACTCCAGTCACACAGTGATCTCGTATGCCGTCTTCTGCTTG	HeLa 0hpi RNH-IP replicate 1
Index Adapter 6	GATCGGAAGAGCACACGTCTGAACTCCAGTCACGCCAATATCTCGTATGCCGTCTTCTGCTTG	HeLa 3hpi RNH-IP replicate 1
Index Adapter 7	GATCGGAAGAGCACACGTCTGAACTCCAGTCACCAGATCATCTCGTATGCCGTCTTCTGCTTG	HeLa 6hpi RNH-IP replicate 1
Index Adapter 8	GATCGGAAGAGCACACGTCTGAACTCCAGTCACACTTGAATCTCGTATGCCGTCTTCTGCTTG	HeLa 12hpi RNH-IP replicate 1
Index Adapter 9	GATCGGAAGAGCACACGTCTGAACTCCAGTCACGATCAGATCTCGTATGCCGTCTTCTGCTTG	HeLa 0hpi RNH+IP replicate 1
Index Adapter 10	GATCGGAAGAGCACACGTCTGAACTCCAGTCACTAGCTTATCTCGTATGCCGTCTTCTGCTTG	HeLa 3hpi RNH+IP replicate 1
Index Adapter 11	GATCGGAAGAGCACACGTCTGAACTCCAGTCACGGCTACATCTCGTATGCCGTCTTCTGCTTG	HeLa 6hpi RNH+IP replicate 1
Index Adapter 12	GATCGGAAGAGCACACGTCTGAACTCCAGTCACCTTGTAATCTCGTATGCCGTCTTCTGCTTG	HeLa 12hpi RNH+IP replicate 1
Index Adapter 28	GATCGGAAGAGCACACGTCTGAACTCCAGTCACCAAAAGATCTCGTATGCCGTCTTCTGCTTG	HeLa 0hpi Input replicate 2
Index Adapter 29	GATCGGAAGAGCACACGTCTGAACTCCAGTCACCAACTAATCTCGTATGCCGTCTTCTGCTTG	HeLa 3hpi Input replicate 2
Index Adapter 30	GATCGGAAGAGCACACGTCTGAACTCCAGTCACCACCGGATCTCGTATGCCGTCTTCTGCTTG	HeLa 6hpi Input replicate 2
Index Adapter 31	GATCGGAAGAGCACACGTCTGAACTCCAGTCACCACGATATCTCGTATGCCGTCTTCTGCTTG	HeLa 12hpi Input replicate 2
Index Adapter 32	GATCGGAAGAGCACACGTCTGAACTCCAGTCACCACTCAATCTCGTATGCCGTCTTCTGCTTG	HeLa 0hpi RNH-IP replicate 2
Index Adapter 33	GATCGGAAGAGCACACGTCTGAACTCCAGTCACCAGGCGATCTCGTATGCCGTCTTCTGCTTG	HeLa 3hpi RNH-IP replicate 2
Index Adapter 34	GATCGGAAGAGCACACGTCTGAACTCCAGTCACCATGGCATCTCGTATGCCGTCTTCTGCTTG	HeLa 6hpi RNH-IP replicate 2
Index Adapter 35	GATCGGAAGAGCACACGTCTGAACTCCAGTCACCATTTATCTCGTATGCCGTCTTCTGCTTG	HeLa 12hpi RNH-IP replicate 2
Index Adapter 36	GATCGGAAGAGCACACGTCTGAACTCCAGTCACCCAACAATCTCGTATGCCGTCTTCTGCTTG	HeLa 0hpi RNH+IP replicate 2
Index Adapter 37	GATCGGAAGAGCACACGTCTGAACTCCAGTCACCGGAATATCTCGTATGCCGTCTTCTGCTTG	HeLa 3hpi RNH+IP replicate 2
Index Adapter 38	GATCGGAAGAGCACACGTCTGAACTCCAGTCACCTAGCTATCTCGTATGCCGTCTTCTGCTTG	HeLa 6hpi RNH+IP replicate 2
Index Adapter 39	GATCGGAAGAGCACACGTCTGAACTCCAGTCACCTATACATCTCGTATGCCGTCTTCTGCTTG	HeLa 12hpi RNH+IP replicate 2

936 **S5 Table. Oligonucleotides used for DRIPc-seq library construction**

Oligonucleotides	Sequence 5' to 3'
P1 Fwd	TTATAAGTCAGCCTCCAGGATCAA
P1 Rev	TTCAGGTCTAGGCAGTCTGA
P2 Fwd	GGA CAG ATG ACA GGG TCG C
P2 Rev	ATG AGG AAG ACC CCC TCG G
P3 Fwd	CTCTGTGTAACGCTGGTGCT
P3 Rev	ACACGCTTCTGACCACTAAGG
N1 Fwd	TTG GCC CTA CTG AAT GAT GGG T
N1 Rev	TTA AGG CAT GCT CAG GCG A
N2 Fwd	TGA GAT TTC AGG TTC CAT GAT TTG
N2 Rev	TGC TCA GTG TTC TAA TTT CCC TGT
β-actin Fwd	AGAGCTACGAGCTGCCTGAC
β-actin Rev	AGCACTGTGTTGGCGTACAG
SH49 (ECFP Fwd)	TGGTTTGTCCAAACTCATCAA
SH40 (mAIRN Fwd)	CGAGAGAGGCTAAGGGTGAA
SH21 (ECFP/mAIRN Rev)	ACATGGTCCTGCTGGAGTTC
RT-qPCR P1 (TOR1AIP2) Fwd	CCTTGGTCTTTCCCACTTGAGTG
RT-qPCR P1 (TOR1AIP2) Rev	GCAGGGTTAAAACCAGCTACTCG
RT-qPCR P2 (DVL1) Fwd	GCATAACCGACTCCACCATGTC
RT-qPCR P2 (DVL1) Rev	GATGGAGCCAATGTAGATGCCG
RT-qPCR P3 (PKN2) Fwd	GCATCACCAACACTAAGTCCACG
RT-qPCR P3 (PKN2) Rev	GCTTTTGACCGTCCAGGGACAT
RT-qPCR N2 (CDK5RAP1) Fwd	AGAGTGGAAGCAGCCGTGTGTT
RT-qPCR N2 (CDK5RAP1) Rev	GATCTTCCTCCGTCTCACCACA

938 **S6 Table. Primers used for qPCR**

Oligonucleotides	Sequence 5' to 3'	Remark
AE5316	TGTGACTCTGGTAACTAGAGATCCCTC	First round LTR primer
AE6380	TAGTCCCTTAAGCGGAG-NH2	replicate 1 5dpi Linker short / replicate 1 pgR-poor DOX- Linker short / CD4+ donor 1 Linker short
AE6381	GTAATACGACTCACTATAGGGCCTCCGCTTAAGGGAC	replicate 1 5dpi Linker long / replicate 1 pgR-poor DOX- Linker long / CD4+ donor 1 Linker long
AE6382	CAAGCAGAAGACGGCATACGAGATCGGGTCTCGGCATTCCTGCTGAACCGCTCTTCCGATCTGTAATACGACTCACTATAGGGC	replicate 1 5dpi Linker primer / replicate 1 pgR-poor DOX- Linker primer / CD4+ donor 1 Linker primer
AE6404	AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCGATCTCGATGTGAGATCCCTCAGACCCTTTAGTCAG	replicate 1 5dpi Second round LTR primer / replicate 1 pgR-poor DOX- Second round LTR primer / CD4+ donor 1 Second round LTR primer
AE6380	TAGTCCCTTAAGCGGAG-NH2	replicate 2 5dpi Linker short / replicate 2 pgR-poor DOX+ Linker short
AE6381	GTAATACGACTCACTATAGGGCCTCCGCTTAAGGGAC	replicate 2 5dpi Linker long / replicate 2 pgR-poor DOX+ Linker long
AE6382	CAAGCAGAAGACGGCATACGAGATCGGGTCTCGGCATTCCTGCTGAACCGCTCTTCCGATCTGTAATACGACTCACTATAGGGC	replicate 2 5dpi Linker primer / replicate 2 pgR-poor DOX+ Linker prime
AE6404-1	AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCGATCTTAGGCGAGATCCCTCAGACCCTTTAGTCAG	replicate 2 5dpi Second round LTR primer / replicate 2 pgR-poor DOX+ Second round LTR primer
AE6386	TACTATGACGGTGACGC-NH2	replicate 1 pgR-rich DOX- Linker short / CD4+ donor 2 Linker short
AE6387	GAGAATCCATGAGTATGCTCACGCGTCACCGTCATAG	replicate 1 pgR-rich DOX- Linker long / CD4+ donor 2 Linker long
AE6388	CAAGCAGAAGACGGCATACGAGATCGGGTCTCGGCATTCCTGCTGAACCGCTCTTCCGATCTGAGAATCCATGAGTAGTGCTCAC	replicate 1 pgR-rich DOX- Linker primer / CD4+ donor 2 Linker primer
AE6406	AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCGATCTACAGTGGAGATCCCTCAGACCCTTTAGTCAG	replicate 1 pgR-rich DOX- Second round LTR primer / CD4+ donor 2 Second round LTR primer
AE6456	TAGACTGACGCAGTCTG-NH2	replicate 1 pgR-poor DOX+ Linker short
AE6457	GACGTACATACTGATCGCATAGCAGACTGCGTCAGTC	replicate 1 pgR-poor DOX+ Linker long
AE6458	CAAGCAGAAGACGGCATACGAGATCGGGTCTCGGCATTCCTGCTGAACCGCTCTTCCGATCTGACGTACATACTGATCGCATAG	replicate 1 pgR-poor DOX+ Linker primer
AE6405	AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCGATCTTGACCAGAGATCCCTCAGACCCTTTAGTCAG	replicate 1 pgR-poor DOX+ Second round LTR primer
AE6386	TACTATGACGGTGACGC-NH2	replicate 2 pgR-rich DOX+ Linker short
AE6387	GAGAATCCATGAGTATGCTCACGCGTCACCGTCATAG	replicate 2 pgR-rich DOX+ Linker long
AE6388	CAAGCAGAAGACGGCATACGAGATCGGGTCTCGGCATTCCTGCTGAACCGCTCTTCCGATCTGAGAATCCATGAGTAGTGCTCAC	replicate 2 pgR-rich DOX+ Linker primer
AE6406-1	AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCGATCTGCCAATGAGATCCCTCAGACCCTTTAGTCAG	replicate 2 pgR-rich DOX+ Second round LTR primer
AE6456	TAGACTGACGCAGTCTG-NH2	replicate 3 pgR-rich DOX- Linker short
AE6457	GACGTACATACTGATCGCATAGCAGACTGCGTCAGTC	replicate 3 pgR-rich DOX- Linker long
AE6458	CAAGCAGAAGACGGCATACGAGATCGGGTCTCGGCATTCCTGCTGAACCGCTCTTCCGATCTGACGTACATACTGATCGCATAG	replicate 3 pgR-rich DOX- Linker primer
AE6411	AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCGATCTAGTTCCGAGATCCCTCAGACCCTTTAGTCAG	replicate 3 pgR-rich DOX- Second round LTR primer
AE6392	TACTGAGACGTCGATGC-NH2	replicate 1 RNH_mut 5dpi Linker short
AE6393	GATCATGCGAGATACATCTCAGGCATCGACGTCTCAG	replicate 1 RNH_mut 5dpi Linker long
AE6394	CAAGCAGAAGACGGCATACGAGATCGGGTCTCGGCATTCCTGCTGAACCGCTCTTCCGATCTGATCATGCGAGATACATCTCAG	replicate 2 RNH_mut 5dpi Linker primer
AE6493	AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCGATCTGGCTACGAGATCCCTCAGACCCTTTAGTCAG	replicate 1 RNH_mut 5dpi Second round LTR primer
AE6493-1	AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCGATCTACTTGAGAGATCCCTCAGACCCTTTAGTCAG	replicate 2 RNH_mut 5dpi Second round LTR primer
AE6462	TAGTAGTCACGAGCCGTC-NH2	replicate 1 RNH_wt 5dpi Linker short / replicate 2 RNH_wt 5dpi Linker short
AE6463	CAGTTAGACTACACGTTAGACGGACGCCTCGTGACTAC	replicate 1 RNH_wt 5dpi Linker long / replicate 2 RNH_wt 5dpi Linker long
AE6464	CAAGCAGAAGACGGCATACGAGATCGGGTCTCGGCATTCCTGCTGAACCGCTCTTCCGATCTCAGTTAGACTACACGTTAGACG	replicate 1 RNH_wt 5dpi Linker primer / replicate 2 RNH_wt 5dpi Linker primer
AE6492	AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCGATCTTAGCTTGAGATCCCTCAGACCCTTTAGTCAG	replicate 1 RNH_wt 5dpi Second round LTR primer
AE6497	AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCGATCTATCACGGAGATCCCTCAGACCCTTTAGTCAG	replicate 2 RNH_wt 5dpi Second round LTR primer

940 **S7 Table. Oligonucleotides used for HIV-1 integration site sequencing library construct**

Oligonucleotides	Sequence 5' to 3'	Remark
R-loop oligo1*	5'-[Cy3]-GCC AGG GAC GAG GTG AAC CTG CAG GTG GGC GGC TAC TAC TTA GAT GTC ATC CGA GGC TTA TTG GTA GAA TTC GGC AGC GTC ATG C GA CGG C-3'	R-loop, R:D+ssDNA, ssDNA
R-loop oligo2*	5'-GCC GTC GCA TGA CGC TGC CGA ATT CTA CCA CGC GAT TCA TAC CTG TCG TGC CAG CTG CTT TGC CCA CCT GCA GGT TCA CCT CGT CCC TGG C-3'	R-loop, dsDNA
R-loop RNA	5'-[Cy5]-GCA GCU GGC ACG ACA GGU AUG AAU C-3'	R-loop, R:D+ssDNA, ssRNA
Homoduplex	5'-[Cy3]-GCC AGG GAC GAG GTG AAC CTG CAG GTG GGC AAA GCA GCT GGC ACG ACA GGT ATG AAT CGC GTG GTA GAA TTC GGC AGC GTC ATG CGA CGG C-3'	dsDNA
Hybrid DNA	5'-CCC ATA CCG TAT AAC CAT TTG GCT GTC CAA GCT CCG GGT-3'	Hybrid
Hybrid RNA	5'-[Cy5]-ACC CGG AGC UUG GAC AGC CAA AUG GUU AUA CGG UAU GGG-3'	Hybrid
oligo 5	5'GCAGTAGCATGACGCTGCTGAATTCTACCACGCTATGCT CTCGTCTAGGTTCACTCCGT CCCTGCGATTCATACCTGTCGTGCCAGCTGC	R:D+ssDNA

942 **S8 Table. Oligonucleotides used for electrophoretic mobility shift assay substrate preparation**

Data	Accession Number/ Website
Jurkat integration site	SRR12322252
TSA-seq_SPAD	SRR3538917, SRR3538918, SRR3538919, SRR3538920
SPIN (Spatial Position Inference of the Nuclear genome) annotation of speckle	https://github.com/ma-compbio/SPIN
LADs	GSE22428

944 **S9 Table. Accession numbers and data sources.**

References

1. W. E. Johnson, Origins and evolutionary consequences of ancient endogenous retroviruses. *Nat Rev Microbiol* **17**, 355-370 (2019).

2. M. Lusic, R. F. Siliciano, Nuclear landscape of HIV-1 infection and integration. *Nat Rev Microbiol* **15**, 69-82 (2017).

3. H. C. Chen, J. P. Martinez, E. Zorita, A. Meyerhans, G. J. Filion, Position effects influence HIV latency reversal. *Nat Struct Mol Biol* **24**, 47-54 (2017).

4. K. B. Einkauf *et al.*, Parallel analysis of transcription, integration, and sequence of single HIV-1 proviruses. *Cell* **185**, 266-282 e215 (2022).

5. C. Jiang *et al.*, Distinct viral reservoirs in individuals with spontaneous control of HIV-1. *Nature* **585**, 261-267 (2020).

6. A. R. Schroder *et al.*, HIV-1 integration in the human genome favors active genes and local hotspots. *Cell* **110**, 521-529 (2002).

7. V. Achuthan *et al.*, Capsid-CPSF6 Interaction Licenses Nuclear HIV-1 Trafficking to Sites of Viral DNA Integration. *Cell Host Microbe* **24**, 392-404 e398 (2018).

8. A. Ciuffi *et al.*, A role for LEDGF/p75 in targeting HIV DNA integration. *Nat Med* **11**, 1287-1289 (2005).

9. G. A. Sowd *et al.*, A critical role for alternative polyadenylation factor CPSF6 in targeting HIV-1 integration to transcriptionally active chromatin. *Proc Natl Acad Sci U S A* **113**, E1054-1063 (2016).

10. B. Lucic *et al.*, Spatially clustered loci with multiple enhancers are frequent targets of HIV-1 integration. *Nat Commun* **10**, 4059 (2019).

11. B. Marini *et al.*, Nuclear architecture dictates HIV-1 integration site selection. *Nature* **521**, 227-231 (2015).

12. M. Kvaratskhelia, A. Sharma, R. C. Larue, E. Serrao, A. Engelman, Molecular mechanisms of retroviral integration site selection. *Nucleic Acids Res* **42**, 10209-10225 (2014).

13. P. Cherepanov *et al.*, HIV-1 integrase forms stable tetramers and associates with LEDGF/p75 protein in human cells. *J Biol Chem* **278**, 372-381 (2003).

14. J. M. Coffin *et al.*, Integration in oncogenes plays only a minor role in determining the in vivo distribution of HIV integration sites before or during suppressive antiretroviral therapy. *PLoS Pathog* **17**, e1009141 (2021).

15. C. Niehrs, B. Luke, Regulatory R-loops as facilitators of gene expression and genome stability. *Nat Rev Mol Cell Biol* **21**, 167-178 (2020).

16. E. Petermann, L. Lan, L. Zou, Sources, resolution and physiological relevance of R-loops and RNA-DNA hybrids. *Nat Rev Mol Cell Biol* **23**, 521-540 (2022).

17. S. Hamperl, M. J. Bocek, J. C. Saldivar, T. Swigut, K. A. Cimprich, Transcription-Replication

Conflict Orientation Modulates R-Loop Levels and Activates Distinct DNA Damage
Responses. *Cell* **170**, 774-786 e719 (2017).

18.   P. A. Ginno, P. L. Lott, H. C. Christensen, I. Korf, F. Chedin, R-loop formation is a distinctive
characteristic of unmethylated human CpG island promoters. *Mol Cell* **45**, 814-825 (2012).

19.   Y. W. Lim, L. A. Sanz, X. Xu, S. R. Hartono, F. Chedin, Genome-wide DNA hypomethylation
and RNA:DNA hybrid accumulation in Aicardi-Goutieres syndrome. *Elife* **4**, (2015).

20.   R. Arora *et al.*, RNaseH1 regulates TERRA-telomeric DNA hybrids and telomere maintenance
in ALT tumour cells. *Nat Commun* **5**, 5220 (2014).

21.   T. Garcia-Muse, A. Aguilera, R Loops: From Physiological to Pathological Roles. *Cell* **179**,
604-618 (2019).

22.   L. A. Sanz *et al.*, Prevalent, Dynamic, and Conserved R-Loop Structures Associate with Specific
Epigenomic Signatures in Mammals. *Mol Cell* **63**, 167-178 (2016).

23.   F. Chedin, Nascent Connections: R-Loops and Chromatin Patterning. *Trends Genet* **32**, 828-
838 (2016).

24.   C. Y. Lee *et al.*, R-loop induced G-quadruplex in non-template promotes transcription by
successive R-loop formation. *Nat Commun* **11**, 3392 (2020).

25.   H. O. Ajoge *et al.*, G-Quadruplex DNA and Other Non-Canonical B-Form DNA Motifs
Influence Productive and Latent HIV-1 Integration and Reactivation Potential. *Viruses* **14**,
(2022).

26.   F. Chedin, C. J. Benham, Emerging roles for R-loop structures in the management of
topological stress. *J Biol Chem* **295**, 4684-4695 (2020).

27.   I. K. Jozwik *et al.*, B-to-A transition in target DNA during retroviral integration. *Nucleic Acids*
Res **50**, 8898-8918 (2022).

28.   A. Ballandras-Colas *et al.*, Multivalent interactions essential for lentiviral integrase function.
Nat Commun **13**, 2416 (2022).

29.   L. A. Sanz, F. Chedin, High-resolution, strand-specific R-loop mapping via S9.6-based DNA-
RNA immunoprecipitation and high-throughput sequencing. *Nat Protoc* **14**, 1734-1755
(2019).

30.   R. B. Jones *et al.*, LINE-1 retrotransposable element DNA accumulates in HIV-1-infected cells.
J Virol **87**, 13307-13320 (2013).

31.   S. Srinivasachar Badarinarayan *et al.*, HIV-1 infection activates endogenous retroviral
promoters regulating antiviral gene expression. *Nucleic Acids Res* **48**, 10890-10908 (2020).

32.   P. Lesbats, A. N. Engelman, P. Cherepanov, Retroviral DNA Integration. *Chem Rev* **116**, 12730-
12757 (2016).

33.   A. Brussel, P. Sonigo, Analysis of early human immunodeficiency virus type 1 DNA synthesis
by use of a new sensitive assay for quantifying integrated provirus. *J Virol* **77**, 10119-10124
(2003).

34.   A. Albanese, D. Arosio, M. Terreni, A. Cereseto, HIV-1 pre-integration complexes selectively

1020		target decondensed chromatin in the nuclear periphery. *PLoS One* **3**, e2413 (2008).
1021	35.	A. Dharan, N. Bachmann, S. Talley, V. Zwikelmaier, E. M. Campbell, Nuclear pore blockade
1022		reveals that HIV-1 completes reverse transcription and uncoating in the nucleus. *Nat*
1023		*Microbiol* **5**, 1088-1095 (2020).
1024	36.	P. A. Ginno, Y. W. Lim, P. L. Lott, I. Korf, F. Chedin, GC skew at the 5' and 3' ends of human
1025		genes links R-loop formation to epigenetic regulation and transcription termination.
1026		*Genome Res* **23**, 1590-1600 (2013).
1027	37.	J. J. Kessl *et al.*, HIV-1 Integrase Binds the Viral RNA Genome and Is Essential during Virion
1028		Morphogenesis. *Cell* **166**, 1257-1268 e1212 (2016).
1029	38.	D. C. van Gent, Y. Elgersma, M. W. Bolk, C. Vink, R. H. Plasterk, DNA binding properties of
1030		the integrase proteins of human immunodeficiency viruses types 1 and 2. *Nucleic Acids Res*
1031		**19**, 3821-3827 (1991).
1032	39.	A. Cristini, M. Groh, M. S. Kristiansen, N. Gromak, RNA/DNA Hybrid Interactome Identifies
1033		DXH9 as a Molecular Player in Transcriptional Termination and R-Loop-Associated DNA
1034		Damage. *Cell Rep* **23**, 1891-1905 (2018).
1035	40.	T. Mosler *et al.*, R-loop proximity proteomics identifies a role of DDX41 in transcription-
1036		associated genomic instability. *Nat Commun* **12**, 7314 (2021).
1037	41.	R. Schrijvers *et al.*, LEDGF/p75-independent HIV-1 replication demonstrates a role for HRP-
1038		2 and remains sensitive to inhibition by LEDGINs. *PLoS Pathog* **8**, e1002558 (2012).
1039	42.	P. C. Stirling, P. Hieter, Canonical DNA Repair Pathways Influence R-Loop-Driven Genome
1040		Instability. *J Mol Biol* **429**, 3132-3138 (2017).
1041	43.	M. L. Garcia-Rubio *et al.*, The Fanconi Anemia Pathway Protects Genome Integrity from R-
1042		loops. *PLoS Genet* **11**, e1005674 (2015).
1043	44.	M. Giannini *et al.*, TDP-43 mutations link Amyotrophic Lateral Sclerosis with R-loop
1044		homeostasis and R loop-mediated DNA damage. *PLoS Genet* **16**, e1009260 (2020).
1045	45.	S. Fu *et al.*, HIV-1 exploits the Fanconi anemia pathway for viral DNA integration. *Cell Rep*
1046		**39**, 110840 (2022).
1047	46.	D. Li, A. Lopez, C. Sandoval, R. Nichols Doyle, O. I. Fregoso, HIV Vpr Modulates the Host
1048		DNA Damage Response at Two Independent Steps to Damage DNA and Repress Double-
1049		Strand DNA Break Repair. *mBio* **11**, (2020).
1050	47.	H. Bauby *et al.*, HIV-1 Vpr Induces Widespread Transcriptomic Changes in CD4(+) T Cells
1051		Early Postinfection. *mBio* **12**, e0136921 (2021).
1052	48.	K. Stopak, C. de Noronha, W. Yonemoto, W. C. Greene, HIV-1 Vif blocks the antiviral activity
1053		of APOBEC3G by impairing both its translation and intracellular stability. *Mol Cell* **12**, 591-
1054		601 (2003).
1055	49.	D. Kmiec, F. Kirchhoff, Antiviral factors and their counteraction by HIV-1: many uncovered
1056		and more to be discovered. *J Mol Cell Biol*, (2024).
1057	50.	J. L. McCann *et al.*, R-loop homeostasis and cancer mutagenesis promoted by the DNA

cytosine deaminase APOBEC3B. 2021.2008.2030.458235 (2021).

51.   S. A. Yukl *et al.*, HIV latency in isolated patient CD4(+) T cells may be due to blocks in HIV
transcriptional elongation, completion, and splicing. *Sci Transl Med* **10**,   (2018).

52.   D. O. Passos *et al.*, Cryo-EM structures and atomic model of the HIV-1 strand transfer
complex intasome. *Science* **355**, 89-92 (2017).

53.   H. D. Nguyen *et al.*, Functions of Replication Protein A as a Sensor of R Loops and a
Regulator of RNaseH1. *Mol Cell* **65**, 832-847 e834 (2017).

54.   S. Andrews. (2010).

55.   F. J. Felix Krueger, Phil Ewels, Ebrahim Afyounian, & Benjamin Schuster-Boeckler,
FelixKrueger/TrimGalore: v0.6.7 - DOI via Zenodo (0.6.7). Zenodo. (2021).

56.   M. Martin, Cutadapt removes adapter sequences from high-throughput sequencing reads.
2011 **17**, 3 %J EMBnet.journal (2011).

57.   H. Li, R. Durbin, Fast and accurate short read alignment with Burrows-Wheeler transform.
Bioinformatics **25**, 1754-1760 (2009).

58.   Y. Zhang *et al.*, Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9**, R137 (2008).

59.   A. R. Quinlan, I. M. Hall, BEDTools: a flexible suite of utilities for comparing genomic features.
Bioinformatics **26**, 841-842 (2010).

60.   W. Li *et al.*, CPSF6-Dependent Targeting of Speckle-Associated Domains Distinguishes
Primate from Nonprimate Lentiviral Integration. *mBio* **11**,   (2020).

61.   H. M. Amemiya, A. Kundaje, A. P. Boyle, The ENCODE Blacklist: Identification of Problematic
Regions of the Genome. *Sci Rep* **9**, 9354 (2019).

62.   A. Frankish *et al.*, Gencode 2021. *Nucleic Acids Res* **49**, D916-D923 (2021).

63.   A. Dobin *et al.*, STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21 (2013).

64.   Y. Jin, O. H. Tam, E. Paniagua, M. Hammell, TEtranscripts: a package for including
transposable elements in differential expression analysis of RNA-seq datasets. *Bioinformatics*
31, 3593-3599 (2015).

1084