The Rhizobial effector NopT targets Nod factor receptors to regulate symbiosis in *Lotus japonicus*

Hanbin Bao[1#], Yanan Wang[1#], Haoxing Li[1], Qiang Wang[1], Yutao Lei[1], Ying Ye[1], Syed F. Wadood[2], Hui Zhu[1], Christian Stahelin[2], Gary Stacey[3], Shutong Xu[1], Yangrong Cao[1*]

[1] National Key Lab of Agricultural Microbiology, Hubei Hongshan Laboratory, Huazhong Agricultural University, Wuhan, Hubei, 430070, China

[2] State Key Laboratory of Biocontrol and Guangdong Key Laboratory of Plant Resources, School of Life Sciences, Sun Yat-sen University, Guangzhou, 510275, China

[3] Divisions of Plant Science and Technology and Biochemistry, Christopher S. Bond Life Sciences Center, University of Missouri, Columbia, MO 65211, USA

# These authors contributed equally to this work.

* Corresponding author: Yangrong Cao, email: yrcao@mail.hzau.edu.cn

Short title: NopT associates with NFR1 and NFR5 to regulate symbiosis

Keywords: Legume-rhizobium symbiosis; Rhizobial effector; Nod factor receptors; NopT.

One-sentence summary: The rhizobial effector NopT and two Nod factor receptors of *Lotus japonicus*, NFR1 and NFR5, mutually interact to precisely regulate rhizobial symbiosis.

Abstract

Abstract

It is well-documented that type-III effectors are required by Gram-negative pathogens to

directly target different host cellular pathways to promote bacterial infection. However,

in the context of legume-rhizobium symbiosis, the role of rhizobial effectors in regulating

plant symbiotic pathways remains largely unexplored. Here, we show that NopT, a

YopT-type cysteine protease of *Sinorhizobium fredii* NGR234 directly targets the plant's

symbiotic signaling pathway by associating with two Nod factor receptors (NFR1 and

NFR5 of *Lotus japonicus*). NopT inhibits cell death triggered by co-expression of

NFR1/NFR5 in *Nicotiana benthamiana*. Full-length NopT physically interacts with

NFR1 and NFR5. NopT proteolytically cleaves NFR5 both *in vitro* and *in vivo*, but can

be inactivated by NFR1 as a result of phosphorylation. NopT plays an essential role in

mediating rhizobial infection in *L. japonicus*. Autocleaved NopT retains the ability to

cleave NFR5 but no longer interacts with NFR1. Interestingly, genomes of certain

Sinorhizobium species only harbor *nopT* genes encoding truncated proteins without the

autocleavage site. These results reveal an intricate interplay between rhizobia and

legumes, in which a rhizobial effector protease targets NFR5 to suppress symbiotic

signaling. NFR1 appears to counteract this process by phosphorylating the effector. This

discovery highlights the role of a bacterial effector in regulating a signaling pathway in

plants and opens up the perspective of developing kinase-interacting proteases to

fine-tune cellular signaling processes in general.

Introduction

The legume–rhizobium symbiosis stands as one of the most crucial and intricate

mutualistic interactions in nature. Its significance goes beyond the supply of nitrogen

fixed by rhizobia, which influences plant growth and ecosystem sustainability, as this symbiosis is a pivotal model for investigating plant-microbe interactions (Yang et al., 2022). The orchestration of this symbiosis involves the perception of rhizobial signal molecules such as lipo–chitooligosaccharidic Nod factors (NFs) and surface polysaccharides (Madsen et al., 2003; Tirichine et al., 2006; Kawaharada et al., 2015), along with the secretion of proteins, including type three protein secretion system (T3SS) effectors translocated into host plants (Sugawara et al., 2018; Zhang et al., 2021).

Plant receptor heterocomplexes consisting of two Lysin-motif receptor kinases (LYKs), play a central role in recognizing various microbial poly- and oligosaccharide molecules. In non–legumes such as *Arabidopsis*, AtLYK5 and AtCERK1 (Chitin elicitor receptor kinase) are required for perception of chitin, and thereby trigger plant immunity to combat fungal pathogens (Cao et al., 2014). In legumes, structurally similar Nod factor receptors (NFRs) such as NFR1 and NFR5 in *Lotus japonicus* are vital for recognition of rhizobial NFs and initiation of symbiotic signaling (Broghammer et al., 2012; Bozsoki et al., 2017). Accordingly, *nfr1* and *nfr5* knockout mutants are almost completely unable to form nodules (Madsen et al., 2003; Radutoiu et al., 2003). Over-expression of either NFR1 or NFR5 can activate NF signaling, resulting in formation of spontaneous nodules in the absence of rhizobia (Ried et al., 2014). The intricate action of NFRs is highlighted by the elicitation of a hypersensitive cell death response when NFR1/NFR5 are simultaneous over-expressed in *Nicotiana benthamiana* leaves (Madsen et al., 2011). Similarly, defense-like responses in *Medicago truncatula* nodules were observed upon over–expression of MtNFP (Nod factor perception, the ortholog of NFR5) (Moling et al., 2014), perhaps reflecting a potential role of MtNFP–interacting LYKs involved in plant–pathogen associations. It can therefore be expected that the protein abundance of NFRs must be tightly controlled to avoid resistance responses, thereby ensuring optimal root infection by rhizobia.

T3SS effectors of phytopathogenic Gram–negative bacteria play a critical role in suppressing pattern-triggered immunity (PTI) in host plants (Jones and Dangl, 2006). To avoid diseases, specific resistance (R) genes are employed to directly or indirectly recognize effectors translocated into host cells. Recognition of such avirulence (Avr) effectors often results in effector-triggered immunity (ETI), a much stronger immune response which is usually associated with programmed cell death (Jones and Dangl, 2006). In contrast to the well–studied effector functions of phytopathogenic bacteria in suppressing immunity through various mechanisms, it is largely unknown how T3SS effectors from symbiotic rhizobia contribute to mutualistic interactions.

T3SS effectors of the YopT–type family, such as YopT of *Yersinia pestis*, AvrPphB of *Pseudomonas syringae* pv. *phaseolicola* and rhizobial NopT proteins, are evolutionary related cysteine proteases (Shao et al., 2002; Dai et al., 2008). YopT cleaves Rho family GTPases leading to their inactivation to disrupt the actin cytoskeleton of human host cells (Shao et al., 2003a). AvrPphB can proteolytically cleave *Arabidopsis* receptor–like cytoplasmic kinase family proteins including PBS1, PBS1-like proteins and BIK1, to suppress PTI (Shao et al., 2003b; Zhang et al., 2010). Cleavage of PBS1 kinase by AvrPphB can be regarded as a "decoy strategy" as it is associated with RPS5 (RESISTANCE TO PSEUDOMONAS SYRINGAE 5)-mediated ETI (Shao et al., 2003b). Depending on the host legume, the NopT effector protease of *Sinorhizobium fredii* NGR234 can have an opposite function in regulating symbiosis (Dai et al., 2008; Kambara et al., 2009). The heterologous expression of NopT from NGR234 in *S. fredii* USDA257 severely blocks nodulation in soybean cultivar Nanfeng 15 but not in other soybean cultivars tested (Khan et al., 2022), implying that NopT might act as an Avr effector. In

contrast, NopT positively regulates nodule formation in the interaction between *S. fredii* HH103 and soybean cultivar Suinong 14 (Li et al., 2023). NopT from *Bradyrhizobium* sp. ORS3257 promotes rhizobial infection in nodules of *Aeschynomene indica* (Teulet et al., 2019). NopT proteases have been biochemically characterized. Some exhibit autocleavage activity and are subsequently acylated at the newly formed N-terminus (Dai et al., 2008; Dowen et al., 2009; Kambara et al., 2009; Fotiadis et al., 2012; Khan et al., 2022). Recently, soybean PBS1-1 and two proteins in *Robinia pseudoacacia* were identified as NopT targets (Luo et al., 2020; Khan et al. 2022; Li et al. 2023). However, at the molecular level, it remains largely unknown how NopT modulates rhizobial infection and whether NopTs from different rhizobial strains differ in their effector activities.

A challenge in the investigation of rhizobial effectors lies in the availability of suitable assays and plants with which effector activities can be characterized. In this study, we took advantage of the observation that ectopic expression of NFR1 and NFR5 in *N. benthamiana* leaves induces programmed cell death (Madsen et al., 2011). We hypothesized that this response could be modulated by co-expressed rhizobial effectors. Among the 15 known or putative T3SS effectors of *S. fredii* NGR234, only NopT could suppress NFR1/NFR5-induced cell death. This observation suggested that NopT is an effector associated with NFRs. Subsequent experiments demonstrated that NopT interacts with both NFR1 and NFR5 at the plasma membrane and proteolytically cleaves NFR5 at the juxtamembrane (JM) domain to suppress NF-mediated host responses. NFR1 phosphorylates the full-length NopT, thereby inactivating its protease activity. However, the autocleaved NopT form of *S. fredii* NGR234 retains the ability to cleave NFR5 but loses its ability to interact with NFR1. Intriguingly, various *Sinorhizobium* strains possess only truncated versions of NopT (related to autocleaved NopT of NGR234)

which might evade the inactivation by NFR1. We present in this study a model of mutual

regulation, in which the rhizobial effector NopT directly targets NFR5 to dampen

symbiotic signaling, whereas NFR1-mediated phosphorylation of NopT counteracts

NFR5 cleavage. Our work provides essential insights into the intricate interplay between

legumes and rhizobia and shows an example of how a rhizobal effector fine-tunes NF

signaling by directly targeting NFRs, and how an NFR inactivates the enzymatic activity

of the effector through phosphorylation.

Results

NopT expression in *N. benthamiana* suppresses the NFR1/NFR5-induced cell death

response

To examine the role of effector proteins in regulating rhizobial infection, 15 effector

genes (Kimbrel et al., 2013) were cloned from *S. fredii* NGR234, a strain with an

exceptionally broad host-range. Initial experiments were performed to screen the effector

proteins for their ability to suppress or enhance the NFR1/NFR5–triggered cell death

response of *N. benthamiana* leaves. In this experiment, each effector was co-expressed

with NFR1/NFR5 (Fig. S1A–S1C). Only NopT, a cysteine protease belonging to the

YopT family (Shao et al., 2002; Dai et al., 2008), could suppress NFR1 and

NFR5-induced cell death (Fig. 1A and S1D–S1F). Interestingly, NopT reduced protein

abundance of the intact NFR5 when co–expressed with NFR1 (Fig. S1E), suggesting that

NopT might promote proteolytic degradation of NFR5. Cys-93 of NopT is an amino acid

residue known to be essential for the protease activity of this effector (Dai et al., 2008;

Kambara et al., 2009). In contrast to the wild-type NopT protein, expression of the

protease-dead NopT^{C93S} form in *N. benthamiana* did not result in suppression of cell

death induced by NFR1/NFR5 (Fig. 1B), indicating that the protease activity of NopT is

essential for the observed cell death-suppressing activity of NopT. In contrast to NopT,

NopM, an effector protein from *S. fredii* NGR234 reported to possess E3 ligase activity

(Xin et al., 2012), induced cell death when expressed in *N. benthamiana* leaves under our

test conditions (Fig. 1C).

We also tested whether the cell death–suppressing effect of NopT was specific for

NFR1/NFR5 expression or whether NopT could act generally to suppress cell death. In

addition to NopM, we examined various proteins known to induce cell death when

expressed in *N. benthamiana* leaves, namely Avr3a and R3a (Bos et al., 2006), apoptosis

regulator BAX (Bcl-2-associated X protein) (Gan et al., 2009), INF1, an elicitor from

Phytophthora infestans (Vleeshouwers et al., 2006) and Arabidopsis AtCERK1 (Cao et

al., 2014) (Fig. 1C). In contrast to the NFR1/NFR5–induced cell death, expression of

NopT was unable to suppress cell death when each of these proteins was expressed in *N.*

benthamiana leaves (Fig. 1D). Hence, the effect of NopT appears to be specific for the

NFRs. These results suggested that the effector protease NopT might act directly on

NFR1 and/or NFR5.

NopT interacts with NFR1 and NFR5

Autocleaved NopT of *S. fredii* NGR234 contains acylation sites required for lipidation

and subsequent plasma membrane localization (Dowen et al., 2009; Khan et al., 2022).

To investigate whether NopT interacts with NFR1 and NFR5, we first used *in vivo*

bimolecular fluorescence complementation (BiFC) analysis. NopT, NFR1 and NFR5

were C-terminally tagged with nYFP and cYFP. Fluorescence signals representing direct

NopT-NFR1, NopT-NFR5 and NopT-NopT interactions were detected at the plasma

membrane of *N. benthamiana* (Fig. 2A and 2B). Interactions between NopT and NFR1

and NFR5 were further verified using a Split-luciferase (Split–LUC) complementation

assay (Fig. 2C). Co-expression of NopT-CLuc and NFR1-NLuc or NFR5-NLuc produced

strong luminescence signals in transformed leaf discs compared with the negative control

expressing AtFLS2-NLuc (Arabidopsis FLAGELLIN SENSING 2) and NopT- CLuc

(Fig. 2C).

It is known that acylation of NopT can alter its subcellular localization (Dowen et al.,

2009; Khan et al., 2022). Here, we performed Split-LUC assays to examine the

interaction between NFR1/NFR5 and NopT$^{G50A/C51A/C52A}$, a modified NopT form lacking

acylation sites. Compared to the wild-type form (NopT^{WT}), the interactions between

NopT$^{G50A/C51A/C52A}$ and NFR1/NFR5 were significantly reduced in Split-LUC assays (Fig.

S3A and S3B). However, the protease-dead version of NopT (NopT^{C93S}) showed

interactions with NFR1 and NFR5, which were different from NopT^{WT} (Fig. S3C). These

results were further confirmed by a co-immunoprecipitation (co-IP) experiment in *N.*

benthamiana leaves co-expressing FLAG-tagged NopT with C-terminally HA-tagged

NFR1 or NFR5. Both NFR1 and NFR5 could be co-precipitated by an anti-FLAG

antibody when the sample contained NopT-FLAG (Fig. 2D and S2). Overall, the

identified physical interactions between NopT and NFR1/NFR5 indicated that NopT

targets NFRs and that NFR5 and/or NFR1 may be proteolytically cleaved by NopT.

NopT proteolytically cleaves NFR5 at the JM domain

A previous study has shown that NopT is autocleaved at its N-terminus to form a

processed protein that lacks the first 49 amino acid residues (Dai et al., 2008). To test

whether NopT could proteolyze NFR1 and/or NFR5, NopT was co-expressed with

NFR1-GFP and NFR5-GFP fusion proteins in *N. benthamiana* leaves. The expressed

proteins were separated by SDS-PAGE and subjected to immunoblotting. Proteolytic

cleavage of NFR5-GFP but not NFR1-GFP was observed in the presence of co-expressed

NopT, whereas protease-dead NopT^{C93S} showed no effect (Fig. 3A and S4A). A similar

in vivo cleavage assay with transgenic *L. japonicus* roots also showed that expression of

NopT but not NopT^{C93S} caused proteolytic cleavage of NFR5 (Fig. 3B). Based on the

molecular weight changes, the cleavage site in NFR5 was predicted to occur within the

cytoplasmic domain (CD) of NFR5. To further explore NFR5 cleavage by NopT, the CD

of NFR5 (Strep-NFR5^{CD}-HA) and NopT-FLAG or protease-dead NopT^{C93S}-FLAG were

co-expressed in the *Escherichia coli* cells. Immunoblotting results showed that the active

NopT protease cleaved NFR5^{CD} resulting in a ~5 kDa smaller protein (Fig. 3C and 3D).

However, when the CD of NFR1 was co-expressed with NopT in the same experiment,

no NFR1 cleavage was detected (Fig. 3C). Migration of the bands representing

full-length NopT or NopT^{C93S} was significantly retarded on the gel when the CD of

NFR1 was co-expressed (Fig. 3C), suggesting that NopT might be phosphorylated by

NFR1 in *E. coli* cells.

The NFR5^{CD} cleavage product with a ~5 kDa lower molecular weight in the cleavage

assay suggested that the cleavage site is located within the juxtamembrane (JM) domain

of NFR5. To test this hypothesis, we replaced the JM with the SUMO tag to creat

SUMO-NFR5^{KC}-HA (KC, kinase domain and C-terminal tail region, a modified NFR5^{CD}

without the JM) for proteolytic assay. Indeed, immunoblot analysis showed that

co-expressed NopT was unable to cleave SUMO-NFR5^{KC}-HA (Fig. S4B). The JM

domain of NFR5 was then fused to SUMO and GFP to create a recombinant protein,

SUMO-NFR5^{JM}-GFP. In an *in vitro* cleavage assay with recombinant proteins from *E.*

coli cells, GFP was immunodetected in the flow though sample when

SUMO-NFR5^{JM}-GFP was incubated with NopT but not with NopT^{C93S} (Fig. 3E). The

interaction between NopT^{C93S} and SUMO-NFR5^{JM}-GFP was confirmed by an *in vitro*

pull-down assay (Fig. S5). Overall, these data strongly indicate that the cleavage site of

NFR5 for NopT is located in its JM domain.

The cleavage of NFR5 by NopT is dependent on multiple residues of the JM domain

The "DKLLSGV" motif in the JM domain of NFR5 (residues Asp-288 to Val-294; Fig.

S6) shows highest similarity with the autocleavage region of NopT (DKMGCCA).

However, a protein variant of NFR5^{CD} in which the "DKLLSGV" motif was replaced by

7 alanine residues could still be cleaved by NopT (Fig. S7A). Similar to NopT, the

AvrPphB effector of *P. syringa* has an autocleavage site (Shao et al., 2003b). We

therefore also examined a version of NFR5^{CD} in which the "DKLLSGV" motif was

replaced by that of AvrPphB. In the cleavage assay with *E. coli* cells, AvrPphB did not

cleave NFR5^{CD}. However, the modified NFR5^{CD} form with the 7 residues from AvrPphB

was cleaved by AvrPphB and the cleavage product had a lower molecular weight than

that formed by NopT (Fig. S7A). These findings indicated that the cleavage site for NopT

is located upstream of the "DKLLSGV" motif in NFR5.

As mentioned above, NopT undergoes autocleavage but it is not clear whether this is due

to intermolecular or intramolecular proteolysis. In addition to Cys-93, His-205 and

Aps-220 of NopT are conserved catalytic residues (Dai et al., 2008; Kambara et al. 2009).

Therefore, NopT forms lacking these catalytic residues were co-expressed with NopT in

E. coli cells. The immunoblot analysis showed that NopT could not proteolyze NopT^{C93S},

NopT^{H205A} or NopT^{D220A} (Fig. S7B), suggesting that NopT autocleavage is due to

intramolecular proteolysis. Hence, the residues at the autocleavage site of NopT may not

help to predict the cleavage site of NopT substrates.

In an attempt to define the NopT cleavage site in NFR5, we created a series of modified

NFR5 forms covering the JM domain of NFR5 (Val-269 to Cys-320) and the first seven

amino acid residues of the kinase domain (Lys-321 to Tyr-327). Initially, 17 variants of

NFR5^{CD} were created in which three adjacent amino acids in the JM domain were replaced with three alanine residues. When these proteins were co-expressed with NopT in *E. coli* cells, all 17 NFR5^{CD} variants could be cleaved by NopT but not NopT^{C93S} (Fig.S7C-S7F). In a similar experiment, we created seven NFR5^{CD} forms which contained a deletion in the JM domain of 10 residues each. Surprisingly, cleavage products were still observed for all seven examined proteins (Fig. S7G). These results suggest that the JM domain may have multiple sites that can be cleaved by NopT, which is different from the specific cleavage site identified for AvrPphB. In another experiment, recombinant His-SUMO-NFR5^{CD}-GST protein was co-expressed with NopT in *E. coli* cells and its cleavage product was subjected to N-terminal protein sequencing using Liquid Chromatography Mass Sprectrometry (LC-MS) analysis. Based on alignment of identified peptides, the NopT cleavage site in NFR5 was mapped to four basic amino acids in the JM domain (RRKK; amino acid residues 271-275) (Fig. S8). However, this result was not consistent with the observation that NopT expressed in *E. coli* cells could still proteolyze a co-expressed NFR5^{CD} form in which the residues from Tyr-268 to Leu-277 were deleted (Fig. S7G). Overall, the experiments performed suggest that NopT preferentially proteolyzes NFR5 at the RRKK motif, but other sites in the JM domain can also be cleaved by NopT. Since NFR5 belongs to a subfamily of LYK family proteins that lacks kinase activity (Yang et al., 2022), we wondered whether NopT could proteolyze other LYK proteins of this subfamily, which might be helpful to characterize the biochemical function of NopT. We therefore investigated whether the CDs from *Arabidopsis* AtLYK5, *L. japonicus* LjLYS11 and *M. truncatula* MtNFP expressed in *E. coli* cells can be proteolyzed by NopT. As shown in Fig. S9, NopT, but not NopT^{C93S}, was able to cleave the CDs of AtLYK5 and LjLYS11. Replacement of the JM domain in NFR5^{CD} with the JM domain from either AtLYK5 or LjLYS11 resulted in cleavage by NopT (Fig. S9). These data show that NopT cleaves AtLYK5 and LjLYS11 at their JM domains. Based on sequence analysis of the JM region of NFR5 and its homologous

proteins, five conserved residues (Ser-283, Gly-294, Tyr-304, Ala-310, Thr-311) were

identified and mutated as NFR5^{CD-5m} for the proteolytic assay (Fig. S6B). As shown in

Fig. 3F, NFR5^{CD-5m}, a NFR5^{CD} variant with point mutations of S283Y, G294Q, Y304S,

A310I, and T311Y, could not be cleaved by NopT. Interestingly, NopT could not

proteolyze MtNFP^{CD} (Fig. 3G). We then swapped the JM domains in NFR5^{CD} and

MtNFP^{CD}, generating NFP^{JM}-NFR5^{KC} and NFR5^{JM}-NFP^{KC} (KC stands for the kinase

domain and the C-terminal tail region). As shown in Fig. 3G, NopT could proteolyze

NFP^{JM}-NFR5^{KC} but not NFR5^{JM}-NFP^{KC} in *E. coli* cells. In an *in vitro* cleavage assay, the

cleavage product of SUMO-NFP^{JM}-GFP proteolyzed by NopT was detected in the flow

though sample (Fig. S10). These data suggest that the cleavage of NFR5 by NopT

depends on the conformational structure of NFR5^{KC} and/or other specific amino acids in

NFR5^{KC}. We then investigated whether two additional recombinant proteins with large

swapping   regions   in   NFR5^{CD} and   NFP^{CD} (NFR5$^{268-445}$-NFP$^{458-595}$ and

NFP$^{270-457}$-NFR5$^{456-595}$) could be cleaved in the *E. coli* proteolysis test (Fig. S11).

Notably, both proteins were cleaved by NopT, suggesting that a specific conformation of

the NopT substrate is required for proteolysis. Taken together, we concluded from our

experiments that NopT can proteolyze NFR5 at the JM domain and that mutiple

nonadjacent residues in NFR5^{JM} in combination with a specific KC conformation are

required for proteolysis.

NopT phosphorylated by NFR1 is proteolytically inactive

The perception of rhizobial NFs by the heterocomplex of NFR1 and NFR5 involves

trans-phosphorylation events from the NFR1 kinase to the pseudo-kinase NFR5 (Madsen

et al., 2011). Since NopT is associated with both NFR1 and NFR5, we hypothesized that

NopT   is   a   phosphorylation   target   of   NFR1,   which   may   interfere   with

transphosphorylation events between the two receptors. NopT^{C93S} was therefore

expressed in *L. japonicus* wild-type and *NFR1* knockout mutant plants (*nfr1-1*) and

analyzed on Zn^{2+}-Phos-tag gels. On such gels, a Zn^{2+}-Phos-tag bound phosphorylated

protein migrates slower than its unbound nonphosphorylated form. Slower migrating

bands corresponding to phosphorylated NopT^{C93S} were observed, particularly when

wild-type plants were inoculated with rhizobia (Fig. 4A and 4B). These findings

indicated that NopT phosphorylation *in planta* is largely dependent on NFR1 and that the

protein was probably phosphorylated at multiple sites.

Next, we performed assays to investigate whether expressed NFR1^{CD} directly

phosphorylates co-expressed NopT. NFR5^{CD} and a kinase-inactive NFR1^{CD} form

(NFR1$^{CD-K351E}$; Lys-351 is a conserved residue required for ATP binding) were included

into these experiments. When NFR1^{CD} was expressed in *E. coli* cells, extracted NopT

(Fig. 3C and 4C) and NFR5^{CD} (Fig. S12) exhibited retarded migration on SDS-PAGE

gels, suggesting that NFR1 was able to phosphorylate both proteins. A band shift was

observed when the phosphorylated full-length NopT was *in vitro* dephosphorylated by

calf intestinal alkaline phosphatase (CIAP). However, the CIAP treatment caused no

obvious band shift of the autocleaved NopT form (Fig. 4C). The phosphorylation sites of

NopT were then identified using an *in vitro* phosphorylation assay followed by LC-MS

analysis. Several serine and threonine residues in NopT were identified to be

phosphorylated by NFR1^{CD} (Table S1). Taken together, these results showed that

full-length NopT is a phosphorylation target of the NFR1 kinase.

Finally, we wondered whether NopT phosphorylation by NFR1 influences its proteolytic

activity. Plasmids encoding NopT variants were constructed in which the phosphorylated

residues were substituted to either alanine (to block phosphorylation) or aspartate (to

mimic the phosphorylation status). The different NopT forms were then expressed in *E.*

coli cells to analyze NopT autocleavage and proteolysis of co-expressed NFR5^{CD}.

Aspartate substitutions at three phosphorylation sites of NopT (Thr-97, Ser-119 and

Thr-257) resulted in the loss of the autoproteolytic activity, as well as the ability to cleave

NFR5^{CD} (Fig. 4D). In contrast, corresponding NopT proteins with alanine substitutions

retained autocleavage activity and the ability to proteolyze NFR5^{CD} (Fig. 4D). These data

indicate that phosphorylation of full-length NopT by NFR1 inhibits its proteolytic

activity, thereby keeping NopT unprocessed and NFR5 uncleaved.

NopT dampens rhizobial infection

As NopT of *S. fredii* NGR234 could directly target and cleave NFR5, we expected that

NopT may be an important player regulating rhizobial infection in legumes. NF signaling

is known to function during early stages of rhizobial infection (Geurts et al., 2005; Wang

et al., 2012; Cai et al., 2018). We focused our experiments on early stages of rhizobial

infection. Inoculation of *L. japonicus* plants with a GFP- or GUS-labelled *nopT* mutant of

NGR234 (NGR234 *ΔnopT*) resulted in a massive infection of root hairs, whereas

infection by the NGR234 wild-type strain was less frequent (Fig. 5A and 5C). We also

performed experiments with *L. japonicus* plants containing *pNIN*:*GUS* which exhibit

β-glucuronidase (GUS) activity when the *NIN* (*NODULE INCEPTION*) promoter

activity is induced by NF signaling (Schauser et al., 1999). GUS staining of roots showed

that the *NIN* promoter activity was stronger in response to NGR234*ΔnopT* inoculation

relative to inoculation by wild-type NGR234 (Fig. 5B and 5D). Likewise, inoculation

with NGR234*ΔnopT* resulted in an increased number of nodule primordia (Fig. 5G). In

contrast, over-expression of NopT under the control of the T7 promoter in the NGR234

parent strain reduced the number of nodule primordia when compared to inoculation with

the wild-type strain (Fig. 5G). These findings indicate that NopT negatively influences

rhizobial infection and nodule initiation. Interestingly, the internal structures of nodules

stained by toluidine blue dye exhibited remarkable similar when inoculated with both

NGR234 and NGR234$\Delta nopT$ (Fig. S13).

To confirm that the massive infection of roots by NGR234$\Delta nopT$ is due to the loss of

NopT, we complemented the NGR234$\Delta nopT$ by expressing wild-type NopT. The *L.*

japonicus plants inoculated with this strain had fewer infection foci, indicating restoration

of the wild-type phenotype (Fig. 5E). We also tested whether the proteolytic activity of

NopT affected rhizobial infection. The protease-dead version of NopT^{C93S} and three

NopT forms with phospho-mimetic residues (i.e., NopT^{T97D}, NopT^{S119D}, and NopT^{T257D}

lacking proteolytic activity) were individually expressed in NGR234$\Delta nopT$. Compared

with the high number of infection foci induced by NGR234$\Delta nopT$ in *L. japonicus* plants,

inoculation with these strains resulted in significantly fewer infection foci, but more than

with the wild-type strain (Fig. 5E).

We also expressed *nopT* under the control of a ubiquitin promoter in hairy roots of *L.*

japonicus and inoculated the transgenic roots with DsRed-labeled *M. loti* MAFF303099,

which does not possess a *nopT* gene in its genome. As shown in Fig. 5F, reduced

numbers of infection foci and infection threads were observed in the NopT expressing

roots, indicating that NopT negatively affects the symbiosis between *L. japonicus* and *M.*

loti strain.

To investigate whether the cleavage of NFR5 by NopT reduces rhizobial infection of *L.*

japonicus roots, *NFR5* and *NFR5^{5m}* (uncleavable variant with 5 amino acid substitutions

in the JM domain; Fig. 3F) under the control of the native *NFR5* promoter, were

expressed in hairy roots of *NFR5* knockout mutant plants (*nfr5-3*). NFR5 expression

resulted in numerous infections and a significant increase of infection foci was observed

upon inoculation with the NGR234$\Delta nopT$ mutant (Fig. 5H). However, NFR5^{5m}

expression showed no effects on rhizobial infection (Fig. S14), suggesting that the amino

acid substitutions in NFR5^{5m} caused a conformational change that rendered the protein

inactive in symbiotic signaling. As the NFR5^{JM}-NFP^{KC} fusion protein was not cleaved by

NopT (Fig. 3G), we also constructed hairy roots of *nfr5-3* mutant plants in which

NFR5-NFP^{KC} (NFR5 variant with its KC domain replaced by the KC of MtNFP) was

expressed under the control of the *NFR5* promoter. Remarkably, inoculation of these

plants showed no difference between NGR234*ΔnopT* and the NGR234 wild-type strain in

terms of rhizobial infection foci (Fig. 5H), suggesting that the cleavage of NFR5 by

NopT reduced the degree of rhizobial infection. Taken together, these data indicate that

the protease activity of NopT, its phosphorylation status and the proteolysis of the NFR5

target are important for the regulation of rhizobial infection.

NopT from other *S. fredii* strains also cleave NFR5

Since NopT of *S. fredii* NGR234 (NopT$_{NGR234}$) cleaves NFR5, we wondered whether

homologs from other rhizobial species also possess such a proteolytic activity.

Phylogenetic analysis showed that NopT proteins of *Sinorhizobium* and *Bradyrhizobium*

are located in different clades (Fig. S15). *B. diazoefficiens* USDA110, a typical

Bradyrhizobium strain used to study nodulation of soybeans, produces two NopT proteins

(Fotiadis et al., 2012). In contrast to NopT$_{NGR234}$, however, neither NopT1$_{USDA110}$ nor

NopT2$_{USDA110}$ were able to cleave co-expressed NFR5 in *E. coli* cells (Fig. 6A). This

finding suggests that NopT homologs from *Bradyrhizbium* species have lost the ability to

cleave NFR5 and probably act on other, unknown host target proteins. The genome of

various *Sinorhizobium* species (e.g., *S. fredii* USDA257 and S. *fredii* HH103) possess

genes that encode truncated NopT forms, which are almost identical to the autocleaved

version of NopT$_{NGR234}$ (Khan et al., 2022). Remarkably, NopT$_{USDA257}$ and NopT$_{HH103}$

were both able to cleave NFR5 *in E. coli* cells (Fig. 6A). However, NopT$_{USDA257}$ were

unable to suppress cell death triggered by NFR1 and NFR5 expression in *N. benthamiana*

leaves (Fig. 6B). Moreover, expression of NopT$_{USDA257}$ in NGR234*ΔnopT* had no effect

on the infection ability of NGR234*ΔnopT,* as observed in inoculation tests with the *L.*

japonicus line containing *pNIN*:*GUS* (Fig. 6C and 6D). These results suggest that

NopT$_{USDA257}$ and NopT$_{HH103}$ exhibit similar proteolytic activity to NopT$_{NGR234}$, but target

different host proteins to regulate rhizobial infection.

Discussion

Plant-microbe interactions are complex, employing distinct "chemical weapons" or

strategies to communicate mutually, thereby securing more resources for their

survival (Jones and Dangl, 2006; Schubert et al., 2020). T3SS effectors of

phytopathogenic bacteria are injected into plant cells to facilitate bacterial

infection by targeting host proteins of the plant immune system, thereby

suppressing PTI (Jones and Dangl, 2006; Tang et al., 2017). However, the role of

rhizobial T3SS effectors in regulating the mutualistic symbiosis between legumes

and rhizobia remains largely unknown. The aim of this study was to gain essential

insights into the regulation mechanisms by which rhizobial effectors act directly on

symbiotic signaling pathways to promote or dampen infection of host cells.

Understanding how plants recognize different microbes as "friends" or "foes" is a

central question in biology. The perception of microbial signals by plant receptor

proteins is essential for triggering specific signaling pathways to resist invading

pathogens or to establish symbiosis with beneficial microbes (Zipfel and Oldroyd,

2017). Direct targeting of these receptors by T3SS effectors represents an

efficient strategy of phytopathogenic bacteria to suppress PTI and promote

bacterial infection. In the well-studied *Arabidopsis-Pseudomonas syringae*

interaction, T3SS effectors target plant flagellin receptor FLS2 and its coreceptor

BAK1 (BRI1-[Brassinosteroid insensitive1]-associated receptor kinase 1) to

suppress flagellin-induced PTI (Zipfel et al., 2004; Sun et al., 2013). For example,

AvrPto targets FLS2 and HopB1 cleaves BAK1 (Xiang et al., 2008; Li et al., 2016).

Likewise, AvrPtoB, an E3 ligase in *P. syringae*, targets FLS2 to promote its

degradation and suppress PTI (Goehre et al., 2008). Despite the evolutionary

importance of T3SS effectors in targeting host receptors, we were surprised to

find that NopT, a rhizobial member of the C58 protease effector family, can

directly target NFRs and manipulate the symbiotic signaling pathway in legumes.

The interaction between NopT of *S. fredii* NGR234 and NFRs of *L. japonicus* was

identified by using a unique screening system, in which co-expression of

NFR1/NFR5 in *N. benthamiana* leaves leads to cell death. NopT, but not other

effectors of NGR234, could suppress this cell death response. Consistent with

these initial observations, inoculation of *L. japonicus* with the *S. fredii*

NGR234Δ*nopT* mutant resulted in increased infection of *L. japonicus* roots.

Moreover, over-expression of NopT in the roots was found to reduce rhizobial

infection by *M. loti*. The interaction between NopT and NFR1/NFR5 was verified

by BiFC, Split-LUC, and co-IP experiments.

Interestingly, using different assays, we found that NopT cleaves NFR5 at the JM

domain. However, the efficiencies of NopT cleavage of NFR5 in *L. japonicus* and

N. benthamiana were slightly different (Fig. S16), which could be related to the

uncontrollable activation of NFRs in *N. benthamiana* or phosphorylation of NopT

by another kinase leading to decreased proteolytic activity. The modification of

five conserved residues in the JM domain and the replacement of NFR5^{KC} by

NFP^{KC} resulted in NFR5^{CD} forms that were resistant to proteolysis by NopT in *E.*

coli cells. These results suggest that cleavage of NFR5 depends on both the JM

domain and the KC region. Although belonging to the same protease family, the

NopT cleavage site in NFR5 was found to be polybasic as in the YopT substrate

(Shao et al., 2003a; Schmidt, 2011), whereas the cleavage site of AvrPphB

substrate is canonical with seven adjacent amino acids involved (Kim et al.,

2016). These differences reflect a broad variety of YopT-type effector proteases

and their substrates in host cells. Our study shows that NopT targets the

NFR1/NFR5 complex thus reduces NF signaling and formation of nodule

primordia in *L. japonicus*. NopT-NFR interactions may reflect a regulation

mechanism to prevent rhizobial hyperinfection in host species other than *L.

japonicus*.

Similar to the *Pseudomonas* effector AvrPphB, NopT of *S. fredii* NGR234 is able

to cleave soybean PBS1-1 and expression of this effector in *S. fredii* USDA257

negatively influences symbiosis with certain soybean cultivars (Khan et al., 2022).

Indeed, NopT of NGR234 acts as an Avr effector in some nonhost plants.

Transient expression of NopT of strain NGR234 in *Arabidopsis* and tobacco

triggers strong immune responses and cell death (Dai et al., 2008; Kimbrel et al.,

2013; Khan et al., 2022; Fig. S17). In *Arabidopsis*, NopT-induced cell death was

found to be dependent on the PBS1-1 kinase and the resistance protein RPS5

(Khan et al., 2022). Likewise, when constitutively expressing NopT and NopT^{C93S}

in *L. japonicus*, we only obtained stable transgenic lines expressing NopT^{C93S},

indicating that the protease activity of NopT had a negative effect on plant

development. In contrast, generation of hairy roots expressing NopT was

possible. Such differences may be explained by different NopT substrates in roots

and aerial parts of the plant. Indeed, our study shows that NopT not only cleaves

NFR5 but is also able to proteolyze *Arabidopsis* AtLYK5 and *L. japonicus*

LjLYS11. These two receptors possess chitin-binding affinities and trigger PTI

responses (Cao et al., 2014; Gysel et al., 2021). NFR5, AtLYK5, and LjLYS11 are

kinase-dead LYKs belonging to the same LYK subfamily. Thus, NopT appears not

only to suppress NF signaling, but may also interfere with signal transduction

pathways related to plant immunity. Depending on the host legume species, NopT

could suppress PTI or induce ETI, thereby modulating rhizobial infection and

nodule formation. Interactions between NopT and proteins related to the plant

immune system may represent an important evolutionary driving force for

host-specific nodulation and explain why the presence of NopT in NGR234 has a

negative effect on symbiosis with *L. japonicus* but a positive one with other

legumes.

The regulation of kinase-dead pseudokinases in plants may be more complex

compared to the well-studied control of protein kinase activities by

phosphorylation. The cleavage of receptor-like protein kinase may be an

overlooked strategy to regulate their function in plant biology. Regulation of

protein activities by cleavage is well studied in animals, e.g., for Notch signaling,

where proteolysis of Notch molecules leads to downstream signaling (Kopan and

Ilagan, 2009). *Arabidopsis* BAK1 undergoes proteolytic cleavage, which is

important for both brassinosteroid signaling and induction of PTI responses (Zhou

et al., 2019). Whereas, HopB1 cleaves immune co-receptor BAK1 at the kinase

domain to inhibit plant defense (Li et al., 2016). NFR5 is a pseudokinase that is

essentially required for the activation of NF signaling. However, the mechanisms

of how the pseudokinase domain of NFR5 mediates symbiotic signaling

transduction remains unclear. Our work indicates that NopT proteolytically

cleaves NFR5 and suppresses NF signaling. Such a fine-tuning process could be

advantageous for the symbiosis in host plants other than *L. japonicus*, for

example to prevent hyperinfection. However, the precise function of NFR5

cleavage by NopT and the fate of the released cytoplasmic domain remain to be

clarified.

Mutual regulation between T3SS effectors and host target proteins has been

investigated for various plant-pathogen interactions. For example, the conserved

Pseudomonas effector AvrPtoB can directly ubiquitinate the key components of

plant immunity to promote bacterial virulence (Goehre et al., 2008). However, the

activity of AvrPtoB in promoting virulence could either be enhanced or dampened

by host proteins. AvrPtoB can be phosphorylated by SnRK2.8 to promote its

virulence activity (Lei et al., 2020). On the other hand, the plant lectin receptor-like

kinase LexRK-IX.2 and the Pto kinase phosphorylate AvrPtoB to inactivate its

ubiquitin E3 ligase activity and undermine the effector's ability to suppress PTI

(Ntoukakis et al., 2009; Xu et al., 2020). A similar strategy is used by NFR1 to

inhibit the protease activity of NopT *via* phosphorylation. NFR1 expressed in *E.*

coli cells phosphorylates full-length NopT of *S. fredii* NGR234 at several residues.

Three NopT forms with a phosphomimic substitution (i.e., NopT^{T97D}, NopT^{S119D},

and NopT^{T257D}) lacked autoproteolytic activity and the ability to cleave NFR5. The

expression of these NopT variants in the NGR234Δ*nopT* mutant showed little

effects in comparison with expression of wild-type NopT. These data suggest that

L. japonicus utilizes NFR1 to phosphorylate NopT in order to dampen its catalytic

activity and protect NFR5 from cleavage, which favors rhizobial infections.

However, NopT^{ΔN50}, which is similar to autocleaved NopT, retained the ability to

interact with NFR5 but not with NFR1 (Fig. S3D). Moreover, full-length NopT, but

not autocleaved NopT, migrated slower on gels when NFR1^{CD} was co-expressed

in *E. coli* cells, suggesting that autocleaved NopT may escape being

phosphorylated and inactivated by NFR1. Future studies are required to explore

the interactions between acylated NopT and NFR1. When expressed in *N.*

benthamiana, GFP-tagged NopT forms lacking their autocleavage site (and thus

non-acylated) were found to be localized within the cytoplasm, whereas NopT

was targeted to the plasma membrane (Khan et al., 2022). Accordingly, NopT

without acylation sites showed weaker interactions with NFRs in our study,

suggesting that acylation of NopT, leading to plasma membrane location,

promotes NopT-NFR interactions.

NopT homologs vary in different *S. fredii* strains. USDA257 and HH103 produce

truncated NopT versions different from NGR234 due to a 19-bp deletion (Khan et

al., 2022). Accordingly, NopT of these strains lacks autocleavage and acylation

sites and possess a different N-terminal secretion signal sequence (Akeda and

Galán, 2005). Nevertheless, NopT$_{HH103}$ is a functional T3SS effector as mutant

analysis indicated a symbiosis-promoting role in certain soybean cultivars (Li et

al., 2023). Similar to NopT$_{NGR234}$, NopT$_{USDA257}$ and NopT$_{HH103}$ were able to cleave

NFR5 in *E. coli* cells. Since NopT$_{USDA257}$ and NopT$_{HH103}$ cannot be acylated, they

likely accumulate in the cytoplasm of host legumes and are therefore less efficient

in cleaving plasma membrane-bound NFR5 in comparison to acylated

NopT$_{NGR234}$. This could also explain why NopT$_{USDA257}$ was unable to suppress cell

death triggered by NFR1/NFR5 in *N. benthamiana* leaves. It can be expected that

NFR1 does not interact with and phosphorylate NopT$_{USDA257}$ or NopT$_{HH103}$, as

these effectors are similar to NopT^{ΔN50} of NGR234, which cannot interact with

NFR1. Consistent with these findings, expression of NopT$_{USDA257}$ in

NGR234*ΔnopT* did not alter the infection phenotype of NGR234*ΔnopT*. It can be

hypothesized that the host targets of these truncated NopT effectors might be

different from those of NopT$_{NGR234}$. *Bradyrhizobium* NopT effectors are also

different from NopT$_{NGR234}$, as NopT of *B. diazoefficiens* USDA110 was unable to

cleave NFR5. NopT effectors may play a crucial role in *Bradyrhizobium*-legume

interactions. A *Bradyrhizobium* sp. ORS3257 mutant deficient in NopT production

induced only ineffective nodules on roots of *Aeschynomene indica* and most of

the formed nodules were not infected (Teulet et al., 2019). Overall, these findings

indicate versatile functions of NopT homologs in different strains and suggest that

NopT controls different regulation processes in specific host cells.

We present here a model of mutual regulation, in which NopT proteolyzes NFR5

to suppress NF signaling, whereas protease activity of full-length NopT is

suppressed by NFR1 *via* phosphorylation (Fig. 6E). NopT impairs the function of

the NFR1/NFR5 receptor complex. Cleavage of NFR5 by NopT reduces its

protein levels. Possible inhibitory effects of NFR5 cleavage products on NF

signaling are unknown but cannot be excluded. Inactivation of NopT protease

activity *via* phosphorylation by kinases such as NFR1 appears to be a

countermeasure of the host to enable symbiotic signaling and rhizobial infection.

Such feedback regulation could be the result of evolutionary pressure to produce

'improved' NFRs, which are resistant to cleavage by NopT and have an increased

capacity to phosphorylate NopT. On the other hand, suppression of NopT

protease activity by NFRs may drive broad-host-range strains such as NGR234 to

produce higher levels and structurally modified NFs.

The negative regulation of plant symbiotic signaling by NopT is detrimental to

rhizobial infection, which appears paradoxical from an evolutionary perspective.

Two possible scenarios will be discussed. One possibility is that NopT may have

originated from its homologs, such as AvrPphB, and possesses a pathogenic

feature by proteolyzing immune receptors to disrupt the plant immune pathway.

However, due to sequence similarity, NFR5 is also proteolyzed by NopT.

Expression of NopT leading to strong cell death in *N. tabacum* and *Arabidopsis*

also suggests that NopT might be recogonized as Avr effector to trigger strong

immunity. Another possibility is that the presence of NopT serves as a strategy for

rhizobia to evade detection by plant immunity, which may be activated by rhizobial

overinfection. The negative effects on symbiosis by cleaving NFR5 is then

inhibited by NFR1 via a direct phosphorylation. In a parallel study, *S. fredii*

NGR234 NopM was also identified to interact with and mediate the ubiquitination

of NFR5, stabilizing NFR5 levels and thereby promoting rhizobial infection and

nodulation (Wang et al., 2024). Therefore, both *S. fredii* NGR234 NopT and NopM

target the same protein and appear to antagonistically regulate rhizobial

symbiosis; however, the exact role of NopT in nodulation needs to be elucidated

in future research.

Taken together, this study provides insights into a legume-rhizobium interaction,

in which the bacterium deploys an effector protease to dampen symbiotic

signaling, while the host plant counteracts by phosphorylating the effector, leading

to its inactivation. Our findings highlight the function of a bacterial effector protease in

regulating a symbiotic signaling pathway in legumes. This opens up the perspective of

developing specific kinase-interacting proteases to reprogram and fine-tune cellular

signaling processes in general.

Materials and Methods

Germination and growth of *Lotus japonicus*

Lotus japonicus (ecotype Gifu B-129) was provided by the Center for Carbohydrate

Recognition and Signaling (https://lotus.au.dk/), and used as wild-type for nodulation

assays. The Gifu *pNIN*:*GUS* transgenic line and *nfr* mutants (*nfr1-1* and *nfr5-3*) were

kindly provided by Dr. Jens Stougaard from the Aarhus University, Denmark (Madsen et

al., 2003; Radutoiu et al. 2003; Heckmann et al., 2011). All *L. japonicus* seeds were

treated with concentrated sulfuric acid for 10 min, followed by surface sterilization in 1%

(w/v) NaClO for 8 min. After incubation at 4°C in the dark for 2 d, the seeds were placed

on 0.8% (w/v) agar containing half-strength Murashige & Skoog (MS) medium for

germination at 22°C for 2 d in the dark. Seedlings were then transferred to growth pots

containing vermiculite supplied with half-strength B&D (Broughton & Dilworth)

medium without nitrogen under long-day conditions (16-h light/8-h dark) at 22°C.

Ten-day-old seedlings were inoculated with rhizobial cultures (OD_{600}=0.02) for infection

experiments.

Plasmid construction

All plasmids used in this study were generated using MultiF Seamless Assembly Mix

(Abclonal, # RK21020). The plasmid pGWB514 was used as an original plasmid for

construction of all binary vectors (Nakagawa et al., 2007). Briefly, different tags

including HA tag, Myc tag, FLAG tag, Strep tag, NLuc, CLuc and GFP, were amplified

using a forward primer containing a *Kpn*I site and ligated into pGWB514 digested with

*Xba*I and *Sac*I using the seamless cloning method described above. The generated

plasmids were named pG5XX-HA, pG5XX-Myc, pG5XX-FLAG, pG5XX-NLuc,

pG5XX-CLuc, pG5XX-GFP, respectively. The coding sequences of *L. japonicus NFR1*

and *NFR5* were cloned into pGWB514 and pG517 between *Xba*I/*Kpn*I to generate

NFR1-HA and NFR5-Myc fusion constructs under the control of the cauliflower mosaic

virus 35S promoter. The DNA fragment containing *pro35S*:NFR5-Myc-NosT was then

amplified and cloned into pG514-NFR1 at *Sbf*I. The final plasmids contained

pro35S:NFR1-HA-NosT and *pro35S*:NFR5-Myc-NosT. Rhizobial genes encoding

effectors from *S. fredii* NGR234 were cloned into pG5XX-FLAG between *Xba*I/*Kpn*I. To

detect the interaction between NFR1/NFR5 and NopT in *N. benthamiana*, the full-length

NFR1/NFR5, *NopT* and *AtFLS2* coding sequences were cloned into pG5XX-NLuc,

pG5XX-CLuc, pSPYCE (MR) and pSPYNE (R)173 (Waadt et al., 2008). For cleavage

assays in *N. benthamiana* leaves, the *NFR5* and *NopT* coding sequences were cloned into

pG5XX-FLAG, pG5XX-GFP and pG5XX-Strep, respectively. For cleavage assays in *E.*

coli cells, the sequence encoding the cytoplasmic domain of NFR5 was cloned into

pACYC duet Strep-HA (Han et al., 2017), and the sequences encoding the kinase domain

of NFR5 fused to the HA tag and the JM domain of NFR5 fused to the GFP tag were

cloned into pSUMO. The point mutations in *NFR5^{CD}* and *NopT* were created by

site-directed mutagenesis PCR using the templates pACYC duet Strep-NFR5^{CD}-HA and

pET28a NopT-FLAG. For NFR/NopT cleavage experiments with *L. japonicus* plants,

NopT-FLAG or NopT^{C93S}-FLAG was cloned into pUB-GFP between XbaI and KpnI,

and 35S-NFR5-Myc-NosT was cloned into pUB-NopT-GFP at PstI. For NopT

phosphorylation experiments with *E. coli*, the sequence encoding the cytoplasmic domain

of NFR1 fused to a Myc tag was cloned into pCDF duet. For complementation

experiments with the NGR234*ΔnopT* mutant (Dai et al., 2008), a 1317-bp DNA fragment

upstream of *NopT* and the coding sequence of a given *NopT* sequence was fused using

overlap PCR and cloned into pHC60 between XhoI and KpnI. For complementation

experiments with the *nfr5-3* mutant, a 1316-bp *NFR5* promoter DNA fragment was fused

to NFR5-NFP^{KC} or NFR5^{5m} using overlap PCR and cloned into pUB-Cherry between

PstI and KpnI.

Transient gene expression in *Nicotiana* leaves

The plasmids of *INF1*, *Avr3*, *BAX*, and *R3a* for transient expression in *Nicotiana*

benthamiana were kindly gifted by Dr. Juan Du from Huazhong Agricultural University.

Agrobacterium tumefaciens strain EHA105 carrying various constructs were mixed with

an Agrobacterium culture harboring the P19 suppressor in an infiltration buffer (10 mM

MgCl$_2$, 10 mM MES-KOH pH=5.8 and 200 μM acetosyringone). After incubation for 2 h

at room temperature, *Agrobacterium* cultures were infiltrated into the leaves of

4-week-old *N. benthamiana* and *N. tabacum* plants. Leaves were harvested and analyzed

two-days post inoculation. The cell death suppression experiments with *N. benthamiana*

plants were performed as described previously (Wang et al., 2011).

Hairy root transformation

Hairy root transformation of *L. japonicus* plants was performed as described previously

(Li et al., 2018). In brief, surface-sterilized seeds were germinated and grown on MS

plates without sucrose (23°C in the dark for the first 3 days and 23°C at a 16-h light/8-h

dark period for the following 2 days). The seeds were then cut at the middle of the

hypocotyl and co-cultivated with *A. rhizogenes* LBA1334 carrying pUB-GFP,

pUB-NopT-FLAG-GFP,   pNFR5:NFR5-mCherry,   pNFR5:NFR5^{5m}-mCherry   and

pNFR5:NFR5-NFP^{KC}-mCherry, respectively (23°C in the dark for the first 3 days and

23°C at a 16-h light/8-h dark period for the following 2 days). The plants were then

transferred onto solid B5 medium. After about 15 days, non-fluorescent hairy roots

(lacking expression of GFP or mCherry) were removed. The plants were inoculated with

rhizobia: *S. fredii* NGR234, NGR234Δ*nopT* or *Mesorhizobium loti* MAFF303099, and

analyzed at indicated time points.

Analysis of NFR5 cleavage by NopT and variants

In experiments with *E. coli*, Strep-NFR5^{CD}-HA or Strep-NFR5^{CD}-HA variants were

co-expressed with NopT-FLAG or FLAG tagged NopT variants or AvrPphB-FLAG by

using Duet vectors (Novagen; Han et al., 2017). Bacterial cultures were treated with 0.5

mM IPTG at 22°C for 15 h. After washing the bacterial cells with PBS buffer, the

bacterial pellet was resuspended in extraction buffer consisting of Tris-HCl (pH=7.3),

150 mM NaCl, 5 mM EDTA, 0.5% (w/v) SDS and 0.5% (v/v) Triton X-100. After

adding SDS loading buffer and boiling at 100°C for 5 min, the extracted proteins were

subjected to immunoblot analysis using anti-HA-peroxidase (Sigma, clone 3F10) or

anti-FLAG (Sigma, F1804) antibodies. An anti-groEL antibody (ABclonal, A0969) was

used to confirm equal loading of proteins on gels.

For *in vitro* cleavage assays, His-SUMO-NFR5^{JM}-GFP and NopT-FLAG-His proteins

were expressed in *E. coli* cells and purified. The reaction mixture contained 3 μg

NopT-FLAG (or NopT^{C93S}-FLAG) and 1 μg SUMO-NFR5^{JM}-GFP in extraction buffer

without SDS and EDTA. After incubation at 22°C for 7 h, 10 μL of a Ni-charged resin

(GenScript, Cat. No. L00223) were added, and the cleavage products in the flow through

were detected by immunoblot analysis using an anti-GFP antibody (ABclonal,

AB_2770402).

To analyze cleavage of NFR5 in *L. japonicus* roots, hairy roots co-expressing

NopT-FLAG or NopT^{C93S}-FLAG with NFR5-Myc were harvested at about 15 days after

induction of hairy roots without rhizobia inoculation and subjected to immunoblot

analysis. Expressed proteins were extract by extraction buffer and immunodetected with

anti-FLAG antibody and anti-Myc (Bio-Legend, # 626808) antibody.

For NFR5/NopT cleavage experiments with *N. benthamiana* plants, leaves expressing

proteins (NopT-FLAG or NopT^{C93S}-FLAG with NFR5-GFP) were analyzed. Leaf discs

were collected at 2 days post infiltration with *agrobacteria* and extracted in the extraction

buffer (Tris-HCl pH=7.3, 150 mM NaCl, 5 mM EDTA, 0.5% (w/v) SDS and 0.5% (v/v)

Triton X-100) and 2% protease inhibitors (Sigma, P9599). NFR5 was immunodetected

with anti-Myc and anti-GFP (ABclonal, AB_2770402) antibodies. The NopT or

NopT^{C93S} were detected with anti-FLAG antibody. The actin proteins were detected with

anti-Actin antibody (Abclonal, # AC009).

Phosphorylation assays

For NopT phosphorylation in *E. coli* cells, His-NFR1^{CD}-Myc or His-NFR1$^{CD–K351E}$-Myc

was co-expressed with NopT-FLAG-His by using Duet vectors (Novagen). Protein

expression was induced by 0.5 mM IPTG and cells were grown at 28°C for 18 h. After

simultaneous purification of NFR1^{CD} and NopT using Ni-charged resin, the proteins were

incubated with CIAP (TaKaRa, 2250B) at 37°C for 3 h. The mobility shifts of NFR1^{CD}

and NopT were detected with anti-Myc (Bio-Legend, 626808) and anti-FLAG (Sigma,

F1804) antibodies by immunoblot analysis.

In phosphorylation experiments with *L. japonicus* plants, NopT^{C93S}-FLAG was expressed

in hairy roots of the *nfr1-1* mutant and wild-type plants, respectively. After removal of

roots without fluorescence signals, plants were kept on half-strength B&D medium for 5

days. Roots were inoculated with *Mesorhizobium loti* MAFF303099 or treated with

water. Total proteins were extracted using the extraction buffer (Tris-HCl pH=7.3, 150

mM NaCl, 5 mM EDTA, 0.5% (w/v) SDS and 0.5% (v/v) Triton X-100) supplemented

with 2% protease inhibitor cocktail (Sigma, P9599) and 2% phosphatase inhibitor

cocktail (Yeasen, 20109-A). Proteins were then precipitated with TCA to remove various

contaminants (e.g., EDTA and surfactants). Phosphorylation of the NopT^{C93S} protein was

analyzed by 50 µM Zn^{2+}-Phos-tag SDS-PAGE gel (Kato & Sakamoto, 2019) and

detected with the anti-FLAG antibody by immunoblot analysis. As loading control, actin

proteins were immunodetected with an anti-Actin antibody.

BiFC and Split–LUC assays

In the BiFC assay with *N. benthamiana* plants, the eYFP fluorescence of expressed

fusion proteins was recorded 2-3 days post infiltration with *A. tumefaciens* using a

confocal microscope (Leica TCS SP8). For the Split–LUC assay, *N. benthamiana* leaves

were collected at 2 days post infiltration and sprayed with a solution of 1 mM D-luciferin

(Promega, E1603) and 0.02% (v/v) Triton X-100. After a dark adaptation of 5 min,

bioluminescence images were acquired using a Tanon Bio–Imaging System (Tanon,

Shanghai, 4600).

LC-MS analysis

For analysis of NopT phosphorylation sites, NopT-FLAG-His was co-expressed with His

labelled NFR1^{CD} or NFR1$^{CD-K351E}$ in *E. coli* cells. The proteins were purified using

Ni-charged resin (GenScript, Cat. No. L00223). The bands containing NopT^P and NopT

(control) proteins, were cut out, destained by 50 mM triethylammonium bicarbonate

(TEAB) solution (50% (v/v) acetonitrile in 50 mM TEAB), and washed with 100%

acetonitrile until the gel turned white. The protein bands were then in-gel digested with

sequencing-grade trypsin. LC-MS analysis was performed by Novogene Co., Ltd.

(Beijing, China).

For identificaiton of the cleavage site in NFR5, NopT and His-SUMO-NFR5^{CD}-GST,

were co-expressed in *E. coli* cells and purified using Glutathione Resin (GenScript, #

L00206). About 80 μg of the cleaved protein was collected for analysis. The protein

bands were digested with trypsin, glu-C, chymotrypsin and pepsin. LC-MS analysis was

performed by Oulu Biotechnology Co., Ltd. (Shanghai, China).

Examination of rhizobial infection

The morphology of rhizobial infection foci and infection threads have been described

previously (Rae et al., 2021). An infection focus is formed when the tip of a root hair

curls around a single bacterium, forming an infection pocket. The infection thread is a

tubular structure forming in the root hair, then extends to cortical cells. The number of

infection foci and infection threads was determined using a Leica fluorescence

microscope (DM2500). Whole roots were mounted on microscope slides and infection

foci and infection thread counts were expressed on a per-root basis.

Histochemical GUS staining of *L. japonicus* Gifu B-129 or Gifu *pNIN*:*GUS* roots

inoculated with different rhizobial strains was performed using the chromogenic

substrate 5-bromo-4-chloro-3-indolyl-β-D-glucuronide as described (Heckmann et al.,

2011). After evacuation of air for 20 min, root samples were incubated in GUS

staining solution overnight (at 37°C in the dark). The GUS-stained tissues were

observed and photographed using a Motic Swift M200D compound microscope

(Gifu B-129 roots inoculated with rhizobial strains harboring pT7-GUS) or a Nikon

SMZ18 stereo microscope (Gifu *pNIN*:*GUS* inoculated with unlabeled rhizobia).

Acknowledgments: We greatly thank Prof. Zhongming Zhang for his valuable

suggestions on this project. We thank Drs. Eric Giraud, Ertao Wang, and Yan

Liang for critical reading of this manuscript. We thank Dr. Jens Stougaard for

kindly providing *L. japonicus pNIN*:*GUS* and *nfr* mutant seeds. Dr. Juan Du is

thanked for providing plasmids for expression of Avr3a and R3a, BAX, and INF1.

The qPCR and microscopy data were acquired from the Core Facility Center at

the National Key Lab of Agricultural Microbiology.

Funding: This work was supported by the National Key R&D Program of China

(2024YFA0918204 and 2019YFA0904700), the National Natural Science Foundation of

China (32090063), and the Fundamental Research Funds for the Central Universities

(2662022SKYJ002 and 2662021JC010). Work in the Stacey's lab was supported by the

NSF Plant Genome Research Program (2048410).

Author Contributions: H.B., Y.C., and G.S. conceived the idea and designed the

research; H.B., Y.W., Y.L., and Y.Y. performed experiments; H.B., Y.W., H. L., H.Z.,

and Y.C. analyzed the data; Q.W. and S.X. identified the cleavage site by MS. H.B., Y.C.

S.F.W., C.S., and G.S. wrote the paper; H.B., Y.W., D.X., and Y.C. revised the paper.

Additional files

Supplementary files

MDAR checklist

Data availability

All data generated or analyzed during this study are included in the manuscript and

supporting files.

References

Akeda, Y., and Galán, J.E. (2005). Chaperone release and unfolding of substrates in
type III secretion. Nature **437,** 911-915.

Bos, J.I., Kanneganti, T.D., Young, C., Cakir, C., Huitema, E., Win, J., Armstrong,
M.R., Birch, P.R., and Kamoun, S. (2006). The C-terminal half of *Phytophthora*
infestans RXLR effector AVR3a is sufficient to trigger R3a-mediated hypersensitivity and
suppress INF1-induced cell death in *Nicotiana benthamiana*. Plant J **48,** 165-176.

Bozsoki, Z., Cheng, J., Feng, F., Gysel, K., Vinther, M., Andersen, K.R., Oldroyd, G.,
Blaise, M., Radutoiu, S., and Stougaard, J. (2017). Receptor-mediated chitin
perception in legume roots is functionally separable from Nod factor perception. Proc Natl
Acad Sci U S A **114,** E8118-E8127.

Broghammer, A., Krusell, L., Blaise, M., Sauer, J., Sullivan, J.T., Maolanon, N.,
Vinther, M., Lorentzen, A., Madsen, E.B., Jensen, K.J., Roepstorff, P., Thirup, S.,
Ronson, C.W., Thygesen, M.B., and Stougaard, J. (2012). Legume receptors perceive
the rhizobial lipochitin oligosaccharide signal molecules by direct binding. Proc Natl Acad
Sci U S A **109,** 13859-13864.

Cao, Y., Liang, Y., Tanaka, K., Nguyen, C.T., Jedrzejczak, R.P., Joachimiak, A., and
Stacey, G. (2014). The kinase LYK5 is a major chitin receptor in *Arabidopsis* and forms a
chitin-induced complex with related kinase CERK1. Elife **3**.e03766.

Cai J, Zhang LY, Liu W, Tian Y, Xiong JS, Wang YH, Li RJ, Li HM, Wen J, Mysore
KS, Boller T, Xie ZP, Staehelin C. (2018). Role of the Nod factor hydrolase MtNFH1 in
regulating Nod factor levels during rhizobial infection and in mature nodules of *Medicago*
truncatula. Plant Cell **30,** 397-414.

Dai, W.J., Zeng, Y., Xie, Z.P., and Staehelin, C. (2008). Symbiosis-promoting and
deleterious effects of NopT, a novel type 3 effector of *Rhizobium* sp. strain NGR234. J
Bacteriol **190,** 5101-5110.

Dowen, R.H., Engel, J.L., Shao, F., Ecker, J.R., and Dixon, J.E. (2009). A family of

bacterial cysteine protease type III effectors utilizes acylation-dependent and

-independent strategies to localize to plasma membranes. J Biol Chem **284,**

15867-15879.

Fotiadis, C.T., Dimou, M., Georgakopoulos, D.G., Katinakis, P., and Tampakaki, A.P.

(2012). Functional characterization of NopT1 and NopT2, two type III effectors of

Bradyrhizobium japonicum. FEMS Microbiol Lett **327,** 66-77.

Gan, Y., Zhang, L., Zhang, Z., Dong, S., Li, J., Wang, Y., and Zheng, X. (2009). The

LCB2 subunit of the sphingolip biosynthesis enzyme serine palmitoyltransferase can

function as an attenuator of the hypersensitive response and Bax-induced cell death.

New Phytol **181,** 127-146.

Geurts, R., Fedorova, E., and Bisseling, T. (2005). Nod factor signaling genes and their

function in the early stages of infection. Curr Opin Plant Biol **8,** 346-352.

Goehre, V., Spallek, T., Haeweker, H., Mersmann, S., Mentzel, T., Boller, T., de

Torres, M., Mansfield, J.W., and Robatzek, S. (2008). Plant pattern-recognition

receptor FLS2 is directed for degradation by the bacterial ubiquitin ligase AvrPtoB. Curr

Biol **18,** 1824-1832.

Gysel, K., Laursen, M., Thygesen, M.B., Lironi, D., Bozsoki, Z., Hjuler, C.T.,

Maolanon, N.N., Cheng, J., Bjork, P.K., Vinther, M., Madsen, L.H., Rubsam, H.,

Muszynski, A., Ghodrati, A., Azadi, P., Sullivan, J.T., Ronson, C.W., Jensen, K.J.,

Blaise, M., Radutoiu, S., Stougaard, J., and Andersen, K.R. (2021). Kinetic

proofreading of lipochitooligosaccharides determines signal activation of symbiotic plant

receptors. Proc Natl Acad Sci U S A **118**.e2111031118.

Han, Y., Sun, J., Yang, J., Tan, Z., Luo, J., and Lu, D. (2017). Reconstitution of the

plant ubiquitination cascade in bacteria using a synthetic biology approach. Plant J **91,**

766-776.

Heckmann, A.B., Sandal, N., Bek, A.S., Madsen, L.H., Jurkiewicz, A., Nielsen, M.W.,

Tirichine, L., and Stougaard, J. (2011). Cytokinin induction of root nodule primordia in

Lotus japonicus is regulated by a mechanism operating in the root cortex. Mol Plant

Microbe Interact **24,** 1385-1395.

Jones, J.D.G., and Dangl, J.L. (2006). The plant immune system. Nature **444,** 323-329.

Kambara, K., Ardissone, S., Kobayashi, H., Saad, M.M., Schumpp, O., Broughton,

W.J., Deakin W.J. (2009). Rhizobia utilize pathogen-like effector proteins during

symbiosis. Mol Microbiol **71,** 92-106.

Kawaharada, Y., Kelly, S., Nielsen, M.W., Hjuler, C.T., Gysel, K., Muszynski, A.,

Carlson, R.W., Thygesen, M.B., Sandal, N., Asmussen, M.H., Vinther, M., Andersen,

S.U., Krusell, L., Thirup, S., Jensen, K.J., Ronson, C.W., Blaise, M., Radutoiu, S.,

and Stougaard, J. (2015). Receptor-mediated exopolysaccharide perception controls

bacterial infection. Nature **523,** 308-312.

Kato, Y., and Sakamoto, W. (2019). Phosphorylation of the chloroplastic

metalloprotease FtsH in *Arabidopsis* characterized by phos-tag SDS-PAGE. Front Plant

Sci **10**, 1080.

Khan, A., Wadood, S.F., Chen, M., Wang, Y., Xie, Z., and Staehelin, C. (2022).

Effector-triggered inhibition of nodulation: A rhizobial effector protease targets soybean

kinase GmPBS1-1. Plant Physiol **189,** 2382-2395.

Kim, S.H., Qi, D., Ashfield, T., Helm, M., and Innes, R.W. (2016). Using decoys to

expand the recognition specificity of a plant disease resistance protein. Science **351,**

684-687.

Kimbrel, J.A., Thomas, W.J., Jiang, Y., Creason, A.L., Thireault, C.A., Sachs, J.L.,

and Chang, J.H. (2013). Mutualistic co-evolution of type III effector genes in

Sinorhizobium fredii and *Bradyrhizobium japonicum*. PLoS Pathog **9,** e1003204.

Kopan, R., and Ilagan, M.X.G. (2009). The canonical notch signaling pathway: Unfolding

the activation mechanism. Cell **137,** 216-233.

Lei, L., Stevens, D.M., and Coaker, G. (2020). Phosphorylation of the *Pseudomonas*

effector AvrPtoB by *Arabidopsis* SnRK2.8 is required for bacterial virulence. Mol Plant **13,**

1513-1522.

Li, D., Zhu, Z., Deng, X., Zou, J., Ma, C., Li, C., Yin, T., Liu, C., Wang, J., Chen, Q.,

and Xin, D. (2023). GmPBS1, a hub gene interacting with rhizobial type-III effectors

NopT and NopP, regulates soybean nodulation. Agronomy **13,** 1242.

Li, H., Chen, M., Duan, L., Zhang, T., Cao, Y., and Zhang, Z. (2018). Domain swap

approach reveals the critical roles of different domains of SYMRK in root nodule

symbiosis in *Lotus japonicus.* Front Plant Sci **9,** 697.

Li, L., Kim, P., Yu, L., Cai, G., Chen, S., Alfano, J.R., and Zhou, J. (2016).

Activation-dependent destruction of a co-receptor by a *Pseudomonas syringae* effector

dampens plant immunity. Cell Host Microbe **20,** 504-514.

Luo, Y., Liu, D., Jiao, S., Liu, S., Wang, X., Shen, X., and Wei, G. (2020) Identification

of *Robinia pseudoacacia* target proteins responsive to *Mesorhizobium amphore*

CCNWGS0123 effector protein NopT. J Exp Bot **71,** 7347-7363.

Madsen, E.B., Madsen, L.H., Radutoiu, S., Olbryt, M., Rakwalska, M., Szczyglowski,

K., Sato, S., Kaneko, T., Tabata, S., Sandal, N., and Stougaard, J. (2003). A receptor

kinase gene of the LysM type is involved in legume perception of rhizobial signals. Nature

425, 637-640.

Madsen, E.B., Antolin-Llovera, M., Grossmann, C., Ye, J., Vieweg, S., Broghammer,

A., Krusell, L., Radutoiu, S., Jensen, O.N., Stougaard, J., and Parniske, M. (2011).

Autophosphorylation is essential for the *in vivo* function of the *Lotus japonicus* Nod factor

receptor 1 and receptor-mediated signalling in cooperation with Nod factor receptor 5.

Plant J **65,** 404-417.

Moling, S., Pietraszewska-Bogiel, A., Postma, M., Fedorova, E., Hink, M.A.,

Limpens, E., Gadella, T.W., and Bisseling, T. (2014). Nod factor receptors form

heteromeric complexes and are essential for intracellular infection in *Medicago* nodules.
Plant Cell **26,** 4188-4199.

Nakagawa, T., Suzuki, T., Murata, S., Nakamura, S., Hino, T., Maeo, K., Tabata, R.,
Kawai, T., Tanaka, K., Niwa, Y., Watanabe, Y., Nakamura, K., Kimura, T., and
Ishiguro, S. (2007). Improved Gateway binary vectors: high-performance vectors for
creation of fusion constructs in transgenic analysis of plants. Biosci Biotechnol Biochem
71, 2095-2100.

Ntoukakis, V., Mucyn, T.S., Gimenez-Ibanez, S., Chapman, H.C., Gutierrez, J.R.,
Balmuth, A.L., Jones, A.M., and Rathjen, J.P. (2009). Host inhibition of a bacterial
virulence effector triggers immunity to infection. Science **324,** 784-787.

Radutoiu S, Madsen LH, Madsen EB, Felle HH, Umehara Y, Grønlund M, Sato S,
Nakamura Y, Tabata S, Sandal N, Stougaard J. (2003). Plant recognition of symbiotic
bacteria requires two LysM receptor-like kinases. Nature **425,** 585-592.

Rae, A.E., Rolland, V., White, R.G., and Mathesius, U. (2021). New methods for
confocal imaging of infection threads in crop and model legumes. Plant Methods **17**, 24.

Ried, MK, Antolín-Llovera, M, and Parniske, M. (2014). Spontaneous symbiotic
reprogramming of plant roots triggered by receptor-like kinases. Elife **3,** e03891.

Schauser, L., Roussis, A., Stiller, J., and Stougaard J. (1999). A plant regulator
controlling development of symbiotic root nodules. Nature **402,** 191-195.

Schmidt, G. (2011). *Yersinia enterocolitica* outer protein T (YopT). Eur J Cell Biol **90,**
955-958.

Schubert, K.A., Xu, Y., Shao, F., and Auerbuch, V. (2020). The type III secretion
system as a tool for studying cytosolic innate immune surveillance. Ann Rev Microbiol **74,**
221-245.

Shao, F., Merritt, P.M., Bao, Z., Innes, R.W., and Dixon, J.E. (2002). A *Yersinia* effector
and a *Pseudomonas* avirulence protein define a family of cysteine proteases functioning
in bacterial pathogenesis. Cell **109,** 575-588.

Shao, F., Vacratsis, P.O., Bao, Z., Bowers, K.E., Fierke, C.A., and Dixon, J.E.

(2003a). Biochemical characterization of the *Yersinia* YopT protease: cleavage site and

recognition elements in Rho GTPases. Proc Natl Acad Sci U S A **100,** 904-909.

Shao, F., Golstein, C., Ade, J., Stoutemyer, M., Dixon, J.E., and Innes, R.W. (2003b).

Cleavage of *Arabidopsis* PBS1 by a bacterial type III effector. Science **301,** 1230-1233.

Sugawara, M., Takahashi, S., Umehara, Y., Iwano, H., Tsurumaru, H., Odake, H.,

Suzuki, Y., Kondo, H., Konno, Y., Yamakawa, T., Sato, S., Mitsui, H., and

Minamisawa, K. (2018). Variation in bradyrhizobial NopP effector determines symbiotic

incompatibility with soybeans via effector-triggered immunity. Nature Commun **9**.3139.

Sun, Y., Li, L., Macho, A.P., Han, Z., Hu, Z., Zipfel, C., Zhou, J., and Chai, J. (2013).

Structural basis for flg22-induced activation of the *Arabidopsis* FLS2-BAK1 immune

complex. Science **342,** 624-628.

Tang, D., Wang, G., and Zhou, J. (2017). Receptor kinases in plant-pathogen

interactions: More than pattern recognition. Plant Cell **29,** 618-637.

Teulet, A., Busset, N., Fardoux, J., Gully, D., Chaintreuil, C., Cartieaux, F., Jauneau,

A., Comorge, V., Okazaki, S., Kaneko, T., Gressent, F., Nouwen, N., Arrighi, J.F.,

Koebnik, R., Mergaert, P., Deslandes, L., and Giraud, E. (2019). The rhizobial type III

effector ErnA confers the ability to form nodules in legumes. Proc Natl Acad Sci U S A

116, 21758-21768.

Tirichine, L., Imaizumi-Anraku, H., Yoshida, S., Murakami, Y., Madsen, L.H., Miwa,

H., Nakagawa, T., Sandal, N., Albrektsen, A.S., Kawaguchi, M., Downie, A., Sato, S.,

Tabata, S., Kouchi, H., Parniske, M., Kawasaki, S., and Stougaard, J. (2006).

Deregulation of a Ca^{2+}/calmodulin-dependent kinase leads to spontaneous nodule

development. Nature **441,** 1153-1156.

Vleeshouwers, V.G., Driesprong, J.D., Kamphuis, L.G., Torto-Alalibo, T., Van't Slot,

K.A., Govers, F., Visser, R.G., Jacobsen, E., and Kamoun, S. (2006).

Agroinfection-based high-throughput screening reveals specific recognition of INF

elicitins in *Solanum*. Mol Plant Pathol **7,** 499-510.

Wang, D., Yang, S., Tang, F., and Zhu, H. (2012). Symbiosis specificity in the legume:

rhizobial mutualism. Cell Microbiol **14,** 334-342.

Wang, Q., Han, C., Ferreira, A.O., Yu, X., Ye, W., Tripathy, S., Kale, S.D., Gu, B.,

Sheng, Y., Sui, Y., Wang, X., Zhang, Z., Cheng, B., Dong, S., Shan, W., Zheng, X.,

Dou, D., Tyler, B.M., and Wang, Y. (2011). Transcriptional programming and functional

interactions within the *Phytophthora sojae* RXLR effector repertoire. Plant Cell **23,**

2064-2086.

Xiang, T., Zong, N., Zou, Y., Wu, Y., Zhang, J., Xing, W., Li, Y., Tang, X., Zhu, L.,

Chai, J., and Zhou, J. (2008). *Pseudomonas syringae* effector AvrPto blocks innate

immunity by targeting receptor kinases. Curr Biol **18,** 74-80.

Xin, D., Liao, S., Xie, Z., Hann, D.R., Steinle, L., Boller, T., Staehelin, C. (2012).

Functional analysis of NopM, a novel E3 ubiquitin ligase (NEL) domain effector of

Rhizobium sp. strain NGR234. PLoS Pathog **8(5),** e1002707.

Xu, N., Luo, X., Wu, W., Xing, Y., Liang, Y., Liu, Y., Zou, H., Wei, H., and Liu, J.

(2020). A plant lectin receptor-like kinase phosphorylates the bacterial effector AvrPtoB to

dampen Iits virulence in *Arabidopsis.* Mol Plant **13,** 1499-1512.

Yang, J., Lan, L., Jin, Y., Yu, N., Wang, D., and Wang, E. (2022). Mechanisms

underlying legume-rhizobium symbioses. J Integr Plant Biol **64,** 244-267.

Zhang, B., Wang, M., Sun, Y., Zhao, P., Liu, C., Qing, K., Hu, X., Zhong, Z., Cheng, J.,

Wang, H., Peng, Y., Shi, J., Zhuang, L., Du, S., He, M., Wu, H., Liu, M., Chen, S.,

Wang, H., Chen, X., Fan, W., Tian, K., Wang, Y., Chen, Q., Wang, S., Dong, F., Yang,

C., Zhang, M., Song, Q., Li, Y., and Wang, X. (2021). *Glycine max* NNL1 restricts

symbiotic compatibility with widely distributed *Bradyrhizobia vi*a root hair infection. Nat

Plants **7,** 73-86.

Wang Y., Bao H., Lei Y., Yuan L., Li H., Zhu H., Xin D., Staehelin C., Cao Y. (2024).

Rhizobial effector NopM ubiquitinates Nod factor receptor NFR5 and promotes rhizobial

infection in *Lotus japonicus.* BioRxiv DOI:10.1101/2024.04.28.591553.

Zhang, J., Li, W., Xiang, T., Liu, Z., Laluk, K., Ding, X., Zou, Y., Gao, M., Zhang, X.,

Chen, S., Mengiste, T., Zhang, Y., and Zhou, J. (2010). Receptor-like cytoplasmic

kinases integrate signaling from multiple plant immune receptors and are targeted by a

Pseudomonas syringae effector. Cell Host Microbe **7,** 290-301.

Zhou, J., Wang, P., Claus, L.A.N., Savatin, D.V., Xu, G., Wu, S., Meng, X., Russinova,

E., He, P., and Shan, L. (2019). Proteolytic processing of SERK3/BAK1 regulates plant

immunity, development, and cell death. Plant Physiol **180,** 543-558.

Zipfel, C., and Oldroyd, G.E.D. (2017). Plant signalling in symbiosis and immunity.

Nature **543,** 328-336.

Zipfel, C., Robatzek, S., Navarro, L., Oakeley, E.J., Jones, J.D.G., Felix, G., and

Boller, T. (2004). Bacterial disease resistance in through flagellin perception. Nature **428,**

764-767.

1030

1031

 Figures

Figure 1. NopT specifically suppresses NFR1/NFR5-triggered cell death in *N. benthamiana*. *Agrobacterium* strains harboring plasmid DNA encoding NopT or the empty vector (EV) were infiltrated into *N. benthamiana* leaves. At 12 hpi, a second infiltration was performed with an *Agrobacterium* strain containing a plasmid with *NFR1/NFR5* genes. Dashed red lines indicate leaf discs where different proteins were expressed. (A, B) NopT (A), but not NopT^{C93S} (B), a protease-inactive version of NopT, suppressed NFR1NFR5-induced cell death. (C) Expression of NopM-induced cell death in *N. benthamiana*. (D) Expression of NopT could not suppress the cell death response triggered by expression of Avr3a/R3a, NopM, BAX, INF1 or AtCERK1. Numbers in (D) represent the number of leaf discs showing cell death and total leaf discs tested. Cell death in leaf disc results in the formation of necrotic plaques, which restrains pathogens within deceased cells. These plaques commonly manifest as leaf dehydration, frequently accompanied by a translucent appearance. Brown and shriveled leaf discs serve as indicators of cell death. The pictures shown in this figure are representative of at least three independent biological replicates.

Figure 2. NopT interacts with NFR1 and NFR5. Interactions between NopT and NFR1 or NFR5 were detected using BiFC (Split-YFP) (A, B), Split-LUC complementation (C) and co-IP (D) assays in *N. benthamiana* leaves. (A, B) For BIFC analysis, nYFP and cYFP tags were fused at the C-terminus of NopT, NFR1, NFR5, and the flagellin receptor FLS2 (negative control). YFP fluorescence signals represent protein-protein interactions. Scale bar=25 μm. (C) For the Split-LUC complementation assay, the NLuc and CLuc tags were fused at the C-terminus of NopT, NFR1, NFR5 and FLS2 (negative control). Luminescence signals represent protein-protein interactions. (D) For the co-IP assay, HA-tagged NFR1 or NFR5 and FLAG-tagged NopT were expressed in *N. benthamiana* cells followed by immunoprecipitation using an anti-FLAG antibody. Immunoblot analysis was performed using anti-HA and anti-FLAG antibodies (NopT^C denotes the truncated version of NopT after autocleavage). The images and immunoblots shown in this figure are representative of three biological replicates

1066
1067

Figure 3. NopT proteolyzes NFR5 at its JM domain. Proteins with indicated tags were expressed in *N. benthamiana* (A), *L. japonicus* (B) or *E. coli* cells (C-G) and detected by immunoblotting. (A) NopT but not NopT[C93S] (a protease-dead version of NopT) cleaves NFR5-GFP protein expressed in *N. benthamiana* cells by releasing NFR5[CD]-GFP (NopT[C] denotes autocleaved NopT). The cleavage efficiency was marked under the lane. (B) NopT but not NopT[C93S] cleaves NFR5-Myc expressed in hairy roots of *L. japonicus* by releasing NFR5[CD]-Myc. The cleavage efficiency was marked under the lane. The asterisk indicates the HA-tagged NFR5 cleavage product containing the kinase domain and a

C-terminal tail region (C) Analysis of the CDs of NFR1 and NFR5 co-expressed

with NopT or NopT^{C93S} in *E. coli* cells. Cleavage of NFR5^{CD} was observed for NopT

but not NopT^{C93S}, while NFR1^{CD} was not proteolyzed by NopT. In the presence of

NFR1^{CD}, a slower migrating band was observed, possibly representing

phosphorylated NopT (NopT^{P}). The cleavage efficiency was marked under the

lane. (D) A repeat experiment confirmed that NopT is able to cleave NFR5^{CD} (the

asterisk indicates the HA-tagged NFR5 cleavage product containing the kinase

domain and a C-terminal tail region). The cleavage efficiency was marked under

the lane. (E) NopT cleaves His-SUMO-NFR5^{JM}-GFP (His-SUMO and GFP linked

by the JM domain of NFR5) *in vitro*. F.T. indicates proteins in flow through

samples after purification with Ni-beads. (F) NopT expressed in *E. coli* cells was

unable to cleave co-expressed NFR5^{CD-5m}-HA, a modified version of NFR5^{CD}-HA

in which five amino acids of the JM were substituted by other residues (S283Y,

G294Q, Y303S, A310I, T311Y). (G) NopT expressed in *E. coli* cells was unable to

cleave co-expressed *M. truncatula* NFP^{CD}-HA and the NFR5^{JM}-NFP^{KC}-HA fusion

protein, while NFP^{JM}-NFR5^{KC} was proteolyzed (KC stands for the kinase domain

and a C-terminal tail region, modified nod factor receptors without the JM).

Figure 4. Phosphorylation of NopT by NFR1 suppresses its proteolytic activity. (A)

In vivo phosphorylation assay with proteins expressed in *L. japonicus* roots

(wild-type and *nfr1-1* mutant plants) using Zn^{2+}-Phos-tag SDS-PAGE.

Phosphorylation of NopT^{C93S} was induced by inoculation with rhizobia

(*Mesorhizobium loti* MAFF303099) and was largely dependent on NFR1. (B) The

relative protein amount of each lane, as shown in (A), was quantified with ImageJ

software (three biological replicates). The value of the control band in each gel

was set to 1 for comparison. Values are means ± SEM. (C) NFR1^{CD} but not

NFR1$^{CD-K351E}$ phosphorylates NopT in *E. coli* cells. The phosphorylated full-length

form of NopT could be dephosphorylated by CIAP (gel band shift). Abbreviations:

NFR1^{CD-P}, autophosphorylated NFR1; NopT^{P}, phosphorylated NopT; NopT,

non-phosphorylated NopT; NopT^{C}, autocleaved NopT. (D) The phosphorylation

sites of NopT identified by LC-MS were either substituted to alanine (A) or

aspartate (D). The indicated NopT variants were subsequently tested for

1110   autocleavage and NFR5^{CD} proteolysis in *E. coli* cells. Wild-type NopT (WT) and

1111   protease-dead NopT^{C93S} were included into the analysis.

1112

Figure 5. NopT regulates rhizobial infection in _L. japonicus_. (A) Analysis of rhizobial infection in _L. japonicus_ roots inoculated with GUS-labelled _S. fredii_ NGR234 (wild-type; WT) or a _nopT_ knockout mutant (NGR234Δ_nopT_; abbreviated as Δ_nopT_ in other panels). Scale bar=100 µm. Infection represents both infection focus and infection thread. The infection threads in the root hairs were shown in the left image in the upper pannel, while the infection foci were shown in the left image in the middle pannel. (B) GUS staining pictures showing roots of _L. japonicus_ plants expressing GUS expression under control of the _NIN_ promoter (_pNIN_:_GUS_). Plants were inoculated with NGR234 or NGR234Δ_nopT_ and analyzed at 7 dpi. Scale bar=1 mm. (C) Infection data (ITs, infection threads) for roots shown in (A) at 7 dpi (n=10, Student's _t_-test; * indicates P < 0.01). (D)

Quantification of GUS staining sites for roots shown in (B) (n=10, Student's *t*-test, P <

0.01). (E) Nodule primordia formation in *L. japonicus* wild-type roots inoculated with

NGR234 (WT), NGR234*ΔnopT* or NGR234 over-expressing *nopT* (*pT7*:NopT). Roots

were analyzed at 14 dpi (n=16, Student's *t*-test: P < 0.01). (F) Infection data for wild-type

roots inoculated with NGR234 (WT), NGR234*ΔnopT* and NGR234*ΔnopT* expressing

indicated NopT variants at 7 dpi (n=5, Student's *t*-test: P < 0.01). (G) Analysis of

rhizobial infection in hairy roots of *L. japonicus* (wild-type) expressing GFP (EV, empty

vector control) or NopT. Plants were inoculated with DsRed-labeled *M. loti*

MAFF303099 and analyzed at 5 dpi (n=8, Student's *t*–test: P < 0.01). (H) Expression of

NFR5 and NFR5-NFP^{KC} in hairy roots of *nfr5-3* mutant plants. Plants were

inoculated with GFP-labelled NGR234 (WT) or NGR234*ΔnopT* and analyzed at 8

dpi. Roots expressing NFR5-NFP^{KC} showed high numbers of infection foci for both

strains whereas significant differences were observed for NFR5 expressing roots

(n=10, Student's *t*-test: P < 0.01).

1138

1139

Figure 6. *S. fredii* NopT proteins cleave NFR5 and working model for NopT of NGR234. (A) NopT of *S. fredii* NGR234 and homologs from other rhizobial strains were co-expressed with NFR5^{CD} in *E. coli* cells (NopT$_{234}$, NopT of NGR234; NopT1$_{110}$ and NopT2$_{110}$, NopT proteins of *B. diazoefficiens* USDA110; NopT$_{257}$, NopT of *S. fredii* USDA257; NopT$_{103}$, NopT of *S. fredii* HH103). Immunoblot analysis indicated NFR5 cleavage by NopT proteins from *S. fredii* strains. (B) Expression of NopT$_{257}$ in *N. benthamiana* could not inhibit the cell death triggered by co-expressed NFR1 and NFR5. (C) *L. japonicus* Gifu *pNIN:GUS* plants were

inoculated with *S. fredii* NGR234, NGR234*ΔnopT* and NGR234*ΔnopT* expressing

NopT$_{257}$ (*ΔnopT*+NopT$_{257}$). Roots were subjected to GUS staining at 7 dpi. Scale

bar=2.5 mm. (D) Quantitative analysis of GUS-stained roots shown in panel C (n=19,

Student's *t*–test: P < 0.01). (E) A proposed model for NopT of NGR234 interacting

with NFRs. NopT and NopT^{C} (autocleaved NopT) proteolytically cleave NFR5 at

the JM domain to release the intracellular domain of NFR5 (cleaved NFR5). NFR1

phosphorylates full-length NopT to block its proteinase activity. NopT^{C} cannot be

phosphorylated by NFR1.