The Rhizobial effector NopT targets Nod factor receptors to regulate symbiosis in *Lotus japonicus*

Hanbin Bao[1#], Yanan Wang[1#], Haoxing Li[1], Qiang Wang[1], Yutao Lei[1], Ying Ye[1], Syed F. Wadood[2], Hui Zhu[1], Christian Stahelin[2], Gary Stacey[3], Shutong Xu[1], Yangrong Cao[1*]

[1] National Key Lab of Agricultural Microbiology, Hubei Hongshan Laboratory, Huazhong Agricultural University, Wuhan, Hubei, 430070, China

[2] State Key Laboratory of Biocontrol and Guangdong Key Laboratory of Plant Resources, School of Life Sciences, Sun Yat-sen University, Guangzhou, 510275, China

[3] Divisions of Plant Science and Technology and Biochemistry, Christopher S. Bond Life Sciences Center, University of Missouri, Columbia, MO 65211, USA

# These authors contributed equally to this work.

* Corresponding author: Yangrong Cao, email: yrcao@mail.hzau.edu.cn

Short title: NopT associates with NFR1 and NFR5 to regulate symbiosis

Keywords: Legume-rhizobium symbiosis; Rhizobial effector; Nod factor receptors; NopT.

One-sentence summary: The rhizobial effector NopT and two Nod factor receptors of *Lotus japonicus*, NFR1 and NFR5, mutually interact to precisely regulate rhizobial symbiosis.

Abstract

It is well-documented that type-III effectors are required by Gram-negative pathogens to directly target different host cellular pathways to promote bacterial infection. However, in the context of legume-rhizobium symbiosis, the role of rhizobial effectors in regulating plant symbiotic pathways remains largely unexplored. Here, we show that NopT, a YopT-type cysteine protease of *Sinorhizobium fredii* NGR234 directly targets the plant's symbiotic signaling pathway by associating with two Nod factor receptors (NFR1 and NFR5 of *Lotus japonicus*). NopT inhibits cell death triggered by co-expression of NFR1/NFR5 in *Nicotiana benthamiana*. Full-length NopT physically interacts with NFR1 and NFR5. NopT proteolytically cleaves NFR5 both *in vitro* and *in vivo*, but can be inactivated by NFR1 as a result of phosphorylation. NopT plays an essential role in mediating rhizobial infection in *L. japonicus*. Autocleaved NopT retains the ability to cleave NFR5 but no longer interacts with NFR1. Interestingly, genomes of certain *Sinorhizobium* species only harbor *nopT* genes encoding truncated proteins without the autocleavage site. These results reveal an intricate interplay between rhizobia and legumes, in which a rhizobial effector protease targets NFR5 to dampen symbiotic signaling. NFR1 appears to counteract this process by phosphorylating the effector. This discovery highlights the role of a bacterial effector in regulating a signaling pathway in plants and opens up the perspective of developing kinase-interacting proteases to fine-tune cellular signaling processes in general.

Introduction

The legume–rhizobium symbiosis stands as one of the most crucial and intricate mutualistic interactions in nature. Its significance goes beyond the supply of nitrogen fixed by rhizobia, which influences plant growth and ecosystem sustainability, as this symbiosis is a pivotal model for investigating plant-microbe interactions (Yang et al., 2022). The orchestration of this symbiosis involves the perception of rhizobial signal molecules such as lipo–chitooligosaccharidic Nod factors (NFs) and surface

polysaccharides (Madsen et al., 2003; Tirichine et al., 2006; Kawaharada et al., 2015),

along with the secretion of proteins, including type three protein secretion system

(T3SS) effectors translocated into host plants (Sugawara et al., 2018; Zhang et al.,

2021).

Plant receptor heterocomplexes consisting of two Lysin-motif receptor kinases

(LYKs), play a central role in recognizing various microbial poly- and

oligosaccharide molecules. In non–legumes such as Arabidopsis, AtLYK5 and

AtCERK1 (Chitin elicitor receptor kinase) are required for perception of chitin, and

thereby trigger plant immunity to combat fungal pathogens (Cao et al., 2014). In

legumes, structurally similar Nod factor receptors (NFRs) such as NFR1 and NFR5 in

Lotus japonicus are vital for recognition of rhizobial NFs and initiation of symbiotic

signaling (Broghammer et al., 2012; Bozsoki et al., 2017). Accordingly, *nfr1* and *nfr5*

knockout mutants are almost completely unable to form nodules (Madsen et al., 2003;

Radutoiu et al., 2003). Over-expression of either NFR1 or NFR5 can activate NF

signaling, resulting in formation of spontaneous nodules in the absence of rhizobia

(Ried et al., 2014). The intricate action of NFRs is highlighted by the elictiation of a

hypersensitive cell death response when NFR1/NFR5 are simultaneous

over-expressed in *Nicotiana benthamiana* leaves (Madsen et al., 2011). Similarly,

defense-like responses in *Medicago truncatula* nodules were observed upon

over–expression of MtNFP (Nod factor perception, the ortholog of NFR5) (Moling et

al., 2014), perhaps reflecting a potential role of MtNFP–interacting LYKs involved in

plant–pathogen associations. It can therefore be expected that the protein abundance

of NFRs must be tightly controlled to avoid resistance responses, thereby ensuring

optimal root infection by rhizobia.

T3SS effectors of phytopathogenic Gram–negative bacteria play a critical role

in suppressing pattern-triggered immunity (PTI) in host plants (Jones and

Dangl, 2006). To avoid diseases, specific resistance (R) genes are employed

to directly or indirectly recognize effectors translocated into host cells.

Recognition of such avirulence (Avr) effectors often results in effector-triggered

immunity (ETI), a much stronger immune response which is usually associated

with programmed cell death (Jones and Dangl, 2006). In contrast to the

well–studied effector functions of phytopathogenic bacteria in suppressing

immunity through various mechanisms, it is largely unknown how T3SS

effectors from symbiotic rhizobia contribute to mutualistic interactions.

T3SS effectors of the YopT–type family, such as YopT of *Yersinia pestis*,

AvrPphB of *Pseudomonas syringae* pv. *phaseolicola* and rhizobial NopT

proteins, are evolutionary related cysteine proteases (Shao et al., 2002; Dai et

al., 2008). YopT cleaves Rho family GTPases leading to their inactivation to

disrupt the actin cytoskeleton of human host cells (Shao et al., 2003a).

AvrPphB can proteolytically cleave Arabidopsis receptor–like cytoplasmic kinase

family proteins including PBS1, PBS1-like proteins and BIK1, to suppress PTI (Shao

et al., 2003b; Zhang et al., 2010). Cleavage of PBS1 kinase by AvrPphB can be

regarded as a "decoy strategy" as it is associated with RPS5 (RESISTANCE TO

PSEUDOMONAS SYRINGAE5)-mediated ETI (Shao et al., 2003b). Depending on

the host legume, the NopT effector protease of *Sinorhizobium fredii* NGR234

can have an opposite function in regulating symbiosis (Dai et al., 2008; Kambara

et al., 2009). The heterologous expression of NopT from NGR234 in *S. fredii*

USDA 257 severely blocks nodulation in soybean Nanfeng 15 but not in other

soybean cultivars (Khan et al., 2022), implying that NopT can act as an Avr

effector. In contrast, NopT positively regulates nodule formation in the

interaction between *S. fredii* HH103 and soybean Suinong 14 (Li et al., 2023).

NopT from *Bradyrhizobium* sp. ORS3257 promotes rhizobial infection in nodules of

Aeschynomene indica (Teulet et al., 2019). NopT proteases have been

biochemically characterized. Some exhibit autocleavage activity and are

subsequently acylated at the newly formed N-terminus (Dai et al., 2008;

Dowen et al., 2009; Kambara et al., 2009; Fotiadis et al., 2012; Khan et al.,

2022). Recently, soybean PBS1-1 and two proteins in *Robinia pseudoacacia*

were identified as NopT targets (Luo et al., 2020; Khan et al. 2022; Li et al.

2023). However, at the molecular level, it remains largely unknown how NopT

modulates rhizobial infection and whether NopTs from different rhizobial

strains differ in their effector activities.

A challenge in the investigation of rhizobial effectors lies in the availability of

suitable assays and plants with which effector activities can be characterized. In this

study, we took advantage of the observation that ectopic expression of NFR1 and

NFR5 in *N. benthamiana* leaves induces programmed cell death (Madsen et al., 2011).

We hypothesized that this response could be modulated by co-expressed rhizobial

effectors. Among the 15 known or putative T3SS effectors of *S. fredii* NGR234, only

NopT could specifically suppress NFR1/NFR5-induced cell death. This observation

suggested that NopT is an effector associated with NFRs. Subsequent experiments

demonstrated that NopT interacts with both NFR1 and NFR5 at the plasma membrane

and proteolytically cleaves NFR5 at the juxtamembrane (JM) domain to suppress

NF-mediated host responses. NFR1 phosphorylates the full-length NopT, thereby

inactivating its protease activity. However, the autocleaved NopT form of *S. fredii*

NGR234 retains the ability to cleave NFR5 but loses its ability to interact with NFR1.

Intriguingly, various *Sinorhizobium* strains possess only truncated versions of NopT

(related to autocleaved NopT of NGR234) which can evade the inactivation by NFR1.

We present in this study a model of mutual regulation, in which the rhizobial effector

NopT directly targets NFR5 to dampen symbiotic signaling, whereas NFR1-mediated

phosphorylation of NopT counteracts NFR5 cleavage. Our work provides essential

insights into the intricate interplay between legumes and rhizobia and shows an

example of how a rhizobal effector fine-tunes NF signaling by directly targeting

NFRs, and how an NFR inactivates the enzymatic activity of the effector through

phosphorylation.

Results

NopT expression in *N. benthamiana* suppresses the NFR1/NFR5-induced cell
147

death response
148

To examine the role of effector proteins in regulating rhizobial infection, 15 effector
150

genes (Kimbrel et al., 2013) were cloned from *S. fredii* NGR234, a strain with an
151

exceptionally broad host-range. Initial experiments were performed to screen the
152

effector proteins for their ability to suppress or enhance the NFR1/NFR5–triggered
153

cell death response of *N. benthamiana* leaves. In this experiment, each effector was
154

co-expressed with NFR1/NFR5 (Fig. S1A–S1C). Only NopT, a cysteine protease
155

belonging to the YopT family (Shao et al., 2002; Dai et al., 2008), could suppress
156

NFR1 and NFR5-induced cell death (Fig. 1A and S1D–S1F). Interestingly, NopT
157

reduced protein abundance of the intact NFR5 when co–expressed with NFR1 (Fig.
158

S1E), suggesting that NopT promotes proteolytic degradation of NFR5. Cys-93 of
159

NopT is an amino acid residue known to be essential for the protease activity of this
160

effector (Dai et al., 2008; Kambara et al., 2009). In contrast to the wild-type NopT
161

protein, expression of the protease-dead NopT^{C93S} form in *N. benthamiana* did not
162

result in suppression of cell death induced by NFR1/NFR5 (Fig. 1B), indicating that
163

the protease activity of NopT is essential for the observed cell death-suppressing
164

activity of NopT. In contrast to NopT, NopM, an effector protein from *S. fredii*
165

NGR234 reported to possess E3 ligase activity (Xin et al., 2012), induced cell death
166

when expressed in *N. benthamiana* leaves under our test conditions (Fig. 1C).
167

We also tested whether the cell death-suppressing effect of NopT was specific for
169

NFR1/NFR5 expression or whether NopT could act generally to suppress cell death.
170

In addition to NopM, we examined various proteins known to induce cell death when
171

expressed in *N. benthamiana* leaves, namely Avr3a and R3a (Bos et al., 2006),
172

apoptosis regulator BAX (Bcl-2-associated X protein) (Gan et al., 2009), INF1, an
173

elicitor from *Phytophthora infestans* (Vleeshouwers et al., 2006) and Arabidopsis

AtCERK1 (Cao et al., 2014) (Fig. 1C). In contrast to the NFR1/NFR5–induced cell

death, expression of NopT was unable to suppress cell death when each of these

proteins was expressed in *N. benthamiana* leaves (Fig. 1D). Hence, the effect of NopT

appears to be specific for the NFRs. These results suggested that the effector protease

NopT could act directly on NFR1 and/or NFR5.

NopT interacts with NFR1 and NFR5

Autocleaved NopT of *S. fredii* NGR234 contains acylation sites required for lipidation

and subsequent plasma membrane localization (Dowen et al., 2009; Khan et al.,

2022). To investigate whether NopT interacts with NFR1 and NFR5, we first used *in*

vivo bimolecular fluorescence complementation (BiFC) analysis. NopT, NFR1 and

NFR5 were C-terminally tagged with nYFP and cYFP. Fluorescence signals

representing direct NopT-NFR1, NopT-NFR5 and NopT-NopT interactions were

detected at the plasma membrane of *N. benthamiana* (Fig. 2A and 2B). Interactions

between NopT and NFR1 and NFR5 were further verified using a Split-luciferase

(Split–LUC) complementation assay (Fig. 2C). Co-expression of NopT-cLUC and

NFR1-nLUC or NFR5-nLUC produced strong luminescence signals in transformed

leaf discs compared with the negative control expressing AtFLS2-nLUC

(Arabidopsis FLAGELLIN SENSING2) and NopT-cLUC (Fig. 2C).

It is known that acylation of NopT can alter its subcellular localization (Dowen et al.,

2009; Khan et al., 2022). Here, we performed Split-LUC assays to examine the

interaction between NFR1/NFR5 and NopT$^{G50A/C51A/C52A}$, a modified NopT form

lacking acylation sites. Compared to the wild-type form (NopT^{WT}), the interactions

between NopT$^{G50A/C51A/C52A}$ and NFR1/NFR5 were significantly reduced in Split-LUC

assays (Fig. S2A and S2B). However, the protease-dead NopT form (NopT^{C93S})

showed interactions with NFR1 and NFR5 that were not different from NopT^{WT} (Fig.

S2C). These results were further confirmed by a co-immunoprecipitation (co-IP)

experiment with *N. benthamiana* leaves co-expressing FLAG-tagged NopT with

C-terminally HA-tagged NFR1 or NFR5.   Both NFR1 and LjNFR5 could be

co-precipitated by an anti-FLAG antibody when the sample contained NopT-FLAG

(Fig. 2D). Overall, the identified physical interactions between NopT and

NFR1/NFR5 indicated that NopT targets NFRs and that NFR5 and/or NFR1 may be

proteolytically cleaved by NopT.

NopT proteolytically cleaves NFR5 at the JM domain

A previous study has shown that NopT is autocleaved at its N-terminus to form a

processed protein that lacks the first 49 amino acid residues (Dai et al., 2008). To test

whether NopT could proteolyze NFR1 and/or NFR5, NopT was co-expressed with

NFR1-GFP and NFR-GFP fusion proteins in *N. benthamiana* leaves. The expressed

proteins were separated by SDS-PAGE and subjected to immunoblotting. Proteolytic

cleavage of NFR5-GFP but not NFR1-GFP was observed in the presence of

co-expressed NopT, whereas protease-dead NopT^{C93S} showed no effect (Fig. 3A and

S3A). A similar *in vivo* cleavage assay with transgenic *L. japonicus* roots also showed

that expression of NopT but not NopT^{C93S} caused proteolytic cleavage of NFR5 (Fig.

3B). Based on the molecular weight changes, the cleavage site in NFR5 was predicted

to occur within the cytoplasmic domain (CD) of NFR5. To further explore NFR5

cleavage by NopT, the CD of NFR5 (Strep-NFR5^{CD}-HA) and NopT-FLAG or

protease-dead NopT^{C93S}-FLAG were co-expressed in *Escherichia coli* cells.

Immunoblotting results with an anti-HA antibody showed that the active NopT

protease cleaved NFR5^{CD} resulting in a ~5 kDa smaller protein. However, when the

CD of NFR1 was co-expressed with NopT in the same experiment, no NFR1 cleavage

was detected (Fig. 3C and 3D). Migration of the bands representing full-length NopT

or NopT^{C93S} was significantly retarded on the gel when the CD of NFR1 was

co-expressed (Fig. 3C), suggesting that NopT was phosphorylated by NFR1 in *E. coli*.

233 The NFR5^{CD} cleavage product with a ~5 kDa lower molecular weight in the cleavage

234 assay suggested that the cleavage site is located within the juxtamembrane (JM)

235 domain of NFR5. To test this hypothesis, we replaced the JM with the SUMO tag to

236 creat SUMO-NFR5^{KC}-HA (KC, kinase domain and C-terminal tail region, a modified

237 NFR5^{CD} without the JM) for proteolytic assay. Indeed, immunoblot analysis showed

238 that co-expressed NopT was unable to cleave SUMO-NFR5^{KC}-HA (Fig. S3B). The

239 JM domain of NFR5 was then fused to SUMO and GFP to create a recombinant

240 protein, SUMO-NFR5^{JM}-GFP. In an *in vitro* cleavage assay with recombinant proteins

241 from *E. coli*, GFP was immunodetected in the flow though sample when

242 SUMO-NFR5^{JM}-GFP was incubated with NopT but not with NopT^{C93S} (Fig. 3E). The

243 interaction between NopT^{C93S} and SUMO-NFR5^{JM}-GFP was confirmed by an *in vitro*

244 pull-down assay (Fig. S4). Overall, these data strongly indicate that the cleavage site

245 of NFR5 for NopT is located in its JM domain.

247 **The cleavage of NFR5 by NopT is dependent on multiple residues of the JM**

248 **domain**

250 The "DKLLSGV" motif in the JM domain of NFR5 (residues Asp-288 to Val-294;

251 Fig. S5) shows highest similarity with the autocleavage region of NopT

252 (DKMGCCA). However, a protein variant of NFR5^{CD} in which the "DKLLSGV"

253 motif was replaced by 7 alanine residues could still be cleaved by NopT (Fig. S6A).

254 Similar to NopT, the AvrPphB effector of *P. syringa* has an autocleavage site (Shao et

255 al., 2003b). We therefore also examined a version of NFR5^{CD} in which the

256 "DKLLSGV" motif was replaced by that of AvrPphB. In the cleavage assay with *E.*

257 *coli* cells, AvrPphB did not cleave NFR5^{CD}. However, the modified NFR5^{CD} form

258 with the 7 residues from AvrPphB was cleaved by AvrPphB and the cleavage product

259 had a lower molecular weight than that formed by NopT (Fig. S6A). These findings

260 indicated that the cleavage site for NopT is located upstream of the "DKLLSGV"

261 motif in NFR5.

As mentioned above, NopT undergoes autocleavage but it is not clear whether this is

due to intermolecular or intramolecular proteolysis. In addition to Cys-93, His-205

and Aps-220 of NopT are conserved catalytic residues (Dai et al., 2008; Kambara et

al. 2009). Therefore, NopT forms lacking these catalytic residues were co-expressed

with NopT in *E. coli*. The immunoblot analysis showed that NopT could not

proteolyze NopT^{C93S}, NopT^{H205A} or NopT^{D220A} (Fig. S6B), suggesting that NopT

autocleavage is due to intramolecular proteolysis. Hence, the residues at the

autocleavage site of NopT may not help to predict the cleavage site of NopT

substrates.

In an attempt to define the NopT cleavage site in NFR5, we created a series of

modified NFR5 forms covering the JM domain of NFR5 (Val-269 to Cys-320) and

the first seven amino acid residues of the kinase domain (Lys-321 to Tyr-327).

Initially, 17 variants of NFR5^{CD} were created in which three adjacent amino acids in

the JM domain were replaced with three alanine residues. When these proteins were

co-expressed with NopT in *E. coli*, all 17 NFR5^{CD} variants could be cleaved by NopT

but not NopT^{C93S} (Fig.S6C-S6F). In a similar experiment, we created seven NFR5^{CD}

forms which contained a deletion in the JM domain of 10 residues each. Surprisingly,

cleavage products were still observed for all seven examined proteins (Fig. S6G).

These results suggest that the JM domain may have multiple sites that can be cleaved

by NopT, which is different from the specific cleavage site identified for AvrPphB. In

another experiment, recombinant His-SUMO-NFR5^{CD}-GST protein was co-expressed

with NopT in *E. coli* and its cleavage product was subjected to N-terminal protein

sequencing using Liquid Chromatography Mass Sprectrometry (LC-MS) analysis.

Based on alignment of identified peptides, the NopT cleavage site in NFR5 was

mapped to four basic amino acids in the JM domain (RRKK; amino acid residues

271-275) (Fig. S7). However, this result was not consistent with the observation that

NopT expressed in *E. coli* could still proteolyze a co-expressed NFR5^{CD} form in

which the residues from Tyr-268 to Leu-277 were deleted (Fig. S6G). Overall, the

experiments performed suggest that NopT preferentially proteolyzes NFR5 at the

RRKK motif, but other sites in the JM domain can also be cleaved by NopT. Since

NFR5 belongs to a subfamily of LYK family proteins that lacks kinase activity (Yang

et al., 2022), we wondered whether NopT could proteolyze other LYK proteins of this

subfamily, which might be helpful to characterize the biochemical function of NopT.

We therefore investigated whether the CDs from Arabidopsis AtLYK5, *L. japonicus*

LjLYS11 and *M. truncatula* MtNFP expressed in *E. coli* can be proteolyzed by NopT.

As shown in Fig. S8, NopT, but not NopT[C93S], was able to cleave the CDs of AtLYK5

and LjLYS11. Replacement of the JM domain in NFR5[CD] with the JM domain from

either AtLYK5 or LjLYS11 resulted in cleavage by NopT (Fig. S8). These data show

that NopT cleaves AtLYK5 and LjLYS11 at their JM domains. Based on sequence

analysis of the JM region of NFR5 and its homologous proteins, five conserved

residues (Ser-283, Gly-294, Tyr-304, Ala-310, Thr-311) were identified and

mutated as NFR5[CD-5m] for the proteolytic assay (Fig. S5B). As shown in Fig. 3F,

NFR5[CD-5m], a NFR5[CD] variant with point mutations of S283Y, G294Q, Y304S,

A310I, and T311Y, could not be cleaved by NopT. Interestingly, NopT could not

proteolyze MtNFP[CD] (Fig. 3G). We then swapped the JM domains in NFR5[CD] and

MtNFP[CD], generating NFP[JM]-NFR5[KC] and NFR5[JM]-NFP[KC] (KC stands for the kinase

domain and the C-terminal tail region). As shown in Fig. 3G, NopT could proteolyze

NFP[JM]-NFR5[KC] but not NFR5[JM]-NFP[KC] in *E. coli*. In an *in vitro* cleavage assay, the

cleavage product of SUMO-NFP[JM]-GFP proteolyzed by NopT was detected in the

flow though sample (Fig. S9). These data suggest that the cleavage of NFR5 by NopT

depends on the conformational structure of NFR5[KC] and/or other specific amino acids

in NFR5[KC]. We then investigated whether two additional recombinant proteins with

large swapping regions in NFR5[CD] and NFP[CD] (NFR5[268-445]-NFP[458-595] and

NFP[270-457]-NFR5[456-595]) could be cleaved in the *E. coli* proteolysis test (Fig. S10).

Notably, both proteins were cleaved by NopT, suggesting that a specific conformation

of the NopT substrate is required for proteolysis. Taken together, we concluded from

our experiments that NopT can proteolyze NFR5 at the JM domain and that mutiple

nonadjacent residues in NFR5[JM] in combination with a specific KC conformation are

required for proteolysis.

NopT phosphorylated by NFR1 is proteolytically inactive

The perception of rhizobial NFs by the heterocomplex of NFR1 and NFR5 involves

trans-phosphorylation events from the NFR1 kinase to the pseudo-kinase NFR5

(Madsen et al., 2011). Since NopT is associated with both NFR1 and NFR5, we

hypothesized that NopT is a phosphorylation target of NFR1, which may interfere

with transphosphorylation events between the two receptors. NopT^{C93S} was therefore

expressed in *L. japonicus* wild-type and *NFR1* knockout mutant plants (*nfr1-1*) and

analyzed on Zn^{2+}-Phos-tag gels. On such gels, a Zn^{2+}-Phos-tag bound

phosphorylated protein migrates slower than its unbound nonphosphorylated form.

Slower migrating bands corresponding to phosphorylated NopT^{C93S} were observed,

particularly when wild-type plants were inoculated with rhizobia (Fig. 4A and 4B).

These findings indicated that NopT phosphorylation *in planta* is largely dependent on

NFR1 and that the protein was probably phosphorylated at multiple sites.

Next, we performed assays to investigate whether expressed NFR1^{CD} directly

phosphorylates co-expressed NopT. NFR5^{CD} and a kinase-inactive NFR1^{CD} form

(NFR1$^{CD-K351E}$; Lys-351 is a conserved residue required for ATP binding) were

included into these experiments. When NFR1^{CD} was expressed in *E. coli*, extracted

NopT (Fig. 3C and 4C) and NFR5^{CD} (Fig. S11) exhibited retarded migration on

SDS-PAGE gels, indicating that NFR1 was able to phosphorylate both proteins. A

band shift was observed when the phosphorylated full-length NopT was *in vitro*

dephosphorylated by calf intestinal alkaline phosphatase (CIAP). However, the CIAP

treatment caused no obvious band shift of the autocleaved NopT form (Fig. 4C). The

phosphorylation sites of NopT were then identified using an *in vitro* phosphorylation

assay followed by LC-MS analysis. Several serine and threonine residues in NopT

were identified to be phosphorylated by NFR1^{CD} (Table S1). Taken together, these

results showed that full-length NopT is a phosphorylation target of the NFR1 kinase.

Finally, we wondered whether NopT phosphorylation by NFR1 influences its

proteolytic activity. Plasmids encoding NopT variants were constructed in which the

phosphorylated residues were substituted to either alanine (to block phosphorylation)

or aspartate (to mimic the phosphorylation status). The different NopT forms were

then expressed in *E. coli* to analyze NopT autocleavage and proteolysis of

co-expressed NFR5^{CD}. Aspartate substitutions at three phosphorylation sites of NopT

(Thr-97, Ser-119 and Thr-257) resulted in the loss of the autoproteolytic activity, as

well as the ability to cleave+ NFR5^{CD} (Fig. 4D). In contrast, corresponding NopT

proteins with alanine substitutions retained autocleavage activity and the ability to

proteolyze NFR5^{CD} (Fig. 4D). These data indicate that phosphorylation of full-length

NopT by NFR1 inhibits its proteolytic activity, thereby keeping NopT unprocessed

and NFR5 uncleaved.

NopT dampens rhizobial infection

As NopT of *S. fredii* NGR234 could directly target and cleave NFR5, we expected

that NopT may be an important player regulating rhizobial infection in legumes. NF

signaling is known to function during early stages of rhizobial infection (Geurts et al.,

2005; Wang et al., 2012; Cai et al., 2018). We focused our experiments on early

stages of rhizobial infection. Inoculation of *L. japonicus* plants with a GFP- or

GUS-labelled *nopT* mutant of NGR234 (NGR234 Δ*nopT*) resulted in a massive

infection of root hairs, whereas infection by the NGR234 wild-type strain was less

frequent (Fig. 5A and 5C). We also performed experiments with *L. japonicus* plants

containing *pNIN*:*GUS* which exhibit β-glucuronidase (GUS) activity when the *NIN*

(NODULE INCEPTION) promoter activity is induced by NF signaling (Schauser et

al., 1999). GUS staining of roots showed that the *NIN* promoter activity was stronger

in response to NGR234Δ*nopT* inoculation relative to inoculatioin by wild-type

NGR234 (Fig. 5B and 5D). Likewise, inoculation with NGR234Δ*nopT* resulted in an

increased number of nodule primordia (Fig. 5G). In contrast, over-expression of NopT

under the control of the T7 promoter in the NGR234 parent strain reduced the number

of nodule primordia when compared to inoculation with the wild-type strain (Fig.

5G). These findings indicate that NopT negatively influences symbiosis at early

symbiotic stages. However, we could not find any uninfected (empty) nodules when

roots were inoculated with these strains.

To confirm that the massive infection of roots by NGR234Δ*nopT* is due to the loss of

NopT, we complemented the NGR234Δ*nopT* by expressing wild-type NopT. The *L.*

japonicus plants inoculated with this complemented strain had fewer infection foci,

indicating restoration of the wild-type phenotype (Fig. 5E). We also tested whether

the proteolytic activity of NopT affected rhizobial infection. The protease-dead

version of NopT^{C93S} and three NopT forms with phospho-mimimetic residues (i.e.,

NopT^{T97D}, NopT^{S119D}, and NopT^{T257D} lacking proteolytic activity) were individually

expressed in NGR234Δ*nopT*. Compared with the high number of infection foci

induced by NGR234Δ*nopT* in *L. japonicus* plants, inoculation with these strains

resulted in significantly fewer infection foci, but more than with the wild-type strain

(Fig. 5E).

We also expressed *nopT* under the control of a ubiquitin promoter in hairy roots of *L.*

japonicus and inoculated the transgenic roots with DsRed-labeled *M. loti*

MAFF303099, which does not possess a *nopT* gene in its genome. As shown in Fig.

5F, reduced numbers of infection foci and infection threads were observed in the

NopT expressing roots, indicating that NopT negatively affects the symbiosis between

L. japonicus and *M. loti* strain.

To investigate whether the cleavage of NFR5 by NopT reduces rhizobial infection of

L. japonicus roots, *NFR5* and *NFR5^{5m}* (uncleavable variant with 5 amino acid

substitutions in the JM domain; Fig. 3F) under the control of the native *NFR5*

promoter, were expressed in hairy roots of *NFR5* knockout mutant plants (*nfr5-3*).

NFR5 expression resulted in numerous infections and a significant increase of

infection foci was observed upon inoculation with the NGR234Δ*nopT* mutant (Fig.

5H). However, NFR5^{5m} expression showed no effects on rhizobial infection (Fig. S12),

suggesting that the amino acid substitutions in NFR5^{5m} caused a conformational

change that rendered the protein inactive in symbiotic signaling. As the

NFR5^{JM}-NFP^{KC} fusion protein was not cleaved by NopT (Fig. 3G), we also

constructed hairy roots of *nfr5-3* mutant plants in which *NFR5-NFP^{KC}* (NFR5 variant

with its KC domain replaced by the KC of MtNFP) was expressed under the control

of the *NFR5* promoter. Remarkably, inoculation of these plants showed no difference

between NGR234Δ*nopT* and the NGR234 wild-type strain in terms of rhizobial

infection foci (Fig. 5H), suggesting that the cleavage of NFR5 by NopT reduced the

degree of rhizobial infection. Taken together, these data indicate that the protease

activity of NopT, its phosphorylation status and the proteolysis of the NFR5 target are

important for the regulation of rhizobial infection.

NopT from other *S. fredii* strains also cleave NFR5

Since NopT of *S. fredii* NGR234 (NopT$_{NGR234}$) cleaves NFR5, we wondered whether

homologs from other rhizobial species also possess such a proteolytic activity.

Phylogenetic analysis showed that NopT proteins of *Sinorhizobium* and

Bradyrhizobium are located in different clades (Fig. S13). *B. diazoefficiens*

USDA110, a typical *Bradyrhizobium* strain used to study nodulation of soybeans,

produces two NopT proteins (Fotiadis et al., 2012). In contrast to NopT$_{NGR234}$,

however, neither NopT1$_{USDA110}$ nor NopT2$_{USDA110}$ were able to cleave co-expressed

NFR5 in *E. coli* (Fig. 6A). This finding suggests that NopT homologs from

Bradyrhizbium species have lost the ability to cleave NFR5 and probably act on other,

unknown host target proteins. The genome of various *Sinorhizobium* species (e.g., *S.*

fredii USDA257 and S. *fredii* HH103) possess genes that encode truncated NopT

forms, which are almost identical to the autocleaved version of NopT$_{NGR234}$ (Khan et

al., 2022). Remarkably, NopT$_{USDA257}$ and NopT$_{HH103}$ were both able to cleave NFR5

in E. coli (Fig. 6A). However, both NopT proteins were unable to suppress cell death

triggered by NFR1 and NFR5 expression in *N. benthamiana* leaves (Fig. 6B).

Moreover, expression of NopT$_{USDA257}$ in NGR234Δ$nopT$ had no effect on the

infection ability of NGR234Δ$nopT$, as observed in inoculation tests with the *L.*

japonicus line containing *pNIN*:*GUS* (Fig. 6C and 6D). These results suggest that

NopT$_{USDA257}$ and NopT$_{HH103}$ exhibit similar proteolytic activity to NopT$_{NGR234}$, but

target different host proteins to regulate rhizobial infection.

Discussion

Plant-microbe interactions are complex,　employing distinct "chemical

weapons" or strategies to communicate mutually, thereby securing more

resources for their survival (Jones and Dangl, 2006; Schubert et al., 2020).

T3SS effectors of phytopathogenic bacteria are injected into plant cells to

facilitate bacterial infection by targeting host proteins of the plant's immune

system, thereby suppressing PTI (Jones and Dangl, 2006; Tang et al., 2017).

However, the role of rhizobial T3SS effectors in regulating the mutualistic

symbiosis between legumes and rhizobia remains largely unknown. The aim of

this study was to gain essential insights into the regulation mechanisms by

which rhizobial effectors act directly on symbiotic signaling pathways to

promote or dampen infection of host cells.

Understanding how plants recognize different microbes as "friends" or "foes" is

a central question in biology. The perception of microbial signals by plant

receptor proteins is essential for triggering specific signaling pathways to resist

invading pathogens or to establish symbiosis with beneficial microbes (Zipfel

and Oldroyd, 2017). Direct targeting of these receptors by T3SS effectors

represents an efficient strategy of phytopathogenic bacteria to suppress PTI

and promote bacterial infection. In the well-studied *Arabidopsis-Pseudomonas*

syringae interaction, T3SS effectors target flagellin receptor FLS2 and its

coreceptor BAK1 (BRI1-[Brassinosteroid insensitive1]-associated receptor

kinase 1) to suppress flagellin-induced PTI (Zipfel et al., 2004; Sun et al.,

2013). For example, AvrPto targets FLS2 and HopB1 cleaves BAK1 (Xiang et

al., 2008; Li et al., 2016). Likewise, AvrPtoB, an E3 ligase in *P. syringae*,

targets FLS2 to promote its degradation and suppress PTI (Goehre et al.,

2008). Despite the evolutionary importance of T3SS effectors in targeting host

receptors, we were surprised to find that NopT, a rhizobial member of the C58

protease effector family, can directly target NFRs and manipulate the

symbiotic signaling pathway in legumes. The interaction between NopT of *S.*

fredii NGR234 and NFRs of *L. japonicus* was identified by using a unique

screening system, in which co-expression of NFR1/NFR5 in *N. benthamiana*

leaves leads to cell death. NopT, but not other effectors of NGR234, could

suppress this cell death response. Consistent with these initial observations,

inoculation of *L. japonicus* with the *S. fredii* NGR234Δ*nopT* mutant resulted in

increased infection of *L. japonicus* roots. Moreover, over-expression of NopT in

the roots was found to reduce rhizobial infection by *M. loti*. The interaction

between NopT and NFR1/NFR5 was verified by BiFC, Split-LUC and co-IP

experiments.

Interestingly, using different assays, we found that NopT cleaves NFR5 at the

JM domain. However, the efficiencies of NopT cleavage of NFR5 in *L.*

japonicus and *N. benthamiana* were different (Fig. S14), which could be

related to the uncontrollable activation of NFRs in *N. benthamiana* or

phosphorylation of NopT by another kinase leading to decreased proteolytic

activity. The modification of five conserved residues in the JM domain and the

replacement of NFR5^{KC} by NFP^{KC} resulted in NFR5^{CD} forms that were

resistant to proteolysis by NopT in *E. coli* cells. These results suggest that

cleavage of NFR5 depends on both the JM domain and the KC region.

Although belonging to the same protease family, the NopT cleavage site in

NFR5 was found to be polybasic as in YopT substrate (Shao et al., 2003a;

Schmidt, 2011), whereas the cleavage site of AvrPphB substrate is canonical

with seven adjacent amino acids involved (Kim et al., 2016). These differences

reflect a broad variety of YopT-type effector proteases and their substrates in

host cells. Our study shows, that NopT targets the NFR1/NFR5 complex thus

reduces NF signaling and formation of nodule primordia in *L. japonicus*.

NopT-NFR interactions may reflect a regulation mechanism to prevent

rhizobial hyperinfection in host species other than *L. japonicus*.

Similar to the *Pseudomonas* effector AvrPphB, NopT of *S. fredii* NGR234 is

able to cleave soybean PBS1-1 and expression of this effector in *S. fredii*

USDA257 negatively influences symbiosis with certain soybean cultivars

(Khan et al., 2022). Indeed, NopT of NGR234 acts as an Avr effector in some

nonhost plants. Transient expression of NopT of strain NGR234 in Arabidopsis

and tobacco triggers strong immune responses and cell death (Dai et al., 2008;

Kimbrel et al., 2013; Khan et al., 2022; Fig. S15). In Arabidopsis,

NopT-induced cell death was found to be dependent on the PBS1-1 kinase

and the resistance protein RPS5 (Khan et al., 2022). Likewise, when

constitutively expressing NopT and NopT^{C93S} in *L. japonicus*, we only obtained

stable transgenic lines expressing NopT^{C93S}, indicating that the protease

activity of NopT had a negative effect on plant development. In contrast,

generation of hairy roots expressing NopT was possible. Such differences may

be explained by different NopT substrates in roots and aerial parts of the plant.

Indeed, our study shows that NopT not only cleaves NFR5 but is also able to

proteolyze Arabidopsis AtLYK5 and *L. japonicus* LjLYS11. These two

receptors possess chitin-binding affinities and trigger PTI responses (Cao et

al., 2014; Gysel et al., 2021). NFR5, AtLYK5, and LjLYS11 are kinase-dead

LYKs belonging to the same LYK subfamily. Thus, NopT appears not only to

suppress NF signaling, but may also interfere with signal transduction

pathways related to plant immunity. Depending on the host legume species,

NopT could suppress PTI or induce ETI, thereby modulating rhizobial infection

and nodule formation. Interactions between NopT and proteins related to the

plant immune system may represent an important evolutionary driving force for

host-specific nodulation and explain why the presence of NopT in NGR234 has

a negative effect on symbiosis with *L. japonicus* but a positive one with other

legumes.

The regulation of kinase-dead pseudokinases in plants may be more complex

compared to the well-studied control of protein kinase activities by

phosphorylation. The cleavage of kinase-dead receptors may be an

overlooked strategy to regulate their function in plant biology. Regulation of

protein activities by cleavage is well studied in animals, e.g., for Notch

signaling, where proteolysis of Notch molecules leads to downstream signaling

(Kopan and Ilagan, 2009). Arabidopsis BAK1 undergoes proteolytic cleavage,

which is important for both brassinosteroid signaling and induction of PTI

responses (Zhou et al., 2019). The cytoplasmic domain of NFR5 is a

pseudokinase that contributes to activation of NF signaling. Our work indicates

that NopT proteolytically cleaves NFR5 and suppresses NF signaling. Such a

fine-tuning process could be advantageous for the symbiosis in host plants

other than *L. japonicus*, for example to prevent hyperinfection. However, the

precise function of NFR5 cleavage by NopT and the fate of the released

cytoplasmic domain remain to be clarified.

Mutual regulation between T3SS effectors and host target proteins has been

investigated for various plant-pathogen interactions. For example, the

conserved *Pseudomonas* effector AvrPtoB can directly ubiquitinate the key

components of plant immunity to promote bacterial virulence (Goehre et al.,

2008). However, the activity of AvrPtoB in promoting virulence could either be

enhanced or dampened by host proteins. AvrPtoB can be phosphorylated by

SnRK2.8 to promote its virulence activity (Lei et al., 2020). On the other hand,

the plant lectin receptor-like kinase LexRK-IX.2 and the Pto kinase

phosphorylate AvrPtoB to inactivate its ubiquitin E3 ligase activity and

undermine the effector's ability to suppress PTI (Ntoukakis et al., 2009; Xu et

al., 2020). A similar strategy is used by NFR1 to inhibit the protease activity of NopT *via* phosphorylation. NFR1 expressed in *E. coli* phosphorylates full-length NopT of *S. fredii* NGR234 at several residues. Three NopT forms with a phosphomimetic substitution (i.e., NopT^{T97D}, NopT^{S119D}, and NopT^{T257D}) lacked autoproteolytic activity and the ability to cleave NFR5. The expression of these NopT variants in the NGR234Δ*nopT* mutant showed little effects in comparison with expression of wild-type NopT. These data suggest that *L. japonicus* utilizes NFR1 to phosphorylate NopT in order to dampen its catalytic activity and protect NFR5 from cleavage, which favors rhizobial infections. However, NopT$^{\Delta N50}$, which is similar to autocleaved NopT, retained the ability to interact with NFR5 but not with NFR1. Moreover, full-length NopT, but not autocleaved NopT, migrated slower on gels when NFR1^{CD} was co-expressed in *E. coli*, suggesting that autocleaved NopT may escape being phosphorylated and inactivated by NFR1. Future studies are required to explore the interactions between acylated NopT and NFR1. When expressed in *N. benthamiana*, GFP-tagged NopT forms lacking their autocleavage site (and thus non-acylated) were found to be localized within the cytoplasm, whereas NopT was targeted to the plasma membrane (Khan et al., 2022). Accordingly, NopT without acylation sites showed weaker interactions with NFRs in our study, suggesting that acylation of NopT, leading to plasma membrane location, promotes NopT-NFR interactions.

NopT homologs vary in different *S. fredii* strains. USDA257 and HH103 produce truncated NopT versions different from NGR234 due to a 19-bp deletion (Khan et al., 2022). Accordingly, NopT of these strains lacks autocleavage and acylation sites and possess a different N-terminal secretion signal sequence (Akeda and Galán, 2005). Nevertheless, NopT$_{HH103}$ is a functional T3SS effector as mutant analysis indicated a symbiosis-promoting role in certain soybean cultivars (Li et al., 2023). Similar to NopT$_{NGR234}$, NopT$_{USDA257}$ and NopT$_{HH103}$ were able to cleave NFR5 in *E. coli*. Since

NopT$_{USDA257}$ and NopT$_{HH103}$ cannot be acylated, they likely accumulate in the

cytoplasm of host legumes and are therefore less efficient in cleaving plasma

membrane-bound NFR5 in comparison to acylated NopT$_{NGR234}$. This could

also explain why NopT$_{USDA257}$ was unable to suppress cell death triggered by

NFR1/NFR5 in *N. benthamiana* leaves. It can be expected that NFR1 does not

interact with and phosphorylate NopT$_{USDA257}$ or NopT$_{HH103}$, as these effectors

are similar to NopT$^{\Delta N50}$ of NGR234, which cannot interact with NFR1.

Consistent with these findings, expression of NopT$_{USDA257}$ in NGR234*ΔnopT*

did not alter the infection phenotype of NGR234*ΔnopT*. It can be hypothesized

that the host targets of these truncated NopT effectors might be different from

those of NopT$_{NGR234}$. *Bradyrhizobium* NopT effectors are also different from

NopT$_{NGR234}$, as NopT of *B. diazoefficiens* USDA110 was unable to cleave

NFR5). NopT effectors may play a crucial role in *Bradyrhizobium*-legume

interactions. A *Bradyrhizobium* sp. ORS3257 mutant deficient in NopT

production induced only ineffective nodules on roots of *Aeschynomene indica*

and most of the formed nodules were not infected (Teulet et al., 2019). Overall,

these findings indicate versatile functions of NopT homologs in different strains

and suggest that NopT controls different regulation processes in specific host

cells.

We present here a model of mutual regulation, in which NopT proteolyzes

NFR5 to suppress NF signaling, whereas protease activity of full-length NopT

is suppressed by NFR1 *via* phosphorylation (Fig. 6E). NopT impairs the

function of the NFR1/NFR5 receptor complex. Cleavage of NFR5 by NopT

reduces its protein levels. Possible inhibitory effects of NFR5 cleavage

products on NF signaling are unknown but cannot be excluded. Inactivation of

NopT protease activity *via* phosphorylation by kinases such as NFR1 appears

to be a countermeasure of the host to enable symbiotic signaling and rhizobial

infection. Such feedback regulation could be the result of evolutionary

pressure to produce 'improved' NFRs, which are resistant to cleavage by NopT

and have an increased capacity to phosphorylate NopT. On the other hand,

suppression of NopT protease activity by NFRs may drive broad-host-range

strains such as NGR234 to produce higher levels and structurally modified

NFs.

Taken together, this study provides insights into a legume-rhizobium

interaction, in which the bacterium deploys an effector protease to dampen

symbiotic signaling, while the host plant counteracts by phosphorylating the

effector, leading to its inactivation. Our findings highlight the function of a

bacterial effector protease in regulating a symbiotic signaling pathway in legumes.

This opens up the perspective of developing specific kinase-interacting proteases to

reprogram and fine-tune cellular signaling processes in general.

Materials and Methods

Germination and growth of *Lotus japonicus*

Lotus japonicus (ecotype Gifu B-129) was provided by the Center for Carbohydrate

Recognition and Signaling (https://lotus.au.dk/), and used as wild-type for nodulation

assays. The Gifu *pNIN*:*GUS* transgenic line and *nfr* mutants (*nfr1-1* and *nfr5-3*) were

kindly provided by Dr. Jens Stougaard from the Aarhus University, Denmark

(Madsen et al., 2003; Radutoiu et al. 2003; Heckmann et al., 2011). All *L. japonicus*

seeds were treated with concentrated sulfuric acid for 10 min, followed by surface

sterilization in 1% (w/v) NaClO for 8 min. After incubation at 4°C in the dark for 2 d,

the seeds were placed on 0.8% (w/v) agar containing half-strength Murashige &

Skoog (MS) medium for germination at 22°C for 2 d in the dark. Seedlings were then

transferred to growth pots containing vermiculite supplied with half-strength B&D

(Broughton & Dilworth) medium without nitrogen under long-day conditions (16-h

light/8-h dark) at 22°C. Ten-day-old seedlings were inoculated with rhizobial cultures

(OD_{600}=0.02) for infection experiments.

Plasmid construction

All plasmids used in this study were generated using MultiF Seamless Assembly Mix

(Abclonal, # RK21020). The plasmid pGWB514 was used as an original plasmid for

construction of all binary vectors (Nakagawa et al., 2007). Briefly, different tags

including HA tag, Myc tag, FLAG tag, Strep tag, nLUC, cLUC and GFP, were

amplified using a forward primer containing a *Kpn*I site and ligated into pGWB514

digested with *Xba*I and *Sac*I using the seamless cloning method described above. The

generated plasmids were named pG5XX-HA, pG5XX-Myc, pG5XX-FLAG,

pG5XX-nLUC, pG5XX-cLUC, pG5XX-GFP, respectively. The coding sequences of

L. japonicus NFR1 and *NFR5* were cloned into pGWB514 and pG517 between

*Xba*I/*Kpn*I to generate NFR1-HA and NFR5-Myc fusion constructs under the control

of the cauliflower mosaic virus 35S promoter. The DNA fragment containing

pro35S:NFR5-Myc-NosT was then amplified and cloned into pG514-NFR1 at *Sbf*I.

The final plasmids contained *pro35S*:NFR1-HA-NosT and *pro35S*:NFR5-Myc-NosT.

Rhizobial genes encoding effectors from *S. fredii* NGR234 were cloned into

pG5XX-FLAG between *Xba*I/*Kpn*I. To detect the interaction between NFR1/NFR5

and NopT in *N. benthamiana*, the full-length *NFR1/NFR5*, *NopT* and *AtFLS2* coding

sequences were cloned into pG5XX-Nluc, pG5XX-Nluc, pSPYCE(MR) and

pSPYNE(R)173 (Waadt et al., 2008). For cleavage assays in *N. benthamiana* leaves,

the *NFR5* and *NopT* coding sequences were cloned into pG5XX-FLAG GFP and

pG5XX-Strep, respectively. For cleavage assays in *E. coli*, the sequence encoding the

cytoplasmic domain of NFR5 was cloned into pACYC duet Strep-HA (Han et al.,

2017), and the sequences encoding the kinase domain of NFR5 fused to the HA tag

and the JM domain of NFR5 fused to the GFP tag were cloned into pSUMO. The

point mutations in $NFR5^{CD}$ and *NopT* were created by site-directed mutagenesis PCR

using the templates pACYC duet Strep-$NFR5^{CD}$-HA and pET28a NopT-FLAG . For

NFR/NopT cleavage experiments with *L. japonicus* plants, NopT-FLAG or

NopT^{C93S}-FLAG was cloned into pUB-GFP between *Xba*I and *Kpn*I, and

35S-NFR5-Myc-NosT was cloned into pUB-NopT-GFP at *Pst*I. For NopT

phosphorylation experiments with *E. coli*, the sequence encoding the cytoplasmic

domain of NFR1 fused to a Myc tag was cloned into pCDF duet. For

complementation experiments with the NGR234Δ*nopT* mutant (Dai et al., 2008), a

1317-bp DNA fragment upstream of *NopT* and the coding sequence of a given *NopT*

sequence was fused using overlap PCR and cloned into pHC60 between *Xho*I and

*Kpn*I. For complementation experiments with the *nfr5-3* mutant, a 1316-bp *NFR5*

promoter DNA fragment was fused to NFR5-NFP^{KC} or NFR5^{5m}using overlap   PCR

and cloned into pUB-Cherry between *Pst*I and *Kpn*I.

Transient gene expression in *Nicotiana* leaves

The plasmids of *INF1*, *Avr3*, *BAX*, and *R3a* for transient expression in *Nicotiana*

benthamiana were kindly gifted by Dr. Juan Du from Huazhong Agricultural

University. *Agrobacterium tumefaciens* strain EHA105 carrying various constructs

were mixed with an *Agrobacterium* culture harboring the P19 suppressor in an

infiltration buffer (10 mM MgCl$_2$, 10 mM MES-KOH pH 5.8 and 200 μM

acetosyringone). After incubation for 2 h at room temperature, *Agrobacterium*

cultures were infiltrated into the leaves of 4-week-old *N. benthamiana* and *N. tabacum*

plants. Leaves were harvested and analyzed two-days post inoculation. The cell death

suppression experiments with *N. benthamiana* plants were performed as described

previously (Wang et al., 2011).

Hairy root transformation

Hairy root transformation of *L. japonicus* plants was performed as described

previously (Li et al., 2018). In brief, surface-sterilized seeds were germinated and

grown on MS plates without sucrose (23°C in the dark for the first 3 days and 23°C at

a 16-h light/8-h dark period for the following 2 days). The seeds were then cut at the

middle of the hypocotyl and co-cultivated with *A. rhizogenes* LBA1334 carrying

pUB-GFP , pUB-NopT-FLAG-GFP, pNFR5:NFR5-mCherry,

pNFR5:NFR5^{5m}-mCherry and pNFR5:NFR5-NFP^{KC}-mCherry, respectively (23°C in

the dark for the first 3 days and 23°C at a 16-h light/8-h dark period for the following

2 days). The plants were then transferred onto solid B5 medium. After about 15 days,

non-fluorescent hairy roots (lacking expression of GFP or mCherry) were removed.

Where indicated, the plants were inoculated with rhizobia: *S. fredii* NGR234,

NGR234Δ*nopT* or *Mesorhizobium loti* MAFF303099, and analyzed at indicated time

points.

Analysis of NFR5 cleavage by NopT and variants

In experiments with *E. coli*, Strep-NFR5^{CD}-HA or Strep-NFR5^{CD}-HA variants were

co-expressed with NopT-FLAG or FLAG tagged NopT variant or AvrPphB-FLAG

by using Duet vectors (Novagen; Han et al., 2017). Bacterial cultures were treated

with 0.5 mM IPTG at 22°C for 15 h. After washing the bacterial cells with PBS buffer,

the bacterial pellet was resuspended in extraction buffer consisting of Tris-HCl (pH

7.3), 150 mM NaCl, 5 mM EDTA, 0.5% (w/v) SDS and 0.5% (v/v) Triton X-100.

After adding SDS loading buffer and boiling at 100°C for 5 min, the extracted

proteins were subjected to immunoblot analysis using anti-HA-peroxidase (Sigma,

clone 3F10) or anti-FLAG (Sigma, F1804) antibodies. An anti-groEL antibody

(ABclonal, A0969) was used to confirm equal loading of proteins on gels.

For *in vitro* cleavage assays, His-SUMO-NFR5^{JM}-GFP and NopT-FLAG-His proteins

were expressed in *E. coli* and purified. The reaction mixture contained 3 μg

NopT-FLAG (or NopT^{C93S}-FLAG) and 1 μg SUMO-NFR5^{JM}-GFP in extraction

buffer without SDS and EDTA. After incubation at 22°C for 7 h, 10 μL of a

Ni-charged resin (GenScript, Cat. No. L00223) were added, and the cleavage products

in the flow through were detected by immunoblot analysis using an anti-GFP antibody

(ABclonal, AB_2770402).

To analyze cleavage of NFR5 in *L. japonicus* roots, hairy roots co-expressing

NopT-FLAG or NopT^{C93S}-FLAG with NFR5-Myc were harvested at about 15 days

after induction of hairy roots without rhizobia inoculation and subjected to

immunoblot analysis. Expressed proteins were extract by extraction buffer and

immunodetected with anti-FLAG antibodies and anti-Myc (Bio-Legend, # 626808)

antibodies.

For NFR5/NopT cleavage experiments with *N. benthamiana* plants, leaves expressing

proteins (NopT-FLAG or NopT^{C93S}-FLAG with NFR5-GFP) were analyzed. Leaf

discs were collected at 2 days post infiltration with agrobacteria and extracted in the

extraction buffer (Tris-HCl (pH 7.3), 150 mM NaCl, 5 mM EDTA, 0.5% (w/v) SDS

and 0.5% (v/v) Triton X-100) and 2% protease inhibitors (Sigma, P9599). NFR5 was

immunodetected with anti-Myc   and anti-GFP (ABclonal, AB_2770402) antibodies.

The NopT or NopT^{C93S} were detected with anti-FLAG antibody . The actin proteins

were detected with anti-Actin antibody (Abclonal, # AC009).

Phosphorylation assays

For NopT phosphorylation in *E. coli*, His-NFR1^{CD}-Myc or His-NFR1$^{CD-K351E}$-Myc

was co-expressed with NopT-FLAG-His by using Duet vectors (Novagen). Protein

expression was induced by 0.5 mM IPTG and cells were grown at 28°C for 18 h.

After simultaneous purification of NFR1^{CD} and NopT using Ni-charged resin, the

proteins were incubated with CIAP (TaKaRa, 2250B) at 37°C for 3 h. The mobility

shifts of NFR1^{CD} and NopT were detected with anti-Myc (Bio-Legend, 626808) and

anti-FLAG (Sigma, F1804) antibodies by immunoblot analysis.

In phosphorylation experiments with *L. japonicus* plants, NopT^{C93S}-FLAG was

expressed in hairy roots of the *nfr1-1* mutant and wild-type plants, respectively. After

removal of roots without fluorescence signals, plants were kept on half-strength B&D

medium for 5 days. Roots were inoculated with *Mesorhizobium loti* MAFF303099

or treated with water. Total proteins were extracted using the extraction buffer

(Tris-HCl (pH 7.3), 150 mM NaCl, 5 mM EDTA, 0.5% (w/v) SDS and 0.5% (v/v)

Triton X-100) supplemented with 2% protease inhibitor cocktail (Sigma, P9599) and

2% phosphatase inhibitor cocktail (Yeasen, 20109-A). Proteins were then precipitated

with TCA to remove various contaminants (e.g., EDTA and surfactant)s.

Phosphorylation of the NopT^{C93S} protein was analyzed by 50 μM Zn^{2+}-Phos-tag

SDS-PAGE gel (Kato & Sakamoto, 2019) and detected with the anti-FLAG antibody

by immunoblot analysis. As loading control, actin proteins were immunodetected with

an anti-Actin antibody.

BiFC and Split–LUC assays

In the BiFC assay with *N. benthamiana* plants, the eYFP fluorescence of expressed

fusion proteins was recorded 2-3 days post infiltration with *A. tumefaciens* using a

confocal microscope (Leica TCS SP8). For the Split–LUC assay, *N. benthamiana*

leaves were collected at 2 days post infiltration and sprayed with a solution of 1 mM

D-luciferin (Promega, E1603) and 0.02% (v/v) Triton X-100. After a dark adaptation

of 5 min, bioluminescence images were acquired using a Tanon Bio–Imaging System

(Tanon, Shanghai, 4600).

LC-MS analysis

For analysis of NopT phosphorylation sites, NopT-FLAG-His was co-expressed with

His labelled NFR1^{CD} or NFR1$^{CD-K351E}$ in *E. coli*. The proteins were purified using

Ni-charged resin (GenScript, Cat. No. L00223). The bands containing NopT^{P} and

NopT (control) proteins, were cut out, destained by 50 mM triethylammonium

bicarbonate (TEAB) solution (50% (v/v) acetonitrile in 50 mM TEAB), and washed

with 100% acetonitrile until the gel turned white. The protein bands were then in-gel

digested with sequencing-grade trypsin. LC-MS analysis was performed by Novogene

Co., Ltd. (Beijing, China).

For identificaiton of the cleavage site in NFR5, NopT and His-SUMO-NFR5^{CD}-GST,

were co-expressed in *E. coli* and purified using Glutathione Resin (GenScript, #

L00206). About 80 μg of the cleaved protein was collected for analysis. The protein

bands were digested with trypsin,glu-C, chymotrypsin and pepsin. LC-MS analysis

was performed by Oulu Biotechnology Co., Ltd. (Shanghai, China).

Examination of rhizobial infection

The morphology of rhizobial infection foci and infection threads have been described

previously (Rae et al., 2021). In brief, an infection focus is formed when the tip of a

root hair curls around a single bacterium, forming an infection pocket. The number of

infection foci and infection threads was determined using a Leica fluorescence

microscope (DM2500). Whole roots were mounted on microscope slides and

infection foci and infection thread counts were expressed on a per-root basis.

Histochemical GUS staining of *L. japonicus* Gifu B-129 or Gifu *pNIN*:*GUS* roots

inoculated with different rhizobial strains was performed using the chromogenic

substrate 5-bromo-4-chloro-3-indolyl-β-D-glucuronide as described (Heckmann et al.,

2011). After evacuation of air for 20 min, root samples were incubated in GUS

staining solution overnight (at 37°C in the dark). The GUS-stained tissues were

observed and photographed using a Motic Swift M200D compound

microscope (Gifu B-129 roots inoculated with rhizobial strains harboring

pT7-GUS) or a Nikon SMZ18 stereo microscope (Gifu *pNIN*:*GUS* inoculated with

unlabeled rhizobia).

Supplemental data

Supplemental Fig. 1. Analyses of the functions of all 15 effectors from *S. fredii*

NGR234 in preventing NFR1- and NFR5-induced cell death in *N. benthamiana*

leaves.

Supplemental Fig. 2. Split–LUC assay testing the interactions between

different NopT mutants and NFRs.

Supplemental Fig. 3. NopT cleaves NFR5 not not NFR1.

Supplemental Fig. 4. NopT interacts with the JM domain of NFR5.

Supplemental Fig. 5. Conserved domains and residues of NFR5 and related

proteins.

Supplemental Fig. 6. NopT cleaves NFR5 at the juxtamembrane domain.

Supplemental Fig. 7. Mass spectrometry analysis of cleavage site of NFR5 by

NopT.

Supplemental Fig. 8. NopT cleaves the NFR5 homolog proteins at the

juxtamembrane domain.

Supplemental Fig. 9. NopT cleaves the JM of MtNFP.

Supplemental Fig. 10. NopT cleaves recombinant proteins

NFR5$^{268-445}$-NFP$^{458-595}$ and NFP$^{270-457}$-NFR$5^{456-595}$.

Supplemental Fig. 11. NFR1^{CD} phosphorylates NFR5^{CD} *in vitro.*

Supplemental Fig. 12. Rhizobial infection in the mutant versions of NFR5.

Supplemental Fig. 13. Phylogenetic tree based on the amino acid sequence of

NopT homologs from different bacterial species.

Supplemental Fig. 14. The cleavage efficiency of NFR5 in different system.

Supplemental Fig. 15. Transient expression of NopT triggers cell death in

Arabidopsis thaliana and *Nicotiana tabacum***.**

Supplemental Table 1. Phosphopeptides of NopT identified by LC-MS.

Supplemental Table 2. Primers used in this study.

Acknowledgments: We greatly thank Prof. Zhongming Zhang for his valuable

suggestions on this project. We thank Drs. Eric Giraud, Ertao Wang, and Yan

Liang for critical reading of this manuscript. We thank Dr. Jens Stougaard for

kindly providing *L. japonicus pNIN:GUS* and *nfr* mutant seeds. Dr. Juan Du is

thanked for providing plasmids for expression of Avr3a and R3a, BAX, and

INF1. The qPCR and microscopy data were acquired from the Core Facility

Center at the National Key Lab of Agricultural Microbiology.

Funding: This work was supported by the National Key R&D Program of China

(2019YFA0904700), the National Natural Science Foundation of China (32090063),

and the Fundamental Research Funds for the Central Universities (2662022SKYJ002

and 2662021JC010). Work in the Stacey's lab was supported by the NSF Plant

Genome Research Program (2048410)..

Author Contributions: H.B., Y.C., and G.S. conceived the idea and designed the

research, H.B., Y.W., Y.L., and Y.Y. performed experiments; H.B., Y.W., H. L.,

H.Z., and Y.C. analyzed the data; Q.W. and S.X. identified the cleavage site by MS.

H.B., Y.C., S.F.W., C.S., and G.S. wrote the paper.

References

Akeda, Y., and Galán, J.E. (2005). Chaperone release and unfolding of substrates in type III secretion. Nature **437,** 911-915.

Bos, J.I., Kanneganti, T.D., Young, C., Cakir, C., Huitema, E., Win, J., Armstrong, M.R., Birch, P.R., and Kamoun, S. (2006). The C-terminal half of *Phytophthora infestans* RXLR effector AVR3a is sufficient to trigger R3a-mediated hypersensitivity and suppress INF1-induced cell death in *Nicotiana benthamiana*. Plant J **48,** 165-176.

Bozsoki, Z., Cheng, J., Feng, F., Gysel, K., Vinther, M., Andersen, K.R., Oldroyd, G., Blaise, M., Radutoiu, S., and Stougaard, J. (2017). Receptor-mediated chitin perception in legume roots is functionally separable from Nod factor perception. Proc Natl Acad Sci U S A **114,** E8118-E8127.

Broghammer, A., Krusell, L., Blaise, M., Sauer, J., Sullivan, J.T., Maolanon, N., Vinther, M., Lorentzen, A., Madsen, E.B., Jensen, K.J., Roepstorff, P., Thirup, S., Ronson, C.W., Thygesen, M.B., and Stougaard, J. (2012). Legume receptors perceive the rhizobial lipochitin oligosaccharide signal molecules by direct binding. Proc Natl Acad Sci U S A **109,** 13859-13864.

Cao, Y., Liang, Y., Tanaka, K., Nguyen, C.T., Jedrzejczak, R.P., Joachimiak, A., and Stacey, G. (2014). The kinase LYK5 is a major chitin receptor in *Arabidopsis* and forms a chitin-induced complex with related kinase CERK1. eLife **3**.e03766.

Cai J, Zhang LY, Liu W, Tian Y, Xiong JS, Wang YH, Li RJ, Li HM, Wen J, Mysore KS, Boller T, Xie ZP, Staehelin C. (2018). Role of the Nod factor hydrolase MtNFH1 in regulating Nod factor levels during rhizobial infection and in mature nodules of *Medicago truncatula*. Plant Cell **30,** 397-414.

Dai, W.J., Zeng, Y., Xie, Z.P., and Staehelin, C. (2008). Symbiosis-promoting and deleterious effects of NopT, a novel type 3 effector of *Rhizobium* sp. strain NGR234. J Bacteriol **190,** 5101-5110.

Dowen, R.H., Engel, J.L., Shao, F., Ecker, J.R., and Dixon, J.E. (2009). A family of bacterial cysteine protease type III effectors utilizes acylation-dependent and

-independent strategies to localize to plasma membranes. J Biol Chem **284,**

15867-15879.

Fotiadis, C.T., Dimou, M., Georgakopoulos, D.G., Katinakis, P., and Tampakaki,

A.P. (2012). Functional characterization of NopT1 and NopT2, two type III effectors of

Bradyrhizobium japonicum. FEMS Microbiol Lett **327,** 66-77.

Gan, Y., Zhang, L., Zhang, Z., Dong, S., Li, J., Wang, Y., and Zheng, X. (2009).

The LCB2 subunit of the sphingolip biosynthesis enzyme serine palmitoyltransferase

can function as an attenuator of the hypersensitive response and Bax-induced cell

death. New Phytol **181,** 127-146.

Geurts, R., Fedorova, E., and Bisseling, T. (2005). Nod factor signaling genes and

their function in the early stages of infection. Curr Opin Plant Biol **8,** 346-352.

Goehre, V., Spallek, T., Haeweker, H., Mersmann, S., Mentzel, T., Boller, T., de

Torres, M., Mansfield, J.W., and Robatzek, S. (2008). Plant pattern-recognition

receptor FLS2 is directed for degradation by the bacterial ubiquitin ligase AvrPtoB.

Curr Biol **18,** 1824-1832.

Gysel, K., Laursen, M., Thygesen, M.B., Lironi, D., Bozsoki, Z., Hjuler, C.T.,

Maolanon, N.N., Cheng, J., Bjork, P.K., Vinther, M., Madsen, L.H., Rubsam, H.,

Muszynski, A., Ghodrati, A., Azadi, P., Sullivan, J.T., Ronson, C.W., Jensen, K.J.,

Blaise, M., Radutoiu, S., Stougaard, J., and Andersen, K.R. (2021). Kinetic

proofreading of lipochitooligosaccharides determines signal activation of symbiotic

plant receptors. Proc Natl Acad Sci U S A **118**.e2111031118.

Han, Y., Sun, J., Yang, J., Tan, Z., Luo, J., and Lu, D. (2017). Reconstitution of the

plant ubiquitination cascade in bacteria using a synthetic biology approach. Plant J

91, 766-776.

Heckmann, A.B., Sandal, N., Bek, A.S., Madsen, L.H., Jurkiewicz, A., Nielsen,

M.W., Tirichine, L., and Stougaard, J. (2011). Cytokinin induction of root nodule

primordia in *Lotus japonicus* is regulated by a mechanism operating in the root cortex.

Mol Plant Microbe Interact **24,** 1385-1395.

Jones, J.D.G., and Dangl, J.L. (2006). The plant immune system. Nature **444,**

323-329.

Kambara, K., Ardissone, S., Kobayashi, H., Saad, M.M., Schumpp, O.,

Broughton, W.J., Deakin W.J. (2009). Rhizobia utilize pathogen-like effector

proteins during symbiosis. Mol Microbiol **71,** 92-106.

Kawaharada, Y., Kelly, S., Nielsen, M.W., Hjuler, C.T., Gysel, K., Muszynski, A.,

Carlson, R.W., Thygesen, M.B., Sandal, N., Asmussen, M.H., Vinther, M.,

Andersen, S.U., Krusell, L., Thirup, S., Jensen, K.J., Ronson, C.W., Blaise, M.,

Radutoiu, S., and Stougaard, J. (2015). Receptor-mediated exopolysaccharide

perception controls bacterial infection. Nature **523,** 308-312.

Kato, Y., and Sakamoto, W. (2019). Phosphorylation of the chloroplastic

metalloprotease FtsH in Arabidopsis characterized by Phos-tag SDS-PAGE. Front

Plant Sci, 10, 1080.

Khan, A., Wadood, S.F., Chen, M., Wang, Y., Xie, Z.P., and Staehelin, C. (2022).

Effector-triggered inhibition of nodulation: A rhizobial effector protease targets

soybean kinase GmPBS1-1. Plant Physiol **189,** 2382-2395.

Kim, S.H., Qi, D., Ashfield, T., Helm, M., and Innes, R.W. (2016). Using decoys to

expand the recognition specificity of a plant disease resistance protein. Science **351,**

684-687.

Kimbrel, J.A., Thomas, W.J., Jiang, Y., Creason, A.L., Thireault, C.A., Sachs,

J.L., and Chang, J.H. (2013). Mutualistic co-evolution of type III effector genes in

Sinorhizobium fredii and *Bradyrhizobium japonicum*. PLoS Pathog **9,** e1003204.

Kopan, R., and Ilagan, M.X.G. (2009). The canonical notch signaling pathway:

Unfolding the activation mechanism. Cell **137,** 216-233.

Lei, L., Stevens, D.M., and Coaker, G. (2020). Phosphorylation of the *Pseudomonas*

effector AvrPtoB by *Arabidopsis* SnRK2.8 is required for bacterial virulence. Mol Plant

13, 1513-1522.

Li, D.D., Zhu, Z.K., Deng, X.M., Zou, J.N., Ma, C., Li, C.D., Yin, T., Liu, C.Y., Wang,

J.H., Chen, Q.S., and Xin, D.W. (2023). GmPBS1, a hub gene interacting with

rhizobial type-III effectors NopT and NopP, regulates soybean nodulation. Agronomy

13, 1242.

Li, H., Chen, M., Duan, L., Zhang, T., Cao, Y., and Zhang, Z. (2018). Domain swap

approach reveals the critical roles of different domains of SYMRK in root nodule

symbiosis in *Lotus japonicus*. Front Plant Sci **9,** 697.

Li, L., Kim, P., Yu, L., Cai, G., Chen, S., Alfano, J.R., and Zhou, J.M. (2016).

Activation-dependent destruction of a co-receptor by a *Pseudomonas syringae*

effector dampens plant immunity. Cell Host Microbe **20,** 504-514.

Luo, Y., Liu, D., Jiao, S., Liu, S., Wang, X., Shen, X., and Wei, G. (2020)

Identification of *Robinia pseudoacacia* target proteins responsive to *Mesorhizobium*

amphore CCNWGS0123 effector protein NopT. J Exp Bot. **71,** 7347-7363.

Madsen, E.B., Madsen, L.H., Radutoiu, S., Olbryt, M., Rakwalska, M.,

Szczyglowski, K., Sato, S., Kaneko, T., Tabata, S., Sandal, N., and Stougaard, J.

(2003). A receptor kinase gene of the LysM type is involved in legume perception of

rhizobial signals. Nature **425,** 637-640.

Madsen, E.B., Antolin-Llovera, M., Grossmann, C., Ye, J., Vieweg, S.,

Broghammer, A., Krusell, L., Radutoiu, S., Jensen, O.N., Stougaard, J., and

Parniske, M. (2011). Autophosphorylation is essential for the *in vivo* function of the

Lotus japonicus Nod factor receptor 1 and receptor-mediated signalling in cooperation

with Nod factor receptor 5. Plant J **65,** 404-417.

Moling, S., Pietraszewska-Bogiel, A., Postma, M., Fedorova, E., Hink, M.A.,

Limpens, E., Gadella, T.W., and Bisseling, T. (2014). Nod factor receptors form

heteromeric complexes and are essential for intracellular infection in *Medicago*

nodules. Plant Cell **26,** 4188-4199.

Nakagawa, T., Suzuki, T., Murata, S., Nakamura, S., Hino, T., Maeo, K., Tabata,

R., Kawai, T., Tanaka, K., Niwa, Y., Watanabe, Y., Nakamura, K., Kimura, T., and

Ishiguro, S. (2007). Improved Gateway binary vectors: high-performance vectors for

creation of fusion constructs in transgenic analysis of plants. Biosci Biotechnol

Biochem **71,** 2095-2100.

Ntoukakis, V., Mucyn, T.S., Gimenez-Ibanez, S., Chapman, H.C., Gutierrez, J.R.,

Balmuth, A.L., Jones, A.M., and Rathjen, J.P. (2009). Host inhibition of a bacterial

virulence effector triggers immunity to infection. Science **324,** 784-787.

Radutoiu S, Madsen LH, Madsen EB, Felle HH, Umehara Y, Grønlund M, Sato S,

Nakamura Y, Tabata S, Sandal N, Stougaard J. (2003). Plant recognition of

symbiotic bacteria requires two LysM receptor-like kinases. Nature **425,** 585-592.

Rae, A.E., Rolland, V., White, R.G., and Mathesius, U. (2021). New methods for

confocal imaging of infection threads in crop and model legumes. Plant Methods **17**,

24.

Ried, MK, Antolín-Llovera, M, and Parniske, M. (2014). Spontaneous symbiotic

reprogramming of plant roots triggered by receptor-like kinases. Elife **3,** e03891.

Schauser, L., Roussis, A., Stiller, J., and Stougaard J. (1999) A plant regulator

controlling development of symbiotic root nodules. Nature **402,** 191-195.

Schmidt, G. (2011). *Yersinia enterocolitica* outer protein T (YopT). Eur J Cell Biol **90,**

955-958.

Schubert, K.A., Xu, Y., Shao, F., and Auerbuch, V. (2020). The type III secretion

system as a tool for studying cytosolic innate immune surveillance. Ann Rev

Microbiol, Vol 74, 2020 **74,** 221-245.

Shao, F., Merritt, P.M., Bao, Z., Innes, R.W., and Dixon, J.E. (2002). A *Yersinia*

effector and a *Pseudomonas* avirulence protein define a family of cysteine proteases

functioning in bacterial pathogenesis. Cell **109,** 575-588.

Shao, F., Vacratsis, P.O., Bao, Z., Bowers, K.E., Fierke, C.A., and Dixon, J.E.

(2003a). Biochemical characterization of the *Yersinia* YopT protease: cleavage site

and recognition elements in Rho GTPases. Proc Natl Acad Sci U S A **100,** 904-909.

Shao, F., Golstein, C., Ade, J., Stoutemyer, M., Dixon, J.E., and Innes, R.W.

(2003b). Cleavage of *Arabidopsis* PBS1 by a bacterial type III effector. Science **301,**

1230-1233.

Sugawara, M., Takahashi, S., Umehara, Y., Iwano, H., Tsurumaru, H., Odake, H.,

Suzuki, Y., Kondo, H., Konno, Y., Yamakawa, T., Sato, S., Mitsui, H., and

Minamisawa, K. (2018). Variation in bradyrhizobial NopP effector determines

symbiotic incompatibility with soybeans via effector-triggered immunity. Nature

Commun **9**.3139.

Sun, Y., Li, L., Macho, A.P., Han, Z., Hu, Z., Zipfel, C., Zhou, J.-M., and Chai, J.

(2013). Structural basis for flg22-induced activation of the *Arabidopsis* FLS2-BAK1

immune complex. Science **342,** 624-628.

Tang, D.Z., Wang, G.X., and Zhou, J.M. (2017). Receptor kinases in plant-pathogen

interactions: More than pattern recognition. Plant Cell **29,** 618-637.

Teulet, A., Busset, N., Fardoux, J., Gully, D., Chaintreuil, C., Cartieaux, F.,

Jauneau, A., Comorge, V., Okazaki, S., Kaneko, T., Gressent, F., Nouwen, N.,

Arrighi, J.F., Koebnik, R., Mergaert, P., Deslandes, L., and Giraud, E. (2019). The

rhizobial type III effector ErnA confers the ability to form nodules in legumes. Proc Natl

Acad Sci U S A **116,** 21758-21768.

Tirichine, L., Imaizumi-Anraku, H., Yoshida, S., Murakami, Y., Madsen, L.H.,

Miwa, H., Nakagawa, T., Sandal, N., Albrektsen, A.S., Kawaguchi, M., Downie, A.,

Sato, S., Tabata, S., Kouchi, H., Parniske, M., Kawasaki, S., and Stougaard, J.

(2006). Deregulation of a Ca2+/calmodulin-dependent kinase leads to spontaneous

nodule development. Nature **441,** 1153-1156.

Vleeshouwers, V.G., Driesprong, J.D., Kamphuis, L.G., Torto-Alalibo, T., Van't

Slot, K.A., Govers, F., Visser, R.G., Jacobsen, E., and Kamoun, S. (2006).

Agroinfection-based high-throughput screening reveals specific recognition of INF

elicitins in *Solanum*. Mol Plant Pathol **7,** 499-510.

Wang, D., Yang, S., Tang, F., and Zhu, H. (2012). Symbiosis specificity in the

legume: rhizobial mutualism. Cell Microbiol **14,** 334-342.

Wang, Q., Han, C., Ferreira, A.O., Yu, X., Ye, W., Tripathy, S., Kale, S.D., Gu, B.,

Sheng, Y., Sui, Y., Wang, X., Zhang, Z., Cheng, B., Dong, S., Shan, W., Zheng, X.,

Dou, D., Tyler, B.M., and Wang, Y. (2011). Transcriptional programming and

functional interactions within the *Phytophthora sojae* RXLR effector repertoire. Plant

Cell **23,** 2064-2086.

Xiang, T., Zong, N., Zou, Y., Wu, Y., Zhang, J., Xing, W., Li, Y., Tang, X., Zhu, L.,

Chai, J., and Zhou, J.M. (2008). *Pseudomonas syringae* effector AvrPto blocks

innate immunity by targeting receptor kinases. Curr Biol **18,** 74-80.

Xin, D.W., Liao, S., Xie, Z.P., Hann, D.R., Steinle, L., Boller, T., Staehelin, C. (2012). Functional analysis of NopM, a novel E3 ubiquitin ligase (NEL) domain effector of *Rhizobium* sp. strain NGR234. PLoS Pathog **8(5),** e1002707.

Xu, N., Luo, X., Wu, W., Xing, Y., Liang, Y., Liu, Y., Zou, H., Wei, H.L., and Liu, J. (2020). A plant lectin receptor-like kinase phosphorylates the bacterial effector AvrPtoB to dampen Iits virulence in *Arabidopsis*. Mol Plant **13,** 1499-1512.

Yang, J., Lan, L., Jin, Y., Yu, N., Wang, D., and Wang, E. (2022). Mechanisms underlying legume-rhizobium symbioses. J Integr Plant Biol **64,** 244-267.

Zhang, B., Wang, M., Sun, Y., Zhao, P., Liu, C., Qing, K., Hu, X., Zhong, Z., Cheng, J., Wang, H., Peng, Y., Shi, J., Zhuang, L., Du, S., He, M., Wu, H., Liu, M., Chen, S., Wang, H., Chen, X., Fan, W., Tian, K., Wang, Y., Chen, Q., Wang, S., Dong, F., Yang, C., Zhang, M., Song, Q., Li, Y., and Wang, X. (2021). *Glycine max* NNL1 restricts symbiotic compatibility with widely distributed *Bradyrhizobia vi*a root hair infection. Nat Plants **7,** 73-86.

Zhang, J., Li, W., Xiang, T., Liu, Z., Laluk, K., Ding, X., Zou, Y., Gao, M., Zhang, X., Chen, S., Mengiste, T., Zhang, Y., and Zhou, J.M. (2010). Receptor-like cytoplasmic kinases integrate signaling from multiple plant immune receptors and are targeted by a *Pseudomonas syringae* effector. Cell Host Microbe **7,** 290-301.

Zhou, J.G., Wang, P., Claus, L.A.N., Savatin, D.V., Xu, G.Y., Wu, S.J., Meng, X.Z., Russinova, E., He, P., and Shan, L.B. (2019). Proteolytic processing of SERK3/BAK1 regulates plant immunity, development, and cell death. Plant Physiol **180,** 543-558.

Zipfel, C., and Oldroyd, G.E.D. (2017). Plant signalling in symbiosis and immunity. Nature **543,** 328-336.

Zipfel, C., Robatzek, S., Navarro, L., Oakeley, E.J., Jones, J.D.G., Felix, G., and Boller, T. (2004). Bacterial disease resistance in through flagellin perception. Nature **428,** 764-767.

Figures

1060

1061

1062

1063

Figure 1. NopT specifically suppresses NFR1/NFR5-triggered cell death

in *N. benthamiana*. *Agrobacterium* strains harboring plasmid DNA encoding

NopT or the empty vector (EV) were infiltrated into *N. benthamiana* leaves. At 12

hpi, a second infiltration was performed with an *Agrobacterium* strain containing a

plasmid with *NFR1*/*NFR5* genes. Dashed red lines indicate leaf discs where different

proteins were expressed. (A, B) NopT (A), but not NopT^{C93S} (B), a protease-inactive

version of NopT, suppressed NFR1NFR5-induced cell death. (C) Expression of

NopM-induced cell death in *N. benthamiana*. (D) Expression of NopT could not

suppress the cell death response triggered by expression of Avr3a/R3a, NopM, BAX,

INF1 or AtCERK1. Numbers in (D) represent the number of leaf discs showing cell

death and total leaf discs tested. The pictures shown in this figure are representative of

at least three independent biological replicates.

1076

1077

1078

1079

1080

Figure 2. NopT interacts with NFR1 and NFR5. Interactions between NopT and

NFR1 or NFR5 were detected using BiFC (Split-YFP) (A, B), Split-LUC

complementation (C) and co-IP (D) assays in *N. benthamiana* leaves. (A, B) For

BIFC analysis, nYFP and cYFP tags were fused at the C-terminus of NopT, NFR1,

NFR5, and the flagellin receptor FLS2 (negative control). YFP fluorescence signals

represent protein-protein interactions. Scale bar=25 µm. (C) For the Split-LUC

complementation assay, the nLUC and cLUC tags were fused at the C-terminus of

NopT, NFR1, NFR5 and FLS2 (negative control). Luminescence signals represent

protein-protein interactions. (D) For the co-IP assay, HA-tagged NFR1 or NFR5 and

FLAG-tagged NopT were expressed in *N. benthamiana* cells followed by

immunoprecipitation using an anti-FLAG antibody. Immunoblot analysis was

performed using anti-HA and anti-FLAG antibodies (NopT^C denotes the truncated

version of NopT after autocleavage). The images and immunoblots shown in this

figure are representative of three biological replicates.

1095

1096

Figure 3. NopT proteolyzes NFR5 at its JM domain. Proteins with indicated tags

were expressed in *N. benthamiana* (A), *L. japonicus* (B) or *E. coli* (C-G) and

detected by immunoblotting. (A) NopT but not NopT^{C93S} cleaves NFR5-GFP

protein expressed in *N. benthamiana* cells by releasing NFR5^{CD}-GFP (NopT^C

denotes autocleaved NopT). (B) NopT but not NopT^{C93S} cleaves NFR5-Myc

expressed in hairy roots of *L. japonicus* by releasing NFR5^{CD}-Myc (marked by

an asterisk). (C) Analysis of the CDs of NFR1 and NFR5 co-expressed with

NopT or NopT^{C93S} in *E. coli*. Cleavage of NFR5^{CD} was observed for NopT but

not NopT^{C93S}, while NFR1^{CD} was not proteolyzed by NopT. In the presence of

NFR1^{CD}, a slower migrating band was observed, possibly representing

phosphorylated NopT (NopT^m). (D) A repeat experiment confirmed that NopT

is able to cleave NFR5^{CD} (the asterisk indicates the HA-tagged NFR5 cleavage

product containing the kinase domain and a C-terminal tail region). (E) NopT

cleaves His-SUMO-NFR5^{JM}-GFP (His-SUMO and GFP linked by the JM

domain of NFR5) *in vitro*. F.T. indicates proteins in flow through samples after

purification with Ni-beads. (F) NopT expressed in *E. coli* was unable to cleave

co-expressed NFR5^{CD-5m}-HA, a modified version of NFR5^{CD}-HA in which five

amino acids of the JM were substituted by other residues (S283Y, G294Q,

Y303S, A310I, T311Y). (G) NopT expressed in *E. coli* was unable to cleave

co-expressed *M. truncatula* NFP^{CD}-HA and the NFR5^{JM}-NFP^{KC}-HA fusion

protein, while NFP^{JM}-NFR5^{KC} was proteolyzed (KC stands for the kinase

domain and a C-terminal tail region).

1119

1120

1121

Figure 4. Phosphorylation of NopT by NFR1 suppresses its proteolytic activity.

(A) *In vivo* phosphorylation assay with proteins expressed in *L. japonicus* roots

(wild-type and *nfr1-1* mutant plants) using Zn^{2+}-Phos-tag SDS-PAGE.

Phosphorylation of NopT^{C93S} was induced by inoculation with rhizobia

(*Mesorhizobium loti* MAFF303099) and was largely dependent on *NFR1*. (B)

The relative protein amount of each lane, as shown in (A), was quantified with

ImageJ software (three biological replicates). The value of the control band in

each gel was set to 1 for comparison. Values are means ± SEM. (C) NFR1^{CD}

but not NFR1$^{CD-K351E}$ phosphorylates NopT in *E. coli*. The phosphorylated

full-length form of NopT could be dephosphorylated by CIAP (gel band shift).

Abbreviations: NFR1^{CD-P}, autophosphorylated NFR1; NopT^P, phosphorylated

NopT; NopT, non-phosphorylated NopT; NopT^C, autocleaved NopT. (D) The

phoshorylation sites of NopT identified by LC-MS were either substituted to

alanine (A) or aspartate (D). The indicated NopT variants were subsequently

tested for autocleavage and NFR5^{CD} proteolysis in *E. coli*. Wild-type NopT

(WT) and protease-dead NopT^{C93S} were included into the analysis.

1138

1139

Figure 5. NopT regulates rhizobial infection in *L. japonicus*. (A) Analysis of

rhizobial infection in *L. japonicus* roots inoculated with *GUS*-labelled *S. fredii*

NGR234 (wild-type; WT) or a *nopT* knockout mutant (NGR234Δ*nopT*; abbreviated

as Δ*nopT* in other panels). Scale bar=100 µm. (B) GUS staining pictures showing

roots of *L. japonicus* plants expressing *GUS* expression under control of the *NIN*

promoter (*pNIN*:*GUS*). Plants were inoculated with NGR234 or NGR234Δ*nopT* and

analyzed at 7 dpi. Scale bar=1 mm. (C) Infection data (ITs, infection threads) for roots

shown in (A) at 7 dpi (n=10, Student's *t*-test; * indicates P < 0.01). (D) Quantification

of GUS staining sites for roots shown in (B) (n=10, Student's *t*-test, P < 0.01). (E)

Nodule primordia formation in *L. japonicus* wild-type roots inoculated with NGR234

(WT), NGR234Δ*nopT* or NGR234 over-expressing *nopT* (*pT7*:NopT). Roots were

analyzed at 14 dpi (n=16, Student's *t*-test: P < 0.01). (F) Infection data for wild-type

roots inoculated with NGR234 (WT), NGR234Δ*nopT* and NGR234Δ*nopT* expressing

indicated NopT variants at 7 dpi (n=5, Student's *t*-test: P < 0.01). (G) Analysis of

rhizobial infection in hairy roots of *L. japonicus* (wild-type) expressing GFP (EV,

empty vector control) or NopT. Plants were inoculated with DsRed-labeled *M. loti*

MAFF303099 and analyzed at 5 dpi (n=8, Student's t–test: P < 0.01). (H)

Expression of NFR5 and NFR5-NFP^{KC} in hairy roots of *nfr5-3* mutant plants.

Plants were inoculated with GFP-labelled NGR234 (WT) or NGR234Δ*nopT*

and analyzed at 8 dpi . Roots expressing NFR5-NFP^{KC} showed high numbers of

infection foci for both strains whereas significant differences were observed for

NFR5 expressing roots (n=10, Student's t-test: P < 0.01).

1162

1163

1164

Figure 6. *S. fredii* **NopT proteins cleave NFR5 and working model for NopT of**

NGR234. (A) NopT of *S. fredii* NGR234 and homologs from other rhizobial

strains were co-expressed with NFR5^{CD} in *E. coli* (NopT$_{234}$, NopT of NGR234;

NopT1$_{110}$ and NopT2$_{110}$, NopT proteins of *B. diazoefficiens* USDA110;

NopT$_{257}$, NopT of *S. fredii* USDA257; NopT$_{103}$, NopT of *S. fredii* HH103).

Immunoblot analysis indicated NFR5 cleavage by NopT proteins from *S. fredii*

strains. (B) Expression of NopT$_{257}$ in *N. benthamiana* could not inhibit the cell

death triggered by co-expressed NFR1 and NFR5. (C) *L. japonicus* Gifu

pNIN:*GUS* plants were inoculated with *S. fredii* NGR234, NGR234Δ*nopT* and

NGR234Δ*nopT* expressing NopT$_{257}$ (Δ*nopT*+NopT$_{257}$). Roots were subjected to

GUS staining at 7 dpi. Scale bar=2.5 mm. (D) Quantitative analysis of GUS-stained

roots shown in panel C (n=19, Student's *t*–test: P < 0.01). (E) A proposed model

for NopT of NGR234 interacting with NFRs. NopT and NopT^{C} (autocleaved

NopT) proteolytically cleave NFR5 at the JM domain to release the intracellular

domain of NFR5 (cleaved NFR5). NFR1 phosphorylates full-length NopT to

block its proteinase activity. NopT^{C} cannot be phosphorylated by NFR1.

1181

1182

1183

1184

1185

1186

1187

1188

1189

1190

1191

1192

1193

1194

1195

1196

1197

 Supplemental Figures

Supplemental Fig. 1. Analyses of the functions of all 15 effectors from *S. fredii* NGR234 in preventing NFR1- and NFR5-induced cell death in *N. benthamiana* leaves.

15 genes from *S. fredii* NGR234 encoding effector proteins were amplified and cloned into binary vector for expression of Strep-tagged proteins in *N. benthamiana*. NFR1 and NFR5 from *L. japonicus* were C-terminally tagged with HA and Myc, respectively, and cloned into binary vectors for expression in *N. benthamiana*. All plasmids were electroporated into *Agrobacterium tumefaciens* EH105.

(A) *Agrobacterium* strain harboring each effector gene or control empty vector (EV) were infiltrated into *N. benthamiana*. 12 hour post infiltration (hpi), agrobacterium strains harboring NFR1 and NFR5 were mixed and hand-infiltrated in the same leaf discs as shown in (B). 48 hpi, cell death phenotypes were evaluated (C). At least three replicates in the same leaf were

performed to test the function of each effector in suppressing cell

death-triggered by NFR1 and LjNFR5. (D) Results of all 15 effector proteins in

suppressing programmed cell death by overexpression of NFR1 and NFR5.

(E) Abundance of NFR1, NFR5, and NopT proteins, as measured by

immunoblot with specific antibodies. Leaf discs expressing NFR1, NFR5, and

NopT or NFR1 and NFR5 and with EV were used for immunoblot analyses.

Actin was used as loading control. (F) Another example of suppression of

programmed cell death by NopT in the leaf discs coexpressing NFR1 and

NFR5. Rep1, Rep2, and Rep3 represent three technical repeats in the same

leave.

1224

1225

1226

1227

1228
1229

1230

Supplemental Fig. 2. Split–LUC assay testing the interactions between different NopT mutants and NFRs.

(A)The interaction between NopT with mutated acylation site and NFR1/NFR5. NopT^{AAA} (G50A, C51A, and C52A). (B) The protein expressing of indicating genes in the leave of (A). (C) Nop^{C93S} interact with NFR1 and NFR5. (D) The interaction between NopT$^{\Delta N50}$ and NFR5. NopT$^{\Delta N50}$ represents a truncated NopT with deletion of about 50 a.a. from its N–terminus.

1248

Supplemental Fig. 3. NopT cleaves NFR5 not not NFR1.

Proteins with indicated tags were expressed in either *N. benthamiana* or *E. coli* and

detected by immunoblotting. (A) NopT did not cleave NFR1-GFP expressed in *N.*

benthamiana leaves. (B) NopT expressed in *E. coli* did not cleave co-expressed

SUMO-NFR5^{KC}-HA. KC: Kinase domain and C-terminal domain

1254

1255

1256

1257

1258

1259

1260

1261

1262

1263 **Supplemental Fig. 4. NopT interacts with the JM domain of NFR5.**

1264 NopT interacts with NFR5^{JM} in *in vitro* pull down assay. IB: immunoblotting, IP:

1265 immunoprecipitation. His-SUMO-NFR5^{JM}-GFP: a recombinant protein containing

1266 His tag, SUMO tag, the JM domain of NFR5, and GFP.

1267

1268

1269

1270

1271

1272

1273

1274

1275

Supplemental Fig. 5. Conserved domains and residues of NFR5 and related proteins.

(A) Domain and motif annotation of NFR5. LysM: Lysin-Motif; TM: transmembrane domain; JM: juxtamembrane domain; KD: kinase domain; CT: C-terminal domain.

(B) Alignment of amino acid residues of AtLYK5, LjLYS11, NFR5 and MtNFP close to the RRKK motif in the JM domain of NFR5. Five highly conserved residues marked by yellow coloration (S283, G294, Y304, A310, T311 in NFR5) and the brown box indicates the start region of kinase domain. The green frame delineates the RRKK motif in NFR5 and the red-framed residues are similar to the NopT autocleavage region.

1286

1287

1288

1289

Supplemental Fig. 6. NopT cleaves NFR5 at the juxtamembrane domain.

(A) Cleavage assay using two mutant versions of NFR5^{CD} in the presence of NopT or

AvrPphB. NFR5$^{CD\ 288-294A}$ represents a mutant NFR5^{CD} with residues 288 to 294

replaced with seven alanines; NFR5$^{CD\ 288-294PphB}$ represents a mutant NFR5^{CD} with

residues 288 to 294 replaced with AvrPphB recognition sites. (B) Autocleavage assay,

using different NopT variants with single amino acid mutations. H205A and D220A

indicate His-205 and Asp-220 replaced with alanine, respectively. (C-F) NopT

cleavage assay using 19 variants of NFR5^{CD} with three adjacent amino acids replaced

with three alanines. The numbers indicate the location of the three residues replaced

with alanines. (G) NopT cleavage assay using seven truncated variants of NFR5^{CD}

with 10 adjacent amino acids deleted. The numbers indicate the location of the 10

deleted residues.

1302

1303

Supplemental Fig. 7. Mass spectrometry analysis of cleavage site of NFR5 by NopT.

(A) The sequence of NFR5 juxtamembrane domain (286-320 a.a.). (B) Mass spectrometry analysis of peptides of NFR5[CD] after proteolysis by NopT. Blue lines indicated the peptides characterized by Mass spectrometry. Red frame delineated cleavage site of NFR5 identified using Mass spectrometry.

1310

Supplemental Fig. 8. NopT cleaves the NFR5 homolog proteins at the juxtamembrane domain.

NopT but not NopT^{C93S} could cleave the CDs of AtLYK5 and LjLYS11 expressed in *E. coli.* NopT could cleave both AtLYK5^{JM}–NFR5^{KC}, LjLYS11^{JM}–NFR5^{KC} and NFR1^{JM}–NFR5^{KC}, three recombinant proteins expressed in *E. coli.* AtLYK5^{JM}–NFR5^{KC}, LjLYS11^{JM}–NFR5^{KC}, and NFR1^{JM}–NFR5^{KC} recombinant NFR5 proteins with the the JM was replaced with the JMs from AtLYK5, LjLYS11 and LjNFR1, respectively. KC: Kinase domain and C-terminal domain as ilustrated in Fig. S5A.

1327

1328 **Supplemental Fig. 9. NopT cleaves the JM of MtNFP.**

1329 NopT cleaves a recombinant protein where His–tagged SUMO and GFP is

1330 bridged with the JM domain of NFP. F.T. represents flow through sample after

1331 Ni–beads purification. IB: immunoblotting.

1332

1333

1334

1335

1336

1337

1338

1339

1340

1341

1342

1343 **Supplemental Fig. 10. NopT cleaves recombinant proteins**

1344 **NFR5$^{268-445}$-NFP$^{458-595}$ and NFP$^{270-457}$-NFR$5^{456-595}$.**

1345 NopT but not NopT^{C93S} could cleave recombinant proteins NFR5$^{268-445}$-NFP$^{458-595}$ and

1346 NFP$^{270-457}$-NFR$5^{456-595}$ detected by immunobltting (IB).

1347

1348

1349

1350

1351

1352

1353

1354

1355

1356

1357

1358 **Supplemental Fig. 11. NFR1^{CD} phosphorylates NFR5^{CD} *in vitro*.**

1359 The cytoplasmic domains (CDs) of NFR1, NFR5, and kinase–dead

1360 NFR1$^{CD-K351E}$ were used for kinase assays in *E. coli*. Proteins were detected by

1361 immunoblotting using anti–HA and anti–Myc antibodies. Asterisk indicated the

1362 band retardation on the gel representing the phosphorylated NFR5^{CD} proteins.

1363

1364

1365

1366

1367

1368

1369

1370

1371

1372

1373

1374

Supplemental Fig. 12. Rhizobial infection in the mutant versions of NFR5.

1375

NFR5[5m] failed to form rhizobial infection (n=7, Student's t–test: P < 0.01). The

1376

transgenic roots expressing NFR5 or NFR5[5m] in the *nfr5-3* mutant under the

1377

control of native promoter were inoculated with GFP-labelled wild type

1378

NGR234 and NGR234Δ*nopT* respectively. The data was collected 9 days post

1379

inoculation. NFR5[5m] represents a mutant version of NFR5 with S283Y, G294Q,

1380

Y304S, A310I, T311Y point mutation.

1381

0.50

 Supplemental Fig. 13. Phylogenetic tree based on the amino acid sequence

1384 **of NopT homologs from different bacterial species.**

1385

1386

1387

1388

1389

1390

1391

1392

1393

1394

1395

1396

1397

1398

1399

Supplemental Fig. 14. The cleavage efficiency of NFR5 in different system.

The relative cleavage efficiency for experiment displayed in Fig. 3A (*N. benthamiana*), Fig. 3B (*L. japonicus*) and Fig. 3D (*E. coli*) quantified from three biological replicates using ImageJ software. Values are means ± SE (n=3, Student t-test, letters represent significant differences, p<0.05).

1415

Supplemental Fig. 15. Transient expression of NopT triggers cell death in

Arabidopsis thaliana **and** *Nicotiana tabacum***.**

(A) *Pseudomonas syringae* pv tomato DC3000 harboring control plasmids (left) or

expressing NopT (right) were infiltrated into Arabidopsis leaves. Pictures were

taken three days post inoculation. (B) *Agrobacterium* strains harboring *NopT*,

NopT with an N-terminal 50–amino acid deletion, *NopT*^{C93S}, or *NopT*^{C93S} with an

N-terminal 50–amino acid deletion were infiltrated into *N. tabacum* leaves.

Pictures were taken three days post inoculation.

Supplemental Table 1.   Phosphopeptides identified by liquid chromatography–mass spectrometry. An *in vitro* kinase assay was performed using the CD of NFR1 and NopT. NopT was separated by SDS-PAGE, and the digested gel slices representing the phosphorylated NopT were used for analysis. The deduced phosphopeptides are listed.

Sequence	Modifications	SITE	PSMs_NFR1$^{CD\text{-}KE}$+NopT	PSMs_NFR1+NopT
ETLANVELRTK	1xPhospho [T10]	37T	-	1
ETLANVELR	1xPhospho [T2]	29T	-	1
ETLANVELRTKSPSAECPDK	1xCarbamidomethyl [C17]; 1xPhospho [T10]	37T	-	1
TKSPSAECPDK	1xCarbamidomethyl [C8]; 1xPhospho [S5]	41S	-	1
SPSAECPDK	1xCarbamidomethyl [C6]; 1xPhospho [S3]	41S		1
SPSAECPDKMGCCASKPQASDPNNPSTSSPARPSTSLFR	3xCarbamidomethyl [C6; C12; C13]; 1xPhospho [S/T]; 1xOxidation [M10]		-	4
MGCCASKPQASDPNNPSTSSPARPSTSLFR	2xCarbamidomethyl [C3; C4]; 1xPhospho [T/S]; 1xOxidation [M1]		-	11
MGCCASKPQASDPNNPSTSSPARPSTSLFR	2xCarbamidomethyl [C3; C4]; 1xPhospho [T/S]		-	9
MGCCASKPQASDPNNPSTSSPARPSTSLFR	2xCarbamidomethyl [C3; C4]; 2xPhospho [T/S]	65S 66T	-	4
MGCCASKPQASDPNNPSTSSPARPSTSLFR	2xCarbamidomethyl [C3; C4]; 1xOxidation [M1]		4	2
MGCCASKPQASDPNNPSTSSPARPSTSLFR	2xCarbamidomethyl [C3; C4]; 2xPhospho [T/S]; 1xOxidation [M1]		-	4
TAELAQANADGICVGLTAEWLR	1xCarbamidomethyl [C13]; 1xPhospho [T17]	97T	-	10
NLNSHPSIR	1xPhospho [S7]	109S	-	2
NLNSHPSIRMEALVPGSQR	1xPhospho [S7]; 1xOxidation [M10]	119S	-	1
MEALVPGSQR	1xPhospho [S8]	119S	-	1
MEALVPGSQR	1xPhospho [S8]; 1xOxidation [M1]	119S	-	1
RQGAGPSEADFAAQNTMLQK	1xPhospho [T/S]; 1xOxidation [M17]		-	3
RQGAGPSEADFAAQNTMLQK	1xPhospho [T16]	156T	-	2
QGAGPSEADFAAQNTMLQK	1xPhospho [T/S]; 1xOxidation [M16]		-	3
QGAGPSEADFAAQNTMLQK	1xPhospho [T15]	156T	-	2
SLANRYSNPNR	1xPhospho [S7]	245S	-	2
QHLTTVTTQK	1xPhospho [T8]	257T	-	1

 Supplemental Table 2. Primers used in this study.

Used for	Target genes	Names	Sequences (from 5' to 3')
Overexpression in Nicotiana benthamiana	NopT or NopT^{C93S}	p5X-NopT-F	agaacacggggggactctagaatgcacagtcccatcag
		p5X-NopT-R	ccgctgttatcggtacctgtcatcttttgggtggtcac
	NopT$^{\Delta N50}$	p5X-NopTdeN50-F	agaacacggggggactctagaatgtgctgcgccagcaag
	NFR1	p5X-NFR1-F	agaacacggggggactctagaatgaagctaaaaactggtctact
		p5X-NFR1-R	ccgctgttatcggtacctcttctcacagac
	NFR5	p5X-NFR5-F	agaacacggggggactctagaatggctgtcttctttcttacc
		p5X-NFR5-R	ccgctgttatcggtaccacgtgcagtaat
		p5X-NFR5gfp-R	tcccgggagcggtaccacgtgcagtaatggaagt
	AvrPphB	p5X-AvrPphB-F	agaacacggggggactctagaatgaaaataggtacgcaggcc
		p5X-AvrPphB-R	ccgctgttatcggtacccgaaactctaaactcgtttacgc
templates for protein expression	NopT$_{USDA257}$	p5X-257T-F	agaacacggggggactctagatgctgcgtgatccgaacaat
		p5X-257T-R	ccgctgttatcggtacctgtcacctttgggtggtcaccg
	NopT1$_{USDA110}$	p5X-110T1-F	agaacacggggggactctagaatgtatgatcgaatcggtgg
		p5X-110T1-R	ccgctgttatcggtaccctgcatcctttgcgtcg
	NopT2$_{USDA110}$	p5X-110T2-F	agaacacggggggactctagaatgtataatcgagtcgatggc
		p5X-110T2-R	ccgctgttatcggtaccccgatgaggttccgc
Protein Expression in E.coli	NopT or NopT^{C93S}	28a NopT-F	tgggtcgcggatccgaattcatgcacagtcccatcag
		28a FLAG-R	tcgagtgcggccgcaagcttcttgtcatcgtcatccttgtag
		28a Strep-R	tcgagtgcggccgcaagctttttttcaaattgaggatgagaccatcc
	AvrPphB	ET28a-pphb-F	tgggtcgcggatccgaattcaaaataggtacgcaggccac
	NopT$_{USDA257}$	ET28a-257T-F	tgggtcgcggatccgaattcctgcgtgatccgaacaa
	NopT1$_{USDA110}$	ET28a-110T1-F	tgggtcgcggatccgaattctatgatcgaatcggtggctc
	NopT2$_{USDA110}$	ET28a-110T2-F	tgggtcgcggatccgaattctataatcgagtcgatggcgaatac
	NFR1^{CD}	duet NFR1-F	accacagccaggatccgagataccagaagaaggaagaagagaaagc
		duet Sac1nost-R	ggcgcgccgagctcgcgatcggggaaattcgagct
	NFR5^{CD}	ACSH-NFR5-F	agaagcgcggatccgaattctatgtatactgccgcagaaagaag
		ACSH-NFR5-R	cgtatgggtagcttccaagcttacgtgcagtaatggaagtc
	NFR5^{KD}	SUMO 5KD-F	ccgcgaacagattggtaaggttggggaatcagtgtac
		SUMO HA-R	gctcgaattcggatcctttcacgcatagtcaggaacatcg
	NFR5^{JM}	SUMO 5JM-F	ccgcgaacagattggttatgtatactgccgcagaaagaag
		SUMO 5JM-R	gctcgaattcggatcctttcacgcatagtcaggaacatcg
	sumo-NFR5^{KD}-HA	duet sumo-F	catcaccacagccaggatgggtccctgcaggac
		duet HA-R	cattatgcggccgcaagcttttacgcatagtcaggaacatcg
	sumo-NFR5^{JM}-GFP	duet sumo-F	catcaccacagccaggatgggtccctgcaggac
		duet GFP-R	atgcggccgcaagctttttacttgtacaactcatccatacc

	AtLYK5	ACSH-LYK5-F	gcgcggatccgaattctacaaacgaaggtctaagaag
		ACSH-LYK5-R	ggtagcttccaagcttgttgccaagagagccg
	LjLYS11	ACSH-LYS11-F	agaagcgcggatccgaattcacttgtctgaggaagagaaag
		ACSH-LYS11-R	cgtatgggtagcttccaagcttacgagctgctatcaaagtt
	LYK5^{JM} NFR5^{KD}	LYK5-NFR5-F	cgaaaacagaaaggttggggaatcagtg
		NFR5 LYK5-R	ccccaacctttctgttttcgtcgctgaaatt
	LYS11^{JM} NFR5KD	LYS11 NFR5-F	gagcagtgtaaggttggggaatcag
		NFR5 LYS11R	ccccaaccttacactgctcattgagg
	NFP^{JM}	ACSUG NFPJ-F	gtggaggcgctagaggatcctattgtctcaaaatgaagagattgaatagaag
		ACSUG NFPJ-R	cttctcccttagagagctcacaattgtcactcagattcattg
	NFR5^{JM}-NFP^{KD}	5JM-PKD-F	gatgagtgcaagattggtgaatcagtttacaaag
		5JM-PKD-R	accaatcttgcactcatcgct
	NFP^{JM}-NFR5^{KD}	PJM-5KD-F	gacaattgtaaggttggggaatcagtg
		PJM-5KD-R	ccccaaccttacaattgtcactcagattcattg
	NFR5$^{268-445}$-NFP^{4}$^{58-595}$	51/2KD-P1/2KD-F	ccatggccagaacttcaaccaactcaat
		51/2KD-P1/2KD-R	gaagttctggccatggcgaa
	NFP$^{270-457}$-NFR^{54}$^{56-595}$	P1/2KD-51/2KD-F	ggatggctagaacttcgaccaac
		P1/2KD-51/2KD-R	cgaagttctagccatcccgaa
site mutantions construction	NopT	T93S-F	gaatctccgtcggcttgactgc
		T93S-F	gacggagattccatctgcgttc
		97D-F	ggcttggatgcggagtggctgcg
		97D-R	ctccgcatccaagccgacgcag
		97A-F	ggcttggctgcggagtggctgc
		97A-R	ctccgcagccaagccgacg
		109D-F	catccggatatccgaatggaggccctag
		109D-R	ttcggatatccggatgactgttgaggttacgc
		109A-F	catccggcaatccgaatggag
		109A-R	ttcggattgccggatgactgttg
		119D-R	cccggagatcaaaggcacgcctcagc
		119A-F	gcctttgatctccgggtactagggcc
		119A-F	cccggagcgcaaaggcac
		119A-R	ctttgcgctccgggtactagg
		156D-F	gcaaaacgatatgttgcagaaagcaggc
		156D-R	gcaacatatcgttttgcgccgcgaagtcggcc
		156A-F	gcaaaacgctatgttgcagaaagcag
		156A-R	ctgcaacatagcgttttgcgccgcgaagtc
		H205A-F	ggcggcaacaccgttgcgacc
		H205A-R	caacggtgttgccgccgccctcagc
		D220A-F	ctcttcgctcctaatttcggcgaatttac
		D220A-R	gccgaaattaggagcgaagagcgtggtgtttc

		245D-F	cgctacgacaatccaaaccggcag
		245D-R	ttggattgtcgtagcgattggctaggct
		257 mutants	changing the sequence of r primer at noptt257when cloned into vectors
	NFR1	K351E-F	gcaattgagaagatggatgtacaagcatc
		K351E-R	catccatcttctcaattgctgttttcttgcc
	NFR5	269-271A-F	ctatgcagctgcacgcagaaagaaggctctg
		269-271A-R	gcgtgcagctgcatagaattcggatccgcgc
		272-274A-F	ctgcgctgcagctaaggctctgaataggactgc
		272-274A-R	cttagctgcagcgcagtatacatagaattcggatcc
		275-277A-F	gaaaggcagctgcaaataggactgcttcatcagc
		275-277A-R	ctatttgcagctgcctttctgcggcagtatacatag
		278-280A-F	ctggctgcagctgcttcatcagctgagactg
		278-280A-R	agcagctgcagccagagccttctttctgcg
		282-283A-F	gcagcagctgagactgctgataaactac
		282-283A-R	gcagtctcagctgctgcagcagtcctattcagag
		285-286A-F	ggggctgctgataaactactttctggag
		285-286A-R	agtagtttatcagcagccccagctgatgaagcagtc
		288-290A-F	gctgcagcactttctggagtttcaggctatgtaag
		288-290A-R	aaactccagaaagtgctgcagcagcagtctcagctgatgaag
		291-294A-F	gagactgctgataaactagctgctgcagtttcaggctatgtaagcaagcc
		291-294A-R	atagcctgaaactgcagcagctagtttatcagcagtctcagctg
		295-298A-F	gagttgcagctgctgtaagcaagccaaacgtgtatg
		295-298A-R	cttacagcagctgcaactccagaaagtagtttatcagcag
		298-300A-F	ctatgcagctgcaccaaacgtgtatgaaatcgacg
		298-300A-R	gtttggtgcagctgcatagcctgaaactccagaaagtag
		301-303A-F	gcaaggcagctgcttatgaaatcgacgagataatggaagc
		301-303A-R	cataagcagctgccttgcttacatagcctgaaactcc
		304-306A-F	gtggctgcagctgacgagataatggaagctacgaag
		304-306A-R	cgtcagctgcagccacgtttggcttgcttacatagc
		307-309A-F	gaaatcgctgcagcaatggaagctacgaaggatttcag
		307-309A-R	ccattgctgcagcgatttcatacacgtttggcttgc
		310-312A-F	cgagatagcagctgctacgaaggatttcagcg
		310-312A-R	gtagcagctgctatctcgtcgatttcatacacgtttg
		313-315A-F	gctgcagctgcattcagcgatgagtgcaagg
		313-315A-R	ctgaatgcagctgcagcttccattatctcgtcgatttc
		316-318A-F	ggatgcagctgctgagtgcaaggttggggaatc
		316-318A-R	ctcagcagctgcatccttcgtagcttccattatctc
		319-321A-F	cgatgctgcagctgttggggaatcagtgtacaagg
		319-321A-R	caacagctgcagcatcgctgaaatccttcgtagc

		322-324A-F	gcaaggcagctgcatcagtgtacaaggccaacatag
		322-324A-R	gatgcagctgccttgcactcatcgctgaaatc
		325-327A-F	gaagcagctgcaaaggccaacatagaaggtcg
		325-327A-R	gcctttgcagctgcttccccaaccttgcactc
		268-277DE-F	agaagcgcggatccgaattcaataggactgct
		268-277DE-R	ctgatgaagcagtcctattgaattcggatccgcgcttc
		278-287DE-F	gcagaaagaaggctctggataaaactactttctggagtttcagg
		278-287DE-R	aaactccagaaagtagtttatccagagccttctttctgcg
		298-307DE-F	caggctatgagataatggaagctacgaaggatttc
		298-307DE-R	cttccattatctcatagcctgaaactccagaaagtag
		308-317DE-F	gaaatcgacgatgagtgcaaggttgggg
		308-317DE-R	ttgcactcatcgtcgatttcatacacgtttggc
		318-327DE-F	gatttcagcaaggccaacatagaaggtcg
		318-327DE-R	gttggccttgctgaaatccttcgtagcttcc
		328-337DE-F	tcagtgtacgtaaagaaaatcaaggaaggtggtgcc
		328-337DE-R	attttctttacgtacactgattccccaaccttg
		288-294A-F	gcagctgccgcagcggctgcctcaggctatgtaagcaagcc
		288-294A-R	gcagccgctgcggcagctgcagcagtctcagctgatgaagc
		288-298PBS1-F	ggagacaaatctcatgtctcctcaggctatgtaagcaagcc
		288-298PBS1-R	ggagacatgagatttgtctccagcagtctcagctgatgaagc
		N5-283S-R	tgaagcagtcctattcagagcc
		283Y-F	tctgaataggactgcttcatacgctgagactgc
		N5-294G	agaaagtagtttatcagcagtctcag
		294Q-F	ctgctgataaaactactttctcaggtttcaggctatgtaag
		N5-303Y-R	cacgtttggcttgcttacatag
		304S-F	gcaagccaaacgtgtccgaaatcgacgag
		N5-310AT-R	ttccattatctcgtcgatttcatacac
		311-2 IY-F	gaaatcgacgagataatggaaatctacaaggatttcagcgatgagtg
Complemental	NopT	HC-proT-F	gaattcgagctcggtacctcatggccttccttggagg
		T-3-F	cacagtcccatcagtggttc
		T-3-R	gaaccactgatgggactgtg
		HC-T-R	cgcgggatcgagatctcgagtcatgtcatcttttgggtggtc
	NFR5	cherry-proNFR5-F	ccaagctgggctgcagggacatgagattgaagctcc
		proNFR5-NFR5-F	cccccacttcacaaacatggctgtcttctttcttacctc
		proNFR5-NFR5-R	gccatgtttgtgaagtgggg
		NFR5-JM-KD-F	gatgagtgcaaggttgggg
		NFR5-JM-KD-R	ccccaaccttgcactcatc
		cherry kpn1	
		Sac1-R	ggcgcgcctaggtaccggggaaattcgagc

1453