**The Rhizobial effector NopT targets Nod factor receptors to regulate symbiosis in**

*Lotus japonicus*

Hanbin Bao[1], Yanan Wang[1], Haoxing Li[1], Qiang Wang[1], Yutao Lei[1], Ying Ye[1], Hui

Zhu[1], Gary Stacey[2], Shutong Xu[1], Yangrong Cao[1*]

[1] National Key Lab of Agricultural Microbiology, Hubei Hongshan Laboratory,

Huazhong Agricultural University, Wuhan, Hubei, 430070, China

[2] Divisions of Plant Science and Technology and Biochemistry, Christopher S. Bond Life

Sciences Center, University of Missouri, Columbia, MO 65211, USA

* Corresponding author: Yangrong Cao, email: yrcao@mail.hzau.edu.cn

**Short title:** NopT associates with NFR1 and NFR5 to regulate symbiosis

**Keywords:** Legume–rhizobial symbiosis; rhizobial effector; Nod Factor Receptors;

NopT.

**One-sentence summary:** The rhizobial effector NopT forms a mutual interaction with

plant's two Nod factor receptors, NFR1 and NFR5, to precisely regulate rhizobial

symbiosis in *Lout japonicus*

**Abstract**

It is well-studied that type-III effectors are essential required by gram-negative bacteria to directly target different cellular pathways from hosts, thus enhancing their infection. However, in the unique context of legume-rhizobium symbiosis, the role of rhizobial effectors in regulating plant symbiotic pathways remains a subject of hypothesized understanding. Here, we show that NopT, a YopT-type cysteine protease from *Sinorhizobium fredii* NGR234 that could directly target symbiotic signaling pathway by associating with two Nod Factor Receptors (NFR1 and NFR5) from *Lotus japonicus*. NopT is the only effector out of 15 effectors from *S. fredii* NGR234 that inhibits cell death triggered by co-overexpression of NFR1 and NFR5 in *Nicotiana benthamiana*. NopT interacts with both NFR1 and NFR5. NopT proteolytically cleaves NFR5 in both *in vitro* and *in vivo* assays, but is later inactivated *via* phosphorylation by NFR1. NopT plays an essential role in mediating rhizobial infection in *L. japonicus*. The truncated NopT, lacking 50 N–terminal amino acids post-autocleavage, retains the ability to cleave NFR5 but no longer interacts with NFR1. Intriguingly, certain *Sinorhizobium* species exclusivley harbor this truncated NopT in their genomes. These findings unveil an intricate regulatory interplay between rhizobia and legumes, wherein rhizobial NopT actively cleaves NFR5, and this cleavage process is subsequently suppressed following NopT phosphorylation by NFR1. This discovery presents a compelling narrative of a nuanced regulatory interplay between rhizobia and legumes, highlighting rhizobial effector as a direct mediator targeting symbiotic pathway from plants.

**Introduction**

The legume–rhizobial symbiosis stands as one of the most crucial and intricate mutualistic interactions in nature. Its significance extends beyond the supply of the substantial amount of ammonium provided by rhizobia, influencing plant growth and

ecosystem sustainability, to serving as a pivotal model for investigating plant–microbe

interactions (Yang et al., 2022). The orchestration of this symbiosis involves the

perception of small molecules, such as nod factors (NFs) and surface polysaccharides

(Madsen et al., 2003; Kawaharada et al., 2015), along with the secretion of proteins,

inlcuding the type three secretion system (T3SS) effectors, to regulate legume–rhizobial

symbiosis (Sugawara et al., 2018; Zhang et al., 2021).

A receptor heterocomplex, consisting of two Lysin–motif receptor kinases (LYKs), plays

a central role in recognizing various polysaccharide molecules. In nonlegumes like

Arabidopsis, AtLYK5 and AtCERK1 (Chitin elicitor receptor kinase) are required for

perception of chitin, triggering plant immunity to combat fungal pathogens (Cao et al.,

2014). Conversely, in legumes, structurally similar receptors like *L. japonicus* NF

receptor1 (LjNFR1) and LjNFR5 are vital for recognizing rhizobial lipo–

chitooligosaccharide NF, initiating symbiotic signaling transduction (Broghammer et al.,

2012; Bozsoki et al., 2017). The intricate regulation of NFRs is highlighted by the

instances of hypersensitive cell death induced by their simultaneous overexpression in

*Nicotiana benthamiana* leaves (Madsen et al., 2011) and defense–like responses in

*Medicago truncatula* nodules upon overexpression of MtNFP (Nod factor perception, the

ortholog of NFR5) (Moling et al., 2014), perhaps reflecting potential conservation of the

plant LYK receptors in interactions with both pathogens and symbionts. Therefore, the

protein abundance of NFRs must be tightly controlled to avoid this resistance response,

allowing rhizobial infection to proceed, emphasizing the need for tight control to enable

rhizobial infection.

T3SS effectors play a critical role in suppressing pattern–triggered immunity (PTI)

during pathogen infections in plants (Jones and Dangl, 2006; Schubert et al., 2020). To

avoid diseases, specific resistance (R) genes are employed to recognize successful

avirulence (Avr) effectors which enable pathogens overcome PTI, known as

effector-triggered immunity (ETI), a much stronger immnune response combined with

programmed cell death (Jones and Dangl, 2006). However, in contrast to the well-studied

fucntion of Avr effectors from phytopathogenic bacteria in mitigating immunity by

different mechanisms, how these effectors from symbiotic bacteria might contribute to

mutualistic interaction is largely unknown.

Some effectors, such as YopT-type cysteine protease family, including *Pseudomonas*

AvrPphB, *Yersinia* YopT, and *Rhizobial* NopT, are well conserved across different

bacteria. AvrPphB from *Pseudomonas syringae* can directly proteolytically cleaves

Arabidopsis receptor-like cytoplasmic kinase (RLCK) family proteins including PBS1,

PBS1-like proteins, and BIK1, to suppress PTI (Zhang et al., 2010). While, the cleavage

of PBS1 serves as a "decoy" in activating RPS5 (RESISTANCE TO PSEUDOMONAS

SYRINGAE5)–mediated ETI (Shao et al., 2003b). *Yersinia* YopT directly cleaves Rho

family leading to their inactivation to disrupt actin cytoskeleton of host cells (Shao et al.,

2003b; Shao et al., 2003a). While, in *Rhizobium* species, NopT have a dual function in

regulating symbiosis. The exogenous expression of NopT from *Sinorhizobium fredii*

NGR234 in *S. fredii* USDA257 almost blocks nodulation in soybean Nanfeng 15 but not

other soybean cultivars (Khan et al., 2022), implying that NopT might act as an Avr

effector. However, NopT positively regulates nodule formation in the interaction betwen

*S. fredii* HH103 and soybean Suinong 14 (Li et al., 2023). NopT from *Bradyrhizobium*

sp. ORS3257 promotes rhizobial infection in nodules of *Aeschynomene indica* (Teulet et

al., 2019). Despite this, little is understood about how NopT precisely modulates

rhizobial infection and the variations among NopTs from different rhizobia strains.

The challenge in studying rhizobial effectors lies in the availability of suitable assays by

which to characterize their phenotypes. In this study, we capitalized on the observation

that ectopic expression of NFR1 and NFR5 in *N. benthamiana* leaves induces

programmed cell death (Madsen et al., 2011), a response that might be modulated by

rhizobial effectors. Among the 15 known T3SS effector proteins encoded within the *S.*

*fredii* NGR234 genome, only NopT, a homolog of the pathogenic effectors, YopT and

AvrPphB (Shao et al., 2002; Dai et al., 2008), could specifically suppress this cell death

response. This suggests that NopT might be an effector associated with NFRs. Our

findings demonstrate that NopT interacts with both NFR1 and NFR5 at the plasma

membrane, and cleaves NFR5 proteolytically at the juxtamembrane position to

suppresses NF–mediated physiological responses. Perhaps as a means to regulate NopT

activity, NFR1 phosphorylates intact NopT, inactivating its protease activity. However,

the autocleaved NopT retains the ability to cleave NFR5 but loses its association with

NFR1. Intriguingly, some *Sinorhizobium* species possess only the truncated versions of

NopT (relative to NGR234 NopT) that might evade suppression by NFR1. This study

enveils a distinct and mutual regulatory model, in which the rhizobial effector NopT

directly targets NFR5 to suppress symbiotic signaling, and NFR1-mediated

phosphorylation acts as a negative feedback mechanism. The model provides essential

insights into the intricate interplay between legumes and rhizobia, revealing a

counteraction where a rhizobal T3SS effector impairs NF signaling by directly targeting

NFR5, and NFR1 phosphorylates and inactivates the rhizobal T3SS effector in response.

**Results**

**NopT suppresses the cell death response from overexpression of NFR1 and NFR5**

To examine the role of effector proteins in regulating rhizobial infection, *S. fredii*

NGR234, a strain with exceptionally broad host–range was chosen. Initial experiments

were performed to screen all 15 effector proteins identified from *S. fredii* NGR234

(Kimbrel et al., 2013), for their ability to suppress or enhance the cell death response in *N.*

*benthamiana* leaves due to co–expression of LjNFR1 and LjNFR5 (Fig. S1A–S1C). Out

of the effectors tested, only NopT, a cysteine protease belonging to the YopT family

(Shao et al., 2002; Dai et al., 2008), could suppress LjNFR1 and LjNFR5–induced cell

death (Fig. 1A and S1D–S1F). Cysteine–93 of NopT is a known residue essential for its

protease activity (Dai et al., 2008). In contrast to the wild–type NopT protein, expression

of NopT$^{C93S}$, an inactive version, was unable to suppress cell death induced by NFR1 and

NFR5 compared with controls (Fig. 1B), indicating the protease activity of NopT is

essential for suppression of cell death in *N. benthamiana*. In contrast to the suppressive

function on cell death, NopM, an effector protein from *S. fredii* NGR234 reported to have

an E3 ligase activity, induced strong cell death when overexpressed in *N. benthamiana*

leaves (Fig. 1C).

As controls, we tested whether the action of NopT was specific for NFR1/NFR5

expression or whether it could act generally to suppress cell death. A variety of other

proteins have been reported to induce cell death when expressed in *N. benthamiana*

leaves; specifically, Avr3a and R3a (Bos et al., 2006), apoptosis regulator BAX

(Bcl-2-associated X protein) (Gan et al., 2009), INF1, an elicitor from *Phytophthora*

*infestans* (Vleeshouwers et al., 2006), Arabidopsis AtCERK1 (Cao et al., 2014), and

NopM (Fig. 1C). In contrast to the case with LjNFR1 and LjNFR5, expression of NopT

was unable to suppress the cell death induced when each of these proteins was expressed

in tobacco leaves (Fig. 1D). Hence, the effect of NopT appears to be specific for the NF

receptors. These results suggested a hypothesis that rhizobial NopT protease might act

directly on the NFR1 and/or NFR5 proteins.

**NopT colocalizes with and interacts with NFR1 and NFR5 at the plasma membrane**

NopT contains an acylation site that appears to localize NopT to cellular membranes

(Dowen et al., 2009). Therefore, we tested the possibility that NopT, NFR1 and NFR5

co–localize on the plasma membrane. *In vivo* bimolecular fluorescence complementation

(BiFC) assays were used to examine the colocalization of NopT with LjNFR1 and

LjNFR5. NopT, LjNFR1 and LjNFR5 were all C–terminally tagged with nYFP and

cYFP. Fluorescence signals representing the associations between NopT and LjNFR1,

LjNFR5, as well as itself, were detected at the plasma membrane (Fig. 2A and 2B),

suggesting that NopT colocalized with and interacted with both LjNFR1 and LjNFR5.

Interactions between NopT and LjNFR1 and LjNFR5 were further verified using a Split–

luciferase (Split–LUC) assay (Fig. 2C). Co–expression of NopT–cLUC and LjNFR1–

nLUC or LjNFR5–nLUC produced strong luminescence signals in leaf discs compared

with the negative control expressing FLS2–nLUC and NopT–cLUC (Fig. 2C).

It has been known that the acylation of NopT could change its subcellular localization

(Dowen et al., 2009). We then exaimed the interaction between NFR1/NFR5 and

NopT$^{G50A/C51A/C52A}$, a mutant version on the acylation sites. The interactions between

NopT$^{G50A/C51A/C52A}$ and NFR1/NFR5 were significantly reduced compared to NopT$^{WT}$ in

Split–LUC assays (Fig. S2A and S2B). While the mutation at catalytic site of NopT

(NopT$^{C93S}$) remained similar interaction with NFR1 and NFR5 (Fig. S2C). The

interaction results were further confimred by co–immunoprecipitation (co-IP) assay. The

C–terminally HA–tagged LjNFR1 and LjNFR5 could be precipitated by anti–FLAG

antibody in the sample expressing NopT–FLAG compared with controls (Fig. 2D). The

interaction between NopT and LjNFR1 and LjNFR5 suggested that LjNFR5 or LjNFR1

might be a target protein for proteolysis by NopT.

**NopT proteolytically cleaves LjNFR5 at the juxtamembrane domain**

A previous study showed that NopT protease activity is detectable upon autocleavage of

about 50 amino acids from its N–terminus (Dai et al., 2008). We next tested whether

NopT could directly proteolyze LjNFR1 and/or LjNFR5 when co–expressed in *N.*

*benthamiana* leaves. The addition of NopT but not NopT$^{C93S}$, an inactive version,

resulted in proteolytic cleavage of LjNFR5–GFP but not LjNFR1–GFP as shown by

immunoblotting of the SDS–PAGE gel (Fig. 3A and S3A). The *in vivo* cleavage assay in

*L. japonicus* transgenic root also showed that NopT but not NopT$^{C93S}$ resulted in

proteolytic cleavage of LjNFR5 (Fig. 3B). Based on the molecular weight change, the

cleavage sites were predicted to occur within the cytoplasmic domain (CD) of LjNFR5.

Therefore, the CD of NFR5 was constructed as a recombinant protein (Strep–NFR5$^{CD}$–

HA) and then co–expressed with NopT–FLAG or the inactive version NopT$^{C93S}$–FLAG

in *Escherichia coli* cells. The results showed that the active NopT protease could cleave

NFR5$^{CD}$ resulting in a band roughly ~5 kDa less than the control as detected using anti–

HA antibody (Fig. 3D). In the same experiment, when the CD of NFR1 was co–

expressed with NopT, no cleavage product was detected in immunoblotting assay (Fig.

3C). However, migration of the bands representing NopT or NopT$^{C93S}$ was significantly

retarded as seen after immunoblotting (Fig. 3C). This retardation of NopT protein band

may be the result of phosphorylation by NFR1.

The cleavage product with ~5 kDa lower molecular weight in the cleavage assay,

suggested that the cleavage site is located within the juxtamembrane (JM) position of

NFR5. To test this hypothesis, SUMO–NFR5$^{KD}$–HA, a recombinant protein with the JM

of NFR5 replaced with a SUMO tag, was constructed and used for the cleavage assay.

Indeed, NopT could not cleave SUMO–NFR5$^{KD}$–HA (Fig. S3B). To confirm this

conclusion, the JM of NFR5 was used to bridge SUMO and GFP tags to create a

recombinant protein, SUMO–JM–GFP. *In vitro* cleavage assay, strong band intensity

representing GFP was detected in the flow though sample incubated with NopT but not

NopT$^{C93S}$ (Fig. 3E). These data strongly indicated that the cleavage site for NopT is

located at the JM of NFR5.

**The cleavage of LjNFR5 at JM by NopT is dependent on multiple residues of NFR5**

NopT has an autocleavage site within its N–terminus, we identified the peptide

(DKLLSGV) from Asp–288 to Val–294 of NFR5 (Fig. S4B) as showing the highest

similarity with the NopT autocleavage region (DKMGCCA). However, the mutant

version of NFR5$^{CD}$ with the peptide (DKLLSGV) replaced with 7 alanines could still be

cleaved by NopT (Fig. S3C). NopT and its homlog AvrPphB (Shao et al., 2003b), an

effector from the pathogen *Pseudomonas syringa*, belong to the same protein family.

Hence, we set up a control experiment by constructing a version of NFR5$^{CD}$, where the

"DKLLSGV" motif was replaced with the cleavage sites of AvrPphB. In the cleavage

assay, we found that this modified NFR5$^{CD}$ was cleaved by both AvrPphB and NopT, and

the cleaved product of modified NFR5$^{CD}$ by AvrPphB shifting faster than by NopT (Fig.

S3C), suggesting that the cleavage site of NFR5 by NopT located ahead of "DKLLSGV"

motif. As mentioned above, NopT undergoes autocleavage but it is not clear whether this

is due to inter–proteolysis or intra–proteolysis. His–205 and Aps–220 of NopT are two

other conserved residues required for the protease activity of YopT, another homolog of

NopT from *Yesinia* whose function has been extensively studied (Shao et al., 2003b).

Therefore, we targeted these residues for site–directed mutagenesis. The cleavage assay

showed that NopT could not proteolyze NopT$^{C93S}$, NopT$^{H205A}$, or NopT$^{D220A}$ (Fig. S3D),

suggesting that intra–proteolysis might occur and the autocleavage site of NopT could not

be used to predict the cleavage site of substrate.

To define the cleavage site of NFR5, we created a variety of point mutations covering the

JM domain of NFR5 (Val-269 to Cys-320) and first seven a.a. of the kinase domain

(Lys-321 to Tyr-327). First, 17 mutant versions of NFR5$^{JM}$ were created with three

adjacent amino acids replaced with three alanines for proteolysis assay in *E. coli* cells.

However, all 17 mutant versions of NFR5$^{CD}$ could be cleaved by NopT but not NopT$^{C93S}$

(Fig. S3E–S3H). An additional 7 truncated versions of NFR5$^{JM}$, with deletion of 10

adjacent residues, were created and tested for cleavage by NopT. Surprisingly, the

cleavage products were still observed from all 7 truncated versions of NFR5$^{JM}$ (Fig. S3I).

The *in vitro* cleavage assay suggested that the JM domain might have multiple sites

susceptible to be cleaved by NopT, which is different from the specific cleavage site

identified for its homologous protein AvrPphB. In a parallel experiment, the recombinant

protein, His-SUMO-NFR5$^{CD}$-GST expressed in *E. coli* was subjected for N-terminal

protein sequencing determined by Liquid Chromatography Mass Spectrometry analysis

after cleavage. Based on allignment using identified peptides, the cleavage site of NFR5

by NopT was mapped to be after four basic amino acids in the JM of NFR5, "RRKK"

from 271 to 275 a.a. of NFR5 (Fig. S4C). However, the result is not consistance with the

data from cleavage assay. When the region from Tyr-268 to Leu-277 was deleted, NopT

could still proteolyze NFR5-CD (Fig. S4I). The cleavage assay suggested that NopT

might prefer to proteolyze NFR5 after four polybasic residues in the NFR5$^{JM}$, but other

sites could also be cleaved by NopT.

Because LjNFR5 belongs to a subfamily of plant LYK family proteins that lacks kinase

activity (Yang et al., 2022), we then asked whether NopT could proteolyze other LYK

proteins in this subfamily, which might be helpful to characterize the biochemical

function of NopT. Hence, the CDs from Arabidopsis AtLYK5, *L. japonicus* LjLYS11

and *M. truncatula* MtNFP were used for cleavage assays in the presence of NopT. As

shown in Fig. S3J, NopT, but not NopT$^{C93S}$, could cleave the CDs of both AtLYK5 and

LjLYS11. Replacement of NFR5$^{JM}$ with the JM domains from either AtLYK5 or

LjLYS11 also resulted in cleavage by NopT (Fig. S3J). The data strongly indicated that

NopT could cleave both AtLYK5 and LjLYS11 at their JM domains. Based on sequence

analysis of the JM region of NFR5 and its homologous proteins, five conserved residues

(Ser-283, Gly-294, Tyr-303, Ala-310, Thr-311) were identified and mutated for the

proteolysis assay (Fig. S4B). As shown in Fig. 3F, the mutant version of NFR5 (i.e.,

NFR5$^{CD}$ with five amino acid mutations) could not be cleaved by NopT. Interestingly,

NopT coud not protelyze MtNFP$^{CD}$ (Fig. 3G). Then, we generated two recombinant

proteins, by switching the JM between NFR5$^{CD}$ and NFP$^{CD}$, generating NFR5$^{JM}$-NFP$^{KC}$

(KC represents kinase domain and C-terminal tail region) and NFP$^{JM}$-NFR5$^{KC}$ for *in vitro*

cleavage assays. As shown in Fig. 3G, NopT could proteolyze NFP$^{JM}$-NFR5$^{KC}$ but not

NFR5$^{JM}$-NFP$^{KC}$. The data suggested that the cleavage of NFR5 by NopT might be

dependent on the confirmational structure of NFR5$^{KC}$ and/or other specific amino acids in

NFR5$^{KC}$. To define this, two additional recombinant proteins with large switch between

NFR5$^{CD}$ and NFP$^{CD}$, NFR5$^{268-445}$-NFP$^{458-595}$ and NFP$^{270-457}$-NFR5$^{456-595}$, were constructed

for proteolysis (Fig. S3K). However, both of them could be cleaved by NopT, suggesting

that a much large region on NFR5$^{KC}$ is required for being proteolyzed by NopT. Based on

all these cleavge results, we concluded that NopT could proteolyze NFR5 at the JM

domain, which nonadjacent mutiple residues within the NFR5$^{JM}$ and the specific KC

region are required for the proteolysis.

**NopT phosphorylation by LjNFR1 suppresses its proteolytic activity**

The perception of rhizobia NFs by the heterocomplex of NFR1 and NFR5 involves the

trans–phosphorylation from the NFR1 kinase to the pseudo–kinase NFR5 (Madsen et al.,

2011). Rhizobial effector NopT associates with both NFR1 and NFR5, which suggests

that NopT might be a phosphorylation target of NFR1 and involved in the

transphosphorylation process between the two receptors. Kinase assays showed that

NFR1–CD but not the kinase–inactive NFR1$^{K351E}$–CD (Lys-351 is the conserved residue

required for ATP binding) could phosphorylate NopT (Fig. 3C and 4B) and NFR5–CD

(Fig. S5) as shown by band retardation on SDS–PAGE gels identified using

immunoblotting. The band representing phosphorylated NopT shifted upon the addition

of a commercial phosphatase, calf intestinal alkaline phosphatase (Fig. 4B), indicating

that NopT as well NFR5$^{CD}$ are phosphorylation targets of the NFR1 kinase. NopT

autocleavage also gives two bands and the site mutant of NopT at C93S did not affect the

interactions between NFRs(Fig. S2C). To avoid multi-bands interference of cleaved

NopT, NopT$^{C93S}$ was expressed in both wild type plants and *nfr1* mutant then treated with

rhizobia or water. After analyzed by $Zn^{2+}$-Phos-tag PAGE gel, several shifted bands were

strengthened after rhizobia treatment in WT roots but not in the *nfr1* mutant roots (Fig.

4A), suggesting that the phosphorylation of NopT could be induced by NFR1 and

multi-sites of NopT were phosphorylated *in planta*. The phosphorylation sites of NopT

were further identified using Liquid Chromatography Mass Spectrometry (Table S1). We

then addressed whether NopT phosphorylation by NFR1 regulates its proteolytic activity

on NFR5. All the potential phosphorylation sites of NopT were substituted into either

alanine in order to block phosphorylation of these sites or aspartate, which mimicks the

acidic condition representing a phosphorylated status. These mutant proteins were then

used in the cleavage assay. Aspartate substitutions on three phosphorylation sites of

NopT, Thr–97, Ser–119, and Thr–257, resulted in the loss of the ability of NopT to

cleave NFR5–CD, as well as the ability to autocleave (Fig. 4C). In contrast, NopT

proteins with alanine substitutions on these three sites retained the ability to cleave

NFR5–CD and to autocleave (Fig. 4C). The data suggest that phosphorylation of NopT

by NFR1 likely inhibits proteolytic activity, retaining a fully functional NFR5 and a full

length NopT.

**NopT dampens rhizobial infection**

The fact that *S. fredii* NGR234 NopT could directly target and cleave NFR5 indicates that

NopT may be an important player regulating rhizobial infection in legumes. Nod factor

signal is known to function in the early stage of rhizobial infection and nodulation

specificity (Geurts et al., 2005; Wang et al., 2012). Prior work showed that type three

secretion system of *S. fredii* NGR234 might not regulate nodule numbers few weeks after

inoculation in *L. japonicus* (Ausmees et al., 2004). Therefore, we focused on the early

stage of rhizobial infection. Indeed, inoculation with GFP–labelled *S. fredii*

NGR234Δ*nopT* induced a massive infection within the root hairs of *L. japonicus* (Gifu)

plants compared with the favorable infection by the NGR234 wild type strain (Fig. 5A

and 5C). Transgenic expression of NopT in *L. japonicus* root hair reduced the infection of

*Mesorhizobium loti* MAFF303099 (Fig. 5G). In *L. japonicus* plants transgenically

expressing *pNIN*:*GUS*, greater range of GUS expression (Fig. 5B and 5D) and formation

of more nodule primordia (Fig. 5E) also reflected a stronger plant response to infection

by NGR234Δ*nopT* relative to the wild–type strain. To confirm the massive infection is

due to the loss of NopT, NopT was expressed in the *S. fredii* NGR234Δ*nopT* strain and

used for inoculation in *L. japonicus*. Complementation with NopT in NGR234Δ*nopT*

reduced the massive infection foci to the similar levels as the WT strain (Fig. 5F). We

then tested whether the proteolytic activity of NopT is required for rhizobial infection.

The protease–dead version of NopT$^{C93S}$ and three phosphorylation mimic versions of

NopT, i.e., NopT$^{T97D}$, NopT$^{S119D}$, and NopT$^{T257D}$, which lack proteolytic activity, were

individually expressed in NGR234Δ*nopT*. Compared with the high levels of infection

foci induced by NGR234Δ*nopT* in *L. japonicus* plants, inoculation with these four mutant

strains induced much lower levels of infection, albeit slightly higher than that induced by

the wild–type strain (Fig. 5F). Inoculation with NGR234Δ*nopT* produced more nodule

primordia compared with wild–type strain inoculation 14 dpi (Fig. 5E). To verify whether

the cleavage of NFR5 reduce rhizobial infection, NFR5–WT and NFR5$^{5m}$, a mutant

version with 5 amino acids substitutions at JM that is resistant to NopT cleavage, were

expressed in the *nfr5-3* roots under the control of its native promoter. However,

expression of NFR5$^{5m}$ was not observed to rescue the infection phenotype of *nfr5-3*

mutant plants (Fig. S5A). The possible reason is that amino acids substitutions at JM

might lead to a conformation change of NFR5 that is required for its function in

symbiotic signaling transduction. Therefore, NFR5-NFP$^{KC}$, the KC domain of NFR5

replaced with KC domain of NFP, was expressed in the roots under the control of native

promoter (Fig. S5B). In the *nfr5-3* roots expressing NFR5-NFP$^{KC}$, no difference of

rhizobial infection was observed between *S. fredii* NGR234Δ*nopT* and *S. fredii* NGR234

wild type strain, suggesting that the cleavage of NFR5 reduced the rhizobial infection

levels. All these data strongly indicated that the protease activity of NopT, the

phosphorylation status of NopT, and the proteolysis of NFR5 are important for rhizobial

infection.

**The truncated version NopT$^{ΔN50}$ cleaves NFR5 but fails to interact with NFR1**

The significance of rhizobial NopT in mediating rhizobial infection led us to analyze its

homologs from other rhizobium species. Sequence comparisons based on the amino acids

of NopT homologs showed that *Sinorhizobium* and *Bradyrhizobium* were located in

different clades (Fig. S7). *B. diazoefficiens* (USDA110) is a typical *Bradyrhizobium*

strain used to study the symbiotic interaction with soybean. Two NopT homologous

genes, named *NopT1$_{USDA110}$* and *NopT2$_{USDA110}$* were identified from the USDA110

genome (Fotiadis et al., 2012). However, when tested, neither NopT1$_{USDA110}$ or

NopT2$_{USDA110}$ could not cleave NFR5 compared to NopT$_{NGR234}$ (Fig. 6A). The possible

scenario is that NopT homologs from *Bradyrhizbium* species might have lost the ability

to cleave NFR5 and/or may act on other, unknown cellular target proteins. The genome

of some *Sinorhizobium* species, for example *S. fredii* USDA257 and S. *fredii* HH103,

encode *NopT* homologous genes predicted to produce same truncated proteins with a 57

aa deletion from its N–terminus, relative to the NGR234 NopT sequence, which is almost

the same as the truncated $NopT_{NGR234}$ after autocleaveage. When these NopT homologs

(i.e., $NopT_{USDA257}$ and $NopT_{HH103}$) were tested, both were observed to have the ability to

cleave NFR5 (Fig. 6A). However, these NopT homologs could not suppress the cell death

triggered by NFR1 and NFR5 in *N. benthamiana* leaves (Fig. 6B) and failed to

complement the massive infection of NGR234Δ*nopT* (Fig. 6C and 6D), suggesting that

$NopT_{USDA257}$ and $NopT_{HH103}$ retained the same catalytic activity as $NopT_{NGR234}$ but might

have different substrates. We then asked whether the truncated version of $NopT^{\Delta N50}$,

which mimics the truncated NopT after autocleavage, retained the ability to interact with

NFR1. However, after repeated tests, the direct interaction between $NopT^{\Delta N50}$ and NFR5

could be detected but not the corresponding interaction with NFR1 using Split–LUC

assays (Fig. S2D). The data suggest that some *Sinorhizobium* species, for example, *S.*

*fredii* NGR234, *S. fredii* USDA257 and *S. fredii* HH103, use a truncated NopT to

regulate the rhizobial infection process in a way that avoids negative regulation by NFR1

phosphorylation, which is not true of all rhizobial strains that possess NopT homologs.

**Discussion**

Plant-microbe interactions involve two sophisticated and intelligent organisms, each

employing distinct "weapons" or strategies to communicate mutually, thereby securing

more nutrients for their survival (Jones and Dangl, 2006; Schubert et al., 2020). A

well-established regulatory model arises from the study of plant-pathogen interactions,

where T3SS effectors from gram-negative bacteria are injected into plant cells to directly

target and suppress the plant's immune pathway, favoring pathogenic bacterium infection

(Jones and Dangl, 2006). However, the unique model of mutual regulation between

legumes and rhizobia in symbiotic interactions remains largely unknown. The broad goal

of this study was to offer essential insights into the mutual regulation mechanisms

402 whereby rhizobial effectors directly target symbiotic signaling pathways to dampen their

403 infection in host cells.

405 Understanding how plants recognize different microbes as "friends" or "foes" is a central

406 question in biology. As the first line of plant response to microbes, the perception of

407 microbial signals by receptor proteins from plants is essential for inducing different

408 signaling pathways to establish various interaction patterns–either symbiotic, pathogenic,

409 or no responses (Zipfel and Oldroyd, 2017). Therefore, it is not surprising to discover that

410 the direct targeting of these receptors by bacterial effectors represents an efficient

411 strategy to manipulate plant signaling pathways for enhancing bacterium infection. In the

412 well-studied *Arabidopsis–Pseudomonas syringa* interaction model, the flagellin receptor

413 FLS2 (Flagellin Sensing 2) and its coreceptor BAK1 (BRI1-[Brassinosteroid insensitive

414 1]-associated receptor kinase 1), triggering plant immunity against pathogenic infection,

415 are targeted by several effectors from bacteria (Zipfel et al., 2004; Sun et al., 2013). For

416 example, AvrPto and HopB1 could suppress the kinase activity and cleave the

417 Arabidopsis receptor complex, respectively (Xiang et al., 2008; Li et al., 2016). AvrPtoB,

418 an E3 ligase in *P. syringae*, targets FLS2 to promote its degradation, resulting in

419 decreased immune responses in plant cells (Goehre et al., 2008). Considering the

420 evolutionary importance of effectors in infection, it is rational to identify that the

421 rhizobial effector NopT, a member of the C58 protease family, can directly target NFRs,

422 manipulating symbiotic signaling pathways in plants. This was demonstrated using a

423 unique screening system in *N. benthamiana* leaves (Fig. S1A-C), in which co-expression

424 of two symbiotic receptors, NFR1 and NFR5, could lead to plant cell death to identify

425 NopT, the only effector from *S. fredii* NGR234. *S. fredii NGR234Δnopt* mutant strains

426 hyperinfect *L. japonicus* roots, and overexpression of NopT in the roots reduces rhizobial

427 infection (Fig. 5A-5D), consistent with the observed suppression of cell death in *N.*

428 *benthamiana* leaves (Fig.1A). The interaction between NopT and NFRs has been verified

and is impaired by the acylation site of NopT (Fig.2 and S2A), suggesting that the plasma

membrane localization of NopT is essential for targeting NFRs.

In the extensively studied context of microbial pathogenesis, various effectors from

pathogens have been proposed to directly suppress the immune system and promote

virulence in host cells (Tang et al., 2017; Schubert et al., 2020). The C58 protease family,

including YopT and AvrPphB, represents such effectors that disarm the host defense

system to favor pathogenic infection in both animals and plants (Shao et al., 2003a;

Zhang et al., 2010). *Yersinia* YopT cleaves the Rho family GTPase to disrupt the actin

cytoskeleton, inhibiting the phagocytosis of pathogens by hosts (Shao et al., 2003a).

Conversely, *P. syringae* AvrPphB cleaves multiple members of the RLCK (receptor-like

cytoplasmic kinase) subfamily VII to suppress the PTI response (Zhang et al., 2010). In

contrast, the cleavage of PBS1 functions as a "decoy" to trigger strong immunity against

bacterial infection (Shao et al., 2003b). We confirmed that NopT cleaves NFR5 at the JM

domain through different assays (Fig.3). Mutants with five dispersed residues (S283,

G294, Y303, A310, T311) at the JM domain (Fig.3F) and the replacement of NFR5$^{\text{KD-CT}}$

with NFP$^{\text{KD-CT}}$ (Fig.3G) resist cleavage by NopT, suggesting a joint decision of cleavage

involving both the JM domain and KD-CT domain. Although belonging to the same

protease family, the cleavage sites of NopT and YopT are polybasic (Shao et al., 2003a;

Schmidt, 2011), in contrast to AvrPphB, whose cleavage site is canonical with seven

adjacent amino acids involved (Kim et al., 2016). The different cleavage sites of

YopT-type effectors indicate that bacteria regulate different physiological pathways to

establish specific interactions with host plants.

Similar to the plant pathogenic effector AvrPphB, the protease activity of NopT appears

to be conserved for this function by cleaving the soybean PBS1 homolog protein to

restrict symbiosis (Khan et al., 2022). Indeed, transient expression of *S. fredii* NopT in *A.*

*thaliana* and *N. tabacum* could induce strong immune responses and hypersensitive cell

death (Fig. S8), indicating that rhizobial NopT might act as an avirulent effector in some

nonhost plant species. Interestingly, we failed to obtain stable transgenic lines

constitutively expressing NopT in *L. japonicus*. However, the generation of transgenic

root hairs was not affected, implying that NopT might act as an Avr effector whose

overexpression leads to whole plant lethality but not root growth. Therefore, given this

conservation of protease function, it is consistent with a successful Avr effector that

NopT has the ability to proteolyze not only NFR5 but also Arabidopsis AtLYK5 and *L.*

*japonicus* LjLYS11, two receptors with chitin-binding affinities that mediate plant PTI

responses (Cao et al., 2014; Gysel et al., 2021). NFR5, AtLYK5, and LjLYS11 are LYKs

belonging to the same subfamily that lacks kinase activities. Thus, NopT may not only

suppress the nod factor–mediated symbiotic signal to avoid excessive infection by *S.*

*fredii* NGR234 but also function in inhibiting the signal triggered by chitin-like

compounds. The cleavage of receptor-like kinases (RLKs) may be an overlooked strategy

to regulate their function in plant biology compared to the well-studied phosphorylation

to control RLK activation. While cleavage regulation is well studied in neurobiology,

e.g., the Notch signal, the proteolysis of each Notch molelue leads to signaling activation

(Kopan and Ilagan, 2009). Arabidopsis BAK1, identified as an universal co-receptor to

regulate plant growth and MAMPs perception, undergoes proteolytic cleavage, which is

important for both BR signaling and PTI response with an unclear mechanism (Zhou et

al., 2019). NFR5 is a pseudokinase that functions as an output of NF signaling. The

cleavage of NFR5 by rhizobial NopT releases the cytoplasmic domain of NFR5, but the

precise function of whom is of interest to be determined.

The legume–rhizobial symbiosis is a unique case in the context of plant–microbe

interactions. Mutual regulation between pathogenic effectors and host proteins has been

reported in plant pathogenesis. For example, the *Pseudomonas* conserved effector

AvrPtoB could directly target key components in the immunity pathway to promote its

virulence (Goehre et al., 2008). However, the virulent function of AvrPtoB could either

be enhanced or dampened by host proteins. AvrPtoB could be phosphorylated by

SnRK2.8 to promote its virulence activity (Lei et al., 2020). Plant lectin receptor-like

kinase LexRK-IX.2, Pto, and Fen kinases could also phosphorylate AvrPtoB to inactivate

its E3 ligase activity and undermine its ability to suppress PTI (Ntoukakis et al., 2009; Xu

et al., 2020). A similar strategy was used by legumes to dampen the "virulence" of the

rhizobial effector NopT, favoring mutualistic infection. NFR1 phosphorylates NopT at

several residues. The phosphomimetic NopTs (i.e., $NopT^{T97D}$, $NopT^{S119D}$, and $NopT^{T257D}$)

are unable to cleave itself and NFR5 (Fig. 4C). The expression of these phosphomimetic

NopTs separately in *S. fredii* NGR234Δnopt mutant strains cannot complement the

hyperinfection phenotype (Fig.5F). These data indicate that *L. japonicus* deploys NFR1

to phosphorylate NopT to dampen its catalytic activity and may protect NFR5 from

cleavage, favoring rhizobial infections. However, $NopT^{\Delta N50}$, mimic version of

autocleaved NopT, retained the ability to interact with NFR5 but not NFR1 (Fig. S2D).

Moreover, intact NopT, but not autocleaved NopT, are migrated by expressing $NFR1^{CD}$

in immunoblotting (Fig. 3C and 4B), suggesting autocleaved NopT may escape from

being phosphorylated and inactivated by NFR1.

In different rhizobium species, a variety of NopT homologs were characterized. NopT

homologs from some *Bradyrhizobium* species lose the ability to cleave NFR5, while

NopT homologs from some *Sinorhizobium*, truncated versions without about 50 a.a. at its

N-terminus, retain the ability to cleave NFR5 (Fig. 6A). The type three secretion signal

depends on the N-terminal of effectors (Akeda and Galán, 2005). However, an earlier

study has shown that NopT from *S. fredii* HH103 exhibited a positive effector on

nodulation in soybean SN14 cultivar (Li et al., 2023), suggesting that another unknown

mechanism might be involved in $NopT_{HH103}$ secretion. Interestingly, these truncated

NopTs still keep the function to cleave NFR5 (Fig. 6A) and autocleaved NopT$_{NGR234}$

cannot interact with NFR1 (Fig. S2D), indicating that NopTs from those strains and

autocleaved NopT$_{NGR234}$ might lose the ability to be suppressed by NFR1. Notably,

natural truncated NopT from *S. fredii* USDA257 and *S. fredii* HH103 are the same in

protein sequence, but they lack the acylation site compared with NopT$_{NGR234}$. Therefore,

NopT$_{USDA257}$ and NopT$_{HH103}$ lose the ability to suppress cell death triggered by NFR1 and

NFR5 in *N. benthamiana* leaves (Fig. 6B) and failed to complement the massive infection

of NGR234*ΔnopT* (Fig. 6C and 6D), implying NopT$_{USDA257}$ and NopT$_{HH103}$ may have a

different target from NopT$_{NGR234}$ in the host plant. In addition, knockout of NopT in

*Bradyrhizobium* sp. ORS3257 was shown to result in uninfected nodule formation,

suggesting that, in this system, NopT$_{ORS3257}$ might play a role in nodule formation (Teulet

et al., 2019). The versatile function of NopT homologs in different rhizobium species

suggests that different regulatory processes might be involved in specific rhizobial

symbiosis.

In this study, we present a mutual regulation model in which the rhizobial effector NopT

directly targets NFR5 to suppress symbiosis signals and is then inactivated by NFR1 *via*

phosphorylation. NopT appears to retain some pathogenic-like features in that it can

cleave the *Arabidopsis* chitin receptor AtLYK5 and induce cell death in some

non-leguminous plants, such as *N. tabacum* and *Arabidopsis*. Thus, the cleavage of NFR5

by NopT might be due to the structure similarity to AtLYK5 targeted by NopT. The

suppression of proteolytic activity of NopT *via* phosphorylation by NFR1 appears to

work as a feedback loop to allow symbiotic signaling in compatible hosts. The model

(Fig. 6E) provides essential insights into the legume–rhizobial symbiosis, where rhizobia

deploy an effector to dampen symbiotic signaling, and legumes slightly recur infection by

suppressing NopT to employthe nod factor signal in response. Even though this couple,

legume and rhizobia, tries to get together and strives to improve their living conditions,

we catch a glimpse into the arguments and the recovery conversations between them.

**Materials and Methods**

Plant materials

*Lotus japonicus* Gifu B–129 cultivar was provided by the Center for Carbohydrate

Recognition and Signaling (https://lotus.au.dk/), and used as wild type for nodulation

assays. The Gifu *pNIN*:*GUS* transgenic line was kindly provided by Dr. J. Stougaard

from the Aarhus University (Heckmann et al., 2011). All *L. japonicus* seeds were treated

with concentrated sulfuric acid for 10 min, followed by surface sterilization in 1% (w/v)

NaClO for 8 min. After synchronization of germination at 4°C in the dark for 2 d, the

seeds were placed on half–strength Murashige & Skoog (MS) medium containing 0.8%

(w/v) agar for germination at 22°C for 2 d in the dark. Seedlings were then transferred to

growth pots containing vermiculite supplied with half–strength B&D (Broughton &

Dilworth) medium without nitrogen under long–day conditions (16–h light/8–h dark) at

22°C. Ten–day–old seedlings were inoculated with *Sinorhizobium fredii* NGR234

($OD_{600}$=0.02) for infection assays.

Plasmid construction

All plasmids used in this study were generated using MultiF Seamless Assembly Mix

(ABclonal, catalog number RK21020). The plasmid pGWB514 was used as an original

plasmid for all binary vector construction (Nakagawa et al., 2007). Briefly, Different tags

including HA tag, Myc tag, FLAG tag, Strep tag, nLUC, cLUC, and GFP were

respectively amplified using a forward primer containing a *Kpn*I site and ligated into

pGWB514 digested with *Xba*I and *Sac*I using seamless cloning method as described

above. The generated plasmids were named pG5XX–HA, pG5XX–Myc, pG5XX–FLAG,

pG5XX–nLUC, pG5XX–cLUC, pG5XX–GFP, respectively. The coding sequences of *L.*

*japonicus NFR1* and *NFR5* were cloned into pGWB514 and pG517 between *Xba*I/*Kpn*I

to generate NFR1–HA and NFR5–Myc fusion constructs under the control of the 35S

promoter. then, the DNA fragment containing *pro35S*:NFR5–Myc–NosT was amplified

and cloned into pG514–NFR1 at *Sbf*I. The final plasmids contained *pro35S*:NFR1–HA–

NosT and *pro35S*:NFR5–Myc–NosT. Rhizobial genes encoding effectors from *S. fredii*

NGR234 were cloned into pG5xx–FLAG between *Xba*I/*Kpn*I. To detect the interaction

between NFR1/NFR5 and NopT in *N. benthamiana*, the full–length *NFR1/NFR5*, *NopT*

and *AtFLS2* coding sequences were cloned into pG5xx–Nluc, pG5xx–Nluc,

pSPYCE(MR) and pSPYNE(R)173 (Waadt et al., 2008). For cleavage assay in *N.*

*benthamiana*, the *NFR5* and *NopT* coding sequences were cloned into pG5XX–FLAG

GFP and pG5xx–Strep respectively. For cleavage assay in *E. coli*, the sequence encoding

the cytoplasmic domain of NFR5 was cloned into pACYC duet Strep–HA, the sequences

encoding the kinase domain of NFR5 fused to HA tag and the juxtamembrane domain of

NFR5 fused to the GFP tag was cloned into pSUMO. The point mutants in NFR5$^{CD}$ and

NopT were created from the pACYC duet Strep–NFR5$^{CD}$–HA and pET28a NopT–FLAG

template using site–directed mutagenesis PCR. For *in vivo* cleavage assay in *L.*

*japonicus*, NopT-FLAG or NopT$^{C93S}$-FLAG was cloned into pUB GFP between *Xba*I and

*Kpn*I, and then 35S-NFR5-Myc-NosT was cloned in pUB NopT GFP at *Pst*I. For

phosphorylation assay, the sequence encoding the cytoplasmic domain of NFR1 fused to

a Myc tag was cloned into pCDF duet. For rhizobium infection assays, 300 bp upstream

of *NopT* and site mutants of *NopT* were fused by overlapping PCR and cloned into

pHC60.

Transient expressing and in *Nicotiana* species

The plasmids for transient expression of INF1, Avr3 and R3a in *N. benthamiana* were

kindly gifted by Dr. Juan Du from HZAU. *Agrobacterium* (*Agrobacterium tumefaciens*)

strain EHA105 carrying various constructs were mixed with an *Agrobacterium* culture

harboring the P19 suppressor in an infiltration buffer (10 mM MgCl$_2$, 10 mM MES–KOH

pH 5.8 and 200 μM acetosyringone). After incubating 2 h at room temperature,

*Agrobacterium* cultures were infiltrated into the leaves of 4–week–old *N. benthamiana*

and *N. tabacum*. Leaves were harvested for analysis at two days post infiltration. The cell

death suppression assay in *N. benthamiana* was performed as described previously

(Wang et al., 2011).

Hairy Root Transformation in *Lotus japonicus*

The hairy root transformation assays were performed as described previously(Li et al.,

2018). In brief, surface-sterilized seeds were germinated and grown in the MS plate

without sucrose (23°C dark for first 3 days and 23°C 16 h light/8 h dark for next two

days). Then seeds were cut at the middle of the hypocotyl and co-cultivation with *A.*

*rhizogenes* carry pUB GFP and pUB NopT-FLAG GFP respectively (23°C dark for first

3 day and 23°C 16 h light/8 h dark for next 2 day). Transfer the plants onto B5 media.

After about 15 days, the plants were picked up for further analysis by cutting the root

without expressing florescent proteins. For rhizobial infection assays, data were collected

at 9dpi for *S.fredii* NGR234 and 6dpi for MAFF303099

NFR5 cytoplasmic domain cleavage assay

For cleavage assay in *E. coli*, Strep–NFR5$^{CD}$–HA or Strep–NFR5$^{CD}$–HA were co–

expressed with NopT–FLAG or NopT$^{C93S}$–FLAG by using Duet expressing vectors

(Novagen) (Han et al., 2017). Bacterial cell cultures were treated with 0.5 mM IPTG at

22°C for 15 h. The bacterial pellet was resuspended in extraction buffer consisting of

Tris–HCl (pH 7.3), 150 mM NaCl, 5 mM EDTA, 0.5% (w/v) SDS and 0.5% (v/v) Triton

X–100. After adding SDS loading buffer and boiling at 100°C for 5 min, total proteins

were detected with anti–HA–peroxidase (Sigma, clone 3F10), anti–FLAG (Sigma,

F1804), and anti–Strep (ABclonal, AB_2863792) antibodies.

For *in vitro* cleavage assays, Sumo–NFR5$^{JM}$–GFP and NopT–FLAG recombinant

proteins were purified, and 3 μg NopT–FLAG and 1 μg Sumo–NFR5$^{JM}$–GFP were

incubated in a test tube at 22°C for 7 h. The cleavage products were detected with anti–

GFP antibody by immunoblot (ABclonal, AB_2770402).

For in vivo cleavage assays, the transgenic root was collected at about 15 days after

induction of hairy roots. 2% protease inhibitors (Sigma, P9599) were added in extraction

buffer. The actin proteins were detected with anti-Actin antibody (ABclonal, AC009).

Phosphorylation assays

For phosphorylation assays in *E. coli*, His–NFR1$^{CD}$–Myc or His–NFR1$^{CD–K351E}$–Myc and

NopT–FLAG–His were co–expressed by using Duet expressing vectors (Novagen) with

0.5 mM IPTG at 28°C for 18 h. After purifying NFR1$^{CD}$ and NopT at the same time,

CIAP (TaKaRa, 2250B) was added and incubated at 37°C for 3 h. The mobility shift of

NFR1$^{CD}$ and NopT was detected with anti–Myc (Bio–Legend, 626808) and anti–FLAG

(Sigma, F1804) antibodies by immunoblot analysis.

For phosphorylation assays in *L. Japonicus* roots, NopT$^{C93S}$-FLAG was expressed in the

*nfr1* mutant and wild type plants, respectively. Transfer the plant with florescent root to

half–strength B&D media for 5 days. After treat with water and rhizobia respectively, the

total proteins were drawn by extraction buffer with addition 2% protease inhibitors

cocktail (Sigma, P9599) and 2% phosphatase inhibitors cocktail (Yeasen,20109-A), then

followed with TCA precipitation to remove various contaminants (e.g. EDTA, surfactant).

Phosphorylation of NopT$^{C93S}$ proteins was analyzed by 50μM Zn$^{2+}$-Phos-tag SDS-PAGE

and detected with anti–FLAG (Sigma, F1804) antibodies by immunoblot analysis.

BiFC and Split–LUC assays in *N. benthamiana*

In BiFC assay, the eYFP fluorescence of fusion proteins was assayed 2–3 days after

infiltration using a confocal microscope (Leica TCS SP8). For the Split–LUC assay, *N.*

*benthamiana* leaves were collected at 2 dpi sprayed with a solution of 1 mM D–luciferin

(Promega, E1603) and 0.02% (v/v) Triton X–100. After a dark–adaptation of 5 min,

bioluminescence images were acquired using a Tanon Bio–Imaging System (Tanon,

Shanghai, 4600).

Liquid chromatography mass spectrometry

For proteomics analysis of NopT phosphorylation sites, *NopT* and *NFR1–CD* or *NFR1–*

*CD^{K351E}* were co–expressed in *E. coli*, then proteins were purified using Ni–Charged

Resin (GenScript, Cat. No. L00223). The bands of NopT$^P$ and NopT, as control, were cut

and destained by 50 mM triethylammonium bicarbonate (TEAB) and 50% (v/v)

acetonitrile (ACN) in 50 mM TEAB followed by washing with 100% ACN until the gel

turned white. The protein bands were then in–gel digested with sequencing–grade trypsin.

LC–MS analysis was performed by Novogene Co., Ltd. (Beijing, China).

For identificaiton of the cleavage site on NFR5, NopT and the recombinant protein,

His-Sumo-NFR5-GST, were co-expressed in *E. coli* followed by protein purification

using Glutathione Resin (GenScript, Cat. No. L00206). About 80μg cleaved NFR5 was

collected for analysis. The protein bands were digested with trypsin,glu-C, chymotrypsin

and pepsin. LC–MS analysis was performed by Oulu Biotechnology Co., Ltd. (Shanghai,

China).

**Supplemental data**

**Supplemental Figure 1.** Analyses of the functions of all 15 effectors from *S. fredii*

NGR234 in preventing LjNFR1- and LjNFR5-induced cell death in *N. benthamiana*

leaves. Supplemental Fig. 2.   Split–LUC assay testing the interactions between different

NopT mutant and NFRs.

**Supplemental Figure 2.**   Split–LUC assay testing the interactions between different

NopT mutants and NFRs.

**Supplemental Figure 3.** NopT cleaves NFR5 at the juxtamembrane domain.

**Supplemental Figure 4.** Characterization of cleavage site of LysM receptor.

**Supplemental Figure 5.** The cytoplasmic domains (CDs) of NFR1, NFR5, and kinase–

dead NFR1CDK351E were used for kinase assays in E. coli.

**Supplemental Figure 6.** Mutant version of NFR5 fails to form infections.

**Supplemental Figure 7.** Phylogenetic tree based on the amino acid sequence of NopT

homologs from different bacterial species.

**Supplemental Figure 8.** Transient expression of NopT triggers cell death in Arabidopsis

and Nicotiana tabacum.

**Supplemental Table 1.** Phosphopeptides identified by Liquid Chromatography–Mass

spectrometry.

**Supplemental Table 2.** Primers used in this study.

**Acknowledgments:** We thank Drs. E. Giraud, E. Wang, and Y. Liang for their critical

reading of this manuscript. We also thank Drs. J. Staugaard and C. Staehelin for kindly

providing *pNIN:GUS* transgenic seed and *S. fredii* NGR234 *ΔnopT* strain, respectively;

Dr. J. Du for providing plasmids expressing of Avr3a and R3a, BAX, and INF1. The

qPCR and microscopy data were acquired from the Core Facility Center run by the

National Key Lab of Agricultural Microbiology.

**Funding:** This work was supported by the National Key R&D Program of China

(2019YFA0904700), the National Natural Science Foundation of China (32090063), and

the Fundamental Research Funds for the Central Universities (2662022SKYJ002). Work

in the Stacey's Lab was supported by a grant from National Science Foundation, Plant

Genome Research Project (IOS–1734145).

**Author Contributions:** H.B., Y.C., and G.S. conceived the idea and designed the

research, H.B., Y.W., Y.L., and Y.Y. performed experiments; H.B., Y.W., H. L., H.Z.,

and Y.C. analyzed the data, Q.W and S.X identified the cleavage site by MS. H.B., Y.C.,

and G.S. wrote the paper.

## References

**Akeda, Y., and Galán, J.E.** (2005). Chaperone release and unfolding of substrates in type III

secretion. Nature **437,** 911-915.

**Ausmees, N., Kobayashi, H., Deakin, W.J., Marie, C., Krishnan, H.B., Broughton, W.J., and

Perret, X.** (2004). Characterization of NopP, a type III secreted effector of sp. strain NGR234. J

Bacteriol **186,** 4774-4780.

**Bos, J.I., Kanneganti, T.D., Young, C., Cakir, C., Huitema, E., Win, J., Armstrong, M.R.,

Birch, P.R., and Kamoun, S.** (2006). The C-terminal half of *Phytophthora infestans* RXLR

effector AVR3a is sufficient to trigger R3a-mediated hypersensitivity and suppress INF1-induced

cell death in *Nicotiana benthamiana*. Plant J **48,** 165-176.

**Bozsoki, Z., Cheng, J., Feng, F., Gysel, K., Vinther, M., Andersen, K.R., Oldroyd, G., Blaise,

M., Radutoiu, S., and Stougaard, J.** (2017). Receptor-mediated chitin perception in legume

roots is functionally separable from Nod factor perception. Proc Natl Acad Sci U S A **114,**

E8118-E8127.

**Broghammer, A., Krusell, L., Blaise, M., Sauer, J., Sullivan, J.T., Maolanon, N., Vinther,

M., Lorentzen, A., Madsen, E.B., Jensen, K.J., Roepstorff, P., Thirup, S., Ronson, C.W.,

Thygesen, M.B., and Stougaard, J.** (2012). Legume receptors perceive the rhizobial lipochitin

oligosaccharide signal molecules by direct binding. Proc Natl Acad Sci U S A **109,** 13859-13864.

**Cao, Y., Liang, Y., Tanaka, K., Nguyen, C.T., Jedrzejczak, R.P., Joachimiak, A., and**

**Stacey, G.** (2014). The kinase LYK5 is a major chitin receptor in *Arabidopsis* and forms a

chitin-induced complex with related kinase CERK1. eLife **3**.e03766.

**Dai, W.J., Zeng, Y., Xie, Z.P., and Staehelin, C.** (2008). Symbiosis-promoting and deleterious

effects of NopT, a novel type 3 effector of *Rhizobium* sp. strain NGR234. J Bacteriol **190,**

5101-5110.

**Dowen, R.H., Engel, J.L., Shao, F., Ecker, J.R., and Dixon, J.E.** (2009). A family of bacterial

cysteine protease type III effectors utilizes acylation-dependent and -independent strategies to

localize to plasma membranes. J Biol Chem **284,** 15867-15879.

**Fotiadis, C.T., Dimou, M., Georgakopoulos, D.G., Katinakis, P., and Tampakaki, A.P.**

(2012). Functional characterization of NopT1 and NopT2, two type III effectors of

*Bradyrhizobium japonicum*. FEMS Microbiol Lett **327,** 66-77.

**Gan, Y., Zhang, L., Zhang, Z., Dong, S., Li, J., Wang, Y., and Zheng, X.** (2009). The LCB2

subunit of the sphingolip biosynthesis enzyme serine palmitoyltransferase can function as an

attenuator of the hypersensitive response and Bax-induced cell death. New Phytol **181,** 127-146.

**Geurts, R., Fedorova, E., and Bisseling, T.** (2005). Nod factor signaling genes and their

function in the early stages of infection. Curr Opin Plant Biol **8,** 346-352.

**Goehre, V., Spallek, T., Haeweker, H., Mersmann, S., Mentzel, T., Boller, T., de Torres, M.,**

**Mansfield, J.W., and Robatzek, S.** (2008). Plant pattern-recognition receptor FLS2 is directed

for degradation by the bacterial ubiquitin ligase AvrPtoB. Curr Biol **18,** 1824-1832.

**Gysel, K., Laursen, M., Thygesen, M.B., Lironi, D., Bozsoki, Z., Hjuler, C.T., Maolanon,**

**N.N., Cheng, J., Bjork, P.K., Vinther, M., Madsen, L.H., Rubsam, H., Muszynski, A.,**

**Ghodrati, A., Azadi, P., Sullivan, J.T., Ronson, C.W., Jensen, K.J., Blaise, M., Radutoiu, S.,**

**Stougaard, J., and Andersen, K.R.** (2021). Kinetic proofreading of lipochitooligosaccharides

determines signal activation of symbiotic plant receptors. Proc Natl Acad Sci U S A

**118**.e2111031118.

**Han, Y., Sun, J., Yang, J., Tan, Z., Luo, J., and Lu, D.** (2017). Reconstitution of the plant

ubiquitination cascade in bacteria using a synthetic biology approach. Plant J **91,** 766-776.

**Heckmann, A.B., Sandal, N., Bek, A.S., Madsen, L.H., Jurkiewicz, A., Nielsen, M.W.,**

**Tirichine, L., and Stougaard, J.** (2011). Cytokinin induction of root nodule primordia in *Lotus*

*japonicus* is regulated by a mechanism operating in the root cortex. Mol Plant Microbe Interact

**24,** 1385-1395.

**Jones, J.D.G., and Dangl, J.L.** (2006). The plant immune system. Nature **444,** 323-329.

**Kawaharada, Y., Kelly, S., Nielsen, M.W., Hjuler, C.T., Gysel, K., Muszynski, A., Carlson,**

**R.W., Thygesen, M.B., Sandal, N., Asmussen, M.H., Vinther, M., Andersen, S.U., Krusell,**

**L., Thirup, S., Jensen, K.J., Ronson, C.W., Blaise, M., Radutoiu, S., and Stougaard, J.**

(2015). Receptor-mediated exopolysaccharide perception controls bacterial infection. Nature **523,**

308-312.

**Khan, A., Wadood, S.F., Chen, M., Wang, Y., Xie, Z.P., and Staehelin, C.** (2022).

Effector-triggered inhibition of nodulation: A rhizobial effector protease targets soybean kinase

GmPBS1-1. Plant Physiol **189,** 2382-2395.

**Kim, S.H., Qi, D., Ashfield, T., Helm, M., and Innes, R.W.** (2016). Using decoys to expand the

recognition specificity of a plant disease resistance protein. Science **351,** 684-687.

**Kimbrel, J.A., Thomas, W.J., Jiang, Y., Creason, A.L., Thireault, C.A., Sachs, J.L., and**

**Chang, J.H.** (2013). Mutualistic co-evolution of type III effector genes in Sinorhizobium fredii

and *Bradyrhizobium japonicum*. PLoS Pathog **9,** e1003204.

**Kopan, R., and Ilagan, M.X.G.** (2009). The canonical notch signaling pathway: Unfolding the

activation mechanism. Cell **137,** 216-233.

**Lei, L., Stevens, D.M., and Coaker, G.** (2020). Phosphorylation of the *Pseudomonas* effector

AvrPtoB by *Arabidopsis* SnRK2.8 is required for bacterial virulence. Mol Plant **13,** 1513-1522.

**Li, D.D., Zhu, Z.K., Deng, X.M., Zou, J.N., Ma, C., Li, C.D., Yin, T., Liu, C.Y., Wang, J.H.,**

**Chen, Q.S., and Xin, D.W.** (2023). GmPBS1, a hub gene interacting with Rhizobial type-III

effectors NopT and NopP, regulates soybean nodulation. Agronomy **13**.1242.

**Li, H., Chen, M., Duan, L., Zhang, T., Cao, Y., and Zhang, Z.** (2018). Domain swap approach

reveals the Ccritical roles of different domains of SYMRK in root nodule symbiosis in *Lotus*

*japonicus*. Front Plant Sci **9,** 697.

**Li, L., Kim, P., Yu, L., Cai, G., Chen, S., Alfano, J.R., and Zhou, J.M.** (2016).

Activation-dependent destruction of a co-receptor by a *Pseudomonas syringae* effector dampens

plant immunity. Cell Host Microbe **20,** 504-514.

**Madsen, E.B., Madsen, L.H., Radutoiu, S., Olbryt, M., Rakwalska, M., Szczyglowski, K.,**

**Sato, S., Kaneko, T., Tabata, S., Sandal, N., and Stougaard, J.** (2003). A receptor kinase gene

of the LysM type is involved in legume perception of rhizobial signals. Nature **425,** 637-640.

**Madsen, E.B., Antolin-Llovera, M., Grossmann, C., Ye, J., Vieweg, S., Broghammer, A.,**

**Krusell, L., Radutoiu, S., Jensen, O.N., Stougaard, J., and Parniske, M.** (2011).

Autophosphorylation is essential for the *in vivo* function of the *Lotus japonicus* Nod factor

receptor 1 and receptor-mediated signalling in cooperation with Nod factor receptor 5. Plant J **65,**

404-417.

**Moling, S., Pietraszewska-Bogiel, A., Postma, M., Fedorova, E., Hink, M.A., Limpens, E.,**

**Gadella, T.W., and Bisseling, T.** (2014). Nod factor receptors form heteromeric complexes and

are essential for intracellular infection in medicago nodules. Plant Cell **26,** 4188-4199.

**Nakagawa, T., Suzuki, T., Murata, S., Nakamura, S., Hino, T., Maeo, K., Tabata, R.,**

**Kawai, T., Tanaka, K., Niwa, Y., Watanabe, Y., Nakamura, K., Kimura, T., and Ishiguro,**

**S.** (2007). Improved Gateway binary vectors: high-performance vectors for creation of fusion

constructs in transgenic analysis of plants. Biosci Biotechnol Biochem **71,** 2095-2100.

**Ntoukakis, V., Mucyn, T.S., Gimenez-Ibanez, S., Chapman, H.C., Gutierrez, J.R., Balmuth,**

**A.L., Jones, A.M., and Rathjen, J.P.** (2009). Host inhibition of a bacterial virulence effector

triggers immunity to infection. Science **324,** 784-787.

**Schmidt, G.** (2011). Yersinia enterocolitica outer protein T (YopT). Eur J Cell Biol **90,** 955-958.

**Schubert, K.A., Xu, Y., Shao, F., and Auerbuch, V.** (2020). The type III secretion system as a

tool for studying cytosolic innate immune surveillance. Ann Rev Microbiol, Vol 74, 2020 **74,**

221-245.

**Shao, F., Merritt, P.M., Bao, Z., Innes, R.W., and Dixon, J.E.** (2002). A *Yersinia* effector and

a *Pseudomonas* avirulence protein define a family of cysteine proteases functioning in bacterial

pathogenesis. Cell **109,** 575-588.

**Shao, F., Vacratsis, P.O., Bao, Z., Bowers, K.E., Fierke, C.A., and Dixon, J.E.** (2003a).

Biochemical characterization of the *Yersinia* YopT protease: cleavage site and recognition

elements in Rho GTPases. Proc Natl Acad Sci U S A **100,** 904-909.

**Shao, F., Golstein, C., Ade, J., Stoutemyer, M., Dixon, J.E., and Innes, R.W.** (2003b).

Cleavage of *Arabidopsis* PBS1 by a bacterial type III effector. Science **301,** 1230-1233.

**Sugawara, M., Takahashi, S., Umehara, Y., Iwano, H., Tsurumaru, H., Odake, H., Suzuki,**

**Y., Kondo, H., Konno, Y., Yamakawa, T., Sato, S., Mitsui, H., and Minamisawa, K.** (2018).

Variation in bradyrhizobial NopP effector determines symbiotic incompatibility with soybeans

via effector-triggered immunity. Nature Commun **9**.3139.

**Sun, Y., Li, L., Macho, A.P., Han, Z., Hu, Z., Zipfel, C., Zhou, J.-M., and Chai, J.** (2013).

Structural basis for flg22-induced activation of the *Arabidopsis* FLS2-BAK1 immune complex.

Science **342,** 624-628.

**Tang, D.Z., Wang, G.X., and Zhou, J.M.** (2017). Receptor kinases in plant-pathogen

interactions: More than pattern recognition. Plant Cell **29,** 618-637.

**Teulet, A., Busset, N., Fardoux, J., Gully, D., Chaintreuil, C., Cartieaux, F., Jauneau, A.,**

**Comorge, V., Okazaki, S., Kaneko, T., Gressent, F., Nouwen, N., Arrighi, J.F., Koebnik, R.,**

**Mergaert, P., Deslandes, L., and Giraud, E.** (2019). The rhizobial type III effector ErnA

confers the ability to form nodules in legumes. Proc Natl Acad Sci U S A **116,** 21758-21768.

**Vleeshouwers, V.G., Driesprong, J.D., Kamphuis, L.G., Torto-Alalibo, T., Van't Slot, K.A.,**

**Govers, F., Visser, R.G., Jacobsen, E., and Kamoun, S.** (2006). Agroinfection-based

high-throughput screening reveals specific recognition of INF elicitins in *Solanum*. Mol Plant

Pathol **7,** 499-510.

**Wang, D., Yang, S., Tang, F., and Zhu, H.** (2012). Symbiosis specificity in the legume:

rhizobial mutualism. Cell Microbiol **14,** 334-342.

**Wang, Q., Han, C., Ferreira, A.O., Yu, X., Ye, W., Tripathy, S., Kale, S.D., Gu, B., Sheng,**

**Y., Sui, Y., Wang, X., Zhang, Z., Cheng, B., Dong, S., Shan, W., Zheng, X., Dou, D., Tyler,**

**B.M., and Wang, Y.** (2011). Transcriptional programming and functional interactions within the

*Phytophthora sojae* RXLR effector repertoire. Plant Cell **23,** 2064-2086.

**Xiang, T., Zong, N., Zou, Y., Wu, Y., Zhang, J., Xing, W., Li, Y., Tang, X., Zhu, L., Chai, J.,**

**and Zhou, J.M.** (2008). *Pseudomonas syringae* effector AvrPto blocks innate immunity by

targeting receptor kinases. Curr Biol **18,** 74-80.

**Xu, N., Luo, X., Wu, W., Xing, Y., Liang, Y., Liu, Y., Zou, H., Wei, H.L., and Liu, J.** (2020).

A plant lectin receptor-like kinase phosphorylates the bacterial effector AvrPtoB to dampen Iits

virulence in *Arabidopsis*. Mol Plant **13,** 1499-1512.

**Yang, J., Lan, L., Jin, Y., Yu, N., Wang, D., and Wang, E.** (2022). Mechanisms underlying

legume-rhizobium symbioses. J Integr Plant Biol **64,** 244-267.

**Zhang, B., Wang, M., Sun, Y., Zhao, P., Liu, C., Qing, K., Hu, X., Zhong, Z., Cheng, J.,**

**Wang, H., Peng, Y., Shi, J., Zhuang, L., Du, S., He, M., Wu, H., Liu, M., Chen, S., Wang, H.,**

**Chen, X., Fan, W., Tian, K., Wang, Y., Chen, Q., Wang, S., Dong, F., Yang, C., Zhang, M.,**

**Song, Q., Li, Y., and Wang, X.** (2021). *Glycine max* NNL1 restricts symbiotic compatibility

with widely distributed *Bradyrhizobia vi*a root hair infection. Nat Plants **7,** 73-86.

**Zhang, J., Li, W., Xiang, T., Liu, Z., Laluk, K., Ding, X., Zou, Y., Gao, M., Zhang, X., Chen,**

**S., Mengiste, T., Zhang, Y., and Zhou, J.M.** (2010). Receptor-like cytoplasmic kinases

integrate signaling from multiple plant immune receptors and are targeted by a *Pseudomonas*

*syringae* effector. Cell Host Microbe **7,** 290-301.

**Zhou, J.G., Wang, P., Claus, L.A.N., Savatin, D.V., Xu, G.Y., Wu, S.J., Meng, X.Z.,**

**Russinova, E., He, P., and Shan, L.B.** (2019). Proteolytic orocessing of SERK3/BAK1 regulates

plant immunity, development, and cell death. Plant Physiol **180,** 543-558.

**Zipfel, C., and Oldroyd, G.E.D.** (2017). Plant signalling in symbiosis and immunity. Nature

**543,** 328-336.

**Zipfel, C., Robatzek, S., Navarro, L., Oakeley, E.J., Jones, J.D.G., Felix, G., and Boller, T.**

(2004). Bacterial disease resistance in through flagellin perception. Nature **428,** 764-767.

**Figures**

**Figure 1. NopT specifically suppresses cell death triggered by NFR1 & NFR5 in** *N.*

**benthamiana.** *Agrobacterium* strains harboring genes were NopT or EV infiltrated into *N.*

*benthamiana*,12 hpi followed by infiltration with *agrobacterium* containing NFR1 and

NFR5. Different leaf discs indicate where different proteins were expressed. NopT (A)

but not NopT$^{C93S}$, the protease–inactive version of NopT (B) suppressed cell death

induced by expression of NFR1 and NFR5. (C) Expression of NopM induced cell death

in *N. benthamiana*. (D) Expression of NopT could not suppress the cell death triggered

by expression of Avr3a&R3a, NopM, BAX, INF1 or AtCERK1. Digital numbers in (D)

represent the number of leaf discs with cell death and total leaf discs tested. All pictures

in the Figure are representative of at least three independent biological replicates.

**Figure 2. NopT interacts with NFR1 and NFR5.** Interactions between NopT and NFR1

and NFR5 were detected using Split–YFP assay (A) and (B), Split–LUC (C) and co–IP

(D) assays in *N. benthamiana*. (A) and (B). nYFP and cYFP tags were fused at the C–

terminus of NopT, NFR1, NFR5, and negative control FLS2. YFP fluorescence signals

represent protein–protein interactions. Scar bar=25 um. (C). nLUC and cLUC tags were

fused at the C–terminus of NopT, NFR1, NFR5, and negative control FLS2.

Luminescence signals represent protein–protein interactions. (D). HA–tagged NFR1 and

NFR5 and FLAG–tagged NopT were expressed in plant cells followed by

immunoprecipitation using anti–FLAG antibody and determined by immunoblot using

anti–HA and anti–FLAG antibodies. All data in Figure are representative of three

biological replicates.

**Figure 3. NopT proteolyzes NFR5 at its juxtamembrane domain.** (A) NopT but not

NopT$^{C93S}$ cleaves NFR5–GFP protein expressed in N. benthamiana cells by releasing

NFR5CD–GFP. (B) NopT but not NopT$^{C93S}$ cleaves NFR5-Myc protein expressed in *L.*

*japonicus* transgenic root by releasing NFR5CD-Myc. Asterisk indicated the cleaved

variant of NFR5. (C) A repeated experiment testing the cleavage of NFR5$^{CD}$ but not

NFR1$^{CD}$ fused with different tags. Asterisk represents the phosphorylated NopT by

inducing a band retardation on SDS–PAGE gel. (D) NopT but not NopT$^{C93S}$ cleaves the

CD of NFR5 protein expressed in *E. coli* cells by releasing the kinase domain of NFR5.

Asterisk indicated the cleaved variant of NFR5. (E) NopT cleaves a recombinant protein

where His–tagged SUMO and GFP is bridged with the JM domain of NFR5. F.T.

represents flow through sample after Ni–beads purification. (F) NopT could not cleave a

mutant version of NFR5CD–HA with 5 residues in the JM substituted into other residues

(S283Y, G294Q, Y303S, A310I, T311Y). (G) NopT could not cleave *Medicago*

*truncatula* NFP and from NFR5$^{JM}$–NFP$^{KD}$. CD represents cytoplasmic domain, KC

represents kinase domain and C-terminal tail region, NopT$^{C}$ indicates the truncated

version of NopT after autocleavage by releasing about 50 a.a. at its N–terminus.

**Figure 4. Phosphorylation of NopT by NFR1 suppresses its proteolytic activity.** (A) *In vivo* phosphorylation assay in *L. japonicus* roots. Phosphorylation of NopT$^{C93S}$ was induced with rhizobia treatment and dependent on *nfr1*. (B) NFR1$^{CD}$ phosphorylates NopT by inducing a gel band shift determined by immunoblotting. (C) The phoshorylation sites on NopT identified by liquid chromatography–mass spectrometry (LC–MS) were either substituted into alaine (A) or asparatate (D) and subsequently tested for their ability to proteolytically cleave NFR5$^{CD}$.

**Figure 5. NopT regulates rhizobial infection in *L. japonicus*.** (A) Rhizobial infection in

*L. japonicus* using GFP-labelled WT and *nopt* mutant strains o*f S. fredii* NGR234.

Fluorescent, GFP-expressing bacteria are imaged in Cyan. Scar bars=12.5µm. (B) GUS

staining pictures from the roots of *L. japonicus* transgenically expressing GUS under

*p*NIN promoter inoculated with WT and *nopT* mutant strains of NGR234. Scar

bars=1mm (C) Statistical analysis of rhizobial infection in A (n=10, Student's *t*–test: P <

0.01). (D) Statistical analysis of GUS straining sites in B (n=10, Student's *t*–test: P <

0.01). (E) Nodule primordia number on the roots 14dpi after inoculation WT or *nopT*

mutant strains (n>20, Student's *t*–test: P < 0.01). (F) Infection foci number from the wild

type plant roots inoculated with WT strains and *nopt* mutant strains expressing NopT

with different variations (n=5, Student's *t*–test: P < 0.01). (G) Statistical analysis of

rhizobial infection in *L. japonicus* transgenic root expressing GFP (EV, empty vector

control) or NopT using DsRed-labeled *M. loti* MAFF303099 (n=8, Student's t–test: P <

0.01).

**Figure 6. Working model of NopT in association with NF receptors.** (A) NopT$_{234}$,

NopT from *S. fredii* NGR234; NopT1$_{110}$ and NopT2$_{110}$, NopT homologs from *B.*

*diazoefficiens* USDA110; NopT$_{257}$, NopT from *S. fredii* USDA257; NopT$_{103}$, NopT from

*S. fredii* HH103. All NopT homologs were tested in the proteolysis assay using NFR5$^{CD}$

as a target. (B) Rhizobial infection $NopT_{257}$ could not inhibit the cell death triggered by

NFR1 and NFR5 in *N. benthamiana*. (C)The indicated rhizobial strains induced *nin::gus*

expressing in *Lotus* root. Scar bars=2.5mm. (D) Statistical analysis of GUS straining in C

(n=19, Student's *t*–test: P < 0.01). (E) A proposed model of NopT in association with NF

receptors. Both NopT and $NopT^C$ after autocleavage exert proteolytic activities to cleave

NFR5 at the juxtamembrane domain to release the kinase domain of NFR5 (cleaved

NFR5). NFR1 phosphorylates intact NopT, but not $NopT^C$, to block its proteinase

activity.

**Supplemental Fig. 1.    Analyses of the functions of all 15 effectors from *S. fredii***

**NGR234 in preventing LjNFR1- and LjNFR5-induced cell death in *N. benthamiana***

**leaves.**

15 genes from *S. fredii* NGR234 encoding effector proteins were amplified and cloned

into binary vector for expression of Strep-tagged proteins in *N. benthamiana*. LjNFR1

and LjNFR5 from *L. japonicus* were C-terminally tagged with HA and Myc, respectively,

and cloned into binary vectors for expression in *N. benthamiana*. All plasmids were

electroporated into *Agrobacterium tumefaciens* EH105.

(A) *Agrobacterium* strain harboring each effector gene or control empty vector (EV)

were infiltrated into *N. benthamiana*. 12 hour post infiltration (hpi), agrobacterium strains

harboring LjNFR1 and LjNFR5 were mixed and hand-infiltrated in the same leaf discs as

shown in (B). 48 hpi, cell death phenotypes were evaluated (C). At least three replicates

in the same leaf were performed to test the function of each effector in suppressing cell

death-triggered by LjNFR1 and LjNFR5. (D) Results of all 15 effector proteins in

suppressing programmed cell death by overexpression of NFR1 and NFR5. (E)

Abundance of LjNFR1, LjNFR5, and NopT proteins, as measured by immunoblot with

specific antibodies. Leaf discs expressing LjNFR1, LjNFR5, and NopT or LjNFR1 and

LjNFR5 and with EV were used for immunoblot analyses. Actin was used as loading

control. (F) Another example of suppression of programmed cell death by NopT in the

leaf discs coexpressing NFR1 and NFR5. Rep1, Rep2, and Rep3 represent three technical

repeats in the same leave.

1013

1014

1015

1016

1017
1018

1019

1020

**Supplemental Fig. 2. Split–LUC assay testing the interactions between different NopT mutants and NFRs.**

(A)The interaction between NopT with mutated acylation site and NFR1/NFR5. $NopT^{AAA}$ (G50A C51A C52A). (B) The protein expressing of indicating genes in the leave of (A). (C) $Nop^{C93S}$ interact with NFR1 and NFR5. (D) The interaction between $NopT^{\Delta N50}$ and NFR5. $NopT^{\Delta N50}$ represents a truncated NopT with deletion of about 50 a.a. from its N–terminus.

1028

1029

1030

1031

1032

1033

**Supplemental Fig. 3. NopT cleaves NFR5 at the juxtamembrane domain.**

(A) NopT cannot cleave NFR1-GFP expressed in *N. benthamiana* leaves. recombinant

Sumo-NFR5$^{KD}$-HA lacking the juxtamembrane (JM) domain. (B) NopT could not cleave

Sumo–NFR5$^{KD}$–HA, a recombinant protein expressed in *E. coli* with the juxtamembrane

domain of NFR5 was replaced with Sumo tag. (C) Cleavage assay using two mutant

versions of NFR5$^{CD}$ in the presence of NopT or AvrPphB. NFR5$^{CD288-294A}$ represents a

mutant NFR5$^{CD}$ with residues 288 to 294 replaced with seven alanines; NFR5$^{CD288-294PphB}$

represents a mutant NFR5$^{CD}$ with residues 288 to 294 replaced with AvrPphB recognition

sites. (D) Autocleavage assay, using different NopT variants with single amino acid

mutations. H205A and D220A indicate His-205 and Asp-220 replaced with alanine,

respectively. (E-H) NopT cleavage assay using 19 variants of NFR5$^{CD}$ with three

adjacent amino acids replaced with three alanines. The numbers indicate the location of

the three residues replaced with alanines. (I) NopT cleavage assay using seven truncated

variants of NFR5$^{CD}$ with 10 adjacent amino acids deleted. The numbers indicate the

location of the 10 deleted residues. (J) NopT but not NopT$^{C93S}$ could cleave the CDs of

AtLYK5 and LjLYS11 expressed in E. coli. NopT could cleave both AtLYK5$^{JM}$–

NFR5$^{KC}$, LjLYS11$^{JM}$–NFR5$^{KC}$ and LjNFR1$^{JM}$–NFR5$^{KC}$, three recombinant proteins

expressed in *E. coli* with the JM of NFR5 was replaced with the JMs from AtLYK5,

LjLYS11 and NFR1, respectively. (K) NopT but not NopT$^{C93S}$ could cleave recombinant

protein NFR5$^{268-445}$-NFP$^{458-595}$ and NFP$^{270-457}$-NFR$^{5456-595}$.

1054

**Supplemental Fig. 4. Characterization of cleavage site of LysM receptor.**

(A) Domain and motif annotation of NFR5. (B) Protein sequence alignment of the
cytoplasmic domain of AtLYK5, LjLYS11, LjNFR5 and MtNFP. Yellow boxes
delineated highly conserved residues (S283, G294, Y303, A310, T311) and brown box
delineated kinase domain. Green frame delineated cleavage site of NFR5 and red frame
delineated residues similar to NopT autocleavage region. (C) Mass spectrometry analysis
of cleaved NFR5. Blue lines indicated the peptides characterized by MS.

1064

1065

**Supplemental Fig. 5. The cytoplasmic domains (CDs) of NFR1, NFR5, and kinase–dead NFR1CD$^{\text{K351E}}$ were used for kinase assays in *E. coli*.** Proteins were detected by immunoblotting using anti–HA and anti–Myc antibodies. Asterisk indicated the band retardation on the gel representing the phosphorylated NFR5$^{\text{CD}}$ proteins.

1070

1071

**Supplemental Fig. 6. Rhizobial infection in the mutant versions of NFR5.**

(A) NFR5$^{5m}$ failed to form rhizobial infection (n=7, Student's t–test: P < 0.01).

(B)NFR5-NFP$^{KD-CT}$ could recuse the rhizobial infection of wild type NGR234 (n=10,

Student's t–test: P < 0.01). The transgenic roots expressing NFR5 or NFR5$^{5m}$ or

NFR5-NFP$^{KCT}$ in nfr5-3 background under the control of native promoter were

inoculated with GFP-labelled wild type NGR234 and NGR234$\Delta nopT$ respectively. The

data was collected 9 days post inoculation.

1079

Sinorhizobium fredii HH103
Sinorhizobium fredii USDA257
Sinorhizobium fredii CCBAU45436
NopT Sinorhizobium fredii NGR234
Ensifer sp. MPMI2T
Bradyrhizobium sp. Ec3.3
Bradyrhizobium sp. NAS96.2
Bradyrhizobium sp. 323S2
Bradyrhizobium sp. CCBAU051011
2 Ensifer sp. MPMI2T
Sinorhizobium sp. BJ1
Sinorhizobium sp. PC2
NopT2 Bradyrhizobium diazoefficiens USDA110
Bradyrhizobium elkanii USDA61
Bradyrhizobium sp. CCBAU53421
Bradyrhizobium sp. CCBAU51765
NopT Bradyrhizobium sp. ORS3257
Bradyrhizobium sp. CIR3A
Bradyrhizobium sp. Rc2d
Bradyrhizobium sp. 2S1
Bradyrhizobium sp. CCBAU51753
Bradyrhizobium sp. CCBAU43298
NopT1 Bradyrhizobium diazoefficiens USDA110
Bradyrhizobium sp. AC87j1
Bradyrhizobium sp. IAR9
Bradyrhizobium sp. CIR18
Bradyrhizobium sp. CIR48
Acidovorax citrulli
Acidovorax avenae
Acidovorax oryzae
Pseudomonas savastanoi pv. phaseolicola
Pseudomonas syringae pv. actinidiae ICMP 9853
Pseudomonas tremae
Ralstonia sp. A12
Xanthomonas oryzae
Xanthomonas bromi
Xanthomonas translucens pv. graminis ART-Xtg29
Xanthomonas translucens pv. graminis
Pseudomonas syringae
Pseudomonas syringae pv. daphniphylli
Pseudomonas amygdali pv. morsprunorum
Pseudomonas caricapapayae
Pantoea agglomerans
Pseudomonas sp. KBS0707
Pseudomonas savastanoi pv. glycinea
Pseudomonas savastanoi pv. nerii
Pseudomonas meliae
Pseudomonas ficuserectae
Pseudomonas amygdali pv. sesami
Pseudomonas cannabina
Ralstonia_solanacearum
Ralstonia pseudosolanacearum
Pasteurella multocida
Vibrio sp. ED002
Vibrio jasicida
Vibrio harveyi
Yersinia enterocolitica
Yersinia pseudotuberculosis
Yersinia pestis
Photorhabdus laumondii subsp. laumondii
Photorhabdus laumondii
Photorhabdus temperata
Photorhabdus sp. APURE
Photorhabdus heterorhabditis
Photorhabdus cinerea
Photorhabdus asymbiotica
Photorhabdus akhurstii
Photorhabdus hindustanensis
Photorhabdus luminescens

0.50

1080

1081 **Supplemental Fig. 7. Phylogenetic tree based on the amino acid sequence of NopT**

1082 **homologs from different bacterial species.**

1083

1084

1085

1086

1087

1088

1089

1090

1091

1092

1093

**Supplemental Fig. 8. Transient expression of NopT triggers cell death in *Arabidopsis***

***thaliana* and *Nicotiana tabacum*.**

(A) *Pseudomonas syringae* pv tomato DC3000 harboring control plasmids (left) or

expressing NopT (right) were infiltrated into Arabidopsis leaves. Note the cell death in

the right-side panel. (B) *Agrobacterium* strains harboring *NopT*, *NopT* with an N-terminal

50–amino acid deletion, *NopT*$^{C93S}$, or *NopT*$^{C93S}$ with an N-terminal 50–amino acid

deletion were infiltrated into *N. tabacum* leaves.

1101

1102 **Supplemental Table 1.   Phosphopeptides identified by liquid chromatography–**

1103 **mass spectrometry.** An *in vitro* kinase assay was performed using the CD of NFR1 and

1104 NopT. NopT was separated by SDS-PAGE, and the digested gel slices representing the

1105 phosphorylated NopT were used for analysis. The deduced phosphopeptides are listed.

| Sequence | Modifications | SITE | PSMs_NFR1KE+NopT | PSMs_NFR1+NopT |
|---|---|---|---|---|
| ETLANVELRTK | 1xPhospho [T10] | 37T | - | 1 |
| ETLANVELR | 1xPhospho [T2] | 29T | - | 1 |
| ETLANVELRTKSPSAECPDK | 1xCarbamidomethyl [C17]; 1xPhospho [T10] | 37T | - | 1 |
| TKSPSAECPDK | 1xCarbamidomethyl [C8]; 1xPhospho [S5] | 41S | - | 1 |
| SPSAECPDK | 1xCarbamidomethyl [C6]; 1xPhospho [S3] | 41S | | 1 |
| SPSAECPDKMGCCASKPQASDPNNPSTSSPARPSTSLFR | 3xCarbamidomethyl [C6; C12; C13]; 1xOxidation [M10] | | 1 | 1 |
| SPSAECPDKMGCCASKPQASDPNNPSTSSPARPSTSLFR | 3xCarbamidomethyl [C6; C12; C13]; 1xPhospho [S/T]; 1xOxidation [M10] | | - | 4 |
| MGCCASKPQASDPNNPSTSSPARPSTSLFR | 2xCarbamidomethyl [C3; C4]; 1xPhospho [T/S]; 1xOxidation [M1] | | - | 11 |
| MGCCASKPQASDPNNPSTSSPARPSTSLFR | 2xCarbamidomethyl [C3; C4]; 1xPhospho [T/S] | | - | 9 |
| MGCCASKPQASDPNNPSTSSPARPSTSLFR | 2xCarbamidomethyl [C3; C4]; 2xPhospho [T/S] | 65S 66T | - | 4 |
| MGCCASKPQASDPNNPSTSSPARPSTSLFR | 2xCarbamidomethyl [C3; C4]; 1xOxidation [M1] | | 4 | 2 |
| MGCCASKPQASDPNNPSTSSPARPSTSLFR | 2xCarbamidomethyl [C3; C4]; 2xPhospho [T/S]; 1xOxidation [M1] | | - | 4 |
| TAELAQANADGICVGLTAEWLR | 1xCarbamidomethyl [C13]; 1xPhospho [T17] | 97T | - | 10 |
| NLNSHPSIR | 1xPhospho [S7] | 109S | - | 2 |
| NLNSHPSIRMEALVPGSQR | 1xPhospho [S7]; 1xOxidation [M10] | 119S | - | 1 |
| MEALVPGSQR | 1xPhospho [S8] | 119S | - | 1 |
| MEALVPGSQR | 1xPhospho [S8]; 1xOxidation [M1] | 119S | - | 1 |
| RQGAGPSEADFAAQNTMLQK | 1xPhospho [T/S]; 1xOxidation [M17] | | - | 3 |
| RQGAGPSEADFAAQNTMLQK | 1xPhospho [T16] | 156T | - | 2 |
| QGAGPSEADFAAQNTMLQK | 1xPhospho [T/S]; 1xOxidation [M16] | | - | 3 |
| QGAGPSEADFAAQNTMLQK | 1xPhospho [T15] | 156T | - | 2 |
| SLANRYSNPNR | 1xPhospho [S7] | 245S | - | 2 |
| QHLTTVTTQK | 1xPhospho [T8] | 257T | - | 1 |

1106

1107

1108

1109

1110

1111

1112

1113

1114

1115

1116

1117

1118

1119 **Supplemental Table 2. Primers used in this study.**

| Used for | Target genes | Names | Sequences (from 5' to 3') |
|---|---|---|---|
| Overexpression in Nicotiana benthamiana | NopT or NopT$^{C93S}$ | p5X-NopT-F | agaacacgggggactctagaatgcacagtcccatcag |
| | | p5X-NopT-R | ccgctgttatcggtacctgtcatcttttgggtggtcac |
| | NopT$^{\Delta N50}$ | p5X-NopTN50-F | agaacacgggggactctagaatgtgctgcgccagcaag |
| | NFR1 | p5X-NFR1-F | agaacacgggggactctagaatgaagctaaaaactggtctact |
| | | p5X-NFR1-R | ccgctgttatcggtacctcttctcacagac |
| | NFR5 | p5X-NFR5-F | agaacacgggggactctagaatggctgtcttctttcttacc |
| | | p5X-NFR5-R | ccgctgttatcggtaccacgtgcagtaat |
| | | p5X-NFR5gfp-R | tcccgggagcggtaccacgtgcagtaatggaagt |
| | AvrPphB | p5X-AvrPphB-F | agaacacgggggactctagaatgaaaataggtacgcaggcc |
| | | p5X-AvrPphB-R | ccgctgttatcggtacccgaaactctaaactcgtttacgc |
| templates for protein expression | NopT$_{USDA257}$ | p5X-257T-F | agaacacgggggactctagatgctgcgtgatccgaacaat |
| | | p5X-257T-R | ccgctgttatcggtacctgtcaccttttgggtggtcaccg |
| | NopT1$_{USDA110}$ | p5X-110T1-F | agaacacgggggactctagaatgtatgatcgaatcggtgg |
| | | p5X-110T1-R | ccgctgttatcggtaccctgcatcctttgcgtcg |
| | NopT2$_{USDA110}$ | p5X-110T2-F | agaacacgggggactctagaatgtataatcgagtcgatggc |
| | | p5X-110T2-R | ccgctgttatcggtaccccgatgaggttccgc |
| Protein Expression in *E.coli* | NopT or NopT$^{C93S}$ | 28a NopT-F | tgggtcgcggatccgaattcatgcacagtcccatcag |
| | | 28a FLAG-R | tcgagtgcggccgcaagcttcttgtcatcgtcatccttgtag |
| | | 28a Strep-R | tcgagtgcggccgcaagcttttttttcaaattgaggatgagaccatcc |
| | AvrPphB | ET28a-pphb-F | tgggtcgcggatccgaattcaaaataggtacgcaggccac |
| | NopT$_{USDA257}$ | ET28a-257T-F | tgggtcgcggatccgaattcctgcgtgatccgaacaa |
| | NopT1$_{USDA110}$ | ET28a-110T1-F | tgggtcgcggatccgaattctatgatcgaatcggtggctc |
| | NopT2$_{USDA110}$ | ET28a-110T2-F | tgggtcgcggatccgaattctataatcgagtcgatggcgaatac |
| | NFR1CD | duet NFR1-F | accacagccaggatccgagataccagaagaaggaagaagagaaagc |
| | | duet Sac1nost-R | ggcgcgccgagctcgcgatcggggaaattcgagct |
| | | MAL-NFR1-F | ggaaggatttcagaattcggatccagataccagaagaaggaagaagag |
| | | MAL-NFR1-R | cgacggccagtgccaagctttcatcttctcacagacagtaaatttatgag |
| | NFR5CD | ACSH-NFR5-F | agaagcgcggatccgaattctatgtatactgccgcagaaagaag |
| | | ACSH-NFR5-R | cgtatgggtagcttccaagcttacgtgcagtaatggaagtc |
| | | MAL-NFR5-F | ggaaggatttcagaattcggatcctatgtatactgccgcagaaagaag |
| | | MAL-NFR5-R | cgacggccagtgccaagctttcaacgtgcagtaatggaagtc |
| | NFR5KD | SUMO 5KD-F | ccgcgaacagattggtaaggttggggaatcagtgtac |
| | | SUMO HA-R | gctcgaattcggatcctttcacgcatagtcaggaacatcg |
| | NFR5JM | SUMO 5JM-F | ccgcgaacagattggttatgtatactgccgcagaaagaag |
| | | SUMO 5JM-R | gctcgaattcggatcctttcacgcatagtcaggaacatcg |
| | sumo-NFR5KD-HA | duet sumo-F | catcaccacagccaggatgggtccctgcaggac |
| | | duet HA-R | cattatgcggccgcaagcttttacgcatagtcaggaacatcg |
| | sumo-NFR5JM-GFP | duet sumo-F | catcaccacagccaggatgggtccctgcaggac |
| | | duet GFP-R | atgcggccgcaagcttttacttgtacaactcatccatacc |

| Category | Gene | Primer | Sequence |
|---|---|---|---|
| | AtLYK5 | ACSH-LYK5-F | gcgcggatccgaattctacaaacgaaggtctaagaag |
| | | ACSH-LYK5-R | ggtagcttccaagcttgttgccaagagagccg |
| | LjLYS11 | ACSH-LYS11-F | agaagcgcggatccgaattcacttgtctgaggaagagaaag |
| | | ACSH-LYS11-R | cgtatgggtagcttccaagcttacgagctgctatcaaagtt |
| | LYK5JM NFR5KD | LYK5-NFR5-F | cgaaaacagaaaggttggggaatcagtg |
| | | NFR5 LYK5-R | ccccaacctttctgttttcgtcgctgaaatt |
| | LYS11JM NFR5KD | LYS11 NFR5-F | gagcagtgtaaggttggggaatcag |
| | | NFR5 LYS11R | ccccaaccttacactgctcattgagg |
| site mutantions construction | NopT | T93S-F | gaatctccgtcggcttgactgc |
| | | T93S-F | gacggagattccatctgcgttc |
| | | 97D-F | ggcttggatgcggagtggctgcg |
| | | 97D-R | ctccgcatccaagccgacgcag |
| | | 97A-F | ggcttggctgcggagtggctgc |
| | | 97A-R | ctccgcagccaagccgacg |
| | | 109D-F | catccggatatccgaatggaggccctag |
| | | 109D-R | ttcggatatccggatgactgttgaggttacgc |
| | | 109A-F | catccggcaatccgaatggag |
| | | 109A-R | ttcggattgccggatgactgttg |
| | | 119D-R | cccggagatcaaaggcacgcctcagc |
| | | 119A-F | gcctttgatctccgggtactagggcc |
| | | 119A-F | cccggagcgcaaaggcac |
| | | 119A-R | ctttgcgctccgggtactagg |
| | | 156D-F | gcaaaacgatatgttgcagaaagcaggc |
| | | 156D-R | gcaacatatcgttttgcgccgcgaagtcggcc |
| | | 156A-F | gcaaaacgctatgttgcagaaagcag |
| | | 156A-R | ctgcaacatagcgttttgcgccgcgaagtc |
| | | H205A-F | ggcggcaacaccgttgcgacc |
| | | H205A-R | caacggtgttgccgccgccctcagc |
| | | D220A-F | ctcttcgctcctaatttcggcgaatttac |
| | | D220A-R | gccgaaattaggagcgaagagcgtggtgtttc |
| | | 245D-F | cgctacgacaatccaaaccggcag |
| | | 245D-R | ttggattgtcgtagcgattggctaggct |
| | | 257 mutants | changing the sequence of R primer at nopt$_{257}$ when cloned into vectors |
| | NFR1 | K351E-F | gcaattgagaagatggatgtacaagcatc |
| | | K351E-R | catccatcttctcaattgctgttttcttgcc |
| | NFR5 | 269-271A-F | ctatgcagctgcacgcagaaagaaggctctg |
| | | 269-271A-R | gcgtgcagctgcatagaattcggatccgcgc |
| | | 272-274A-F | ctgcgctgcagctaaggctctgaataggactgc |
| | | 272-274A-R | cttagctgcagcgcagtatacatagaattcggatcc |

| | | | |
|---|---|---|---|
| | | 275-277A-F | gaaaggcagctgcaaataggactgcttcatcagc |
| | | 275-277A-R | ctatttgcagctgcctttctgcggcagtatacatag |
| | | 278-280A-F | ctggctgcagctgcttcatcagctgagactg |
| | | 278-280A-R | agcagctgcagccagagccttctttctgcg |
| | | 282-283A-F | gcagcagctgagactgctgataaactac |
| | | 282-283A-R | gcagtctcagctgctgcagcagtcctattcagag |
| | | 285-286A-F | ggggctgctgataaactactttctggag |
| | | 285-286A-R | agtagtttatcagcagccccagctgatgaagcagtc |
| | | 288-290A-F | gctgcagcactttctggagtttcaggctatgtaag |
| | | 288-290A-R | aaactccagaaagtgctgcagcagcagtctcagctgatgaag |
| | | 291-294A-F | gagactgctgataaactagctgctgcagtttcaggctatgtaagcaagcc |
| | | 291-294A-R | atagcctgaaactgcagcagctagtttatcagcagtctcagctg |
| | | 295-298A-F | gagttgcagctgctgtaagcaagccaaacgtgtatg |
| | | 295-298A-R | cttacagcagctgcaactccagaaagtagtttatcagcag |
| | | 298-300A-F | ctatgcagctgcaccaaacgtgtatgaaatcgacg |
| | | 298-300A-R | gtttggtgcagctgcatagcctgaaactccagaaagtag |
| | | 301-303A-F | gcaaggcagctgcttatgaaatcgacgagataatggaagc |
| | | 301-303A-R | cataagcagctgccttgcttacatagcctgaaactcc |
| | | 304-306A-F | gtggctgcagctgacgagataatggaagctacgaag |
| | | 304-306A-R | cgtcagctgcagccacgtttggcttgcttacatagc |
| | | 307-309A-F | gaaatcgctgcagcaatggaagctacgaaggatttcag |
| | | 307-309A-R | ccattgctgcagcgatttcatacacgtttggcttgc |
| | | 310-312A-F | cgagatagcagctgctacgaaggatttcagcg |
| | | 310-312A-R | gtagcagctgctatctcgtcgatttcatacacgtttg |
| | | 313-315A-F | gctgcagctgcattcagcgatgagtgcaagg |
| | | 313-315A-R | ctgaatgcagctgcagcttccattatctcgtcgatttc |
| | | 316-318A-F | ggatgcagctgctgagtgcaaggttggggaatc |
| | | 316-318A-R | ctcagcagctgcatccttcgtagcttccattatctc |
| | | 319-321A-F | cgatgctgcagctgttggggaatcagtgtacaagg |
| | | 319-321A-R | caacagctgcagcatcgctgaaatccttcgtagc |
| | | 322-324A-F | gcaaggcagctgcatcagtgtacaaggccaacatag |
| | | 322-324A-R | gatgcagctgccttgcactcatcgctgaaatc |
| | | 325-327A-F | gaagcagctgcaaaggccaacatagaaggtcg |
| | | 325-327A-R | gcctttgcagctgcttccccaaccttgcactc |
| | | 268-277DE-F | agaagcgcggatccgaattcaataggactgct |
| | | 268-277DE-R | ctgatgaagcagtcctattgaattcggatccgcgcttc |
| | | 278-287DE-F | gcagaaagaaggctctggataaactactttctggagtttcagg |
| | | 278-287DE-R | aaactccagaaagtagtttatccagagccttctttctgcg |
| | | 298-307DE-F | caggctatgagataatggaagctacgaaggatttc |
| | | 298-307DE-R | cttccattatctcatagcctgaaactccagaaagtag |

| | | 308-317DE-F | gaaatcgacgatgagtgcaaggttgggg |
|---|---|---|---|
| | | 308-317DE-R | ttgcactcatcgtcgatttcatacacgtttggc |
| | | 318-327DE-F | gatttcagcaaggccaacatagaaggtcg |
| | | 318-327DE-R | gttggccttgctgaaatccttcgtagcttcc |
| | | 328-337DE-F | tcagtgtacgtaaagaaaatcaaggaaggtggtgcc |
| | | 328-337DE-R | attttctttacgtacactgattccccaaccttg |
| | | 288-294A-F | gcagctgccgcagcggctgcctcaggctatgtaagcaagcc |
| | | 288-294A-R | gcagccgctgcggcagctgcagcagtctcagctgatgaagc |
| | | 288-298PBS1-F | ggagacaaatctcatgtctcctcaggctatgtaagcaagcc |
| | | 288-298PBS1-R | ggagacatgagatttgtctccagcagtctcagctgatgaagc |
| | | N5-283S-R | tgaagcagtcctattcagagcc |
| | | 283Y-F | tctgaataggactgcttcatacgctgagactgc |
| | | N5-294G | agaaagtagtttatcagcagtctcag |
| | | 294Q-F | ctgctgataaactactttctcaggtttcaggctatgtaag |
| | | N5-303Y-R | cacgtttggcttgcttacatag |
| | | 304S-F | gcaagccaaacgtgtccgaaatcgacgag |
| | | N5-310AT-R | ttccattatctcgtcgatttcatacac |
| | | 311-2 IY-F | gaaatcgacgagataatggaaatctacaaggatttcagcgatgagtg |

1120