The Rhizobial effector NopT targets Nod factor receptors to regulate symbiosis in *Lotus japonicus*

Hanbin Bao[1], Yanan Wang[1], Haoxing Li[1], Qiang Wang[1], Yutao Lei[1], Ying Ye[1], Syed F. Wadood[2], Hui Zhu[1], Christian Staehelin[2], Gary Stacey[3], Shutong Xu[1], Yangrong Cao[1*]

[1] National Key Lab of Agricultural Microbiology, Hubei Hongshan Laboratory, Huazhong Agricultural University, Wuhan, Hubei, 430070, China

[2] State Key Laboratory of Biocontrol and Guangdong Key Laboratory of Plant Resources, School of Life Sciences, Sun Yat-sen University, East Campus, 510006 Guangzhou, China

[3] Divisions of Plant Science and Technology and Biochemistry, Christopher S. Bond Life Sciences Center, University of Missouri, Columbia, MO 65211, USA

* Corresponding author: Yangrong Cao, email: yrcao@mail.hzau.edu.cn

Short title: NopT associates with NFR1 and NFR5 to regulate symbiosis

Keywords: Legume–rhizobial symbiosis; rhizobial effector; Nod Factor Receptors; NopT.

One-sentence summary: The rhizobial effector NopT forms a mutual interaction with plant's two Nod factor receptors, NFR1 and NFR5, to precisely regulate rhizobial symbiosis in *Lout japonicus*

Abstract

It is well-studied that type-III effectors are essential required by gram-negative pathogens to directly target different cellular pathways from hosts, thus enhancing bacterial infection. However, in the unique context of legume-rhizobium symbiosis, the role of rhizobial effectors in regulating plant symbiotic pathways remains a subject of hypothesized understanding. Here, we show that NopT, a YopT-type cysteine protease from *Sinorhizobium fredii* NGR234 that could directly target symbiotic signaling pathway by associating with two plants' Nod factor receptors (NFR1 and NFR5). NopT inhibits cell death triggered by co-overexpression of NFR1/NFR5 in *Nicotiana benthamiana*. NopT interacts with both NFR1 and NFR5. NopT proteolytically cleaves NFR5 in both *in vitro* and *in vivo* assays, but is later inactivated *via* phosphorylation by NFR1. NopT plays an essential role in mediating rhizobial infection in *L. japonicus*. The truncated NopT, lacking 50 N–terminal amino acids post-autocleavage, retains the ability to cleave NFR5 but no longer interacts with NFR1. Intriguingly, certain *Sinorhizobium* species exclusivley harbor this truncated NopT in their genomes. These findings unveil an intricate regulatory interplay between rhizobia and legumes, wherein rhizobial NopT actively cleaves NFR5, and this cleavage process is subsequently suppressed following NopT phosphorylation by NFR1. This discovery presents a compelling narrative of a nuanced regulatory interplay between rhizobia and legumes, highlighting rhizobial effector as a direct mediator targeting symbiotic pathway from plants.

Introduction

The legume–rhizobial symbiosis stands as one of the most crucial and intricate mutualistic interactions in nature. Its significance extends beyond the supply of the substantial amount of ammonium provided by rhizobia, influencing plant growth and ecosystem sustainability, to serving as a pivotal model for investigating plant–microbe interactions (Yang et al., 2022). The orchestration of this symbiosis involves the perception of small molecules,

such as nod factors (NFs) and surface polysaccharides (Madsen et al., 2003; Kawaharada et

al., 2015), along with the secretion of proteins, inlcuding the type three secretion system

(T3SS) effectors, to regulate legume–rhizobial symbiosis (Sugawara et al., 2018; Zhang et

al., 2021).

A receptor heterocomplex, consisting of two Lysin–motif receptor kinases (LYKs), plays a

central role in recognizing various polysaccharide molecules. In nonlegumes like

Arabidopsis, AtLYK5 and AtCERK1 (Chitin elicitor receptor kinase) are required for

perception of chitin, triggering plant immunity to combat fungal pathogens (Cao et al.,

2014). Conversely, in legumes, structurally similar receptors like *L. japonicus* NF

receptor1 (LjNFR1) and LjNFR5 are vital for recognizing rhizobial

lipo–chitooligosaccharide NF, initiating symbiotic signaling transduction (Broghammer et

al., 2012; Bozsoki et al., 2017). The intricate regulation of NFRs is highlighted by the

instances of hypersensitive cell death induced by their simultaneous overexpression in

Nicotiana benthamiana leaves (Madsen et al., 2011) and defense–like responses in

Medicago truncatula nodules upon overexpression of MtNFP (Nod factor perception, the

ortholog of NFR5) (Moling et al., 2014), perhaps reflecting potential conservation of the

plant LYK receptors in interactions with both pathogens and symbionts. Therefore, the

protein abundance of NFRs must be tightly controlled to avoid this resistance response,

allowing rhizobial infection to proceed, emphasizing the need for tight control to enable

rhizobial infection.

T3SS effectors play a critical role in suppressing pattern–triggered immunity (PTI)

during pathogen infections in plants (Jones and Dangl, 2006; Schubert et al.,

2020). To avoid diseases, specific resistance (R) genes are employed to

recognize successful avirulence (Avr) effectors which enable pathogens overcome

PTI, known as effector-triggered immunity (ETI), a much stronger immnune

response combined with programmed cell death (Jones and Dangl, 2006).

However, in contrast to the well-studied fucntion of Avr effectors from

phytopathogenic bacteria in mitigating immunity by different mechanisms, how

these effectors from symbiotic bacteria might contribute to mutualistic interaction is

largely unknown.

Some effectors, such as YopT-type cysteine protease family, including

Pseudomonas AvrPphB, *Yersinia* YopT, and *Rhizobial* NopT, are well conserved

across different bacteria. AvrPphB from *Pseudomonas syringae* can directly

proteolytically cleaves Arabidopsis receptor-like cytoplasmic kinase (RLCK) family

proteins including PBS1, PBS1-like proteins, and BIK1, to suppress PTI (Zhang et al.,

2010). While, the cleavage of PBS1 serves as a "decoy" in activating RPS5

(RESISTANCE TO PSEUDOMONAS SYRINGAE5)–mediated ETI (Shao et al., 2003b).

Yersinia YopT directly cleaves Rho family leading to their inactivation to disrupt

actin cytoskeleton of host cells (Shao et al., 2003b; Shao et al., 2003a). While, in

rhizobium species, NopT have a dual function in regulating symbiosis. The

exogenous expression of NopT from *Sinorhizobium fredii* NGR234 in *S. fredii*

USDA257 almost blocks nodulation in soybean Nanfeng 15 but not other soybean

cultivars (Khan et al., 2022), implying that NopT might act as an Avr effector.

However, NopT positively regulates nodule formation in the interaction betwen *S.*

fredii HH103 and soybean Suinong 14 (Li et al., 2023). NopT from *Bradyrhizobium*

sp. ORS3257 promotes rhizobial infection in nodules of *Aeschynomene indica* (Teulet

et al., 2019). Despite this, little is understood about how NopT precisely modulates

rhizobial infection and the variations among NopTs from different rhizobia strains.

The challenge in studying rhizobial effectors lies in the availability of suitable assays by

which to characterize their phenotypes. In this study, we capitalized on the observation that

ectopic expression of NFR1 and NFR5 in *N. benthamiana* leaves induces programed cell

death (Madsen et al., 2011), a response that might be modulated by rhizobial effectors.

Among the 15 known T3SS effector proteins encoded within the *S. fredii* NGR234 genome,

only NopT, a homolog of the pathogenic effectors, YopT and AvrPphB (Shao et al., 2002;

Dai et al., 2008), could specifically suppress this cell death response. This suggests that

NopT might be an effector associated with NFRs. Our findings demonstrate that NopT

interacts with both NFR1 and NFR5 at the plasma membrane, and cleaves NFR5

proteolytically at the juxtamembrane position to suppresses NF–mediated physiological

responses. Perhaps as a means to regulate NopT activity, NFR1 phosphorylates intact

NopT, inactivating its protease activity. However, the autocleaved NopT retains the ability

to cleave NFR5 but loses its association with NFR1. Intriguingly, some *Sinorhizobium*

species possess only the truncated versions of NopT (relative to NGR234 NopT) that might

evade suppression by NFR1. This study enveils a distinct and mutual regulatory model, in

which the rhizobial effector NopT directly targets NFR5 to suppress symbiotic signaling,

and NFR1-mediated phosphorylation acts as a negative feedback mechanism. The model

provides essential insights into the intricate interplay between legumes and rhizobia,

revealing a counteraction where a rhizobal T3SS effector impairs NF signaling by directly

targeting NFR5, and NFR1 phosphorylates and inactivates the rhizobal T3SS effector in

response.

Results

NopT suppresses the cell death response from overexpression of NFR1 and NFR5

To examine the role of effector proteins in regulating rhizobial infection, *S. fredii* NGR234,

a strain with exceptionally broad host–range was chosen. Initial experiments were

performed to screen all 15 effector proteins identified from *S. fredii* NGR234 (Kimbrel et

al., 2013), for their ability to suppress or enhance the cell death response in *N. benthamiana*

leaves due to co–expression of LjNFR1 and LjNFR5 (Fig. S1A–S1C). Out of the effectors

tested, only NopT, a cysteine protease belonging to the YopT family (Shao et al., 2002; Dai

et al., 2008), could suppress LjNFR1 and LjNFR5–induced cell death (Fig. 1A and

S1D–S1F). Cysteine–93 of NopT is a known residue essential for its protease activity (Dai

et al., 2008). In contrast to the wild–type NopT protein, expression of NopT^{C93S}, an

inactive version, was unable to suppress cell death induced by NFR1 and NFR5 compared

with controls (Fig. 1B), indicating the protease activity of NopT is essential for suppression

of cell death in *N. benthamiana*. In contrast to the suppressive function on cell death,

NopM, an effector protein from *S. fredii* NGR234 reported to have an E3 ligase activity,

induced strong cell death when overexpressed in *N. benthamiana* leaves (Fig. 1C).

As controls, we tested whether the action of NopT was specific for NFR1/NFR5

expression or whether it could act generally to suppress cell death. A variety of other

proteins have been reported to induce cell death when expressed in *N. benthamiana* leaves;

specifically, Avr3a and R3a (Bos et al., 2006), apoptosis regulator BAX (Bcl-2-associated

X protein) (Gan et al., 2009), INF1, an elicitor from *Phytophthora infestans* (Vleeshouwers

et al., 2006), Arabidopsis AtCERK1 (Cao et al., 2014), and NopM (Fig. 1C). In contrast to

the case with LjNFR1 and LjNFR5, expression of NopT was unable to suppress the cell

death induced when each of these proteins was expressed in tobacco leaves (Fig. 1D).

Hence, the effect of NopT appears to be specific for the NF receptors. These results

suggested a hypothesis that rhizobial NopT protease might act directly on the NFR1 and/or

NFR5 proteins.

	Avr3a&R3a	NopM	BAX	INF1	AtCERK1	NFR1&NFR5
Death/Total	10/10	57/57	16/16	24/24	75/78	2/97

Figure 1. NopT specifically suppresses cell death triggered by NFR1 & NFR5 in *N. benthamiana*.
Agrobacterium strains harboring genes were NopT or EV infiltrated into *N. benthamiana*,12 hpi followed by infiltration with *agrobacterium* containing NFR1 and NFR5. Different leaf discs indicate where different proteins were expressed. NopT (A) but not NopT[C93S], the protease–inactive version of NopT (B) suppressed cell death induced by expression of NFR1 and NFR5. (C) Expression of NopM induced cell death in *N. benthamiana*. (D) Expression of NopT could not suppress the cell death triggered by expression of Avr3a&R3a, NopM, BAX, INF1 or AtCERK1. Digital numbers in (D) represent the number of leaf discs with cell death and total leaf discs tested. All pictures in the Figure are representative of at least three independent biological replicates.

NopT colocalizes with and interacts with NFR1 and NFR5 at the plasma membrane

NopT contains an acylation site that appears to localize NopT to cellular membranes (Dowen et al., 2009). Therefore, we tested the possibility that NopT, NFR1 and NFR5 co–localize on the plasma membrane. *In vivo* bimolecular fluorescence complementation (BiFC) assays were used to examine the colocalization of NopT with LjNFR1 and LjNFR5. NopT, LjNFR1 and LjNFR5 were all C–terminally tagged with nYFP and cYFP. Fluorescence signals representing the associations between NopT and LjNFR1, LjNFR5, as well as itself, were detected at the plasma membrane (Fig. 2A and 2B), suggesting that NopT colocalized with and interacted with both LjNFR1 and LjNFR5. Interactions

between NopT and LjNFR1 and LjNFR5 were further verified using a Split–luciferase

(Split–LUC) assay (Fig. 2C). Co–expression of NopT–cLUC and LjNFR1–nLUC or

LjNFR5–nLUC produced strong luminescence signals in leaf discs compared with the

negative control expressing FLS2–nLUC and NopT–cLUC (Fig. 2C).

It has been known that the acylation of NopT could change its subcellular localization

(Dowen et al., 2009). We then exaimed the interaction between NFR1/NFR5 and

NopT$^{G50A/C51A/C52A}$, a mutant version on the acylation sites. The interactions between

NopT$^{G50A/C51A/C52A}$ and NFR1/NFR5 were significantly reduced compared to NopT^{WT} in

Split–LUC assays (Fig. S2A and S2B). While the mutation at catalytic site of NopT

(NopT^{C93S}) remained similar interaction with NFR1 and NFR5 (Fig. S2C). The interaction

results were further confimred by co–immunoprecipitation (co-IP) assay. The

C–terminally HA–tagged LjNFR1 and LjNFR5 could be precipitated by anti–FLAG

antibody in the sample expressing NopT–FLAG compared with controls (Fig. 2D). The

interaction between NopT and LjNFR1 and LjNFR5 suggested that LjNFR5 or LjNFR1

might be a target protein for proteolysis by NopT.

Figure 2. NopT interacts with NFR1 and NFR5. Interactions between NopT and NFR1 and NFR5

were detected using Split–YFP assay (A) and (B), Split–LUC (C) and co–IP (D) assays in *N.*

benthamiana. (A) and (B). nYFP and cYFP tags were fused at the C–terminus of NopT, NFR1, NFR5,

and negative control FLS2. YFP fluorescence signals represent protein–protein interactions. Scar

bar=25 um. (C). nLUC and cLUC tags were fused at the C–terminus of NopT, NFR1, NFR5, and
negative control FLS2. Luminescence signals represent protein–protein interactions. (D). HA–tagged
NFR1 and NFR5 and FLAG–tagged NopT were expressed in plant cells followed by
immunoprecipitation using anti–FLAG antibody and determined by immunoblot using anti–HA and
anti–FLAG antibodies. All data in Figure are representative of three biological replicates.

NopT proteolytically cleaves LjNFR5 at the juxtamembrane domain

A previous study showed that NopT protease activity is detectable upon autocleavage of

about 50 amino acids from its N–terminus (Dai et al., 2008). We next tested whether NopT

could directly proteolyze LjNFR1 and/or LjNFR5 when co–expressed in *N. benthamiana*

leaves. The addition of NopT but not NopT^{C93S}, an inactive version, resulted in proteolytic

cleavage of LjNFR5–GFP but not LjNFR1–GFP as shown by immunoblotting of the

SDS–PAGE gel (Fig. 3A and S3A). The *in vivo* cleavage assay in *L. japonicus* transgenic

root also showed that NopT but not NopT^{C93S} resulted in proteolytic cleavage of LjNFR5

(Fig. 3B). Based on the molecular weight change, the cleavage sites were predicted to

occur within the cytoplasmic domain (CD) of LjNFR5. Therefore, the CD of NFR5 was

constructed as a recombinant protein (Strep–NFR5^{CD}–HA) and then co–expressed with

NopT–FLAG or the inactive version NopT^{C93S}–FLAG in *Escherichia coli* cells. The

results showed that the active NopT protease could cleave NFR5^{CD} resulting in a band

roughly ~5 kDa less than the control as detected using anti–HA antibody (Fig. 3D). In the

same experiment, when the CD of NFR1 was co–expressed with NopT, no cleavage

product was detected in immunoblotting assay (Fig. 3C). However, migration of the bands

representing NopT or NopT^{C93S} was significantly retarded as seen after immunoblotting

(Fig. 3C). This retardation of NopT protein band may be the result of phosphorylation by

NFR1.

The cleavage product with ~5 kDa lower molecular weight in the cleavage assay,

suggested that the cleavage site is located within the juxtamembrane (JM) position of

NFR5. To test this hypothesis, SUMO–NFR5^{KD}–HA, a recombinant protein with the JM of

NFR5 replaced with a SUMO tag, was constructed and used for the cleavage assay. Indeed,

NopT could not cleave SUMO–NFR5^{KD}–HA (Fig. S3B). To confirm this conclusion, the

JM of NFR5 was used to bridge SUMO and GFP tags to create a recombinant protein,

SUMO–JM–GFP. *In vitro* cleavage assay, strong band intensity representing GFP was

detected in the flow though sample incubated with NopT but not NopT^{C93S} (Fig. 3E). These

data strongly indicated that the cleavage site for NopT is located at the JM of NFR5.

Figure 3. NopT proteolyzes NFR5 at its juxtamembrane domain. (A) NopT but not NopT^{C93S}

cleaves NFR5–GFP protein expressed in N. benthamiana cells by releasing NFR5CD–GFP. (B) NopT

but not NopT^{C93S} cleaves NFR5-Myc protein expressed in *L. japonicus* transgenic root by releasing

NFR5CD-Myc. Asterisk indicated the cleaved variant of NFR5. (C) A repeated experiment testing the

cleavage of NFR5^{CD} but not NFR1^{CD} fused with different tags. Asterisk represents the phosphorylated

NopT by inducing a band retardation on SDS–PAGE gel. (D) NopT but not NopT^{C93S} cleaves the CD of

NFR5 protein expressed in *E. coli* cells by releasing the kinase domain of NFR5. Asterisk indicated the

cleaved variant of NFR5. (E) NopT cleaves a recombinant protein where His–tagged SUMO and GFP is

bridged with the JM domain of NFR5. F.T. represents flow through sample after Ni–beads purification.

(F) NopT could not cleave a mutant version of NFR5CD–HA with 5 residues in the JM substituted into

other residues (S283Y, G294Q, Y303S, A310I, T311Y). (G) NopT could not cleave *Medicago*

truncatula NFP and from NFR5^{JM}–NFP^{KD}. CD represents cytoplasmic domain, KC represents kinase

domain and C-terminal tail region, NopT^{C} indicates the truncated version of NopT after autocleavage by

releasing about 50 a.a. at its N–terminus.

The cleavage of LjNFR5 at JM by NopT is dependent on multiple residues of NFR5

NopT has an autocleavage site within its N–terminus, we identified the peptide

(DKLLSGV) from Asp–288 to Val–294 of NFR5 (Fig. S4B) as showing the highest

similarity with the NopT autocleavage region (DKMGCCA). However, the mutant version

of NFR5^{CD} with the peptide (DKLLSGV) replaced with 7 alanines could still be cleaved by

NopT (Fig. S3C). NopT and its homlog AvrPphB (Shao et al., 2003b), an effector from the

pathogen *Pseudomonas syringa*, belong to the same protein family. Hence, we set up a

control experiment by constructing a version of NFR5^{CD}, where the "DKLLSGV" motif

was replaced with the cleavage sites of AvrPphB. In the cleavage assay, we found that this

modified NFR5^{CD} was cleaved by both AvrPphB and NopT, and the cleaved product of

modified NFR5^{CD} by AvrPphB shifting faster than by NopT (Fig. S3C), suggesting that the

cleavage site of NFR5 by NopT located ahead of "DKLLSGV" motif. As mentioned

above, NopT undergoes autocleavage but it is not clear whether this is due to

inter–proteolysis or intra–proteolysis. His–205 and Aps–220 of NopT are two other

conserved residues required for the protease activity of YopT, another homolog of NopT

from *Yesinia* whose function has been extensively studied (Shao et al., 2003b). Therefore,

we targeted these residues for site–directed mutagenesis. The cleavage assay showed that

NopT could not proteolyze NopT^{C93S}, NopT^{H205A}, or NopT^{D220A} (Fig. S3D), suggesting

that intra–proteolysis might occur and the autocleavage site of NopT could not be used to

predict the cleavage site of substrate.

Figure 4. Phosphorylation of NopT by NFR1 suppresses its proteolytic activity. (A) *In vivo* phosphorylation assay in *L. japonicus* roots. Phosphorylation of NopT^{C93S} was induced with rhizobia treatment and dependent on *nfr1*. (B) NFR1^{CD} phosphorylates NopT by inducing a gel band shift determined by immunoblotting. (C) The phoshorylation sites on NopT identified by liquid chromatography–mass spectrometry (LC–MS) were either substituted into alaine (A) or asparatate (D) and subsequently tested for their ability to proteolytically cleave NFR5^{CD}.

To define the cleavage site of NFR5, we created a variety of point mutations covering the JM domain of NFR5 (Val-269 to Cys-320) and first seven a.a. of the kinase domain (Lys-321 to Tyr-327). First, 17 mutant versions of NFR5^{JM} were created with three adjacent amino acids replaced with three alanines for proteolysis assay in *E. coli* cells. However, all 17 mutant versions of NFR5^{CD} could be cleaved by NopT but not NopT^{C93S} (Fig. S3E–S3H). An additional 7 truncated versions of NFR5^{JM}, with deletion of 10 adjacent residues, were created and tested for cleavage by NopT. Surprisingly, the cleavage products were still observed from all 7 truncated versions of NFR5^{JM} (Fig. S3I). The *in vitro* cleavage assay suggested that the JM domain might have multiple sites susceptible to be cleaved by NopT, which is different from the specific cleavage site

identified for its homologous protein AvrPphB. In a parallel experiment, the recombinant

protein, His-SUMO-NFR5^{CD}-GST expressed in *E. coli* was subjected for N-terminal

protein sequencing determined by Liquid Chromatography Mass Spectrometry analysis

after cleavage. Based on allignment using identified peptides, the cleavage site of NFR5 by

NopT was mapped to be after four basic amino acids in the JM of NFR5, "RRKK" from

271 to 275 a.a. of NFR5 (Fig. S4C). However, the result is not consistance with the data

from cleavage assay. When the region from Tyr-268 to Leu-277 was deleted, NopT could

still proteolyze NFR5-CD (Fig. S4I). The cleavage assay suggested that NopT might prefer

to proteolyze NFR5 after four polybasic residues in the NFR5^{JM}, but other sites could also

be cleaved by NopT.

Because LjNFR5 belongs to a subfamily of plant LYK family proteins that lacks kinase

activity (Yang et al., 2022), we then asked whether NopT could proteolyze other LYK

proteins in this subfamily, which might be helpful to characterize the biochemical function

of NopT. Hence, the CDs from Arabidopsis AtLYK5, *L. japonicus* LjLYS11 and *M.*

truncatula MtNFP were used for cleavage assays in the presence of NopT. As shown in

Fig. S3J, NopT, but not NopT^{C93S}, could cleave the CDs of both AtLYK5 and LjLYS11.

Replacement of NFR5^{JM} with the JM domains from either AtLYK5 or LjLYS11 also

resulted in cleavage by NopT (Fig. S3J). The data strongly indicated that NopT could

cleave both AtLYK5 and LjLYS11 at their JM domains. Based on sequence analysis of the

JM region of NFR5 and its homologous proteins, five conserved residues (Ser-283,

Gly-294, Tyr-303, Ala-310, Thr-311) were identified and mutated for the proteolysis

assay (Fig. S4B). As shown in Fig. 3F, the mutant version of NFR5 (i.e., NFR5^{CD} with five

amino acid mutations) could not be cleaved by NopT. Interestingly, NopT coud not

protelyze MtNFP^{CD} (Fig. 3G). Then, we generated two recombinant proteins, by switching

the JM between NFR5^{CD} and NFP^{CD}, generating NFR5^{JM}-NFP^{KC} (KC represents kinase

domain and C-terminal tail region) and NFP^{JM}-NFR5^{KC} for *in vitro* cleavage assays. As

shown in Fig. 3G, NopT could proteolyze NFP^{JM}-NFR5^{KC} but not NFR5^{JM}-NFP^{KC}. The

data suggested that the cleavage of NFR5 by NopT might be dependent on the

confirmational structure of NFR5^{KC} and/or other specific amino acids in NFR5^{KC}. To

define this, two additional recombinant proteins with large switch between NFR5^{CD} and

NFP^{CD}, NFR5$^{268\text{-}445}$-NFP$^{458\text{-}595}$ and NFP$^{270\text{-}457}$-NFR5$^{456\text{-}595}$, were constructed for

proteolysis (Fig. S3K). However, both of them could be cleaved by NopT, suggesting that

a much large region on NFR5^{KC} is required for being proteolyzed by NopT. Based on all

these cleavge results, we concluded that NopT could proteolyze NFR5 at the JM domain,

which nonadjacent mutiple residues within the NFR5^{JM} and the specific KC region are

required for the proteolysis.

NopT phosphorylation by LjNFR1 suppresses its proteolytic activity

The perception of rhizobia NFs by the heterocomplex of NFR1 and NFR5 involves the

trans–phosphorylation from the NFR1 kinase to the pseudo–kinase NFR5 (Madsen et al.,

2011). Rhizobial effector NopT associates with both NFR1 and NFR5, which suggests that

NopT might be a phosphorylation target of NFR1 and involved in the transphosphorylation

process between the two receptors. Kinase assays showed that NFR1–CD but not the

kinase–inactive NFR1^{K351E}–CD (Lys-351 is the conserved residue required for ATP

binding) could phosphorylate NopT (Fig. 3C and 4B) and NFR5–CD (Fig. S5) as shown by

band retardation on SDS–PAGE gels identified using immunoblotting. The band

representing phosphorylated NopT shifted upon the addition of a commercial phosphatase,

calf intestinal alkaline phosphatase (Fig. 4B), indicating that NopT as well NFR5^{CD} are

phosphorylation targets of the NFR1 kinase. NopT autocleavage also gives two bands and

the site mutant of NopT at C93S did not affect the interactions between NFRs(Fig. S2C).

To avoid multi-bands interference of cleaved NopT, NopT^{C93S} was expressed in both wild

type plants and *nfr1* mutant then treated with rhizobia or water. After analyzed by

Zn^{2+}-Phos-tag PAGE gel, several shifted bands were strengthened after rhizobia treatment

in WT roots but not in the *nfr1* mutant roots (Fig. 4A), suggesting that the phosphorylation

of NopT could be induced by NFR1 and multi-sites of NopT were phosphorylated *in*

planta. The phosphorylation sites of NopT were further identified using Liquid

Chromatography Mass Spectrometry (Table S1). We then addressed whether NopT

phosphorylation by NFR1 regulates its proteolytic activity on NFR5. All the potential

phosphorylation sites of NopT were substituted into either alanine in order to block

phosphorylation of these sites or aspartate, which mimicks the acidic condition

representing a phosphorylated status. These mutant proteins were then used in the cleavage

assay. Aspartate substitutions on three phosphorylation sites of NopT, Thr–97, Ser–119,

and Thr–257, resulted in the loss of the ability of NopT to cleave NFR5–CD, as well as the

ability to autocleave (Fig. 4C). In contrast, NopT proteins with alanine substitutions on

these three sites retained the ability to cleave NFR5–CD and to autocleave (Fig. 4C). The

data suggest that phosphorylation of NopT by NFR1 likely inhibits proteolytic activity,

retaining a fully functional NFR5 and a full length NopT.

Figure 5. NopT regulates rhizobial infection in *L. japonicus*. (A) Rhizobial infection in *L. japonicus*

using GFP-labelled WT and *nopt* mutant strains of *S. fredii* NGR234. Fluorescent, GFP-expressing

bacteria are imaged in Cyan. Scar bars=12.5µm. (B) GUS staining pictures from the roots of *L.*
japonicus transgenically expressing GUS under *p*NIN promoter inoculated with WT and *nopT* mutant
strains of NGR234. Scar bars=1mm (C) Statistical analysis of rhizobial infection in A (n=10, Student's
t–test: P < 0.01). (D) Statistical analysis of GUS straining sites in B (n=10, Student's *t*–test: P < 0.01).
(E) Nodule primordia number on the roots 14dpi after inoculation WT, *nopT* mutant strains or *pT7*:
NopT (n=16, Student's *t*–test: P < 0.01). (F) Infection foci number from the wild type plant roots
inoculated with WT strains and *nopt* mutant strains expressing NopT with different variations (n=5,
Student's *t*–test: P < 0.01). (G) Statistical analysis of rhizobial infection in *L. japonicus* transgenic root
expressing GFP (EV, empty vector control) or NopT using DsRed-labeled *M. loti* MAFF303099 (n=8,
Student's t–test: P < 0.01).

NopT dampens rhizobial infection

The fact that *S. fredii* NGR234 NopT could directly target and cleave NFR5 indicates that

NopT may be an important player regulating rhizobial infection in legumes. Nod factor

signal is known to function in the early stage of rhizobial infection and nodulation

specificity (Geurts et al., 2005; Wang et al., 2012). Prior work showed that type three

secretion system of *S. fredii* NGR234 might not regulate nodule numbers few weeks after

inoculation in *L. japonicus* (Ausmees et al., 2004). Therefore, we focused on the early

stage of rhizobial infection. Indeed, inoculation with GFP–labelled *S. fredii*

NGR234Δ*nopT* induced a massive infection within the root hairs of *L. japonicus* (Gifu)

plants compared with the favorable infection by the NGR234 wild type strain (Fig. 5A and

5C). Transgenic expression of NopT in *L. japonicus* root hair reduced the infection of

Mesorhizobium loti MAFF303099 (Fig. 5G). In *L. japonicus* plants transgenically

expressing *pNIN*:*GUS*, greater range of GUS expression (Fig. 5B and 5D) and formation of

more nodule primordia (Fig. 5E) also reflected a stronger plant response to infection by

NGR234Δ*nopT* relative to the wild–type strain. To confirm the massive infection is due to

the loss of NopT, NopT was expressed in the *S. fredii* NGR234Δ*nopT* strain and used for

inoculation in *L. japonicus*. Complementation with NopT in NGR234Δ*nopT* reduced the

massive infection foci to the similar levels as the WT strain (Fig. 5F). We then tested

whether the proteolytic activity of NopT is required for rhizobial infection. The

protease–dead version of NopT^{C93S} and three phosphorylation mimic versions of NopT,

i.e., NopT^{T97D}, NopT^{S119D}, and NopT^{T257D}, which lack proteolytic activity, were

individually expressed in NGR234Δ*nopT*. Compared with the high levels of infection foci

induced by NGR234Δ*nopT* in *L. japonicus* plants, inoculation with these four mutant

strains induced much lower levels of infection, albeit slightly higher than that induced by

the wild–type strain (Fig. 5F). Inoculation with NGR234Δ*nopT* produced more nodule

primordia, while over-expression of NopT under the control of T7 promoter in *S. fredii*

NGR234 WT strain reduced the numer of nodule primordia, compared with wild–type

strain inoculation 14 dpi (Fig. 5E). To verify whether the cleavage of NFR5 reduce

rhizobial infection, NFR5–WT and NFR5^{5m}, a mutant version with 5 amino acids

substitutions at JM that is resistant to NopT cleavage, were expressed in the *nfr5-3* roots

under the control of its native promoter. However, expression of NFR5^{5m} was not observed

to rescue the infection phenotype of *nfr5-3* mutant plants (Fig. S5A). The possible reason is

that amino acids substitutions at JM might lead to a conformation change of NFR5 that is

required for its function in symbiotic signaling transduction. Therefore, NFR5-NFP^{KC}, the

KC domain of NFR5 replaced with KC domain of NFP, was expressed in the roots under

the control of native promoter (Fig. S5B). In the *nfr5-3* roots expressing NFR5-NFP^{KC}, no

difference of rhizobial infection was observed between *S. fredii* NGR234Δ*nopT* and *S.*

fredii NGR234 wild type strain, suggesting that the cleavage of NFR5 reduced the

rhizobial infection levels. All these data strongly indicated that the protease activity of

NopT, the phosphorylation status of NopT, and the proteolysis of NFR5 are important for

rhizobial infection.

The truncated version NopT^{ΔN50} cleaves NFR5 but fails to interact with NFR1

The significance of rhizobial NopT in mediating rhizobial infection led us to analyze its

homologs from other rhizobium species. Sequence comparisons based on the amino acids

of NopT homologs showed that *Sinorhizobium* and *Bradyrhizobium* were located in

different clades (Fig. S7). *B. diazoefficiens* (USDA110) is a typical *Bradyrhizobium* strain

used to study the symbiotic interaction with soybean. Two NopT homologous genes,

named $NopT1_{USDA110}$ and $NopT2_{USDA110}$ were identified from the USDA110 genome

(Fotiadis et al., 2012). However, when tested, neither $NopT1_{USDA110}$ or $NopT2_{USDA110}$

could not cleave NFR5 compared to $NopT_{NGR234}$ (Fig. 6A). The possible scenario is that

NopT homologs from *Bradyrhizbium* species might have lost the ability to cleave NFR5

and/or may act on other, unknown cellular target proteins. The genome of some

Sinorhizobium species, for example *S. fredii* USDA257 and S. *fredii* HH103, encode *NopT*

homologous genes predicted to produce same truncated proteins with a 57 aa deletion from

its N–terminus, relative to the NGR234 NopT sequence, which is almost the same as the

truncated $NopT_{NGR234}$ after autocleaveage. When these NopT homologs (i.e., $NopT_{USDA257}$

and $NopT_{HH103}$) were tested, both were observed to have the ability to cleave NFR5 (Fig.

6A). However, these NopT homologs could not suppress the cell death triggered by NFR1

and NFR5 in *N. benthamiana* leaves (Fig. 6B) and failed to complement the massive

infection of NGR234Δ*nopT* (Fig. 6C and 6D), suggesting that $NopT_{USDA257}$ and $NopT_{HH103}$

retained the same catalytic activity as $NopT_{NGR234}$ but might have different substrates. We

then asked whether the truncated version of $NopT^{\Delta N50}$, which mimics the truncated NopT

after autocleavage, retained the ability to interact with NFR1. However, after repeated

tests, the direct interaction between $NopT^{\Delta N50}$ and NFR5 could be detected but not the

corresponding interaction with NFR1 using Split–LUC assays (Fig. S2D). The data

suggest that some *Sinorhizobium* species, for example, *S. fredii* NGR234, *S. fredii*

USDA257 and *S. fredii* HH103, use a truncated NopT to regulate the rhizobial infection

process in a way that avoids negative regulation by NFR1 phosphorylation, which is not

true of all rhizobial strains that possess NopT homologs.

Figure 6. Working model of NopT in association with NF receptors. (A) $NopT_{234}$, NopT from *S.*
fredii NGR234; $NopT1_{110}$ and $NopT2_{110}$, NopT homologs from *B. diazoefficiens* USDA110; $NopT_{257}$,
NopT from *S. fredii* USDA257; $NopT_{103}$, NopT from *S. fredii* HH103. All NopT homologs were tested
in the proteolysis assay using $NFR5^{CD}$ as a target. (B) Rhizobial infection $NopT_{257}$ could not inhibit the
cell death triggered by NFR1 and NFR5 in *N. benthamiana*. (C)The indicated rhizobial strains induced
nin::gus expressing in *Lotus* root. Scar bars=2.5mm. (D) Statistical analysis of GUS straining in C
(n=19, Student's *t*–test: P < 0.01). (E) A proposed model of NopT in association with NF receptors.
Both NopT and $NopT^{C}$ after autocleavage exert proteolytic activities to cleave NFR5 at the

459 juxtamembrane domain to release the kinase domain of NFR5 (cleaved NFR5). NFR1 phosphorylates
460 intact NopT, but not NopT^C, to block its proteinase activity.

Discussion

Plant-microbe interactions involve two sophisticated and intelligent organisms,

each employing distinct "weapons" or strategies to communicate mutually, thereby

securing more nutrients for their survival (Jones and Dangl, 2006; Schubert et al.,

2020). A well-established regulatory model arises from the study of plant-pathogen

interactions, where T3SS effectors from gram-negative bacteria are injected into

plant cells to directly target and suppress the plant's immune pathway, favoring

pathogenic bacterium infection (Jones and Dangl, 2006). However, the unique

model of mutual regulation between legumes and rhizobia in symbiotic interactions

remains largely unknown. The broad goal of this study was to offer essential

insights into the mutual regulation mechanisms whereby rhizobial effectors directly

target symbiotic signaling pathways to dampen their infection in host cells.

Understanding how plants recognize different microbes as "friends" or "foes" is a

central question in biology. As the first line of plant response to microbes, the

perception of microbial signals by receptor proteins from plants is essential for

inducing different signaling pathways to establish various interaction

patterns–either symbiotic, pathogenic, or no responses (Zipfel and Oldroyd, 2017).

Therefore, it is not surprising to discover that the direct targeting of these receptors

by bacterial effectors represents an efficient strategy to manipulate plant signaling

pathways for enhancing bacterium infection. In the well-studied

Arabidopsis–Pseudomonas syringa interaction model, the flagellin receptor FLS2

(Flagellin Sensing 2) and its coreceptor BAK1 (BRI1-[Brassinosteroid insensitive

1]-associated receptor kinase 1), triggering plant immunity against pathogenic

infection, are targeted by several effectors from bacteria (Zipfel et al., 2004; Sun et

al., 2013). For example, AvrPto and HopB1 could suppress the kinase activity and

cleave the Arabidopsis receptor complex, respectively (Xiang et al., 2008; Li et al.,

2016). AvrPtoB, an E3 ligase in *P. syringae*, targets FLS2 to promote its

degradation, resulting in decreased immune responses in plant cells (Goehre et

al., 2008). Considering the evolutionary importance of effectors in infection, it is

rational to identify that the rhizobial effector NopT, a member of the C58 protease

family, can directly target NFRs, manipulating symbiotic signaling pathways in

plants. This was demonstrated using a unique screening system in *N.*

benthamiana leaves (Fig. S1A-C), in which co-expression of two symbiotic

receptors, NFR1 and NFR5, could lead to plant cell death to identify NopT, the only

effector from *S. fredii* NGR234. *S. fredii NGR234Δnopt* mutant strains hyperinfect

L. japonicus roots, and overexpression of NopT in the roots reduces rhizobial

infection (Fig. 5A-5D), consistent with the observed suppression of cell death in *N.*

benthamiana leaves (Fig.1A). The interaction between NopT and NFRs has been

verified and is impaired by the acylation site of NopT (Fig.2 and S2A), suggesting

that the plasma membrane localization of NopT is essential for targeting NFRs.

In the extensively studied context of microbial pathogenesis, various effectors from

pathogens have been proposed to directly suppress the immune system and

promote virulence in host cells (Tang et al., 2017; Schubert et al., 2020). The C58

protease family, including YopT and AvrPphB, represents such effectors that

disarm the host defense system to favor pathogenic infection in both animals and

plants (Shao et al., 2003a; Zhang et al., 2010). *Yersinia* YopT cleaves the Rho

family GTPase to disrupt the actin cytoskeleton, inhibiting the phagocytosis of

pathogens by hosts (Shao et al., 2003a). Conversely, *P. syringae* AvrPphB

cleaves multiple members of the RLCK (receptor-like cytoplasmic kinase)

subfamily VII to suppress the PTI response (Zhang et al., 2010). In contrast, the

cleavage of PBS1 functions as a "decoy" to trigger strong immunity against

bacterial infection (Shao et al., 2003b). We confirmed that NopT cleaves NFR5 at

the JM domain through different assays (Fig.3). Mutants with five dispersed

residues (S283, G294, Y303, A310, T311) at the JM domain (Fig.3F) and the

replacement of NFR5^{KD-CT} with NFP^{KD-CT} (Fig.3G) resist cleavage by NopT,

suggesting a joint decision of cleavage involving both the JM domain and KD-CT

domain. Although belonging to the same protease family, the cleavage sites of

NopT and YopT are polybasic (Shao et al., 2003a; Schmidt, 2011), in contrast to

AvrPphB, whose cleavage site is canonical with seven adjacent amino acids

involved (Kim et al., 2016). The different cleavage sites of YopT-type effectors

indicate that bacteria regulate different physiological pathways to establish specific

interactions with host plants.

Similar to the plant pathogenic effector AvrPphB, the protease activity of NopT

appears to be conserved for this function by cleaving the soybean PBS1 homolog

protein to restrict symbiosis (Khan et al., 2022). Indeed, transient expression of *S.*

fredii NopT in *A. thaliana* and *N. tabacum* could induce strong immune responses

and hypersensitive cell death (Fig. S8), indicating that rhizobial NopT might act as

an avirulent effector in some nonhost plant species. Interestingly, we failed to

obtain stable transgenic lines constitutively expressing NopT in *L. japonicus.*

However, the generation of transgenic root hairs was not affected, implying that

NopT might act as an Avr effector whose overexpression leads to whole plant

lethality but not root growth. Therefore, given this conservation of protease

function, it is consistent with a successful Avr effector that NopT has the ability to

proteolyze not only NFR5 but also Arabidopsis AtLYK5 and *L. japonicus* LjLYS11,

two receptors with chitin-binding affinities that mediate plant PTI responses (Cao

et al., 2014; Gysel et al., 2021). NFR5, AtLYK5, and LjLYS11 are LYKs belonging

to the same subfamily that lacks kinase activities. Thus, NopT may not only

suppress the nod factor–mediated symbiotic signal to avoid excessive infection by

S. fredii NGR234 but also function in inhibiting the signal triggered by chitin-like

compounds. The cleavage of receptor-like kinases (RLKs) may be an overlooked

strategy to regulate their function in plant biology compared to the well-studied

phosphorylation to control RLK activation. While cleavage regulation is well

studied in neurobiology, e.g., the Notch signal, the proteolysis of each Notch

molelue leads to signaling activation (Kopan and Ilagan, 2009). Arabidopsis BAK1,

identified as an universal co-receptor to regulate plant growth and MAMPs

perception, undergoes proteolytic cleavage, which is important for both BR

signaling and PTI response with an unclear mechanism (Zhou et al., 2019). NFR5

is a pseudokinase that functions as an output of NF signaling. The cleavage of

NFR5 by rhizobial NopT releases the cytoplasmic domain of NFR5, but the precise

function of whom is of interest to be determined.

The legume–rhizobial symbiosis is a unique case in the context of plant–microbe

interactions. Mutual regulation between pathogenic effectors and host proteins

has been reported in plant pathogenesis. For example, the *Pseudomonas*

conserved effector AvrPtoB could directly target key components in the immunity

pathway to promote its virulence (Goehre et al., 2008). However, the virulent

function of AvrPtoB could either be enhanced or dampened by host proteins.

AvrPtoB could be phosphorylated by SnRK2.8 to promote its virulence activity (Lei

et al., 2020). Plant lectin receptor-like kinase LexRK-IX.2, Pto, and Fen kinases

could also phosphorylate AvrPtoB to inactivate its E3 ligase activity and undermine

its ability to suppress PTI (Ntoukakis et al., 2009; Xu et al., 2020). A similar

strategy was used by legumes to dampen the "virulence" of the rhizobial effector

NopT, favoring mutualistic infection. NFR1 phosphorylates NopT at several

residues. The phosphomimetic NopTs (i.e., NopT^{T97D}, NopT^{S119D}, and NopT^{T257D})

are unable to cleave itself and NFR5 (Fig. 4C). The expression of these

phosphomimetic NopTs separately in *S. fredii* NGR234Δnopt mutant strains

cannot complement the hyperinfection phenotype (Fig.5F). These data indicate

that *L. japonicus* deploys NFR1 to phosphorylate NopT to dampen its catalytic

activity and may protect NFR5 from cleavage, favoring rhizobial infections.

However, NopT^{ΔN50}, mimic version of autocleaved NopT, retained the ability to

interact with NFR5 but not NFR1 (Fig. S2D). Moreover, intact NopT, but not

autocleaved NopT, are migrated by expressing NFR1^{CD} in immunoblotting (Fig. 3C

and 4B), suggesting autocleaved NopT may escape from being phosphorylated

and inactivated by NFR1.

In different rhizobium species, a variety of NopT homologs were characterized.

NopT homologs from some *Bradyrhizobium* species lose the ability to cleave

NFR5, while NopT homologs from some *Sinorhizobium*, truncated versions

without about 50 a.a. at its N-terminus, retain the ability to cleave NFR5 (Fig. 6A).

The type three secretion signal depends on the N-terminal of effectors (Akeda and

Galán, 2005). However, an earlier study has shown that NopT from *S. fredii* HH103

exhibited a positive effector on nodulation in soybean SN14 cultivar (Li et al.,

2023), suggesting that another unknown mechanism might be involved in

NopT$_{HH103}$ secretion. Interestingly, these truncated NopTs still keep the function to

cleave NFR5 (Fig. 6A) and autocleaved NopT$_{NGR234}$ cannot interact with NFR1

(Fig. S2D), indicating that NopTs from those strains and autocleaved NopT$_{NGR234}$

might lose the ability to be suppressed by NFR1. Notably, natural truncated NopT

from *S. fredii* USDA257 and *S. fredii* HH103 are the same in protein sequence, but

they lack the acylation site compared with $NopT_{NGR234}$. Therefore, $NopT_{USDA257}$

and $NopT_{HH103}$ lose the ability to suppress cell death triggered by NFR1 and NFR5

in *N. benthamiana* leaves (Fig. 6B) and failed to complement the massive infection

of NGR234*ΔnopT* (Fig. 6C and 6D), implying $NopT_{USDA257}$ and $NopT_{HH103}$ may

have a different target from $NopT_{NGR234}$ in the host plant. In addition, knockout of

NopT in *Bradyrhizobium* sp. ORS3257 was shown to result in uninfected nodule

formation, suggesting that, in this system, $NopT_{ORS3257}$ might play a role in nodule

formation (Teulet et al., 2019). The versatile function of NopT homologs in different

rhizobium species suggests that different regulatory processes might be involved

in specific rhizobial symbiosis.

In this study, we present a mutual regulation model in which the rhizobial effector

NopT directly targets NFR5 to suppress symbiosis signals and is then inactivated

by NFR1 *via* phosphorylation. NopT appears to retain some pathogenic-like

features in that it can cleave the *Arabidopsis* chitin receptor AtLYK5 and induce

cell death in some non-leguminous plants, such as *N. tabacum* and *Arabidopsis*.

Thus, the cleavage of NFR5 by NopT might be due to the structure similarity to

AtLYK5 targeted by NopT. The suppression of proteolytic activity of NopT *via*

phosphorylation by NFR1 appears to work as a feedback loop to allow symbiotic

signaling in compatible hosts. The model (Fig. 6E) provides essential insights into

the legume–rhizobial symbiosis, where rhizobia deploy an effector to dampen

symbiotic signaling, and legumes slightly recur infection by suppressing NopT to

employthe nod factor signal in response. Even though this couple, legume and

rhizobia, tries to get together and strives to improve their living conditions, we

catch a glimpse into the arguments and the recovery conversations between them.

Materials and Methods

Plant materials

Lotus japonicus Gifu B–129 cultivar was provided by the Center for Carbohydrate

Recognition and Signaling (https://lotus.au.dk/), and used as wild type for nodulation

assays. The Gifu *pNIN*:*GUS* transgenic line was kindly provided by Dr. J. Stougaard from

the Aarhus University (Heckmann et al., 2011). All *L. japonicus* seeds were treated with

concentrated sulfuric acid for 10 min, followed by surface sterilization in 1% (w/v) NaClO

for 8 min. After synchronization of germination at 4°C in the dark for 2 d, the seeds were

placed on half–strength Murashige & Skoog (MS) medium containing 0.8% (w/v) agar for

germination at 22°C for 2 d in the dark. Seedlings were then transferred to growth pots

containing vermiculite supplied with half–strength B&D (Broughton & Dilworth) medium

without nitrogen under long–day conditions (16–h light/8–h dark) at 22°C. Ten–day–old

seedlings were inoculated with *Sinorhizobium fredii* NGR234 (OD_{600}=0.02) for infection

assays.

Plasmid construction

All plasmids used in this study were generated using MultiF Seamless Assembly Mix

(ABclonal, catalog number RK21020). The plasmid pGWB514 was used as an original

plasmid for all binary vector construction (Nakagawa et al., 2007). Briefly, Different tags

including HA tag, Myc tag, FLAG tag, Strep tag, nLUC, cLUC, and GFP were respectively

amplified using a forward primer containing a *Kpn*I site and ligated into pGWB514

digested with *Xba*I and *Sac*I using seamless cloning method as described above. The

generated plasmids were named pG5XX–HA, pG5XX–Myc, pG5XX–FLAG,

pG5XX–nLUC, pG5XX–cLUC, pG5XX–GFP, respectively. The coding sequences of *L.*

japonicus NFR1 and *NFR5* were cloned into pGWB514 and pG517 between *Xba*I/*Kpn*I to

generate NFR1–HA and NFR5–Myc fusion constructs under the control of the 35S

promoter. then, the DNA fragment containing *pro35S*:NFR5–Myc–NosT was amplified

and cloned into pG514–NFR1 at *Sbf*I. The final plasmids contained

pro35S:NFR1–HA–NosT and *pro35S*:NFR5–Myc–NosT. Rhizobial genes encoding

effectors from *S. fredii* NGR234 were cloned into pG5xx–FLAG between *Xba*I/*Kpn*I. To

detect the interaction between NFR1/NFR5 and NopT in *N. benthamiana*, the full–length

NFR1/NFR5, *NopT* and *AtFLS2* coding sequences were cloned into pG5xx–Nluc,

pG5xx–Nluc, pSPYCE(MR) and pSPYNE(R)173 (Waadt et al., 2008). For cleavage

assay in *N. benthamiana*, the *NFR5* and *NopT* coding sequences were cloned into

pG5XX–FLAG GFP and pG5xx–Strep respectively. For cleavage assay in *E. coli*, the

sequence encoding the cytoplasmic domain of NFR5 was cloned into pACYC duet

Strep–HA, the sequences encoding the kinase domain of NFR5 fused to HA tag and the

juxtamembrane domain of NFR5 fused to the GFP tag was cloned into pSUMO. The point

mutants in NFR5^{CD} and NopT were created from the pACYC duet Strep–NFR5^{CD}–HA and

pET28a NopT–FLAG template using site–directed mutagenesis PCR. For *in vivo* cleavage

assay in *L. japonicus*, NopT-FLAG or NopT^{C93S}-FLAG was cloned into pUB GFP between

*Xba*I and *Kpn*I, and then 35S-NFR5-Myc-NosT was cloned in pUB NopT GFP at *Pst*I.

For phosphorylation assay, the sequence encoding the cytoplasmic domain of NFR1 fused

to a Myc tag was cloned into pCDF duet. For rhizobium infection assays, 300 bp upstream

of *NopT* and site mutants of *NopT* were fused by overlapping PCR and cloned into pHC60.

Transient expressing and in *Nicotiana* species

The plasmids for transient expression of INF1, Avr3 and R3a in *N. benthamiana* were

kindly gifted by Dr. Juan Du from HZAU. Agrobacterium (*Agrobacterium tumefaciens*)

strain EHA105 carrying various constructs were mixed with an Agrobacterium culture

harboring the P19 suppressor in an infiltration buffer (10 mM $MgCl_2$, 10 mM MES–KOH

pH 5.8 and 200 μM acetosyringone). After incubating 2 h at room temperature,

Agrobacterium cultures were infiltrated into the leaves of 4–week–old *N. benthamiana*

and *N. tabacum*. Leaves were harvested for analysis at two days post infiltration. The cell

death suppression assay in *N. benthamiana* was performed as described previously (Wang

et al., 2011).

Hairy Root Transformation in *Lotus japonicus*

The hairy root transformation assays were performed as described previously(Li et al.,

2018). In brief, surface-sterilized seeds were germinated and grown in the MS plate

without sucrose (23°C dark for first 3 days and 23°C 16 h light/8 h dark for next two days).

Then seeds were cut at the middle of the hypocotyl and co-cultivation with *A. rhizogenes*

carry pUB GFP and pUB NopT-FLAG GFP respectively (23°C dark for first 3 day and

23°C 16 h light/8 h dark for next 2 day). Transfer the plants onto B5 media. After about 15

days, the plants were picked up for further analysis by cutting the root without expressing

florescent proteins. For rhizobial infection assays, data were collected at 9dpi for *S.fredii*

NGR234 and 6dpi for MAFF303099

NFR5 cytoplasmic domain cleavage assay

For cleavage assay in *E. coli*, Strep–NFR5^{CD}–HA or Strep–NFR5^{CD}–HA were

co–expressed with NopT–FLAG or NopT^{C93S}–FLAG by using Duet expressing vectors

(Novagen) (Han et al., 2017). Bacterial cell cultures were treated with 0.5 mM IPTG at

22°C for 15 h. The bacterial pellet was resuspended in extraction buffer consisting of

Tris–HCl (pH 7.3), 150 mM NaCl, 5 mM EDTA, 0.5% (w/v) SDS and 0.5% (v/v) Triton

X–100. After adding SDS loading buffer and boiling at 100°C for 5 min, total proteins

were detected with anti–HA–peroxidase (Sigma, clone 3F10), anti–FLAG (Sigma, F1804),

and anti–Strep (ABclonal, AB_2863792) antibodies.

For *in vitro* cleavage assays, Sumo–NFR5^{JM}–GFP and NopT–FLAG recombinant proteins

were purified, and 3 μg NopT–FLAG and 1 μg Sumo–NFR5^{JM}–GFP were incubated in a

test tube at 22°C for 7 h. The cleavage products were detected with anti–GFP antibody by

immunoblot (ABclonal, AB_2770402).

For in vivo cleavage assays, the transgenic root was collected at about 15 days after

induction of hairy roots. 2% protease inhibitors (Sigma, P9599) were added in extraction

buffer. The actin proteins were detected with anti-Actin antibody (ABclonal, AC009).

Phosphorylation assays

For phosphorylation assays in *E. coli*, His–NFR1^{CD}–Myc or His–NFR1$^{CD–K351E}$–Myc and

NopT–FLAG–His were co–expressed by using Duet expressing vectors (Novagen) with

0.5 mM IPTG at 28°C for 18 h. After purifying NFR1^{CD} and NopT at the same time, CIAP

(TaKaRa, 2250B) was added and incubated at 37°C for 3 h. The mobility shift of NFR1^{CD}

and NopT was detected with anti–Myc (Bio–Legend, 626808) and anti–FLAG (Sigma,

F1804) antibodies by immunoblot analysis.

For phosphorylation assays in *L. Japonicus* roots, NopT^{C93S}-FLAG was expressed in the

nfr1 mutant and wild type plants, respectively. Transfer the plant with florescent root to

half–strength B&D media for 5 days. After treat with water and rhizobia respectively, the

total proteins were drawn by extraction buffer with addition 2% protease inhibitors

cocktail (Sigma, P9599) and 2% phosphatase inhibitors cocktail (Yeasen,20109-A), then

followed with TCA precipitation to remove various contaminants (e.g. EDTA, surfactant).

Phosphorylation of NopT^{C93S} proteins was analyzed by 50μM Zn^{2+}-Phos-tag SDS-PAGE

and detected with anti–FLAG (Sigma, F1804) antibodies by immunoblot analysis.

BiFC and Split–LUC assays in *N. benthamiana*

In BiFC assay, the eYFP fluorescence of fusion proteins was assayed 2–3 days after

infiltration using a confocal microscope (Leica TCS SP8). For the Split–LUC assay, *N.*

benthamiana leaves were collected at 2 dpi sprayed with a solution of 1 mM D–luciferin

(Promega, E1603) and 0.02% (v/v) Triton X–100. After a dark–adaptation of 5 min,

bioluminescence images were acquired using a Tanon Bio–Imaging System (Tanon,

Shanghai, 4600).

Liquid chromatography mass spectrometry

For proteomics analysis of NopT phosphorylation sites, *NopT* and *NFR1–CD* or

NFR1–CD^{K351E} were co–expressed in *E. coli*, then proteins were purified using

Ni–Charged Resin (GenScript, Cat. No. L00223). The bands of NopT^P and NopT, as

control, were cut and destained by 50 mM triethylammonium bicarbonate (TEAB) and

50% (v/v) acetonitrile (ACN) in 50 mM TEAB followed by washing with 100% ACN until

the gel turned white. The protein bands were then in–gel digested with sequencing–grade

trypsin. LC–MS analysis was performed by Novogene Co., Ltd. (Beijing, China).

For identificaiton of the cleavage site on NFR5, NopT and the recombinant protein,

His-Sumo-NFR5-GST, were co-expressed in *E. coli* followed by protein purification using

Glutathione Resin (GenScript, Cat. No. L00206). About 80μg cleaved NFR5 was collected

for analysis. The protein bands were digested with trypsin,glu-C, chymotrypsin and pepsin.

LC–MS analysis was performed by Oulu Biotechnology Co., Ltd. (Shanghai, China).

Supplemental data

Supplemental Figure 1. Analyses of the functions of all 15 effectors from *S. fredii*

NGR234 in preventing LjNFR1- and LjNFR5-induced cell death in *N. benthamiana*

leaves. Supplemental Fig. 2.   Split–LUC assay testing the interactions between

different NopT mutant and NFRs.

Supplemental Figure 2.   Split–LUC assay testing the interactions between

different NopT mutants and NFRs.

Supplemental Figure 3. NopT cleaves NFR5 at the juxtamembrane domain.

Supplemental Figure 4. Characterization of cleavage site of LysM receptor.

Supplemental Figure 5. The cytoplasmic domains (CDs) of NFR1, NFR5, and

kinase–dead NFR1CDK351E were used for kinase assays in E. coli.

Supplemental Figure 6. Mutant version of NFR5 fails to form infections.

Supplemental Figure 7. Phylogenetic tree based on the amino acid sequence of NopT homologs from different bacterial species.

Supplemental Figure 8. Transient expression of NopT triggers cell death in Arabidopsis and Nicotiana tabacum.

Supplemental Table 1. Phosphopeptides identified by Liquid Chromatography–Mass spectrometry.

Supplemental Table 2. Primers used in this study.

Acknowledgments: We greatly thank Prof. Zhongming Zhang for his valuable suggestions in this project. We thank Drs. E. Giraud, E. Wang, and Y. Liang for their critical reading of this manuscript. We thank Dr. J. Staugaard for kindly providing *pNIN:GUS* transgenic seed; Dr. J. Du for providing plasmids expressing of Avr3a and R3a, BAX, and INF1. The qPCR and microscopy data were acquired from the Core Facility Center run by the National Key Lab of Agricultural Microbiology.

Funding: This work was supported by the National Key R&D Program of China (2019YFA0904700), the National Natural Science Foundation of China (32090063), and the Fundamental Research Funds for the Central Universities (2662022SKYJ002). Work in the Stacey's Lab was supported by a grant from National Science Foundation, Plant Genome Research Project (IOS–1734145).

Author Contributions: H.B., Y.C., and G.S. conceived the idea and designed the research, S.F.W. and C.S. provided the NGR234 *nopt* strain; H.B., Y.W., Y.L., and Y.Y.

771 performed experiments; H.B., Y.W., H. L., H.Z., and Y.C. analyzed the data, Q.W and S.X

772 identified the cleavage site by MS. H.B., Y.C., and G.S. wrote the paper; S.F.W. and C.S.

773 helped to revise the MS.

References

Akeda, Y., and Galán, J.E. (2005). Chaperone release and unfolding of substrates in
type III secretion. Nature **437,** 911-915.

Ausmees, N., Kobayashi, H., Deakin, W.J., Marie, C., Krishnan, H.B., Broughton,
W.J., and Perret, X. (2004). Characterization of NopP, a type III secreted effector of sp.
strain NGR234. J Bacteriol **186,** 4774-4780.

Bos, J.I., Kanneganti, T.D., Young, C., Cakir, C., Huitema, E., Win, J., Armstrong,
M.R., Birch, P.R., and Kamoun, S. (2006). The C-terminal half of *Phytophthora*
infestans RXLR effector AVR3a is sufficient to trigger R3a-mediated hypersensitivity and
suppress INF1-induced cell death in *Nicotiana benthamiana*. Plant J **48,** 165-176.

Bozsoki, Z., Cheng, J., Feng, F., Gysel, K., Vinther, M., Andersen, K.R., Oldroyd, G.,
Blaise, M., Radutoiu, S., and Stougaard, J. (2017). Receptor-mediated chitin
perception in legume roots is functionally separable from Nod factor perception. Proc Natl
Acad Sci U S A **114,** E8118-E8127.

Broghammer, A., Krusell, L., Blaise, M., Sauer, J., Sullivan, J.T., Maolanon, N.,
Vinther, M., Lorentzen, A., Madsen, E.B., Jensen, K.J., Roepstorff, P., Thirup, S.,
Ronson, C.W., Thygesen, M.B., and Stougaard, J. (2012). Legume receptors perceive
the rhizobial lipochitin oligosaccharide signal molecules by direct binding. Proc Natl Acad
Sci U S A **109,** 13859-13864.

Cao, Y., Liang, Y., Tanaka, K., Nguyen, C.T., Jedrzejczak, R.P., Joachimiak, A., and
Stacey, G. (2014). The kinase LYK5 is a major chitin receptor in *Arabidopsis* and forms a
chitin-induced complex with related kinase CERK1. eLife **3**.e03766.

Dai, W.J., Zeng, Y., Xie, Z.P., and Staehelin, C. (2008). Symbiosis-promoting and

deleterious effects of NopT, a novel type 3 effector of *Rhizobium* sp. strain NGR234. J

Bacteriol **190,** 5101-5110.

Dowen, R.H., Engel, J.L., Shao, F., Ecker, J.R., and Dixon, J.E. (2009). A family of

bacterial cysteine protease type III effectors utilizes acylation-dependent and

-independent strategies to localize to plasma membranes. J Biol Chem **284,**

15867-15879.

Fotiadis, C.T., Dimou, M., Georgakopoulos, D.G., Katinakis, P., and Tampakaki, A.P.

(2012). Functional characterization of NopT1 and NopT2, two type III effectors of

Bradyrhizobium japonicum. FEMS Microbiol Lett **327,** 66-77.

Gan, Y., Zhang, L., Zhang, Z., Dong, S., Li, J., Wang, Y., and Zheng, X. (2009). The

LCB2 subunit of the sphingolip biosynthesis enzyme serine palmitoyltransferase can

function as an attenuator of the hypersensitive response and Bax-induced cell death.

New Phytol **181,** 127-146.

Geurts, R., Fedorova, E., and Bisseling, T. (2005). Nod factor signaling genes and their

function in the early stages of infection. Curr Opin Plant Biol **8,** 346-352.

Goehre, V., Spallek, T., Haeweker, H., Mersmann, S., Mentzel, T., Boller, T., de

Torres, M., Mansfield, J.W., and Robatzek, S. (2008). Plant pattern-recognition

receptor FLS2 is directed for degradation by the bacterial ubiquitin ligase AvrPtoB. Curr

Biol **18,** 1824-1832.

Gysel, K., Laursen, M., Thygesen, M.B., Lironi, D., Bozsoki, Z., Hjuler, C.T.,

Maolanon, N.N., Cheng, J., Bjork, P.K., Vinther, M., Madsen, L.H., Rubsam, H.,

Muszynski, A., Ghodrati, A., Azadi, P., Sullivan, J.T., Ronson, C.W., Jensen, K.J.,

Blaise, M., Radutoiu, S., Stougaard, J., and Andersen, K.R. (2021). Kinetic

proofreading of lipochitooligosaccharides determines signal activation of symbiotic plant

receptors. Proc Natl Acad Sci U S A **118**.e2111031118.

Han, Y., Sun, J., Yang, J., Tan, Z., Luo, J., and Lu, D. (2017). Reconstitution of the plant ubiquitination cascade in bacteria using a synthetic biology approach. Plant J **91,** 766-776.

Heckmann, A.B., Sandal, N., Bek, A.S., Madsen, L.H., Jurkiewicz, A., Nielsen, M.W., Tirichine, L., and Stougaard, J. (2011). Cytokinin induction of root nodule primordia in *Lotus japonicus* is regulated by a mechanism operating in the root cortex. Mol Plant Microbe Interact **24,** 1385-1395.

Jones, J.D.G., and Dangl, J.L. (2006). The plant immune system. Nature **444,** 323-329.

Kawaharada, Y., Kelly, S., Nielsen, M.W., Hjuler, C.T., Gysel, K., Muszynski, A., Carlson, R.W., Thygesen, M.B., Sandal, N., Asmussen, M.H., Vinther, M., Andersen, S.U., Krusell, L., Thirup, S., Jensen, K.J., Ronson, C.W., Blaise, M., Radutoiu, S., and Stougaard, J. (2015). Receptor-mediated exopolysaccharide perception controls bacterial infection. Nature **523,** 308-312.

Khan, A., Wadood, S.F., Chen, M., Wang, Y., Xie, Z.P., and Staehelin, C. (2022). Effector-triggered inhibition of nodulation: A rhizobial effector protease targets soybean kinase GmPBS1-1. Plant Physiol **189,** 2382-2395.

Kim, S.H., Qi, D., Ashfield, T., Helm, M., and Innes, R.W. (2016). Using decoys to expand the recognition specificity of a plant disease resistance protein. Science **351,** 684-687.

Kimbrel, J.A., Thomas, W.J., Jiang, Y., Creason, A.L., Thireault, C.A., Sachs, J.L., and Chang, J.H. (2013). Mutualistic co-evolution of type III effector genes in Sinorhizobium fredii and *Bradyrhizobium japonicum*. PLoS Pathog **9,** e1003204.

Kopan, R., and Ilagan, M.X.G. (2009). The canonical notch signaling pathway: Unfolding the activation mechanism. Cell **137,** 216-233.

Lei, L., Stevens, D.M., and Coaker, G. (2020). Phosphorylation of the *Pseudomonas* effector AvrPtoB by *Arabidopsis* SnRK2.8 is required for bacterial virulence. Mol Plant **13,** 1513-1522.

Li, D.D., Zhu, Z.K., Deng, X.M., Zou, J.N., Ma, C., Li, C.D., Yin, T., Liu, C.Y., Wang,

J.H., Chen, Q.S., and Xin, D.W. (2023). GmPBS1, a hub gene interacting with Rhizobial

type-III effectors NopT and NopP, regulates soybean nodulation. Agronomy **13**.1242.

Li, H., Chen, M., Duan, L., Zhang, T., Cao, Y., and Zhang, Z. (2018). Domain swap

approach reveals the Ccritical roles of different domains of SYMRK in root nodule

symbiosis in *Lotus japonicus.* Front Plant Sci **9,** 697.

Li, L., Kim, P., Yu, L., Cai, G., Chen, S., Alfano, J.R., and Zhou, J.M. (2016).

Activation-dependent destruction of a co-receptor by a *Pseudomonas syringae* effector

dampens plant immunity. Cell Host Microbe **20,** 504-514.

Madsen, E.B., Madsen, L.H., Radutoiu, S., Olbryt, M., Rakwalska, M., Szczyglowski,

K., Sato, S., Kaneko, T., Tabata, S., Sandal, N., and Stougaard, J. (2003). A receptor

kinase gene of the LysM type is involved in legume perception of rhizobial signals. Nature

425, 637-640.

Madsen, E.B., Antolin-Llovera, M., Grossmann, C., Ye, J., Vieweg, S., Broghammer,

A., Krusell, L., Radutoiu, S., Jensen, O.N., Stougaard, J., and Parniske, M. (2011).

Autophosphorylation is essential for the *in vivo* function of the *Lotus japonicus* Nod factor

receptor 1 and receptor-mediated signalling in cooperation with Nod factor receptor 5.

Plant J **65,** 404-417.

Moling, S., Pietraszewska-Bogiel, A., Postma, M., Fedorova, E., Hink, M.A.,

Limpens, E., Gadella, T.W., and Bisseling, T. (2014). Nod factor receptors form

heteromeric complexes and are essential for intracellular infection in medicago nodules.

Plant Cell **26,** 4188-4199.

Nakagawa, T., Suzuki, T., Murata, S., Nakamura, S., Hino, T., Maeo, K., Tabata, R.,

Kawai, T., Tanaka, K., Niwa, Y., Watanabe, Y., Nakamura, K., Kimura, T., and

Ishiguro, S. (2007). Improved Gateway binary vectors: high-performance vectors for

creation of fusion constructs in transgenic analysis of plants. Biosci Biotechnol Biochem

71, 2095-2100.

Ntoukakis, V., Mucyn, T.S., Gimenez-Ibanez, S., Chapman, H.C., Gutierrez, J.R.,

Balmuth, A.L., Jones, A.M., and Rathjen, J.P. (2009). Host inhibition of a bacterial

virulence effector triggers immunity to infection. Science **324,** 784-787.

Schmidt, G. (2011). Yersinia enterocolitica outer protein T (YopT). Eur J Cell Biol **90,**

955-958.

Schubert, K.A., Xu, Y., Shao, F., and Auerbuch, V. (2020). The type III secretion

system as a tool for studying cytosolic innate immune surveillance. Ann Rev Microbiol,

Vol 74, 2020 **74,** 221-245.

Shao, F., Merritt, P.M., Bao, Z., Innes, R.W., and Dixon, J.E. (2002). A *Yersinia* effector

and a *Pseudomonas* avirulence protein define a family of cysteine proteases functioning

in bacterial pathogenesis. Cell **109,** 575-588.

Shao, F., Vacratsis, P.O., Bao, Z., Bowers, K.E., Fierke, C.A., and Dixon, J.E.

(2003a). Biochemical characterization of the *Yersinia* YopT protease: cleavage site and

recognition elements in Rho GTPases. Proc Natl Acad Sci U S A **100,** 904-909.

Shao, F., Golstein, C., Ade, J., Stoutemyer, M., Dixon, J.E., and Innes, R.W. (2003b).

Cleavage of *Arabidopsis* PBS1 by a bacterial type III effector. Science **301,** 1230-1233.

Sugawara, M., Takahashi, S., Umehara, Y., Iwano, H., Tsurumaru, H., Odake, H.,

Suzuki, Y., Kondo, H., Konno, Y., Yamakawa, T., Sato, S., Mitsui, H., and

Minamisawa, K. (2018). Variation in bradyrhizobial NopP effector determines symbiotic

incompatibility with soybeans via effector-triggered immunity. Nature Commun **9**.3139.

Sun, Y., Li, L., Macho, A.P., Han, Z., Hu, Z., Zipfel, C., Zhou, J.-M., and Chai, J.

(2013). Structural basis for flg22-induced activation of the *Arabidopsis* FLS2-BAK1

immune complex. Science **342,** 624-628.

Tang, D.Z., Wang, G.X., and Zhou, J.M. (2017). Receptor kinases in plant-pathogen

interactions: More than pattern recognition. Plant Cell **29,** 618-637.

Teulet, A., Busset, N., Fardoux, J., Gully, D., Chaintreuil, C., Cartieaux, F., Jauneau,

A., Comorge, V., Okazaki, S., Kaneko, T., Gressent, F., Nouwen, N., Arrighi, J.F.,

Koebnik, R., Mergaert, P., Deslandes, L., and Giraud, E. (2019). The rhizobial type III

effector ErnA confers the ability to form nodules in legumes. Proc Natl Acad Sci U S A

116, 21758-21768.

Vleeshouwers, V.G., Driesprong, J.D., Kamphuis, L.G., Torto-Alalibo, T., Van't Slot,

K.A., Govers, F., Visser, R.G., Jacobsen, E., and Kamoun, S. (2006).

Agroinfection-based high-throughput screening reveals specific recognition of INF

elicitins in *Solanum*. Mol Plant Pathol **7,** 499-510.

Wang, D., Yang, S., Tang, F., and Zhu, H. (2012). Symbiosis specificity in the legume:

rhizobial mutualism. Cell Microbiol **14,** 334-342.

Wang, Q., Han, C., Ferreira, A.O., Yu, X., Ye, W., Tripathy, S., Kale, S.D., Gu, B.,

Sheng, Y., Sui, Y., Wang, X., Zhang, Z., Cheng, B., Dong, S., Shan, W., Zheng, X.,

Dou, D., Tyler, B.M., and Wang, Y. (2011). Transcriptional programming and functional

interactions within the *Phytophthora sojae* RXLR effector repertoire. Plant Cell **23,**

2064-2086.

Xiang, T., Zong, N., Zou, Y., Wu, Y., Zhang, J., Xing, W., Li, Y., Tang, X., Zhu, L.,

Chai, J., and Zhou, J.M. (2008). *Pseudomonas syringae* effector AvrPto blocks innate

immunity by targeting receptor kinases. Curr Biol **18,** 74-80.

Xu, N., Luo, X., Wu, W., Xing, Y., Liang, Y., Liu, Y., Zou, H., Wei, H.L., and Liu, J.

(2020). A plant lectin receptor-like kinase phosphorylates the bacterial effector AvrPtoB to

dampen lits virulence in *Arabidopsis*. Mol Plant **13,** 1499-1512.

Yang, J., Lan, L., Jin, Y., Yu, N., Wang, D., and Wang, E. (2022). Mechanisms

underlying legume-rhizobium symbioses. J Integr Plant Biol **64,** 244-267.

Zhang, B., Wang, M., Sun, Y., Zhao, P., Liu, C., Qing, K., Hu, X., Zhong, Z., Cheng, J.,

Wang, H., Peng, Y., Shi, J., Zhuang, L., Du, S., He, M., Wu, H., Liu, M., Chen, S.,

Wang, H., Chen, X., Fan, W., Tian, K., Wang, Y., Chen, Q., Wang, S., Dong, F., Yang,

C., Zhang, M., Song, Q., Li, Y., and Wang, X. (2021). *Glycine max* NNL1 restricts

symbiotic compatibility with widely distributed *Bradyrhizobia via* root hair infection. Nat

Plants **7,** 73-86.

Zhang, J., Li, W., Xiang, T., Liu, Z., Laluk, K., Ding, X., Zou, Y., Gao, M., Zhang, X.,

Chen, S., Mengiste, T., Zhang, Y., and Zhou, J.M. (2010). Receptor-like cytoplasmic

kinases integrate signaling from multiple plant immune receptors and are targeted by a

Pseudomonas syringae effector. Cell Host Microbe **7,** 290-301.

Zhou, J.G., Wang, P., Claus, L.A.N., Savatin, D.V., Xu, G.Y., Wu, S.J., Meng, X.Z.,

Russinova, E., He, P., and Shan, L.B. (2019). Proteolytic orocessing of SERK3/BAK1

regulates plant immunity, development, and cell death. Plant Physiol **180,** 543-558.

Zipfel, C., and Oldroyd, G.E.D. (2017). Plant signalling in symbiosis and immunity.

Nature **543,** 328-336.

Zipfel, C., Robatzek, S., Navarro, L., Oakeley, E.J., Jones, J.D.G., Felix, G., and

Boller, T. (2004). Bacterial disease resistance in through flagellin perception. Nature **428,**

764-767.

967   **Supplemental Figures**

Effector	Leaf Discs (Cell death/total)	
	Ctrl.	NFR1&NFR5
NopA	0/3	9/9
NopB	0/3	9/9
NopC	0/3	9/9
NopJ	0/3	9/9
NopL	0/3	9/9
NopM	3/3	9/9
NopP	0/3	9/9
NopT	0/3	0/9
NopU	0/3	9/9
NopV	0/3	9/9
NopX	0/3	9/9
NopY	0/3	9/9
NopZ	0/3	4/9
NopBE	0/3	9/9
NopBI	0/3	9/9

Supplemental Fig. 1. Analyses of the functions of all 15 effectors from *S. fredii* NGR234 in preventing LjNFR1- and LjNFR5-induced cell death in *N. benthamiana* leaves.

15 genes from *S. fredii* NGR234 encoding effector proteins were amplified and cloned into binary vector for expression of Strep-tagged proteins in *N. benthamiana*. LjNFR1 and LjNFR5 from *L. japonicus* were C-terminally tagged with HA and Myc, respectively, and cloned into binary vectors for expression in *N. benthamiana*. All plasmids were electroporated into *Agrobacterium tumefaciens* EH105.

(A) *Agrobacterium* strain harboring each effector gene or control empty vector (EV) were infiltrated into *N. benthamiana*. 12 hour post infiltration (hpi), agrobacterium strains harboring LjNFR1 and LjNFR5 were mixed and hand-infiltrated in the same leaf discs as shown in (B). 48 hpi, cell death

phenotypes were evaluated (C). At least three replicates in the same leaf were

performed to test the function of each effector in suppressing cell death-triggered

by LjNFR1 and LjNFR5. (D) Results of all 15 effector proteins in suppressing

programmed cell death by overexpression of NFR1 and NFR5. (E) Abundance of

LjNFR1, LjNFR5, and NopT proteins, as measured by immunoblot with specific

antibodies. Leaf discs expressing LjNFR1, LjNFR5, and NopT or LjNFR1 and

LjNFR5 and with EV were used for immunoblot analyses. Actin was used as

loading control. (F) Another example of suppression of programmed cell death by

NopT in the leaf discs coexpressing NFR1 and NFR5. Rep1, Rep2, and Rep3

represent three technical repeats in the same leave.

996
997

Supplemental Fig. 2. Split–LUC assay testing the interactions between different NopT mutants and NFRs.

(A)The interaction between NopT with mutated acylation site and NFR1/NFR5. NopT^{AAA} (G50A C51A C52A). (B) The protein expressing of indicating genes in the leave of (A). (C) Nop^{C93S} interact with NFR1 and NFR5. (D) The interaction between NopT$^{\Delta N50}$ and NFR5. NopT$^{\Delta N50}$ represents a truncated NopT with deletion of about 50 a.a. from its N–terminus.

1000

1001

1002

1003

1004

1005

1006

1007

1008

1009

1010

1011

1012

Supplemental Fig. 3. NopT cleaves NFR5 at the juxtamembrane domain.

(A) NopT cannot cleave NFR1-GFP expressed in *N. benthamiana* leaves. recombinant

Sumo-NFR5^{KD}-HA lacking the juxtamembrane (JM) domain. (B) NopT could not cleave

Sumo–NFR5^{KD}–HA, a recombinant protein expressed in *E. coli* with the juxtamembrane

domain of NFR5 was replaced with Sumo tag. (C) Cleavage assay using two mutant

versions of NFR5^{CD} in the presence of NopT or AvrPphB. NFR5$^{CD288-294A}$ represents a

mutant NFR5^{CD} with residues 288 to 294 replaced with seven alanines; NFR5$^{CD288-294PphB}$

represents a mutant NFR5^{CD} with residues 288 to 294 replaced with AvrPphB recognition

sites. (D) Autocleavage assay, using different NopT variants with single amino acid

mutations. H205A and D220A indicate His-205 and Asp-220 replaced with alanine,

respectively. (E-H) NopT cleavage assay using 19 variants of NFR5^{CD} with three adjacent

amino acids replaced with three alanines. The numbers indicate the location of the three

residues replaced with alanines. (I) NopT cleavage assay using seven truncated variants of

NFR5^{CD} with 10 adjacent amino acids deleted. The numbers indicate the location of the 10

deleted residues. (J) NopT but not NopT^{C93S} could cleave the CDs of AtLYK5 and

LjLYS11 expressed in E. coli. NopT could cleave both AtLYK5^{JM}–NFR5^{KC},

LjLYS11^{JM}–NFR5^{KC} and LjNFR1^{JM}–NFR5^{KC}, three recombinant proteins expressed in *E.*

coli with the JM of NFR5 was replaced with the JMs from AtLYK5, LjLYS11 and NFR1,

respectively. (K) NopT but not NopT^{C93S} could cleave recombinant protein

NFR5$^{268\text{-}445}$-NFP$^{458\text{-}595}$ and NFP$^{270\text{-}457}$-NFR$^{5456\text{-}595}$.

1033

Supplemental Fig. 4. Characterization of cleavage site of LysM receptor.

(A) Domain and motif annotation of NFR5. (B) Protein sequence alignment of the

cytoplasmic domain of AtLYK5, LjLYS11, LjNFR5 and MtNFP. Yellow boxes delineated

highly conserved residues (S283, G294, Y303, A310, T311) and brown box delineated

kinase domain. Green frame delineated cleavage site of NFR5 and red frame delineated

residues similar to NopT autocleavage region. (C) Mass spectrometry analysis of cleaved

NFR5. Blue lines indicated the peptides characterized by MS.

1042

1043

1044

1045 **Supplemental Fig. 5. The cytoplasmic domains (CDs) of NFR1, NFR5, and**

1046 **kinase–dead NFR1CD^{K351E} were used for kinase assays in *E. coli*.** Proteins

1047 were detected by immunoblotting using anti–HA and anti–Myc antibodies.

1048 Asterisk indicated the band retardation on the gel representing the

1049 phosphorylated NFR5^{CD} proteins.

1050

1051

Supplemental Fig. 6. Rhizobial infection in the mutant versions of NFR5.

1052

(A) NFR5^{5m} failed to form rhizobial infection (n=7, Student's t–test: $P < 0.01$).

1053

(B)NFR5-NFP^{KD-CT} could recuse the rhizobial infection of wild type NGR234

1054

(n=10, Student's t–test: $P < 0.01$). The transgenic roots expressing NFR5 or NFR5^{5m}

1055

or NFR5-NFP^{KCT} in the *nfr5-3* mutant under the control of native promoter were

1056

inoculated with GFP-labelled wild type NGR234 and NGR234Δ*nopT* respectively.

1057

The data was collected 9 days post inoculation.

1058

1059

Sinorhizobium fredii HH103
Sinorhizobium fredii USDA257
Sinorhizobium fredii CCBAU45436
NopT Sinorhizobium fredii NGR234
Ensifer sp. MPMI2T
Bradyrhizobium sp. Ec3.3
Bradyrhizobium sp. NAS96.2
Bradyrhizobium sp. 32362
Bradyrhizobium sp. CCBAU051011
2 Ensifer sp. MPMI2T
Sinorhizobium sp. BJ1
Sinorhizobium sp. PC2
NopT2 Bradyrhizobium diazoefficiens USDA110
Bradyrhizobium elkanii USDA61
Bradyrhizobium sp. CCBAU53421
Bradyrhizobium sp. CCBAU51765
NopT Bradyrhizobium sp. ORS3257
Bradyrhizobium sp. CIR3A
Bradyrhizobium sp. Rc2d
Bradyrhizobium sp. 281
Bradyrhizobium sp. CCBAU51753
Bradyrhizobium sp. CCBAU43298
NopT1 Bradyrhizobium diazoefficiens USDA110
Bradyrhizobium sp. AC87j1
Bradyrhizobium sp. IAR9
Bradyrhizobium sp. CIR18
Bradyrhizobium sp. CIR48
Acidovorax citrulli
Acidovorax avenae
Acidovorax oryzae
Pseudomonas savastanoi pv. phaseolicola
Pseudomonas syringae pv. actinidiae ICMP 9853
Pseudomonas tremae
Ralstonia sp. A12
Xanthomonas oryzae
Xanthomonas bromi
Xanthomonas translucens pv. graminis ART-Xtg29
Xanthomonas translucens pv. graminis
Pseudomonas syringae
Pseudomonas syringae pv. daphniphylli
Pseudomonas amygdali pv. morsprunorum
Pseudomonas caricapapayae
Pantoea agglomerans
Pseudomonas sp. KBS0707
Pseudomonas savastanoi pv. glycinea
Pseudomonas savastanoi pv. nerii
Pseudomonas meliae
Pseudomonas ficuserectae
Pseudomonas amygdali pv. sesami
Pseudomonas cannabina
Ralstonia_solanacearum
Ralstonia pseudosolanacearum
Pasteurella multocida
Vibrio sp. ED002
Vibrio jasicida
Vibrio harveyi
Yersinia enterocolitica
Yersinia pseudotuberculosis
Yersinia pestis
Photorhabdus laumondii subsp. laumondii
Photorhabdus laumondii
Photorhabdus temperata
Photorhabdus sp. APURE
Photorhabdus heterorhabditis
Photorhabdus cinerea
Photorhabdus asymbiotica
Photorhabdus akhurstii
Photorhabdus hindustanensis
Photorhabdus luminescens

1060

0.50

1061 **Supplemental Fig. 7. Phylogenetic tree based on the amino acid sequence of**

1062 **NopT homologs from different bacterial species.**

1063

1064

1065

1066

1067

1068

1069

1070

1071

Supplemental Fig. 8. Transient expression of NopT triggers cell death in

Arabidopsis thaliana **and** *Nicotiana tabacum***.**

(A) *Pseudomonas syringae* pv tomato DC3000 harboring control plasmids (left) or

expressing NopT (right) were infiltrated into Arabidopsis leaves. Note the cell

death in the right-side panel. (B) *Agrobacterium* strains harboring *NopT*, *NopT*

with an N-terminal 50–amino acid deletion, *NopT*^{C93S}, or *NopT*^{C93S} with an

N-terminal 50–amino acid deletion were infiltrated into *N. tabacum* leaves.

1079

1080

1081 **Supplemental Table 1. Phosphopeptides identified by liquid chromatography–mass**

1082 **spectrometry.** An *in vitro* kinase assay was performed using the CD of NFR1 and NopT.

1083 NopT was separated by SDS-PAGE, and the digested gel slices representing the

1084 phosphorylated NopT were used for analysis. The deduced phosphopeptides are listed.

Sequence	Modifications	SITE	PSMs_NFR1KE+NopT	PSMs_NFR1+NopT
ETLANVELRTK	1xPhospho [T10]	37T	-	1
ETLANVELR	1xPhospho [T2]	29T	-	1
ETLANVELRTKSPSAECPDK	1xCarbamidomethyl [C17]; 1xPhospho [T10]	37T	-	1
TKSPSAECPDK	1xCarbamidomethyl [C8]; 1xPhospho [S5]	41S	-	1
SPSAECPDK	1xCarbamidomethyl [C6]; 1xPhospho [S3]	41S		1
SPSAECPDKMGCCASKPQASDPNNPSTSSPARPSTSLFR	3xCarbamidomethyl [C6; C12; C13]; 1xOxidation [M10]		1	1
SPSAECPDKMGCCASKPQASDPNNPSTSSPARPSTSLFR	3xCarbamidomethyl [C6; C12; C13]; 1xPhospho [S/T]; 1xOxidation [M10]		-	4
MGCCASKPQASDPNNPSTSSPARPSTSLFR	2xCarbamidomethyl [C3; C4]; 1xPhospho [T/S]; 1xOxidation [M1]		-	11
MGCCASKPQASDPNNPSTSSPARPSTSLFR	2xCarbamidomethyl [C3; C4]; 1xPhospho [T/S]		-	9
MGCCASKPQASDPNNPSTSSPARPSTSLFR	2xCarbamidomethyl [C3; C4]; 2xPhospho [T/S]	65S 66T	-	4
MGCCASKPQASDPNNPSTSSPARPSTSLFR	2xCarbamidomethyl [C3; C4]; 1xOxidation [M1]		4	2
MGCCASKPQASDPNNPSTSSPARPSTSLFR	2xCarbamidomethyl [C3; C4]; 2xPhospho [T/S]; 1xOxidation [M1]		-	4
TAELAQANADGICVGLTAEWLR	1xCarbamidomethyl [C13]; 1xPhospho [T17]	97T	-	10
NLNSHPSIR	1xPhospho [S7]	109S	-	2
NLNSHPSIRMEALVPGSQR	1xPhospho [S7]; 1xOxidation [M10]	119S	-	1
MEALVPGSQR	1xPhospho [S8]	119S	-	1
MEALVPGSQR	1xPhospho [S8]; 1xOxidation [M1]	119S	-	1
RQGAGPSEADFAAQNTMLQK	1xPhospho [T/S]; 1xOxidation [M17]		-	3
RQGAGPSEADFAAQNTMLQK	1xPhospho [T16]	156T	-	2
QGAGPSEADFAAQNTMLQK	1xPhospho [T/S]; 1xOxidation [M16]		-	3
QGAGPSEADFAAQNTMLQK	1xPhospho [T15]	156T	-	2
SLANRYSNPNR	1xPhospho [S7]	245S	-	2
QHLTTVTTQK	1xPhospho [T8]	257T	-	1

1085

1086

1087

1088

1089

1090

1091

1092

1093

1094

1095

1096

1097

1098 **Supplemental Table 2. Primers used in this study.**

Used for	Target genes	Names	Sequences (from 5' to 3')
Overexpression in Nicotiana benthamiana	NopT or NopT^{C93S}	p5X-NopT-F	agaacacggggggactctagaatgcacagtcccatcag
		p5X-NopT-R	ccgctgttatcggtacctgtcatcttttgggtggtcac
	NopT$^{\Delta N50}$	p5X-NopTN50-F	agaacacgggggactctagaatgtgctgcgccagcaag
	NFR1	p5X-NFR1-F	agaacacgggggactctagaatgaagctaaaaactggtctact
		p5X-NFR1-R	ccgctgttatcggtacctcttctcacagac
	NFR5	p5X-NFR5-F	agaacacgggggactctagaatggctgtcttctttcttacc
		p5X-NFR5-R	ccgctgttatcggtaccacgtgcagtaat
		p5X-NFR5gfp-R	tcccgggagcggtaccacgtgcagtaatggaagt
	AvrPphB	p5X-AvrPphB-F	agaacacgggggactctagaatgaaaataggtacgcaggcc
		p5X-AvrPphB-R	ccgctgttatcggtacccgaaactctaaactcgtttacgc
templates for protein expression	NopT$_{USDA257}$	p5X-257T-F	agaacacgggggactctagatgctgcgtgatccgaacaat
		p5X-257T-R	ccgctgttatcggtacctgtcacctttgggtggtcaccg
	NopT1$_{USDA110}$	p5X-110T1-F	agaacacgggggactctagaatgtatgatcgaatcggtgg
		p5X-110T1-R	ccgctgttatcggtaccctgcatcctttgcgtcg
	NopT2$_{USDA110}$	p5X-110T2-F	agaacacgggggactctagaatgtataatcgagtcgatggc
		p5X-110T2-R	ccgctgttatcggtaccccgatgaggttccgc
Protein Expression in *E.coli*	NopT or NopT^{C93S}	28a NopT-F	tgggtcgcggatccgaattcatgcacagtcccatcag
		28a FLAG-R	tcgagtgcggccgcaagcttcttgtcatcgtcatccttgtag
		28a Strep-R	tcgagtgcggccgcaagctttttttcaaattgaggatgagaccatcc
	AvrPphB	ET28a-pphb-F	tgggtcgcggatccgaattcaaaataggtacgcaggccac
	NopT$_{USDA257}$	ET28a-257T-F	tgggtcgcggatccgaattcctgcgtgatccgaacaa
	NopT1$_{USDA110}$	ET28a-110T1-F	tgggtcgcggatccgaattctatgatcgaatcggtggctc
	NopT2$_{USDA110}$	ET28a-110T2-F	tgggtcgcggatccgaattctataatcgagtcgatggcgaatac
	NFR1CD	duet NFR1-F	accacagccaggatccgagataccagaagaaggaagaagagaaagc
		duet Sac1nost-R	ggcgcgccgagctcgcgatcggggaaattcgagct
		MAL-NFR1-F	ggaaggatttcagaattcggatccagataccagaagaaggaagaagag
		MAL-NFR1-R	cgacggccagtgccaagctttcatcttctcacagacagtaaatttatgag
	NFR5CD	ACSH-NFR5-F	agaagcgcggatccgaattctatgtatactgccgcagaaagaag
		ACSH-NFR5-R	cgtatgggtagcttccaagcttacgtgcagtaatggaagtc
		MAL-NFR5-F	ggaaggatttcagaattcggatcctatgtatactgccgcagaaagaag
		MAL-NFR5-R	cgacggccagtgccaagctttcaacgtgcagtaatggaagtc
	NFR5KD	SUMO 5KD-F	ccgcgaacagattggtaaggttggggaatcagtgtac
		SUMO HA-R	gctcgaattcggatcctttcacgcatagtcaggaacatcg
	NFR5JM	SUMO 5JM-F	ccgcgaacagattggttatgtatactgccgcagaaagaag
		SUMO 5JM-R	gctcgaattcggatcctttcacgcatagtcaggaacatcg
	sumo-NFR5KD-HA	duet sumo-F	catcaccacagccaggatgggtccctgcaggac
		duet HA-R	cattatgcggccgcaagcttttacgcatagtcaggaacatcg
	sumo-NFR5JM-GFP	duet sumo-F	catcaccacagccaggatgggtccctgcaggac
		duet GFP-R	atgcggccgcaagcttttacttgtacaactcatccatacc

	AtLYK5	ACSH-LYK5-F	gcgcggatccgaattctacaaacgaaggtctaagaag
		ACSH-LYK5-R	ggtagcttccaagcttgttgccaagagagccg
	LjLYS11	ACSH-LYS11-F	agaagcgcggatccgaattcacttgtctgaggaagagaaag
		ACSH-LYS11-R	cgtatgggtagcttccaagcttacgagctgctatcaaagtt
	LYK5JM NFR5KD	LYK5-NFR5-F	cgaaaacagaaaggttggggaatcagtg
		NFR5 LYK5-R	ccccaacctttctgttttcgtcgctgaaatt
	LYS11JM NFR5KD	LYS11 NFR5-F	gagcagtgtaaggttggggaatcag
		NFR5 LYS11R	ccccaaccttacactgctcattgagg
site mutantions construction	NopT	T93S-F	gaatctccgtcggcttgactgc
		T93S-F	gacggagattccatctgcgttc
		97D-F	ggcttggatgcggagtggctgcg
		97D-R	ctccgcatccaagccgacgcag
		97A-F	ggcttggctgcggagtggctgc
		97A-R	ctccgcagccaagccgacg
		109D-F	catccggatatccgaatggaggccctag
		109D-R	ttcggatatccggatgactgttgaggttacgc
		109A-F	catccggcaatccgaatggag
		109A-R	ttcggattgccggatgactgttg
		119D-R	cccggagatcaaaggcacgcctcagc
		119A-F	gcctttgatctccgggtactagggcc
		119A-F	cccggagcgcaaaggcac
		119A-R	ctttgcgctccgggtactagg
		156D-F	gcaaaacgatatgttgcagaaagcaggc
		156D-R	gcaacatatcgttttgcgccgcgaagtcggcc
		156A-F	gcaaaacgctatgttgcagaaagcag
		156A-R	ctgcaacatagcgttttgcgccgcgaagtc
		H205A-F	ggcggcaacaccgttgcgacc
		H205A-R	caacggtgttgccgccgccctcagc
		D220A-F	ctcttcgctcctaatttcggcgaatttac
		D220A-R	gccgaaattaggagcgaagagcgtggtgtttc
		245D-F	cgctacgacaatccaaaccggcag
		245D-R	ttggattgtcgtagcgattggctaggct
		257 mutants	changing the sequence of R primer at nopt$_{257}$when cloned into vectors
	NFR1	K351E-F	gcaattgagaagatggatgtacaagcatc
		K351E-R	catccatcttctcaattgctgttttcttgcc
	NFR5	269-271A-F	ctatgcagctgcacgcagaaagaaggctctg
		269-271A-R	gcgtgcagctgcatagaattcggatccgcgc
		272-274A-F	ctgcgctgcagctaaggctctgaataggactgc
		272-274A-R	cttagctgcagcgcagtatacatagaattcggatcc

275-277A-F	gaaaggcagctgcaaataggactgcttcatcagc
275-277A-R	ctatttgcagctgcctttctgcggcagtatacatag
278-280A-F	ctggctgcagctgcttcatcagctgagactg
278-280A-R	agcagctgcagccagagccttctttctgcg
282-283A-F	gcagcagctgagactgctgataaactac
282-283A-R	gcagtctcagctgctgcagcagtcctattcagag
285-286A-F	ggggctgctgataaactactttctggag
285-286A-R	agtagtttatcagcagccccagctgatgaagcagtc
288-290A-F	gctgcagcactttctggagtttcaggctatgtaag
288-290A-R	aaactccagaaagtgctgcagcagcagtctcagctgatgaag
291-294A-F	gagactgctgataaactagctgctgcagtttcaggctatgtaagcaagcc
291-294A-R	atagcctgaaactgcagcagctagtttatcagcagtctcagctg
295-298A-F	gagttgcagctgctgtaagcaagccaaacgtgtatg
295-298A-R	cttacagcagctgcaactccagaaagtagtttatcagcag
298-300A-F	ctatgcagctgcaccaaacgtgtatgaaatcgacg
298-300A-R	gtttggtgcagctgcatagcctgaaactccagaaagtag
301-303A-F	gcaaggcagctgcttatgaaatcgacgagataatggaagc
301-303A-R	cataagcagctgccttgcttacatagcctgaaactcc
304-306A-F	gtggctgcagctgacgagataatggaagctacgaag
304-306A-R	cgtcagctgcagccacgtttggcttgcttacatagc
307-309A-F	gaaatcgctgcagcaatggaagctacgaaggatttcag
307-309A-R	ccattgctgcagcgatttcatacacgtttggcttgc
310-312A-F	cgagatagcagctgctacgaaggatttcagcg
310-312A-R	gtagcagctgctatctcgtcgatttcatacacgtttg
313-315A-F	gctgcagctgcattcagcgatgagtgcaagg
313-315A-R	ctgaatgcagctgcagcttccattatctcgtcgatttc
316-318A-F	ggatgcagctgctgagtgcaaggttggggaatc
316-318A-R	ctcagcagctgcatccttcgtagcttccattatctc
319-321A-F	cgatgctgcagctgttggggaatcagtgtacaagg
319-321A-R	caacagctgcagcatcgctgaaatccttcgtagc
322-324A-F	gcaaggcagctgcatcagtgtacaaggccaacatag
322-324A-R	gatgcagctgccttgcactcatcgctgaaatc
325-327A-F	gaagcagctgcaaaggccaacatagaaggtcg
325-327A-R	gcctttgcagctgcttccccaaccttgcactc
268-277DE-F	agaagcgcggatccgaattcaataggactgct
268-277DE-R	ctgatgaagcagtcctattgaattcggatccgcgcttc
278-287DE-F	gcagaaagaaggctctggataaactactttctggagtttcagg
278-287DE-R	aaactccagaaagtagtttatccagagccttctttctgcg
298-307DE-F	caggctatgagataatggaagctacgaaggatttc
298-307DE-R	cttccattatctcatagcctgaaactccagaaagtag

308-317DE-F	gaaatcgacgatgagtgcaaggttggggg
308-317DE-R	ttgcactcatcgtcgatttcatacacgtttggc
318-327DE-F	gatttcagcaaggccaacatagaaggtcg
318-327DE-R	gttggccttgctgaaatccttcgtagcttcc
328-337DE-F	tcagtgtacgtaaagaaaatcaaggaaggtggtgcc
328-337DE-R	attttctttacgtacactgattccccaaccttg
288-294A-F	gcagctgccgcagcggctgcctcaggctatgtaagcaagcc
288-294A-R	gcagccgctgcggcagctgcagcagtctcagctgatgaagc
288-298PBS1-F	ggagacaaatctcatgtctcctcaggctatgtaagcaagcc
288-298PBS1-R	ggagacatgagatttgtctccagcagtctcagctgatgaagc
N5-283S-R	tgaagcagtcctattcagagcc
283Y-F	tctgaataggactgcttcatacgctgagactgc
N5-294G	agaaagtagtttatcagcagtctcag
294Q-F	ctgctgataaactactttctcaggtttcaggctatgtaag
N5-303Y-R	cacgtttggcttgcttacatag
304S-F	gcaagccaaacgtgtccgaaatcgacgag
N5-310AT-R	ttccattatctcgtcgatttcatacac
311-2 IY-F	gaaatcgacgagataatggaaatctacaaggatttcagcgatgagtg

1099