Circular RNA HMGCS1 sponges miR-4521 to aggravate type 2 diabetes-induced vascular endothelial dysfunction

Ming Zhang[1,2], Guangyi Du[1], Lianghua Xie[1], Yang Xu[1,3], Wei Chen[1,2,3,*]

[1] Department of Food Science and Nutrition, College of Biosystems Engineering and Food Science, Zhejiang University, Hangzhou, 310058, China.

[2] Sir Run Run Shaw Hospital, School of Medicine, Zhejiang University, Hangzhou, 310016, China.

[3] Ningbo Innovation Center, Zhejiang University, Ningbo 315100, China.

[*]Corresponding author: Wei Chen. Department of Food Science and Nutrition, Zhejiang University, No.866 Yuhangtang Road, Xihu District, Hangzhou 310058, China. Phone: +86 0571 88982191; E-mail: zjuchenwei@zju.edu.cn.

Abstract

Noncoding RNA plays a pivotal role as novel regulators of endothelial cell function. Type 2 diabetes, acknowledged as a primary contributor to cardiovascular diseases, plays a vital role in vascular endothelial cell dysfunction due to induced abnormalities of glucolipid metabolism and oxidative stress. In this study, aberrant expression levels of circHMGCS1 and miR-4521 were observed in diabetes-induced vascular endothelial dysfunction. The overexpression of circHMGCS1 or silencing of miR-4521 expedited the onset of diabetes and aggravated vascular endothelial dysfunction. Mechanistically, circHMGCS1 upregulated arginase 1 by sponging miR-4521, leading to decrease in vascular nitric oxide secretion and inhibition of endothelial nitric oxide synthase activity, and an increase in the expression of adhesion molecules and generation of cellular reactive oxygen species, reduced vasodilation and accelerated the impairment of vascular endothelial function. Collectively, these findings illuminate the physiological role and interacting mechanisms of circHMGCS1 and miR-4521 in diabetes-induced cardiovascular diseases, suggesting that modulating the expression of circHMGCS1 and miR-4521 could serve as a potential strategy to prevent diabetes-associated cardiovascular diseases. Furthermore, our findings provide a novel technical avenue for unraveling ncRNAs regulatory roles of ncRNAs in diabetes and its associated complications.

Impact Statement:

Cardiovascular disease, the top cause of diabetic deaths, progresses via vascular endothelial dysfunction. This study highlights the role of circHMGCS1 and miR-4521 in diabetic vascular function and may be markers for the development of diabetic vascular endothelial dysfunction.

Introduction

Cardiovascular disease (CVD) is the leading cause of morbidity and mortality in patients with type 2 diabetes mellitus (T2DM). The incidence of CVD in T2DM individuals was estimated to be at least two to four times higher than that in non-diabetic individuals (Gregg et al, 2016; Yun & Ko, 2021). T2DM gives rise to abnormal metabolic conditions, including chronic hyperglycemia, dyslipidemia, insulin resistance, inflammation, and oxidative stress (Roden & Shulman, 2019). These processes ultimately lead to vascular endothelial damage, which disrupts the maintenance of vascular homeostasis, thereby promoting the development of CVD (Lundberg & Weitzberg, 2022; Niemann et al, 2017). Vascular endothelial dysfunction (VED) is considered the underlying basis for vascular complications at all stages of T2DM (Eelen et al, 2015; Meigs et al, 2004). However, the mechanism by which the key molecules trigger endothelial dysfunction in diabetes-induced vascular impairment remains unknown.

Endothelial cells form a continuous monolayer lining the arterial, venous, and lymphatic vessels, playing a crucial physiological role in maintaining vascular homeostasis (Bazzoni & Dejana, 2004). Moreover, endothelial cells actively regulate vascular tone, preserve endothelial integrity, and modulate platelet activity, contributing to hemostasis and endothelial repair (Li et al, 2019). The leading cause of T2DM is a prolonged high-sugar, high-fat diet (HFHG), commonly known as a Western diet, which inevitably leads to VED, characterized by reduced content of nitric oxide (NO), uncoupling of endothelial nitric oxide synthase (eNOS), increased formation of reactive oxygen species (ROS) and upregulation of endothelial-related adhesion molecules such as intercellular adhesion molecule 1 (ICAM1), vascular cell adhesion molecule 1 (VCAM1), and endothelin 1 (ET-1). These alterations ultimately lead to elevated blood pressure and impaired vascular arteriolar relaxation, contributing to CVD development (Xu et al, 2021; Yeh et al, 2022; Zheng et al, 2018). Despite this understanding, the identification of pivotal regulatory factors

governing gene expression in diabetes-induced VED remains a formidable

challenge.

circRNAs are generated through back-splicing of precursor messenger RNAs

(pre-mRNAs), exhibiting features such as a covalently closed loop structure, high

stability, conservation, and tissue-specific expression. These molecules primarily

function as miRNA sponges, regulators of transcription, and interactors with proteins

(Kristensen et al, 2019; Liu & Chen, 2022). Conversely, miRNAs are single-stranded

small RNA molecules of 21–23 nucleotides, characterized by high conservation,

temporal and tissue-specific expression, and relatively poor stability.

miRNAs—critical regulators of gene expression—control a wide array of cellular

processes (Gebert & MacRae, 2019). The interplay between miRNAs and circRNAs

forms a complex regulatory network, profoundly influencing diverse biological

pathways and disease states. miRNAs serve as guides, targeting specific mRNAs to

induce their degradation or translational repression, thereby suppressing gene

expression post-transcriptionally (Treiber et al, 2019). By contrast, circRNAs contain

miRNA binding sites, competitively inhibiting miRNA activity and thereby reducing

miRNA binding to other mRNAs. This competitive adsorption effect modulates

intracellular miRNA levels, affecting miRNA-mediated regulation of other target

genes (Cheng et al, 2019; Huang et al, 2019; Shen et al, 2019; Zeng et al, 2021).

The expression levels of circRNA in diabetic patients exhibit abnormalities, and

its potential as a biomarker for early diagnosis or therapeutic target has been

gradually being substantiated (Jiang et al, 2022; Stoll et al, 2020; Tian et al, 2018).

However, research on circRNA in the context of diabetes-induced VED is scarce.

Current research has predominantly focused on the regulatory roles of circRNA in

diabetic cardiomyopathy (Yuan et al, 2023), diabetic nephropathy (Yang & Liu,

2022), gestational diabetes (Du et al, 2022), and diabetic retinopathy (Zhu et al,

2019). Investigations into circRNA regulation of VED have only described the

phenomenon and analyzed the correlation of the results, lacking the precision to infer

causation accurately (Jiang et al, 2020; Liu et al, 2017; Ma et al, 2023). Moreover,

comprehension of VED induced by the complex multifactorial processes of diabetes

pathogenesis remains relatively inadequate. Additionally, the interaction between

circRNA and miRNA in diabetes-induced VED remains a subject of research

controversy (Cheng et al, 2019; Liu et al, 2019; Pan et al, 2018). Therefore, a more

comprehensive research approach is required to elucidate the specific mechanisms

by which circRNA operates in diabetes-related CVDs.

The present study elucidated the mechanisms underlying the interaction

between circRNA and miRNA in regulating diabetes-associated VED. Through the

screening strategy, we successfully identified circHMGCS1—a circRNA originating

from the pre-mRNA of HMGCS1. We demonstrated that the upregulation of

circHMGCS1 and downregulation of miR-4521 significantly promoted

diabetes-induced VED. Moreover, in-depth mechanistic investigations revealed that

circHMGCS1 functioned as a miR-4521 sponge, increasing arginase 1 (ARG1)

expression in vascular endothelial cells and accelerating VED progression.

Furthermore, the expression patterns and interaction mechanisms between

circHMGCS1 and miR-4521 in endothelial cells were identified for the first time in

these findings, which also highlight the circHMGCS1/miR-4521/ARG1 axis as a

novel therapeutic target for interventions designed to protect patients from

diabetes-induced VED.

Results

Global expression analysis of endotheliocyte circRNAs

To explore the potential role of circRNAs in regulating endotheliocyte function, we characterized the transcriptome of circRNA derived from total RNA in human umbilical vein endothelial cells (HUVECs) stimulated by high palmitate and high glucose (PAHG), which allowed us to investigate the potential role of circRNAs in diabetes-induced VED (Figures S1A-S1C). A total of 17,179 known circRNAs were identified, with 66 of them exhibiting differential expression (GEO submission: GSE237597). Our main focus was on circRNAs that shared identical genomic and splicing lengths that exhibited fold changes greater than 2 or less than -2 ($p < 0.01$) in PAHG-treated HUVECs (Figure 1A). Specifically, 48 circRNAs were upregulated, whereas 18 were downregulated (Figure S1D). Furthermore, approximately 66.65% of circRNAs were found to be produced through reverse splicing of exons (Figure S1E).

The observed changes in circRNA levels were confirmed through the real-time quantitative reverse transcription PCR (qRT-PCR) analysis of the five most upregulated circRNAs, suggesting that the results of the RNA-seq data are credible. Among them, circHMGCS1 (hsa_circ_0008621, 899 nt in length, identified as circHMGCS1 in subsequent studies because of its host gene being HMGCS1) exhibited significant upregulation compared with the non-PAHG-treated condition (Figure 1B). The circHMGCS1 sequence is situated on chromosome 5 of the human genome, specifically at 43292575-43297268 with no homology to mouse sequences. Subsequently, divergent primers were designed to amplify circHMGCS1, whereas convergent primers were designed to amplify the corresponding linear mRNA from cDNA and genomic DNA (gDNA) in HUEVCs. amplification of circHMGCS1 was observed in cDNA but not in gDNA (Figure 1C), which confirmed the circularity of circHMGCS1. Sanger sequencing of the amplified product of circHMGCS1 confirmed its sequence alignment with the annotated circHMGCS1 in circBase

(Figure 1D). This observation confirms the origin of circHMGCS1 from exons 2 to 7

of the HMGCS1 gene (Figure 1E).

RNA-Fluorescence *in situ* hybridization (RNA-FISH) confirmed the robust

expression of cytoplasmic circHMGCS1 in HUVECs (Figure 1F). The qRT-PCR

analysis of nuclear and cytoplasmic RNA revealed the predominant cytoplasmic

expression of circHMGCS1 in HUVECs (Figure 1G). qRT-PCR also demonstrated

that circHMGCS1 displayed a stable half-life exceeding 24 h, whereas the linear

transcript HMGCS1 mRNA had a half-life of less than 8 h (Figure 1H). Furthermore,

circHMGCS1 exhibited resistance to digestion by ribonuclease R (an exonuclease

that selectively degrades linear RNA (Xiao & Wilusz, 2019), also known as RNase

R), whereas linear HMGCS1 mRNA was easily degraded upon RNase R treatment

(Figure 1I). These findings indicate that the higher expression of circHMGCS1 in

VED is more than just a byproduct of splicing and suggestive of functionality.

Upregulation of circHMGCS1 promotes diabetes-induced VED

To explore the potential role of circHMGCS1 in regulating endothelial cell function,

we cloned exons 2–7 of HMGCS1 into lentiviral vectors for ectopic overexpression

of circHMGCS1 (Figure S2). We found that circHMGCS1 was successfully

overexpressed in HUVECs without significant changes in HMGCS1 mRNA

expression, and the level of the circular transcripts increased nearly 60-fold than the

level of the linear transcripts (Figure 2A). These results confirm that the

circularization is efficient and that circHMGCS1 has no effect on HMGCS1

expression. Further experiments demonstrated that the overexpression of

circHMGCS1 stimulated the expression of adhesion molecules (VCAM1, ICAM1,

and ET-1) (Figures 2B and 2C), suggesting that circHMGCS1 is involved in VED

promotion. We then used PAHG to stimulate HUVEC-overexpressing circHMGCS1.

NO levels were significantly decreased after PAHG stimulation for 24 h, and

circHMGCS1 overexpression further decreased NO levels (Figure 2D), indicating

that increased circHMGCS1 expression inhibits endothelial diastolic function. eNOS

activity was inhibited by PAHG, and upregulated circHMGCS1 expression further decreased eNOS activity (Figure 2E). Superoxide is one of the major ROS known to promote atherosclerosis (Griendling et al, 2016). Therefore, we used DHE to assess superoxide levels in endothelial cells. The overexpression of circHMGCS1 enhanced PAHG-induced ROS generation (Figure 2F), increasing oxidative stress in endothelial cells. The expression levels of adhesion molecules (VCAM1, ICAM1, and ET-1) were further elevated with circHMGCS1 overexpression (Figure 2G and 2H). These findings suggest that elevated expression of circHMGCS1 may contribute to the development of endothelial cell dysfunction.

miR-4521 protects endothelial cells from PAHG-induced dysfunction

circRNAs may function as ceRNAs to sponge miRNAs, thereby modulating miRNA target depression and imposing an additional level of post-transcriptional regulation in human diseases (Zhong et al, 2018). Accordingly, we conducted miRNA sequencing in PAHG-stimulated HUVECs to identify the potential miRNA targets of circHMGCS1. The deep sequencing analysis revealed 98 significantly differentially expressed miRNAs (Figures 3A and 3B, GEO submission: GSE237295). We then used the miRanda database and ceRNA theory to obtain four candidate miRNAs (miR-4521, miR-3143, miR-98-5p, and miR-181a-2-3p) (Figures 3C and 3D). qRT-PCR confirmed that all four miRNAs were downregulated (Figure 3E). Next, we observed that each of the four miRNA mimics induced a significantly increase in the expression of their corresponding miRNAs in HUVECs through the application of synthetic miRNA mimics (Figure S3A). Relative to the other three miRNAs, miR-4521 mimics significantly inhibited adhesion molecules (ICAM1, VCAM1, and ET-1) expression in HUVECs (Figures 3F and 3G), making it a suitable candidate for further endothelial function studies. We then investigate the effect of miR-4521 on PAHG-stimulated endothelial cell function using synthetic miR-4521 mimics and miR-4521 inhibitor in HUVECs (Figures S3B and S3C). Enhanced miR-4521 levels effectively restored PAHG-induced reductions in NO content and eNOS activity in

HUEVCs, while miR-4521 inhibition led to opposite results (Figures 3H and 3I).

whereas miR-4521 mimics significantly inhibited PAHG-induced expression of ROS

and adhesion molecules (VCAM1, ICAM1, and ET-1) (Figure 3J), whereas the

miR-4521 inhibitor increased the PAHG-induced expression of ROS and adhesion

molecules (VCAM1, ICAM1, and ET-1) (Figures 3K and 3L, and Figure S3D and

S3E). These findings suggest that miR-4521 is involved in PAHG-induced

endothelial cell dysfunction.

circHMGCS1 acts as a miR-4521 sponge to regulate endothelial dysfunction

We then explored the interaction mechanism between circHMGCS1 and miR-4521.

As displayed in Figure S4A, the overexpression of circHMGCS1 reduced the

expression level of miR-4521. Meanwhile, bioinformatics analysis revealed the

predicted binding sites between miR-4521 and circHMGCS1 (Figure S4B). We

constructed a dual-luciferase reporter by inserting the wild-type (WT) or mutant

(MUT) linear sequence of circHMGCS1 into the pmirGLO luciferase vector. The

co-transfection of miR-4521 mimics with the circHMGCS1 luciferase reporter gene

in HEK293T cells, reducing luciferase activity; mutations in the binding sites

between circHMGCS1 and miR-4521 abrogated the effect of miR-4521 mimics on

the activity of the circHMGCS1 luciferase reporter gene mutant (Figure 4A).

Furthermore, AGO2 immunoprecipitation in HUVECs transfected with miR-4521 or

its mutants demonstrated that miR-4521 facilitates the association of AGO2 with

circHMGCS1 in HUVECs (Figure 4B). Meanwhile, we used biotinylated miR-4521

mimics in HUVECs stably overexpressing circHMGCS1 to further validate the

direct binding of miR-4521 and circHMGCS1. The qRT-PCR results revealed that

miR-4521 captured endogenous circHMGCS1, and the negative control with a

disrupting putative binding sequence failed to coprecipitate circHMGCS1 (Figure

4C). Moreover, the RNA-FISH assay showed colocalization of miR-4521 and

circHMGCS1 in the cytoplasm (Figure 4D). Altogether, these findings indicate that

circHMGCS1 functions as a molecular sponge for miR-4521.

Rescue experiments were conducted to investigate whether circHMGCS1

regulates endothelial cell function through miR-4521 sponging. We achieved

miR-4521 inhibition by transfecting HUVECs with a recombinant adeno-associated

virus 9 (AAV9) vector carrying miR-4521 sponges. Notably, miR-4521 sponges

significantly promoted the PAHG-induced reduction in NO content and eNOS

activity. Under this condition, increased circHMGCS1 expression did not interfere

with the changes in NO content and eNOS activity (Figures 4E and 4F). Moreover,

miR-4521 sponges elevated ROS expression in PAHG treated HUVECs, whereas no

further additional effect on ROS expression was observed upon the overexpression

of circHMGCS1 when miR-4521 was sponged (Figure 4G). Further evaluation of

endothelial functional molecules revealed that circHMGCS1 failed to stimulate the

expression of these factors when miR-4521 was sponged (Figures 4H-4J).

Furthermore, the reduction in NO content and eNOS activity induced by

circHMGCS1 overexpression was reversed by miR-4521 mimics (Figures 4K and

4L), and the circHMGCS1-mediated increase in ROS content was inhibited by

miR-4521 mimics (Figure 4M). Furthermore, the expression of adhesion molecules

(ICAM1, VCAM1, and ET-1) was increased in circHMGCS1-transfected HUVECs,

and this effect was significantly ameliorated by miR-4521 mimics (Figures 4N-4P).

These findings demonstrate that circHMGCS1 acts as a sponge for miR-4521,

thereby regulating endothelial function.

circHMGCS1 serves as a miR-4521 sponge to regulate diabetes-induced VED

To further explore the interaction between miR-4521 and circHMGCS1 in

diabetes-induced VED, we administered exogenous miR-4521 agomir (mimics) via

tail vein injection to mice, whereas agomir NC, expressing random sequence, served

as the negative control in HFHG-induced diabetes to examine its effect on diabetes

and associated VED (Figure S5A). Compared with to non-diabetic WT (Control)

mice, DM mice exhibited a significant increase in body weight, upon miR-4521

agomir application, the body weight of diabetic mice significantly decreased

compared with the untreated diabetic mice (Figure 5A), and elevated fasting blood glucose levels in DM mice were likewise substantially inhibited (Figure 5B). By contrast, the combined treatment of circHMGCS1 and miR-4521 agomir did not significantly affect the body weight and blood glucose levels. OGTT and ITT experiments demonstrated that miR-4521 agomir considerably enhanced glucose tolerance and insulin resistance in diabetic mice (Figures 5C, 5D, and Figure S5B and S5C). Blood lipid biochemistry analysis revealed that miR-4521 agomir restored abnormal blood lipid levels in diabetic mice (Figures S5D-S5G). Notably, the abundant presence of circHMGCS1 in the body can negate the inhibitory effect of miR-4521 on diabetes development. These results suggest that miR-4521 can inhibit the occurrence of diabetes, whereas circHMGCS1 specifically dampens the function of miR-4521, weakening its protective effect against diabetes.

To investigate the roles of miR-4521 and circHMGCS1 in mouse vascular relaxation, we conducted an *ex vivo* culture of thoracic aortas from mice subjected to different treatments. A significant impairment in acetylcholine (Ach)-induced vascular relaxation response in the thoracic aorta of diabetic mice was markedly improved by 4521 agomir, emphasizing the critical involvement of miR-4521 in vascular relaxation. Conversely, circHMGCS1 inhibited the protective function of miR-4521 on vascular relaxation (Figure 5E). Across all experimental groups, the smooth muscle exhibited a normal endothelium-dependent relaxation response to the nitric oxide donor sodium nitroprusside (SNP) (Figure S5H), revealing intact smooth muscle function. Meanwhile, diabetic mice displayed a significant increase in SBP compared with normal mice, which was significantly restrained by miR-4521 agomir treatment. Nevertheless, this beneficial effect was nullified in the presence of circHMGCS1 (Figure 5F). Furthermore, miR-4521 agomir inhibited intimal thickening and smooth muscle cell proliferation in diabetic mice, whereas upregulation of circHMGCS1 abrogated this effect (Figure 5G). These findings demonstrate that circHMGCS1 and miR-4521 play specific roles in regulating vascular endothelial function in diabetic mice.

We measured the NO level to further investigate the interaction between

miR-4521 and circHMGCS1 in regulating diabetes-induced VED. Compared with

the control group, the NO level decreased in the thoracic aorta of diabetic mice;

however, miR-4521 agomir treatment increased its content (Figure 5H). Furthermore,

miR-4521 treatment effectively enhanced eNOS activity (Figure 5I), whereas

circHMGCS1 hindered miR-4521 regulatory function on NO content and eNOS

activity. Furthermore, miR-4521 reduced ROS production in the blood vessels

(Figures 5J and 5K) and inhibited the expression of endothelial adhesion molecules

(ICAM1, ET-1, and VCAM1) (Figure 5L). Nonetheless, in the presence of

circHMGCS1, the effect of miR-4521 was abrogated. These results indicated that

circHMGCS1 acts as a miR-4521 sponge, participating in diabetes-induced VED.

ARG1 is a direct target of miR-4521 to accelerate endothelial dysfunction

To investigate whether circHMGCS1-associated miR-4521 regulated VED-related

gene expression by targeting the 3'-untranslated region (UTR), we conducted

predictions of mRNAs containing miR-4521 binding sites using the GenCard,

mirDIP, miRWalk, and Targetscan databases. Through bioinformatics analyses, we

identified binding sites for miR-4521 on KLF6, ARG1, and MAPK1 genes (Figure

6A). Previous studies have demonstrated the involvement of elevated arginase

activity in diabetes-induced VED. Arginase can modulate the production of NO

synthesized by eNOS through competitive utilization of the shared substrate

L-arginine (Jung et al, 2010; Romero et al, 2008). Moreover, miR-4521 possesses

the capability to bind to the 3' untranslated region of ARG1 in both human and

mouse genomes (Figure S6A). The protein expression of ARG1 was increased in the

aorta of diabetic mice and PAHG-treated HUVECs (Figures 6B and Figure S6B).

The luciferase reporter assay was applied to verify the targeting ability through the

pmirGLO vector, which included either the WT or MUT 3'-UTR of ARG1 (Figure

S6C). The overexpression of miR-4521 reduced the luciferase activities of the WT

reporter vector but not the MUT reporter vector (Figure 6C). We next investigated

the function of miR-4521 to regulate the mRNA and protein levels of adhesion

molecules (ET-1, ICAM1, and VCAM1) through ARG1 in HUVECs. The rescue

experiment results demonstrated that miR-4521 mimics reversed the promotional

effect of ARG1 on the mRNA and protein expression of adhesion molecules *in vitro*

(Figures 6D and 6E). miR-4521 mimics recovered the reduced levels of NO and the

compromised activity of eNOS triggered by ARG1 overexpression (Figures S6D and

S6E). Meanwhile, it inhibited the upregulation in ROS levels induced by ARG1

overexpression (Figures S6F and S6G).

To verify the regulatory effect of ARG1 in diabetes-induced VED, we

transfected HUVECs with lentiviral vectors containing cloned ARG1 sequences to

achieve ARG1 overexpression (pLV-ARG1). The NO content assay revealed that

elevated ARG1 expression significantly suppressed the reduction of NO content

induced by PAHG in HUVECs (Figure 6F). Moreover, we measured eNOS

activity—a significant regulator of NO production—and observed its inhibition by

ARG1 overexpression (Figure 6G). Besides, upregulated ARG1 expression increased

the production of ROS in endothelial cells, elevating oxidative stress in HUVECs

(Figure 6H). These findings suggest ARG1 involvement in promoting VED

development, and miR-4521 mediates the participation of ARG1 in diabetes-induced

VED.

To explore the regulatory effect of the interaction between circHMGCS1 and

miR-4521 on ARG1 expression, we transfected HUVECs with either miR-4521

mimics or miR-4521 inhibitors. Subsequently, we evaluated the gene and protein

expression levels of ARG1 using qRT-PCR and western blotting. miR-4521 mimics

repressed ARG1 transcription and protein expression, whereas miR-4521 inhibition

upregulated ARG1 expression (Figures 6I-6K). Additionally, circHMGCS1

overexpression significantly further promoted ARG1 transcription and protein

expression during the PAHG treatment (Figures 6L and 6M), confirming the

regulatory function of circHMGCS1 in VED-related ARG1 expression. We next

studied the function of miR-4521, which acts as a sponge for circHMGCS1 to regulate ARG1 in HUVECs. The mRNA and protein levels of ARG1 were significantly increased when HUVECs were transfected with circHMGCS1 (Figures 6N and 6O). However, miR-4521 overexpression counteracted the effects of circHMGCS1 on ARG1 mRNA and protein expression. Besides, the inhibition of miR-4521 using a miR-4521 sponge in HUVECs further increased ARG1 expression under the PAHG treatment. Notably, elevated circHMGCS1 expression did not affect ARG1 regulation (Figure 6P). Moreover, compared with diabetic mice, circHMGCS1 overexpression dampened the function of miR-4521 agomir on ARG1 (Figure 6Q). These findings indicate that circHMGCS1 serves as a sponge for miR-4521, facilitating ARG1 regulation and VED promotion.

ARG1 is essential for circHMGCS1 and miR-4521 to regulate diabetes-induced VED

To delve further into how circHMGCS1 and miR-4521 regulated endothelial cell function via ARG1, we used AAV9 expressing ARG1 shRNA (ARG1 shRNA) or GFP (NC shRNA, used as the control) to reduce ARG1 levels. ARG1 shRNA mitigated the effects of PAHG on NO content and eNOS activity (Figures 7A and 7B), and inhibited PAHG-induced ROS in endothelial cells (Figure 7C). Additionally, ARG1 shRNA significantly reduced the elevated expression of adhesion molecules (ICAM1, VCAM1, and ET-1) induced by PAHG (Figures 7D and 7E). F Further rescue experiments revealed that miR-4521 and circHMGCS1 exhibited no significant regulatory effect on endothelial function in the presence of ARG1 shRNA. These findings demonstrate that ARG1 plays a crucial regulatory effect on miR-4521 and circHMGCS1 in modulating endothelial cell function.

Considering the role of ARG1 in regulating endothelial cell function through circHMGCS1 and miR-4521, we downregulated ARG1 expression in mice by administering AAV9 ARG1 shRNA through tail vein injection (Figure S7A). As anticipated, AAV9 ARG1 shRNA-infused mice exhibited decreased body weight and

lower fasting blood glucose levels compared with diabetic mice (Figures 7F and 7G).

Moreover, OGTT and ITT demonstrated that reduced ARG1 levels in AAV9 ARG1

shRNA-treated mice preserved glucose tolerance and insulin sensitivity (Figures 7H,

7I, and Figures S7B and S7C). The lowered ARG1 levels also reversed serum lipid

levels (Figure S7D-S7G). However, the overexpression of circHMGCS1 or

miR-4521 had no significant effect on glucose and lipid metabolism in AAV9 ARG1

shRNA-expressing mice. These findings confirm that ARG1 is an indispensable

regulator for circHMGCS1 and miR-4521 in regulating diabetes.

We next investigated whether ARG1 regulates vascular endothelial cell function

in the context of diabetes. Knocking down ARG1 in the thoracic aorta of mice

expressing AAV9 ARG1 shRNA restored diastolic function of the thoracic aorta in

diabetic mice (Figure 7J), emphasizing the vital role of ARG1 in VED. Furthermore,

all experimental groups exhibited a normal endothelium-dependent relaxation

response to the NO donor SNP, suggesting intact smooth muscle function (Figure

S7H). Moreover, AAV9 ARG1 shRNA effectively inhibited diabetes-induced intima

thickening and inhibited smooth muscle cell proliferation (Figure 7K), and which

also counteracted the elevation of SBP induced by diabetes (Figure 7L). However,

the regulatory effects of circHMGCS1 and miR-4521 on vascular dilation were

ineffective in the presence of ARG1 shRNA. We next measured VED-related

functional factors. Compared with diabetic mice, the thoracic aorta of mice

expressing AAV9 ARG1 shRNA exhibited restored NO content and eNOS activity

(Figures 7M and 7N). Furthermore, knocking down ARG1 reduced the high

expression levels of vascular endothelial adhesion molecules (ICAM1, ET-1, and

VCAM1) induced by diabetes (Figure 7O). Immunofluorescence analysis

demonstrated a significant decrease in ROS content in the thoracic aorta of mice

expressing AAV9 ARG1 shRNA compared with diabetic mice (Figures 7P and 7Q).

Notably, circHMGCS1 and miR-4521 lost their regulatory capacity on vascular

endothelial cell function in the presence of ARG1 shRNA. These results indicate that

ARG1 plays a critical role in circHMGCS1 and miR-4521 to modulate

diabetes-induced VED.

Discussion

The field of ncRNA biology has garnered considerable attention and has witnessed intensive research over the last few years. ncRNAs possess the capacity to function as novel regulators in various physiological systems and disease contexts (Esteller, 2011; Matsui & Corey, 2017; Statello et al, 2021). However, most studies have explored the expression patterns and functions of ncRNAs within single disease settings (Arcinas et al, 2019; Shan et al, 2017). The present study contributes to the understanding of ncRNAs and their involvement in diabetes-associated CVD, characterized by disruption of lipid metabolic homeostasis and redox balance in vascular endothelial cells under diabetic conditions. A comprehensive genome-wide analysis revealed 17,179 known circRNA loci transcribed in diabetic endothelial cells. Based on conservation, endothelial specificity, and abundance criteria, five circRNAs were selected for validation, indicating their predominant upregulation in the diabetic endothelial environment. Among them, we discovered a novel circRNA, circHMGCS1, exhibiting high abundance, stability, and cell specificity in diabetic endothelial cells and contributing to endothelial function regulation. Our findings establish that circHMGCS1 promotes VED progression, and its overexpression exacerbates endothelial cell dysfunction. Our study highlights the pivotal role of circHMGCS1 in regulating endothelial cell function during diabetes.

circRNAs play regulatory roles by acting as microRNA sponges or interacting with proteins to regulate alternative splicing, transcription, and epigenetic modifications (Arcinas et al, 2019; Hansen et al, 2013; Memczak et al, 2013). To investigate the mechanism of circHMGCS1 in VED regulation, we analyzed miRNA sequencing results and prediction software to identify miR-4521 as a novel endothelial-expressed miRNA. miR-4521 was found to possess a stable binding site with circHMGCS1, and its expression was significantly decreased in endothelial cells within a diabetic environment. Our functional studies revealed that miR-4521 mimics attenuated VED development in diabetes, whereas its inhibition exacerbated

VED progression. These findings preliminarily demonstrate the regulatory ability of

miR-4521 in VED. The interaction between circHMGCS1 and miR-4521 was further

characterized through luciferase reporter gene assays, RNA pull-down, and RNA

immunoprecipitation. AAV9-mediated circHMGCS1 overexpression in endothelial

cells dampened miR-4521 expression and accelerated diabetes-induced VED.

Conversely, restoring miR-4521 expression effectively alleviated VED progression,

highlighting the critical role of miR-4521 in counteracting circHMGCS1-mediated

promotion of diabetes-induced VED. Moreover, the absence of miR-4521 aggravated

diabetes-induced VED, whereas exogenous circHMGCS1 addition did not affect

VED development. Intriguingly, exogenous miR-4521 significantly restrained

diabetes-induced VED exacerbation, which was attenuated upon the subsequent

addition of exogenous circHMGCS1. These findings underscore the essential

regulatory role of the circHMGCS1 and miR-4521 interaction in maintaining

diabetic vascular homeostasis.

Finally, we investigated the downstream targets crucial for

circHMGCS1-mediated endothelial cell function. Arginase is a dual-nucleus

manganese metalloenzyme that catalyzes the hydrolysis of L-arginine, primarily

producing L-ornithine and urea (Kanyo et al, 1996). ARG1 and ARG2—two

isoforms of arginase—exist in vertebrates, including mammals. Although ARG1 and

ARG2 share the same catalytic reaction, they are encoded by different genes and

exhibit distinct immunological characteristics (Hara et al, 2020; Pudlo et al, 2017; Su

et al, 2021). Elevated ARG1 activity in endothelial cells has been identified as a

significant contributor to VED in T2DM individuals. This dysfunction is attributed

to the excessive formation of ROS caused by the competition between ARG1 and

eNOS for L-arginine. Consequently, oxidases or mitochondrial complexes become

overactivated, resulting in reduced NO levels and subsequent VED (Caldwell et al,

2018). By contrast, inhibiting arginase has been shown to considerably enhance

endothelial function in T2DM patients (Mahdi et al, 2018; Shemyakin et al, 2012;

Zhou et al, 2018). Our findings suggest that T2DM enhances ARG1 activity and

expression in vascular endothelial cells. Conversely, ARG1 inhibition in endothelial

cells delayed the onset of diabetes and preserved NO metabolite levels in the aorta.

Additionally, Ach-induced vasodilation remains intact in the absence of ARG1 in the

aortic ring, as ARG1 inhibition promotes eNOS-dependent relaxation, thereby

preventing VED. Meanwhile, ROS content in the aorta of diabetic mice could be

reduced under ARG1 inhibition, thereby suppressing the occurrence of oxidative

stress. AAV9-mediated overexpression of circHMGCS1 increases ARG1 expression,

which is inhibited by exogenous miR-4521 intervention. In the absence of miR-4521,

circHMGCS1 cannot regulate ARG1 expression, whereas exogenous miR-4521

intervention inhibits the diabetes-induced increase in ARG1 expression *in vivo*, and

this effect is counteracted by circHMGCS1 overexpression. Both *in vitro* and *in vivo*

data suggest that the miR-4521 or circHMGCS1 fails to regulate the effect of

diabetes-induced VED in the absence of ARG1. Therefore, ARG1 may serve as a

promising VED biomarker, and circHMGCS1 and miR-4521 play a key role in

regulating diabetes-induced VED by ARG1.

It would be intriguing to elucidate the mechanisms through which

circHMGCS1 and miR-4521 exert their regulatory roles in endothelial cell function.

The present study elucidated the potential VED-associated mechanisms in diabetes,

with a specific focus on the complicated mutual effects between circHMGCS1 and

miR-4521. The ceRNA network involving circHMGCS1-miR-4521-ARG1 can

become a novel and significant regulatory pathway for preventing and potentially

treating diabetes-induced VED. Our findings suggest that targeted inhibition of

circHMGCS1 or the overexpression of miR-4521 could serve as effective strategies

in mitigating diabetes-induced VED. These insights underscore the promising

function of ncRNAs in developing therapeutic interventions for diabetes-associated

VED.

Experimental Section

Cell Culture

HUVECs were cultured in high glucose Dulbecco's Modified Eagle Medium (DMEM; Hyclone, USA) supplemented with 10% fetal bovine serum (FBS; Gibco, USA), 100 μg/mL of streptomycin, and 100 U/mL of penicillin. The cells were cultured at 37 °C in a humidified incubator with 95% air and 5% CO_2. To simulate the elevated glucose levels observed in diabetes, we cultured HUVECs in a PAHG medium. The medium contained 25 mM glucose and 250 μM saturated free fatty acid (FFA) palmitate (16:C; Sigma, USA) and was incubated for 24 h. For *in vitro* experiments, HUVECs were subcultured in six-well plates. The cells were then transfected with miR-4521 mimics, miR-inhibitor, or miR-NC and incubated for 24 h. Subsequently, the cells were treated with PAHG for another 24 h. Samples were collected to determine the corresponding indices.

Animal design

Male C57BL/6J mice (6–8 weeks old) were obtained from SLAC Laboratory Animal Co., Ltd. (SLAC ANIMAL, China). The mice were housed at a temperature of 22 ± 1 °C with a 12-h light/dark cycle. To induce diet-induced diabetes, we fed wild type littermates either a standard chow (Control) or a high fat-high sucrose (HFHG) diet, where the diet composition consisted of 60% fat, 20% protein, and 20% carbohydrate (H10060, Hfkbio, China). The dietary regimen was maintained for 14 weeks. Throughout this period, body weight and fasting blood glucose (FBG) levels were measured on a weekly basis. An FBG level of ≥11.1 mmol/L was used as the criterion for a successful diabetic model. To study the role of ARG1 in diabetes-induced endothelial dysfunction, we established an HFHG-induced diabetic model (DM), and then pAAV9-ARG1 shRNA (1×10^{12} vg/mL, 100 μL) was injected into the tail vein. Subsequently, we observed changes in relevant indicators. To investigate the regulatory relationship among ARG1, miR-4521, and circHMGCS1 in diabetes-induced endothelial dysfunction, we injected miR-4521 agomir or

pAAV9-circHMGCS1 via the tail vein, following the procedure described in Figure

S5A. The mice were sacrificed after a 12-h fast, and blood and other tissues were

collected and stored at –80 °C until further analysis. All animal experiments were

conducted in strict accordance with the guidelines and laws governing the use and care

of laboratory animals in China (GB/T 35892-2018 and GB/T 35823-2018) and NIH

Guide for the Care and Use of Laboratory Animals. The experimental procedures

involving animals were performed at the Animal Experiment Center of Zhejiang

Chinese Medical University in Hangzhou, China. The animal protocol was conducted

in compliance with the guidelines set forth by the Ethics Committee for Laboratory

Animal Care at Zhejiang Chinese Medical University (Approval No. 2022101345).

Vascular function

After sacrificing the mice, the thoracic aortae and mesenteric arteries were carefully

removed and dissected in an oxygenated ice-cold Krebs solution. The changes in the

isometric tone of the aortic rings and second-order mesenteric arteries were recorded

using a wire myograph (Danish Myo, DK). The arterial segments were stretched to

achieve an optimal baseline tension (3 mN for the aorta) and allowed to equilibrate

for 1 h. Subsequently, the segments were contracted using 60 mmol/L of KCl and

rinsed with Krebs solution. Endothelium-dependent relaxation (EDR) was evaluated

by assessing the concentration-responses to cumulative additions of acetylcholine

(ACh) in noradrenaline (NE, 10^{-3} mol/L) precontracted rings.

Blood pressure measurements

An intelligent non-invasive blood pressure monitor (BP-2010AUL) was used to

measure blood pressure in different mouse groups simultaneously. Prior to

measurement, the mice were placed in a blood pressure room until reaching a

temperature of 25 ± 1 °C. After a 15-min equilibration period, the mice were gently

positioned in a mice jacket set at a constant temperature of 40 °C, with their tail

inserted into a pulse sensor. Throughout the measurements, a quiet environment was

maintained to minimize external disturbances. The instrument automatically

recorded the stabilized sensor signal to obtain the systolic blood pressure (SBP).

Serum lipid assays

After 5 weeks of blood collection from mouse orbits and 12 weeks of drug

administration, blood samples were obtained from mouse arteries. The collected

blood was centrifuged at 1,500 g for 15 min to separate the serum. Commercial kits

and an automatic biochemical analyzer (Biobase BK-600) were used to measure

triglycerides (TG), total cholesterol (TC), high-density lipoprotein cholesterol

(HDL-c), and low-density lipoprotein cholesterol (LDL-c).

Construction of circHMGCS1 and ARG1 plasmids and stable transfection using

lentivirus

circHMGCS1-overexpressing lentiviruses (pLV-circHMGCS1) were obtained from

Hangzhou Ruipu Biotechnology Co., Ltd (Guangzhou, China). Lentiviruses

containing an empty vector and expressing GFP were used as the negative control

(pLV-NC). ARG1-overexpressing lentiviruses (pLV-ARG1 OE) were acquired from

Genomeditech (China). HUVECs were cultured in 6-well plates until reaching 60%

confluence and then infected with lentivirus particles in the presence of 5 μg/ml of

polybrene at a multiplicity of infection of 30. The cells were cultured for at least 3

days before further experiments were conducted. The overexpression efficiency of

circHMGCS1 or ARG1 was evaluated using quantitative real-time PCR or western

blottings.

Cloning and production of circHMGCS1, miR-4521 sponge, and ARG1 shRNA

adeno-associated Virus

The pAAV-circHMGCS1-ZsGreen1 vector was constructed by inserting the

full-length circHMGCS1 into the pAAV-circRNA-ZsGreen1 vector. To achieve

AAV-mediated gene silencing, we obtained a shRNA sequence targeting mouse

ARG1 from Sigma-Aldrich Mission RNAi (TRCN0000101796). The oligo

sequences

5'GATCCGCCTTTGTTGATGTCCCTAATCTCGAGATTAGGGACATCAACAAA

GGCTTTTTA-3'   and

5'-AGCTTAAAAGCCTTTGTTGATGTCCCTAATCTCGAG

ATTAGGGACATCAACAAAGGCG-3' were synthesized, annealed, and ligated to

the pAAV-ZsGreen-shRNA shuttle vector, resulting in the construction of

pAAV-shARG1-ZsGreen1. The miR-4521 sponge cassette was cloned into the

pAAV-ZsGreen1 expression vector to generate pAAV-miR-4521 sponge-ZsGreen1.

Recombinant AAVs (rAAV9s) were produced by transfecting HEK293T cells plated

at a density of 1×10^7 cells/15 cm plate one day prior to transfection using

polyethyleneimine (Linear PEI, MW 25 kDa, Sigma-Aldrich) as the transfection

reagent. For each 15-cm plate, the following reagents were added: 20 μg of pHelper,

10 μg of pAAV-RC9, and 10 μg of pAAV-circHMGCS1-ZsGreen1,

pAAV-shARG1-ZsGreen1, or pAAV-miR-4521 sponge-ZsGreen1. These plasmids

were combined with 500 μL of serum- and antibiotic-free DMEM and 100 μL of PEI

reagent (1 mg/mL, pH 5.0). The DNA-PEI reagent was added drop-wise to the cells

without changing the media on the 15-cm plates. After three days, the cells were

collected and lysed through repeated cycles of freezing and thawing at −80 °C and

37 °C, respectively. The rAAV9s were purified using the ViraTrap™ Adenovirus

Purification Maxiprep Kit (BW-V1260-02, Biomiga). The virus titer was determined

using qPCR and adjusted to 1×10^{12} viral genomes (vg)/mL in PBS containing 4%

sucrose.

Transfection of miRNA mimic or inhibitor

miR-4521 mimics, miRNA negative control (miR-NC), miR-4521 inhibitor, and

miRNA inhibitor negative control (miR-NC inhibitor) were obtained from Tingske

(China). To investigate the functional role of miR-4521 in HUVECs, we performed

transfections using Hieff Trans® *in vitro* siRNA/miRNA Transfection Reagent with

miR-4521 mimics, miR-NC, miR-NC inhibitor, or miR-NC inhibitor at a

concentration of 50 nM, following the manufacturer's protocols. After 48 h of

transfection, proteins and RNA were collected for further analysis. The sequences

used in the experiments are listed as follows: miRNA negative control (miR-NC):

5'-UUCUCCGAACGUGUCACGUTT-3',

5'-ACGUGACACGUUCGGAGAATT-3';

hsa-miR-98-5p-mimics: 5'-UGAGGUAGUAAGUUGUAUUGUU-3',

5'-CAAUACAACUUACUACCUCAUU-3'; Hsa-miR-3143-mimics:

5'-UAACAUUGUAAAGCGCUUCUUU-3',

5'-AGAAGCGCUUUACAAUGUUAUU-3'; Hsa-miR-181a-2-3p-mimics:

5'-ACCACUGACCGUUGACUGUACC-3',

5'-UACAGUCAACGGUCAGUGGUUU-3'; hsa-miR-4521-mimics:

5'-GAGCACAGGACUUCCUUAGCUU-3',

5'-GCUAAGGAAGUCCUGUGCUCAG-3'; hsa-miR-4521-inhibitor:

5'-CUGAGCACAGGACUUCCUUAGC-3'; miRNA inhibitor negative control

(miR-NC inhibitor): 5'-CAGUACUUUUGUGUAGUACAA-3'; miR-4521 agomir:

5'-GAGCACAGGACUUCCUUAGCUU:

5'-GCUAAGGAAGUCCUGUGCUCAG-3'.

NO assays

The supernatant fluid from HUVECs treated with the indicated reagents and a tissue

homogenate of aortic rings were collected. NO concentrations were determined using

the Griess reagent and a total nitric oxide assay kit (Beyotime, China) following the

manufacturer's instructions.

eNOS activity assay

The cellular eNOS activity was assessed as per previously described protocol.

(Leopold et al, 2007). Briefly, Cells were exposed to a PBS buffer (400 μL/well)

containing 1.5 Ci/ml [3H] L-arginine for 15 minutes to initiate eNOS activity

measurement. The reaction was stopped using 1 N ice-cold TCA (500 μL/well).

After freeze-thawing in liquid nitrogen, cells were scraped, treated with

water-saturated ether three times, and neutralized with 1.5 ml of 25 mM pH 8.0

HEPES. Dowex AG50WX8 columns (Tris form) were used for elution with a 1 ml

solution of 40 mM pH 5.5 HEPES supplemented with 2 mM EDTA and 2 mM

EGTA. The eluate was collected for [3H] L-citrulline quantification using liquid

scintillation spectroscopy. Additionally, the assessment of eNOS activity in cell

lysates and tissues was performed through the previously outlined

immunoprecipitation approach (Du et al, 2001). Briefly, the samples were divided

into two tubes: one for protein blotting and the other for eNOS activity measurement.

The eNOS immunocomplexes, linked to protein A-Sepharose beads, were

reconstituted in assay buffer, and eNOS activity was gauged by tracking the

transformation of [3H] L-arginine into [3H] L-citrulline. Equal enzyme quantities in

each incubation were validated via Western blotting.

Detection of ROS for tissue and cell

To assess the level of ROS in tissues, we used a red fluorescent reactive oxygen

probe called DHE (dihydroethidium) (Karim et al, 2022; Su et al, 2018). In brief, 10

μL of the DHE probe was combined with 190 μL of tissue homogenate supernatant,

thoroughly mixed, and subsequently incubated in darkness at 37 °C for 30 min.

Following incubation, the samples were promptly imaged under a fluorescent

microscope.

CircRNA sequencing

Total RNA was extracted from HUVECs, both with and without PAHG stimulation,

utilizing Trizol (Invitrogen, Carlsbad, USA) following the manufacturer's

instructions. The extracted total RNA was then assessed for quality and quantity

using a denaturing agarose gel, Nano Drop, and Agilent 2100 bioanalyzer (Thermo

Fisher Scientific, USA). For the construction of RNA-seq libraries to obtain

sequence information of linear transcripts, 3 μg of total RNA samples underwent

treatment with the epicenter Ribo-ZeroTM Kit (Illumina, San Diego, USA) to

remove rRNA. Linear RNA was subsequently eliminated using RNase R (Epicentre

Technologies, USA). The resulting cleaved RNA fragments were reverse-transcribed to generate cDNA, which served as the template for the synthesis of U-labeled second-stranded DNAs. This synthesis step involved the use of E. coli DNA polymerase I, RNase H, and dUTP, facilitated by the PrimeScript RT regent Kit (TaKaRa, Japan). Purification of the synthesized second-stranded DNAs was accomplished using AMPureXP beads. All subsequent steps were conducted in accordance with the manufacturer's protocols. To assess the quality of the libraries, an Agilent 2100 Bioanalyzer was employed, and the paired-end sequencing was performed using an Illumina Hiseq 4000 (LC Bio, China), adhering to the recommended protocol provided by the vendor.

Treatment of circRNA raw sequencing data

The sequencing data was filtered with Cutadapt by removing reads containing adaptor contamination, poly-N or low quality and undetermined bases, and then sequence quality was verified using FastQC (http://www.bioinformatics.babraham.ac.uk/projects/fastqc/) (Kechin et al, 2017). Bowtie2 and Hisat2 to map reads to the genome of species (Kim et al, 2015; Langmead & Salzberg, 2012). CIRCExplorer2 and CIRI were used to *denovo* assemble the mapped reads to circular RNAs (Kim & Salzberg, 2011; Zhang et al, 2016). Then, back splicing reads were identified in unmapped reads by tophat-fusion. All samples were generated unique circular RNAs. The expression level was calculated according to the junction reads per billion mapped reads (RPB). Differentially expressed circRNAs analysis was performed with log2 (fold change) >1 or log2 (fold change) <-1 and with statistical significance (p value < 0.05) by R package–edgeR (Robinson et al, 2010).

microRNA microarray assay

Total RNA was extracted from HUVECs treated with either PAHG or PBS using TRIzol reagent (Invitrogen, USA). The μParaflo MicroRNA microarray Assay by LC Sciences (LC Sciences, USA) using miRBase version 22.0 was performed. After

3'-extension with a poly(A) tail and ligating an oligonucleotide tag, hybridization was

done overnight on a µParaflo microfluidic chip. Cy3 dye facilitated dye staining after

RNA hybridization. Fluorescence images were acquired using the GenePix 400B laser

scanner (Molecular Devices, USA). Data were analyzed by background subtraction

and signal normalization using a LOWESS filter. miRNA expression with a two-tailed

p-value of <0.05 was considered significant. Intensities below 500 were considered

false positives.

Network analysis of competing endogenous RNA

The miRNA targets of circHMGCS1 were computationally predicted using three

bioinformatic tools: miRanda (http://www. microrna.org/microrna/home.do), whereas

the potential target genes of miR-4521 were envisaged using four distinct databases:

TargetScan (http://www.targetscan.org/), GenCard (https://www.genecards.org/),

miRWalk   (http://mirwalk.umm.uni-heidelberg.de/),   and   mirDIP

(http://ophid.utoronto.ca/mirDIP/).

Isolation of RNA and miRNA for quantitative Real Time-PCR (qRT-PCR)

analysis

Total RNA was isolated using TRIzol reagent (Life Technologies, USA) following the

manufacturer's protocol. The RNA concentration was determined at 260 nm using a

spectrophotometer. Reverse transcludion was conducted using a reverse transcription

kit (TAKARA, Japan) according to the manufacturer's protocol to synthesize cDNA.

Quantitative real-time PCRs (qRT-PCRs) were conducted on the CFX96 Touch

Real-Time PCR Detection System (Bio-Rad, USA) using the standard protocol

provided by the SYBR Green PCR Kit (Yeason, China). For miRNA expression

detection, reverse transcription was executed, and microRNAs were detected using

stem-loop primers procured from Tsingke (China). The relative gene expression was

calculated using the $2^{-\Delta\Delta Ct}$ method. All qRT-PCR assays were repeated thrice. The

primers used for the

qRT-PCR   assays   are   listed   as   follows: Hsa-miR-98-5p:

5'-CGCGCGTGAGGTAGTAAGTTGT-3',

5'-AGTGCAGGGTCCGAGGTATT-3';   Hsa-miR-3143:

5'-GCGATAACATTGTAAAGCGCTT-3', 5'-AGTGCAGGGTCCGAGGTATT-3';

Hsa-miR-181a-2-3p: 5'-GCGACCACTGACCGTTGAC-3',

5'-AGTGCAGGGTCCGAGGTATT-3';   Has-miR-4521:

5'-CGGCTAAGGAAGTCCTGTGC-3', 5'-CGCAGGGTCCGAGGTATTC-3'; U6:

5'-ATTGGAACGATACAGAGAAGATT-3'   ,

5'-GGAACGCTTCACGAATTTG-3';   ET-1:

5'-GCCTGCCTTTTCTCCCCGTTAAA-3',

5'-CAAGCCACAAACAGCAGAGA-3';   ICAM1:

5'-TGACCGTGAATGTGCTCTCC-3',   5'-TCCCTTTTTGGGCCTGTTGT-3';

ARG1:   5'-TTCTCAAAGGGACAGCCACG-3',

5'-CGCTTGCTTTTCCCACAGAC-3';   ICAM1:

5'-ATGCCCAGACATCTGTGTCC-3',   5'-GGGGTCTCTATGCCCAACAA-3';

VCAM1:   5'-AAATCGAGACCACCCCAGAA-3',

5'-AGGAAAAGAGCCTGTGGTGC-3';   β-actin:

5'-CTCACCATGGATGATGATATCGC-3',

5'-CACATAGGAATCCTTCTGACCCA-3';   HMGCS1:

5'-TCGTGGGACACATATGCAAC-3',   5'-TGGGCATGGATCTTTTTGCAG-3';

hsa_circ_0000992:   5'-TCACACGGGAGAGTGTTACC-3',

5'-GTCCATCGAGAAAAGCTGATGC-3';   hsa_circ_0004036:

5'-GGCTCACAAGCAGCCTTTAC-3',   5'-GGTGATACAGGAGCGGGTAG-3';

hsa_circ_0004889:   5'-GTCATATATGGGAAAGAGGGCTT-3',

5'-GGACAAGCCTTCATGGGCTC-3';   hsa_circ_0006719:

5'-CCTCCCTCGGTGTTGCCTTC-3',   5'-TCCAGGCCAGGTAGACAGAAC-3';

hsa_circ_0008621:   5'-CAAACATAGCAACTGAGGGCTTC-3',

5'-TAATGCACTGAGGTAGCACTG-3';   GAPDH:

5'-GAAAGCCTGCCGGTGACTAA-3', 5'-GCATCACCCGGAGGAGAAAT-3'.

circRNA Validation by PCR

DNA and gDNA templates were PCR amplified using BioRad Mastercyclers following the manufacturer's protocol. Subsequently, the PCR products were visualized using a 1% agarose gel stained with GelRed. To validate the PCR results, we purified the products using the EZ-10 Column DNA Purification Kit (Sangon Biotech, China), and direct PCR product Sanger sequencing was performed.

Cell nuclear/cytoplasmic fractionation

RNA extracts from HUVECs were subjected to nuclear/cytoplasmic isolation using the NE-PER Nuclear and Cytoplasmic Extraction Reagent (Thermo Fisher Scientific, USA) following the manufacturer's protocol.

RNase R treatment

For RNase R treatment, approximately 2 μg of total RNAs from HUVECs were incubated with or without 20 U of RNase R (Geneseed, China) at 37°C for 30 min. The resulting RNAs were then purified using an RNA Purification Kit (Qiagen, USA).

Actinomycin D assay

HUVECs were seeded equally in 24-well plates at a density of 5×10^4 cells per well. Subsequently, the cells were exposed to actinomycin D (2 μg/mL; Abcam, USA) for different time intervals: 0, 4, 8, 12, and 24 h. After the respective exposure times, the cells were collected for RNA extraction. The relative RNA levels of circHMGCS1 and HMGCS1 mRNA were analyzed using qRT-PCR and normalized to the values obtained from the control group (0 h).

RNA immunoprecipitation

RIP assay was conducted using the PureBinding®RNA Immunoprecipitation Kit (GENESEED, China) as per the manufacturer's guidelines. HUVECs were lysed in complete RNA immunoprecipitation lysis buffer containing 1% proteinase inhibitor

and 1% RNase inhibitor after being infected with circNC or circHMGCS1. The cell extract was then incubated with magnetic beads conjugated with anti-Argonaute 2 (AGO2) or anti-IgG antibody (Abcam, USA) overnight at 4 °C. The beads were subsequently washed, and the proteins were removed using columns. Eventually, the isolated RNA was extracted using TRIzol Reagent, and the purified RNA was utilized for subsequent qRT-PCR analysis.

Pull-down Assay with biotinylated miR-4521

A pull-down assay was conducted using the PureBinding®RNA-Protein pull-down Kit (GENESEED, P0201, China) following the manufacturer's protocol. The pull-down assay with biotinylated miR-4521 was performed as previously described (Wang et al, 2020). Briefly, HUVECs were transfected with biotinylated miR-4521 mimics or mutants (Tingke, 50 nmol/L) using the Hieff Trans® *in vitro* siRNA/miRNA Transfection Reagent (Yeason, China). Following a 48-h transfection period, the cells were collected, PBS-washed, and incubated for 10 min in a capture buffer on ice. A 10% fraction of the cell lysates was retained as input. The remaining lysates were subsequently exposed to streptavidin magnetic beads for 30 min at 4 °C. After treatment with wash buffer and the RNeasy Mini Kit (Qiagen), the captured RNAs were extracted for subsequent qRT-PCR analysis.

The sequences are listed as follows: Biotin-miR-4521-MUT: 5'-GCAUUCCUUCAGGAGUGCUCAG-3', Biotin-miR-4521-WT: 5'-GCUAAGGAAGUCCUGUGCUCAG-3'.

Western blot analysis

Western blot was performed following previously described procedures (Xu et al, 2022). Protein extraction from cells or tissues was carried out using protein lysis buffer containing protease and phosphatase inhibitor cocktail (Beyotime, China). The concentrations of the protein lysates were determined using the Beyotime BCA Protein Assay Kit (Beyotime, China) according to the manufacturer's instructions.

Equal quantities of proteins (20 μg/lane) were loaded onto 8% or 10% SDS-PAGE

gels and subsequently transferred to PVDF membranes (Millipore, USA). The

membranes were then incubated with primary antibodies overnight at 4°C. The

primary antibodies used were as follows: anti-ARG1 (diluted 1:1,000; Abcam,

ab133543), anti-ARG2 (diluted 1:1,000; Abcam, ab264066), anti-ET-1 (diluted

1:1,000; Abcam, ab2786), anti-VCAM1 (diluted 1:1,000; Abcam, ab134047),

anti-ICAM1 (diluted 1:1,000; Abcam, ab222736), anti-β-actin (diluted 1:2,000;

Abcam, ab5694), and anti-GAPDH (diluted 1:2,000; Abcam, ab8245). After

incubation with the corresponding secondary antibodies for 2 hours at room

temperature, the membranes were washed. Antibody binding was detected using an

ECL detection reagent (Millipore, USA). Digital images were captured using a Gel

DocTM XR+ System with ImageLab software (BioRad, USA). The quantification of

the labeled bands was performed using ImageJ 1.55.

RNA Fluorescent *In Situ* Hybridization (RNA-FISH)

The RNA-FISH assay was conducted in HUVECs. Cy3-labeled circHMGCS1 probe

and FAM-labeled miR-4521 probe were custom-designed and synthesized by

RiboBio (China). The signals emanating from the probes were detected using the

Ribo^{TM} FISH Kit (RiboBio) in accordance with the manufacturer's protocol. Briefly,

frozen sections were fixed with 4% paraformaldehyde for 10 minutes, followed by

PBS washing. The fixed sections were treated with PBS containing 0.5% triton

X-100 at 4°C for 5 min, followed by incubation in pre-dehydration buffer at 37°C

for 30 min. Subsequently, probes targeting circMAP3K5 and miR-4521 were applied

to the sections. Hybridization was performed in a humid chamber at 37°C overnight.

Post-hybridization washing was initially conducted with 4× saline sodium citrate

containing 0.1% tween-20 at 42°C, followed by further washing with 2× saline

sodium citrate at 42°C to eliminate non-specific and repetitive RNA hybridization.

The slides were counterstained with DAPI (beyotime, China) and examined using an

Olympus FV1200 (Olympus, Japan) confocal microscopy system. The mean

fluorescent intensity (MFI) was determined using Image J software and subjected to

statistical analysis. The probe

sequences   are   detailed   as   follows: Cy3-circHMGCS1:

5'-UCCCACGAAGCCCUCAGUUGCUAUG-3',   FAM-miR-4521:

5'-UUUGACUCGUGUCCUGAAGGAAUCG-3'.

Luciferase reporter assay

The complete sequence of circHMGCS1 and a mutant circHMGCS1 sequence

lacking the miRNA binding site were chemically synthesized. These sequences were

inserted into the Nhel and SaII sites of the pmirGLO vector to generate expression

vectors denoted as circHMGCS1-WT and circHMGCS1-MUT. Transfection was

carried out using lipofectamine 2000 (Invitrogen, USA) with 50 nM miR-4521

mimics, miR-NC, miR-4521 inhibitor, and miR-NC inhibitor. HEK293T cells were

seeded in 96-well plates and cultured until reaching 50%-70% confluence prior to

transfection. After 48-h of transfection, cells were harvested, and luciferase activity

was quantified utilizing the Dual Luciferase Reporter Gene Assay Kit (Yeason,

China). The pmirGLO vector expressing Renilla luciferase was utilized as an

internal control for transfection, while the empty pmirGLO vector served as the

negative control. Firefly luciferase activity of the pmirGLO vector was normalized

with Renilla luciferase activity for comparison. Luciferase reporter assay was also

used to investigate the regulation of miR-4521 on the expression of its target genes.

For ARG1 and miR-4521, either wild-type or mutant ARG1 3'UTR fragments (433

bp) were inserted into NheI/SalI restriction sites of pmirGLO. 1 μg plasmids of

ARG1 3'UTR-WT and ARG1 3'UTR-MUT, 50 nM miR-4521 mimics, miR-NC,

miR-4521 inhibitor and miR-NC inhibitor were transfected. Luciferase activity was

evaluated 48 hours post-transfection using the Dual Luciferase Reporter Gene Assay

Kit (Yeason).

Histological and morphometric analysis

Aortic tissues were prepared by fixing them in 4% paraformaldehyde in PBS and

subsequently embedding them in paraffin. The resulting paraffin-embedded tissues

were then sectioned into 4-μm-thick slices. Hematoxylin-eosin (H&E) staining was

performed on these sections to visualize the vascular tissue structures and assess the

degree of thickness in the thoracic aorta. The average thickness of five randomly

selected regions of the thoracic aorta was measured using OlyVIA software

(Olympus, Japan).

Statistical analysis

Statistical analyses were conducted using Prism software (GraphPad 8.0). Data are

presented as mean ± standard deviation (SD) unless otherwise stated. Prior to

statistical analysis, the normality of data distribution was evaluated using the

Shapiro-Wilk test. For comparisons between the two groups, an unpaired two-tailed

Student's t-test was used. When comparing more than the two groups, either a

one-way or two-way analysis of variance (ANOVA) was conducted, followed by the

Bonferroni multiple comparison post hoc test. Group differences were considered

statistically significant at a p-value of <0.05. The level of significance was indicated

by asterisks, with *, **, and *** denoting p-values lower than 0.05, 0.01, and 0.001,

respectively.

Author contributions

Conceptualization: M.Z. and W.C. Methodology: M.Z., G.D., Y.X., W.C., and L.X. Investigation: M.Z., L.X., Y.X., and G.D. Formal analysis: M.Z. Validation: M.Z. Data curation: M.Z. and G.D. Writing (original draft): W.C. and M.Z. Writing (review and editing): W.C. and M.Z. Funding acquisition: W.C. Resources: W.C. Supervision: W.C. All authors contributed, read and approved the final version.

Acknowledgments

The authors would like to thank core facilities of medicine in Zhejiang university for technical support.

Funding

This work was supported by the National Natural Science Foundation of China (32172192) and Zhejiang Provincial Key R&D Program of China (2021C02018).

Conflict of interest

The authors have declared that no conflict of interest exists.

Data availability

All data needed to evaluate the conclusions in the paper are present in the paper and/or the Supplementary Materials. Unprocessed original image data have been deposited to Mendely Data and are available at: https://doi.org/10.17632/dxzvhnj7pm.1. Profiling by circRNA and miRNA Array in our work has been deposited in the GEO database as GSE237295 (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE237295) and GSE237597 (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE237597). Additional data related to this paper may be requested from the authors.

References

Arcinas, C., W. Tan, W. Fang, T.P. Desai, D.C.S. Teh, U. Degirmenci, D. Xu, R. Foo, and L. Sun. 2019. Adipose circular RNAs exhibit dynamic regulation in obesity and functional role in adipogenesis. *Nat Metab.* 1: 688-703.

Arcinas, C., W. Tan, W. Fang, T.P. Desai, D.C.S. Teh, U. Degirmenci, D. Xu, R. Foo, and L. Sun. 2019. Adipose circular RNAs exhibit dynamic regulation in obesity and functional role in adipogenesis. *Nature Metabolism.* 1: 688-703.

Bazzoni, G., and E. Dejana. 2004. Endothelial cell-to-cell junctions: molecular organization and role in vascular homeostasis. *Physiol Rev.* 84: 869-901.

Caldwell, R.W., P.C. Rodriguez, H.A. Toque, S.P. Narayanan, and R.B. Caldwell. 2018. Arginase: A Multifaceted Enzyme Important in Health and Disease. *Physiol Rev.* 98: 641-665.

Cheng, J., Q. Liu, N. Hu, F. Zheng, X. Zhang, Y. Ni, and J. Liu. 2019. Downregulation of hsa_circ_0068087 ameliorates TLR4/NF-κB/NLRP3 inflammasome-mediated inflammation and endothelial cell dysfunction in high glucose conditioned by sponging miR-197. *Gene.* 709: 1-7.

Cheng, Z., C. Yu, S. Cui, H. Wang, H. Jin, C. Wang, B. Li, M. Qin, C. Yang, J. He, Q. Zuo, S. Wang, J. Liu, W. Ye, Y. Lv, F. Zhao, M. Yao, L. Jiang, and W. Qin. 2019. circTP63 functions as a ceRNA to promote lung squamous cell carcinoma progression by upregulating FOXM1. *Nat Commun.* 10: 3200-3213.

Du, R., N. Wu, Y. Bai, L. Tang, and L. Li. 2022. circMAP3K4 regulates insulin resistance in trophoblast cells during gestational diabetes mellitus by modulating the miR-6795-5p/PTPN1 axis. *J Transl Med.* 20: 180-196.

Du, X.L., D. Edelstein, S. Dimmeler, Q. Ju, C. Sui, and M. Brownlee. 2001. Hyperglycemia inhibits endothelial nitric oxide synthase activity by

posttranslational modification at the Akt site. *J Clin Invest.* 108: 1341-1348.

Eelen, G., P. De Zeeuw, M. Simons, and P. Carmeliet. 2015. Endothelial cell
metabolism in normal and diseased vasculature. *Circulation research.* 116:
1231-1244.

Esteller, M. 2011. Non-coding RNAs in human disease. *Nat Rev Genet.* 12: 861-874.

Gebert, L.F.R., and I.J. Macrae. 2019. Regulation of microRNA function in animals.
Nat Rev Mol Cell Biol. 20: 21-37.

Gregg, E.W., N. Sattar, and M.K. Ali. 2016. The changing face of diabetes
complications. *Lancet Diabetes Endocrinol.* 4: 537-547.

Griendling, K.K., R.M. Touyz, J.L. Zweier, S. Dikalov, W. Chilian, Y.R. Chen, D.G.
Harrison, and A. Bhatnagar. 2016. Measurement of Reactive Oxygen Species,
Reactive Nitrogen Species, and Redox-Dependent Signaling in the
Cardiovascular System: A Scientific Statement From the American Heart
Association. *Circ Res.* 119: 39-75.

Hansen, T.B., T.I. Jensen, B.H. Clausen, J.B. Bramsen, B. Finsen, C.K. Damgaard,
and J. Kjems. 2013. Natural RNA circles function as efficient microRNA
sponges. *Nature.* 495: 384-388.

Hara, M., K. Torisu, K. Tomita, Y. Kawai, K. Tsuruya, T. Nakano, and T. Kitazono.
2020. Arginase 2 is a mediator of ischemia-reperfusion injury in the kidney
through regulation of nitrosative stress. *Kidney Int.* 98: 673-685.

Huang, S., X. Li, H. Zheng, X. Si, B. Li, G. Wei, C. Li, Y. Chen, Y. Chen, W. Liao, Y.
Liao, and J. Bin. 2019. Loss of Super-Enhancer-Regulated circRNA Nfix
Induces Cardiac Regeneration After Myocardial Infarction in Adult Mice.
Circulation. 139: 2857-2876.

Jiang, B., J. Zhang, X. Sun, C. Yang, G. Cheng, M. Xu, S. Li, and L. Wang. 2022.
Circulating exosomal hsa_circRNA_0039480 is highly expressed in

gestational diabetes mellitus and may be served as a biomarker for early
diagnosis of GDM. *J Transl Med.* 20: 5-18.

Jiang, Q., C. Liu, C.P. Li, S.S. Xu, M.D. Yao, H.M. Ge, Y.N. Sun, X.M. Li, S.J.
Zhang, K. Shan, B.H. Liu, J. Yao, C. Zhao, and B. Yan. 2020. Circular
RNA-ZNF532 regulates diabetes-induced retinal pericyte degeneration and
vascular dysfunction. *J Clin Invest.* 130: 3833-3847.

Jung, C., A.T. Gonon, P.O. Sjöquist, J.O. Lundberg, and J. Pernow. 2010. Arginase
inhibition mediates cardioprotection during ischaemia-reperfusion.
Cardiovasc Res. 85: 147-154.

Kanyo, Z.F., L.R. Scolnick, D.E. Ash, and D.W. Christianson. 1996. Structure of a
unique binuclear manganese cluster in arginase. *Nature.* 383: 554-557.

Karim, N., M.R.I. Shishir, Y. Li, O.Y. Zineb, J. Mo, J. Tangpong, and W. Chen. 2022.
Pelargonidin-3-O-Glucoside Encapsulated Pectin-Chitosan-Nanoliposomes
Recovers Palmitic Acid-Induced Hepatocytes Injury. *Antioxidants (Basel).* 11:
1-17.

Kechin, A., U. Boyarskikh, A. Kel, and M. Filipenko. 2017. cutPrimers: A New Tool
for Accurate Cutting of Primers from Reads of Targeted Next Generation
Sequencing. *J Comput Biol.* 24: 1138-1143.

Kim, D., B. Langmead, and S.L. Salzberg. 2015. HISAT: a fast spliced aligner with
low memory requirements. *Nat Methods.* 12: 357-360.

Kim, D., and S.L. Salzberg. 2011. TopHat-Fusion: an algorithm for discovery of
novel fusion transcripts. *Genome Biol.* 12: 1-15.

Kristensen, L.S., M.S. Andersen, L.V.W. Stagsted, K.K. Ebbesen, T.B. Hansen, and J.
Kjems. 2019. The biogenesis, biology and characterization of circular RNAs.
Nat Rev Genet. 20: 675-691.

Langmead, B., and S.L. Salzberg. 2012. Fast gapped-read alignment with Bowtie 2.

957 *Nat Methods.* 9: 357-359.

958 Leopold, J.A., A. Dam, B.A. Maron, A.W. Scribner, R. Liao, D.E. Handy, R.C.
959   Stanton, B. Pitt, and J. Loscalzo. 2007. Aldosterone impairs vascular
960   reactivity by decreasing glucose-6-phosphate dehydrogenase activity. *Nat*
961   *Med.* 13: 189-197.

962 Li, X., X. Sun, and P. Carmeliet. 2019. Hallmarks of Endothelial Cell Metabolism in
963   Health and Disease. *Cell Metab.* 30: 414-433.

964 Liu, C., H.M. Ge, B.H. Liu, R. Dong, K. Shan, X. Chen, M.D. Yao, X.M. Li, J. Yao,
965   R.M. Zhou, S.J. Zhang, Q. Jiang, C. Zhao, and B. Yan. 2019. Targeting
966   pericyte-endothelial cell crosstalk by circular RNA-cPWWP2A inhibition
967   aggravates diabetes-induced microvascular dysfunction. *Proc Natl Acad Sci*
968   *U S A.* 116: 7455-7464.

969 Liu, C., M.D. Yao, C.P. Li, K. Shan, H. Yang, J.J. Wang, B. Liu, X.M. Li, J. Yao, Q.
970   Jiang, and B. Yan. 2017. Silencing Of Circular RNA-ZNF609 Ameliorates
971   Vascular Endothelial Dysfunction. *Theranostics.* 7: 2863-2877.

972 Liu, C.X., and L.L. Chen. 2022. Circular RNAs: Characterization, cellular roles, and
973   applications. *Cell.* 185: 2016-2034.

974 Lundberg, J.O., and E. Weitzberg. 2022. Nitric oxide signaling in health and disease.
975   *Cell.* 185: 2853-2878.

976 Ma, C., X. Wang, L. Zhang, X. Zhu, J. Bai, S. He, J. Mei, J. Jiang, X. Guan, X.
977   Zheng, L. Qu, and D. Zhu. 2023. Super Enhancer-Associated Circular
978   RNA-CircKrt4 Regulates Hypoxic Pulmonary Artery Endothelial Cell
979   Dysfunction in Mice. *Arterioscler Thromb Vasc Biol.* 43: 1179-1198.

980 Mahdi, A., O. Kövamees, A. Checa, C.E. Wheelock, M. Von Heijne, M. Alvarsson,
981   and J. Pernow. 2018. Arginase inhibition improves endothelial function in
982   patients with type 2 diabetes mellitus despite intensive glucose-lowering

therapy. *J Intern Med.* 284: 388-398.

Matsui, M., and D.R. Corey. 2017. Non-coding RNAs as drug targets. *Nat Rev Drug*
Discov. 16: 167-179.

Meigs, J.B., F.B. Hu, N. Rifai, and J.E. Manson. 2004. Biomarkers of endothelial
dysfunction and risk of type 2 diabetes mellitus. *Jama.* 291: 1978-1986.

Memczak, S., M. Jens, A. Elefsinioti, F. Torti, J. Krueger, A. Rybak, L. Maier, S.D.
Mackowiak, L.H. Gregersen, M. Munschauer, A. Loewer, U. Ziebold, M.
Landthaler, C. Kocks, F. Le Noble, and N. Rajewsky. 2013. Circular RNAs
are a large class of animal RNAs with regulatory potency. *Nature.* 495:
333-338.

Niemann, B., S. Rohrbach, M.R. Miller, D.E. Newby, V. Fuster, and J.C. Kovacic.
2017. Oxidative Stress and Cardiovascular Risk: Obesity, Diabetes, Smoking,
and Pollution: Part 3 of a 3-Part Series. *J Am Coll Cardiol.* 70: 230-251.

Pan, L., W. Lian, X. Zhang, S. Han, C. Cao, X. Li, and M. Li. 2018. Human circular
RNA-0054633 regulates high glucose-induced vascular endothelial cell
dysfunction through the microRNA-218/roundabout 1 and
microRNA-218/heme oxygenase-1 axes. *Int J Mol Med.* 42: 597-606.

Pudlo, M., C. Demougeot, and C. Girard-Thernier. 2017. Arginase Inhibitors: A
Rational Approach Over One Century. *Med Res Rev.* 37: 475-513.

Robinson, M.D., D.J. Mccarthy, and G.K. Smyth. 2010. edgeR: a Bioconductor
package for differential expression analysis of digital gene expression data.
Bioinformatics. 26: 139-179.

Roden, M., and G.I. Shulman. 2019. The integrative biology of type 2 diabetes.
Nature. 576: 51-60.

Romero, M.J., D.H. Platt, H.E. Tawfik, M. Labazi, A.B. El-Remessy, M. Bartoli, R.B.
Caldwell, and R.W. Caldwell. 2008. Diabetes-induced coronary vascular

dysfunction involves increased arginase activity. *Circ Res.* 102: 95-102.

Shan, K., C. Liu, B.H. Liu, X. Chen, R. Dong, X. Liu, Y.Y. Zhang, B. Liu, S.J.
Zhang, J.J. Wang, S.H. Zhang, J.H. Wu, C. Zhao, and B. Yan. 2017. Circular
Noncoding RNA HIPK3 Mediates Retinal Vascular Dysfunction in Diabetes
Mellitus. *Circulation.* 136: 1629-1642.

Shemyakin, A., O. Kövamees, A. Rafnsson, F. Böhm, P. Svenarud, M. Settergren, C.
Jung, and J. Pernow. 2012. Arginase inhibition improves endothelial function
in patients with coronary artery disease and type 2 diabetes mellitus.
Circulation. 126: 2943-2950.

Shen, S., Y. Wu, J. Chen, Z. Xie, K. Huang, G. Wang, Y. Yang, W. Ni, Z. Chen, P. Shi,
Y. Ma, and S. Fan. 2019. CircSERPINE2 protects against osteoarthritis by
targeting miR-1271 and ETS-related gene. *Ann Rheum Dis.* 78: 826-836.

Statello, L., C.J. Guo, L.L. Chen, and M. Huarte. 2021. Gene regulation by long
non-coding RNAs and its biological functions. *Nat Rev Mol Cell Biol.* 22:
96-118.

Stoll, L., A. Rodríguez-Trejo, C. Guay, F. Brozzi, M.B. Bayazit, S. Gattesco, V.
Menoud, J. Sobel, A.C. Marques, M.T. Venø, J.L.S. Esguerra, M. Barghouth,
M. Suleiman, L. Marselli, J. Kjems, L. Eliasson, E. Renström, K. Bouzakri,
M. Pinget, P. Marchetti, and R. Regazzi. 2020. A circular RNA generated
from an intron of the insulin gene controls insulin secretion. *Nat Commun.* 11:
5611-5624.

Su, H., Y. Li, D. Hu, L. Xie, H. Ke, X. Zheng, and W. Chen. 2018. Procyanidin B2
ameliorates free fatty acids-induced hepatic steatosis through regulating
TFEB-mediated lysosomal pathway and redox state. *Free Radic Biol Med.*
126: 269-286.

Su, X., Y. Xu, G.C. Fox, J. Xiang, K.A. Kwakwa, J.L. Davis, J.I. Belle, W.C. Lee,
W.H. Wong, F. Fontana, L.F. Hernandez-Aya, T. Kobayashi, H.M. Tomasson,

J. Su, S.J. Bakewell, S.A. Stewart, C. Egbulefu, P. Karmakar, M.A. Meyer, D.J. Veis, D.G. Denardo, G.M. Lanza, S. Achilefu, and K.N. Weilbaecher. 2021. Breast cancer-derived GM-CSF regulates arginase 1 in myeloid cells to promote an immunosuppressive microenvironment. *J Clin Invest.* 131: 1-17.

Tian, Y., J. Xu, X. Du, and X. Fu. 2018. The interplay between noncoding RNAs and insulin in diabetes. *Cancer Lett.* 419: 53-63.

Treiber, T., N. Treiber, and G. Meister. 2019. Regulation of microRNA biogenesis and its crosstalk with other cellular pathways. *Nat Rev Mol Cell Biol.* 20: 5-20.

Wang, Y., D. Han, T. Zhou, J. Zhang, C. Liu, F. Cao, and N. Dong. 2020. Melatonin ameliorates aortic valve calcification via the regulation of circular RNA CircRIC3/miR-204-5p/DPP4 signaling in valvular interstitial cells. *J Pineal Res.* 69: e12666.

Xiao, M.S., and J.E. Wilusz. 2019. An improved method for circular RNA purification using RNase R that efficiently removes linear RNAs containing G-quadruplexes or structured 3' ends. *Nucleic Acids Res.* 47: 8755-8769.

Xu, S., I. Ilyas, P.J. Little, H. Li, D. Kamato, X. Zheng, S. Luo, Z. Li, P. Liu, J. Han, I.C. Harding, E.E. Ebong, S.J. Cameron, A.G. Stewart, and J. Weng. 2021. Endothelial Dysfunction in Atherosclerotic Cardiovascular Diseases and Beyond: From Mechanism to Pharmacotherapies. *Pharmacol Rev.* 73: 924-967.

Xu, Y., Y. Li, J. Li, and W. Chen. 2022. Ethyl carbamate triggers ferroptosis in liver through inhibiting GSH synthesis and suppressing Nrf2 activation. *Redox Biol.* 53: 102349-102352.

Yang, J., and Z. Liu. 2022. Mechanistic Pathogenesis of Endothelial Dysfunction in Diabetic Nephropathy and Retinopathy. *Front Endocrinol (Lausanne).* 13: 816400-816419.

Yeh, C.F., S.H. Cheng, Y.S. Lin, T.P. Shentu, R.T. Huang, J. Zhu, Y.T. Chen, S.
Kumar, M.S. Lin, H.L. Kao, P.H. Huang, E. Roselló-Sastre, F. Garcia, H. Jo,
Y. Fang, and K.C. Yang. 2022. Targeting mechanosensitive endothelial
TXNDC5 to stabilize eNOS and reduce atherosclerosis in vivo. *Sci Adv.* 8:
eabl8096.

Yuan, Q., Y. Sun, F. Yang, D. Yan, M. Shen, Z. Jin, L. Zhan, G. Liu, L. Yang, Q.
Zhou, Z. Yu, X. Zhou, Y. Yu, Y. Xu, Q. Wu, J. Luo, X. Hu, and C. Zhang.
2023. CircRNA DICAR as a novel endogenous regulator for diabetic
cardiomyopathy and diabetic pyroptosis of cardiomyocytes. *Signal Transduct*
Target Ther. 8: 99-111.

Yun, J.S., and S.H. Ko. 2021. Current trends in epidemiology of cardiovascular
disease and cardiovascular risk management in type 2 diabetes. *Metabolism.*
123: 154838-154848.

Zeng, Z., L. Xia, S. Fan, J. Zheng, J. Qin, X. Fan, Y. Liu, J. Tao, Y. Liu, K. Li, Z.
Ling, Y. Bu, K.A. Martin, J. Hwa, R. Liu, and W.H. Tang. 2021. Circular
RNA CircMAP3K5 Acts as a MicroRNA-22-3p Sponge to Promote
Resolution of Intimal Hyperplasia Via TET2-Mediated Smooth Muscle Cell
Differentiation. *Circulation.* 143: 354-371.

Zhang, X.O., R. Dong, Y. Zhang, J.L. Zhang, Z. Luo, J. Zhang, L.L. Chen, and L.
Yang. 2016. Diverse alternative back-splicing and alternative splicing
landscape of circular RNAs. *Genome Res.* 26: 1277-1287.

Zheng, Y., S.H. Ley, and F.B. Hu. 2018. Global aetiology and epidemiology of type 2
diabetes mellitus and its complications. *Nat Rev Endocrinol.* 14: 88-98.

Zhong, Y., Y. Du, X. Yang, Y. Mo, C. Fan, F. Xiong, D. Ren, X. Ye, C. Li, Y. Wang, F.
Wei, C. Guo, X. Wu, X. Li, Y. Li, G. Li, Z. Zeng, and W. Xiong. 2018.
Circular RNAs function as ceRNAs to regulate and control human cancer
progression. *Mol Cancer.* 17: 79-90.

Zhou, Z., A. Mahdi, Y. Tratsiakovich, S. Zahorán, O. Kövamees, F. Nordin, A.E.

Uribe Gonzalez, M. Alvarsson, C.G. Östenson, D.C. Andersson, U. Hedin, E.

Hermesz, J.O. Lundberg, J. Yang, and J. Pernow. 2018. Erythrocytes From

Patients With Type 2 Diabetes Induce Endothelial Dysfunction Via Arginase I.

J Am Coll Cardiol. 72: 769-780.

Zhu, K., X. Hu, H. Chen, F. Li, N. Yin, A.L. Liu, K. Shan, Y.W. Qin, X. Huang, Q.

Chang, G.Z. Xu, and Z. Wang. 2019. Downregulation of circRNA DMNT3B

contributes to diabetic retinal vascular dysfunction through targeting

miR-20b-5p and BAMBI. *EBioMedicine.* 49: 341-353.

1099

Figure legends

Figure 1. **circHMGCS1 Upregulation and its Association with PAHG-induced Endothelial Dysfunction.** (A) Heat map illustrating the differential expression of circRNAs in HUVECs treated with PAHG for 24 hours (n=3). (B) qRT-PCR validation of five circRNAs in HUVECs treated with PAHG compared to normal cells, normalized to GAPDH (n=6). (C) PCR validation of circHMGCS1 presence in HUVECs, GAPDH was utilized as a negative control (n=6). (D) Alignment of circHMGCS1 sequence in CircBase (circular RNA database, upper) in agreement with Sanger sequencing results (lower). (E) Schematic representation of circularization of exons 2 to 7 of HMGCS1 (red arrow). (F) RNA-FISH analysis detecting circHMGCS1 expression in HUVECs using Cy3-labeled probes, Nuclei were counterstained with DAPI (red represents circHMGCS1, blue represents nucleus, scale bar = 50 μm, n=4). (G) qRT-PCR quantification of circHMGCS1 and HMGCS1 mRNA levels in the cytoplasm or nucleus of HUVECs. circHMGCS1 and HMGCS1 mRNA levels were normalized to cytoplasmic values (n=4). (H) qRT-PCR measure circHMGCS1 and HMGCS1 mRNA levels after actinomycin D treatment (n=4 at different time points). (I) circHMGCS1 expression was detected in RNase R-treated total RNA by qRT-PCR (n=4). $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, All significant difference was determined by unpaired two-tailed Student's t-test or one-way ANOVA followed by Bonferroni multiple comparison post hoc test, error bar indicates SD.

Figure 2. circHMGCS1 Overexpression Aggravates PAHG-induced Endothelial Dysfunction. (A) HUVECs were transfected with either circHMGCS1 overexpression lentivirus (pLV-circHMGCS1) or lentiviral circular RNA negative control vector (pLV-circNC). After 48 hours of infection, circHMGCS1 and HMGCS1 expression levels were assessed by qRT-PCR and normalized to GAPDH (n=4). (B and C) Western blot and qRT-PCR were conducted to detect the expressions of adhesion molecules (VCAM1, ICAM1, and ET-1) in HUVECs

between pLV-circHMGCS1 and pLV-circNC groups (n=4). (D) NO content in pLV-circHMGCS1-infected HUVECs after PAHG treatment (n=4). (E) eNOS activity in pLV-circHMGCS1-infused HUVECs after PAHG treatment (n=4). (F) DHE fluorescence was used to characterize ROS expression in different groups. Scale bar = 50 μm (n=8). (G and H) Relative expression of adhesion molecules (ICAM1, VCAM1 and ET-1) in HUVECs infected with pLV-circNC or pLV-circHMGCS1 after PAHG treatment were determined by Western blot and qRT-PCR (n=4). $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, All significant difference was determined by one-way ANOVA followed by Bonferroni multiple comparison post hoc test, error bar indicates SD.

Figure 3. miR-4521 Prevents PAHG-induced Endothelial Dysfunction. (A) Heat map depicting differentially expressed miRNAs in HUVECs with or without PAHG treatment for 24 hours (n=3). (B) Volcano plot illustrating significant changes in miRNAs between control and PAHG-treated groups (Fold-change > 2 or < 0.5, $p \leq 0.01$). (C) Schematic illustration showing overlapping target miRNAs of circHMGCS1 predicated by miRanda and sequencing results. (D) The binding sites of circHMGCS1 were predicted using miRanda and involve hsa-miR-98-5p, hsa-miR-3143, hsa-miR-4521, and hsa-miR-181a-2-3p. (E) Relative expression of four miRNA candidates in HUVECs treated with PAHG was assessed using qRT-PCR (n=4). (F and G) The regulatory effects of four miRNA mimics on the protein expression levels of adhesion molecules (ICAM1, VCAM1, and ET-1) (n=4). (H) Regulation of NO content by miR-4521 mimic or miR-4521 inhibitor under PAHG treatment (n=4). (I) Impact of miR-4521 mimic or miR-4521 inhibitor on eNOS activity during PAHG treatment. (n=4). (J) The ROS expression in miR-4521 mimic or miR-4521 inhibitor combined with PAHG treatment (n=8). (K) Adhesion molecules (ICAM1, VCAM1, and ET-1) expression in PAHG-treated HUVECs transfected with miR-4521 mimics was determined by Western blot (n=4). (L) Determination of adhesion molecules (ICAM1, VCAM1, and ET-1) expression via Western blot in PAHG-treated HUVECs transfected with miR-4521 inhibitor

(n=4). *p < 0.05, **p < 0.01, ***p < 0.001, All significant difference was determined by one-way ANOVA followed by Bonferroni multiple comparison post hoc test, error bar indicates SD.

Figure 4. **circHMGCS1 Regulates PAHG-induced Endothelial Dysfunction by Targeting and Sponging miR-4521**. (A) Luciferase reporter constructs containing wild-type (WT) or mutant (MUT) circHMGCS1 were cotransfected with miR-4521 mimics, miR-NC, miR-4521 inhibitor, or miR-NC inhibitor in HEK293T cells (n=4). (B) Immunoprecipitation shows AGO2-mediated binding of circHMGCS1 and miR-4521 (n=4). (C) Biotin-coupled miR-4521 or its mutant probe was employed for circHMGCS1 pull-down, and captured circHMGCS1 level was quantified by qRT-PCR (n=4). (D) RNA-FISH showing the colocalization of circHMGCS1 and miR-4521 in HUVECs (red represents circHMGCS1, green represents miR-4521, blue represents nucleus, scale bar = 50 μm, n=4). (E and F) circHMGCS1 exhibited no impact on NO content and eNOS activity in the presence of miR-4521 sponge (n=4). (G) circHMGCS1 demonstrated no influence on ROS expression with miR-4521 sponge (n=8). (H-J) circHMGCS1 had no effect on the expression of adhesion molecules (ICAM1, VCAM1, and ET-1) in the presence of miR-4521 sponge which were determined by Western blot and qRT-PCR (n=4). (K and L) miR-4521 attenuated the reduction of NO content and eNOS activity induced by circHMGCS1 (n=4). (M) miR-4521 inhibited the increase of ROS expression caused by circHMGCS1 (n=8). (N-P) miR-4521 attenuated the increased expression of adhesion molecules (ICAM1, VCAM1 and ET-1) induced by circHMGCS1 which were determined by Western blot and qRT-PCR, respectively (n=4). *p < 0.05, **p < 0.01, ns means no significant. All significant difference was determined by one-way ANOVA followed by Bonferroni multiple comparison post hoc test, error bar indicates SD.

Figure 5. AAV9-mediated circHMGCS1 Overexpression Attenuates the Protective Effect of miR-4521 against Diabetes-induced VED. (A) Changes in

body weight of mice from 0 to 14 weeks (n=8, ***p<0.001 DM versus control, #p<0.05 DM + agomir miR-4521 versus DM). (B) Alterations in fasting blood glucose in mice from 0 to 14 weeks (n=8, ***p<0.001 DM versus control, #p<0.05 DM + agomir miR-4521 versus DM). (C and D) ITT and OGTT were performed in the fasted mice (n=8, *p<0.05 DM versus control, #p<0.05 DM + agomir miR-4521 versus DM). (E) Relaxation responses of aortic rings from control, miR-4521 agomir, or miR-4521 agomir+AAV9 circHMGCS1 mice on DM diet (n=6). (F) SBP measured via tail-cuff method in different groups (n=8). (G) Hematoxylin and eosin staining (H&E) was performed on serial cross-sections of thoracic aortas from differently treated mice to assess vessel wall thickness. Scale bar: 100 μm (n=6). (H and I) NO content and eNOS activity in thoracic aorta after 6 weeks of DM diet with miR-4521 agomir or miR-4521 agomir + circHMGCS1 treatment (n=6). (J and K) Detection of ROS expression in thoracic aorta using DHE after miR-4521 agomir or miR-4521 agomir+AAV9 circHMGCS1 injection (red represents ROS, green represents GFP, blue represents nucleus, scale bar = 100 μm, n=6). (L) Expression levels of adhesion molecules (ICAM1, VCAM1, and ET-1) in thoracic aorta after 6 weeks of DM diet with miR-4521 agomir or miR-4521 agomir + circHMGCS1 treatment (n=6). *p < 0.05, **p < 0.01, ***p < 0.001. All significant difference was determined by one-way ANOVA followed by Bonferroni multiple comparison post hoc test, error bar indicates SD.

Figure 6. circHMGCS1 Functions as a miR-4521 Sponge in HUVECs to Modulate ARG1 Expression. (A) Schematic depiction illustrating the intersection of predicted target genes of miR-4521 from TargetScan, miRWalk, mirDIP, and GeneCard. (B) ARG1 expression in thoracic aorta of DM by western blot (n=6). (C) HEK293T cells were cotransfected with miR-4521 mimics, miR-NC, miR-4521 inhibitor, or miR-NC inhibitor, along with luciferase reporter constructs containing WT or MUT 3′-untranslated region of ARG1 (n=4). (D and E) ARG1-overexpressing HUVECs were generated using lentivirus and transfected with miR-4521 mimics for 48h to evaluate adhesion molecule expression (ICAM1,

VCAM1, and ET-1) by Western blot and qRT-PCR (n=4). (F and G) HUVECs were treated as shown. the intracellular NO content and eNOS activity in HUVECs were analyzed (n=4). (H) DHE fluorescence was used to assess ROS expression in different groups. Scale bar = 100 μm (n=8). (I-K) ARG1 expression was significantly decreased by miR-4521 mimics and increased by miR-4521 inhibitor, as determined by qRT-PCR and Western blot (n=4). (L and M) ARG1 expression was significantly increased by circHMGCS1 overexpression, as determined by Western blot and qRT-PCR (n≥4). (N and O) circHMGCS1-overexpressed HUVECs, created with lentivirus and transfected with miR-4521 mimics for 48h, were examined for ARG1 expression by Western blot and qRT-PCR (n=4). (P) AAV9 miR-4521 sponge was used to inhibit the miR-4521 expression in HUVECs, followed by circHMGCS1 transfection, and then treated with PAHG for 24 hours to evaluate the expression of ARG1 by Western blot (n=4). (Q) Detection of ARG1 expression in the thoracic aorta of DM treated with AAV9 miR-4521 agomir combined with AAV9 circHMGCS1 by Western blot (n=4). *p < 0.05, **p < 0.01, ns means no significant. All significant difference was determined by one-way ANOVA followed by Bonferroni multiple comparison post hoc test, error bar indicates SD.

Figure 7. ARG1 is Inseparable from circHMGCS1 and miR-4521 Regulating Diabetes-Induced VED. (A and B) circHMGCS1 and miR-4521 had no effect on NO content and eNOS activity expression in the absence of ARG1 (n=4). (C) ROS expression remained unaffected by circHMGCS1 and miR-4521 in the absence of ARG1 (n=4). (D and E) Relative expression of adhesion molecules (ICAM1, VCAM1, and ET-1) remained unchanged in the absence of ARG1, despite the presence of circHMGCS1 and miR-4521, as determined by qRT-PCR and Western blot (n=4). (F) Changes in mice body weight over the experimental period (n=8, **p<0.01 HFHG versus control, #p<0.05 HFHG + AAV9 ARG1 shRNA versus HFHG). (G) Fasting blood glucose levels in mice over time (n=8, **p<0.01 HFHG versus control, #p<0.05 HFHG + AAV9 ARG1 shRNA versus HFHG). (H and I)

Blood glucose levels measured at week 13 (ITT) and week 14 (OGTT) (n=8,

**$p<0.01$ HFHG versus control, #$p<0.05$ HFHG + AAV9 ARG1 shRNA versus

HFHG). (J) Endothelium-dependent relaxations in aortic rings from different groups

(n=8). (K) H&E performed on serial cross-sections of thoracic aortas from

differently treated mice to evaluate vessel wall thickness. Scale bar = 100 μm (n=6).

(L) SBP measured by the tail-cuff method in different groups (n=8). (M and N) NO

content and eNOS activity expression in thoracic aort (n=6). (O) Relative expression

of adhesion molecules (ICAM1, VCAM1, and ET-1) in thoracic aorta assessed by

Western blot (n=6). (P and Q) ROS expression in the thoracic aorta using the DHE

probe (Red represents ROS, Green represents GFP, Blue represents nucleus, scale

bar: 100 μm, n=6). *$p < 0.05$, **$p < 0.01$, ns means no significant. All significant

difference was determined by one-way ANOVA followed by Bonferroni multiple

comparison post hoc test, error bar indicates SD.

Supplemental   Figure legend

Figure S1. Distinct expression profiles of circRNAs in endothelial dysfunction of HUVECs. HUVECs exposed to palmitic acid and high glucose (PAHG) treatment for 24 hours were compared with untreated Control. (**A**) Alterations in NO content between the control and PAHG-treated groups (n=4). (**B**) Changes in eNOS activity between the control and PAHG-treated groups (n=4). (**C**) Relative expression levels of adhesion molecules (ICAM1, VCAM1, and ET-1) in HUVECs were assessed by Western blotting between the control and PAHG-treated groups (n=4). (**D**) Volcano plot illustrating significantly altered circRNAs between the control and PAHG-treated groups. Fold-change > 2 or < 0.5, $p \leq 0.001$. (**E**) Number of identified circRNAs in Control groups and PAHG-treated groups (n=3). $*p < 0.05$, $**p < 0.01$, All significant difference was determined by unpaired two-tailed Student's t-test, error bar indicates SD.

Figure S2. A sketch map of circHMGCS1 plasmid construction.

Figure S3. miR-4521 inhibition aggravates diabetes-induced endothelial dysfunction. (A) Relative expression of hsa-miR-98-5p, hsa-miR-3143, hsa-miR-181a-2-3p and hsa-miR-4521 were detected in HUVECs after corresponding miRNA mimics treatment by qRT-PCR (n=4). (**B** and **C**) Relative expression level of miR-4521 was determined by qRT-PCR in HUVECs transfected with miR-4521 mimics or miR-4521 inhibitor (n=4). (**D**) miR-4521 mimic inhibited the expression of adhesion molecules (ICAM1, VCAM1, and ET-1) in PAHG-treated HUVEC as detected by qRT-PCR (n=4). (**E**) miR-4521 inhibitor exacerbated the expression of adhesion molecules (ICAM1, VCAM1, and ET-1) in PAHG-treated HUVEC as detected by qRT-PCR (n=4). $*p < 0.05$, $**p < 0.01$, $*** p < 0.001$, $**** p < 0.0001$ All significant difference was determined by one-way ANOVA followed by Bonferroni multiple comparison post hoc test, error bar indicates SD.

Figure S4. Verification of the binding site of miR-4521 and circHMGCS1. HEK293T cells were used to package lentivirus overexpressing circHMGCS1, and then infected HUVECs cells, (**A**) Expression level of miR-4521 was assessed by qRT-PCR in pLV-circHMGCS1-transfected HEK293T cells (n=4). (**B**) Schematic representation of the predicted binding sites of miR-4521 on the circHMGCS1 transcript in human cells. $**p < 0.01$, significant difference was determined by

unpaired two-tailed Student's t-test, error bar indicates SD.

Figure S5. circHMGCS1 counteracts the protective role of miR-4521 in diabetes-induced
VED. (**A**) Timeline illustrating the HFHG feeding, AAV9 circHMGCS1, and miR-4521 agomir
treatment schedule. (**B** and **C**) Quantification of the area under the curve (AUC) for glucose and
insulin tests (n=8). (**D**) Triglyceride (TG), (**E**) Total cholesterol (T-CHO), (**F**) High-density
lipoprotein cholesterol (HDL-C), and (**G**) Low-density lipoprotein cholesterol (LDL-C) levels in
the serum were measured using Automatic Biochemical Analyzer (n=6). (**H**) Effects on
endothelium-independent vasorelaxation to SNP (n=6). *p < 0.05, **p < 0.01. All significant
difference was determined by one-way ANOVA followed by Bonferroni multiple comparison post
hoc test, error bar indicates SD.

Figure S6. miR-4521 modulates the expression of ARG1. (**A**) Schematic representation of the
predicted binding sites and mutated sites for miR-4521 on ARG1 transcripts in both human cells
and mice. (**B**) Western blot analysis assessing ARG1 and ARG2 expression in HUVECs after
PAHG treatment (n=6). (**C**) Sequencing results displaying the mutation site in the dual luciferase
assay. (**D** and **E**) Modulatory effect of miR-4521 mimic on NO content and eNOS activity under
ARG1 overexpression (n=4). (**F** and **G**) Regulatory impact of miR-4521 mimic on ROS content
under ARG1 overexpression (n=4). *p < 0.05, **p < 0.01, ns means no significant. All significant
difference was determined by one-way ANOVA followed by Bonferroni multiple comparison
post hoc test, error bar indicates SD.

Figure S7. Loss of functional regulation of circHMGCS1 and miR-4521 on HUVECs in the
absence of ARG1 in vivo. (**A**) Schematic outlining the timeline of HFHG feeding, AAV9
circHMGCS1 and miR-4521 agomir treatment, AAV9 ARG1 shRNA treatment, and experiment
conclusion. (**B** and **C**) Blood glucose levels measured at week 13 (ITT) and week 14 (OGTT), the
corresponding AUC for glucose or insulin tests was calculated (n=8, **p<0.01 HFHG versus
control, #p<0.05 HFHG + AAV9 ARG1 shRNA versus HFHG). (**D**) TG, (**E**) T-CHO, (**F**) HDL-C,
and (**G**) LDL-C levels in the serum of different treated groups (n=6). (**H**) Assessment of
endothelium-independent vasorelaxation to SNP (n=6). *p < 0.05, **p < 0.01, *** p < 0.001, ns
means no significant. All significant difference was determined by one-way ANOVA followed by

Bonferroni multiple comparison post hoc test, error bar indicates SD.

A

miR-4521 hARG1 3'-UTR binding site

hsa-ARG1 wt: 5' taACTTGGCAAAAGACTTATCCTTAGa 3'
                     ||| :|||| | |||| |||||||
Hsa-miR-4521:   3' ttTGACTCGTGTCCTGA--AGGAATCg 5'
                     ××× :××× × ××××   ×××××××
hsa-ARG1 mut: 5' taTGGTGCGTATACTGGTAGGGTTTCa 3'

miR-4521 mARG1 3'-UTR binding site

mmu-ARG1 wt: 5' catTTTAGCATAAGTTATCCTTAag 3'
                     | :| ||||:| | : ||||||
Hsa-miR-4521:   3' tttGACUCGUGUCCUGAAGGAAUcg 5'
                     ××× :××× × ××××   ×××××××
mmu-ARG1 mut: 5' catGGCTGGAGTCCTTGGGGAATag 3'

C

Sequencing results of fluorescent reporter vector h-pmirGLO-ARG1-3'UTR-WT

Sequencing results of fluorescent reporter vector h-pmirGLO-ARG1-3'UTR-MUT

F

pLV-NC   pLV-ARG1

pLV-ARG1+   pLV-ARG1+
miR-NC   miR-4521 mimic