Circular RNA *HMGCS1* sponges *MIR4521* to aggravate type 2 diabetes-induced

vascular endothelial dysfunction

Ming Zhang[1], Guangyi Du[1], Lianghua Xie[1], Yang Xu[1,2], Wei Chen[1,2*]

[1] Department of Food Science and Nutrition, College of Biosystems Engineering and

Food Science, Zhejiang University, Hangzhou, 310058, China.

[2] Ningbo Innovation Center, Zhejiang University, Ningbo 315100, China.

[*]Corresponding author: Wei Chen. Department of Food Science and Nutrition,

Zhejiang University, No.866 Yuhangtang Road, Xihu District, Hangzhou 310058,

China. Phone: +86 0571 88982191; E-mail: zjuchenwei@zju.edu.cn.

Abstract

Noncoding RNA plays a pivotal role as novel regulators of endothelial cell function. Type 2 diabetes, acknowledged as a primary contributor to cardiovascular diseases, plays a vital role in vascular endothelial cell dysfunction due to induced abnormalities of glucolipid metabolism and oxidative stress. In this study, aberrant expression levels of *circHMGCS1* and *MIR4521* were observed in diabetes-induced human umbilical vein endothelial cell dysfunction. Persistent inhibition of *MIR4521* accelerated development and exacerbated vascular endothelial dysfunction in diabetic mice. Mechanistically, *circHMGCS1* upregulated arginase 1 by sponging *MIR4521*, leading to decrease in vascular nitric oxide secretion and inhibition of endothelial nitric oxide synthase activity, and an increase in the expression of adhesion molecules and generation of cellular reactive oxygen species, reduced vasodilation and accelerated the impairment of vascular endothelial function. Collectively, these findings illuminate the physiological role and interacting mechanisms of *circHMGCS1* and *MIR4521* in diabetes-induced cardiovascular diseases, suggesting that modulating the expression of *circHMGCS1* and *MIR4521* could serve as a potential strategy to prevent diabetes-associated cardiovascular diseases. Furthermore, our findings provide a novel technical avenue for unraveling ncRNAs regulatory roles of ncRNAs in diabetes and its associated complications.

Introduction

Cardiovascular disease (CVD) is the leading cause of morbidity and mortality in patients with type 2 diabetes mellitus (T2DM). The incidence of CVD in T2DM individuals was estimated to be at least two to four times higher than that in non-diabetic individuals (Gregg et al, 2016; Yun & Ko, 2021). T2DM gives rise to abnormal metabolic conditions, including chronic hyperglycemia, dyslipidemia, insulin resistance, inflammation, and oxidative stress (Roden & Shulman, 2019). These processes ultimately lead to vascular endothelial damage, which disrupts the maintenance of vascular homeostasis, thereby promoting the development of CVD (Lundberg & Weitzberg, 2022; Niemann et al, 2017). Vascular endothelial dysfunction (VED) is considered the underlying basis for vascular complications at all stages of T2DM (Eelen et al, 2015; Meigs et al, 2004). However, the mechanism by which the key molecules trigger endothelial dysfunction in diabetes-induced vascular impairment remains unknown.

Endothelial cells form a continuous monolayer lining the arterial, venous, and lymphatic vessels, playing a crucial physiological role in maintaining vascular homeostasis (Bazzoni & Dejana, 2004). Moreover, endothelial cells actively regulate vascular tone, preserve endothelial integrity, and modulate platelet activity, contributing to hemostasis and endothelial repair (Li et al, 2019). The leading cause of T2DM is a prolonged high-sugar, high-fat diet (HFHG), commonly known as a Western diet, which inevitably leads to VED, characterized by reduced content of nitric oxide (NO), uncoupling of endothelial nitric oxide synthase (*ENOS*), increased formation of reactive oxygen species (ROS) and upregulation of endothelial-related adhesion molecules such as intercellular adhesion molecule 1 (*ICAM1*), vascular cell adhesion molecule 1 (*VCAM1*), and endothelin 1 (*ET-1*). These alterations ultimately lead to elevated blood pressure and impaired vascular arteriolar relaxation, contributing to CVD development (Xu et al, 2021; Yeh et al, 2022; Zheng et al, 2018). Despite this understanding, the identification of pivotal regulatory factors

governing gene expression in diabetes-induced VED remains a formidable

challenge.

circRNAs are generated through back-splicing of precursor messenger RNAs

(pre-mRNAs), exhibiting features such as a covalently closed loop structure, high

stability, conservation, and tissue-specific expression. These molecules primarily

function as miRNA sponges, regulators of transcription, and interactors with proteins

(Kristensen et al, 2019; Liu & Chen, 2022). Conversely, miRNAs are single-stranded

small RNA molecules of 21–23 nucleotides, characterized by high conservation,

temporal and tissue-specific expression, and relatively poor stability.

miRNAs—critical regulators of gene expression—control a wide array of cellular

processes (Gebert & MacRae, 2019). The interplay between miRNAs and circRNAs

forms a complex regulatory network, profoundly influencing diverse biological

pathways and disease states. miRNAs serve as guides, targeting specific mRNAs to

induce their degradation or translational repression, thereby suppressing gene

expression post-transcriptionally (Treiber et al, 2019). By contrast, circRNAs contain

miRNA binding sites, competitively inhibiting miRNA activity and thereby reducing

miRNA binding to other mRNAs. This competitive adsorption effect modulates

intracellular miRNA levels, affecting miRNA-mediated regulation of other target

genes (Cheng et al, 2019; Huang et al, 2019; Shen et al, 2019; Zeng et al, 2021).

The expression levels of circRNA in diabetic patients exhibit abnormalities, and

its potential as a biomarker for early diagnosis or therapeutic target has been

gradually being substantiated (Jiang et al, 2022; Stoll et al, 2020; Tian et al, 2018).

However, research on circRNA in the context of diabetes-induced VED is scarce.

Current research has predominantly focused on the regulatory roles of circRNA in

diabetic cardiomyopathy (Yuan et al, 2023), diabetic nephropathy (Yang & Liu,

2022), gestational diabetes (Du et al, 2022), and diabetic retinopathy (Zhu et al,

2019). Investigations into circRNA regulation of VED have only described the

phenomenon and analyzed the correlation of the results, lacking the precision to infer

causation accurately (Jiang et al, 2020; Liu et al, 2017; Ma et al, 2023). Moreover,

comprehension of VED induced by the complex multifactorial processes of diabetes

pathogenesis remains relatively inadequate. Additionally, the interaction between

circRNA and miRNA in diabetes-induced VED remains a subject of research

controversy (Cheng et al, 2019; Liu et al, 2019; Pan et al, 2018). Therefore, a more

comprehensive research approach is required to elucidate the specific mechanisms

by which circRNA operates in diabetes-related CVDs.

The present study elucidated the mechanisms underlying the interaction

between circRNA and miRNA in regulating diabetes-associated VED. Through the

screening strategy, we successfully identified *circHMGCS1*—a circRNA originating

from the pre-mRNA of *HMGCS1*. We demonstrated that the upregulation of

circHMGCS1 and downregulation of *MIR4521* significantly promoted

diabetes-induced VED. Moreover, in-depth mechanistic investigations revealed that

circHMGCS1 functioned as a *MIR4521* sponge, increasing arginase 1 (*ARG1*)

expression in vascular endothelial cells and accelerating VED progression.

Furthermore, the expression patterns and interaction mechanisms between

circHMGCS1 and *MIR4521* in endothelial cells were identified for the first time in

these findings, which also highlight the *circHMGCS1/MIR4521/ARG1* axis as a

novel therapeutic target for interventions designed to protect patients from

diabetes-induced VED.

Results

Global expression analysis of endotheliocyte circRNAs

To explore the potential role of circRNAs in regulating endotheliocyte function, we characterized the transcriptome of circRNA derived from total RNA in human umbilical vein endothelial cells (HUVECs) stimulated by high palmitate and high glucose (PAHG), which allowed us to investigate the potential role of circRNAs in diabetes-induced VED (**Figure 1-figure supplement 1A-C**). A total of 17,179 known circRNAs were identified, with 66 of them exhibiting differential expression (GEO submission: GSE237597). Our main focus was on circRNAs that shared identical genomic and splicing lengths that exhibited fold changes greater than 2 or less than -2 (p < 0.01) in PAHG-treated HUVECs (**Figure 1A**). Specifically, 48 circRNAs were upregulated, whereas 18 were downregulated (**Figure 1-figure supplement 1D**). Furthermore, approximately 66.65% of circRNAs were found to be produced through reverse splicing of exons (**Figure 1-figure supplement 1E**).

The observed changes in circRNA levels were confirmed through the real-time quantitative reverse transcription PCR (qRT-PCR) analysis of the five most upregulated circRNAs, suggesting that the results of the RNA-seq data are credible. Among them, *circHMGCS1* (hsa_circ_0008621, 899 nt in length, identified as *circHMGCS1* in subsequent studies because of its host gene being *HMGCS1* (**Figure 1-figure supplement 1F**) (Liang et al, 2021)) exhibited significant upregulation compared with the non-PAHG-treated condition (**Figure 1B**). The *circHMGCS1* sequence is situated on chromosome 5 of the human genome, specifically at 43292575-43297268 with no homology to mouse sequences. Subsequently, divergent primers were designed to amplify *circHMGCS1*, whereas convergent primers were designed to amplify the corresponding linear mRNA from cDNA and genomic DNA (gDNA) in HUEVCs. amplification of *circHMGCS1* was observed in cDNA but not in gDNA (**Figure 1C**), which confirmed the circularity of *circHMGCS1*. Sanger sequencing of the amplified product of *circHMGCS1*

confirmed its sequence alignment with the annotated *circHMGCS1* in circBase

(**Figure 1D**). This observation confirms the origin of *circHMGCS1* from exons 2 to

7 of the *HMGCS1* gene (**Figure 1E and Figure 1-figure supplement 1G, H**).

RNA-Fluorescence *in situ* hybridization (RNA-FISH) confirmed the robust

expression of cytoplasmic *circHMGCS1* in HUVECs (**Figure 1F**). The qRT-PCR

analysis of nuclear and cytoplasmic RNA revealed the predominant cytoplasmic

expression of *circHMGCS1* in HUVECs (**Figure 1G**). qRT-PCR also demonstrated

that *circHMGCS1* displayed a stable half-life exceeding 24 h, whereas the linear

transcript *HMGCS1* mRNA had a half-life of less than 8 h (**Figure 1H**). Furthermore,

circHMGCS1 exhibited resistance to digestion by ribonuclease R (an exonuclease

that selectively degrades linear RNA (Xiao & Wilusz, 2019), also known as RNase

R), whereas linear *HMGCS1* mRNA was easily degraded upon RNase R treatment

(**Figure 1I**). These findings indicate that the higher expression of *circHMGCS1* in

VED is more than just a byproduct of splicing and suggestive of functionality.

Upregulation of circHMGCS1 promotes diabetes-induced VED

To explore the potential role of *circHMGCS1* in regulating endothelial cell function,

we cloned exons 2–7 of *HMGCS1* into lentiviral vectors for ectopic overexpression

of *circHMGCS1* (**Figure 2-figure supplement 1**). We found that *circHMGCS1* was

successfully overexpressed in HUVECs without significant changes in *HMGCS1*

mRNA expression, and the level of the circular transcripts increased nearly 60-fold

than the level of the linear transcripts (**Figure 2A**). These results confirm that the

circularization is efficient and that *circHMGCS1* has no effect on *HMGCS1*

expression. Further experiments demonstrated that the overexpression of

circHMGCS1 stimulated the expression of adhesion molecules (*VCAM1*, *ICAM1*,

and *ET-1*) (**Figure 2B, C**), suggesting that *circHMGCS1* is involved in VED

promotion. We then used PAHG to stimulate HUVEC-overexpressing *circHMGCS1*.

NO levels were significantly decreased after PAHG stimulation for 24 h, and

circHMGCS1 overexpression further decreased NO levels (**Figure 2D**), indicating

that increased *circHMGCS1* expression inhibits endothelial diastolic function. *ENOS*

activity was inhibited by PAHG, and upregulated *circHMGCS1* expression further

decreased *ENOS* activity (**Figure 2E**). Superoxide is one of the major ROS known to

promote atherosclerosis (Griendling et al, 2016). Therefore, we used DHE to assess

superoxide levels in endothelial cells. The overexpression of *circHMGCS1* enhanced

PAHG-induced ROS generation (**Figure 2F**), increasing oxidative stress in

endothelial cells. The expression levels of adhesion molecules (*VCAM1*, *ICAM1*, and

ET-1) were further elevated with *circHMGCS1* overexpression (**Figure 2G, H**).

These findings suggest that elevated expression of *circHMGCS1* may contribute to

the development of endothelial cell dysfunction.

***MIR4521* protects endothelial cells from PAHG-induced dysfunction**

circRNAs may function as ceRNAs to sponge miRNAs, thereby modulating miRNA

target depression and imposing an additional level of post-transcriptional regulation

in human diseases (Zhong et al, 2018). Accordingly, we conducted miRNA

sequencing in PAHG-stimulated HUVECs to identify the potential miRNA targets of

circHMGCS1. The deep sequencing analysis revealed 98 significantly differentially

expressed miRNAs (**Figure 3A, B**, GEO submission: GSE237295). We then used

the miRanda database and ceRNA theory to obtain four candidate miRNAs

(*MIR4521*, *MIR3143*, *MIR98-5P*, and *MIR181A-2-3P*) (**Figure 3C, D**). qRT-PCR

confirmed that all four miRNAs were downregulated (**Figure 3E**). Next, we

observed that each of the four miRNA mimics induced a significantly increase in the

expression of their corresponding miRNAs in HUVECs through the application of

synthetic miRNA mimics (**Figure 3-figure supplement 1A**). Relative to the other

three miRNAs, *MIR4521* mimics significantly inhibited adhesion molecules (*ICAM1*,

VCAM1, and *ET-1*) expression in HUVECs (**Figure 3F, G**), making it a suitable

candidate for further endothelial function studies. We then investigate the effect of

MIR4521 on PAHG-stimulated endothelial cell function using synthetic *MIR4521*

mimics and *MIR4521* inhibitor in HUVECs (**Figure 3-figure supplement 1B, C**).

Enhanced *MIR4521* levels effectively restored PAHG-induced reductions in NO

content and *ENOS* activity in HUEVCs, while *MIR4521* inhibition led to opposite

results (**Figure 3H, I**). whereas *MIR4521* mimics significantly inhibited

PAHG-induced expression of ROS and adhesion molecules (*VCAM1*, *ICAM1*, and

ET-1) (**Figure 3J**), whereas the *MIR4521* inhibitor increased the PAHG-induced

expression of ROS and adhesion molecules (*VCAM1*, *ICAM1*, and *ET-1*) (**Figure 3K,**

L and **Figure 3-figure supplement 1D, E**). These findings suggest that *MIR4521* is

involved in PAHG-induced endothelial cell dysfunction.

***circHMGCS1* acts as a *MIR4521* sponge to regulate endothelial dysfunction**

We then explored the interaction mechanism between *circHMGCS1* and *MIR4521*.

As displayed in **Figure 4-figure supplement 1A, B**, the overexpression of

circHMGCS1 reduced the expression level of *MIR4521*, whereas knockdown of

circHMGCS1 resulted in an upregulation of *MIR4521*. Meanwhile, bioinformatics

analysis revealed the predicted binding sites between *MIR4521* and *circHMGCS1*

(**Figure 4-figure supplement 1C**). We constructed a dual-luciferase reporter by

inserting the wild-type (WT) or mutant (MUT) linear sequence of circHMGCS1 into

the pmirGLO luciferase vector. The co-transfection of *MIR4521* mimics with the

circHMGCS1 luciferase reporter gene in HEK293T cells, reducing luciferase activity;

mutations in the binding sites between *circHMGCS1* and *MIR4521* abrogated the

effect of *MIR4521* mimics on the activity of the *circHMGCS1* luciferase reporter

gene mutant (**Figure 4A**). Furthermore, AGO2 immunoprecipitation in HUVECs

transfected with *MIR4521* or its mutants demonstrated that *MIR4521* facilitates the

association of *AGO2* with *circHMGCS1* in HUVECs (**Figure 4B**). Meanwhile, we

used biotinylated *MIR4521* mimics in HUVECs stably overexpressing *circHMGCS1*

to further validate the direct binding of *MIR4521* and *circHMGCS1*. The qRT-PCR

results revealed that *MIR4521* captured endogenous *circHMGCS1*, and the negative

control with a disrupting putative binding sequence failed to coprecipitate

circHMGCS1 (**Figure 4C**). Consistently, the use of biotin-labeled *circHMGCS1*

effectively captured both *MIR4521* and *AGO2* (**Figure 4-figure supplement 1D, E**). Moreover, the RNA-FISH assay showed colocalization of *MIR4521* and *circHMGCS1* in the cytoplasm (**Figure 4D**). Altogether, these findings indicate that *circHMGCS1* functions as a molecular sponge for *MIR4521*.

Rescue experiments were conducted to investigate whether *circHMGCS1* regulates endothelial cell function through *MIR4521* sponging. We achieved *MIR4521* inhibition by transfecting HUVECs with a recombinant adeno-associated virus 9 (AAV9) vector carrying *MIR4521* sponges. Notably, *MIR4521* sponges significantly promoted the PAHG-induced reduction in NO content and ENOS activity. Under this condition, increased circHMGCS1 expression did not interfere with the changes in NO content and *ENOS* activity (**Figure 4E, F**). Moreover, *MIR4521* sponges elevated ROS expression in PAHG treated HUVECs, whereas no further additional effect on ROS expression was observed upon the overexpression of *circHMGCS1* when *MIR4521* was sponged (**Figure 4G**). Further evaluation of endothelial functional molecules revealed that circHMGCS1 failed to stimulate the expression of these factors when *MIR4521* was sponged (**Figure 4H-J**). Furthermore, the reduction in NO content and *ENOS* activity induced by *circHMGCS1* overexpression was reversed by *MIR4521* mimics (**Figure 4K, L**), and the *circHMGCS1*-mediated increase in ROS content was inhibited by *MIR4521* mimics (**Figure 4M**). Furthermore, the expression of adhesion molecules (*ICAM1*, *VCAM1*, and *ET-1*) was increased in *circHMGCS1*-transfected HUVECs, and this effect was significantly ameliorated by *MIR4521* mimics (**Figure 4N-P**). These findings demonstrate that *circHMGCS1* acts as a sponge for *MIR4521*, thereby regulating endothelial function.

circHMGCS1 serves as a *MIR4521* sponge to regulate diabetes-induced VED

To further explore the interaction between *MIR4521* and *circHMGCS1* in diabetes-induced VED, we administered exogenous *MIR4521* agomir (mimics) via tail vein injection to mice, whereas agomir NC, expressing random sequence, served

as the negative control in HFHG-induced diabetes to examine its effect on diabetes and associated VED (**Figure 5-figure supplement 1A**). Compared with to non-diabetic WT (Control) mice, DM mice exhibited a significant increase in body weight, upon *MIR4521* agomir application, the body weight of diabetic mice significantly decreased compared with the untreated diabetic mice (**Figure 5A**), and elevated fasting blood glucose levels in DM mice were likewise substantially inhibited (**Figure 5B**). By contrast, the combined treatment of *circHMGCS1* and *MIR4521* agomir did not significantly affect the body weight and blood glucose levels. OGTT and ITT experiments demonstrated that *MIR4521* agomir considerably enhanced glucose tolerance and insulin resistance in diabetic mice (**Figures 5C, D** and **Figure 5-figure supplement 1B, C**). Blood lipid biochemistry analysis revealed that *MIR4521* agomir restored abnormal blood lipid levels in diabetic mice (**Figure 5-figure supplement 1D-G**). Notably, the abundant presence of *circHMGCS1* in the body can negate the inhibitory effect of *MIR4521* on diabetes development. These results suggest that *MIR4521* can inhibit the occurrence of diabetes, whereas *circHMGCS1* specifically dampens the function of *MIR4521*, weakening its protective effect against diabetes.

To investigate the roles of *MIR4521* and *circHMGCS1* in mouse vascular relaxation, we conducted an *ex vivo* culture of thoracic aortas from mice subjected to different treatments. A significant impairment in acetylcholine (Ach)-induced vascular relaxation response in the thoracic aorta of diabetic mice was markedly improved by *MIR4521* agomir, emphasizing the critical involvement of *MIR4521* in vascular relaxation. Conversely, *circHMGCS1* inhibited the protective function of *MIR4521* on vascular relaxation (**Figure 5E**). Across all experimental groups, the smooth muscle exhibited a normal endothelium-dependent relaxation response to the nitric oxide donor sodium nitroprusside (SNP) (**Figure 5-figure supplement 1H**), revealing intact smooth muscle function. Meanwhile, diabetic mice displayed a significant increase in SBP compared with normal mice, which was significantly restrained by *MIR4521* agomir treatment. Nevertheless, this beneficial effect was

nullified in the presence of *circHMGCS1* (**Figure 5F**). Furthermore, *MIR4521* agomir inhibited intimal thickening and smooth muscle cell proliferation in diabetic mice, whereas upregulation of *circHMGCS1* abrogated this effect (**Figure 5G**). These findings demonstrate that *circHMGCS1* and *MIR4521* play specific roles in regulating vascular endothelial function in diabetic mice.

We measured the NO level to further investigate the interaction between *MIR4521* and *circHMGCS1* in regulating diabetes-induced VED. Compared with the control group, the NO level decreased in the thoracic aorta of diabetic mice; however, *MIR4521* agomir treatment increased its content (**Figure 5H**). Furthermore, *MIR4521* treatment effectively enhanced *Enos* activity (**Figure 5I**), whereas *circHMGCS1* hindered *MIR4521* regulatory function on NO content and *ENOS* activity. Furthermore, *MIR4521* reduced ROS production in the blood vessels (**Figure 5J, K**) and inhibited the expression of endothelial adhesion molecules (*Icam1*, *Et-1*, and *Vcam1*) (**Figure 5L**). Nonetheless, in the presence of *circHMGCS1*, the effect of *MIR4521* was abrogated. These results indicated that *circHMGCS1* acts as a *MIR4521* sponge, participating in diabetes-induced VED.

ARG1 is a direct target of *MIR4521* to accelerate endothelial dysfunction

To investigate whether *circHMGCS1*-associated *MIR4521* regulated VED-related gene expression by targeting the 3'-untranslated region (UTR), we conducted predictions of mRNAs containing *MIR4521* binding sites using the GenCard, mirDIP, miRWalk, and Targetscan databases. Through bioinformatics analyses, we identified binding sites for *MIR4521* on *KLF6*, *ARG1*, and *MAPK1* genes (**Figure 6A**). Previous studies have demonstrated the involvement of elevated arginase activity in diabetes-induced VED. Arginase can modulate the production of NO synthesized by *ENOS* through competitive utilization of the shared substrate L-arginine (Jung et al, 2010; Romero et al, 2008). Moreover, *MIR4521* possesses the capability to bind to the 3' untranslated region of *ARG1* in both human and mouse genomes (**Figure 6-figure supplement 1A**). The protein expression of *Arg1* was increased in the aorta

of diabetic mice and PAHG-treated HUVECs (**Figure 6B** and **Figure 6-figure supplement 1B**). The luciferase reporter assay was applied to verify the targeting ability through the pmirGLO vector, which included either the WT or MUT 3'-UTR of *ARG1* (**Figure 6-figure supplement 1C**). The overexpression of *MIR4521* reduced the luciferase activities of the WT reporter vector but not the MUT reporter vector (**Figure 6C**). We next investigated the function of *MIR4521* and *ARG1* to regulate the function of HUVECs. *MIR4521* mimics recovered the reduced levels of NO and the compromised activity of *ENOS* triggered by *ARG1* overexpression (**Figure 6D, E and Figure 6-figure supplement 1D, E**). Meanwhile, it inhibited the upregulation in ROS levels induced by *ARG1* overexpression (**Figure 6F and Figure 6-figure supplement 1F, G**). The rescue experiment results demonstrated that *MIR4521* mimics reversed the promotional effect of *ARG1* on the mRNA and protein expression of adhesion molecules *in vitro* (**Figure 6G, H**). Subsequently, we evaluated the gene and protein expression levels of ARG1 using qRT-PCR and western blotting. *MIR4521* mimics repressed *ARG1* transcription and protein expression, whereas MIR4521 inhibition upregulated *ARG1* expression (**Figure 6I-K**). Taken together, these results suggest that *MIR4521* inhibits VED through sponging *ARG1*.

To explore the regulatory effect of the interaction between *circHMGCS1* and *MIR4521* on *ARG1* expression. we firstly used HUVECs employing overexpression of *circHMGCS1*. *circHMGCS1* overexpression significantly further promoted *ARG1* transcription and protein expression during the PAHG treatment (**Figure 6L, M**), confirming the regulatory function of *circHMGCS1* in VED-related *ARG1* expression. We next studied the function of *MIR4521*, which acts as a sponge for *circHMGCS1* to regulate *ARG1* in HUVECs. The mRNA and protein levels of *ARG1* were significantly increased when HUVECs were transfected with *circHMGCS1*. However, *MIR4521* overexpression counteracted the effects of *circHMGCS1* on *ARG1* mRNA and protein expression (**Figure 6N, O**). Conversely, knockdown of *circHMGCS1* reduced *ARG1* expression and increased *MIR4521* expression in

HUVECs (**Figure 6P, Q** and **Figure 4-figure supplement 1B**). Besides, the inhibition of *MIR4521* using a *MIR4521* sponge in HUVECs further increased *ARG1* expression under the PAHG treatment. Notably, elevated *circHMGCS1* expression did not affect *ARG1* regulation (**Figure 6R**). Moreover, compared with diabetic mice, *circHMGCS1* overexpression dampened the function of *MIR4521* agomir on *Arg1* (**Figure 6S**). These findings indicate that *circHMGCS1* serves as a sponge for *MIR4521*, facilitating *ARG1* regulation and VED promotion.

ARG1 is essential for *circHMGCS1* and *MIR4521* to regulate diabetes-induced VED

To delve further into how *circHMGCS1* and *MIR4521* regulated endothelial cell function via *ARG1*, we used AAV9 expressing *ARG1* shRNA (*ARG1* shRNA) or GFP (NC shRNA, used as the control) to reduce *ARG1* levels. *ARG1* shRNA mitigated the effects of PAHG on NO content and *ENOS* activity (**Figure 7A, B**), and inhibited PAHG-induced ROS in endothelial cells (**Figure 7C**). Additionally, *ARG1* shRNA significantly reduced the elevated expression of adhesion molecules (*ICAM1*, *VCAM1*, and *ET-1*) induced by PAHG (**Figure 7D, E**). F Further rescue experiments revealed that *MIR4521* and *circHMGCS1* exhibited no significant regulatory effect on endothelial function in the presence of ARG1 shRNA. These findings demonstrate that ARG1 plays a crucial regulatory effect on *MIR4521* and *circHMGCS1* in modulating endothelial cell function.

Considering the role of *ARG1* in regulating endothelial cell function through *circHMGCS1* and *MIR4521*, we downregulated *ARG1* expression in mice by administering AAV9 *ARG1* shRNA through tail vein injection (**Figure 7-figure supplement 1A**). As anticipated, AAV9 *ARG1* shRNA-infused mice exhibited decreased body weight and lower fasting blood glucose levels compared with diabetic mice (**Figure 7F, G**). Moreover, OGTT and ITT demonstrated that reduced *ARG1* levels in AAV9 *ARG1* shRNA-treated mice preserved glucose tolerance and insulin sensitivity (**Figure 7H, I** and **Figure 7-figure supplement 1B, C**). The

lowered *ARG1* levels also reversed serum lipid levels (**Figure 7-figure supplement**

1D-G). However, the overexpression of *circHMGCS1* or *MIR4521* had no

significant effect on glucose and lipid metabolism in AAV9 *ARG1*

shRNA-expressing mice. These findings confirm that *ARG1* is an indispensable

regulator for *circHMGCS1* and *MIR4521* in regulating diabetes.

We next investigated whether *ARG1* regulates vascular endothelial cell function

in the context of diabetes. Knocking down *ARG1* in the thoracic aorta of mice

expressing AAV9 *ARG1* shRNA restored diastolic function of the thoracic aorta in

diabetic mice (**Figure 7J**), emphasizing the vital role of *ARG1* in VED. Furthermore,

all experimental groups exhibited a normal endothelium-dependent relaxation

response to the NO donor SNP, suggesting intact smooth muscle function (**Figure**

7-figure supplement 1H). Moreover, AAV9 *ARG1* shRNA effectively inhibited

diabetes-induced intima thickening and inhibited smooth muscle cell proliferation

(**Figure 7K**), and which also counteracted the elevation of SBP induced by diabetes

(**Figure 7L**). However, the regulatory effects of *circHMGCS1* and *MIR4521* on

vascular dilation were ineffective in the presence of *ARG1* shRNA. We next

measured VED-related functional factors. Compared with diabetic mice, the thoracic

aorta of mice expressing AAV9 *ARG1* shRNA exhibited restored NO content and

Enos activity (**Figure 7M, N**). Furthermore, knocking down *ARG1* reduced the high

expression levels of vascular endothelial adhesion molecules (*Icam1*, *Et-1*, and

Vcam1) induced by diabetes (**Figure 7O**). Immunofluorescence analysis

demonstrated a significant decrease in ROS content in the thoracic aorta of mice

expressing AAV9 *ARG1* shRNA compared with diabetic mice (**Figure 7P, Q**).

Notably, *circHMGCS1* and *MIR4521* lost their regulatory capacity on vascular

endothelial cell function in the presence of *ARG1* shRNA. These results indicate that

ARG1 plays a critical role in *circHMGCS1* and *MIR4521* to modulate

diabetes-induced VED.

Discussion

The field of ncRNA biology has garnered considerable attention and has witnessed intensive research over the last few years. ncRNAs possess the capacity to function as novel regulators in various physiological systems and disease contexts (Esteller, 2011; Matsui & Corey, 2017; Statello et al, 2021). However, most studies have explored the expression patterns and functions of ncRNAs within single disease settings (Arcinas et al, 2019; Shan et al, 2017). The present study contributes to the understanding of ncRNAs and their involvement in diabetes-associated CVD, characterized by disruption of lipid metabolic homeostasis and redox balance in vascular endothelial cells under diabetic conditions. A comprehensive genome-wide analysis revealed 17,179 known circRNA loci transcribed in diabetic endothelial cells. Based on conservation, endothelial specificity, and abundance criteria, five circRNAs were selected for validation, indicating their predominant upregulation in the diabetic endothelial environment. Among them, we discovered a novel circRNA, *circHMGCS1*, exhibiting high abundance, stability, and cell specificity in diabetic endothelial cells and contributing to endothelial function regulation. Our findings establish that *circHMGCS1* promotes VED progression, and its overexpression exacerbates endothelial cell dysfunction. Our study highlights the pivotal role of *circHMGCS1* in regulating endothelial cell function during diabetes.

circRNAs play regulatory roles by acting as microRNA sponges or interacting with proteins to regulate alternative splicing, transcription, and epigenetic modifications (Arcinas et al, 2019; Hansen et al, 2013; Memczak et al, 2013). To investigate the mechanism of circHMGCS1 in VED regulation, we analyzed miRNA sequencing results and prediction software to identify *MIR4521* as a novel endothelial-expressed miRNA. *MIR4521* was found to possess a stable binding site with *circHMGCS1*, and its expression was significantly decreased in endothelial cells within a diabetic environment. Our functional studies revealed that *MIR4521* mimics attenuated VED development in diabetes, whereas its inhibition exacerbated

VED progression. These findings preliminarily demonstrate the regulatory ability of

MIR4521 in VED. The interaction between *circHMGCS1* and *MIR4521* was further

characterized through luciferase reporter gene assays, RNA pull-down, and RNA

immunoprecipitation. AAV9-mediated *circHMGCS1* overexpression in endothelial

cells dampened *MIR4521* expression and accelerated diabetes-induced VED.

Conversely, restoring *MIR4521* expression effectively alleviated VED progression,

highlighting the critical role of *MIR4521* in counteracting *circHMGCS1*-mediated

promotion of diabetes-induced VED. Moreover, the absence of *MIR4521* aggravated

diabetes-induced VED, whereas exogenous *circHMGCS1* addition did not affect

VED development. Intriguingly, exogenous *MIR4521* significantly restrained

diabetes-induced VED exacerbation, which was attenuated upon the subsequent

addition of exogenous *circHMGCS1*. These findings underscore the essential

regulatory role of the *circHMGCS1* and *MIR4521* interaction in maintaining diabetic

vascular homeostasis.

Finally, we investigated the downstream targets crucial for

circHMGCS1-mediated endothelial cell function. Arginase is a dual-nucleus

manganese metalloenzyme that catalyzes the hydrolysis of L-arginine, primarily

producing L-ornithine and urea (Kanyo et al, 1996). *ARG1* and *ARG2*—two

isoforms of arginase—exist in vertebrates, including mammals. Although *ARG1* and

ARG2 share the same catalytic reaction, they are encoded by different genes and

exhibit distinct immunological characteristics (Hara et al, 2020; Pudlo et al, 2017; Su

et al, 2021). Elevated *ARG1* activity in endothelial cells has been identified as a

significant contributor to VED in T2DM individuals. This dysfunction is attributed

to the excessive formation of ROS caused by the competition between *ARG1* and

ENOS for L-arginine. Consequently, oxidases or mitochondrial complexes become

overactivated, resulting in reduced NO levels and subsequent VED (Caldwell et al,

2018). By contrast, inhibiting arginase has been shown to considerably enhance

endothelial function in T2DM patients (Mahdi et al, 2018; Shemyakin et al, 2012;

Zhou et al, 2018). Our findings suggest that T2DM enhances *ARG1* activity and

expression in vascular endothelial cells. Conversely, *ARG1* inhibition in endothelial

cells delayed the onset of diabetes and preserved NO metabolite levels in the aorta.

Additionally, Ach-induced vasodilation remains intact in the absence of *ARG1* in the

aortic ring, as *ARG1* inhibition promotes *ENOS*-dependent relaxation, thereby

preventing VED. Meanwhile, ROS content in the aorta of diabetic mice could be

reduced under *ARG1* inhibition, thereby suppressing the occurrence of oxidative

stress. AAV9-mediated overexpression of *circHMGCS1* increases *ARG1* expression,

which is inhibited by exogenous *MIR4521* intervention. In the absence of *MIR4521*,

circHMGCS1 cannot regulate *ARG1* expression, whereas exogenous *MIR4521*

intervention inhibits the diabetes-induced increase in *ARG1* expression *in vivo*, and

this effect is counteracted by *circHMGCS1* overexpression. Both *in vitro* and *in vivo*

data suggest that the *MIR4521* or *circHMGCS1* fails to regulate the effect of

diabetes-induced VED in the absence of *ARG1*. Therefore, *ARG1* may serve as a

promising VED biomarker, and *circHMGCS1* and *MIR4521* play a key role in

regulating diabetes-induced VED by *ARG1*.

It would be intriguing to elucidate the mechanisms through which *circHMGCS1*

and *MIR4521* exert their regulatory roles in endothelial cell function. The present

study elucidated the potential VED-associated mechanisms in diabetes, with a

specific focus on the complicated mutual effects between *circHMGCS1* and

MIR4521. The ceRNA network involving *circHMGCS1-MIR4521-ARG1* can

become a novel and significant regulatory pathway for preventing and potentially

treating diabetes-induced VED. Our findings suggest that targeted inhibition of

circHMGCS1 or the overexpression of *MIR4521* could serve as effective strategies

in mitigating diabetes-induced VED. These insights underscore the promising

function of ncRNAs in developing therapeutic interventions for diabetes-associated

VED.

Materials and Methods

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
Cell line (H. sapiens)	HUVEC	the Shanghai Cell Bank of the Chinese Academy of Sciences	RRID：CVCL_2959	
Cell line (H. sapiens)	HEK-293T	the Shanghai Cell Bank of the Chinese Academy of Sciences	RRID: CVCL_0063	
Gene (H. sapiens)	circHMGCS1	circbase	circRNA ID: hsa_circ_0008621	
Gene (H. sapiens)	HMGCS1	GeneBank	Gene ID: 3157	
Gene (H. sapiens)	MIR4521	miRbase	miRNA ID: MIMAT0019058	
Gene (M. musculus)	ARG1	GeneBank	Gene ID: 383	
Gene (M. musculus)	Arg1	GeneBank	Gene ID: 11846	

Antibody	Anti- Liver Arginase (ARG1) (Rabbit monoclonal)	Abcam	ab133543	WB (1: 1000)
Antibody	Anti- Arg2 (Rabbit monoclonal)	Abcam	ab264066	WB (1: 1000)
Antibody	anti-Endothelin 1(ET-1)(Mouse monoclonal)	Abcam	ab2786	WB (1: 1000)
Antibody	Anti- VCAM1 (Rabbit monoclonal)	Abcam	ab134047	WB (1: 1000)
Antibody	Anti-ICAM1 (Rabbit monoclonal)	Abcam	ab222736	WB (1: 1000)
Antibody	Anti-AGO2 (Mouse monoclonal)	Proteintech	67934-1-Ig	WB (1: 2000)
Antibody	Anti-beta Actin (Mouse monoclonal)	Abcam	ab8226	WB (1: 2000)
Antibody	Anti-GAPDH (Mouse monoclonal)	Abcam	ab8245	WB (1: 2000)
Recombinant DNA reagent	pCDH-CMV-MCS-EF1-Puro (plasmid)	Addgene	RRID: Addgene_73030	
Recombinant DNA reagent	pLKO.1-GFP-shRNA (plasmid)	Addgene	RRID: Addgene_30323	

Recombinant DNA reagent	psPAX2(plasmid)	Addgene	RRID: Addgene_12260	
Recombinant DNA reagent	pMD2G(plasmid)	Addgene	RRID: Addgene_12259	
Recombinant DNA reagent	pAAV-MCS (plasmid)	Ruipute	Cat#: 2212E4	
Recombinant DNA reagent	pAAV-RC9 (plasmid)	Ruipute	Cat#: 23011	
Recombinant DNA reagent	pHelper (plasmid)	Ruipute	Cat#: 230112	
Recombinant DNA reagent	pAAV-MIR4521 sponge-zsGREEn1-shRNA(plasmid)	Ruipute	Cat#: 230113	
Recombinant DNA reagent	pAAV-Arg1-zsGREEn1-shRNA(plasmid)	Ruipute	Cat#: 2212E1	
Recombinant DNA reagent	pAAV-ARG1-zsGREEn1-shRNA(plasmid)	Ruipute	Cat#: 2212E2	
Recombinant DNA reagent	pAAV-ZsGreen1 (plasmid)	Takara	Cat#: 6231	
Recombinant DNA reagent	pAAV-ZsGreen1-shRNA (plasmid)	youbio	Cat#: VT8093	
Commercial assay or kit	Fluorescent In Situ Hybridization Kit	RIBOBIO	Cat#: C10910	
Commercial assay or kit	RNA pull down kit	GENESEED	Cat#: P0202	

Commercial assay or kit	RNA Immunoprecipitation Kit	GENESEED	Cat#: P0102	
Software, algorithm	ImageJ	National Institutes of Health	https: //imagej.nih.gov/ij/	
Software, algorithm	Prism	GraphPad v8.0	RRID: SCR_002798	

Cell Culture

HUVECs were cultured in high glucose Dulbecco's Modified Eagle Medium (DMEM; Hyclone, USA) supplemented with 10% fetal bovine serum (FBS; Gibco, USA), 100 µg/mL of streptomycin, and 100 U/mL of penicillin. The cells were cultured at 37 °C in a humidified incubator with 95% air and 5% CO_2. To simulate the elevated glucose levels observed in diabetes, we cultured HUVECs in a PAHG medium. The medium contained 25 mM glucose and 250 µM saturated free fatty acid (FFA) palmitate (16:C; Sigma, USA) and was incubated for 24 h. For *in vitro* experiments, HUVECs were subcultured in six-well plates. The cells were then transfected with *MIR4521* mimics, MIR-inhibitor, or MIR-NC and incubated for 48 h. Subsequently, the cells were treated with PAHG for another 24 h. Samples were collected to determine the corresponding indices.

Animal design

Male C57BL/6J mice (6–8 weeks old) were obtained from SLAC Laboratory Animal Co., Ltd. (SLAC ANIMAL, China). The mice were housed at a temperature of 22 ± 1 °C with a 12-h light/dark cycle. To induce diet-induced diabetes, we fed wild type littermates either a standard chow (Control) or a high fat-high sucrose (HFHG) diet, where the diet composition consisted of 60% fat, 20% protein, and 20% carbohydrate (H10060, Hfkbio, China). The dietary regimen was maintained for 14 weeks. Throughout this period, body weight and fasting blood glucose (FBG) levels were

measured on a weekly basis. An FBG level of ≥11.1 mmol/L was used as the criterion

for a successful diabetic model. To study the role of *ARG1* in diabetes-induced

endothelial dysfunction, we established an HFHG-induced diabetic model (DM), and

then pAAV9-*Arg1* shRNA (1×10^{12} vg/mL, 100 μL) was injected into the tail vein.

Subsequently, we observed changes in relevant indicators. To investigate the

regulatory relationship among *Arg1*, *MIR4521*, and *circHMGCS1* in diabetes-induced

endothelial dysfunction, we injected *MIR4521* agomir or pAAV9-*circHMGCS1* via

the tail vein, following the procedure described in **Figure 5-figure supplement 1A**.

The mice were sacrificed after a 12-h fast, and blood and other tissues were collected

and stored at –80 °C until further analysis. All animal experiments were conducted in

strict accordance with the guidelines and laws governing the use and care of

laboratory animals in China (GB/T 35892-2018 and GB/T 35823-2018) and NIH

Guide for the Care and Use of Laboratory Animals. The experimental procedures

involving animals were performed at the Animal Experiment Center of Zhejiang

Chinese Medical University in Hangzhou, China. The animal protocol was conducted

in compliance with the guidelines set forth by the Ethics Committee for Laboratory

Animal Care at Zhejiang Chinese Medical University (Approval No. 2022101345).

Vascular function

After sacrificing the mice, the thoracic aortae and mesenteric arteries were carefully

removed and dissected in an oxygenated ice-cold Krebs solution. The changes in the

isometric tone of the aortic rings and second-order mesenteric arteries were recorded

using a wire myograph (Danish Myo, DK). The arterial segments were stretched to

achieve an optimal baseline tension (3 mN for the aorta) and allowed to equilibrate

for 1 h. Subsequently, the segments were contracted using 60 mmol/L of KCl and

rinsed with Krebs solution. Endothelium-dependent relaxation (EDR) was evaluated

by assessing the concentration-responses to cumulative additions of acetylcholine

(ACh) in noradrenaline (NE, 10^{-3} mol/L) precontracted rings.

Blood pressure measurements

An intelligent non-invasive blood pressure monitor (BP-2010AUL) was used to measure blood pressure in different mouse groups simultaneously. Prior to measurement, the mice were placed in a blood pressure room until reaching a temperature of 25 ± 1 °C. After a 15-min equilibration period, the mice were gently positioned in a mice jacket set at a constant temperature of 40 °C, with their tail inserted into a pulse sensor. Throughout the measurements, a quiet environment was maintained to minimize external disturbances. The instrument automatically recorded the stabilized sensor signal to obtain the systolic blood pressure (SBP).

Serum lipid assays

After 5 weeks of blood collection from mouse orbits and 12 weeks of drug administration, blood samples were obtained from mouse arteries. The collected blood was centrifuged at 1,500 g for 15 min to separate the serum. Commercial kits and an automatic biochemical analyzer (Biobase BK-600) were used to measure triglycerides (TG), total cholesterol (TC), high-density lipoprotein cholesterol (HDL-c), and low-density lipoprotein cholesterol (LDL-c).

Construction of *circHMGCS1* and *ARG1* plasmids and stable transfection using lentivirus

circHMGCS1-overexpressing lentiviruses (pLV-*circHMGCS1*) were obtained from Hangzhou Ruipu Biotechnology Co., Ltd (Guangzhou, China). The *circHMGCS1* sequence [NM_001098272: 43292575-43297268], the splice site AG/GT and ALU elements were inserted into the pCDH-circRNA-GFP vector (upstream ALU: AAAGTGCTGAGATTACAGGCGTGAGCCACCACCCCCGGCCCACTTTTTGT AAAGGTACGTACTAATGACTTTTTTTTTATACTTCAG, downstream ALU: GTAAGAAGCAAGGAAAAGAATTAGGCTCGGCACGGTAGCTCACACCTGT AATCCCAGCA). The restriction enzyme sites selected were EcoRI and NotI. Lentiviruses containing an empty vector and expressing GFP were used as the negative control (pLV-NC) (Liang & Wilusz, 2014). *ARG1*-overexpressing lentiviruses (pLV-*ARG1* OE) were acquired from Genomeditech (China). HUVECs

were cultured in 6-well plates until reaching 60% confluence and then infected with

lentivirus particles in the presence of 5 μg/ml of polybrene at a multiplicity of

infection of 30. The cells were cultured for at least 3 days before further experiments

were conducted. The overexpression efficiency of *circHMGCS1* or *ARG1* was

evaluated using quantitative real-time PCR or western blottings.

Cloning and production of *circHMGCS1*, *MIR4521* sponge, *circHMGCS1* shRNA

and *ARG1* shRNA

The pAAV-*circHMGCS1*-ZsGreen1 vector was constructed by inserting the

full-length circHMGCS1 into the pAAV-circRNA-ZsGreen1 vector. *circHMGCS1*

shRNA vector was constructed by inserting *circHMGCS1* shRNA

(5'-ACAUAGCAACUGAGGGCUUCG-3') into pLKO.1 GFP shRNA plasmid. To

achieve AAV-mediated *ARG1* gene silencing, we obtained a shRNA sequence

targeting *ARG1* from Sigma-Aldrich Mission RNAi (TRCN0000101796). The oligo

sequences

5'GATCCGCCTTTGTTGATGTCCCTAATCTCGAGATTAGGGACATCAACAAA

GGCTTTTTA-3'   and

5'-AGCTTAAAAAGCCTTTGTTGATGTCCCTAATCTCGAG

ATTAGGGACATCAACAAAGGCG-3' were synthesized, annealed, and ligated to

the pAAV-ZsGreen-shRNA shuttle vector, resulting in the construction of

pAAV-shARG1-ZsGreen1. The *MIR4521* sponge cassette was cloned into the

pAAV-ZsGreen1 expression vector to generate pAAV-*MIR4521* sponge-ZsGreen1.

Recombinant AAVs (rAAV9s) were produced by transfecting HEK293T cells plated

at a density of 1×10^7 cells/15 cm plate one day prior to transfection using

polyethyleneimine (Linear PEI, MW 25 kDa, Sigma-Aldrich) as the transfection

reagent. For each 15-cm plate, the following reagents were added: 20 μg of pHelper,

10 μg of pAAV-RC9, and 10 μg of pAAV-circHMGCS1-ZsGreen1,

pAAV-shARG1-ZsGreen1, or pAAV-*MIR4521* sponge-ZsGreen1. These plasmids

were combined with 500 μL of serum-free and antibiotic-free DMEM and 100 μL of

PEI reagent (1 mg/mL, pH 5.0). The DNA-PEI reagent was added drop-wise to the

cells without changing the media on the 15-cm plates. After three days, the cells

were collected and lysed through repeated cycles of freezing and thawing at −80 °C

and 37 °C, respectively. The rAAV9s were purified using the ViraTrap™ Adenovirus

Purification Maxiprep Kit (BW-V1260-02, Biomiga). The virus titer was determined

using qPCR and adjusted to 1×10^{12} viral genomes (vg)/mL in PBS containing 4%

sucrose.

Transfection of miRNA mimic or inhibitor

MIR4521 mimics, miRNA negative control (MIR-NC), *MIR4521* inhibitor, and

miRNA inhibitor negative control (MIR-NC inhibitor) were obtained from Tingske

(China). To investigate the functional role of *MIR4521* in HUVECs, we performed

transfections using Hieff Trans® *in vitro* siRNA/miRNA Transfection Reagent with

MIR4521 mimics, MIR-NC, MIR-NC inhibitor, or MIR-NC inhibitor at a

concentration of 50 nM, following the manufacturer's protocols. After 48 h of

transfection, proteins and RNA were collected for further analysis. The sequences

used in the experiments are listed as follows: miRNA negative control (MIR-NC):

5'-UUCUCCGAACGUGUCACGUTT-3',

5'-ACGUGACACGUUCGGAGAATT-3';

MIR98-5P-mimics: 5'-UGAGGUAGUAAGUUGUAUUGUU-3',

5'-CAAUACAACUUACUACCUCAUU-3'; *MIR3143*-mimics:

5'-UAACAUUGUAAAGCGCUUCUUU-3',

5'-AGAAGCGCUUUACAAUGUUAUU-3'; *MIR181A-2-3P*-mimics:

5'-ACCACUGACCGUUGACUGUACC-3',

5'-UACAGUCAACGGUCAGUGGUUU-3'; *MIR4521*-mimics:

5'-GAGCACAGGACUUCCUUAGCUU-3',

5'-GCUAAGGAAGUCCUGUGCUCAG-3'; *MIR4521*-inhibitor:

5'-CUGAGCACAGGACUUCCUUAGC-3'; miRNA inhibitor negative control

(MIR-NC inhibitor): 5'-CAGUACUUUUGUGUAGUACAA-3'; *MIR4521* agomir:

5'-GAGCACAGGACUUCCUUAGCUU:

5'-GCUAAGGAAGUCCUGUGCUCAG-3'.

NO assays

611   The supernatant fluid from HUVECs treated with the indicated reagents and a tissue
612   homogenate of aortic rings were collected. NO concentrations were determined using
613   the Griess reagent and a total nitric oxide assay kit (Beyotime, China) following the
614   manufacturer's instructions.

ENOS **activity assay**

The cellular *ENOS* activity was assessed as per previously described protocol.
(Leopold et al, 2007). Briefly, Cells were exposed to a PBS buffer (400 μL/well)
containing 1.5 Ci/ml [3H] L-arginine for 15 minutes to initiate *ENOS* activity
measurement. The reaction was stopped using 1 N ice-cold TCA (500 μL/well).
After freeze-thawing in liquid nitrogen, cells were scraped, treated with
water-saturated ether three times, and neutralized with 1.5 ml of 25 mM pH 8.0
HEPES. Dowex AG50WX8 columns (Tris form) were used for elution with a 1 ml
solution of 40 mM pH 5.5 HEPES supplemented with 2 mM EDTA and 2 mM
EGTA. The eluate was collected for [3H] L-citrulline quantification using liquid
scintillation spectroscopy. Additionally, the assessment of ENOS activity in cell
lysates and tissues was performed through the previously outlined
immunoprecipitation approach (Du et al, 2001). Briefly, the samples were divided
into two tubes: one for protein blotting and the other for *ENOS* activity measurement.
The *ENOS* immunocomplexes, linked to protein A-Sepharose beads, were
reconstituted in assay buffer, and *ENOS* activity was gauged by tracking the
transformation of [3H] L-arginine into [3H] L-citrulline. Equal enzyme quantities in
each incubation were validated via Western blotting.

Detection of ROS for tissue and cell

To assess the level of ROS in tissues, we used a red fluorescent reactive oxygen

probe called DHE (dihydroethidium) (Karim et al, 2022; Su et al, 2018). In brief, 10

μL of the DHE probe was combined with 190 μL of tissue homogenate supernatant,

thoroughly mixed, and subsequently incubated in darkness at 37 °C for 30 min.

Following incubation, the samples were promptly imaged under a fluorescent

microscope.

For ROS detection in cells, the treated cells were washed once by PBS, then 20 μM

DHE was added, and incubated at 37°C for 30 min away from light, then washed

three times by PBS and then colorless DMEM medium was added, followed by

fluorescence microscopy for observation.

CircRNA sequencing

Total RNA was extracted from HUVECs, both with and without PAHG stimulation,

utilizing Trizol (Invitrogen, Carlsbad, USA) following the manufacturer's

instructions. The extracted total RNA was then assessed for quality and quantity

using a denaturing agarose gel, Nano Drop, and Agilent 2100 bioanalyzer (Thermo

Fisher Scientific, USA). For the construction of RNA-seq libraries to obtain

sequence information of linear transcripts, 3 μg of total RNA samples underwent

treatment with the epicenter Ribo-ZeroTM Kit (Illumina, San Diego, USA) to

remove rRNA. Linear RNA was subsequently eliminated using RNase R (Epicentre

Technologies, USA). The resulting cleaved RNA fragments were reverse-transcribed

to generate cDNA, which served as the template for the synthesis of U-labeled

second-stranded DNAs. This synthesis step involved the use of E. coli DNA

polymerase I, RNase H, and dUTP, facilitated by the PrimeScript RT regent Kit

(TaKaRa, Japan). Purification of the synthesized second-stranded DNAs was

accomplished using AMPureXP beads. All subsequent steps were conducted in

accordance with the manufacturer's protocols. To assess the quality of the libraries,

an Agilent 2100 Bioanalyzer was employed, and the paired-end sequencing was

performed using an Illumina Hiseq 4000 (LC Bio, China), adhering to the

recommended protocol provided by the vendor.

Treatment of circRNA raw sequencing data

The sequencing data was filtered with Cutadapt by removing reads containing adaptor contamination, poly-N or low quality and undetermined bases, and then sequence quality was verified using FastQC (http://www.bioinformatics.babraham.ac.uk/projects/fastqc/) (Kechin et al, 2017). Bowtie2 and Hisat2 to map reads to the genome of species (Kim et al, 2015; Langmead & Salzberg, 2012). CIRCExplorer2 and CIRI were used to *denovo* assemble the mapped reads to circular RNAs (Kim & Salzberg, 2011; Zhang et al, 2016). Then, back splicing reads were identified in unmapped reads by tophat-fusion. All samples were generated unique circular RNAs. The expression level was calculated according to the junction reads per billion mapped reads (RPB). Differentially expressed circRNAs analysis was performed with log2 (fold change) >1 or log2 (fold change) <-1 and with statistical significance (p value < 0.05) by R package–edgeR (Robinson et al, 2010).

microRNA microarray assay

Total RNA was extracted from HUVECs treated with either PAHG or PBS using TRIzol reagent (Invitrogen, USA). The µParaflo MicroRNA microarray Assay by LC Sciences (LC Sciences, USA) using miRBase version 22.0 was performed. After 3'-extension with a poly(A) tail and ligating an oligonucleotide tag, hybridization was done overnight on a µParaflo microfluidic chip. Cy3 dye facilitated dye staining after RNA hybridization. Fluorescence images were acquired using the GenePix 400B laser scanner (Molecular Devices, USA). Data were analyzed by background subtraction and signal normalization using a LOWESS filter. miRNA expression with a two-tailed p-value of <0.05 was considered significant. Intensities below 500 were considered false positives.

Network analysis of competing endogenous RNA

The miRNA targets of *circHMGCS1* were computationally predicted using three

690 bioinformatic tools: miRanda (http://www. microrna.org/microrna/home.do), whereas

691 the potential target genes of *MIR4521* were envisaged using four distinct databases:

692 TargetScan (http://www.targetscan.org/), GenCard (https://www.genecards.org/),

693 miRWalk (http://mirwalk.umm.uni-heidelberg.de/), and mirDIP

694 (http://ophid.utoronto.ca/mirDIP/).

Isolation of RNA and miRNA for quantitative Real Time-PCR (qRT-PCR)
695
analysis

RNA was extracted utilizing a total RNA extraction kit (Yeason, 19221ES50, China)

and a miRNA extraction kit (Yeason, 19331ES50, China) according to the

corresponding manufacturer's protocol. The RNA concentration was determined at

260 nm using a spectrophotometer. Reverse transcription was conducted using a

reverse transcription kit (TAKARA, Japan) with random primers according to the

manufacturer's protocol to synthesize cDNA and circRNA. Quantitative real-time

PCRs (qRT-PCRs) were conducted on the CFX96 Touch Real-Time PCR Detection

System (Bio-Rad, USA) using the standard protocol provided by the SYBR Green

PCR Kit (Yeason, China). For miRNA expression detection, reverse transcription was

executed, and microRNAs were detected using stem-loop primers procured from

Tsingke (China). The relative gene expression was calculated using the $2^{-\Delta\Delta Ct}$ method.

All qRT-PCR assays were repeated thrice. The primers used for the

qRT-PCR assays are listed as follows: *MIR98-5P*:

5'-CGCGCGTGAGGTAGTAAGTTGT-3',

5'-AGTGCAGGGTCCGAGGTATT-3'; *MIR3143*:

5'-GCGATAACATTGTAAAGCGCTT-3', 5'-AGTGCAGGGTCCGAGGTATT-3';

MIR181A-2-3P: 5'-GCGACCACTGACCGTTGAC-3',

5'-AGTGCAGGGTCCGAGGTATT-3'; *MIR4521*:

5'-CGGCTAAGGAAGTCCTGTGC-3', 5'-CGCAGGGTCCGAGGTATTC-3'; *U6*:

5'-ATTGGAACGATACAGAGAAGATT-3' ,

5'-GGAACGCTTCACGAATTTG-3'; *ET-1*:

5'-GCCTGCCTTTTCTCCCCGTTAAA-3',

5'-CAAGCCACAAACAGCAGAGA-3';   ICAM1:

5'-TGACCGTGAATGTGCTCTCC-3',   5'-TCCCTTTTTGGGCCTGTTGT-3';

ARG1:   5'-TTCTCAAAGGGACAGCCACG-3',

5'-CGCTTGCTTTTCCCACAGAC-3';   *ICAM1*:

5'-ATGCCCAGACATCTGTGTCC-3',   5'-GGGGTCTCTATGCCCAACAA-3';

VCAM1:   5'-AAATCGAGACCACCCCAGAA-3',

5'-AGGAAAAGAGCCTGTGGTGC-3';   *β-actin*:

5'-CTCACCATGGATGATGATATCGC-3',

5'-CACATAGGAATCCTTCTGACCCA-3';   *HMGCS1*:

5'-TCGTGGGACACATATGCAAC-3',   5'-TGGGCATGGATCTTTTTGCAG-3';

hsa_circ_0000992:   5'-TCACACGGGAGAGTGTTACC-3',

5'-GTCCATCGAGAAAAGCTGATGC-3';   *hsa_circ_0004036*:

5'-GGCTCACAAGCAGCCTTTAC-3',   5'-GGTGATACAGGAGCGGGTAG-3';

hsa_circ_0004889:   5'-GTCATATATGGGAAAGAGGGCTT-3',

5'-GGACAAGCCTTCATGGGCTC-3';   *hsa_circ_0006719*:

5'-CCTCCCTCGGTGTTGCCTTC-3',   5'-TCCAGGCCAGGTAGACAGAAC-3';

hsa_circ_0008621:   5'-CAAACATAGCAACTGAGGGCTTC-3',

5'-TAATGCACTGAGGTAGCACTG-3';   *GAPDH*:

5'-GAAAGCCTGCCGGTGACTAA-3', 5'-GCATCACCCGGAGGAGAAAT-3'.

circRNA Validation by PCR

DNA and gDNA templates were PCR amplified using BioRad Mastercyclers

following the manufacturer's protocol. Subsequently, the PCR products were

visualized using a 1% agarose gel stained with GelRed. To validate the PCR results,

we purified the products using the EZ-10 Column DNA Purification Kit (Sangon

Biotech, China), and direct PCR product Sanger sequencing was performed.

circHMGCS1-full   length: 5'-AGCAACTGAGGGCTTCGT-3',

5'-ATGTTTGAATGCACAAGTCCTAC-3';   *circHMGCS1*-divergent   primers:

5′-TTATGGTTCTGGTTTGGCTGC-3′, 5′-TTCAGCGAAGACATCTGGTGC-3′;

circHMGCS1-convergent primers: 5′-ATAGCAACTGAGGGCTTCGTG-3′,

5′-GCGGTCTAATGCACTGAGGT-3′.

Cell nuclear/cytoplasmic fractionation

750 RNA extracts from HUVECs were subjected to nuclear/cytoplasmic isolation using

751 the NE-PER Nuclear and Cytoplasmic Extraction Reagent (Thermo Fisher Scientific,

752 USA) following the manufacturer's protocol.

RNase R treatment

754 For RNase R treatment, approximately 2 μg of total RNAs from HUVECs were

755 incubated with or without 20 U of RNase R (Geneseed, China) at 37°C for 30 min.

756 The resulting RNAs were then purified using an RNA Purification Kit (Qiagen,

757 USA).

Actinomycin D assay

HUVECs were seeded equally in 24-well plates at a density of 5×10^4 cells per well.

Subsequently, the cells were exposed to actinomycin D (2 μg/mL; Abcam, USA) for

different time intervals: 0, 4, 8, 12, and 24 h. After the respective exposure times, the

cells were collected for RNA extraction. The relative RNA levels of circHMGCS1

and HMGCS1 mRNA were analyzed using qRT-PCR and normalized to the values

obtained from the control group (0 h).

RNA immunoprecipitation

RIP assay was conducted using the PureBinding®RNA Immunoprecipitation Kit

(GENESEED, China) as per the manufacturer's guidelines. HUVECs were lysed in

complete RNA immunoprecipitation lysis buffer containing 1% proteinase inhibitor

and 1% RNase inhibitor after being infected with circNC or *circHMGCS1*. The cell

extract was then incubated with magnetic beads conjugated with anti-Argonaute 2

(*AGO2*) or anti-IgG antibody (Abcam, USA) overnight at 4 °C. The beads were

subsequently washed, and the proteins were removed using columns. Eventually, the isolated RNA was extracted using TRIzol Reagent, and the purified RNA was utilized for subsequent qRT-PCR analysis.

Pull-down Assay

A pull-down assay was conducted using the PureBinding®RNA-Protein pull-down Kit (GENESEED, P0201, China) following the manufacturer's protocol. The pull-down assay with biotinylated *MIR4521* or *circHMGCS1* was performed as previously described (Wang et al, 2020). Briefly, HUVECs were transfected with biotinylated RNA or mutants (Tingke, 50 nmol/L) using the Hieff Trans® *in vitro* siRNA/miRNA Transfection Reagent (Yeason, China). Following a 48-h transfection period, the cells were collected, PBS-washed, and incubated for 10 min in a capture buffer on ice. A 10% fraction of the cell lysates was retained as input. The remaining lysates were subsequently exposed to streptavidin magnetic beads for 30 min at 4 °C. After treatment with wash buffer and the RNeasy Mini Kit (Qiagen), the captured RNAs were extracted for subsequent qRT-PCR analysis , while the associated proteins were processed for Western blot analysis.

The sequences are listed as follows: Biotin-*MIR4521*-MUT: 5'-GCAUUCCUUCAGGAGUGCUCAG-3', Biotin-*MIR4521*-WT: 5'-GCUAAGGAAGUCCUGUGCUCAG-3'. Biotin-*circHMGCS1*-WT: 5'-ATGTGTCCCACGAAGCCCTCAGTTGCTATGTTTGAA-3', Biotin-*circHMGCS1*-MUT: 5'-GACGTCGTGTGCGTCGGTGCTAAGCTTCACAGATAC-3'.

Western blot analysis

Western blot was performed following previously described procedures (Xu et al, 2022). Protein extraction from cells or tissues was carried out using protein lysis buffer containing protease and phosphatase inhibitor cocktail (Beyotime, China). The concentrations of the protein lysates were determined using the Beyotime BCA

Protein Assay Kit (Beyotime, China) according to the manufacturer's instructions.

Equal quantities of proteins (20 μg/lane) were loaded onto 8% or 10% SDS-PAGE

gels and subsequently transferred to PVDF membranes (Millipore, USA). The

membranes were then incubated with primary antibodies overnight at 4°C. The

primary antibodies used were as follows: anti-*ARG1* (diluted 1:1,000; Abcam,

ab133543), anti-*ARG2* (diluted 1:1,000; Abcam, ab264066), anti-*ET-1* (diluted

1:1,000; Abcam, ab2786), anti-*VCAM1* (diluted 1:1,000; Abcam, ab134047),

anti-*ICAM1* (diluted 1:1,000; Abcam, ab222736), anti-AGO2(diluted 1:2,000;

proteintech, 67934-1-Ig), anti-*β-ACTIN* (diluted 1:2,000; Abcam, ab8226), and

anti-*GAPDH* (diluted 1:2,000; Abcam, ab8245). After incubation with the

corresponding secondary antibodies for 2 hours at room temperature, the membranes

were washed. Antibody binding was detected using an ECL detection reagent

(Millipore, USA). Digital images were captured using a Gel DocTM XR+ System

with ImageLab software (BioRad, USA). The quantification of the labeled bands

was performed using ImageJ 1.55.

RNA Fluorescent *In Situ* Hybridization (RNA-FISH)

The RNA-FISH assay was conducted in HUVECs. Cy3-labeled *circHMGCS1* probe

and FAM-labeled *MIR4521* probe were custom-designed and synthesized by

RiboBio (China). The signals emanating from the probes were detected using the

Ribo^{TM} FISH Kit (RiboBio) in accordance with the manufacturer's protocol. Briefly,

frozen sections were fixed with 4% paraformaldehyde for 10 minutes, followed by

PBS washing. The fixed sections were treated with PBS containing 0.5% triton

X-100 at 4°C for 5 min, followed by incubation in pre-dehydration buffer at 37°C

for 30 min. Subsequently, probes targeting *circHMGCS1* and *MIR4521* were applied

to the sections. Hybridization was performed in a humid chamber at 37°C overnight.

Post-hybridization washing was initially conducted with 4× saline sodium citrate

containing 0.1% tween-20 at 42°C, followed by further washing with 2× saline

sodium citrate at 42°C to eliminate non-specific and repetitive RNA hybridization.

The slides were counterstained with DAPI (beyotime, China) and examined using an

Olympus FV1200 (Olympus, Japan) confocal microscopy system. The mean

fluorescent intensity (MFI) was determined using Image J software and subjected to

statistical analysis. The probe

sequences   are   detailed   as   follows: Cy3-*circHMGCS1*:

5'-UCCCACGAAGCCCUCAGUUGCUAUG-3',   FAM-*MIR4521*:

5'-UUUGACUCGUGUCCUGAAGGAAUCG-3'.

Luciferase reporter assay

The complete sequence of *circHMGCS1* and a mutant *circHMGCS1* sequence

lacking the miRNA binding site were chemically synthesized. These sequences were

inserted into the Nhel and SaII sites of the pmirGLO vector to generate expression

vectors denoted as *circHMGCS1*-WT and *circHMGCS1*-MUT. Transfection was

carried out using lipofectamine 2000 (Invitrogen, USA) with 50 nM *MIR4521*

mimics, MIR-NC, *MIR4521* inhibitor, and MIR-NC inhibitor. HEK293T cells were

seeded in 96-well plates and cultured until reaching 50%-70% confluence prior to

transfection. After 48-h of transfection, cells were harvested, and luciferase activity

was quantified utilizing the Dual Luciferase Reporter Gene Assay Kit (Yeason,

China). The pmirGLO vector expressing Renilla luciferase was utilized as an

internal control for transfection, while the empty pmirGLO vector served as the

negative control. Firefly luciferase activity of the pmirGLO vector was normalized

with Renilla luciferase activity for comparison. Luciferase reporter assay was also

used to investigate the regulation of *MIR4521* on the expression of its target genes.

For ARG1 and *MIR4521*, either wild-type or mutant *ARG1* 3'UTR fragments (433

bp) were inserted into NheI/SalI restriction sites of pmirGLO. 1 μg plasmids of

ARG1 3'UTR-WT and *ARG1* 3'UTR-MUT, 50 nM *MIR4521* mimics, MIR-NC,

MIR4521 inhibitor and MIR-NC inhibitor were transfected. Luciferase activity was

evaluated 48 hours post-transfection using the Dual Luciferase Reporter Gene Assay

Kit (Yeason).

Histological and morphometric analysis

Aortic tissues were prepared by fixing them in 4% paraformaldehyde in PBS and subsequently embedding them in paraffin. The resulting paraffin-embedded tissues were then sectioned into 4-μm-thick slices. Hematoxylin-eosin (H&E) staining was performed on these sections to visualize the vascular tissue structures and assess the degree of thickness in the thoracic aorta. The average thickness of five randomly selected regions of the thoracic aorta was measured using OlyVIA software (Olympus, Japan).

Statistical analysis

Statistical analyses were conducted using Prism software (GraphPad 8.0). Data are presented as mean $\pm$ standard deviation (SD) unless otherwise stated. Prior to statistical analysis, the normality of data distribution was evaluated using the Shapiro-Wilk test. For comparisons between the two groups, an unpaired two-tailed Student's t-test was used. When comparing more than the two groups, either a one-way or two-way analysis of variance (ANOVA) was conducted, followed by the Bonferroni multiple comparison post hoc test. Group differences were considered statistically significant at a p-value of <0.05. The level of significance was indicated by asterisks, with *, **, and *** denoting p-values lower than 0.05, 0.01, and 0.001, respectively.

Acknowledgments

The authors would like to thank core facilities of medicine in Zhejiang university for technical support.

Funding

This work was supported by the National Natural Science Foundation of China (32172192) and Zhejiang Provincial Key R&D Program of China (2021C02018).

Conflict of interest

The authors have declared that no conflict of interest exists.

Data availability

All data needed to evaluate the conclusions in the paper are present in the paper and/or the Supplementary Materials. Unprocessed original image data have been deposited to Mendely Data and are available at: https://doi.org/10.17632/dxzvhnj7pm.1. Profiling by circRNA and miRNA Array in our work has been deposited in the GEO database as GSE237295 (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE237295) and GSE237597 (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE237597). Additional data related to this paper may be requested from the authors.

References

Arcinas, C., W. Tan, W. Fang, T.P. Desai, D.C.S. Teh, U. Degirmenci, D. Xu, R. Foo,
and L. Sun. 2019. Adipose circular RNAs exhibit dynamic regulation in
obesity and functional role in adipogenesis. *Nature Metabolism.* 1: 688-703.

Arcinas, C., W. Tan, W. Fang, T.P. Desai, D.C.S. Teh, U. Degirmenci, D. Xu, R. Foo,
and L. Sun. 2019. Adipose circular RNAs exhibit dynamic regulation in
obesity and functional role in adipogenesis. *Nat Metab.* 1: 688-703.

Bazzoni, G., and E. Dejana. 2004. Endothelial cell-to-cell junctions: molecular
organization and role in vascular homeostasis. *Physiol Rev.* 84: 869-901.

Caldwell, R.W., P.C. Rodriguez, H.A. Toque, S.P. Narayanan, and R.B. Caldwell.
2018. Arginase: A Multifaceted Enzyme Important in Health and Disease.
Physiol Rev. 98: 641-665.

Cheng, J., Q. Liu, N. Hu, F. Zheng, X. Zhang, Y. Ni, and J. Liu. 2019.
Downregulation of hsa_circ_0068087 ameliorates TLR4/NF-κB/NLRP3
inflammasome-mediated inflammation and endothelial cell dysfunction in
high glucose conditioned by sponging miR-197. *Gene.* 709: 1-7.

Cheng, Z., C. Yu, S. Cui, H. Wang, H. Jin, C. Wang, B. Li, M. Qin, C. Yang, J. He, Q.
Zuo, S. Wang, J. Liu, W. Ye, Y. Lv, F. Zhao, M. Yao, L. Jiang, and W. Qin.
2019. circTP63 functions as a ceRNA to promote lung squamous cell
carcinoma progression by upregulating FOXM1. *Nat Commun.* 10:
3200-3213.

Du, R., N. Wu, Y. Bai, L. Tang, and L. Li. 2022. circMAP3K4 regulates insulin
resistance in trophoblast cells during gestational diabetes mellitus by
modulating the miR-6795-5p/PTPN1 axis. *J Transl Med.* 20: 180-196.

Du, X.L., D. Edelstein, S. Dimmeler, Q. Ju, C. Sui, and M. Brownlee. 2001.
Hyperglycemia inhibits endothelial nitric oxide synthase activity by
posttranslational modification at the Akt site. *J Clin Invest.* 108: 1341-1348.

Eelen, G., P. De Zeeuw, M. Simons, and P. Carmeliet. 2015. Endothelial cell

metabolism in normal and diseased vasculature. *Circulation research.* 116: 1231-1244.

Esteller, M. 2011. Non-coding RNAs in human disease. *Nat Rev Genet.* 12: 861-874.

Gebert, L.F.R., and I.J. Macrae. 2019. Regulation of microRNA function in animals. *Nat Rev Mol Cell Biol.* 20: 21-37.

Gregg, E.W., N. Sattar, and M.K. Ali. 2016. The changing face of diabetes complications. *Lancet Diabetes Endocrinol.* 4: 537-547.

Griendling, K.K., R.M. Touyz, J.L. Zweier, S. Dikalov, W. Chilian, Y.R. Chen, D.G. Harrison, and A. Bhatnagar. 2016. Measurement of Reactive Oxygen Species, Reactive Nitrogen Species, and Redox-Dependent Signaling in the Cardiovascular System: A Scientific Statement From the American Heart Association. *Circ Res.* 119: 39-75.

Hansen, T.B., T.I. Jensen, B.H. Clausen, J.B. Bramsen, B. Finsen, C.K. Damgaard, and J. Kjems. 2013. Natural RNA circles function as efficient microRNA sponges. *Nature.* 495: 384-388.

Hara, M., K. Torisu, K. Tomita, Y. Kawai, K. Tsuruya, T. Nakano, and T. Kitazono. 2020. Arginase 2 is a mediator of ischemia-reperfusion injury in the kidney through regulation of nitrosative stress. *Kidney Int.* 98: 673-685.

Huang, S., X. Li, H. Zheng, X. Si, B. Li, G. Wei, C. Li, Y. Chen, Y. Chen, W. Liao, Y. Liao, and J. Bin. 2019. Loss of Super-Enhancer-Regulated circRNA Nfix Induces Cardiac Regeneration After Myocardial Infarction in Adult Mice. *Circulation.* 139: 2857-2876.

Jiang, B., J. Zhang, X. Sun, C. Yang, G. Cheng, M. Xu, S. Li, and L. Wang. 2022. Circulating exosomal hsa_circRNA_0039480 is highly expressed in gestational diabetes mellitus and may be served as a biomarker for early diagnosis of GDM. *J Transl Med.* 20: 5-18.

Jiang, Q., C. Liu, C.P. Li, S.S. Xu, M.D. Yao, H.M. Ge, Y.N. Sun, X.M. Li, S.J. Zhang, K. Shan, B.H. Liu, J. Yao, C. Zhao, and B. Yan. 2020. Circular RNA-ZNF532 regulates diabetes-induced retinal pericyte degeneration and

948   vascular dysfunction. *J Clin Invest.* 130: 3833-3847.

949   Jung, C., A.T. Gonon, P.O. Sjöquist, J.O. Lundberg, and J. Pernow. 2010. Arginase

950   inhibition mediates cardioprotection during ischaemia-reperfusion.

951   *Cardiovasc Res.* 85: 147-154.

952   Kanyo, Z.F., L.R. Scolnick, D.E. Ash, and D.W. Christianson. 1996. Structure of a

953   unique binuclear manganese cluster in arginase. *Nature.* 383: 554-557.

954   Karim, N., M.R.I. Shishir, Y. Li, O.Y. Zineb, J. Mo, J. Tangpong, and W. Chen. 2022.

955   Pelargonidin-3-O-Glucoside Encapsulated Pectin-Chitosan-Nanoliposomes

956   Recovers Palmitic Acid-Induced Hepatocytes Injury. *Antioxidants (Basel).* 11:

957   1-17.

958   Kechin, A., U. Boyarskikh, A. Kel, and M. Filipenko. 2017. cutPrimers: A New Tool

959   for Accurate Cutting of Primers from Reads of Targeted Next Generation

960   Sequencing. *J Comput Biol.* 24: 1138-1143.

961   Kim, D., B. Langmead, and S.L. Salzberg. 2015. HISAT: a fast spliced aligner with

962   low memory requirements. *Nat Methods.* 12: 357-360.

963   Kim, D., and S.L. Salzberg. 2011. TopHat-Fusion: an algorithm for discovery of

964   novel fusion transcripts. *Genome Biol.* 12: 1-15.

965   Kristensen, L.S., M.S. Andersen, L.V.W. Stagsted, K.K. Ebbesen, T.B. Hansen, and J.

966   Kjems. 2019. The biogenesis, biology and characterization of circular RNAs.

967   *Nat Rev Genet.* 20: 675-691.

968   Langmead, B., and S.L. Salzberg. 2012. Fast gapped-read alignment with Bowtie 2.

969   *Nat Methods.* 9: 357-359.

970   Leopold, J.A., A. Dam, B.A. Maron, A.W. Scribner, R. Liao, D.E. Handy, R.C.

971   Stanton, B. Pitt, and J. Loscalzo. 2007. Aldosterone impairs vascular

972   reactivity by decreasing glucose-6-phosphate dehydrogenase activity. *Nat*

973   *Med.* 13: 189-197.

974   Li, X., X. Sun, and P. Carmeliet. 2019. Hallmarks of Endothelial Cell Metabolism in

975   Health and Disease. *Cell Metab.* 30: 414-433.

976   Liang, D., and J.E. Wilusz. 2014. Short intronic repeat sequences facilitate circular

977   RNA production. *Genes Dev.* 28: 2233-2247.

978   Liang, J., X. Li, J. Xu, G.M. Cai, J.X. Cao, and B. Zhang. 2021. hsa_circ_0072389,

979   hsa_circ_0072386,   hsa_circ_0008621,   hsa_circ_0072387,   and

980   hsa_circ_0072391 aggravate glioma via miR-338-5p/IKBIP. *Aging (Albany*

981   *NY).* 13: 25213-25240.

982   Liu, C., H.M. Ge, B.H. Liu, R. Dong, K. Shan, X. Chen, M.D. Yao, X.M. Li, J. Yao,

983   R.M. Zhou, S.J. Zhang, Q. Jiang, C. Zhao, and B. Yan. 2019. Targeting

984   pericyte-endothelial cell crosstalk by circular RNA-cPWWP2A inhibition

985   aggravates diabetes-induced microvascular dysfunction. *Proc Natl Acad Sci*

986   *U S A.* 116: 7455-7464.

987   Liu, C., M.D. Yao, C.P. Li, K. Shan, H. Yang, J.J. Wang, B. Liu, X.M. Li, J. Yao, Q.

988   Jiang, and B. Yan. 2017. Silencing Of Circular RNA-ZNF609 Ameliorates

989   Vascular Endothelial Dysfunction. *Theranostics.* 7: 2863-2877.

990   Liu, C.X., and L.L. Chen. 2022. Circular RNAs: Characterization, cellular roles, and

991   applications. *Cell.* 185: 2016-2034.

992   Lundberg, J.O., and E. Weitzberg. 2022. Nitric oxide signaling in health and disease.

993   *Cell.* 185: 2853-2878.

994   Ma, C., X. Wang, L. Zhang, X. Zhu, J. Bai, S. He, J. Mei, J. Jiang, X. Guan, X.

995   Zheng, L. Qu, and D. Zhu. 2023. Super Enhancer-Associated Circular

996   RNA-CircKrt4 Regulates Hypoxic Pulmonary Artery Endothelial Cell

997   Dysfunction in Mice. *Arterioscler Thromb Vasc Biol.* 43: 1179-1198.

998   Mahdi, A., O. Kövamees, A. Checa, C.E. Wheelock, M. Von Heijne, M. Alvarsson,

999   and J. Pernow. 2018. Arginase inhibition improves endothelial function in

1000   patients with type 2 diabetes mellitus despite intensive glucose-lowering

1001   therapy. *J Intern Med.* 284: 388-398.

1002   Matsui, M., and D.R. Corey. 2017. Non-coding RNAs as drug targets. *Nat Rev Drug*

1003   *Discov.* 16: 167-179.

1004   Meigs, J.B., F.B. Hu, N. Rifai, and J.E. Manson. 2004. Biomarkers of endothelial

1005   dysfunction and risk of type 2 diabetes mellitus. *Jama.* 291: 1978-1986.

Memczak, S., M. Jens, A. Elefsinioti, F. Torti, J. Krueger, A. Rybak, L. Maier, S.D.
Mackowiak, L.H. Gregersen, M. Munschauer, A. Loewer, U. Ziebold, M.
Landthaler, C. Kocks, F. Le Noble, and N. Rajewsky. 2013. Circular RNAs
are a large class of animal RNAs with regulatory potency. *Nature.* 495:
333-338.

Niemann, B., S. Rohrbach, M.R. Miller, D.E. Newby, V. Fuster, and J.C. Kovacic.
2017. Oxidative Stress and Cardiovascular Risk: Obesity, Diabetes, Smoking,
and Pollution: Part 3 of a 3-Part Series. *J Am Coll Cardiol.* 70: 230-251.

Pan, L., W. Lian, X. Zhang, S. Han, C. Cao, X. Li, and M. Li. 2018. Human circular
RNA☐0054633 regulates high glucose☐induced vascular endothelial cell
dysfunction   through   the   microRNA☐218/roundabout   1   and
microRNA☐218/heme oxygenase☐1 axes. *Int J Mol Med.* 42: 597-606.

Pudlo, M., C. Demougeot, and C. Girard-Thernier. 2017. Arginase Inhibitors: A
Rational Approach Over One Century. *Med Res Rev.* 37: 475-513.

Robinson, M.D., D.J. Mccarthy, and G.K. Smyth. 2010. edgeR: a Bioconductor
package for differential expression analysis of digital gene expression data.
Bioinformatics. 26: 139-179.

Roden, M., and G.I. Shulman. 2019. The integrative biology of type 2 diabetes.
Nature. 576: 51-60.

Romero, M.J., D.H. Platt, H.E. Tawfik, M. Labazi, A.B. El-Remessy, M. Bartoli, R.B.
Caldwell, and R.W. Caldwell. 2008. Diabetes-induced coronary vascular
dysfunction involves increased arginase activity. *Circ Res.* 102: 95-102.

Shan, K., C. Liu, B.H. Liu, X. Chen, R. Dong, X. Liu, Y.Y. Zhang, B. Liu, S.J.
Zhang, J.J. Wang, S.H. Zhang, J.H. Wu, C. Zhao, and B. Yan. 2017. Circular
Noncoding RNA HIPK3 Mediates Retinal Vascular Dysfunction in Diabetes
Mellitus. *Circulation.* 136: 1629-1642.

Shemyakin, A., O. Kövamees, A. Rafnsson, F. Böhm, P. Svenarud, M. Settergren, C.
Jung, and J. Pernow. 2012. Arginase inhibition improves endothelial function
in patients with coronary artery disease and type 2 diabetes mellitus.

Circulation. 126: 2943-2950.

Shen, S., Y. Wu, J. Chen, Z. Xie, K. Huang, G. Wang, Y. Yang, W. Ni, Z. Chen, P. Shi,
Y. Ma, and S. Fan. 2019. CircSERPINE2 protects against osteoarthritis by
targeting miR-1271 and ETS-related gene. *Ann Rheum Dis.* 78: 826-836.

Statello, L., C.J. Guo, L.L. Chen, and M. Huarte. 2021. Gene regulation by long
non-coding RNAs and its biological functions. *Nat Rev Mol Cell Biol.* 22:
96-118.

Stoll, L., A. Rodríguez-Trejo, C. Guay, F. Brozzi, M.B. Bayazit, S. Gattesco, V.
Menoud, J. Sobel, A.C. Marques, M.T. Venø, J.L.S. Esguerra, M. Barghouth,
M. Suleiman, L. Marselli, J. Kjems, L. Eliasson, E. Renström, K. Bouzakri,
M. Pinget, P. Marchetti, and R. Regazzi. 2020. A circular RNA generated
from an intron of the insulin gene controls insulin secretion. *Nat Commun.* 11:
5611-5624.

Su, H., Y. Li, D. Hu, L. Xie, H. Ke, X. Zheng, and W. Chen. 2018. Procyanidin B2
ameliorates free fatty acids-induced hepatic steatosis through regulating
TFEB-mediated lysosomal pathway and redox state. *Free Radic Biol Med.*
126: 269-286.

Su, X., Y. Xu, G.C. Fox, J. Xiang, K.A. Kwakwa, J.L. Davis, J.I. Belle, W.C. Lee,
W.H. Wong, F. Fontana, L.F. Hernandez-Aya, T. Kobayashi, H.M. Tomasson,
J. Su, S.J. Bakewell, S.A. Stewart, C. Egbulefu, P. Karmakar, M.A. Meyer,
D.J. Veis, D.G. Denardo, G.M. Lanza, S. Achilefu, and K.N. Weilbaecher.
2021. Breast cancer-derived GM-CSF regulates arginase 1 in myeloid cells to
promote an immunosuppressive microenvironment. *J Clin Invest.* 131: 1-17.

Tian, Y., J. Xu, X. Du, and X. Fu. 2018. The interplay between noncoding RNAs and
insulin in diabetes. *Cancer Lett.* 419: 53-63.

Treiber, T., N. Treiber, and G. Meister. 2019. Regulation of microRNA biogenesis
and its crosstalk with other cellular pathways. *Nat Rev Mol Cell Biol.* 20:
5-20.

Wang, Y., D. Han, T. Zhou, J. Zhang, C. Liu, F. Cao, and N. Dong. 2020. Melatonin

ameliorates aortic valve calcification via the regulation of circular RNA CircRIC3/miR-204-5p/DPP4 signaling in valvular interstitial cells. *J Pineal Res.* 69: e12666.

Xiao, M.S., and J.E. Wilusz. 2019. An improved method for circular RNA purification using RNase R that efficiently removes linear RNAs containing G-quadruplexes or structured 3' ends. *Nucleic Acids Res.* 47: 8755-8769.

Xu, S., I. Ilyas, P.J. Little, H. Li, D. Kamato, X. Zheng, S. Luo, Z. Li, P. Liu, J. Han, I.C. Harding, E.E. Ebong, S.J. Cameron, A.G. Stewart, and J. Weng. 2021. Endothelial Dysfunction in Atherosclerotic Cardiovascular Diseases and Beyond: From Mechanism to Pharmacotherapies. *Pharmacol Rev.* 73: 924-967.

Xu, Y., Y. Li, J. Li, and W. Chen. 2022. Ethyl carbamate triggers ferroptosis in liver through inhibiting GSH synthesis and suppressing Nrf2 activation. *Redox Biol.* 53: 102349-102352.

Yang, J., and Z. Liu. 2022. Mechanistic Pathogenesis of Endothelial Dysfunction in Diabetic Nephropathy and Retinopathy. *Front Endocrinol (Lausanne).* 13: 816400-816419.

Yeh, C.F., S.H. Cheng, Y.S. Lin, T.P. Shentu, R.T. Huang, J. Zhu, Y.T. Chen, S. Kumar, M.S. Lin, H.L. Kao, P.H. Huang, E. Roselló-Sastre, F. Garcia, H. Jo, Y. Fang, and K.C. Yang. 2022. Targeting mechanosensitive endothelial TXNDC5 to stabilize ENOS and reduce atherosclerosis in vivo. *Sci Adv.* 8: eabl8096.

Yuan, Q., Y. Sun, F. Yang, D. Yan, M. Shen, Z. Jin, L. Zhan, G. Liu, L. Yang, Q. Zhou, Z. Yu, X. Zhou, Y. Yu, Y. Xu, Q. Wu, J. Luo, X. Hu, and C. Zhang. 2023. CircRNA DICAR as a novel endogenous regulator for diabetic cardiomyopathy and diabetic pyroptosis of cardiomyocytes. *Signal Transduct Target Ther.* 8: 99-111.

Yun, J.S., and S.H. Ko. 2021. Current trends in epidemiology of cardiovascular disease and cardiovascular risk management in type 2 diabetes. *Metabolism.*

1093    123: 154838-154848.

1094 Zeng, Z., L. Xia, S. Fan, J. Zheng, J. Qin, X. Fan, Y. Liu, J. Tao, Y. Liu, K. Li, Z.

1095    Ling, Y. Bu, K.A. Martin, J. Hwa, R. Liu, and W.H. Tang. 2021. Circular

1096    RNA CircMAP3K5 Acts as a MicroRNA-22-3p Sponge to Promote

1097    Resolution of Intimal Hyperplasia Via TET2-Mediated Smooth Muscle Cell

1098    Differentiation. *Circulation.* 143: 354-371.

1099 Zhang, X.O., R. Dong, Y. Zhang, J.L. Zhang, Z. Luo, J. Zhang, L.L. Chen, and L.

1100    Yang. 2016. Diverse alternative back-splicing and alternative splicing

1101    landscape of circular RNAs. *Genome Res.* 26: 1277-1287.

1102 Zheng, Y., S.H. Ley, and F.B. Hu. 2018. Global aetiology and epidemiology of type 2

1103    diabetes mellitus and its complications. *Nat Rev Endocrinol.* 14: 88-98.

1104 Zhong, Y., Y. Du, X. Yang, Y. Mo, C. Fan, F. Xiong, D. Ren, X. Ye, C. Li, Y. Wang, F.

1105    Wei, C. Guo, X. Wu, X. Li, Y. Li, G. Li, Z. Zeng, and W. Xiong. 2018.

1106    Circular RNAs function as ceRNAs to regulate and control human cancer

1107    progression. *Mol Cancer.* 17: 79-90.

1108 Zhou, Z., A. Mahdi, Y. Tratsiakovich, S. Zahorán, O. Kövamees, F. Nordin, A.E.

1109    Uribe Gonzalez, M. Alvarsson, C.G. Östenson, D.C. Andersson, U. Hedin, E.

1110    Hermesz, J.O. Lundberg, J. Yang, and J. Pernow. 2018. Erythrocytes From

1111    Patients With Type 2 Diabetes Induce Endothelial Dysfunction Via Arginase I.

1112    *J Am Coll Cardiol.* 72: 769-780.

1113 Zhu, K., X. Hu, H. Chen, F. Li, N. Yin, A.L. Liu, K. Shan, Y.W. Qin, X. Huang, Q.

1114    Chang, G.Z. Xu, and Z. Wang. 2019. Downregulation of circRNA DMNT3B

1115    contributes to diabetic retinal vascular dysfunction through targeting

1116    miR-20b-5p and BAMBI. *EBioMedicine.* 49: 341-353.

1117

 Figure legends

 Figure 1. *circHMGCS1* **Upregulation and its Association with PAHG-induced**

 Endothelial Dysfunction. (A) Heat map illustrating the differential expression of

 circRNAs in HUVECs treated with PAHG for 24 hours (n=3). (B) qRT-PCR

 validation of five circRNAs in HUVECs treated with PAHG compared to normal

 cells, normalized to GAPDH (n=6). (C) PCR validation of *circHMGCS1* presence in

 HUVECs, *GAPDH* was utilized as a negative control (n=6). (D) Alignment of

 circHMGCS1 sequence in CircBase (circular RNA database, upper) in agreement

 with Sanger sequencing results (lower). (E) Schematic representation of

 circularization of exons 2 to 7 of *HMGCS1* (red arrow). (F) RNA-FISH analysis

 detecting *circHMGCS1* expression in HUVECs using Cy3-labeled probes, Nuclei

 were counterstained with DAPI (red represents *circHMGCS1*, blue represents

 nucleus, scale bar = 50 μm, n=4). (G) qRT-PCR quantification of *circHMGCS1* and

 HMGCS1 mRNA levels in the cytoplasm or nucleus of HUVECs. *circHMGCS1* and

 HMGCS1 mRNA levels were normalized to cytoplasmic values (n=4). (H)

 qRT-PCR measure *circHMGCS1* and *HMGCS1* mRNA levels after actinomycin D

 treatment (n=4 at different time points). (I) *circHMGCS1* expression was detected in

 RNase R-treated total RNA by qRT-PCR (n=4). $*p < 0.05$, $**p < 0.01$, $***p <$

 0.001, All significant difference was determined by unpaired two-tailed Student's

 t-test or one-way ANOVA followed by Bonferroni multiple comparison post hoc

 test, error bar indicates SD.

 Figure 1-source data 1. Uncropped and labelled gels for (Figure 1).

 Figure 1-source data 2. Raw unedited gels for (Figure 1).

 Figure 2. *circHMGCS1* **Overexpression Aggravates PAHG-induced Endothelial**

 Dysfunction. (A) HUVECs were transfected with either *circHMGCS1*

 overexpression lentivirus (pLV-*circHMGCS1*) or lentiviral circular RNA negative

control vector (pLV-circNC). After 48 hours of infection, *circHMGCS1* and *HMGCS1* expression levels were assessed by qRT-PCR and normalized to *GAPDH* (n=4). (B and C) Western blot and qRT-PCR were conducted to detect the expressions of adhesion molecules (*VCAM1*, *ICAM1*, and *ET-1*) in HUVECs between pLV-*circHMGCS1* and pLV-circNC groups (n=4). (D) NO content in pLV-circHMGCS1-infected HUVECs after PAHG treatment (n=4). (E) *ENOS* activity in pLV-*circHMGCS1*-infused HUVECs after PAHG treatment (n=4). (F) DHE fluorescence was used to characterize ROS expression in different groups. Scale bar = 50 μm (n=8). (G and H) Relative expression of adhesion molecules (ICAM1, VCAM1 and ET-1) in HUVECs infected with pLV-circNC or pLV-*circHMGCS1* after PAHG treatment were determined by Western blot and qRT-PCR (n=4). *p < 0.05, **p < 0.01, ***p < 0.001, All significant difference was determined by one-way ANOVA followed by Bonferroni multiple comparison post hoc test, error bar indicates SD.

Figure 2-source data 1. Uncropped and labelled gels for (Figure 2).

Figure 2-source data 2. Raw unedited gels for (Figure 2).

Figure 3. *MIR4521* Prevents PAHG-induced Endothelial Dysfunction. (A) Heat map depicting differentially expressed miRNAs in HUVECs with or without PAHG treatment for 24 hours (n=3). (B) Volcano plot illustrating significant changes in miRNAs between control and PAHG-treated groups (Fold-change > 2 or < 0.5, p ≤ 0.01). (C) Schematic illustration showing overlapping target miRNAs of circHMGCS1 predicated by miRanda and sequencing results. (D) The binding sites of *circHMGCS1* were predicted using miRanda and involve *MIR98-5P*, *MIR3143*, *MIR4521*, and *MIR181A-2-3P*. (E) Relative expression of four miRNA candidates in HUVECs treated with PAHG was assessed using qRT-PCR (n=4). (F and G) The regulatory effects of four miRNA mimics on the protein expression levels of adhesion molecules (*ICAM1*, *VCAM1*, and *ET-1*) (n=4). (H) Regulation of NO

content by *MIR4521* mimic or *MIR4521* inhibitor under PAHG treatment (n=4). (I)

Impact of *MIR4521* mimic or *MIR4521* inhibitor on *ENOS* activity during PAHG

treatment. (n=4). (J) The ROS expression in *MIR4521* mimic or *MIR4521* inhibitor

combined with PAHG treatment (n=8). (K) Adhesion molecules (*ICAM1*, *VCAM1*,

and *ET-1*) expression in PAHG-treated HUVECs transfected with *MIR4521* mimics

was determined by Western blot (n=4). (L) Determination of adhesion molecules

(*ICAM1*, *VCAM1*, and *ET-1*) expression via Western blot in PAHG-treated HUVECs

transfected with *MIR4521* inhibitor (n=4). $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, All

significant difference was determined by one-way ANOVA followed by Bonferroni

multiple comparison post hoc test, error bar indicates SD.

Figure 3-source data 1. Uncropped and labelled gels for (Figure 3).

Figure 3-source data 2. Raw unedited gels for (Figure 3).

Figure 4. *circHMGCS1* **Regulates PAHG-induced Endothelial Dysfunction by**
Targeting and Sponging *MIR4521*. (A) Luciferase reporter constructs containing
wild-type (WT) or mutant (MUT) circHMGCS1 were cotransfected with *MIR4521*
mimics, MIR-NC, *MIR4521* inhibitor, or MIR-NC inhibitor in HEK293T cells
(n=4). (B) Immunoprecipitation shows *AGO2*-mediated binding of *circHMGCS1*
and *MIR4521* (n=4). (C) Biotin-coupled *MIR4521* or its mutant probe was employed
for *circHMGCS1* pull-down, and captured *circHMGCS1* level was quantified by
qRT-PCR (n=4). (D) RNA-FISH showing the colocalization of *circHMGCS1* and
MIR4521 in HUVECs (red represents *circHMGCS1*, green represents *MIR4521*, blue
represents nucleus, scale bar = 50 μm, n=4). (E and F) *circHMGCS1* exhibited no
impact on NO content and *ENOS* activity in the presence of *MIR4521* sponge (n=4).
(G) *circHMGCS1* demonstrated no influence on ROS expression with *MIR4521*
sponge (n=8). (H-J) *circHMGCS1* had no effect on the expression of adhesion
molecules (*ICAM1*, *VCAM1*, and *ET-1*) in the presence of *MIR4521* sponge which
were determined by Western blot and qRT-PCR (n=4). (K and L) *MIR4521*

attenuated the reduction of NO content and *ENOS* activity induced by *circHMGCS1*

(n=4). (M) *MIR4521* inhibited the increase of ROS expression caused by

circHMGCS1 (n=8). (N-P) *MIR4521* attenuated the increased expression of adhesion

molecules (*ICAM1*, *VCAM1* and *ET-1*) induced by *circHMGCS1* which were

determined by Western blot and qRT-PCR, respectively (n=4). *p < 0.05, **p <

0.01, ns means no significant. All significant difference was determined by one-way

ANOVA followed by Bonferroni multiple comparison post hoc test, error bar

indicates SD.

Figure 4-source data 1. Uncropped and labelled gels for (Figure 4).

Figure 4-source data 2. Raw unedited gels for (Figure 4).

1211

Figure 5. AAV9-mediated *circHMGCS1* Overexpression Attenuates the

Protective Effect of *MIR4521* against Diabetes-induced VED. (A) Changes in

body weight of mice from 0 to 14 weeks (n=8, ***p<0.001 DM versus control,

#p<0.05 DM + agomir MIR4521 versus DM). (B) Alterations in fasting blood

glucose in mice from 0 to 14 weeks (n=8, ***p<0.001 DM versus control, #p<0.05

DM + agomir *MIR4521* versus DM). (C and D) ITT and OGTT were performed in

the fasted mice (n=8, *p<0.05 DM versus control, #p<0.05 DM + agomir *MIR4521*

versus DM). (E) Relaxation responses of aortic rings from control, *MIR4521* agomir,

or *MIR4521* agomir+AAV9 *circHMGCS1* mice on DM diet (n=6). (F) SBP

measured via tail-cuff method in different groups (n=8). (G) Hematoxylin and eosin

staining (H&E) was performed on serial cross-sections of thoracic aortas from

differently treated mice to assess vessel wall thickness. Scale bar: 100 μm (n=6). (H

and I) NO content and *Enos* activity in thoracic aorta after 6 weeks of DM diet with

MIR4521 agomir or *MIR4521* agomir + *circHMGCS1* treatment (n=6). (J and K)

Detection of ROS expression in thoracic aorta using DHE after *MIR4521* agomir or

MIR4521 agomir+AAV9 *circHMGCS1* injection (red represents ROS, green

represents GFP, blue represents nucleus, scale bar = 100 μm, n=6). (L) Expression

levels of adhesion molecules (*Icam1*, *Vcam1*, and *Et-1*) in thoracic aorta after 6 weeks of DM diet with *MIR4521* agomir or *MIR4521* agomir + *circHMGCS1* treatment (n=6). *p < 0.05, **p < 0.01, ***p < 0.001. All significant difference was determined by one-way ANOVA followed by Bonferroni multiple comparison post hoc test, error bar indicates SD.

Figure 5-source data 1. Uncropped and labelled gels for (Figure 5).

Figure 5-source data 2. Raw unedited gels for (Figure 5).

Figure 6. *circHMGCS1* Functions as a *MIR4521* Sponge in HUVECs to Modulate *ARG1* Expression. (A) Schematic depiction illustrating the intersection of predicted target genes of *MIR4521* from TargetScan, miRWalk, mirDIP, and GeneCard. (B) *Arg1* expression in thoracic aorta of DM by western blot (n=6). (C) HEK293T cells were cotransfected with *MIR4521* mimics, MIR-NC, *MIR4521* inhibitor, or MIR-NC inhibitor, along with luciferase reporter constructs containing WT or MUT 3′-untranslated region of *ARG1* (n=4). (D and E) Modulatory effect of *MIR4521* mimic on NO content and *ENOS* activity under ARG1 overexpression (n=4). (F) Regulatory impact of *MIR4521* mimic on ROS content under *ARG1* overexpression (n=4). (G and H) *ARG1*-overexpressing HUVECs were generated using lentivirus and transfected with *MIR4521* mimics for 24h to evaluate adhesion molecule expression (*ICAM1*, *VCAM1*, and *ET-1*) by Western blot and qRT-PCR (n=4). (I-K) *ARG1* expression was significantly decreased by *MIR4521* mimics and increased by *MIR4521* inhibitor, as determined by qRT-PCR and Western blot (n=4). (L and M) *ARG1* expression was significantly increased by *circHMGCS1* overexpression, as determined by Western blot and qRT-PCR (n≥4). (N and O) *circHMGCS1*-overexpressed HUVECs, created with lentivirus and transfected with *MIR4521* mimics for 48h, were examined for *ARG1* expression by Western blot and qRT-PCR (n=4). (P and Q) *ARG1* expression was significantly reduced by *circHMGCS1* shRNA, as determined by Western blot and qRT-PCR (n=3). (R)

AAV9 *MIR4521* sponge was used to inhibit the *MIR4521* expression in HUVECs,

followed by *circHMGCS1* transfection, and then treated with PAHG for 24 hours to

evaluate the expression of *ARG1* by Western blot (n=4). (S) Detection of *Arg1*

expression in the thoracic aorta of DM treated with AAV9 *MIR4521* agomir

combined with AAV9 *circHMGCS1* by Western blot (n=4). *$p < 0.05$, **$p < 0.01$,

ns means no significant. All significant difference was determined by one-way

ANOVA followed by Bonferroni multiple comparison post hoc test, error bar

indicates SD.

Figure 6-source data 1. Uncropped and labelled gels for (Figure 6).

Figure 6-source data 2. Raw unedited gels for (Figure 6).

1267

Figure 7. *ARG1* is Inseparable from *circHMGCS1* and *MIR4521* Regulating

Diabetes-Induced VED. (A and B) *circHMGCS1* and *MIR4521* had no effect on

NO content and *ENOS* activity expression in the absence of *ARG1* (n=4). (C) ROS

expression remained unaffected by *circHMGCS1* and *MIR4521* in the absence of

ARG1 (n=4). (D and E) Relative expression of adhesion molecules (*ICAM1*,

VCAM1, and *ET-1*) remained unchanged in the absence of *ARG1*, despite the

presence of *circHMGCS1* and *MIR4521*, as determined by qRT-PCR and Western

blot (n=4). (F) Changes in mice body weight over the experimental period (n=8,

**$p<0.01$, DM versus control, #$p<0.05$, DM + AAV9 *ARG1* shRNA versus DM).

(G) Fasting blood glucose levels in mice over time (n=8, **$p<0.01$, DM versus

control, #$p<0.05$, DM + AAV9 *ARG*1 shRNA versus DM). (H and I) Blood glucose

levels measured at week 13 (ITT) and week 14 (OGTT) (n=8, **$p<0.01$ DM versus

control, #$p<0.05$, DM + AAV9 *ARG1* shRNA versus DM). (J)

Endothelium-dependent relaxations in aortic rings from different groups (n=8). (K)

H&E performed on serial cross-sections of thoracic aortas from differently treated

mice to evaluate vessel wall thickness. Scale bar = 100 μm (n=6). (L) SBP measured

by the tail-cuff method in different groups (n=8). (M and N) NO content and *Enos*

activity expression in thoracic aort (n=6). (O) Relative expression of adhesion molecules (*Icam1*, *Vcam1*, and *Et-1*) in thoracic aorta assessed by Western blot (n=6). (P and Q) ROS expression in the thoracic aorta using the DHE probe (Red represents ROS, Green represents GFP, Blue represents nucleus, scale bar: 100 μm, n=6). $p < 0.05$, **$p < 0.01$, ns means no significant. All significant difference was determined by one-way ANOVA followed by Bonferroni multiple comparison post hoc test, error bar indicates SD.

Figure 7-source data 1. Uncropped and labelled gels for (Figure 7).

Figure 7-source data 2. Raw unedited gels for (Figure 7).

Figure Supplements

Figure 1-figure supplement 1. Distinct expression profiles of circRNAs in endothelial dysfunction of HUVECs. HUVECs exposed to palmitic acid and high glucose (PAHG) treatment for 24 hours were compared with untreated Control. (**A**) Alterations in NO content between the control and PAHG-treated groups (n=4). (**B**) Changes in *ENOS* activity between the control and PAHG-treated groups (n=4). (**C**) Relative expression levels of adhesion molecules (*ICAM1*, *VCAM1*, and *ET-1*) in HUVECs were assessed by Western blotting between the control and PAHG-treated groups (n=4). (**D**) Volcano plot illustrating significantly altered circRNAs between the control and PAHG-treated groups. Fold-change > 2 or < 0.5, $p \leq 0.001$. (**E**) Number of identified circRNAs in Control groups and PAHG-treated groups (n=3). (**F**) PCR amplification confirming the full-length *circHMGCS1* (n=3). (**G**) Sanger sequencing validation of the full-length *circHMGCS1* sequence (n=3). (**H**) Comparison results of *circHMGCS1* sequences in Sanger sequencing and circbase database. $*p < 0.05$, $**p < 0.01$, All significant difference was determined by unpaired two-tailed Student's t-test, error bar indicates SD.

Figure 1-figure supplement 1-source data 1. Uncropped and labelled gels for (Figure 1-figure supplement 1).

Figure 1-figure supplement 1-source data 2. Raw unedited gels for (Figure 1-figure supplement 1).

Figure 2-figure supplement 1. A sketch map of *circHMGCS1* plasmid construction.

Figure 3-figure supplement 1. *MIR4521* inhibition aggravates diabetes-induced endothelial dysfunction. (**A**) Relative expression of *MIR98-5P*, *MIR3143*, *MIR181A-2-3P* and *MIR4521* were detected in HUVECs after corresponding

miRNA mimics treatment by qRT-PCR (n=4). (**B** and **C**) Relative expression level of *MIR4521* was determined by qRT-PCR in HUVECs transfected with *MIR4521* mimics or *MIR4521* inhibitor (n=4). (**D**) *MIR4521* mimic inhibited the expression of adhesion molecules (*ICAM1*, *VCAM1*, and *ET-1*) in PAHG-treated HUVEC as detected by qRT-PCR (n=4). (**E**) *MIR4521* inhibitor exacerbated the expression of adhesion molecules (*ICAM1*, *VCAM1*, and *ET-1*) in PAHG-treated HUVEC as detected by qRT-PCR (n=4). *$p < 0.05$, **$p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$ All significant difference was determined by one-way ANOVA followed by Bonferroni multiple comparison post hoc test, error bar indicates SD.

Figure 4-figure supplement 1. Verification of the binding site of *MIR4521* and *circHMGCS1*. HEK293T cells were used to package lentivirus overexpressing *circHMGCS1*, and then infected HUVECs cells, (**A**) Expression level of *MIR4521* was assessed by qRT-PCR in pLV-*circHMGCS1*-transfected HEK293T cells (n=4). (**B**) The expression level of *MIR4521* was assessed by qRT-PCR in HUVEC cells transfected with *circHMGCS1* shRNA (n=3). (**C**) Schematic illustration of the predicted binding sites of *MIR4521* on the *circHMGCS1* transcript in human cells. (**D**) Biotin-coupled *circHMGCS1* or its mutant probe was employed for *MIR4521* pull-down assay, and captured *MIR4521* level was quantified by qRT-PCR (n=4). (**E**) Western blot analysis of *AGO2* protein expression captured by biotin-coupled *circHMGCS1*. **$p < 0.01$, ***$p < 0.001$, significant difference was determined by unpaired two-tailed Student's t-test, error bar indicates SD.

Figure 4-figure supplement 1-source data 1. Uncropped and labelled gels for (Figure 4-figure supplement 1).

Figure 4-figure supplement 1-source data 2. Raw unedited gels for (Figure 4-figure supplement 1).

Figure 5-figure supplement 1. *circHMGCS1* counteracts the protective role of

***MIR4521* in diabetes-induced VED.** (**A**) Timeline illustrating the HFHG feeding,

AAV9 *circHMGCS1*, and *MIR4521* agomir treatment schedule. (**B** and **C**)

Quantification of the area under the curve (AUC) for glucose and insulin tests (n=8).

(**D**) Triglyceride (TG), (**E**) Total cholesterol (T-CHO), (**F**) High-density lipoprotein

cholesterol (HDL-C), and (**G**) Low-density lipoprotein cholesterol (LDL-C) levels in

the serum were measured using Automatic Biochemical Analyzer (n=6). (**H**) Effects

on endothelium-independent vasorelaxation to SNP (n=6). *p < 0.05, **p < 0.01. All

significant difference was determined by one-way ANOVA followed by Bonferroni

multiple comparison post hoc test, error bar indicates SD.

1358

Figure 6-figure supplement 1. *MIR4521* modulates the expression of *ARG1*. (**A**)

Schematic representation of the predicted binding sites and mutated sites for *MIR4521*

on *ARG1* transcripts in both human cells and mice. (**B**) Western blot analysis assessing

ARG1 and *ARG2* expression in HUVECs after PAHG treatment (n=3). (**C**)

Sequencing results displaying the mutation site in the dual luciferase assay. *p < 0.05,

**p < 0.01, ns means no significant. All significant difference was determined by

one-way ANOVA followed by Bonferroni multiple comparison post hoc test, error

bar indicates SD.

Figure 6-figure supplement 1-source data 1. Uncropped and labelled gels for

(Figure 6-figure supplement 1).

Figure 6-figure supplement 1-source data 2. Raw unedited gels for (Figure

6-figure supplement 1).

1371

Figure 7-figure supplement 1. Loss of functional regulation of *circHMGCS1* and

***MIR4521* on HUVECs in the absence of *ARG1* *in vivo*.** (**A**) Schematic outlining the

timeline of HFHG feeding, AAV9 *circHMGCS1* and *MIR4521* agomir treatment,

AAV9 *ARG1* shRNA treatment, and experiment conclusion. (**B** and **C**) Blood glucose

levels measured at week 13 (ITT) and week 14 (OGTT), the corresponding AUC for

glucose or insulin tests was calculated (n=8, $**p<0.01$, DM versus control, $\#p<0.05$,

DM + AAV9 *ARG1* shRNA versus HFHG). (**D**) TG, (**E**) T-CHO, (**F**) HDL-C, and (**G**)

LDL-C levels in the serum of different treated groups (n=6). (**H**) Assessment of

endothelium-independent vasorelaxation to SNP (n=6). $*p < 0.05$, $**p < 0.01$, $*** p <$

0.001, ns means no significant. All significant difference was determined by one-way

ANOVA followed by Bonferroni multiple comparison post hoc test, error bar

indicates SD.