Strategies for Genetic Manipulation of the Halotolerant Black Yeast *H. werneckii*: Ectopic DNA Integration and Marker-free CRISPR/Cas9 Transformation

Yainitza Hernandez-Rodriguez[1], A. Makenzie Bullard[2,3], Rebecca J. Busch[4], and José M. Vargas-Muñiz[4,5.]

[1] Department of Biological Sciences, Florida Gulf Coast University, Ft. Myers, FL

[2] Molecular Biology, Microbiology, and Biochemistry Program, School of Biological Science, Southern Illinois University, Carbondale, IL

[3] Microbiology Program, School of Biological Science, Southern Illinois University, Carbondale, IL

[4] Department of Biological Sciences, Virginia Tech, Blacksburg, VA

[5] Early Career Whitman Fellow, Marine Biological Laboratory, Woods Hole, MA

Running title: *H. werneckii* genetic transformation

Word Count: 1876 (text), 186 (abstract)

[#] Corresponding Author:

José Vargas-Muñiz
Department of Biological Sciences
Steger Hall
1015 Life Science Cir.
Virginia Tech, Blacksburg, VA 24061
E-mail: j.vargas.muniz@vt.edu

Keywords: *Hortaea werneckii,* transformation, morphology, ectopic integration, CRISPR/Cas9

Abstract

Hortaea werneckii is a halotolerant black yeast commonly found in hypersaline environments. This yeast is also the causative agent of tinea nigra, a superficial mycosis of the palm of the hand and soles of the feet of humans. In addition to their remarkable halotolerance, this black yeast exhibits an unconventional cell division cycle. *H. werneckii* alternates between fission and budding cell division. This cell division cycle is regulated by cell density and the salt concentration in their media. *H. werneckii* have been extensively studied due to their ability to grow in high salinity environments and unconventional cell biology. Nonetheless, deciphering the underlying mechanisms behind these remarkable phenotypes has been limited due to the lack of genetic tools available. Here, we report a new ectopic integration protocol for *H. werneckii* using PEG-CaCl$_2$ mediated protoplast transformation. This approach relies on a selectable marker (hygromycin B) to select for successful integration of the genetic construct. The same construct was used to express cytosolic green fluorescent protein. Finally, we also developed a marker-free CRISPR/Cas9 protocol for targeted gene deletion. These transformation strategies will allow testing hypotheses related to *H. werneckii* cell biology and physiology.

Introduction

Hortaea werneckii (*Dothideales, Ascomycota*) is a black yeast of particular interest due to its ability to grow in high salinity (1). This black yeast is commonly isolated from hypersaline environments, including seawater and solar salterns, and can grow in saturated sodium chloride solution. Not only does this yeast display a remarkable halotolerance, but it also exhibits an unconventional cell division cycle (2). *H. werneckii* first grows in a pattern similar to that of fission yeast. After the first septation, it switches to budding from the poles. This division pattern and cell morphology depend on the growth environment and cell density (3, 4). *H. werneckii* is also of clinical interest due to its ability to cause a superficial mycosis of the hands and feet known as tinea nigra (5– 7). Tinea nigra frequently occurs in countries located in the tropics (6). On rare occasions, *H. werneckii* can cause a systemic infection known as disseminated phaeohyphomycosis (7). *H. werneckii* isolates exhibit great phenotypic diversity, including drug susceptibility and pathogenicity variation (3, 8, 9). This phenotypic diversity also correlates with the genetic diversity of *H. werneckii*. *H. werneckii* isolates are intraspecific hybrids of two divergent isolates, contributing to their genetic and phenotypic diversity (10, 11). For these reasons, *H. werneckii* is an emerging model for understanding eukaryotic adaptation to hypersaline environments and how hybridization events contribute to fungal adaptation to extreme environments (9).

Most of our understanding of *H. werneckii* biology and adaptation to hypersaline conditions have been derived from genomics, metabolic, and physiological analyses (1, 9, 18–22, 10–17). However, the lack of reliable genetic tools has limited the ability to

test mechanistic hypotheses related to *H. werneckii* physiology and cell biology.

Different methods exist to genetically transform fungi, including protoplast-mediated,

Agrobacterium-mediate, electroporation, and biolistic (23, 24). These strategies require

selectable markers, usually a drug-resistance gene or nutritional gene, to isolate cells

that contain the desired construct. Recently, *in vitro* assembled CRISPR/Cas9

transformation systems have been utilized for highly efficient genetic manipulation of

fungi (25–30). The CRISPR/Cas9 system has been adapted as a "marker-free" system

to perform genome editing without needing a selectable marker (27).

Here, we adapted a protoplast-mediated transformation protocol to integrate a plasmid

into the genome of *H. werneckii* ectopically. This ectopic integration strategy can be

used for expressing green fluorescent protein (GFP) in the cytoplasm of *H. werneckii.*

We also develop a marker-free CRISPR/Cas9 protocol for knocking out genes in *H.*

werneckii. These new strategies will further our understanding of the biology of this

remarkable yeast.

Materials and Methods

Strains, media, culture conditions

H. werneckii EXF-2000 was use for these studies. Yeast cells were grown in Glucose

Minimal Media (GMM [Dextrose 10g/L, Trace Elements, Salt Solution]). Cells were

grown at 30˚C for 5 days unless otherwise specified.

Protoplast generation

Hortaea werneckii cells were inoculated in GMM agar plates and incubated at 37˚C for

5-7 days. Cells were harvested and washed once with osmotic media (1.2 M

magnesium sulfate, 10 mM sodium phosphate buffer, pH 5.8). Digest the *H. werneckii*

cell wall using osmotic media with 200 mg of Vinotaste for 4-8 hours at 30°C and

shaking at 75 rpm. After digestion, two tubes were combined and overlayed with

trapping buffer (0.6M sorbitol, 0.1 M Tris-HCl pH 7). Tubes then were centrifuged for 15

minutes at 3,500 rpm at 4°C. The cloudy layer (protoplast) at the interface of the

osmotic media and trapping buffer was carefully moved into a new 15 mL conical and

washed once with ice-cold STC (1.2M sorbitol, 10 mM $CaCl_2$, 10 mM Tris-HCl pH 7.5).

Protoplasts were resuspended with 1 mL of STC.

PEG-CaCl₂-mediated transformation of pUCGH plasmid

200 µL of protoplast was transferred into a 1.5 mL and incubated with 5-15 µg of the

pUCGH plasmid for 50 minutes to 1 hour. 1.25 mL of PEG-$CaCl_2$ was added to the

protoplast and incubated for 20 minutes. STC was added to protoplast up to 3 mL, and

300 µL were plated to SMM agar plates. Protoplasts were allowed to recover for 24

hours at room temperature. After 24 hours, the protoplast plates were overlayed with

SMM top agar with 450 µg/ml of hygromycin B. Colonies appeared between 7-15 days

of incubation. Individual colonies were transferred into fresh GMM agar with hygromycin

B (150 µg/mL) and incubated at 30°C.

Imaging EXF-2000

A wet mount of each strain was made by smearing *H. werneckii* cells in 20 µl of distilled

water. Cells were then imaged using a widefield microscope (Leica DMi8) using a 100 X

oil apochromat objective, and images were captured using a Leica K5 microscope

camera.

Assembly of CRISPR/Cas9 RNPs

5 µL of one crRNA, tracrRNA, and nuclease-free duplex buffer (Integrated DNA

Technology, IDT) were mixed and heated at 95°C for 5 minutes. Then, it was cool-down

at room temperature to assemble the gRNA. 1.5 µL of each 33 µM gRNA, 1.5 µL of

Cas9 nuclease (1µg/µL) (Integrated DNA Technology, IDT), and 22 µl of Cas9 working

buffer were mixed and incubated for 5 minutes at room temperature. CRISPR/Cas9

complex was osmotically stabilized by adding an equal volume of 2X STC.

Marker-free CRISPR/Cas9-mediated transformation

Osmotically stabilized CRISPR/Cas9 complex, 70µL of protoplast, and 200 µl of PEG-

$CaCl_2$ were gently mixed in a 50 ml conical tube. The protoplast mixture was incubated

on ice for at least 30 minutes. 1 ml of PEG-$CaCl_2$ was added and incubated for 15

minutes at room temperature. 325 µl of the protoplast were inoculated into 3 large SMM

agar plates and incubated overnight at room temperature. Then, plates were incubated

at 30°C until colonies started appearing on the plates.

Results

Polyethylene glycol (PEG) calcium chloride protoplast-mediated transformation

allows for ectopic integration of the pUCGH plasmid.

H. werneckii susceptibility to drugs is determined by the environment (3, 4). We

discovered that *H. werneckii* is susceptible to hygromycin B when grown on glucose

minimum media, the same media commonly used to grow *Aspergillus fumigatus* (28).

Based on this, we decided to use hygromycin B as our selectable marker and used a

sorbitol-stabilized glucose minimum media (SMM) to perform our transformations. We

consistently obtained protoplast after 4-8 hours of digesting the cell wall using vinotaste

(Novozymes). We then transformed the protoplast using a pUCGH vector, which

contains the hygromycin B resistant gene (*hph*) under the control of *Aspergillus nidulans*

gpdA promoter and eGFP under the control of the *Aspergillus oryzae* TEF1 promoter

(31). We used a high quantity of plasmid to induce ectopic integration (5-15µg) of the

plasmid. Protoplasts were allowed to recover for 24 hours before overlaying them with

SMM top agar containing hygromycin B (450µg/ml). Hygromycin B-resistant colonies

emerged after 7 days of incubation at 30°C (Fig 1A). Colonies were picked and streaked

into a small petri dish containing GMM supplemented with hygromycin B (150µg/ml).

Transformants expressed eGFP even after 5 passages in GMM+hygromycin B (Fig 1B).

Thus, this approach can be utilized for ectopic integration of markers for cell biology

studies.

CRISPR/Cas9 marker-free transformation.

More targeted genetic approaches are needed to further our understanding of *H.*

werneckii biology. Similar to other ascomycetes, adapting the PEG-CaCl2 mediated

transformation protocol has proven challenging for targeted gene manipulation due to

the low homologous recombination rate (32). This challenge is further exacerbated due

to the EXF-2000 strain having a significant portion of its genome duplicated due to an

intraspecific hybridization event (10). Due to this challenge, we decided to adapt a

CRISPR/Cas9 marker-free approach and target the *H. werneckii* melanin synthesis

pathway to have a visual phenotype we could screen (27). We targeted a conserved

region between *H. werneckii's* two *alb1* alleles: *alb1-1* (BTJ68_00107) and *alb1-2*

(BTJ68_01291) using 2 gRNAs (Table 1). Using this approach, approximately 3-5

colonies exhibited an albino phenotype (Figure 2) per plate. These albino colonies were picked and streaked into GMM plates. The albino mutants could still not produce melanin after 5 passages, indicating that we had a stable deletion of the *alb1* alleles. These albino mutants retain the same cell morphology as the wild-type when grown in GMM (Fig 2B), indicating that melanin does not impact cell morphology when grown in GMM Agar.

Discussion

Hortaea werneckii has a unique cell cycle division and can grow at high salt concentrations (3, 4, 33). Due to most isolates being intraspecific hybrids with significant phenotypic variation, *H. werneckii* can potentially become a model for understanding how hybridization leads to novel phenotypes (3, 9). However, the lack of genetic tools has limited the ability to study this organism. Here, we develop two different genetic transformation approaches to further our studies of this unconventional yeast. First, we have developed a plasmid-based ectopic integration protocol. This system can be used to express cellular markers for live cell imaging. For example, we could use it to insert a histone-GFP plasmid and monitor nuclear dynamics. Second, we develop a marker-free CRISPR/Cas9 protocol to target specific genes. One caveat is that our current CRISPR/Cas9 protocol is still largely ineffective and would not be practical for genes that would not have a visual readout. On average, we got 3-5 albino colonies, while other organisms have a 50% to 90% success rate (27–29). However, these organisms were haploid, and only one copy of the gene needed to be mutated, not the 2 copies in *H. werneckii.* It is still possible that we utilized too many protoplasts, and the protoplast

number will need to be titrated to increase efficiency (27). Lastly, as there is a marker

system and CRISPR/Ca9 can overcome the homologous recombination rate, we could

combine both methods to develop a marker-based CRISPR/Cas9 approach and select

those mutants with successful integration of our cassette of interest (28). At the moment

of this publication, only one marker has been developed (hygromycin) for *H. werneckii,*

but other drugs can be explored to expand the available range of selectable markers.

Acknowledgments

The authors thank Dr. William Steinbach and Dr. Praveen Juvaadi for sharing the

pUCGH plasmid. We also want to thank Dr. Rachel Whitaker, Dr. Scott Dawson, and

the Marine Biological Laboratory's (MBL) Microbial Diversity course for allowing us to

use their microscope and resources. The author wants to thank Norman van Rhijn

(Manchester Fungal Infection Group) and Robb Cramer's Lab (Dartmouth) for sharing

their CRISPR/Cas9 protocols that were the basis for developing our protocol. This

project was supported through the L. & A. Colwin Summer Research Fellowship

awarded to JVM. JVM wants to thank William Wolf (Virginia Tech), and the Vargas-

Muñiz Lab for critically reading this manuscript.

Figure Legends

Figure 1. Integration of the pUCGH plasmid in *H. werneckii*. (A) *H. werneckii* EXF-

200 is susceptible to hygromycin in GMM media, and hygromycin allows for the

selection of ectopic integration of the pUCGH plasmid. Plates were streaked into the

respective media, incubated at 37˚C, and photographed after 5 days. **(B)** *H. werneckii*

with integrated pUCGH expresses eGFP under the *A. oryzae* tef1 promoter.

Micrographs were obtained using a 100X objective. Scale bar, 20 μm.

Figure 2. Generation of albino mutants using CRISPR/Cas9. (A) The *alb1Δ/Δ* fails to

melanize after prolonged incubation in GMM. Plates were streaked and incubated at

30˚C for 5 days before being photographed. **(B)** *alb1Δ/Δ* have similar morphology as the

parent EXF-2000 strain when grown in GMM agar. Micrographs were obtained using a

100X objective. Scale bar, 20 μm.

Table 1: crRNAs used in this study

crRNA	Target	Sequence
alb1-gRNA-5-1	5' of *alb1*	AAGCGCUUGACGGUGGUUGGCGG
alb1-gRNA-5-2	*5' of alb1*	AGCGUUGCGCGAAAGCGCUGCGG

References

1.   Petrovič U, Gunde-Cimerman N, Plemenitaš A. 2002. Cellular responses to
environmental salinity in the halophilic black yeast Hortaea werneckii. Mol
Microbiol 45:665–672.
2.   Mitchison-Field LMY, Vargas-Muñiz JM, Stormo BM, Vogt EJD, Van Dierdonck S,
Pelletier JF, Ehrlich C, Lew DJ, Field CM, Gladfelter AS. 2019. Unconventional
Cell Division Cycles from Marine-Derived Yeasts. Curr Biol 29:3439-3456.e5.
3.   Anthonies S, Vargas-Muñiz JM. 2022. Hortaea werneckii isolates exhibit different
pathogenic potential in the invertebrate infection model Galleria mellonella. Front
fungal Biol 3:941691.
4.   Goshima G. 2022. Growth and division mode plasticity is dependent on cell
density in marine-derived black yeasts. Genes to Cells 27:124–137.
5.   Giordano Lorca MC, Maria De la Fuente Larrain A, Lorca Jungmann MB, Kramer
Hepp DB. 2018. Tinea nigra: Report of three pediatrics cases. Rev Chil Pediatr
89:506–510.
6.   Bonifaz A, Badali H, de Hoog GS, Cruz M, Araiza J, Cruz MA, Fierro L, Ponce
RM. 2008. Tinea nigra by Hortaea werneckii, a report of 22 cases from Mexico.
Stud Mycol 61:77–82.
7.   Ng KP, Soo-Hoo TS, Na SL, Tay ST, Hamimah H, Lim PC, Chong PP, Seow HF,
Chavez AJ, Messer SA. 2005. The mycological and molecular study of Hortaea
werneckii isolated from blood and splenic abscess. Mycopathologia 159:495–500.

8.   Badali H, Al-Hatmi AMS, Fakhim H, Moghaddasi A, Khodavaisy S, Vaezi A,
243   Ahangarkani F, de Hoog GS, Meis JF. 2019. In vitro activity of nine antifungal
244   agents against a global collection of Hortaea werneckii isolates, the agent of tinea
245   nigra. Int J Antimicrob Agents https://doi.org/10.1016/j.ijantimicag.2019.05.006.
246   9.   Zalar P, Zupančič J, Gostinčar C, Zajc J, de Hoog GS, De Leo F, Azua-Bustos A,
247   Gunde-Cimerman N. 2019. The extremely halotolerant black yeast Hortaea
248   werneckii - a model for intraspecific hybridization in clonal fungi. IMA Fungus
249   10:1–27.
250   10.   Gostinčar C, Stajich JE, Zupančič J, Zalar P, Gunde-Cimerman N. 2018. Genomic
251   evidence for intraspecific hybridization in a clonal and extremely halotolerant
252   yeast. BMC Genomics 19:1–12.
253   11.   Sinha S, Flibotte S, Neira M, Formby S, Plemenitas A, Cimerman NG, Lenassi M,
254   Gostinčar C, Stajich JE, Nislow C. 2017. Insight into the recent genome
255   duplication of the halophilic yeast Hortaea werneckii: Combining an improved
256   genome with gene expression and chromatin structure. G3 Genes, Genomes,
257   Genet 7:2015–2022.
258   12.   Gostinčar C, Sun X, Černoša A, Fang C, Gunde-Cimerman N, Song Z. 2022.
259   Clonality, inbreeding, and hybridization in two extremotolerant black yeasts.
260   Gigascience 11:1–16.
261   13.   Turk M, Méjanelle L, Šentjurc M, Grimalt JO, Gunde-Cimerman N, Plemenitaš A.
262   2004. Salt-induced changes in lipid composition and membrane fluidity of
263   halophilic yeast-like melanized fungi. Extremophiles 8:53–61.
264   14.   Plemenitaš A, Lenassi M, Konte T, Kejžar A, Zajc J, Gostinčar C, Gunde-
265   Cimerman N. 2014. Adaptation to high salt concentrations in
266   halotolerant/halophilic fungi: A molecular perspective. Front Microbiol 5:1–12.
267   15.   Lenassi M, Gostinčar C, Jackman S, Turk M, Sadowski I, Nislow C, Jones S, Birol
268   I, Cimerman NG, Plemenitaš A. 2013. Whole Genome Duplication and
269   Enrichment of Metal Cation Transporters Revealed by De Novo Genome
270   Sequencing of Extremely Halotolerant Black Yeast Hortaea werneckii. PLoS One
271   8.
272   16.   Romeo O, Marchetta A, Giosa D, Giuffrè L, Urzì C, De Leo F. 2020. Whole
273   Genome Sequencing and Comparative Genome Analysis of the Halotolerant
274   Deep Sea Black Yeast Hortaea werneckii. Life 10:229.
275   17.   Marchetta A, van den Ende BG, Al-Hatmi AMS, Hagen F, Zalar P, Sudhadham M,
276   Gunde-Cimerman N, Urzì C, de Hoog S, De Leo F. 2018. Global molecular
277   diversity of the halotolerant fungus Hortaea werneckii. Life 8:1–12.
278   18.   Lenassi M, Plemenitaš A. 2007. Novel group VII histidine kinase HwHhk7B from
279   the halophilic fungi Hortaea werneckii has a putative role in osmosensing. Curr
280   Genet 51:393–405.
281   19.   Kejžar A, Grötli M, Tamás MJ, Plemenitaš A, Lenassi M. 2015. HwHog1 kinase
282   activity is crucial for survival of Hortaea werneckii in extremely hyperosmolar
283   environments. Fungal Genet Biol 74:45–58.
284   20.   Lenassi M, Vaupotic T, Gunde-Cimerman N, Plemenitas A. 2007. The MAP
285   kinase HwHog1 from the halophilic black yeast Hortaea werneckii: Coping with
286   stresses in solar salterns. Saline Systems 3:1–11.
21.   Turk M, Plemenitaš A. 2002. The HOG pathway in the halophilic black yeast

Hortaea werneckii: Isolation of the HOG1 homolog gene and activation of
HwHog1p. FEMS Microbiol Lett 216:193–199.
22.   Vaupotix̌ T, Plemenitaš A. 2007. Differential gene expression and Hog1
interaction with osmoresponsive genes in the extremely halotolerant black yeast
Hortaea werneckii. BMC Genomics 8:1–15.
23.   Li D, Tang Y, Lin J, Cai W. 2017. Methods for genetic transformation of
filamentous fungi. Microb Cell Fact 16:1–13.
24.   Malavia D, Gow NAR, Usher J. 2020. Advances in molecular tools and in vivo
models for the study of human fungal pathogenesis. Microorganisms 8.
25.   Grahl N, Demers EG, Crocker AW, Hogan DA. 2017. Use of RNA-Protein
Complexes for Genome Editing in Non- albicans Candida Species. mSphere 2:1–
9.
26.   Santana DJ, O'Meara TR. 2021. Forward and reverse genetic dissection of
morphogenesis identifies filament-competent Candida auris strains. Nat Commun
12.
27.   van Rhijn N, Furukawa T, Zhao C, McCann BL, Bignell E, Bromley MJ. 2020.
Development of a marker-free mutagenesis system using CRISPR-Cas9 in the
pathogenic mould Aspergillus fumigatus. Fungal Genet Biol 145:103479.
28.   Al Abdallah Q, Ge W, Fortwendel JR. 2017. A Simple and Universal System for
Gene Manipulation in Aspergillus fumigatus: In Vitro -Assembled Cas9-Guide
RNA Ribonucleoproteins Coupled with Microhomology Repair Templates .
mSphere 2.
29.   Lightfoot JD, Fuller KK. 2019. Crispr/cas9-mediated gene replacement in the
fungal keratitis pathogen fusarium solani var. Petroliphilum. Microorganisms 7.
30.   Souza ACO, Martin-Vicente A, Nywening A V., Ge W, Lowes DJ, Peters BM,
Fortwendel JR. 2021. Loss of Septation Initiation Network (SIN) kinases blocks
tissue invasion and unlocks echinocandin cidal activity against Aspergillus
fumigatus. PLOS Pathog 17:e1009806.
31.   Langfelder K, Philippe B, Jahn B, Latgé J-P, Brakhage AA. 2001. Differential
Expression of the Aspergillus fumigatus pksP Gene Detected In Vitro and In Vivo
with Green Fluorescent Protein. Infect Immun 69:6411–6418.
32.   Kress MRVZ, Savoldi M, Goldman MHS, Ha A, Heinekamp T, Brakhage A a,
Goldman GH. 2006. Mutant De cient for Nonhomologous End Joining Is a
Powerful Tool for Analyzing Pathogenicity in. Society 5:207–211.
33.   Kogej T, Stein M, Volkmann M, Gorbushina AA, Galinski EA, Gunde-Cimerman
N. 2007. Osmotic adaptation of the halophilic fungus Hortaea werneckii: Role of
osmolytes and melanization. Microbiology 153:4261–4273.
325

Figure 1

A

| GMM | GMM+Hygromicin | GMM+Hygromicin |
| EXF-2000 | EXF-2000 | EXF-2000+pUCGH |

B

| DIC | GFP | Merged |

EXF-2000

EXF-2000 + pUCGH

Figure 2