1

High Frequency Terahertz Stimulation Alleviates Neuropathic Pain by Inhibiting the Pyramidal Neuron Activity in the Anterior Cingulate Cortex of mice

Wenyu Peng[1#], Pan Wang[2#], Chaoyang Tan[2#], Han Zhao[2], Kun Chen[2], Huaxing Si[2], Yuchen Tian[1], Anxin Lou[2], Zhi Zhu[4], Yifang Yuan[3], Kaijie Wu[5,6*], Chao Chang[3,7*], Yuanming Wu[1*], Tao Chen[2*]

[1]Department of Biochemistry and Molecular Biology, School of Basic Medicine, The Fourth Military Medical University, 710032 Xi'an, China.

[2]Department of Anatomy, Histology and Embryology and K.K. Leung Brain Research Centre, The Fourth Military Medical University, 710032 Xi'an, China.

[3]Innovation Laboratory of Terahertz Biophysics, National Innovation Institute of Defense Technology, 100071 Beijing, China.

[4]Laboratory of Optical Technology and Instrument for Medicine, Ministry of Education, College of Optical-Electrical and Computer Engineering, University of Shanghai for Science and Technology, 200093 Shanghai, China.

[5]Information Materials and Intelligent Sensing Laboratory of Anhui Province, Anhui University, 230601 Hefei, China

[6]School of Electronic and Information Engineering, Anhui University, 230601 Hefei, China

[7]School of Physics, Peking University, 100871 Beijing, China.

[#]Co-first authors, contribute equally to this work

[*]Corresponding authors: Tao Chen (chtkkl@fmmu.edu.cn), Yuanming Wu (wuym@fmmu.edu.cn), Chao Chang (gwyzlzssb@pku.edu.cn), Kaijie Wu (23109@ahu.edu.cn)

Lead contact: Tao Chen (chtkkl@fmmu.edu.cn)

Abstract

Neuropathic pain (NP) is caused by a lesion or disease of the somatosensory system and is characterized by abnormal hypersensitivity to stimuli and nociceptive responses to non-noxious stimuli, affecting approximately 7–10% of the general population. However, current first-line drugs like non-steroidal anti-inflammatory agents and opioids have limitations, including dose-limiting side effects, dependence, and tolerability issues. Therefore, developing new interventions for the management of NP is urgent. In this study, we discovered that the high-frequency terahertz stimulation (HFTS) at approximate 36 THz effectively alleviates NP symptoms in mice with spared nerve injury. *In vivo* and *in vitro* results demonstrate that HFTS reduces the excitability of pyramidal neurons in the anterior cingulate cortex through enhancing the voltage-gated K^+ (Kv) conductance. Computational simulation suggests that the frequency resonates with the carbonyl group in the filter region of Kv1.2 channels, facilitating the translocation of potassium ions. This research presents a novel optical intervention strategy with terahertz waves for the treatment of NP and holds promising application in other nervous system diseases.

Introduction

Neuropathic pain (NP) refers to a debilitating chronic pain condition, which is often a consequence of nerve injury or of the diseases such as cancer, diabetes mellitus, infection, autoimmune disease, and trauma[1, 2]. The symptoms of NP include spontaneous pain, hyperalgesia and mechanical allodynia. Unfortunately, NP is often resistant to currently available drug treatments, including non-steroidal anti-inflammatory drugs and even opioids[3.] More evidences reveal that NP is not merely a symptom of a disease but rather an expression of pathological operations of the nervous system[4]. Therefore, developing new therapeutic technology aimed at these underlying mechanisms for pain relief represents a considerable challenge.

Compared with the limitations of chemical-based drugs research, physics-based treatment offers a new concept and opportunity for intervening in NP. Optogenetics, as an interdisciplinary approach, has demonstrated therapeutic potential in NP. However, the limitations of viral vector delivery systems in humans are well-known[5]. Recently, evidence has emerged suggesting that high frequency terahertz (THz) photons directly resonate with molecules, thereby regulating corresponding biological functions. For instance, our previous study demonstrated that a 34.88 THz wave resonates with Aβ protein, disrupting the process of fibril formation[6]. Li et al. discover that the band of 42.55 THz resonates with the stretching mode of either the –COO- or the –C=O group significantly enhancing the Ca^{2+} conductance[7]. Zhu et al. conclude that 48.2 THz photons greatly increase the permeability of sodium channel by a factor of 33.6 through breaking the hydrated hydrogen bonding network between the hydrosphere layer of the ions and the carboxylate groups[8]. Additionally, the frequency of 53.5 THz has been reported to enhance the voltage-gated K^+ currents, which modulate the startle response and associative learning[9, 10]. These studies strongly prompt us the potential application of THz photons in the treatment of neuropathic pain by targeting the ion channels[11].

The anterior cingulate cortex plays a crucial role in pain regulation[12, 13]. Our previous research has demonstrated that nociceptive information resulting from nerve injury is transmitted to the ACC[14]. This region exhibits pre- and postsynaptic long-term plasticity (LTP), which contributes to chronic pain and associated negative emotions[15, 16]. Furthermore, descending projection pathways from the ACC enhance the neuronal activity of the spinal dorsal horn (SDH) and regulate nociceptive sensory transmission[17]. Brain-imaging and MRI studies also provide evidence of hyperexcitability in the ACC during both acute and chronic pain[12, 18]. Specifically, the activity of pyramidal cells in the ACC is directly correlated with the expression of chronic pain[19, 20]. Optogenetic excitation of ACC pyramidal cells induces pain, while their inhibition leads to analgesia[21]. Therefore, targeting the cortical regions of the ACC and inhibiting the activity of ACC pyramidal neurons may hold promise as a strategy for treating NP[22-25].

Neuronal excitability is influenced by various types of voltage-gated ion channels and among them, voltage-dependent potassium (Kv) channels, as one of the important physiological regulators of neuronal membrane potentials, has been proposed as potential target candidates for pain therapy[26-28]. Zhao et al. have reported that enhancing Kv currents in injured dorsal root ganglion (DRG) neurons alleviates neuropathic pain[29.] Additionally, Fan et al. have demonstrated that lumbar $(L)_5$ spinal nerve ligation (SNL) leads to a time-dependent decrease in Kv1.2-positive neurons in the ipsilateral L_5 DRG. However, rescuing Kv1.2 expression in the injured L5 DRG attenuates the development and persistence of pain hypersensitivity[30]. These findings highlight the potential of targeting Kv channels as a therapeutic approach for managing pain.

In this study, we investigated the effects of high-frequency terahertz stimulation (HFTS) on the Kv model. By analyzing the absorbance spectra of Kv channels, we observed a significant response to photons with a frequency of approximately 36 THz. Subsequently, we conducted *in vivo* multi-channel recordings and *in vitro* patch recordings to confirm the activation effect of HFTS on Kv channel conductance and its inhibition of neuronal activity in the ACC pyramidal cells. Importantly, the application of HFTS resulted in a significant reduction in pain

behavior in mice with spared nerve injury (SNI). To understand the underlying mechanism, we performed computational simulations that suggest HFTS modulate the resonance of the carbonyl group of Kv1.2, which is crucial for the action potential waveform and frequency.

Results

HFTS attenuates the generation of action potential through molecular dynamics simulation

To identify a specific terahertz (THz) frequency capable of modulating a major subset of voltage-gated potassium (Kv) channels, we developed an integrated model comprising both mouse Kv channels (Protein Data Bank [PDB] ID: 3LUT) and Na^+ channels (PDB ID: 3RVY) (Fig. 1a and Supplement Fig. S1). We conducted a comprehensive analysis of the spectral absorption characteristics within the THz frequency range. Our results revealed a pronounced absorption peak at approximately 36 THz for the potassium channel, which exhibited a considerable board band compared to the absence of a corresponding peak for the sodium channel (Fig. 1a). This indicates a preferential and resonant absorption of photons at the ~36 THz frequency by the potassium channel. We then tested the possible kinetic changes of Kv1.2, the typical and widely distributed Kv channel in the central nervous system [31], following the absorption of these THz photons. Our findings indicated a significant kinetic change in the -C=O groups of the channel filter structure, as evidenced by an expansion of its van der Waals radius by approximately 0.5 Å (Fig. 1b). Furthermore, during exposure to THz photons, the conductance of the potassium ion channel exhibited an almost linear increase with the intensity of the THz field, while the conductance of sodium ions remained largely unchanged (Fig. 1c). Interestingly, under terahertz photonic influence, the cortical neurons model showed significantly decreased in discharge (Fig. 1d). We performed a detailed waveform analysis of the action potentials, including parameters such as full width at half maximum (FWHM) and firing frequency (Fig. 1e). Our observations revealed that the FWHM of action potentials in the THz-exposed group decreased to 95% of the control group (Fig. 1f, red column), and the firing frequency experienced a reduction of approximately 70% after THz photon stimulation (Fig. 1f, green column). These results collectively suggest that THz photons primarily attenuate neuronal firing activity by increasing potassium ion conductance, thereby modulating neuronal excitability.

HFTS enhances voltage-gated K^+ currents of pyramidal neurons in the ACC

To investigate the impact of high-frequency terahertz stimulation (HFTS) on voltage-gated potassium/sodium (Kv/Nav) channels, which play a crucial role in action potential generation and waveform, we conducted whole-cell patch recording from layer-5 pyramidal neurons (PYR^{ACC}) in acute slices of anterior cingulate cortex (ACC) in mice with spared nerve injury (SNI) (Fig. 2a). Initially, we examined the Nav current by applying a series of test pulses (from -80 to -10 mV) with a command voltage ranging from -100 mV (50 ms) (Fig. 2b). Upon illumination with ~36 THz photons (0.3 ± 0.05 mW), we observed that HFTS had no significant effect on the activation curve slope, half-activation voltage, inactivation slope and half-inactivation voltage (Figs. 2c-e). These findings indicate that HFTS does not affect Nav channel-mediated currents. Subsequently, we investigated the impact of HFTS on Kv currents by applying a series of test pulses (100 ms) ranging from -70 to +100 mV with a command voltage of -100 mV (20 ms) (Fig. 2f). Our results demonstrated that the application of HFTS induced a significant increase in the amplitude of K^+ currents and an increased slope of the current-voltage characteristic (I-V) curve (Figs. 2g-i) (Table S1), without affecting the half-activation voltage (Fig. 2j). These experiments revealed the influence of HFTS on Kv channel activity, leading to an acceleration of potassium ion flow and an increase in potassium conductance in PYR^{ACC} neurons. Importantly, these experimental findings were consistent with the results obtained from molecular dynamics analysis.

HFTS reduces the spike frequency of pyramidal neurons in the ACC

We proceeded to investigate the impact of the specific resonant frequency of THz photons on the excitability of

PYR[ACC] neurons in SNI mice. Using whole-cell current-clamp recording, we compared the input-output curves of
evoked action potentials before and after HFTS. Our findings revealed a significant increase in the spike frequency
in SNI mice (Fig. 3a)[32], which was effectively rescued by the application of HFTS (Fig. 3b), but not by 465 nm
blue light stimulation (BLS) (Figs. 3c) (Tables S2-4). To further analyze the properties of single action potentials,
we induced them by applying a depolarizing current pulse of an appropriate suprathreshold magnitude (Figs. 3d-i).
In SNI mice, we observed a decrease in the rheobase and an elevation in the resting membrane potential (RMP)
compared to those in sham mice. However, these alterations were reversed by the application of HFTS, while BLS
had no effects (Figs. 3e-f). Other parameters such as voltage threshold, amplitude, and half-width of the action
potentials were not different between SNI and sham mice. Given that the spike firing, rheobase and RMP are
closely related to low-threshold Kv channels[26, 31, 33], these results suggest that HFTS affects the activity of PYR[ACC]
neurons through its specific impact on Kv channels.

136   HFTS decreases the excitability of pyramidal neurons in the ACC *in vivo*
We then investigate the effect of HFTS on the activities of PYR[ACC] in head-fixed awake SNI mice. One-week prior
to the illumination experiment, a 16-channel electrode was implanted into the ACC. We applied THz photon
stimulation for 15 minutes and compared the neuronal activities before and after HFTS (Fig. 4a). Our findings
revealed a significant decrease in the mean firing rate of ACC neurons after HFTS application in both the sham and
SNI groups (Figs. 4b and f). To further analyze the effect of HFTS on the PYR[ACC], we classified them along with
interneurons in the ACC (INT[ACC]) based on their firing rate, trough-to-peak duration and half width (Fig. 4c), as
described in our previous study[19]. We assessed the internal-spiking interval (ISI) and waveform characteristics of
the isolated neurons in each channel to ensure that the pre- and post-HFTS units originated from the same neuron
(Fig. 4d). In the sham group, we observed that 63.4% of PYR[ACC] neurons exhibited a decrease in firing rate, 10.8%
of PYR[ACC] showed an increase, and 25.8% of PYR[ACC] remained unchanged (93 well-isolated PYR[ACC] neurons
were recognized out of 108 total recorded units). In the SNI group, we found that 61.8% of PYR[ACC] neurons
exhibited decreased activity, 20.3% of PYR[ACC] showed increased activity, and 17.9% of PYR[ACC] remained
unchanged (123 well-isolated PYR neurons were recognized out of 130 total recorded units) (Fig. 4e). Consistently,
the increased mean firing rate of PYR[ACC] neurons in SNI mice was significantly inhibited by the application of
HFTS (Fig. 4g). The activity of INT[ACC] also tended to decrease after HFTS, especially in the SNI group (Figs. 4e
and h). In contrast, blue light stimulation (BLS) has no effect on the mean firing rate on the PYR[ACC] and INT[ACC] in
both sham and SNI mice (Supplement Fig. S2). These results indicate that HFTS reduces the spike firing of ACC
neurons, whereas BLS does not have the same effect.

155   HFTS alleviates mechanical allodynia of SNI mice
Finally, we tested whether application of HFTS into the ACC induced analgesic effects. The SNI surgery and optic
fiber implantation into the ACC were performed one week before pain behavioral tests, which included the
mechanical pain threshold test and Catwalk analysis (Fig. 5a). We compared the paw withdrawal mechanical
thresholds (PWMTs) before and after HFTS (0.3 ± 0.05 mW at the tip of the optic fiber) application and found that
SNI treatment decreased the PWMTs compared to the sham group. However, after the application of HFTS, the
PWMTs significantly increased, even surpassing those in the sham group, and this analgesic effect lasted for 60
minutes. In contrast, the PWMTs did not significantly change in the SNI group with the application of 465 nm blue
light (Fig. 5b).
Furthermore, we performed the Catwalk gait analysis (Figs. 5c-k), which provides exquisite and reliable
observations for evaluating the spontaneous pain behaviors[34]. We focused on the print intensity and print area
related parameters of the left hind paw (ipsilateral side of the injured nerve). We found that SNI treatment
significantly altered the stand time, the stand index, the max contact area, the mean print area, the mean intensity

and the duty cycle (Figs. 5f-k). This suggests that the SNI mice tend to avoid standing and walking on their injured

hind paw due to pain hyper-sensitivity. The application of HFTS but not BLS rescued most of the above parameters,

indicating HFTS' strong analgesic effect.

Discussion

In the present study, we provide evidence that high frequency terahertz photons alleviate neuropathic pain in the

SNI mice by decreasing the excitability of pyramidal neurons in the ACC. The mechanism underlying this effect is

that HFTS increases voltage-gated potassium ion conductance through resonance with the carbonyl group in the

potassium channel filter region (Fig. 6). Unlike optogenetic technology, HFTS can directly regulate the

conformation of ion channel without delivering a transgene that encodes a light-response protein. It exhibits

frequency selectivity and dependence to channel structure. This research suggests that HFTS has potential to serve

as a novel optical technology for the treatment of NP pathology.

Neuropathic pain is closely associated with nociceptor excitability in the ACC[12, 35-37], with ion channels

playing a fundamental role in determining neuronal excitability, particularly in the hyperexcitability of pyramidal

neurons[11]. Excitatory Nav channels, responsible for initiating and depolarizing the action potential, can be targeted

by inhibitors to effectively decrease or eliminate electrical excitability. These inhibitors are commonly used in

neurology as antiepileptic drugs. On the other hand, inhibitory Kv channels, responsible for repolarization,

contribute to the initiation of action potentials in diverse ways[33]. Enhancing Kv conductivity could have a similar

effect to Nav channel blockers. For instance, retigabine, an activator for Kv7, has recently been approved as a

first-in-class antiepileptic drug[38, 39]. Among the 12 subfamilies of Kv channels (Kv1-12), Kv1.2 is the most

prevalent in neuronal membranes[40] and has been reported to be significant associated with neuropathic pain[30, 41, 42].

Although the response frequency of Kv1.2 at ~53 THz [10] or ~34 THz [43] and the corresponding modulation function

have been verified due to the broad absorption band, this study highlights the significant resonance of Kv1.2 filter

structure with photons at 36 THz. By optically stimulating ACC neurons with the frequency of ~36 THz, we

observed a significant reduction in the firing rate of pyramidal neurons' action potentials in SNI mice, accompanied

by a notable enhancement of K^+ conductance. This confirms the effect of THz photons on Kv channels, including

Kv1.2. Moreover, the application of HFTS resulted in significant changes in the rheobase and RMP of pyramidal

cells. Although RMP is primarily stabilized by two-pore K^+ channels and Kv7 background conductance[31], it has

been reported that basal excitability is influenced by the opening of low-threshold Kv1.2 channels, which filter out

small depolarizations and thus control the number of triggered APs[40]. However, we acknowledge the possibility

that THz photons affect other K channel functions, but further research is required to confirm this in the future.

During our research, we focus on studying of pyramidal neurons in the ACC[44]. It has been reported that the

firing rate of glutamatergic pyramidal cells, rather than inhibitory interneurons, increases in the ACC after chronic

pain, suggesting an imbalance of excitatory/inhibitory (E/I) ratio[19, 35.] In the local circuits of the ACC, inhibitory

neurons release GABA and inhibit the activities of pyramidal cells. Different studies by Kang et al.[45] and Joseph

Cichon et al.[46] have reported that specific activation of interneurons in the ACC or in the somatosensory cortex

reduces pyramidal neuron hyperactivity and alleviate mechanical allodynia. Thus, the application of THz may

induce complicated results through affecting the Kv channels distributed on both pyramidal cells and interneurons.

However, as shown in our *in vivo* recording data (Figs. 4G-H), although THz illumination slightly decreased the

activity of interneurons, which could potentially lead to an enhanced activity of local pyramidal cells, the direct and

significant decrease in pyramidal cell activity caused by the illumination would overcome this disinhibitory effect,

ultimately resulting in a net decrease in pyramidal cell activity. The behavioral analgesic effect caused by THz

illumination also confirmed this conclusion.

There are several limitations in this study that should be acknowledged. Firstly, we did not investigate the

thermal effect of HFTS on the Kv channel. Previous research has demonstrated the non-thermal, long-distance

stimulation of high-frequency terahertz stimulation on neuronal activity[10]. Nevertheless, the specific interaction
between ~36 THz and Kv channels in terms of thermal effects remains unexplored. Additionally, in this study, we
used blue light as a comparison, and found no significant changes in potassium current and the excitability of
pyramidal cells. This finding suggests the specificity of the terahertz frequency and supports the existence of
non-thermal effects. Another limitation of our research is the use of an optic fiber to deliver the HFTS into the ACC
region. This invasive approach may pose challenges for potential noninvasive applications. However, we believe
that with the advancement of terahertz enhancement techniques, such as the use of metasurfaces or nanomaterials[47,
48], high-frequency terahertz waves show promising potential for broad applications in regulating diverse brain
diseases, such as episodic ataxia[38, 49], benign familial neonatal convulsions[50], Alzheimer's disease[51], and others.

Material and methods

Animals

Male adult (8–10 weeks) C57BL/6 were used for all experiments. Mice were housed on a 12 h light–dark cycle with food and water freely available. The living conditions were carefully controlled, with temperatures maintained at 22-26℃ and humidity at 40%. All animal procedures in the present experiments were in accordance with protocols approved by the Animal Care Committee of the Fourth Military Medical University. All efforts were made to minimize animal suffering and the number of animals used.

Neuropathic Pain Model

We used spared nerve injury (SNI) model to establish neuropathic pain. The detailed process has been described in our previous study[19]. In brief, mice were generally anaesthetized by 2% isoflurane. Three terminal branches of the left sciatic nerve were exposed by making a direct incision in the skin and a section of the biceps femoris muscle in the left thigh. The tibial nerve and the common peroneal nerve were ligated using 6-0 silk sutures and then sectioned distal to the ligation. After ligating and cutting the nerves, they were carefully put back into their original positions, and the muscle and skin were sutured in two layers. For the sham mice, animals only received an operation that exposed the branches of the left sciatic nerve but without any nerve injury. Following a week accommodation period, pain behaviors were assessed using the *von Frey* filament test and CatWalk gait analysis to confirm the successful establishment of the NP model.

Molecular Dynamics Simulation

The simulation was conducted to gain a deeper understanding of the interaction between terahertz photons and ion channels. A composite model of mouse eukaryotic voltage-gated K^+ channels (PDB ID: 3LUT) and eukaryotic Na^+ channels (PDB ID: 3RVY) was built using the Charmm-GUI website. The model consisted of intact proteins, phospholipid bilayers, and saline solution (with a concentration of 0.15 M). Kinetic calculations were performed using GROMACS 5.1.2 software. The CHARMM 36 force field and periodic boundary conditions were applied to the proteins. Electrostatic interactions were handled using the connected element algorithm Ewald. During the simulation, the Rattle algorithm was used to constrain key lengths. The motion equation was solved using the Velocity-Verlet algorithm with a time step of 2 fs. Initially, the simulation was carried out at a room temperature of 303.5 K to observe the ion transport process within the channels at the molecular level. Subsequently, conductivity values (gNa, gK) for potassium and sodium ions and their corresponding absorption spectrum were calculated. To investigate the effect of ion transport under the influence of terahertz radiation, time-varying electric fields of THz radiation were added to the system. In the interaction of terahertz radiation with biological systems, electrical components play a crucial role. The electric field was used to simulate terahertz radiation, and its formula is as follows:

$$E(t) = A \cdot u \cdot \cos(\omega t + phi)$$

Where A represents the terahertz radiation intensity, u and phi represent the polarization direction and phase of the radiated photon, which set to (0, 0, 1) and 0, respectively. The terahertz radiation frequency v is related to the angular frequency ω by the equation:

$$v = \omega/2\pi$$

The cortical neuron Hodgkin-Huxley (H-H) model links the microscopic level of ion channels to the macroscopic level of currents and action potentials. The model consists of two distinct components: a rapid inward current carried by sodium ions and a slower activating outward current carried by potassium ions. These currents result from independent permeability mechanisms for Na^+ and K^+, where the conductance changes over time and membrane potential. Consequently, the model can replicate and explain a wide range of phenomena, including the shape and propagation of action potentials, the sharp threshold, refractory period, anode-break excitation,

265 accommodation, and subthreshold oscillations. Minor adjustments in key conductance and stimulus current
266 parameters enable the model to describe various action potential phenomena[52]. The formula shows as follows:

$$C\frac{dv}{dt} = G_{stim} - g_{Na}(THz)m^3h(v-v_{Na}) - g_K(THz)n(v-v_K) - g_Ln(v-v_L)$$

$$\frac{dy}{dt} = \alpha_y(1-y) - \beta_y y, \qquad y = m, n$$

$$\frac{dh}{dt} = (\frac{1}{1+\exp((v+60)/6.2)} - h)(\alpha_h + \beta_h),$$

268 Where v, m, h and n represent the membrane voltage, probability of opening or closing of potassium-sodium ion
269 channel. V_{Na}, V_K and V_L are the sodium ion reverse potential, potassium ion reverse potential and resting membrane
270 potential, respectively. g_{Na}, g_K are the maximum conductivity of sodium and potassium ion, respectively. C is
271 membrane capacitive reactance with 0.75 uF/cm^2, the G_{stim} is the stimulation by an external current.

High Frequency Terahertz and Blue Light Stimulation

For high frequency of terahertz stimulation (HFTS), we used a quantum cascade laser with a center frequency of
35.93 ± 0.1 THz. The laser beam was coupled into a coupler, supported by the Innovation Laboratory of Terahertz
Biophysics. We then connected the coupler to a Polycrystalline fiber (PIR) infrared fiber (Art photonics) with a
core composition of AgCl/Br. This fiber has excellent transmittance in the range of 3-18 μm, with a core refractive
index of 2.15 and an effective numerical aperture (NA) of 0.35 ± 0.05. At the distal end of the fiber, we left
approximately 3-4 cm of bare fiber to allow for the insertion of a hollow tube with an inner diameter of 620 μm.
This tube was pre-implanted into the ACC region of the SNI and sham group mice brain (*53*). The duration of
HFTS was 15 minutes, with a pulse width of 2 μs, a repetition frequency of 10 kHz, and a duty cycle of 40%. The
average output power at the tip of the fiber, measured by a MIR detector (NOVA II-3A, Israel), was 0.3 ± 0.05 mW.
For comparison purposes, we also used a blue laser to stimulate the same brain region for 15 minutes, with a
frequency of 1 Hz and an average output power of 10 mW.

In Vitro Patch Clamp Recording

The experimental procedures were based on our previous reports[54]. Briefly, mice were anesthetized and then
decapitated to sacrifice. Brain slices (300 μm thick) containing the ACC were cut on a vibrating microtome (Leica
VT 1200s, Heidelberger, Nussloch, Germany) at 0−4°C in oxygenated (95% O_2 and 5% CO_2) artificial
cerebrospinal fluid (ACSF) consisting of (in mM) 124 NaCl, 25 NaHCO$_3$, 2.5 KCl, 1 NaH$_2$PO$_4$, 2 CaCl$_2$, 1 MgSO$_4$
and 10 glucose. Slices were then transferred to a room temperature-submerged recovery chamber containing
oxygenated ACSF and incubated for at least one hour before patch clamp recording. The neurons were then
visualized under a microscope with infrared differential interference contrast or fluorescent optics video
microscopy (BX51W1, Olympus, Tokyo, Japan). The recording pipettes (3-5 MΩ) were filled with a solution
composed of (in mM) 124 K-gluconate, 5 NaCl, 1 MgCl$_2$, 0.2 EGTA, 2 MgATP, 0.1 Na$_3$GTP, 10 HEPES and 10
phosphocreatine disodium (adjusted to pH 7.2 with KOH, 290 mOsmol). Biocytin (0.2%) were added into pippette
solution for verifying neurons and visualized through biocytin-avidin reaction. To examine the properties of
voltage-gated K$^+$ currents, tetrodotoxin (TTX, 1 μM) and CdCl$_2$ (100 μM) were added into the ACSF. For
recording of voltage-gated Na$^+$ current, 3 mM 4-AP and 0.1 mM CdCl$_2$ were added into the ACSF. Electrical
signals were filtered at 1 kHz by a Multiclamp 700B amplifier (Molecular Devices, USA), and digitized by an
Axon DigiData 1550A converter with a sampling frequency of 10 kHz. Data analyses were performed with the
Clampfit 10.02.

In Vivo Multi-Channel Recording

302 Before the SNI operation, we implanted an electrode into the right ACC as in our previous reports[19], following

stereotaxic coordinates: 1.1 mm anterior to the bregma, 0.3 mm lateral to the midline and 1.8 mm vertical to the skull surface. The electrodes were secured to the exposed skull using the dental adhesive resin cement Super-bond C&B (Japan)[53]. This electrode consisted of 16-channel wire electrodes and included a hollow tube. During the optical stimulation, we employed multi-channel recording technology by Neurolego system (Nanjing Greathink Medical Technology, Nanjing, China). Subsequently, single-unit spike sorting was performed using the MClust-v4.4 toolbox in MATLAB software (MathWorks, USA). In the ACC region, the two main cell types are pyramidal neurons and interneuron cells, which are gamma-aminobutyric acid (GABA) neurons. Pyramidal neurons were primarily classified based on a trough-to-peak duration above 430 us, indicating long-duration action potentials. Interneuron cells, on the other hand, were identified based on a duration time below 430 μs[55].

Behavioral Assays

Mechanical allodynia

Briefly, the paw withdrawal mechanical threshold (PWMT) was evaluated by using von Frey filaments (Stoelting, Kiel, WI, USA) as reported in our previous works[54]. Mice were habituated to the testing environment for 3 days before baseline testing and then placed under inverted plastic boxes ($7 \times 7 \times 10$ cm) on an elevated mesh floor and allowed to habituate for 30 min before threshold testing. A logarithmic series of 8 calibrated Semmes-Weinstein monofilaments (von Frey hairs; Stoelting, Kiel, WI, USA) (0.008, 0.02, 0.04, 0.16, 0.4, 0.6, 1, 1.4, and 2 g) with various bending forces (0.078, 0.196, 0.392, 1.568, 3.92, 5.88, 9.8, 13.72, and 19.6 mN) was applied to the plantar surface of the hind paw until the mice withdrew from the stimulus. Positive responses included licking, biting, and sudden withdrawal of the hind paws. A von Frey filament was applied 5 times (3 seconds for each stimulus) to each tested area. The minimum bending force of the von Frey filament able to evoke 3 occurrences of the paw withdrawal reflex was considered the paw withdrawal threshold. All tests were performed in a blinded manner.

CatWalk gait analysis

Gait analysis was conducted using the CatWalk XT system (Noldus, the Netherlands) to measure pain-related parameters. The experimental setup involved placing the mouse on a glass platform with open ends, allowing the mouse to walk voluntarily. Simultaneously, a high-speed camera positioned underneath the platform captured images of each step, which were then transmitted to the analysis software (version 10.6, CatWalk XT, Noldus) for further processing. In this study, eight parameters were identified to assess dynamic behaviors relevant to neuropathic pain: (1) Stand: this parameter represents the duration (in seconds) of a paw touching the glass plate; (2) Stand index: it describes the speed at which the paw moves away from the glass plate; (3) Max contact area: it describes the maximum contact area of the paw or leg with the glass plate; (4) Mean print area: it represents the average area of the paw print during locomotion; (5) Mean intensity: this parameter denotes the average intensity value of the running stage; (6) Duty cycle This parameter denotes the average intensity value of the running stage.

Statistical Analysis

GraphPad Prism 5 (Graph Pad Software, Inc.) was used for the statistical analyses and graphing. Statistical significance was assessed by unpaired *t*-test, paired *t*-test, one-way and two-way ANOVA followed by *post hoc* comparison. All data in the experiment are expressed in mean ± S.E.M. Statistical significance was indicated as *, $P < 0.05$, **, $P < 0.01$, ***, $P < 0.001$ and ****, $P < 0.0001$.

Competing interests

The authors declare there are no conflicts of interest/competing interests related to this work.

Author contributions

W.-Y.P., Data curtation, Formal analysis, Investigation, Visualization, Writing-original draft, Writing-review and

editing, Funding acquisition; P.W., Data curation, Investigation, Methodology; C.-Y.T., Data curation, Investigation, Methodology, Writing-original draft; H.Z., Investigation, Methodology; K.C., Investigation; H.-X.S., Methodology; Y.-C.T., Methodology; A.-X.L., Methodology; Z.Z., Investigation, software; Y.-F.Y., Investigation; K.-J.W., Data curation, Investigation, Methodology, Writing-original draft; C.C., Conceptualization, Supervision; Y.-M.W., Supervision, Investigation, Methodology; T.C., Conceptualization, Supervision, Funding acquisition, Investigation, Visualization, Methodology, Writing-review and editing, Project administration, Funding acquisition.

Acknowledgments

We thank the support the teaching center of Air force medical university for the invaluable technical assistance and J.-X.G. of our laboratory for the helpful comments on an earlier version of the manuscript and discussions. This work was supported by grants from the National Natural Science Foundation of China (32192410, 32071000 to T. C., 82271893 to Y.-M.W., 8230071226 to W.-Y.P., 82302090 to K.C.), National Science Fund for Distinguished Young Scholars (12225511 to C.C.), National Science Fund of China Major Project (T2241002 to C.C.), Innovation Laboratory of Terahertz Biophysics (23-163-00-GZ-001-001-02-04 to W.-Y.P.), The Key Research and Development Plan of Shaanxi Province (S2024-YF-YBSF-0277 to W.-Y.P.)

References

1. R. Baron, G. Binder A Fau - Wasner, G. Wasner, Neuropathic pain: diagnosis, pathophysiological mechanisms, and treatment. *Neurology* vol. 9(8), 807-19 (2010).

2. L. Shan, D. Selvarajah, Z. Wang, Editorial: Insights in neuropathic pain: 2022. *Frontiers in pain research (Lausanne, Switzerland)* 4, 1232025 (2023).

3. T. S. Jensen, N. B. Finnerup, Allodynia and hyperalgesia in neuropathic pain: clinical manifestations and mechanisms. *Lancet Neurol.* 13(9), 924-35 (2014).

4. M. Costigan, C. J. Scholz J Fau - Woolf, C. J. Woolf, Neuropathic pain: a maladaptive response of the nervous system to damage. *Annu. Rev. Neurosci.* 32, 1-32 (2009).

5. K. Liu, L. Wang, Optogenetics: Therapeutic spark in neuropathic pain. *Bosnian J. Basic Med. Sci.* 19(4), 321-327 (2019).

6. W. Peng *et al.*, High-frequency terahertz waves disrupt Alzheimer's β-amyloid fibril formation. *eLight* **3**, 18 (2023).

7. Y. A.-O. Li, C. A.-O. Chang, Z. A.-O. Zhu, L. Sun, C. Fan, Terahertz Wave Enhances Permeability of the Voltage-Gated Calcium Channel. *J. Am. Chem. Soc.* 143 (11), 4311-4318 (2021).

8. Y. Zhao, L. Wang, Y. Li, Z. Zhu, Terahertz Waves Enhance the Permeability of Sodium Channels. *Symmetry* 15(2), 427 (2023).

9. J. Zhang *et al.*, Non-invasive, opsin-free mid-infrared modulation activates cortical neurons and accelerates associative learning. *Nat. Commun.* **12**, 2730 (2021).

10. X. A.-O. Liu *et al.*, Nonthermal and reversible control of neuronal signaling and behavior by midinfrared stimulation. *Proc Natl Acad Sci U S A.* 118(10) (2021).

11. J. S. Trimmer, Ion channels and pain: important steps towards validating a new therapeutic target for neuropathic pain. *Exp. Neurol.* 254, 190-194 (2014).

12. T. V. Bliss, G. L. Collingridge, B. K. Kaang, M. Zhuo, Synaptic plasticity in the anterior cingulate cortex in acute and chronic pain. *Nat Rev Neurosci.* 17(8), 485-496 (2016)

13.   T. V. P. Bliss, G. L. Collingridge, B.-K. Kaang, M. Zhuo, Synaptic plasticity in the anterior cingulate cortex in
acute and chronic pain. *Nat. Rev. Neurosci.* **17**, 485-496 (2016).

14.   M. A.-O. Tsuda, K. Koga, T. Chen, M. Zhuo, Neuronal and microglial mechanisms for neuropathic pain in the
spinal dorsal horn and anterior cingulate cortex. *Neurochem.* 141(4), 486-498 (2017).

15.   W. Chen T Fau - Wang *et al.*, Postsynaptic insertion of AMPA receptor onto cortical pyramidal neurons in the
anterior cingulate cortex after peripheral nerve injury. *Mol Brain.* 7, 76 (2014).

16.   K. Koga *et al.*, Coexistence of two forms of LTP in ACC provides a synaptic mechanism for the interactions
between anxiety and chronic pain. *Neuron.* 85(2), 377-389 (2015).

17.   T. Chen *et al.*, Top-down descending facilitation of spinal sensory excitatory transmission from the anterior
cingulate cortex. *Nat. Commun.* 9(1), 1886 (2018).

18.   S. R. A. Alles, P. A. Smith, Etiology and Pharmacology of Neuropathic Pain. *Pharmacol Rev,* 70(2), 315-347
(2018)

19.   D. Y. Zhu *et al.*, The increased in vivo firing of pyramidal cells but not interneurons in the anterior cingulate
cortex after neuropathic pain. *Mol Brain.* 15(1), 12 (2022).

20.   X.-Y. Li *et al.*, Alleviating Neuropathic Pain Hypersensitivity by Inhibiting PKMζ in the Anterior Cingulate
Cortex. *Science* **330**, 1400-1404 (2010).

21.   S. J. Kang *et al.*, Bidirectional modulation of hyperalgesia via the specific control of excitatory and inhibitory
neuronal activity in the ACC. *Mol Brain.* **8**, 81 (2015).

22.   P. N. Fuchs, Y. B. Peng, J. A. Boyette-Davis, M. L. Uhelski, The anterior cingulate cortex and pain processing.
Front Integr Neurosci. 8, 35 (2014).

23.   K. Lançon, C. Qu, E. Navratilova, F. Porreca, P. Séguéla, Decreased dopaminergic inhibition of pyramidal
neurons in anterior cingulate cortex maintains chronic neuropathic pain. *Cell Rep.* 37(9), 109933 (2021).

24.   S. W. Um, M. J. Kim, J. W. Leem, S. J. Bai, B. H. Lee, Pain-Relieving Effects of mTOR Inhibitor in the
Anterior Cingulate Cortex of Neuropathic Rats. *Mol. Neurobiol.* **56**, 2482-2494 (2019).

25.   W. Tan *et al.*, Alleviating neuropathic pain mechanical allodynia by increasing Cdh1 in the anterior cingulate
cortex. *Mol Pain.* 11, 56 (2015).

26.   M. Takeda *et al.*, Potassium channels as a potential therapeutic target for trigeminal neuropathic and
inflammatory pain. *Mol Pain.* 7, 5 (2011).

27.   X. Du, N. Gamper, Potassium channels in peripheral pain pathways: expression, function and therapeutic
potential.   *Curr. Neuropharmacol.* 11(6), 621-40 (2013).

28.   A. Sakai *et al.*, MicroRNA cluster miR-17-92 regulates multiple functionally related voltage-gated potassium
channels in chronic neuropathic pain. *Nat. Commun.* **8**, 16079 (2017).

29.   J.-Y. Zhao *et al.*, DNA methyltransferase DNMT3a contributes to neuropathic pain by repressing Kcna2 in
primary afferent neurons. *Nat. Commun.* **8**, 14712 (2017).

30.   L. Fan *et al.*, Impaired neuropathic pain and preserved acute pain in rats overexpressing voltage-gated potassium
channel subunit Kv1.2 in primary afferent neurons. *Mol. Pain* 10, 8 (2014).

31.   C. Tsantoulas, S. B. McMahon, Opening paths to novel analgesics: the role of potassium channels in chronic
pain. *Trends Neurosci.* 37(3), 146-158 (2014).

32.   X. Y. Cao *et al.*, Characterization of intrinsic properties of cingulate pyramidal neurons in adult mice after nerve
injury. *Mol. Pain.* 5, 73 (2009).

33.   C. González *et al.*, K(+) channels: function-structural overview. *Compr Physiol.* 2(3), 2087-2149 (2012).

34.   K. L. Zhang *et al.*, Targeted up-regulation of Drp1 in dorsal horn attenuates neuropathic pain hypersensitivity by
increasing mitochondrial fission. *Redox Biol.* 49, 102216 (2022).

35.   R. Zhao *et al.*, Neuropathic Pain Causes Pyramidal Neuronal Hyperactivity in the Anterior Cingulate Cortex.

Front. Cell. Neurosci. 12 (2018).

36.   Z. Chen, X. Shen, L. Huang, H. Wu, M. Zhang, Membrane potential synchrony of neurons in anterior cingulate
cortex plays a pivotal role in generation of neuropathic pain. *Sci. Rep.* **8**, 1691 (2018).

37.   H. Xu *et al.*, Presynaptic and postsynaptic amplifications of neuropathic pain in the anterior cingulate cortex. *J*
Neurosci. 28(29), 7445-7453 (2008).

38.   K. Gao, Z. Lin, S. Wen, Y. Jiang, Potassium channels and epilepsy. *Acta Neurol Scand.* 146(6), 699-707 (2022).

39.   S. Maljevic, H. Lerche, Potassium channels: a review of broadening therapeutic possibilities for neurological
diseases. *J Neurol.* 260(9), 2201-2211 (2013).

40.   A. Al-Sabi, Seshu K. Kaza, J. O. Dolly, J. Wang, Pharmacological characteristics of Kv1.1- and
Kv1.2-containing channels are influenced by the stoichiometry and positioning of their α subunits. *Biochem. J.*
454, 101-108 (2013).

41.   Y. Liang *et al.*, Transcription factor EBF1 mitigates neuropathic pain by rescuing Kv1.2 expression in primary
sensory neurons. *Transl Res.* **263**, 15-27 (2024).

42.   J. Zhang *et al.*, Epigenetic restoration of voltage-gated potassium channel Kv1.2 alleviates nerve injury-induced
neuropathic pain. *J. Neurochem.* **156**, 367-378 (2021).

43.   T. A.-O. Xiao *et al.*, Sensory input-dependent gain modulation of the optokinetic nystagmus by mid-infrared
stimulation in pigeons. *Elife.* 12:e78729 (2023).

44.   X. Xiao, Y.-Q. Zhang, A new perspective on the anterior cingulate cortex and affective pain. *Neurosci. Biobehav.*
Rev. **90**, 200-211 (2018).

45.   S. J. Kang *et al.*, Bidirectional modulation of hyperalgesia via the specific control of excitatory and inhibitory
neuronal activity in the ACC. *Mol. Brain* vol. 8,1 81. 2 (2015).

46.   J. Cichon, T. J. J. Blanck, W.-B. Gan, G. Yang, Activation of cortical somatostatin interneurons prevents the
development of neuropathic pain. *Nat. Neurosci.* **20**, 1122-1132 (2017).

47.   R. Zhou, W. Wang C Fau - Xu, L. Xu W Fau - Xie, L. Xie, Biological applications of terahertz technology based
on nanomaterials and nanostructures. *Nanoscale*, 11, 3445-3457 (2019).

48.   Q. Wang, Y. Chen, J. Mao, F. Yang, N. A.-O. Wang, Metasurface-Assisted Terahertz Sensing. *Sensors* 23(13),
5902 (2023).

49.   B. A. Cole, S. J. Clapcote, S. P. Muench, J. D. Lippiat, Targeting K(Na)1.1 channels in KCNT1-associated
epilepsy. *Pharmacol Sci.* 42(8), 700-713 (2021).

50.   T. J. Jentsch, Neuronal KCNQ potassium channels: physiology and role in disease. *Nat Rev Neurosci* 1, 21–30
(2000).

51.   J. L. Taylor *et al.*, Functionally linked potassium channel activity in cerebral endothelial and smooth muscle
cells is compromised in Alzheimer's disease. *Proc Natl Acad Sci U S A.* 119(26), e2204581119 (2022).

52.   M. Häusser, The Hodgkin-Huxley theory of the action potential. *Nature Neuroscience* **3**, 1165 (2000/11/01,
2000).

53.   Z. V. Guo *et al.*, Procedures for behavioral experiments in head-fixed mice. *PLoS One* 9(6), e101397 (2014).

54.   M. M. Zhang *et al.*, Glutamatergic synapses from the insular cortex to the basolateral amygdala encode
observational pain. *Neuron*, 110(12), 1993-2008.e6 (2022).

55.   A. V. Apkarian, R.-D. Bushnell Mc Fau - Treede, J.-K. Treede Rd Fau - Zubieta, J. K. Zubieta, Human brain
mechanisms of pain perception and regulation in health and disease. *Eur J Pain.* 9(4), 463-484 (2005).

Fig. 1 Specific frequency THz photons resonate K_v channel and decrease the AP firing rate in cortical neurons through molecular dynamics simulation. (a) Absorbance spectra of voltage-gated potassium/sodium ion channels and the bulk water. (b) The dynamic attributes of the Kv1.2 filter structure in pre- and post-exposure to HFTS. Purple balls represent the K^+, blue balls represent the Cl^-. (c) The alterations in potassium/sodium ion conductance consequent to the influence of HFTS. (d) Changes of the firing rate of APs of cortical neuron models before and after HFTS. (e) The FWHM of an AP pre- and post HFTS. (f) Changes in FWHM and firing frequency with or without HFTS. HFTS, high frequency terahertz stimulation. AP, action potential. FWHM, Full Wide of Half Maximum.

Fig. 2 HFTS enhances Kv currents of pyramidal neurons in SNI mouse *in vitro*. (a) Anatomical location of ACC region in mice and a recorded PYR neuron (biocytin-labeled, green). (b) Representative Nav currents without (orange) or with HFTS (red) under the given step voltage protocol. (c) The application of HFTS does not change the activation and inactivation curves of Nav currents. (d-e) The corresponding slopes of the activation and inactivation curves (d) and the comparison of the half-activation and inactivation voltages (e). (f) Representative Kv currents evoked by a series of step voltages (inset) without (orange) or with HFTS (red). (g) I-V plots constructed from the values of traces shown in (f) display a stronger K^+ current with HFTS. (SNI *vs.* SNI + HFTS: $F_{(1, 66)} = 67.97$, $P < 0.0001$, $n_{SNI} = 7$, $n_{SNI+HFTS} = 7$; Two-way ANOVA followed by *post hoc* comparison using the Šídák's multiple comparisons test (Supplement Table 1). (h) The application of HFTS does not change the activation curves of the Kv currents. (i-j) The corresponding slopes of the activation curves (i) and the half-activation voltages (j). SNI *vs.* SNI + HFTS: $t = 5.872$, $P = 0.0011$, n = 7, unpaired *t*-test. **, $P < 0.01$, ****, $P < 0.0001$.

Fig. 3 HFTS reduces the APs firing rate of pyramidal neurons in SNI mice *in vitro*. (a-c) Representative traces (upper panels) and line-charts (lower panels) showing the changes of evoked spikes of pyramidal neurons in different groups Sham *vs*. SNI: $F_{(1, 40)} = 124.2$, $P < 0.001$, $n_{sham} = 6$, $n_{SNI} = 6$; SNI *vs*. SNI + HFTS: $F_{(1, 40)} = 23.13$, $P < 0.0001$, $n_{SNI} = 6$, $n_{SNI+HFTS} = 6$; SNI *vs*. SNI + BLS: $F_{(1, 40)} = 0.1401$, $P = 0.7101$, $n_{SNI} = 6$, $n_{SNI+BLS} = 6$. Two-way ANOVA followed by *post hoc* comparison using the Šídák's multiple comparisons test (Supplement Table 2-4). (d) Superimposed traces showing the single AP evoked by threshold current stimulation in different groups. (e) Histograms showing the statistical comparison of rheobase in each group. (Sham *vs*. SNI: $P < 0.001$, $n_{sham} = 6$, $n_{SNI} = 19$; SNI *vs*. SNI + HFTS: $P < 0.01$, $n_{SNI} = 19$, $n_{SNI+HFTS} = 6$, one-way ANOVA followed by *post hoc* comparison using the Tukey's multiple comparisons test). (f) The RMP in each group (Sham *vs*. SNI: q = 4.5, $P < 0.05$, $n_{sham} = 6$, $n_{SNI} = 19$; SNI *vs*. SNI + HFTS: $P < 0.05$, $n_{SNI} = 19$, $n_{SNI+HFTS} = 6$, one-way ANOVA followed by *post hoc* comparison using the Tukey's multiple comparisons test). (g-i) HFTS has no significant effect on the threshold, amplitude and half-width of APs in pyramidal neurons. *, $P < 0.05$, **, $P < 0.01$, ***, $P < 0.001$, ****, $P < 0.0001$. BLS, blue light stimulation.

Fig. 4 HFTS decreases the mean firing rate of pyramidal neurons in the ACC in both sham and SNI awake mice. (a) Schematic showing the timeline and the stimulating pattern of HFTS on an awake mouse, as well as the single-unit recording of the ACC using an *in vivo* multi-channel recording technique. (b) Example recording signals of ACC neurons before and after HFTS application in sham and SNI groups, respectively. (c) ACC neurons are classified as pyramidal (PYR) cells and interneurons (INT) using *k*-means cluster-separation algorithm based on their electrophysiological properties. (d) Histograms of the inter-spike intervals (ISI) from the spikes of a PYR and an INT in pre- and post-HFTS recording period. Insets at the top right corner show the waveforms of the detected single unit. (e) Pie charts summarize the changes in firing rate of PYR and INT in sham and SNI groups. Pre *vs.* post HFTS, Wilcoxon rank-sum test. (f) The mean firing rate of all recorded neurons in sham and SNI groups before and after HFTS. Sham ($P < 0.0001$, n = 73, paired *t*-test,), SNI ($P < 0.0001$, n = 130, paired *t*-test). (g) The mean firing rate of PYR neurons in sham and SNI groups before and after HFTS. Sham group ($P < 0.0001$, n = 93, paired *t*-test), SNI group ($P < 0.0001$, n = 123, paired *t*-test). (h) The mean firing rate of INT neurons in sham and SNI groups before and after HFTS. Sham group ($P = 0.065$, n = 15, paired *t*-test), SNI group ($P = 0.0115$, n = 7, paired *t*-test). *, $P < 0.05$, ****, $P < 0.0001$.

Fig. 5 HFTS alleviates neuropathic pain of SNI mice through pain behavior tests. (a) Schematic of the establishment of NP model, the application of HFTS in ACC region and the following behavior tests including *Von Frey* test and Catwalk analysis. (b) HFTS increases the paw withdrawal mechanical thresholds (PWMTs) compared to the SNI model ($F_{(18, 140)}$ = 12.65. $P < 0.0001$. Sham *vs.* SNI: $P < 0.0001$; SNI *vs.* SNI + HFTS: $P < 0.0001$; n = 6 in each group. Two-way ANOVA repeated measures followed by *post hoc* comparison using the Šídák's multiple comparisons test). (c-d) The print view of a mouse (c) and the step sequence of a sham mouse who passing through the glass pane (d). The red line represents the maximum intensity of each foot, the color box represents the mean intensity of the corresponding print during walking. (e) The feet view of the left front (LF), left hind (LH), right front (RF) and right hind (RH) in the groups of sham, SNI, SNI + HFTS and SNI + BLS, respectively. (f) HFTS

increases the LH stand time of SNI mice (sham *vs.* SNI: $P < 0.0001$; SNI *vs.* SNI + HFTS: $P < 0.01$). (g) HFTS increases the LH stand index of SNI mice (sham *vs.* SNI: $P < 0.001$; SNI *vs*. SNI + HFTS: $P < 0.05$). (h) HFTS increases the LH max contact area of SNI mice (sham *vs.* SNI: $P < 0.0001$; SNI *vs*. SNI + HFTS: $P < 0.05$). (i) HFTS increases the LH mean print area of SNI mice (sham *vs.* SNI: $P < 0.0001$; SNI *vs*. SNI + HFTS: $P < 0.05$; SNI vs. SNI + BLS: $P < 0.05$). (j) HFTS increases the LH mean intensity of SNI mice (sham *vs.* SNI: $P < 0.05$; SNI *vs*. SNI + HFTS: $P < 0.05$). (k) HFTS have no significant for the pain behavior parameter of the duty cycle. *, $P < 0.05$, **, $P < 0.01$, ***, $P < 0.001$, ****, $P < 0.0001$. One-way ANOVA (f-k) followed by *post hoc* comparison using the Tukey's multiple comparisons test. $n_{Sham} = 38$, $n_{SNI} = 35$, $n_{SNI+HFTS} = 34$, $n_{SNI+BLS} = 9$.

Fig. 6 Schematic diagram shows the mechanism of HFTS in alleviating neuropathic pain. Left panel shows the
544 group with BLS and the right panel shows the group with HFTS.