Cell-cycle and Age-Related Modulations in Mouse Chromosome Stiffness

Ning Liu[1], Wenan Qiang[2], Philip Jordan[3,4], John Marko[5,6 *,] Huanyu Qiao[1 *]

1. Department of Comparative Biosciences, University of Illinois at Urbana-Champaign, Urbana, IL, USA
2. Center for Developmental Therapeutics, Northwestern University, Evanston, IL, USA
3. Biochemistry and Molecular Biology Departments, Johns Hopkins University, Baltimore, MD, USA
4. School of Medicine, Uniformed Services University of the Health Sciences, Bethesda, MD, USA
5. Department of Molecular Biosciences, Northwestern University, Evanston, IL, USA
6. Department of Physics and Astronomy, Northwestern University, Evanston, IL, USA

*Corresponding authors

Keywords: meiosis, spermatocyte, oocyte, cohesin protein, chromosome stiffness, cell cycle, age

Abstract

The intricate structure of chromosomes is complex, and many aspects of chromosome

configuration/organization remain to be fully understood. Measuring chromosome stiffness can

provide valuable insights into their structure. However, the nature of chromosome stiffness,

whether static or dynamic, remains elusive. In this study, we analyzed chromosome stiffness in

MI and MII oocytes. We revealed that MI oocytes had a ten-fold increase in stiffness compared

to mitotic chromosomes, whereas chromosome stiffness in MII oocytes was relatively low

chromosome. We then investigated the contribution of meiosis-specific cohesin complexes to

chromosome stiffness in MI and MII oocytes. Surprisingly, the Young's modulus of

chromosomes from the three meiosis-specific cohesin mutants did not exhibit significant

differences compared to the wild type, indicating that these proteins may not play a substantial

role in determining chromosome stiffness. Additionally, our findings revealed an age-related

increase in chromosome stiffness in MI oocytes. Age correlates with elevated DNA damage

levels, so we investigated the impact of etoposide-induced DNA damage on chromosome

stiffness, discovering a reduction in stiffness in response to such damage in MI oocytes. Overall,

our study underscores the dynamic nature of chromosome stiffness, subject to changes

influenced by the cell cycle and age.

Introduction

DNA serves as the genetic carrier of the genome and binds with various types of protein

to form chromatin at interphase. Chromatin condenses into a thick thread-like structure known as

a chromosome during cell division, which plays a critical role in ensuring the even distribution of

genetic material [1]. This process is essential for genome integrity, as any issues with

chromosomes during meiosis or mitosis can cause defects such as cancer, infertility, miscarriage,

or congenital diseases [2,3]. The chromosome structure has not been fully understood despite

being discovered for several centuries.

The condensation of chromosomes is not a random process, but rather a precisely

regulated one. Several-inches-long chromatin is compacted into a chromosome less than 10 μm

in size in a stepwise manner [4]. The process involves wrapping of histone octamers to form

nucleosome units after DNA replication, known as the "beads on a string" model [5]. After

adding histone H1, the 10nm fiber turns into a 30nm fiber [6]. H1, a linker histone, is located at

the site of DNA entry and exit base of the nucleosome and binds to the linker region of the DNA

segments. The 30nm fiber has two classic models, the "Solenoid" model and the "Zig-zag"

model. The 10nm fiber and 30nm fiber structures could be repeated *in vitro* using different

concentrations of salt solution [7,8].

However, further condensation cannot be solely explained by the interaction between

histone and DNA. Many structure proteins must be involved in this process [9]. There are

different models for chromosome condensation, with the most popular being the

"Scaffold/Radial-Loop" model [10,11] and the "Chromatin Network" model [12]. The first

model proposes a continuous central protein core with chromatin loops stacked around it, based

on electron microscopy studies. The second model suggests that structure proteins link chromatin

segments but do not form a continuous core. Experiments on chromosomes stiffness

measurement found that MNase degraded them entirely and ablated their stiffness, contradicting

the Scaffold/Radial-Loop model[9]. DNase with different cutting frequencies affected

chromosome stiffness, supporting the Chromatin Network model. Studying chromosome

mechanics helps us understand their components and unravel their true structure.

The chromosome condensation and decondensation occur during the cell cycle [10].

Therefore, the chromosomes in different cell cycles have different shapes and sizes. It is rational

to postulate that their chromosome stiffness is also different, especially mitotic and meiotic

chromosomes even though they looked similar to each other [11,12]. By measuring chromosome

stiffness, we found that the chromosomes from prophase I spermatocytes are approximately ten

times stiffer than those in mitosis [13], indicating that the chromosome structure and components

are different between meiosis I and mitosis. During meiosis I, a unique railroad-track-like protein

structure, the synaptonemal complex (SC), forms between the two homologous chromosomes

[14,15]. Although SC is a natural candidate to explain the structure difference, we have

confirmed that one SC component SYCP1 does not play a role in chromosome stiffness [13].

SYCP1 connects the two "rails" of the SC, axial elements (AEs), and holds them laterally. It may

not contribute to the longitudinal stiffness. However, the chromosome axis, AE, most likely

provides strength to the meiotic chromosomes longitudinally. One of the basic components of

AE are cohesin proteins, which connect sister chromatids in both mitosis and meiosis [16,17].

There are four meiosis-specific cohesin proteins, SMC1β, RAD21L, REC8, and STAG3, which

may play a role in chromosome structure and stiffness [18–20].

Aging can also bring about significant changes in chromosomes, potentially leading to

cell apoptosis, senescence, or cancer, impacting the lifespan and well-being of both animals and

humans [21][22]. Moreover, protein levels on chromosomes are dynamic, either increasing or

decreasing during the aging process [23,24]. For instance, the cohesin along chromosome axes

decreases with age, potentially contributing to higher rates of unsuccessful gamete production

[25][26]. There is an elevated aneuploidy rate for oocytes in older age in oocytes, possibly linked

to alterations in chromosome structure and organization. Therefore, it is essential to explore how

aging may impact chromosome mechanics and gain a better understanding of the underlying

mechanisms.

Results

Stiffness measurement of the chromosomes isolated from MI and MII mouse oocytes

To measure the stiffness of oocyte chromosomes, we first need to isolate chromosomes

from the oocytes collected from 3 to 4 week-old mice that were cultured for 6 h (hours) to reach

metaphase I (MI) stage or 14 h to reach metaphase II (MII) stage. The zona pellucida was

removed by Tyrode's solution treatment for around 3 min (Figure 1A). The oocyte membrane

was then lysed by microspraying Triton X-100, allowing the contents inside of the oocyte to flow

out spontaneously. Using this technique, we successfully isolated the spindle from the oocytes

(Figure 1B Left) and separated the chromosomes from the spindle (Figure 1B Middle). Next, we

used two pipettes with small openings to grab the two ends of the chromosome (Figure 1B Right).

This approach allowed us to measure the stiffness of the isolated chromosome using a

micromanipulation system (see Materials and Methods for further details). We stretched and

relaxed the chromosome to monitor its length change with applied force to determine the

Young's modulus (a measure of the stiffness of a material) of the isolated chromosome. Our

experiments were carried for extensions of less than twice native length, where the chromosome

mechanical response was reversible, i.e., the same force versus extension was obtained during

extension and subsequent retraction.

Chromosome stiffness from MI oocytes is about 10 times higher than that in mitotic cells

Chromosomes stiffness has been extensively studied in various mitotic cells, revealing

similarities and differences in chromosomes stiffness among different cell types [27–29].

However, the chromosome stiffness at different cell cycles have not been completely studied. In

this project, we measured the chromosome stiffness of MEFs (Mouse Embryonic Fibroblasts)

and found that the average Young's Modulus of MEF chromosomes was 340 ± 80 Pa (Figure

2B), which is consistent with our previously published data that MEF chromosomes have a

modulus of 370 ± 70 Pa [13]. Next, we isolated chromosomes from mouse MI oocytes and

measured their stiffness (Figure 2A). Our results showed that chromosome stiffness from MI

mouse oocytes was 3790 ± 700 Pa, which was significantly higher than that of MEF

chromosomes (Figure 2B). This finding was also consistent with prior results which

demonstrated that spermatocyte prophase I chromosomes are approximately 10 times stiffer than

MEF chromosomes [13]. These results suggest that the high chromosome stiffness observed in

meiotic cells is a common feature in gametes, rather than a sex-specific phenomenon. We then

sought to investigate the chromosome stiffness of MII oocytes and possible factors that

contribute to the high chromosome stiffness in gametes.

To confirm the consistency of chromosome measurements, we conducted a comparative

analysis with data previously published [13], in terms of the "doubling force", or force needed to

double chromosome length. MEF chromosomes exhibited a doubling force of 190 ± 40 pN,

while WT spermatocytes had a doubling force of 2130 ± 440 pN [13]. Our results showed that

MEF chromosomes have a doubling force of 210 ± 40 pN and WT Spermatocytes of 1690 ± 450

pN, which is very similar to the published data (Supplement Fig.1). This close correspondence

across experiments establishes the consistency of our chromosome measurements, supporting

their utility in evaluating chromosome stiffness and furthering our understanding of chromosome

mechanics.

The stiffness of chromosomes in MI mouse oocytes is significantly higher compared to MII

oocytes

To study the effect of cell stages on chromosome stiffness, we measured the chromosome

stiffness of the MII oocytes, as we did for the MI chromosomes. As postulated, we found that the

chromosome stiffness of MII oocytes was significantly lower than that of MI oocytes.

Specifically, the Young's Modulus of MII oocytes was 670 ± 130 Pa, while that of MI oocytes is

3790 ± 700 Pa (P<0.001; Figure 2B). Surprisingly, the stiffness of MII oocytes' chromosomes is

slightly higher than that of mitotic cells (Figure 2B). Hence, this finding suggests that meiosis II

is not merely a special analogous event to mitosis, which is contrary to the previous belief that

meiosis II is similar to mitosis [30]. In summary, these results affirm that chromosome stiffness

varies dynamically across different cell stages.

Meiosis-specific cohesins do not contribute to chromosome stiffness

We previously demonstrated that the central elements of the synaptonemal complex do

not contribute to chromosome stiffness, and so we shifted our focus to the role meiotic cohesins

during the meiosis I stage [13]. Cohesins can still load onto chromosomes even when an intact

synaptonemal complex is not formed [31]. During mammalian mitosis, cohesin proteins load

along the chromosome axis during S phase and remain bound to centromeres until they are

cleaved by separase during anaphase. Specifically, majority of cohesin bound to chromosome

arms are already removed by a separase-independent pathway as early as metaphase [32].

However, in the first division of meiosis, anaphase I, cohesin proteins are removed from

chromosome arms; and only a small amount of cohesin proteins is retained at the centromere

until anaphase II [33]. Thus, at mitotic metaphase and MII, cohesin proteins disappear from

chromosome arms but remain at the centromere. In contrast, the majority of cohesin proteins

remain along the chromosome arms at MI.

We hypothesized that MI oocytes, which have more cohesin proteins along the

chromosome axis, would have higher chromosome stiffness than MII oocytes. Cohesins are

components of axial elements during meiosis and there are some meiosis-specific cohesins, such

as REC8, STAG3, RAD21L [34]. We sought to investigate the effects of these cohesins on

chromosome stiffness by utilizing $Rec8^{-/-}$, $Stag3^{-/-}$, and $Rad21l^{-/-}$ mutant mice. These cohesins are

essential for sister chromatid cohesion, and their absence inhibits spermatogenesis, resulting in

arrested spermatocytes at the prophase I stage that cannot mature into sperm [35–37]. We

isolated chromosomes from spermatocytes at the prophase I stage and measured the stiffness of

chromosomes.

First, chromosomes were isolated from $Rec8^{-/-}$ spermatocytes. Under phase-contrast

microscopy, we can identify spermatocytes at the prophase I stage by their larger size, round

shape, and the presence of thick chromosome threads (Supplement Fig.2 Left). Thick

chromosome threads on the spermatocytes indicated that the chromosomes finished chromosome

condensation during prophase I. At prophase I stage, the chromosomes formed a bundle,

allowing identifying the spermatocyte stage (Supplement Fig.2 Middle). We then isolated

chromosomes from the bundle and held it with two pipettes (Supplement Fig.2 Right). Following

the micromanipulation protocols, we stretched the chromosome and measured its stiffness [38].

Surprisingly, we did not observe a significant difference in chromosome stiffness between wild

type (WT) control and $Rec8^{-/-}$ mutant (2710 ± 610 Pa in WT spermatocyte versus 2580 ± 620 Pa

in $Rec8^{-/-}$ spermatocyte, P=0.8884) (Figure 3B). Therefore, we concluded that $Rec8^{-/-}$ did not

contribute to chromosome stiffness. To further investigate the effects of meiosis-specific

cohesins, we tested $Stag3^{-/-}$ and $Rad21l^{-/-}$ mutants using the same method (Figure 3A and

Supplement Fig.3). Similar to the results of $Rec8$ mutant, we did not observe a significant

difference in chromosome stiffness for either $Stag3$ knockout mutant (2710 ± 610 Pa in WT

spermatocyte versus 2240 ± 210 Pa in $Stag3$ mutant spermatocyte, P=0.4533) or $Rad21l^{-/-}$ (2710

± 610 Pa in WT spermatocyte versus 2050 ± 370 Pa in $Rad21l^{-/-}$ spermatocyte, P=0.3514)

(Figure 3B). We concluded that meiosis-specific cohesins do not contribute to chromosome

stiffness.

Chromosomes from old MI oocytes have higher stiffness than those from young MI oocytes

To further investigate other factors influencing chromosome stiffness, we also examined

the effects of age on chromosome stiffness. Initially, we hypothesized that chromosome from

aged oocytes would be less stiff because it has been previously shown that aging is associated

with decreased levels of cohesin, such as REC8, in chromosomes [39,40]. To test this hypothesis,

we isolated chromosomes from MI oocytes of 48 week-old mice (equivalent to 40 years old in

humans) and compared them to chromosomes from 3 to 4 week-old mice (Figure 4A). We

measured the stiffness of chromosomes from both young and old mice and we found that

chromosomes from old mice were much stiffer than those from young mice (8150 ± 1590 Pa in

old MI oocyte versus 3790 ± 700 Pa in young MI oocyte, P=0.0150) (Figure 4B). This result

again supports the conclusion that cohesins are not a main contributor to chromosome stiffness.

We also compared our findings with previous research which identified an increased

chromosome stiffness in aged MII oocytes when compared to their younger counterparts, which

is consistent with our own observations in MI oocytes [27]. Our study revealed that the doubling

force of 3-4 weeks old MII oocytes is measured at 510 ± 50 pN (see Supplemental Figure 1).

This should be contrasted with a doubling force of 830 ± 100 pN for 6–8 weeks old MII oocytes

[27], reinforcing the trend of increased chromosome stiffness with advancing age.

At the MII stage, most cohesin proteins have already disassociated from chromosome

arms, and only a small amount of cohesin proteins remain to connect sister chromosomes at

centromeres until anaphase II. Therefore, the changes in chromosome stiffness observed in aged

oocytes are not due to alterations in cohesin levels but rather to other factors associated with cell

age. Future investigations are needed to determine these age-related factors that impact

chromosome stiffness.

DNA damage reduces chromosome stiffness in oocytes

The oocytes from older individuals are known to exhibit higher levels of DNA damage

compared to those from younger individuals [41,42]. In response to DNA damage on

chromosomes, numerous DNA repair mechanisms are activated, which recruit various DNA

repair proteins to DNA damage sites [43]. Given this background, we hypothesized that DNA

damage could recruit DNA damage repairing proteins affecting chromosome stiffness. To test

this hypothesis, we used etoposide, a chemotherapy drug that has been used treat a variety of

cancers including testicular and ovarian cancer [44,45]. We treated the oocytes with etoposide to

introduce DNA damage and investigated its impact on chromosome stiffness [41]. We cultured

oocytes from the GV (germinal vesicle) stage for 6 hours to MI stage with etoposide at a

concentration of 50μg/ml. After lysing the MI oocytes and isolating the spindle, we observed that

the etoposide-treated chromosomes were not evenly distributed in the middle of the spindle panel

but rather accumulated into several large clusters (Figure 5A). Similarly, DAPI staining showed

that the etoposide-treated oocytes exhibited disrupted chromosome condensation and alignment

compared to the control group, where well-aligned and individualized chromosomes were

observed (Figure 5B).

We then isolated the chromosomes and measured their stiffness, finding that the stiffness

of the chromosomes from etoposide-treated MI oocytes was significantly lower than that from

the control MI oocytes (1710 ± 430 Pa versus 3780 ± 700 Pa, respectively, P=0.0245) (Figure

5C). Therefore, DNA damage can reduce chromosome stiffness in oocytes instead of making it

stiffer. This result also suggests that the high chromosome stiffness of old oocytes does not

attribute to DNA damage induced factors.

Discussion

Despite many years of research, our understanding of the chromosome structure is not yet

fully understood. Measuring mechanics is a useful approach for studying the physical properties

of chromosomes. In the past, mitotic chromosome stiffness has been measured and it was found

that mitotic chromosomes had a chromatin network structure [9]. Several factors, including

condensin, have been found to affect chromosome stiffness [46].

We aimed to identify whether cell stages and aging could influence chromosome

stiffness. Despite extensive research on mitotic chromosomes, meiotic chromosomes have been

relatively understudied. Recently, we found a huge stiffness difference between mitotic and

meiotic chromosomes in spermatocytes [13]. Specifically, meiotic chromosomes were found to

be ten times stiffer than their mitotic counterparts in spermatocytes, which may be due to

different chromosome condensation or folding mechanisms. During prophase I of meiosis, the

synaptonemal complex zips up the two homologous chromosomes, and most of its components

are meiosis-specific [47]. Previously, we investigated whether SYCP1, an essential component

of the synaptonemal complex that connects the lateral and central elements, contributes to the

high chromosome stiffness in spermatocytes [13]. However, we found that the transverse

filament SYCP1 does not impact chromosome stiffness in spermatocytes.

In this study, we first confirmed that high chromosome stiffness also occurred in MI

oocytes. The central element has already dissociated from chromosomes before the MI stage,

further supporting our conclusion that SYCP1 does not contribute to chromosome stiffness in

meiotic cells [13]. Next, we also measured the chromosome stiffness of MII oocytes and found

that MII chromosomes are significantly less stiff than MI chromosomes. This indicates that some

factors in meiosis I but not in meiosis II and mitosis contribute to chromosome stiffness. Hence,

we focused on meiosis I-specific factors.

We know that axial elements of the SC still can load on chromosomes even in $Sycp1^{-/-}$

spermatocytes [13]. Cohesin proteins are components of the axial element and load onto

chromosomes during DNA replication. Some meiosis-specific cohesins, such as REC8, could

persist on chromosome arms until meiosis I anaphase. We hypothesized that the cohesin rather

than the central element increases the meiotic chromosome stiffness. To test this, we measured

the stiffness of chromosomes isolated from cohesin mutant spermatocytes. However, we found

no difference in chromosome stiffness between the wild type and the *Rec8*[-/-], *Rad21l*[-/-] and *Stag3*[-/-]

mutants. Therefore, we concluded that the meiosis-specific cohesins do not contribute to high

chromosome stiffness.

It is also interesting to investigate the effect of age on chromosome stiffness. We isolated

chromosomes from old MI oocytes and observed that aged oocytes had higher chromosome

stiffness compared to the young oocytes. This result indicated that the chromosome stiffness is

also altered during aging. It also provided further support for our conclusion that cohesins do not

make meiotic chromosomes stiffer as cohesin protein levels reduce with age. However, how can

meiotic chromosomes become stiffer even in the presence of hundreds of DNA breaks *in vivo*?

The answer may lie in the DNA-damage-repairing proteins that are recruited to or near the DNA

damage sites to repair the DNA damage and maintain genome integrity. Therefore, we

hypothesized that DNA repair proteins contribute to meiotic chromosome stiffness. To test this,

we used etoposide to induce the DNA damage in MI oocytes. Etoposide could increase the levels

of TOP2–DNA covalent complexes, which generate DNA damage [48]. Surprisingly, we found

that chromosome stiffness decreased instead of increased. This result is consistent with previous

study that DNA double-strand breaks can reduce chromosome stiffness *in vitro* [9]. Again it

confirmed that meiotic chromosomes integrity relies on DNA continuity rather than the linkage

of chromosome axis components [38].

Since DNA repair proteins were not able to increase chromosome stiffness, there must be

other factors that result in high chromosome stiffness. During senescence, the amount of nuclear

protein increases by around two-fold, even though the cohesin level is decreased [40,49]. Thus, it

is possible that some nuclear proteins regulate chromosome stiffness with age. However, further

investigation is needed to determine which nuclear proteins contribute to chromosome stiffness.

It is also possible that histone methylation can influence chromosome stiffness because the level

of histone methylation is proportional to chromosome stiffness [38]. Given that histone

methylation levels are altered during aging, it is also plausible that some histone

methyltransferases and demethylases might regulate chromosome stiffness [50,51]. Moreover,

the methylation plays a very important role during synapsis and recombination especially H3K4,

H3K9 and H3K36 which have been found on meiotic chromatin from leptotene stage [52–54].

Therefore, it is very likely that histone methylation is involved in regulating chromosome

stiffness in different cells.

No matter what factors involved, they must alter chromosome organization to regulate

chromosome stiffness. Therefore, chromosome stiffness is an important parameter for studying

chromosome structure. Defective chromosome organization is often related to various diseases,

such as cancer, infertility, and senesce [55–57]. By using the micromanipulation system to study

chromosome stiffness, we can potentially identify the underlying mechanisms that lead to or

associate with these chromosome defects. This knowledge can then be used to develop new

treatments and therapies for these conditions.

One well-established characteristic of aged oocytes is a higher rate of aneuploidy

compared to young oocytes [58,59]. The majority of aneuploidy can be traced to meiosis I

because anaphase I is an error-prone process [60]. It would be intriguing to investigate whether

the heightened chromosome stiffness plays a role in contributing to this issue. One common

cause of aneuploidy at this stage is lagging chromosomes, which refers to one or more

chromosomes lagging behind others when homologous chromosomes separate during anaphase

[61]. Chromosome stiffness may be related to the occurrence of lagging chromosomes, as

chromosomes are supposed to be separated by pulling forces from microtubules [62,63]. During

anaphase I, homologous chromosomes, rather than sister chromatids, segregate. Thus, the

chromosome fragment between the centromere and crossover must withstand the pulling force

applied to the centromere by the spindle [62]. Chromosome fragments must be stiff enough to

avoid chromosome breakage caused by the pulling force from the spindle. If the chromosome is

too stiff, the tightly connected homologous chromosomes may be hard to separate, causing

aneuploidy consequently. In this case, the high rate of aneuploidy may be related to chromosome

stiffness change and minimizing the factors altering chromosome stiffness may help to reduce

the aneuploidy rate. Moreover, investigating the relationship between chromosome stiffness and

the occurrence of lagging chromosomes may provide further insights into the mechanisms

underlying aneuploidy in oocytes in the future.

Materials and Methods

Animals

Wild-type CD-1 mice (Charles River Laboratories, Wilmington, MA) were used for all

chromosome measurements except for the *Rec8*, *Stag3*, and *Rad21l* mutants. All the mutants are

in C57BL6/J background. All mice were housed in Pancoe CCM rooms of Northwestern

University under 12 h dark/12 h light cycles at $22 \pm 1°C$. The mice were provided with sufficient

food and water. All the handling and procedures of animal experiments were approved by the

Institutional Animal Care and Use Committee at Northwestern University.

Mouse oocyte *in vitro* culture

Oocytes from 3-4 weeks old (48 weeks old for aging study) female mice were used for *in*

vitro culture. For the oocyte *in vitro* culture, both ovaries were dissected from mice immediately

and washed with M2 medium. Then ovaries were set in 100 μM IBMX in M2 medium that was

prewarmed to 37 °C. Ovaries were punched by sterilized needles to release cumulus-oocyte

complexes (COCs). Next, COCs were pipetted in and out several times using a mouth pipette to

remove the cumulus cells surrounding the oocyte under dissection microscope. In this case, the

denuded oocytes at germinal vesicle (GV) stage could be obtained and set in prewarmed M2

medium with 100 μM IBMX. The oocytes with irregular shape and abnormal size were

discarded. The healthy oocytes were selected and washed three times in culture medium (M16)

before transferring. In the end, oocytes were transferred to a small drop of M16 culture medium

covered with mineral oil in a petri dish. The petri dish was incubated at 37°C in a 5% CO2

incubator. Following the experimental plan, the oocytes were cultured for 6 h to reach the MI

stage or 14 h to reach the MII stage, respectively. After culture, the oocytes were rinsed 3 times

with M2 medium and transferred to Tyrode's solution to remove zona pellucida. The oocytes

were transferred to PBS (phosphate buffered saline) solution for chromosome stiffness

measurement.

Mitotic cell culture

Mouse embryonic fibroblasts (MEFs) were selected for mitotic chromosome

measurement. Cells were cultured in DMEM (Corning) with 10% fetal bovine serum (FBS)

(HyClone) and 1% 100x penicillin streptomycin (Corning). The plates were incubated at 37 °C

and 5% CO_2 and passaged every 3-5 days, no more than 20 generations. Before the

measurement, cells would be transferred to prepared culture wells that were made by fixing a

rubber ring on a coverslip using wax. Culture media (2ml) was added into each culture dish well.

The cells were cultured in culture dish wells for 1-3 days to allow for cells to attach to the

coverslip. The chromosome measurement was done in the culture dish well.

Spermatocyte preparation

Testes were dissected from the adult mice [13]. After removing tunica albuginea, a small

cluster of seminiferous tubules were dissected and rinsed in a culture well containing 2 ml of

PBS so that some spermatocytes could release from the seminiferous tubules and enter PBS.

Spermatocytes would settle down to the bottom of the culture well and were used for

chromosome measurement.

Chromosome isolation

Hold pipette, force pipette, and stiff pipette were prepared to grab chromosomes. They

were forged by using a micropipette puller (Sutter P-97) and were cut into suitable sizes [13].

Chromosomes were isolated and measured under the inverted microscope (IX-70; Olympus)

with a 60x 1.42 NA oil immersion objective and a 1.5x magnification pullout. The whole

experiment was conducted at room temperature within 3 h after the culture well was set up on

the microscope due to water evaporation. MEF cells and spermatocytes were identified by eyes

under the brightfield of a microscope. Then these cells were lysed with 0.05% Triton X-100 in

PBS using a spray pipette to remove the cell membrane. After lysis, meiotic nuclei or mitotic

chromosome bundles came out of the cell and were grabbed by a pipette filled with PBS. The

force pipette was used to attach a single chromosome and pull it out of the meiotic nuclei or

mitotic chromosome bundles. Then, the stiff pipette was used to grab the other end of the

chromosome and the meiotic nuclei and mitotic chromosome bundles was moved away from the

grabbed chromosome. One difference between oocytes and other types of cells was that the

spindle with chromosomes could be isolated as a whole unit. A force pipette was inserted into

the spindle to capture chromosomes and drag them out from the spindle by the stiff pipette. After

isolation, the chromosomes were ready to conduct chromosome measurement.

Chromosome stiffness measurement and calculation

After holding the chromosome between the force and stiff pipettes, the chromosome was

stretched by moving the stiff pipettes perpendicularly and the process was recorded by

LabVIEW software[13]. Before stretching, the picture of the chromosome was captured and thus

deflection of the force pipette was calculated. The stiff pipette would move around 6.0 μm and

return to its starting position at a constant rate of 0.20 μm/sec in 0.04 μm steps controlled by the

LabVIEW program. The whole measurement process was repeated 6 times. The location of the

stiff pipette and force pipette were also recorded. From the picture, the original length and

diameter "r" would be obtained using ImageJ scan. Cross section area was calculated as equation

$\pi r^2/2$. The force constant of the force pipette was calibrated by a pipette whose spring constant

was already measured[13]. The chromosome spring constant was obtained by deflection of the

force pipette multiplied by force pipette constant, then divided by the distance between the

pipettes. Young's modulus was calculated according to the formula $E=(F/A)/(\Delta L/L_0)$. E was the

Young's modulus, F was the force given to stretch the chromosome, A was the cross section

area, ΔL was the length change of the chromosome, and L_0 was the original length of the

chromosome.

Chromosome staining

After harvest, oocytes were fixed using 4% (W/V) paraformaldehyde in PBS (phosphate

buffered saline) solution for 30 min at room temperature. Then the oocytes were washed three

times in washing buffer (0.1% tween-20 and 0.01% Triton X-100 in PBS). Next, oocytes were

permeabilized using 0.5% Triton X-100 in PBS for 20 min at RT. After that, the oocytes were

transferred into 3% BSA (bovine serum albumin) for blocking for one hour at RT. After

blocking, the oocytes were washed three times, then oocyte were counter-stained with 1 μg/ml of

(DAPI) 4′,6-diamidine-2′-phenylindole dihydrochloride for 10 min at RT. Finally, after washing

oocytes twice with washing buffer, they were mounted on glass slides with 80% glycerol. The

prepared slides were examined with a Nikon A1R confocal microscope and processed using

NIS-Elements software.

Acknowledgments

We would like to thank all the members of Qiao and Marko groups for their technical support

and comments on this manuscript.

This work was supported by National Institutions of Health (NIH) R00 HD082375 and R01

GM135549, UM1 HG011536, R01 GM105847, R01 GM117155, U54 CA193419, and U54

CA203000.

Author contributions: H. Qiao and J. Marko conceived the project and designed the experiments;

N. Liu analyzed the data and performed all the experiments; P. Jordan gave experimental

direction and provided cohesin mutant mice for the study. W. Qiang provided technical

assistance in the experiments. P. Jordan and W. Qiang helped manuscript editing; and N. Liu, H.

Qiao and J. Marko wrote the manuscript. All authors read and approved the final manuscript.

Disclosures: The authors declare no competing interests exist.

References

[1]   P. Vagnarelli, Chromatin reorganization through mitosis, 1st ed., Elsevier Inc., 2013.

        https://doi.org/10.1016/B978-0-12-410523-2.00006-7.

[2]   T. Potapova, G.J. Gorbsky, The consequences of chromosome segregation errors in

        mitosis and meiosis, Biology (Basel). 6 (2017) 1–33.

        https://doi.org/10.3390/biology6010012.

[3]   L. Shaffer, Theisen, Disorders caused by chromosome abnormalities, Appl. Clin. Genet.

        (2010) 159. https://doi.org/10.2147/tacg.s8884.

[4]   M. Kschonsak, C.H. Haering, Shaping mitotic chromosomes: From classical concepts to

        molecular mechanisms, BioEssays. 37 (2015) 755–766.

        https://doi.org/10.1002/bies.201500020.

[5]   B. Alberts, A. Johnson, J. Lewis, M. Raff, K. Roberts, P. Walter, Chromosomal DNA and

        Its Packaging in the Chromatin Fiber, (2002).

        https://www.ncbi.nlm.nih.gov/books/NBK26834/ (accessed February 20, 2023).

[6]   S.A. Grigoryev, C.L. Woodcock, Chromatin organization - The 30nm fiber, Exp. Cell

        Res. 318 (2012) 1448–1455. https://doi.org/10.1016/j.yexcr.2012.02.014.

[7]   F. Song, P. Chen, D. Sun, M. Wang, L. Dong, D. Liang, R.M. Xu, P. Zhu, G. Li, Cryo-

EM study of the chromatin fiber reveals a double helix twisted by tetranucleosomal units,

Science (80-.). 344 (2014) 376–380. https://doi.org/10.1126/science.1251413.

[8]   T.O. Tolsma, J.C. Hansen, Post-translational modifications and chromatin dynamics,

Essays Biochem. 63 (2019) 89–96. https://doi.org/10.1042/EBC20180067.

[9]   M.G. Poirier, J.F. Marko, Mitotic chromosomes are chromatin networks without a

mechanically contiguous protein scaffold, Proc. Natl. Acad. Sci. U. S. A. 99 (2002)

15393–15397. https://doi.org/10.1073/pnas.232442599.

[10]   W. Antonin, H. Neumann, Chromosome condensation and decondensation during mitosis,

Curr. Opin. Cell Biol. 40 (2016) 15–22. https://doi.org/10.1016/j.ceb.2016.01.013.

[11]   J.F. Marko, E.D. Siggia, Polymer models of meiotic and mitotic chromosomes, Mol. Biol.

Cell. 8 (1997) 2217–2231. https://doi.org/10.1091/mbc.8.11.2217.

[12]   D. Koshland, A. Strunnikov, Mitotic chromosome condensation, Annu. Rev. Cell Dev.

Biol. 12 (1996) 305–333. https://doi.org/10.1146/annurev.cellbio.12.1.305.

[13]   R.J. Biggs, N. Liu, Y. Peng, J.F. Marko, H. Qiao, Micromanipulation of prophase I

chromosomes from mouse spermatocytes reveals high stiffness and gel-like chromatin

organization, Commun. Biol. 3 (2020) 1–7. https://doi.org/10.1038/s42003-020-01265-w.

[14]   C. Heyting, Synaptonemal complexes: Structure and function, Curr. Opin. Cell Biol. 8

(1996) 389–396. https://doi.org/10.1016/S0955-0674(96)80015-9.

[15]   N.M. Hollingsworth, A new role for the synaptonemal complex in the regulation of

meiotic recombination, Genes Dev. 34 (2020) 1562–1564.

https://doi.org/10.1101/gad.345488.120.

[16]   J.M. Peters, A. Tedeschi, J. Schmitz, The cohesin complex and its roles in chromosome

biology, Genes Dev. 22 (2008) 3089–3114. https://doi.org/10.1101/gad.1724308.

[17]   K.I. Ishiguro, J. Kim, S. Fujiyama-Nakamura, S. Kato, Y. Watanabe, A new meiosis-

500   specific cohesin complex implicated in the cohesin code for homologous pairing, EMBO

501   Rep. 12 (2011) 267–275. https://doi.org/10.1038/embor.2011.2.

502   [18]   H. Zheng, W. Xie, The role of 3D genome organization in development and cell

503   differentiation, Nat. Rev. Mol. Cell Biol. 20 (2019) 535–550.

504   https://doi.org/10.1038/s41580-019-0132-4.

505   [19]   M. Rong, A. Matsuda, Y. Hiraoka, J. Lee, Meiotic cohesin subunits RAD21l and REC8

506   are positioned at distinct regions between lateral elements and transverse filaments in the

507   synaptonemal complex of mouse spermatocytes, J. Reprod. Dev. 62 (2016) 623–630.

508   https://doi.org/10.1262/jrd.2016-127.

509   [20]   M.R. Gyuricza, K.B. Manheimer, V. Apte, B. Krishnan, E.F. Joyce, B.D. McKee, K.S.

510   McKim, Dynamic and Stable Cohesins Regulate Synaptonemal Complex Assembly and

511   Chromosome Segregation, Curr. Biol. 26 (2016) 1688–1698.

512   https://doi.org/10.1016/j.cub.2016.05.006.

513   [21]   M.A. Shammas, Telomeres, lifestyle, cancer, and aging, Curr. Opin. Clin. Nutr. Metab.

514   Care. 14 (2011) 28–34. https://doi.org/10.1097/MCO.0b013e32834121b1.

515   [22]   F. Faggioli, J. Vijg, C. Montagna, Chromosomal aneuploidy in the aging brain, Mech.

516   Ageing Dev. 132 (2011) 429–436. https://doi.org/10.1016/j.mad.2011.04.008.

517   [23]   M.K. Thakur, Covalent modifications of chromosomal proteins during aging, Arch.

518   Gerontol. Geriatr. 2 (1983) 1–10. https://doi.org/10.1016/0167-4943(83)90012-2.

519   [24]   G. Oliviero, S. Kovalchuk, A. Rogowska-Wrzesinska, V. Schwämmle, O.N. Jensen,

520   Distinct and diverse chromatin proteomes of ageing mouse organs reveal protein

521   signatures that correlate with physiological functions, Elife. 11 (2022) 1–27.

https://doi.org/10.7554/eLife.73524.

[25] L.M. Lister, A. Kouznetsova, L.A. Hyslop, D. Kalleas, S.L. Pace, J.C. Barel, A. Nathan,

V. Floros, C. Adelfalk, Y. Watanabe, R. Jessberger, T.B. Kirkwood, C. Höög, M. Herbert,

Age-related meiotic segregation errors in mammalian oocytes are preceded by depletion

of cohesin and Sgo2, Curr. Biol. 20 (2010) 1511–1521.

https://doi.org/10.1016/j.cub.2010.08.023.

[26] S. Nakagawa, G. FitzHarris, Intrinsically Defective Microtubule Dynamics Contribute to

Age-Related Chromosome Segregation Errors in Mouse Oocyte Meiosis-I, Curr. Biol. 27

(2017) 1040–1047. https://doi.org/10.1016/j.cub.2017.02.025.

[27] J.E. Hornick, F.E. Duncan, M. Sun, R. Kawamura, J.F. Marko, T.K. Woodruff, Age-

associated alterations in the micromechanical properties of chromosomes in the

mammalian egg, J. Assist. Reprod. Genet. 32 (2015) 765–769.

https://doi.org/10.1007/s10815-015-0453-y.

[28] M. Sun, R. Biggs, J. Hornick, J.F. Marko, Condensin controls mitotic chromosome

stiffness and stability without forming a structurally contiguous scaffold, BioRxiv. (2018)

277–295. https://doi.org/10.1101/384982.

[29] A.R. Strom, R.J. Biggs, E.J. Banigan, X. Wang, K. Chiu, C. Herman, J. Collado, F. Yue,

J.C.R. Politz, L.J. Tait, D. Scalzo, A. Telling, M. Groudine, C.P. Brangwynne, J.F. Marko,

A.D. Stephens, Hp1α is a chromatin crosslinker that controls nuclear and mitotic

chromosome mechanics, Elife. 10 (2021) 1–30. https://doi.org/10.7554/eLife.63972.

[30] A. Hochwagen, Meiosis, Curr. Biol. 18 (2008) 641–645.

https://doi.org/10.1016/j.cub.2008.06.013.

[31] F.A.T. De Vries, E. De Boer, M. Van Den Bosch, W.M. Baarends, M. Ooms, L. Yuan,

J.G. Liu, A.A. Van Zeeland, C. Heyting, A. Pastink, Mouse Sycp1 functions in

synaptonemal complex assembly, meiotic recombination, and XY body formation, Genes

Dev. 19 (2005) 1376–1389. https://doi.org/10.1101/gad.329705.

[32] B.E. McGuinness, T. Hirota, N.R. Kudo, J.M. Peters, K. Nasmyth, Shugoshin prevents

dissociation of cohesin from centromeres during mitosis in vertebrate cells, PLoS Biol. 3

(2005) 0433–0449. https://doi.org/10.1371/journal.pbio.0030086.

[33] J. Lee, T. Iwai, T. Yokota, M. Yamashita, Temporally and spatially selective loss of Rec8

protein from meiotic chromosomes during mammalian meiosis, J. Cell Sci. 116 (2003)

2781–2790. https://doi.org/10.1242/jcs.00495.

[34] A. Ward, J. Hopkins, M. Mckay, S. Murray, P.W. Jordan, Genetic Interactions Between

the Meiosis-Specific Cohesin Components, STAG3, REC8, and RAD21L., G3

(Bethesda). 6 (2016) 1713–24. https://doi.org/10.1534/g3.116.029462.

[35] H. Xu, M.D. Beasley, W.D. Warren, G.T.J. van der Horst, M.J. McKay, Absence of

Mouse REC8 Cohesin Promotes Synapsis of Sister Chromatids in Meiosis, Dev. Cell. 8

(2005) 949–961. https://doi.org/10.1016/j.devcel.2005.03.018.

[36] J. Hopkins, G. Hwang, J. Jacob, N. Sapp, R. Bedigian, K. Oka, P. Overbeek, S. Murray,

P.W. Jordan, Meiosis-Specific Cohesin Component, Stag3 Is Essential for Maintaining

Centromere Chromatid Cohesion, and Required for DNA Repair and Synapsis between

Homologous Chromosomes, PLoS Genet. 10 (2014) e1004413.

https://doi.org/10.1371/journal.pgen.1004413.

[37] Y. Herrán, C. Gutiérrez-Caballero, M. Sánchez-Martín, T. Hernández, A. Viera, J.L.

Barbero, E. de Álava, D.G. de Rooij, J.Á. Suja, E. Llano, A.M. Pendás, The cohesin

subunit RAD21L functions in meiotic synapsis and exhibits sexual dimorphism in

fertility., EMBO J. 30 (2011) 3091–105. https://doi.org/10.1038/emboj.2011.222.

[38]   R. Biggs, P.Z. Liu, A.D. Stephens, J.F. Marko, Effects of altering histone posttranslational

570   modifications on mitotic chromosome structure and mechanics, Mol. Biol. Cell. 30 (2019)

571   820–827. https://doi.org/10.1091/mbc.E18-09-0592.

572   [39]   L.Y. Tian, L. Zhang, C.Q. Tao, X.Q. Lin, F. Zhang, B. Zhang, Age-cumulative effect of

573   REC8 reduction on meiotic chromosome segregation errors in mice, Reprod. Dev. Med. 5

574   (2021) 193–198. https://doi.org/10.4103/2096-2924.327880.

575   [40]   M. Tsutsumi, R. Fujiwara, H. Nishizawa, M. Ito, H. Kogo, H. Inagaki, T. Ohye, T. Kato,

576   T. Fujii, H. Kurahashi, Age-related decrease of meiotic cohesins in human oocytes, PLoS

577   One. 9 (2014). https://doi.org/10.1371/journal.pone.0096710.

578   [41]   P. Marangos, M. Stevense, K. Niaka, M. Lagoudaki, I. Nabti, R. Jessberger, J. Carroll,

579   DNA damage-induced metaphase i arrest is mediated by the spindle assembly checkpoint

580   and maternal age, Nat. Commun. 6 (2015) 1–10. https://doi.org/10.1038/ncomms9706.

581   [42]   F. Horta, S. Catt, P. Ramachandran, B. Vollenhoven, P. Temple-Smith, Female ageing

582   affects the DNA repair capacity of oocytes in IVF using a controlled model of sperm

583   DNA damage in mice, Hum. Reprod. 35 (2020) 529–544.

584   https://doi.org/10.1093/humrep/dez308.

585   [43]   E. Marcon, P.B. Moens, The evolution of meiosis: Recruitment and modification of

586   somatic DNA-repair proteins, BioEssays. 27 (2005) 795–808.

587   https://doi.org/10.1002/bies.20264.

588   [44]   P.J. Hoskins, K.D. Swenerton, Oral etoposide is active against platinum-resistant

589   epithelial ovarian cancer, J. Clin. Oncol. 12 (1994) 60–63.

590   https://doi.org/10.1200/JCO.1994.12.1.60.

[45] J.D. Hainsworth, F.A. Greco, Etoposide: Twenty years later, Ann. Oncol. 6 (1995) 325–

341. https://doi.org/10.1093/oxfordjournals.annonc.a059180.

[46] M. Sun, R. Biggs, J. Hornick, J.F. Marko, Condensin controls mitotic chromosome

stiffness and stability without forming a structurally contiguous scaffold, Chromosom.

Res. 26 (2018) 277–295. https://doi.org/10.1007/s10577-018-9584-1.

[47] S.G. Gordon, L.E. Kursel, K. Xu, O. Rog, Synaptonemal Complex dimerization regulates

chromosome alignment and crossover patterning in meiosis, PLoS Genet. 17 (2021) 1–23.

https://doi.org/10.1371/journal.pgen.1009205.

[48] D. Menendez, J.R. Anand, C.C. Murphy, W.J. Bell, J. Fu, N. Slepushkina, E. Buehler,

S.E. Martin, M. Lal-Nag, J.L. Nitiss, M.A. Resnick, Etoposide-induced DNA damage is

increased in p53 mutants: identification of ATR and other genes that influence effects of

p53 mutations on Top2-induced cytotoxicity, Oncotarget. 13 (2022) 332–346.

https://doi.org/10.18632/ONCOTARGET.28195.

[49] M. De Cecco, J. Jeyapalan, X. Zhao, M. Tamamori-Adachi, J.M. Sedivy, Nuclear protein

accumulation in cellular senescence and organismal aging revealed with a novel single-

cell resolution fluorescence microscopy assay, Aging (Albany. NY). 3 (2011) 955–967.

https://doi.org/10.18632/aging.100372.

[50] C. Soriano-Tárraga, J. Jiménez-Conde, J. Roquer, Epigenetics and aging, Handb. Nutr.

Diet, Epigenetics. 2 (2019) 1413–1433. https://doi.org/10.1007/978-3-319-55530-0_123.

[51] K. Wang, H. Liu, Q. Hu, L. Wang, J. Liu, Z. Zheng, W. Zhang, J. Ren, F. Zhu, G.H. Liu,

Epigenetic regulation of aging: implications for interventions of aging and diseases, Signal

Transduct. Target. Ther. 7 (2022). https://doi.org/10.1038/s41392-022-01211-8.

[52] L. Wang, Z. Xu, M.B. Khawar, C. Liu, W. Li, The histone codes for meiosis,

Reproduction. 154 (2017) R65–R79. https://doi.org/10.1530/REP-17-0153.

[53]   A. Hochwagen, G.A.A.B. Marais, Meiosis: A PRDM9 Guide to the Hotspots of

Recombination, Curr. Biol. 20 (2010) 271–274. https://doi.org/10.1016/j.cub.2010.01.048.

[54]   N.R. Powers, E.D. Parvanov, C.L. Baker, M. Walker, P.M. Petkov, K. Paigen, The

Meiotic Recombination Activator PRDM9 Trimethylates Both H3K36 and H3K4 at

Recombination Hotspots In Vivo, PLoS Genet. 12 (2016) 1–24.

https://doi.org/10.1371/journal.pgen.1006146.

[55]   S.L. Thompson, D.A. Compton, Chromosomes and cancer cells, Chromosom. Res. 19

(2011) 433–444. https://doi.org/10.1007/s10577-010-9179-y.

[56]   G.L. Harton, H.G. Tempest, Chromosomal disorders and male infertility, Asian J. Androl.

14 (2012) 32–39. https://doi.org/10.1038/aja.2011.66.

[57]   Q. He, B. Au, M. Kulkarni, Y. Shen, K.J. Lim, J. Maimaiti, C.K. Wong, M.N.H. Luijten,

H.C. Chong, E.H. Lim, G. Rancati, I. Sinha, Z. Fu, X. Wang, J.E. Connolly, K.C. Crasta,

Chromosomal instability-induced senescence potentiates cell non-autonomous

tumourigenic effects, Oncogenesis. 7 (2018). https://doi.org/10.1038/s41389-018-0072-4.

[58]   K. Selesniemi, H.J. Lee, A. Muhlhauser, J.L. Tilly, Prevention of maternal aging-

associated oocyte aneuploidy and meiotic spindle defects in mice by dietary and genetic

strategies, Proc. Natl. Acad. Sci. U. S. A. 108 (2011) 12319–12324.

https://doi.org/10.1073/pnas.1018793108.

[59]   J.Y. Ma, S. Li, L.N. Chen, H. Schatten, X.H. Ou, Q.Y. Sun, Why is oocyte aneuploidy

increased with maternal aging?, J. Genet. Genomics. 47 (2020) 659–671.

https://doi.org/10.1016/j.jgg.2020.04.003.

[60]   A. Kolano, S. Brunet, A.D. Silk, D.W. Cleveland, M.H. Verlhac, Error-prone mammalian

female meiosis from silencing the spindle assembly checkpoint without normal

interkinetochore tension, Proc. Natl. Acad. Sci. U. S. A. 109 (2012).

https://doi.org/10.1073/pnas.1204686109.

[61]   K.M. Godek, D.A. Compton, Quantitative methods to measure aneuploidy and

chromosomal instability, 1st ed., Elsevier Inc., 2018.

https://doi.org/10.1016/bs.mcb.2018.03.002.

[62]   E. Duro, A.L. Marston, From equator to pole: Splitting chromosomes in mitosis and

meiosis, Genes Dev. 29 (2015) 109–122. https://doi.org/10.1101/gad.255554.114.

[63]   O. Cohen-Fix, Sister chromatid separation: Falling apart at the seams, Curr. Biol. 10

(2000) 816–819. https://doi.org/10.1016/S0960-9822(00)00799-5.

Tables, Figures and Legends

667
668

669
670
Figure 1. Chromosome isolation from oocytes. (A) Oocytes after zona pellucida removal. Left

panel: MI oocyte. Right panel: MII oocyte with polar body. Scale bar = 10 μm. (B) Spindle

673   isolation process. Left panel: Spindle flowing out of the oocyte after oocyte lysis. Middle panel:

674   isolate a chromosome from a spindle-chromosome complex. Right panel: chromosome captured

675   by two pipettes. Scale bar = 10 μm.

681   **Figure 2. Chromosome stiffness measurement.** (A) Example images of chromosome isolation.

682   Left: a MII oocyte chromosome. Right: a MI oocyte chromosome. Scale bar = 10 μm. (B)

683   Chromosome stiffness comparison between mitotic cells (n=8), MI oocytes(n=8), MII oocytes

684   (n=8) and WT spermatocytes at prophase I stage (n=8).

692   **Figure 3. Chromosome stiffness measurement from meiosis-specific cohesin mutant.** (A)

693   Chromosomes isolation images of *Stag3*[-/-] spermatocytes. Scale bar = 10 μm. (B) Chromosome

694   stiffness comparison between WT spermatocytes at prophase I stage (n=8), $Rec8^{-/-}$ spermatocytes

695   at prophase I stage (n=8), $Stag3^{-/-}$ spermatocytes at prophase I stage (n=9) and $Rad21l^{-/-}$

696   spermatocytes (n=10).

701   **Figure 4. Aged chromosomes have higher chromosome stiffness.** (A) Chromosomes isolation

702   images of an old MI oocyte. Left: old MI oocyte images. Middle: a spindle isolated from the old

703 MI oocyte. Right: a chromosome isolated from the spindle-chromosome complex. Scale bar = 10

704 μm. (B) Chromosome stiffness comparison between the 3-4 weeks old MI oocytes (n=8) and 48

705 weeks old MI oocytes (n=5).

Figure 5. Etoposide treatment reduces chromosome stiffness. (A) Chromosomes isolation

images of a MI oocyte treated with 50μg/ml etoposide. Left: a spindle after cell lysis. Middle: a

spindle captured with pipettes. Right: chromosome isolation. Scale bar = 10 μm. (B)Staining of

control and 50μg/ml etoposide-treated MI oocytes. Scale bar = 10 μm. (C) Chromosome stiffness

comparison between mitotic cells (n=8), MI oocyte control (n=8) and 50µg/ml etoposide-treated

MI oocyte (n=8).

Supplement Fig.1

720 **Supplement Fig.1 Chromosomes stiffness of different types of cells.** Chromosome doubling

721 force comparison between mitotic cells (n=8), MI oocytes (n=8), MII oocytes (n=8) and WT

722 spermatocytes at prophase I stage (n=8).

Supplement Fig.2

730   **Supplement Fig.2 Chromosomes isolation images of *Rec8*^{-/-} spermatocytes.** Chromosomes

731   were isolated from *Rec8*^{-/-} spermatocytes and captured with two pipettes. Scale bars = 10 μm.

Supplement Fig.3

739   **Supplement Fig.3 Chromosomes isolation images of *Rad21l*^{-/-} spermatocytes.** Chromosomes

740   were isolated from *Rad21l*^{-/-} spermatocytes and captured with two pipettes. Scale bars = 10 μm.

749