# De novo detection of somatic variants in long-read single-cell RNA sequencing data

**Arthur Dondi[1,2], Nico Borgsmüller[1,2], Pedro F. Ferreira[1,2], Brian J. Haas[3], Francis Jacob[4], Viola Heinzelmann-Schwarz[4], Tumor Profiler Consortium, Niko Beerenwinkel[1,2,*]**

\* Corresponding author

[1] ETH Zurich, Department of Biosystems Science and Engineering, Schanzenstrasse 44, 4056 Basel, Switzerland

[2] SIB Swiss Institute of Bioinformatics, Schanzenstrasse 44, 4056 Basel, Switzerland

[3] Broad Institute of Massachusetts Institute of Technology and Harvard, Cambridge, Massachusetts, USA

[4] University Hospital Basel and University of Basel, Ovarian Cancer Research, Department of Biomedicine, Hebelstrasse 20, 4031 Basel, Switzerland

# Abstract

In cancer, genetic and transcriptomic variations generate clonal heterogeneity, possibly leading to treatment resistance. Long-read single-cell RNA sequencing (LR scRNA-seq) has the potential to detect genetic and transcriptomic variations simultaneously. Here, we present LongSom, a computational workflow leveraging LR scRNA-seq data to call de novo somatic single-nucleotide variants (SNVs), copy-number alterations (CNAs), and gene fusions to reconstruct the tumor clonal heterogeneity. For SNV calling, LongSom distinguishes somatic SNVs from germline polymorphisms by reannotating marker gene expression-based cell types using called variants and applying strict filters. Applying LongSom to ovarian cancer samples, we detected clinically relevant somatic SNVs that were validated against single-cell and bulk panel DNA-seq data and could not be detected with short-read (SR) scRNA-seq. Leveraging somatic SNVs and fusions, LongSom found subclones with different predicted treatment outcomes. In summary, LongSom enables de novo SNVs, CNAs, and fusions detection, thus enabling the study of cancer evolution, clonal heterogeneity, and treatment resistance.

# Introduction

Cancer cells accumulate genomic variations, such as single-nucleotide variants (SNVs), copy number alterations (CNAs), and gene fusions during their lifetime, leading to subpopulations with distinct genotypes. Together with changes in the tumor microenvironment, genomic variations result in distinct phenotypes, such as expression patterns (Lappalainen et al. 2013). Intratumor heterogeneity, i.e., the existence of cancer subpopulations with distinct genotypes and phenotypes, is presumed to be a leading cause of therapy resistance and one of the main reasons for poor overall survival in cancer patients with metastatic disease (Jamal-Hanjani et al. 2015; Ramón Y Cajal et al. 2020). The adaptive mechanisms underlying therapy resistance are of both genetic (SNVs, CNAs, gene fusions, etc.) and non-genetic (epigenetic, transcriptomic, microenvironment, etc.) origin. The first step to identifying therapy-resistant subclones is to capture those genetic and transcriptomic variants through sequencing (Mansoori et al. 2017; Marine et al. 2020). Unraveling different subpopulations is particularly challenging with bulk techniques; however, the advent of single-cell sequencing technologies has significantly improved our ability to decipher intratumor heterogeneity within complex tissue samples (Dagogo-Jack and Shaw 2018).

In scDNA-seq data, cancer cell subpopulations are inferred from SNVs and CNAs, which are conventionally obtained from exome or whole-genome sequencing approaches (Roth et al. 2016; Duan et al. 2018). In scRNA-seq, gene expression patterns are commonly used to differentiate between cell types or cancer cell subpopulations. However, relying solely on gene-level expression may be insufficient, as cells can express different isoforms, resulting in different phenotypes (Ding et al. 2020). Isoform-specific cancer resistance can be induced, for example, through alternative splicing (Mitra et al. 2009; Chen et al. 2022), polyadenylation (Guo et al. 2022), or large genomic rearrangements leading to gene fusions (Amatu et al. 2016; Lei et al. 2018; (Cesi et al. 2018). These interlinked features need to be examined together, thus requiring complete isoform coverage (Foord et al. 2023). High-throughput

droplet-based scRNA-seq protocols (10X Genomics Chromium) capture reads via their 3'

polyA tails. In short-read (SR) scRNA-seq, this results in a heavy coverage bias towards the

3' ends as only a few hundred base pairs of each molecule are sequenced. Long-read (LR)

scRNA-seq, in contrast, sequences full-length RNA molecules, and thus can access gene

expression at the isoform level (Joglekar et al. 2021; Al'Khafaji et al. 2023; Dondi et al. 2023).

Linking genetic to transcriptomic variations is crucial to understanding treatment resistance in

cancer (Vasan et al. 2019). However, this is challenging with SR sequencing, as genetic

variations are difficult to recover from SR scRNA-seq data due to capture bias, while scDNA-

seq cannot assess gene expression. Recently, DNA-free de novo scRNA SNV (Muyas et al.

2023; Zhang et al. 2023) and CNA (Serin Harmanci et al. 2020); (Gao et al. 2021, 2023) calling

methods were developed for SR sequencing, compensating the 3' capture bias by pooling

large amounts of cells or sequencing at very high read depths per cell. However, SR

sequencing is unsuited to detect isoforms or gene fusions. Because it is less sensitive to

capture bias, we have shown in recent work that LR scRNA-seq is more suited to detect

genetic variations than SR scRNA-seq (Dondi et al. 2023). Furthermore, LR scRNA-seq can

simultaneously detect SNVs, CNAs, fusions, and gene isoform expression in the same cells

(Dondi et al. 2023; Shiau et al. 2023).

In this study, we present LongSom, a computational workflow for calling de novo somatic

SNVs, fusions, and CNAs in LR scRNA-seq, and integrating them to reconstruct the samples'

clonal heterogeneity. Applied to omentum metastasis samples obtained from three chemo-

naive high-grade serous ovarian cancer (HGSOC) patients, we show that LongSom can detect

clinically relevant somatic SNVs validated against scDNA and panel data, whereas SR

scRNA-seq fails to do so. We demonstrate that by leveraging somatic SNVs and fusions,

LongSom can detect subclones with different predicted treatment outcomes, and those

subclones were highly concordant with gene expression clusters and CNAs subclones.

Additionally, we find that tumor microenvironment cells are contaminated by cancer cell-

derived mitochondria.

# Results

## Overview of LongSom

We developed LongSom, a workflow for detecting genetic variants in LR scRNA-seq data without requiring matched DNA sequencing and finding cancer subclones based on these. Briefly, LongSom takes BAM files as input, calls SNVs in pseudo-bulk and fusions and CNAs in single cells with the Trinity Cancer Transcriptome Analysis Toolkit (CTAT, https://github.com/NCIP/Trinity_CTAT), and then reconstructs the clonal heterogeneity using the Bayesian non-parametric clustering method BnpC (Borgsmüller et al. 2020).

LongSom first calls candidate SNV loci in a pseudo-bulk generated by aggregating LR scRNA-seq data from all cells, using CTAT-Mutations (https://github.com/NCIP/ctat-mutations), which we enhanced here for scRNA-seq and long isoform reads (see **Methods**). Next, to distinguish between somatic and germline variants, the variant allele frequency (VAF) is calculated for each candidate locus and each cell, and cells are grouped into cancer or non-cancer cells based on marker-gene expression. SNVs detected across multiple cell types are considered germline polymorphisms. Accordingly, if cancer cells are misannotated as non-cancer cells, SNVs will wrongly be filtered out as germline variants (false negatives). To account for this, LongSom first defines a set of cancer-specific variants (SNVs and fusions). SNVs are defined as cancer-specific if their VAF is high in cancer, low in non-cancer, and, when available, zero in normal sample cells (**Methods**). Fusions are detected using CTAT-LR-fusion (https://github.com/TrinityCTAT/CTAT-LR-fusion) (Qin et al. 2024) and cancer-specific fusions are those expressed in more than 5% of cancer cells and less than 1% of non-cancer cells. LongSom reannotates cells as cancer cells if they harbor at least two cancer-specific variants (**Figure 1, Methods**).

**Figure 1: Overview of LongSom.**

LongSom's methodology for detecting somatic SNVs, fusions, and CNAs and subsequently inferring cancer subclones in LR scRNA-seq individual patients data. (1) SNV candidates are detected from pseudo-bulk samples. (2) Population germline SNVs and SNVs present in normal samples (optional) are filtered out. (3) A cell-SNV matrix based on the remaining SNV candidates is computed. (4) A cell-fusion matrix is computed. (5) Using high-confidence cancer fusions and SNVs, cells are reannotated. (6) Following reannotation, SNVs present in non-cancer cells (germlines) are filtered out. (7) cells are clustered based on somatic fusions and SNVs. In parallel, (8) gene expression per cell is computed, (9) CNAs are detected, (10) cells are clustered based on CNAs, and (11) CNA clones are incorporated to the fusions and SNVs clustered matrix.

After cell reannotation, LongSom performs germline SNV filtering in five steps: (A) It filters SNV loci detected in the matched normal, when available. (B) It filters SNV loci from the gnomAD database (Chen et al. 2024) with a frequency of at least 0.01% in the total population. (C) After cell-type reannotation, it filters SNV loci that were called in more than 1% of the non-cancer cells. (D) SNV loci where less than 1% of the non-cancer cells are covered by at least one read are filtered. This step helps to filter germline SNVs not detected due to low

expression in non-cancer cells. (E) Finally, adjacent SNV loci within a 10,000 bp distance are

filtered, as these are likely to be misalignment artifacts in low-complexity regions. Of note,

steps (C) and (E) are not applied to mitochondrial SNVs. Finally, LongSom keeps somatic loci

that are mutated in a minimum of five cancer cells or 5% of cancer cells (user-defined

parameters) and filters loci matching known RNA-editing sites.

Finally, LongSom infers the clonal structure of the samples using two different approaches.

One approach leverages the detected SNVs and fusions as input for the Bayesian non-

parametric clustering method BnpC (Borgsmüller et al. 2020). The other approach predicts

CNAs based on gene expression in cancer cells and defines subclusters using inferCNV

([https://github.com/broadinstitute/infercnv](https://github.com/broadinstitute/infercnv)) (**Methods**).

## Cell-type reannotation improves somatic SNV detection sensitivity

We applied LongSom to previously published (Dondi et al. 2023) SR and LR scRNA-seq data

of five omentum metastasis samples obtained from three chemo-naive HGSOC patients: P1,

P2, and P3. Three samples were derived from HGSOC omental metastases and two from

matching distal tumor-free omental tissues (normal). After cell-type reannotation (**Methods**),

the reannotated cells were always more similar to the expression-based clustering (Jaccard

similarity score in patient P1: 0.97, P2: 0.99, P3: 0.97) than the previous annotation derived

from marker-gene expression (Jaccard similarity score in patient P1: 0.94, P2: 0.98, P3: 0.76),

supporting the reannotation (**Figure 2a**). We found that 6, 2, and 21% of the cells that we

annotated as cancer were previously annotated as non-cancer cells in the tumor biopsy

samples of patients P1, P2, and P3, respectively (**Figure 2b**). The tumor biopsy of patient P3

had only 10% cancer cells (Dondi et al. 2023), which could explain the high level of cell

misannotation. In the following, cancer or non-cancer cells refer to the reannotated cell types.

After cell-type reannotation and germline polymorphism filtering, we found 32, 50, and 62

somatic SNVs and 4, 16, and 2 somatic fusions in patients P1, P2, and P3, respectively

(**Supplementary Tables 1, 2**). In patient P1, a variant at locus chr21:8455886 was manually

detected as a technical false positive due to mismapping in a highly repetitive region, and it

was excluded from further analyses. Without cell type reannotation, we only found 23/32

(72%), 48/50 (96%), and 19/62 (31%) of those SNVs, in P1, P2, and P3, respectively, and no

additional SNV was discovered (**Figure 2c**). In patient P3, numerous cancer cells were

misannotated as non-cancer cells before reannotation (**Figure 2a,b**), leading to 69% of false

negative somatic SNVs during germline SNV filtering (**Figure 2c**). Cells reannotated from non-

cancer to cancer cells showed a mean VAF across somatic SNVs significantly different from

cells annotated as non-cancer cells in both methods ($P<0.001$ in all patients, two-tailed two-

sample t-test), but not from cells annotated as cancer in both methods ($P>0.05$ in all patients),

thus further supporting the cell-type reannotation (**Figure 2d**). Out of the 144 somatic SNVs

identified, we found 32.6% of variants in or affecting coding regions (2.1% start or stop codon

gain (n=3), 1.4% splice region (n=2), 22.2% missense (n=32), and 6.9% synonymous variants

(n=10)) and 67.4% in non-coding regions (17.4% 3' or 5' UTR (n=25), 40.3% intron (n=48) and

9.7% intergenic variants (n=14)) (**Figure 2e**).

**Figure 2: Cell-type reannotation improves somatic SNVs detection sensitivity**

**a.** UMAP embeddings of LR scRNA-seq expression per patient. Cells are colored by annotation status; light-red cells show cells predicted as non-cancer using marker gene expression based annotation, and cancer using high-confidence somatic variants reannotation **b.** Confusion matrices of cells predicted as cancer or non-cancer using marker genes, and cells reannotated as cancer or non-cancer by LongSom, colored and annotated by the percentage of the total number of cells in each category. E.g. the bottom left square represents cells previously annotated as non-cancer that were reannotated as cancer (false negative cancer cells). **c.** Number of SNVs found per patient, with or without cell type reannotation before filtering germline SNPs. **d.** Boxplots of the mean VAF per covered SNV loci of each cell, per patient, colored by their annotation status. Boxes display the first to third quartile with median as horizontal line, whiskers encompass 1.5 times the interquartile range, and data beyond that threshold

182 is indicated as outliers. P values were calculated using a two-sided Student's t-test between groups

183 and are described with the following symbols: n.s : P > 0.05, *: P ≤ 0.05, **: P ≤ 0.01, ***: P ≤ 0.001. **e.**

184 Waffle plot representing each somatic SNV detected, colored by their genomic region and effect on the

185 coding sequence.

## Validation of LR scRNA-seq-derived SNVs using scDNA-seq data

For validation of the SNVs detected using LongSom, we employed scDNA-seq data from

matched omental metastases for each patient. First, we inferred the cellular copy number

profiles based on the scDNA-seq data and identified fully diploid subclones (likely non-cancer)

and aneuploid subclones (likely cancer) from these (Kuipers et al. 2020) (**Methods**). We found

two aneuploid clones in patient P1, one in P2, and two in P3 (**Figure 3a-c**). In each somatic

locus detected by LongSom, we estimated the mean VAF of diploid and aneuploid scDNA

subclones by generating pseudo-bulks. We assumed that a scDNA subclone supported an

SNV if the mean VAF was greater than 10% at the respective locus.

Overall, 55% (n=79) of the somatic SNVs detected in LR scRNA were found exclusively in

scDNA aneuploid subclones and were therefore likely somatic (**Figure 3d-f**). In all cases

where SNVs were not detected in scDNA aneuploid subclones, the scDNA-seq coverage was

<10x (**Figure 3g**). The 10% (n=15) of SNVs detected in diploid scDNA subclones (suggesting

germline polymorphism) were all in patient P2 except one in patient P1 which was only

supported by one read (**Figure 3h**). No normal LR scRNA-seq sample was available for

Patient P2, and non-cancer cells were mainly T-cells with an overall low gene

expression(Joglekar et al. 2021), possibly explaining why germline SNVs were insufficiently

filtered. This finding highlights the utility of matched normal samples to filter germlines

sufficiently.

**Figure 3: Somatic SNVs detected in scRNA are validated as somatic in scDNA**.

**a,b,c.** scDNA-seq copy number values per subclone in **a.** patient 1, **b.** patient 2, and **c.** patient 3 data. Subclones with multiple copy number alterations are aneuploid (likely cancer), while copy number neutral subclones are diploid (likely non-cancer). **d,e,f** Venn diagrams of somatic SNVs supported (VAF>10%) in scDNA cancer subclones (purple), scDNA non-cancer subclones (green), and both (brown). **g.** scDNA cancer subclones coverage per somatic locus identified in scRNA, categorized by whether they are found mutated in cancer subclones (Yes) or not (No). **h.** Number of mutated reads in

scDNA subclones per somatic loci identified in LR scRNA, categorized by cancer and non-cancer scDNA subclones.

## Somatic mitochondrial reads contaminate tumor microenvironment cells in scRNA-seq and scDNA-seq data

As somatic mitochondrial SNVs can also be used for genotype and clonal reconstruction (Miller et al. 2022), LongSom also detects them. Due to the high mitochondrial RNA expression in cells(Osorio and Cai 2021), somatic mitochondrial SNVs (mtSNVs) were amongst the most covered across all cell types and patients in the HGSOC dataset. We found three somatic mitochondrial SNVs in patient P1 (chrM:3092:T>C, chrM:5179:T>C, chrM:16192:C>T), three in patient P2 (chrM:2573:G>A, chrM:4308:G>A, chrM:16065:G>A), and none in patient P3 (**Supplementary Table 1**).

In patient P1, at locus chrM:3092, all covered cancer cells exhibited a >99% VAF in scRNA data, while non-cancer cells showed heteroplasmy (VAF ranging between 0-40%, with 28% of cells mutated, median VAF when mutated 4%) (**Figure 4a)**. However, all cells from distal samples exhibited a VAF <1% (>99% cells covered), ruling out a germline SNV. We detected the same mutational profile in matching scDNA-seq data: amongst the diploid subclones, the average VAF was 5%, while the average VAF in aneuploid subclones was >99% (**Figure 3a**). At locus chrM:5179, cancer cells were either mutated (n = 44, median VAF = 49.7%) or not (n = 30, median VAF <0.1%) in scRNA-seq data, suggesting the presence of two subclones. In non-cancer cells, the VAF ranged from 0 to 33% (11% cells mutated, median VAF when mutated 3%), and all cells from distal samples exhibited again a VAF <1% (>99% cells covered, **Figure 4b**). In the matched scDNA-seq data, at locus chrM:5179, the VAF of aneuploid subclone C3 (**Figure 3a**) was 42%, while it was <1% in the second aneuploid subclone C1 and 2% in the diploid subclones, confirming the subclone specificity of this somatic SNV (**Figure 4c**).

In Patient P2, locus chrM:2573 showed the same pattern with a mean VAF of 75% in cancer

cells compared to a mean VAF of 2% in non-cancer cells. This SNV was observed in 20% of

the non-cancer cells, with a mean VAF of 5% (SD=3.6)) when mutated (**Figure 4d**). No

matching normal sample was available for this patient. In scDNA, the aneuploid subclone had

a VAF of 19% while the diploid subclone had a VAF of 81%. At locus chrM:4308, cancer cells

had a mean VAF of 97%, and non-cancer cells had a VAF ranging between 0 and 100% (19%

cells mutated, mean VAF when mutated 88% (SD=25)). All cells mutated at this locus had

only one variant allele read (**Figure 4e**).

In summary, we observed mitochondrial SNVs with high VAF in cancer cells and lower (but

non-zero) VAF in non-cancer cells from the same scRNA-seq and scDNA-seq samples.

Remarkably, cells from distal (normal) scRNA samples had a VAF of zero in those loci,

suggesting that the mutated mitochondrial reads found in non-cancer cells originate from

cancer cells. This phenomenon could be explained via biological mechanisms such as

intercellular mitochondrial transfer (Liu et al. 2021), or via technical contaminations such as

mitochondria from dead cancer cells being captured together with non-cancer cells during

single-cell encapsulation. The correlation between the amount of mutated mitochondrial SNV

reads found in cancer and in non-cancer supports the contamination hypothesis (**Figure 4f**).

To account for those contaminations, LongSom does not apply step (E) of germline filtering

(filtering of loci that were called in more than 1% of the non-cancer cells).

**Figure 4: Non-cancer cells are contaminated by cancer cells mitochondria.**

**a,b,** VAF of cells in patient P1 at loci **a.** chrM:3092:C and **b.** chrM:5179:C, categorized by reannotated cell types. Color gradient represents the number of variant allele reads per cell. Cells with more than 100 mutated reads are represented with 100 mutated reads. N refers to the number of cells with at least one read covering the locus. **c.** Number of reads supporting the reference or alternative allele in patient P1's scDNA aneuploid (cancer) subclones C1 and C3 at locus chrM:5179, normalized by the number of cells per subclone. **d,e,** VAF of cells in patient P2 at loci **d.** chrM:2573:C and **e.** chrM:4308:C. **f.** Log aggregated mutated reads in non-cancer cells, as a function of log aggregated mutated reads in cancer cells for loci chrM:3092:C, chrM:5179:C and chrM:16192:T in patient P1, and chrM:2573:C, chrM:4308:C and chrM:16065:A in patient 2.

## LR scRNA-seq enables somatic SNV detection with higher sensitivity than SR scRNA-seq

Next, we aimed to compare LR to SR scRNA-seq to detect SNVs. The HGSOC study had LR and SR scRNA-seq data from the same cells available, and while the SR dataset had 4.3 times more sequenced reads compared to LR (mean 117.4k vs. 26.9k reads per cell), it had 3.5 times fewer mapped bases (mean 16Gb mapped vs. 33Gb mapped) due to shorter read length (**Supplementary Figure 1a,b**). When we applied LongSom to SR scRNA-seq, we found only 4/32 (13%), 9/50 (18%), and 9/62 (15%) somatic SNVs in patients P1, P2, and P3 respectively, and no new SNV was detected (**Supplementary Figure 1c**). Additionally, only 1/4 (25%), 9/17 (53%), and 1/2 (50%) fusions were detected in SR scRNA-seq data from patients P1, P2, and P3, respectively (Qin et al. 2024).

We computed cell-variants sparse matrices from each patient's LR and SR data, using cells as columns and somatic SNVs and fusions as rows (**Methods, Figure 5a-c**). For comparison, we computed the same matrix in each patient using SR scRNA-seq data (**Figure 5d-f**). On average, 13.7% (standard deviation (SD) = 1.2) of the matrix positions had at least one LR coverage, whereas only 4.7% (SD = 1.2) had at least one SR coverage (**Supplementary Figure 1d-f**). However, the coverage depends on the cell type expression, and certain cell types, for example, T cells, rarely express mutated genes (**Figure 5a-f**). In cancer cells, on average 27.9% (SD = 7.5) of the matrix positions were covered by at least one LR, whereas only 8.1% (SD = 0.8) had at least one SR coverage (**Figure 5g-i**). On average, LR covered 94.8% (SD=3) of the positions covered by SR and covered an additional 3.4 times more positions (SD = 0.6), in line with the 3.5 times more bases mapped in LR.

**Figure 5: Patient-specific cell-variant matrices created from LR and SR scRNA-seq.**

**a-f.** Matrices of somatic SNVs and fusions (rows) by single cells (columns) computed using LR sc-RNA-seq from the tumor biopsy of **(a)** patient P1, **(b)** P2 and **(c)** P3, and using SR sc-RNA-seq of **(d)** patient P1, **(e)** P2, and **(f)** P3, ordered by gene expression-derived cell types. VAF is depicted as a gradient

from white (no mutated reads, VAF=0) to red (only mutated reads, VAF=1). Grey indicates no coverage

in the cell at a given locus. Rows are colored by the scDNA VAF of aggregated diploid and aneuploid

cells at the loci: SNVs with high aneuploid VAF and low diploid VAF are somatic in scDNA data. RNA

fusions do not give a direct indication of the DNA breakpoint, thus we could not assess their presence

in scDNA data, and they appear in pink. Columns are colored by marker-expression-derived cell-types

(top row) and cell-types reannotated by LongSom (bottom row) **d,e,f.** Venn diagram of matrices'

positions covered in the cancer cells in **(h)** patient P1, **(i)** patient P2, and **(i)** patient P3, colored by

sequencing data modality. Total positions equal n variants x m cancer cells. Blue indicates positions

with coverage >0 in LR and 0 in SR. Red indicates positions with coverage 0 in LR and coverage >0 in

SR. Purple indicates positions with coverage >0 in both LR and SR. Grey indicates positions with

coverage 0 in both LR and SR.

## LongSom detects panel-validated resistance-associated variants

The three patients also underwent bulk panel DNA sequencing (Methods), where 29 SNVs

were found (**Supplementary Table 3**). All three patients had at least one somatic SNV called

in *TP53* (including a variant introducing a stop codon in patient P3) with a VAF >30%, and

patient P1 had a second *TP53* SNV detected with VAF 1%. LongSom detected all *TP53*

somatic SNVs with VAF >30% in LR scRNA-seq. The remaining SNVs detected in the panel

were not retained with our method for the following reasons: they were either germline variants

found in normal and non-cancer cells (n=5), detected in cancer but with insufficient coverage

in non-cancer cells (n=3), detected but not in enough cancer cells (n=7), not detected despite

sufficient coverage (n=3), or not covered (n=8) (**Supplementary Table 3**). Overall, 62% of the

SNVs detected in the panel also found support in scRNA data. Of note, two deletions were

found in panel sequencing, and they were detected manually in the LR scRNA-seq data.

LongSom detected none of the panel SNVs in SR scRNA-seq data.

In addition to *TP53*, LongSom was able to detect SNVs predicted as clinically relevant in genes

not included in the bulk panel (Methods). In patient P1, we found missense variants predicted

as pathogenic in the apoptosis regulator genes *CCAR2* (Arg722Trp) and *FAM129B*

*(*Leu508Pro*)*, and another missense variant in the ferroptosis regulator *ALDH3A2* (Val321Leu)

(**Supplementary Table 4**). *ALDH3A2* is a tumor suppressor in multiple cancers (Xia et al.

2023) and *ALDH3A2, CCAR2,* and *FAM129B* are all associated with treatment resistance in

ovarian cancer (Cheng et al. 2019; Iyer et al. 2022; Dong et al. 2023).  In patient P2, the

chrM:4308 G>A variant was predicted as likely pathogenic. In patient P3, we detected a

missense variant in *AHCY*, as well as a pathogenic *NIF3L1* variant and a missense variant in

the resistance-associated gene *KDM6B (He et al. 2019)*. In SR scRNA-seq data, LongSom

found no clinically relevant variants in patients P1 and P3, and only found chrM:4308 G>A in

patient P2.

## LongSom identifies subclones in LR scRNA-seq data

Next, LongSom inferred the clonal structure of the tumors based on the SNVs and fusions it

detected using BnpC. LongSom also inferred the clonal structure from CNA profiles in the

same cells, using inferCNV (**Supplementary Figure 2a-c, Methods**). We also clustered the

cells based on their gene expression, manually annotated the cancer clusters, and used those

clusters as transcriptomic validation. Finally, we used the subclones inferred from scDNA as

external validation (**Figure 3**).

In patient P2, LongSom found one cancer clone using mutations and fusions, and this clone

coincided very well with the aneuploid CNA clone found in scRNA (Jaccard similarity = 98%)

and the gene-expression-based cancer cluster (Jaccard similarity = 97%, **Supplementary**

**Figures 2b, 3a**). Similarly, in scDNA-seq data we only found one aneuploid CNA clone

(**Figure 3b**). Therefore, all available data modalities point toward a monoclonal cancer

population in this patient. Using SR scRNA-seq, LongSom reconstructed the clonal structure

with lower accuracy than LR due to low coverage (Jaccard similarity BnpC clone - cancer

cluster = 85%, **Supplementary Figure 3b**).

In patient P3, LongSom found one clone, coinciding with the scRNA gene expression-based

cancer cluster (Jaccard similarity = 93%, **Supplementary Figure 4a**), however, two aneuploid

subclones were detected in both scDNA-seq and scRNA-seq data using CNA analysis (**Figure**

**2c, Supplementary Figures 2c,4a**). This difference could be due to the difficulty of calling

subclones in a low number of cancer cells (n=42 after reannotation) or due to inter-sample

heterogeneity. Using SR data, the clustering resulted in a subclone only partially covering the

expression-based cancer cluster, and many individual cells formed singleton subclones

(Jaccard similarity BnpC subclone - cancer cluster = 36%, **Supplementary Figure 4b**).

In patient P1, LongSom found two cancer subclones, referred to as A and B, as well as a

subclone composed of one cell that we assigned to subclone B (**Figure 6a**). The larger

subclone A (n = 40 cells) was predominantly defined by an SNV at locus chrM:5179 and the

smaller subclone B  (n = 34 cells) was mainly defined by the fusion SMG7--CH507-513H4.1.

In expression-based UMAP embedding, cancer cells formed two distinct expression clusters

that highly overlapped the genotypic cancer subclones found based on SNVs and fusions

(Jaccard similarity BnpC subclone A - Expression cluster 1 = 79%, BnpC subclone B -

expression cluster 2 = 72%, **Figure 6 a-d**). CNAs subclones and expression clusters were

also very similar (Jaccard similarity inferCNV subclone A - expression cluster 2 = 89%,

inferCNV subclone B - expression cluster 2 = 85%), likely because they are both derived from

gene expression (**Figure 6a,e, Supplementary Figure 3a).** Clonal assignments based on

SNVs and fusions and on CNA data were also similar (Jaccard similarity subclone A = 72%,

subclone B = 68%. In patient P1's matched scDNA data, we also found two aneuploid (cancer)

subclones based on CNA profiles (**Figure 3a**), and only one of the subclones harbored the

SNV chrM:5179:T>C (**Figure 4c**), concordantly with LR scRNA-seq data. Unfortunately, the

three other subclone-defining SNVs had insufficient scDNA-seq coverage, and we could not

detect any reads supporting the variant allele at those loci, therefore we could not confirm their

subclonality. Using SR scRNA-seq, LongSom also identified cancer subclones in patient P1,

mainly based on chrM:5179 status. However, as the fusion SMG7--CH507-513H4.1 was not

detected in SR, multiple cancer cells clustered with non-cancer cells (Jaccard similarity BnpC

SR subclone A - Expression cluster 1 = 70%, BnpC SR subclone B - expression cluster 2 =

57%, **Supplementary Figure 5**).

## Subclones identified in patient P1 have differing predicted treatment outcomes

To explore the potential therapeutic resistance of the subclones identified in Patient P1 by

LongSom, we investigated the genomic and transcriptomic variations between them. The

*ALDH3A2* pathogenic variant identified earlier was exclusively expressed in subclone A, while

the *CCAR2* pathogenic variant was exclusive to subclone B (**Supplementary Table 4**).

Remarkably, *ALDH3A2* is a ferroptosis inhibitor and its loss of function would lower cisplatin

resistance (Dong et al. 2023), while *CCAR2* is a suppressor of homologous recombination,

and its loss would lead to resistance against PARP inhibitors (Iyer et al. 2022). Therefore,

based on SNVs, subclone A is more likely to be treatment-sensitive, while subclone B is more

likely to be treatment-resistant. Fusions SMG7--CH507-513H4.1 and  GS1-279B7.2--GNG4

were exclusively expressed in subclone B (**Figure 6a**), however, their pathogenicity is difficult

to predict. On the transcriptomic level, Subclone B had notably downregulated expression of

keratin genes *KRT8* and *KRT18*, two epithelial markers used to differentiate HGSOC cells

from non-cancer cells (**Figure 6f**). When compared to cancer subclones in patients P2 and

P3, *KRT8* and *KRT18* were downregulated in subclone B, but not upregulated in subclone A,

thus confirming a downregulation in subclone B  (**Supplementary Figure 6a,b**). It has been

shown in vitro that *KRT8* and *KRT18* have a protective effect against cell death (Bozza et al.

2018), and loss of *KRT8* and *KRT18* leads to increased invasiveness but also cisplatin

sensitivity (Fortier et al. 2013). Subclone B is therefore more likely to be chemosensitive than

subclone A. We additionally investigated differential isoform usage, and while both subclones

were mostly similar, we found a significant difference in *CHPF* (**Figure 6g**), *MYL6*, the tumor

suppressor *BTG2,* and *NUTM2B-AS1* (**Supplementary Figure 6c-e**), however, we could not

predict their pathogenicity. None of the subclone-exclusive variants or isoforms were detected

in Patient P1 SR scRNA-seq data.

**Figure 6: Analysis of intra-tumor heterogeneity using somatic variants detected in LR**

**scRNA-seq in Patient P1.**

**a.** BnpC clustering of single cells from the tumor biopsy of patient P1 (columns) by somatic SNVs and

fusions (rows). VAF is depicted as a gradient from white (no mutated reads, VAF=0) to red (only mutated

reads, VAF=1). Grey indicates no coverage in the cell at a given locus. Rows are colored by the scDNA VAF of aggregated diploid and aneuploid cells at the loci: SNVs with high aneuploid VAF and low diploid VAF are somatic in scDNA data. Fusions appear in pink. Columns are colored from top to bottom by cell types reannotated by LongSom, CNAs subclones, expression clusters, and BnpC clusters **b,c.** UMAP embedding of patient P1 gene expression data, colored by (**b**) cell types reannotated by LongSom and (**c**) subclones inferred from somatic SNVs and fusions. The dashed line indicates the manual separation between cancer clusters 1 and 2. **d,e.** Confusion matrix of cells in each expression-derived cancer cluster (rows) and (**d**) cells in the subclones inferred from BnpC, and (**e**) cells in the subclones inferred from inferCNV (columns), colored by the percentage of the total number of cells in each subclone and annotated by the absolute numbers. **f.** Volcano plot of differentially expressed genes identified between subclones B and A. Keratin genes downregulated in subclone B are annotated. **g.** ScisorWiz representation of CHPF isoforms in subclones A and B. Colored areas are exons, whitespace areas are intronic space, not drawn to scale, and each horizontal line represents a single read colored according to subclones.

# Discussion

SNVs, CNAs, fusions, gene expression, isoforms expression, and the micro-environment composition can all affect cancer treatment outcomes (Marine et al. 2020). Assessing all of these variations simultaneously from a single patient sample is particularly relevant in a clinical setting, where biological material is limited. Here, we show for the first time that this is possible using LR scRNA-seq data and we introduce LongSom, a workflow for detecting de novo somatic SNVs, fusions, and CNAs in LR scRNA-seq. When applied to data from three HGSOC patients, it detected panel- and scDNA-seq-validated SNVs, including clinically relevant *TP53, ALDH3A2,* and *CCAR2* SNVs. By integrating SNVs and fusions, LongSom successfully reconstructed the clonal heterogeneity and linked variants-defined subclones to CNA-defined subclones and gene expression clusters. Finally, in each subclone, we identified differentially expressed genes as well as subclone-specific SNVs with different implications for treatment

resistance. Thus, we demonstrated that LR scRNA-seq is suitable for predicting treatment
outcomes.

The cell-type reannotation step implemented in LongSom, based on the somatic variation
profile of cells, led to the detection of up to 2.4 times more somatic SNVs (patient P3) and
significantly reduced the false-negative rate without sacrificing sensitivity for precision. In
general, cell-type annotation is an open challenge due to overlapping, poorly expressed, or
incomplete marker genes, e.g., in ovarian cancer omentum metastases (Lähnemann et al.
2020; Van Egeren et al. 2021, 2022). Our proposed reannotation can improve existing
methods and shows the potential of combining genomic variants with transcriptomic cell
typing.

To our knowledge, LongSom is the first method combining de novo detection of SNVs and
fusions from the same cell to reconstruct clonal heterogeneity. Besides nuclear SNVs, which
are commonly obtained from RNA (Muyas et al. 2023; Zhang et al. 2023) and DNA seq (Zafar
et al. 2016), LongSom also calls mitochondrial SNVs. In the analyzed HGSOC dataset, the
mitochondrial SNVs were called in most cancer and non-cancer cells, and some high-
confidence fusion calls were expressed in most clones or subclones (P2: IGF2BP2::TESPA1,
P1: SMG7::CH507-513H4.1, etc.), making them ideal variations for cell-type reannotation and
clustering. Furthermore, both can be clinically relevant (Amatu et al. 2016; Lei et al. 2018; Cesi
et al. 2018; Dentro et al. 2021), as we demonstrated in Patient P2. Mitochondrial RNA is
particularly abundant in cancer cells, especially HGSOC (Yuan et al. 2023), and an increasing
number of methods are leveraging them for clonal reconstruction or validation (Kwok et al.
2022; Miller et al. 2022; Gao et al. 2023). However, we demonstrated that mitochondrial SNVs
require special filtering thresholds, as non-cancer cells were frequently contaminated by
cancer mitochondrial reads. We assume that entire cancer mitochondria might contaminate
non-cancer cells, as we observed mitochondrial SNVs in both scRNA and scDNA-seq data.
Whether these mitochondria originate from a biological mechanism, e.g. intercellular transfer

from cancer cells into non-cancer cells for microenvironment revitalization (Liu et al. 2021;

Zampieri et al. 2021), or technical contaminations, e.g. cancer mitochondria encapsulated

jointly with non-cancer cells during single-cell preparation, requires further investigation.

One limitation of this study is the lack of isoform and fusion annotation in the literature,

resulting from the difficulty of detecting them in SR scRNA-seq (Dentro et al. 2021), making it

challenging to explore the biological implications of subclone-specific isoforms or fusions. To

fully exhaust the possibilities of LR scRNA-seq, characterizing more isoforms and fusions will

be necessary in the future. Furthermore, a population-level database dedicated to fusions,

similar to gnomAD (Chen et al. 2024) for SNVs, would be beneficial to filter germline fusions.

We believe that the reliable detection of isoforms and fusions with LR scRNA-seq is the first

step toward this goal.

Despite rapid progress in the LR scRNA-seq field (Al'Khafaji et al. 2023; Dondi et al. 2023;

Joglekar et al. 2023; Marx 2023), multiple technical limitations remain unsolved, limiting the

potential of downstream analysis. First, variant detection remains challenging due to the

sparsity and low coverage of scRNA-seq assays. LongSom excludes SNVs called in non-

cancer cells to filter germline SNVs, possibly leading to false negative somatic SNVs, as

shown by the matched panel-seq. Second, read coverage is also uneven within a transcript,

as transcripts produced by 10X Genomics Chromium remain incomplete on the 5' end (Hsu

et al. 2022). Third, RNA-seq is inherently limited to detecting only expressed SNVs and

fusions. Nevertheless, LongSom detected a large fraction of variants in intronic or even

intergenic regions. Last, indels are the most common source of errors in LR scRNA-seq data,

whereby they are frequently excluded from the analyses, ours included (Shiau et al. 2023). To

further improve the genomic analyses of scRNA-seq data, algorithms for filtering technical

indels while detecting somatic indels are required, especially as technical indels can lead to

false positive somatic SNVs (Ahsan et al. 2021).

In summary, we proposed a workflow for detecting multiple genetic variants (SNVs, CNAs,

fusions) in LR scRNA-seq, enabling clonal heterogeneity reconstruction and clonal genotypes

to treatment-resistance phenotype linkage. LR scRNA-seq provides a unique snapshot of the

cellular mechanisms by capturing multiple genomic and transcriptomic readouts from the

same cell, including expressed isoforms, fusion transcripts, SNVs, and CNAs, more effectively

than with any other sequencing technique. With decreasing costs and increasing data size in

LR scRNA-seq, we envision that LR scRNA-seq will become more common, potentially

facilitating a better understanding of the processes underlying cancer treatment resistance.

LongSom can be a valuable first step in guiding these analyses.

# 509 Methods

## 510 scRNA expression analysis

### 511 Gene expression counts

LR gene expression counts were generated as described in (Dondi et al. 2023). Briefly, we

preprocessed the BAM files using scIsoPrep (https://github.com/cbg-

ethz/scIsoPrep/tree/master) and generated a gene expression-cell matrix. UMI counts were

quality-controlled and cells and genes were filtered to remove mitochondrial and ribosomal

contaminants. Cells for which over 50% of the reads mapped to mitochondrial genes and cells

with fewer than 400 genes expressed were removed. By default, all non-protein-coding genes,

genes coding for ribosomal and mitochondrial proteins, and genes that were expressed in less

than 20 cells were removed. Subsequently, counts were normalized with sctransform

(Hafemeister and Satija 2019), regressing out cell cycle effects and library size as non-

regularized dependent variables.

### Marker gene expression-based annotation

Cells were annotated with scROSHI (Prummer et al. 2023) using ovarian cancer marker gene lists. Marker genes are available at https://github.com/ETH-NEXUS/scAmpi_single_cell_RNA/blob/master/required_files/ovarian/celltype_list_ovarian.gm). We used "HGSOC" labels as cancer cells, and "Mesothelial.cells", "Fibroblast", "T.NK.cells", "B.cells", "Myeloid.cells", "Endothelial.cells" labels as non-cancer cells.

### Clustering and visualization

Similar cells were grouped using Seurat FindClusters (Hao et al. 2024), and clusters with a majority (>90%) of non-cancer cells were grouped together as "non-cancer". The results of the clustering and cell typing are visualized in a low-dimensional representation using Uniform Manifold Approximation and Projection (UMAP).

### Differential gene expression analysis

Differential expression was computed using Seurat FindMarkers (Hao et al. 2024), which uses a Wilcoxon test, corrected for multiple testing using the Bonferroni correction. A threshold of corrected P-value <0.01 and abs(log2(fold change)) >1 was used for significance.

### Differential isoform usage analysis

Isoform classification and quantification were performed using scIsoPrep. Differential isoform testing was performed using a $\chi 2$ test as previously described in Scisorseqr (Joglekar et al. 2021). Differentially used isoforms were visualized using ScisorWiz (Stein et al. 2022).

## Somatic variants calling in LR scRNA-seq data with LongSom

To call somatic variants in LR scRNA-seq, we developed LongSom, a workflow implemented in python3 using Snakemake (Köster and Rahmann 2012) and available at https://github.com/cbg-ethz/LongSom.

### Preprocessing

Long reads with minimal quality Q20 were de-concatenated, adapters were trimmed, demultiplexed, polyA tails were trimmed and finally, UMIs were deduplicated using scIsoPrep (https://github.com/cbg-ethz/scIsoPrep/tree/master) as described in (Dondi et al. 2023). All deduplicated reads belonging to a cell passing filter (cells for which under 50% of the reads mapped to mitochondrial genes and cells with more than 400 genes expressed, see (Dondi et al. 2023), were then pooled together in a pseudo bulk fashion. Gene expression-based cell types were derived from the same work (Dondi et al. 2023).

### SNV calling in LR scRNA-seq data using CTAT-Mutations

First, LongSom calls somatic SNVs in the tumor and (when available) normal biopsy pseudo bulks, using the CTAT mutations pipeline v4.0.0 (https://github.com/NCIP/ctat-mutations/releases/tag/CTAT-Mutations-v4.0.0), which we enhanced to enable compatibility with long reads and report variants according to single cell barcodes. When executed with option --is_long_reads, minimap2 (Li 2018) is used to align long isoform reads to the reference genome hg38 (instead of the STAR aligner used with shorter Illumina RAN-seq), followed by our implementation of the GATK best practices for variant calling using RNA-seq (https://gatk.broadinstitute.org/hc/en-us/articles/360035531192-RNAseq-short-variant-discovery-SNPs-Indels). Loci flagged as RNA-editing sites or with less than 5 reads mutated are filtered out. For generating variant reports at single-cell resolution, allele-supporting reads annotated with cell barcodes and UMIs were captured from the aligned reads, tallied, and reported for downstream integration with cell typing and related metadata.

 Fusion calling in LR scRNA-seq data using CTAT-LR-Fusion

LongSom detects fusions on the single cell level using CTAT-LR-fusion v0.13.0 (

https://github.com/TrinityCTAT/CTAT-LR-fusion/releases/tag/ctat-LR-fusion-v0.13.0) with

standard options (Qin et al. 2024).

Cell-variant matrices construction

LongSom defines three groups based on the marker-expression-based cell types: cancer cells

in the tumor biopsy (in this study, HGSOC cells), non-cancer cells in the tumor biopsy (in this

study: mesothelial cells, fibroblasts, T cells, myeloid cells, B cells, and endothelial cells) and,

if available, normal cells from the normal biopsy. For each of those groups, LongSom builds a

cell-variant matrix with n cells (columns) and m SNVs + p fusions (rows). For SNV rows, the

matrices are filled as follows: if at least one read is covering the locus in a cell, a VAF is

computed for this cell (with a value ranging from 0 to 1), otherwise, the position is a missing

value. A cell is defined as "mutated" at an SNV locus if it has a VAF >= 0.3. For fusion rows,

the matrices are filled as follows: a cell with at least one fusion read is considered "mutated"

for this fusion (value = 1), otherwise, it is a missing value.

Cell type reannotation

To improve the cell type annotation, LongSom defines a set of "high-confidence cancer

variants". To be a "high-confidence cancer variant", an SNV needs to (1) be mutated in more

than 5% of cancer cells, (2) be mutated in >20% of the cancer cells covering the locus, (3)

have >1% of non-cancer cells covering the locus, (4) be mutated in less than 5% of the non-

cancer cells covering the locus, and (5) be mutated in 0 normal cells (optional). For

mitochondrial SNVs, due to the contaminations observed, LongSom does not follow those

rules. Instead, a mitochondrial SNV is a "high-confidence cancer variant" if:

$\% \, of \, cancer \, cells \; mutated \; - \; \% \, of \, noncancer \, cells \; mutated > 20\%$.

To be a "high-confidence cancer variant", a fusion needs to be found in more than 5 cancer

cells and less than 5% of the non-cancer cells. We then reannotated the cell types by defining

as "cancer" any cell mutated in more than two of the "high-confidence cancer variants", and

as "non-cancer" all the other cells in the tumor biopsy.

Final somatic variants call set and matrix

After cell reannotation, LongSom rebuilds two cell-variants matrices using the annotated

cancer and non-cancer labels. Longsom then filters germline polymorphisms (rows) from the

variant matrices in five steps: (A) It filters SNV loci detected in the matched normal, when

available. (B) It filters SNV loci from the gnomAD database (Chen et al. 2024) with a frequency

of at least 0.01% in the total population. (C) After cell-type reannotation, it filters SNV loci that

were called in more than 1% of the non-cancer cells. (D) SNV loci where less than 1% of the

non-cancer cells are covered by at least one read are filtered. This step helps to filter germline

SNVs not detected due to low expression in non-cancer cells. (E) Finally, adjacent SNV loci

within a 10,000 bp distance are filtered, as these are likely to be misalignment artifacts in low-

complexity regions. Of note, steps (C) and (E) are not applied to mitochondrial SNVs. Finally,

LongSom keeps somatic loci that are mutated in a minimum of five cancer cells or 5% of

cancer cells (user-defined parameters).

Cancer and non-cancer cell-variant matrices containing only somatic SNVs and fusions are

then concatenated to create the final cell-variant matrix. SNVs are sorted in decreasing order

by:

$$Diff = mean(\% \ of \ covered \ cancer \ cells \ mutated)$$

$$- \ mean(\% \ of \ covered \ noncancer \ cells \ mutated)$$

Clonal detection based on SNVs and fusions

LongSom uses the cell-variant matrices as input for Bayesian non-parametric clustering

(BnpC) (Borgsmüller et al. 2020) to detect subclones in cancer samples, with arguments -n 16

--steps 1000 --DPa_prior [1,1] --conc_update_prob 0 --param_prior [1,1].

## Clonal detection based on CNAs

LongSom first computes cell-gene matrices using featureCounts from Subread v2.0.6 (https://subread.sourceforge.net/) with parameters -L, using hg38 and gencode v36 as reference. It then uses those matrices as input for inferCNV to detect CNA subclones (https://github.com/broadinstitute/infercnv). For running CreateInfercnvObject, reannotated non-cancer cells are used as a reference, and the parameter min_max_counts_per_cell=c(1e3,1e7) is used. For running inferCNV, the parameters cutoff=0.1 and leiden_resolution=0.01 are used. The CNA profiles displayed in this study are the ones obtained from the Hidden Markov Model learned by inferCNV.

## scDNA analysis

### Preprocessing and clonal reconstruction

Using annotated cell types, we re-computed the cell-variant matrices as well as the percentage of cells mutated, the percentage of cells covered, and the percentage of covered cells mutated, for each locus. We then called the final somatic SNVs set at all loci mutated in more than 5% of cancer cells, mutated in less than 1% of the non-cancer cells (min. 1% non-cancer cells covered), and mutated in no normal cells. We obtained copy number profiles and detected the main clonal structure of samples using SCICoNE (Kuipers et al. 2020). Subclones were considered as cancer subclones if they had an aneuploid CNA profile, and as non-cancer subclones if they had a fully diploid CNA profile.

We investigated all loci from the final somatic SNV set in scDNA subclones in a pseudobulk manner. Cancer subclones were pooled together as well as non-cancer subclones because the coverage was low (<10x per subclone). scDNA subclones with a mean VAF>10% in an SNV locus were considered as supporting the SNV.

## Clinically relevant SNVs

Clinically relevant SNVs were detected using the CTAT-Mutations pipeline (https://github.com/NCIP/ctat-mutations/releases/tag/CTAT-Mutations-v4.0.0). Briefly, an SNV was considered clinically relevant if it completed one of these conditions: it was flagged as pathogenic by ClinVar (Landrum et al. 2014), the CHASMplus (Tokheim and Karchin 2019) P-value was <0.05, the VEST (Carter et al. 2013) P-value was <0.05, or FATHMM (Rogers et al. 2018) flagged it as "CANCER", or "PATHOGENIC".

## Panel validation

To investigate LongSom somatic SNV calls, we used the FoundationOne®CDx targeted NGS panel (Milbury et al. 2022) in matched bulk DNA samples. SNVs detected in the bulk DNA panel but not by LongSom were independently investigated in scRNA-seq data to detect variant allele read support.

# Data availability

The raw sequencing files, as well as the associated analysis files reported in this study are available in the European Genome-phenome Archive (EGA) under the accession number EGAS00001006807. Gencode v36 gene annotation used in this study is available at https://ftp.ebi.ac.uk/pub/databases/gencode/Gencode_human/release_36/gencode.v36.annotation.gtf.gz. All additional information will be made available upon reasonable request to the

authors. Marker genes for cancer and non-cancer cells are available at https://github.com/ETH-NEXUS/scAmpi_single_cell_RNA/blob/master/required_files/ovarian/celltype_list_ovarian.gmx.

# Code availability

LongSom is available at https://github.com/cbg-ethz/LongSom.

# Bibliography

Ahsan MU, Liu Q, Fang L, Wang K. 2021. NanoCaller for accurate detection of SNPs and indels in difficult-to-map regions from long-read sequencing by haplotype-aware deep neural networks. *Genome Biol* **22**: 261.

Al'Khafaji AM, Smith JT, Garimella KV, Babadi M, Popic V, Sade-Feldman M, Gatzen M, Sarkizova S, Schwartz MA, Blaum EM, et al. 2023. High-throughput RNA isoform sequencing using programmed cDNA concatenation. *Nat Biotechnol*.

Amatu A, Sartore-Bianchi A, Siena S. 2016. NTRK gene fusions as novel targets of cancer therapy across multiple tumour types. *ESMO Open* **1**: e000023.

Borgsmüller N, Bonet J, Marass F, Gonzalez-Perez A, Lopez-Bigas N, Beerenwinkel N. 2020. BnpC: Bayesian non-parametric clustering of single-cell mutation profiles. *Bioinformatics* **36**: 4854–4859.

Bozza WP, Zhang Y, Zhang B. 2018. Cytokeratin 8/18 protects breast cancer cell lines from TRAIL-induced apoptosis. *Oncotarget* **9**: 23264–23273.

Carter H, Douville C, Stenson PD, Cooper DN, Karchin R. 2013. Identifying Mendelian disease genes with the variant effect scoring tool. *BMC Genomics* **14 Suppl 3**: S3.

Cesi G, Philippidou D, Kozar I, Kim YJ, Bernardin F, Van Niel G, Wienecke-Baldacchino

A, Felten P, Letellier E, Dengler S, et al. 2018. A new ALK isoform transported by

extracellular vesicles confers drug resistance to melanoma cells. *Mol Cancer* **17**: 145.

Cheng K-C, Lin R-J, Cheng J-Y, Wang S-H, Yu J-C, Wu J-C, Liang Y-J, Hsu H-M, Yu J,

Yu AL. 2019. FAM129B, an antioxidative protein, reduces chemosensitivity by

competing with Nrf2 for Keap1 binding. *EBioMedicine* **45**: 25–38.

Chen C, Zhao S, Zhao X, Cao L, Karnad A, Kumar AP, Freeman JW. 2022.

Gemcitabine resistance of pancreatic cancer cells is mediated by IGF1R dependent

upregulation of CD44 expression and isoform switching. *Cell Death Dis* **13**: 682.

Chen S, Francioli LC, Goodrich JK, Collins RL, Kanai M, Wang Q, Alföldi J, Watts NA,

Vittal C, Gauthier LD, et al. 2024. A genomic mutational constraint map using variation

in 76,156 human genomes. *Nature* **625**: 92–100.

Dagogo-Jack I, Shaw AT. 2018. Tumour heterogeneity and resistance to cancer

therapies. *Nat Rev Clin Oncol* **15**: 81–94.

Dentro SC, Leshchiner I, Haase K, Tarabichi M, Wintersinger J, Deshwar AG, Yu K,

Rubanova Y, Macintyre G, Demeulemeester J, et al. 2021. Characterizing genetic intra-

tumor heterogeneity across 2,658 human cancer genomes. *Cell* **184**: 2239-2254.e39.

Ding S, Chen X, Shen K. 2020. Single-cell RNA sequencing in breast cancer:

Understanding tumor heterogeneity and paving roads to individualized therapy. *Cancer*

*Commun (Lond)* **40**: 329–344.

Dondi A, Lischetti U, Jacob F, Singer F, Borgsmüller N, Coelho R, Tumor Profiler

Consortium, Heinzelmann-Schwarz V, Beisel C, Beerenwinkel N. 2023. Detection of

isoforms and genomic alterations by high-throughput full-length single-cell RNA

sequencing in ovarian cancer. *Nat Commun* **14**: 7780.

Dong H, He L, Sun Q, Zhan J, Li J, Xiong X, Zhuang L, Wu S, Li Y, Yin C, et al. 2023.

Inhibit ALDH3A2 reduce ovarian cancer cells survival via elevating ferroptosis

sensitivity. *Gene* **876**: 147515.

Duan M, Hao J, Cui S, Worthley DL, Zhang S, Wang Z, Shi J, Liu L, Wang X, Ke A, et

al. 2018. Diverse modes of clonal evolution in HBV-related hepatocellular carcinoma

revealed by single-cell genome sequencing. *Cell Res* **28**: 359–373.

Foord C, Hsu J, Jarroux J, Hu W, Belchikov N, Pollard S, He Y, Joglekar A, Tilgner HU.

2023. The variables on RNA molecules: concert or cacophony? Answers in long-read

sequencing. *Nat Methods* **20**: 20–24.

Fortier A-M, Asselin E, Cadrin M. 2013. Keratin 8 and 18 loss in epithelial cancer cells

increases collective cell migration and cisplatin sensitivity through claudin1 up-

regulation. *J Biol Chem* **288**: 11555–11571.

Gao R, Bai S, Henderson YC, Lin Y, Schalck A, Yan Y, Kumar T, Hu M, Sei E, Davis A,

et al. 2021. Delineating copy number and clonal substructure in human tumors from

single-cell transcriptomes. *Nat Biotechnol* **39**: 599–608.

Gao T, Soldatov R, Sarkar H, Kurkiewicz A, Biederstedt E, Loh P-R, Kharchenko PV.

2023. Haplotype-aware analysis of somatic copy number variations from single-cell

transcriptomes. *Nat Biotechnol* **41**: 417–426.

Guo Q, Wang H, Duan J, Luo W, Zhao R, Shen Y, Wang B, Tao S, Sun Y, Ye Q, et al.

2022. An alternatively spliced p62 isoform confers resistance to chemotherapy in breast

cancer. *Cancer Res* **82**: 4001–4015.

Hafemeister C, Satija R. 2019. Normalization and variance stabilization of single-cell

RNA-seq data using regularized negative binomial regression. *Genome Biol* **20**: 296.

Hao Y, Stuart T, Kowalski MH, Choudhary S, Hoffman P, Hartman A, Srivastava A,

Molla G, Madad S, Fernandez-Granda C, et al. 2024. Dictionary learning for integrative,

multimodal and scalable single-cell analysis. *Nat Biotechnol* **42**: 293–304.

He C, Sun J, Liu C, Jiang Y, Hao Y. 2019. Elevated H3K27me3 levels sensitize

osteosarcoma to cisplatin. *Clin Epigenetics* **11**: 8.

Hsu J, Jarroux J, Joglekar A, Romero JP, Nemec C, Reyes D, Royall A, He Y, Belchikov N, Leo K, et al. 2022. Comparing 10x Genomics single-cell 3' and 5' assay in short-and long-read sequencing. *BioRxiv*.

Iyer DR, Harada N, Clairmont C, Jiang L, Martignetti D, Nguyen H, He YJ, Chowdhury D, D'Andrea AD. 2022. CCAR2 functions downstream of the Shieldin complex to promote double-strand break end-joining. *Proc Natl Acad Sci USA* **119**: e2214935119.

Jamal-Hanjani M, Quezada SA, Larkin J, Swanton C. 2015. Translational implications of tumor heterogeneity. *Clin Cancer Res* **21**: 1258–1266.

Joglekar A, Foord C, Jarroux J, Pollard S, Tilgner HU. 2023. From words to complete phrases: insight into single-cell isoforms using short and long reads. *Transcription* 1–13.

Joglekar A, Prjibelski A, Mahfouz A, Collier P, Lin S, Schlusche AK, Marrocco J, Williams SR, Haase B, Hayes A, et al. 2021. A spatially resolved brain region- and cell type-specific isoform atlas of the postnatal mouse brain. *Nat Commun* **12**: 463.

Köster J, Rahmann S. 2012. Snakemake--a scalable bioinformatics workflow engine. *Bioinformatics* **28**: 2520–2522.

Kuipers J, Tuncel MA, Ferreira P, Jahn K, Beerenwinkel N. 2020. Single-cell copy number calling and event history reconstruction. *BioRxiv*.

Kwok AWC, Qiao C, Huang R, Sham M-H, Ho JWK, Huang Y. 2022. MQuad enables clonal substructure discovery using single cell mitochondrial variants. *Nat Commun* **13**: 1205.

Lähnemann D, Köster J, Szczurek E, McCarthy DJ, Hicks SC, Robinson MD, Vallejos CA, Campbell KR, Beerenwinkel N, Mahfouz A, et al. 2020. Eleven grand challenges in single-cell data science. *Genome Biol* **21**: 31.

Landrum MJ, Lee JM, Riley GR, Jang W, Rubinstein WS, Church DM, Maglott DR. 2014. ClinVar: public archive of relationships among sequence variation and human phenotype. *Nucleic Acids Res* **42**: D980-5.

Lappalainen T, Sammeth M, Friedländer MR, 't Hoen PAC, Monlong J, Rivas MA,

Gonzàlez-Porta M, Kurbatova N, Griebel T, Ferreira PG, et al. 2013. Transcriptome and

genome sequencing uncovers functional variation in humans. *Nature* **501**: 506–511.

Lei JT, Shao J, Zhang J, Iglesia M, Chan DW, Cao J, Anurag M, Singh P, He X, Kosaka

Y, et al. 2018. Functional Annotation of ESR1 Gene Fusions in Estrogen Receptor-

Positive Breast Cancer. *Cell Rep* **24**: 1434-1444.e7.

Liu D, Gao Y, Liu J, Huang Y, Yin J, Feng Y, Shi L, Meloni BP, Zhang C, Zheng M, et al.

2021. Intercellular mitochondrial transfer as a means of tissue revitalization. *Signal*

*Transduct Target Ther* **6**: 65.

Li H. 2018. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**:

3094–3100.

Mansoori B, Mohammadi A, Davudian S, Shirjang S, Baradaran B. 2017. The different

mechanisms of cancer drug resistance: A brief review. *Adv Pharm Bull* **7**: 339–348.

Marine J-C, Dawson S-J, Dawson MA. 2020. Non-genetic mechanisms of therapeutic

resistance in cancer. *Nat Rev Cancer* **20**: 743–756.

Marx V. 2023. Method of the year: long-read sequencing. *Nat Methods* **20**: 6–11.

Milbury CA, Creeden J, Yip W-K, Smith DL, Pattani V, Maxwell K, Sawchyn B, Gjoerup

O, Meng W, Skoletsky J, et al. 2022. Clinical and analytical validation of

FoundationOne®CDx, a comprehensive genomic profiling assay for solid tumors. *PLoS*

*ONE* **17**: e0264138.

Miller TE, Lareau CA, Verga JA, DePasquale EAK, Liu V, Ssozi D, Sandor K, Yin Y,

Ludwig LS, El Farran CA, et al. 2022. Mitochondrial variant enrichment from high-

throughput single-cell RNA sequencing resolves clonal populations. *Nat Biotechnol* **40**:

1030–1034.

Mitra D, Brumlik MJ, Okamgba SU, Zhu Y, Duplessis TT, Parvani JG, Lesko SM, Brogi

E, Jones FE. 2009. An oncogenic isoform of HER2 associated with locally disseminated

788  breast cancer and trastuzumab resistance. *Mol Cancer Ther* **8**: 2152–2162.

789  Muyas F, Sauer CM, Valle-Inclán JE, Li R, Rahbari R, Mitchell TJ, Hormoz S, Cortés-

790  Ciriano I. 2023. De novo detection of somatic mutations in high-throughput single-cell

791  profiling data sets. *Nat Biotechnol.*

792  Osorio D, Cai JJ. 2021. Systematic determination of the mitochondrial proportion in

793  human and mice tissues for single-cell RNA-sequencing data quality control.

794  *Bioinformatics* **37**: 963–967.

795  Prummer M, Bertolini A, Bosshard L, Barkmann F, Yates J, Boeva V, Tumor Profiler

796  Consortium, Stekhoven D, Singer F. 2023. scROSHI: robust supervised hierarchical

797  identification of single cells. *NAR Genom Bioinform* **5**: lqad058.

798  Qin Q, Popic V, Yu H, White E, Khorgade A, Shin A, Wienand K, Dondi A, Beerenwinkel

799  N, Vazquez F, et al. 2024. CTAT-LR-fusion: accurate fusion transcript identification from

800  long and short read isoform sequencing at bulk or single cell resolution. *BioRxiv.*

801  Ramón Y Cajal S, Sesé M, Capdevila C, Aasen T, De Mattos-Arruda L, Diaz-Cano SJ,

802  Hernández-Losa J, Castellví J. 2020. Clinical implications of intratumor heterogeneity:

803  challenges and opportunities. *J Mol Med* **98**: 161–177.

804  Rogers MF, Shihab HA, Mort M, Cooper DN, Gaunt TR, Campbell C. 2018. FATHMM-

805  XF: accurate prediction of pathogenic point mutations via extended features.

806  *Bioinformatics* **34**: 511–513.

807  Roth A, McPherson A, Laks E, Biele J, Yap D, Wan A, Smith MA, Nielsen CB, McAlpine

808  JN, Aparicio S, et al. 2016. Clonal genotype and population structure inference from

809  single-cell tumor sequencing. *Nat Methods* **13**: 573–576.

810  Serin Harmanci A, Harmanci AO, Zhou X. 2020. CaSpER identifies and visualizes CNV

811  events by integrative analysis of single-cell or bulk RNA-sequencing data. *Nat Commun*

812  **11**: 89.

813  Shiau C-K, Lu L, Kieser R, Fukumura K, Pan T, Lin H-Y, Yang J, Tong EL, Lee G, Yan

Y, et al. 2023. High throughput single cell long-read sequencing analyses of same-cell

genotypes and phenotypes in human tumors. *Nat Commun* **14**: 4124.

Stein AN, Joglekar A, Poon C-L, Tilgner HU. 2022. ScisorWiz: visualizing differential

isoform expression in single-cell long-read data. *Bioinformatics* **38**: 3474–3476.

Tokheim C, Karchin R. 2019. Chasmplus reveals the scope of somatic missense

mutations driving human cancers. *Cell Syst* **9**: 9-23.e8.

Van Egeren D, Escabi J, Nguyen M, Liu S, Reilly CR, Patel S, Kamaz B, Kalyva M,

DeAngelo DJ, Galinsky I, et al. 2021. Reconstructing the lineage histories and

differentiation trajectories of individual cancer cells in myeloproliferative neoplasms. *Cell*

*Stem Cell* **28**: 514-523.e9.

Van Egeren D, Kamaz B, Liu S, Nguyen M, Reilly CR, Kalyva M, DeAngelo DJ,

Galinsky I, Wadleigh M, Winer ES, et al. 2022. Transcriptional differences between

JAK2-V617F and wild-type bone marrow cells in patients with myeloproliferative

neoplasms. *Exp Hematol* **107**: 14–19.

Vasan N, Baselga J, Hyman DM. 2019. A view on drug resistance in cancer. *Nature*

**575**: 299–309.

Xia J, Li S, Liu S, Zhang L. 2023. Aldehyde dehydrogenase in solid tumors and other

diseases: Potential biomarkers and therapeutic targets. *MedComm* **4**: e195.

Yuan Y, Ju YS, Kim Y, Li J, Wang Y, Yoon CJ, Yang Y, Martincorena I, Creighton CJ,

Weinstein JN, et al. 2023. Author Correction: Comprehensive molecular characterization

of mitochondrial genomes in human cancers. *Nat Genet* **55**: 1078.

Zafar H, Wang Y, Nakhleh L, Navin N, Chen K. 2016. Monovar: single-nucleotide

variant detection in single cells. *Nat Methods* **13**: 505–507.

Zampieri LX, Silva-Almeida C, Rondeau JD, Sonveaux P. 2021. Mitochondrial transfer

in cancer: A comprehensive review. *Int J Mol Sci* **22**.

Zhang T, Jia H, Song T, Lv L, Gulhan DC, Wang H, Guo W, Xi R, Guo H, Shen N. 2023.

De novo identification of expressed cancer somatic mutations from single-cell RNA

sequencing data. *Genome Med* **15**: 115.

# Acknowledgments

We thank Joanna Hård for her help with mitochondrial mutation contamination. We thank Ivan Topolsky for his support with cloud computing. We thank Lara Fuhrmann for naming LongSom. A.D. and N.Bo were supported by the European Union's Horizon 2020 research and innovation program under the Marie Sklodowska-Curie grant agreement (#766030 to N.Be). B.J.H. was supported by the National Cancer Institute grant U24CA180922.

# Author information

## Corresponding author

Correspondence to Niko Beerenwinkel: <niko.beerenwinkel@bsse.ethz.ch>

## Author contributions

N.Be. acquired the funding for this project. A.D. conceptualized the idea of this project. A.D. designed the LongSom pipeline with the help of N.Bo. The Tumor Profiler Consortium provided scDNA-seq data and NGS panel results. A.D. conducted all computational analyses except scDNA subclones inference (performed by P.F.). A.D. did all the visualizations and figures. A.D., N.Bo., F.J., and N.Be interpreted the results. A.D. and N.Bo. wrote the manuscript with contributions from all authors. B.J.H integrated single cell and long read isoform support into CTAT-Mutations and assisted with fusion transcript identification via CTAT-LR-fusion. All authors read and approved the final manuscript.

## Tumor Profiler Consortium authors list

Rudolf Aebersold[2], Melike Ak[28], Faisal S Al-Quaddoomi[9,17], Silvana I Albert[7], Jonas Albinus[7], Ilaria Alborelli[24], Sonali Andani[6,17,26,31], Per-Olof Attinger[11], Marina Bacac[16], Daniel Baumhoer[24], Beatrice Beck-Schimmer[39], Niko Beerenwinkel[4,17], Christian Beisel[4], Lara Bernasconi[27], Anne Bertolini[9,17], Bernd Bodenmiller[8,35], Ximena Bonilla[6,17,26], Lars Bosshard[9,17], Byron Calgua[24], Ruben Casanova[35], Stéphane Chevrier[35], Natalia Chicherova[9,17], Ricardo Coelho[18], Maya D'Costa[10], Esther Danenberg[37], Natalie Davidson[6,17,26], Monica-Andreea Drăgan[4], Reinhard Dummer[28], Stefanie Engler[35], Martin Erkens[14], Katja Eschbach[4], Cinzia Esposito[37], André Fedier[18], Pedro Ferreira[4], Joanna Ficek[6,17,26], Anja L Frei[31], Bruno Frey[13], Sandra Goetze[7], Linda Grob[9,17], Gabriele Gut[37], Detlef Günther[5], Martina Haberecker[31], Pirmin Haeuptle[1], Viola Heinzelmann-Schwarz[18,23], Sylvia Herter[16], Rene Holtackers[37], Tamara Huesser[16], Alexander Immer[6,12], Anja Irmisch[28], Francis Jacob[18], Andrea Jacobs[35], Tim M Jaeger[11], Katharina Jahn[4], Alva R James[6,17,26], Philip M Jermann[24], André Kahles[6,17,26], Abdullah Kahraman[17,31], Viktor H Koelzer[31], Werner Kuebler[25], Jack Kuipers[4,17], Christian P Kunze[22], Christian Kurzeder[21], Kjong-Van Lehmann[6,17,26], Mitchell Levesque[28], Ulrike Lischetti[18], Sebastian Lugert[10], Gerd Maass[13], Markus G Manz[30], Philipp Markolin[6,17,26], Martin Mehnert[7], Julien Mena[2], Julian M Metzler[29], Nicola Miglino[1], Emanuela S Milani[7], Holger Moch[31], Simone Muenst[24], Riccardo Murri[38], Charlotte KY Ng[24,34], Stefan Nicolet[24], Marta Nowak[31], Monica Nunez Lopez[18], Patrick GA Pedrioli[3], Lucas Pelkmans[37], Salvatore Piscuoglio[18,24], Michael Prummer[9,17], Natalie Rimmer[18], Mathilde Ritter[18], Christian Rommel[14], María L Rosano-González[9,17], Gunnar Rätsch[3,6,17,26], Natascha Santacroce[4], Jacobo Sarabia del Castillo[37], Ramona Schlenker[15], Petra C Schwalie[14], Severin Schwan[11], Tobias Schär[4], Gabriela Senti[27], Wenguang Shao[7], Franziska Singer[9,17], Sujana Sivapatham[35], Berend Snijder[2,17], Bettina Sobottka[31], Vipin T Sreedharan[9,17], Stefan Stark[6,17,26], Daniel J Stekhoven[9,17], Tanmay Tanna[4,6], Alexandre PA Theocharides[30], Tinu M Thomas[6,17,26], Markus Tolnay[24], Vinko Tosevski[16], Nora C Toussaint[9,17], Mustafa A Tuncel[4,17], Marina Tusup[28], Audrey Van Drogen[7], Marcus Vetter[20],

Tatjana Vlajnic[24], Sandra Weber[27], Walter P Weber[19], Rebekka Wegmann[2], Michael

Weller[33], Fabian Wendt[7], Norbert Wey[31], Andreas Wicki[30,36], Mattheus HE Wildschut[2,30],

Bernd Wollscheid[7], Shuqing Yu[9,17], Johanna Ziegler[28], Marc Zimmermann[6,17,26], Martin

Zoche[31], Gregor Zuend[32]

[1]Cantonal Hospital Baselland, Medical University Clinic, Rheinstrasse 26, 4410 Liestal,

Switzerland, [2]ETH Zurich, Department of Biology, Institute of Molecular Systems Biology,

Otto-Stern-Weg 3, 8093 Zurich, Switzerland, [3]ETH Zurich, Department of Biology, Wolfgang-

Pauli-Strasse 27, 8093 Zurich, Switzerland, [4]ETH Zurich, Department of Biosystems Science

and Engineering, Mattenstrasse 26, 4058 Basel, Switzerland, [5]ETH Zurich, Department of

Chemistry and Applied Biosciences, Vladimir-Prelog-Weg 1-5/10, 8093 Zurich, Switzerland,

[6]ETH Zurich, Department of Computer Science, Institute of Machine Learning,

Universitätstrasse 6, 8092 Zurich, Switzerland, [7]ETH Zurich, Department of Health Sciences

and Technology, Otto-Stern-Weg 3, 8093 Zurich, Switzerland, [8]ETH Zurich, Institute of

Molecular Health Sciences, Otto-Stern-Weg 7, 8093 Zurich, Switzerland, [9]ETH Zurich,

NEXUS Personalized Health Technologies, John-von-Neumann-Weg 9, 8093 Zurich,

Switzerland, [10]F. Hoffmann-La Roche Ltd, Grenzacherstrasse 124, 4070 Basel, Switzerland,

[11]F. Hoffmann-La Roche Ltd, Grenzacherstrasse 124, 4070 Basel, Switzerland, , [12]Max

Planck ETH Center for Learning Systems, , [13]Roche Diagnostics GmbH, Nonnenwald 2,

82377 Penzberg, Germany, [14]Roche Pharmaceutical Research and Early Development,

Roche Innovation Center Basel, Grenzacherstrasse 124, 4070 Basel, Switzerland, [15]Roche

Pharmaceutical Research and Early Development, Roche Innovation Center Munich, Roche

Diagnostics GmbH, Nonnenwald 2, 82377 Penzberg, Germany, [16]Roche Pharmaceutical

Research and Early Development, Roche Innovation Center Zurich, Wagistrasse 10, 8952

Schlieren, Switzerland, [17]SIB Swiss Institute of Bioinformatics, Lausanne, Switzerland,

[18]University Hospital Basel and University of Basel, Department of Biomedicine,

Hebelstrasse 20, 4031 Basel, Switzerland, [19]University Hospital Basel and University of

Basel, Department of Surgery, Brustzentrum, Spitalstrasse 21, 4031 Basel, Switzerland,

[20]University Hospital Basel, Brustzentrum & Tumorzentrum, Petersgraben 4, 4031 Basel,

Switzerland, [21]University Hospital Basel, Brustzentrum, Spitalstrasse 21, 4031 Basel,

Switzerland, [22]University Hospital Basel, Department of Information- and Communication

Technology, Spitalstrasse 26, 4031 Basel, Switzerland, [23]University Hospital Basel,

Gynecological Cancer Center, Spitalstrasse 21, 4031 Basel, Switzerland, [24]University

Hospital Basel, Institute of Medical Genetics and Pathology, Schönbeinstrasse 40, 4031

Basel, Switzerland, [25]University Hospital Basel, Spitalstrasse 21/Petersgraben 4, 4031

Basel, Switzerland, [26]University Hospital Zurich, Biomedical Informatics,

Schmelzbergstrasse 26, 8006 Zurich, Switzerland, [27]University Hospital Zurich, Clinical

Trials Center, Rämistrasse 100, 8091 Zurich, Switzerland, [28]University Hospital Zurich,

Department of Dermatology, Gloriastrasse 31, 8091 Zurich, Switzerland, [29]University

Hospital Zurich, Department of Gynecology, Frauenklinikstrasse 10, 8091 Zurich,

Switzerland, [30]University Hospital Zurich, Department of Medical Oncology and Hematology,

Rämistrasse 100, 8091 Zurich, Switzerland, [31]University Hospital Zurich, Department of

Pathology and Molecular Pathology, Schmelzbergstrasse 12, 8091 Zurich, Switzerland,

[32]University Hospital Zurich, Rämistrasse 100, 8091 Zurich, Switzerland, [33]University

Hospital and University of Zurich, Department of Neurology, Frauenklinikstrasse 26, 8091

Zurich, Switzerland, [34]University of Bern, Department of BioMedical Research,

Murtenstrasse 35, 3008 Bern, Switzerland, [35]University of Zurich, Department of Quantitative

Biomedicine, Winterthurerstrasse 190, 8057 Zurich, Switzerland, [36]University of Zurich,

Faculty of Medicine, Zurich, Switzerland, [37]University of Zurich, Institute of Molecular Life

Sciences, Winterthurerstrasse 190, 8057 Zurich, Switzerland, [38]University of Zurich,

Services and Support for Science IT, Winterthurerstrasse 190, 8057 Zurich, Switzerland,

[39]University of Zurich, VP Medicine, Künstlergasse 15, 8001 Zurich, Switzerland

# 939    Conflict of interest

940    The authors declare no competing interests.