Road density simplifies regional food webs

Abstract

Roads stands as major threats to biodiversity, affecting the functioning of ecosystems and their provision of ecosystem services. Understanding how road-related impacts affect the dynamics of ecological interactions is essential to help manage human impacts on biodiversity, but such studies remain largely unexplored. We investigated the intricate relationship between roads, species vulnerability to road density, and the ensuing effects on food webs across Europe. Utilizing species-specific road density thresholds and trophic interaction data, we constructed regional food webs to assess the potential loss of trophic interactions due to roadkill. Our analysis, encompassing 551 species across three trophic levels, revealed spatially varied impacts, with areas surrounding major cities facing drastic losses of trophic interaction exceeding 90%. Notably, 191 species may be affected by trophic interaction losses (loss of prey or predator). Apex predators exhibited lower direct impacts from road density, while basal-level species seem to be more exposed to direct road-related effects, potentially triggering a cascade of interaction disruptions. Our findings emphasize the need for informed road infrastructure development and targeted conservation strategies to mitigate the negative impacts of roads and traffic, thereby preserving the integrity of ecological networks. This research offers valuable insights for policymakers and conservation practitioners globally by identifying critical areas where road-induced cascade effects may be most pronounced and highlighting trophic groups of species that may be at higher risk.

Introduction

Roads can play a vital role in human development by allowing access to important services, such as education and health care, further contributing to economic growth by enabling trade (Gachassin et al., 2010; Iimi et al., 2015; Sewell et al., 2019). However, roads and traffic stand as a major threat to biodiversity, leading to habitat fragmentation and degradation (Forman and Alexander, 1998; Maxwell et al., 2016; Van der Ree et al., 2015). Notably, the expansion of road networks and the associated increase in traffic drastically affects wildlife populations, primarily due to animal-vehicle collisions (Grilo et al., 2020; Morelli et al., 2020). In developed regions like Europe, where the density of roads rank among the highest globally (Meijer et al., 2018), roads exert a considerable impact on wildlife. For example, recent estimates point to over 200 million individual birds and mammals being road-killed on European paved roads every year (Grilo et al., 2020).

Road-related impacts are likely to ripple through regional ecological communities when species' interactions are weakened or even lost following reductions in population abundance or local extinctions (Barrientos et al., 2020). A simplification of ecological network structure such as food webs (Dunne et al., 2002; Mestre et al., 2022), can change fluxes of matter and energy (Barnes et al., 2018), affecting the functioning of ecosystems, ultimately disrupting the provision of ecosystem services (Barbosa et al., 2017; Tylianakis et al., 2008). This simplification of trophic chains can further impact species indirectly by altering prey density or reducing predator pressure (Leighton et al., 2023). Therefore, understanding how roads and traffic affect the dynamics of ecological interactions, particularly food webs, is crucial for managing human impacts on biodiversity, especially those associated with transport infrastructure. However, the study of road-induced impacts on ecological networks remains largely unexplored, likely due to the logistical and financial

complexities of monitoring numerous regional food webs across varying road density

scenarios and over large spatial extents.

Previous research established the generalized theoretical population model for

deriving species-specific vulnerability to roads based on road density thresholds, beyond

which local populations may not persist (Borda-de-Água et al., 2011). This theoretical

framework was recently used to calculate thresholds of road density for a collection of

European birds and mammals (Grilo et al., 2020) (see Methods for further details). Some

examples of the thresholds of road density include 0.91 km/km^2 for Iberian lynx (*Lynx*

pardinus), 1.43 km/km^2 for golden eagle (*Aquila chrysaetos*), 2.51 km/km^2 for wild cat (*Felis*

silvestris), and 3.00 km/km^2 for the wild rabbit (*Oryctolagus cuniculus*) (all species-specific

threshold values are available in Grilo et al. (2020)). We used these estimated thresholds to

infer where potential losses in species' occurrences may occur and, consequently, the

interactions they were involved in (Fig. 1).

Using the combined information on road density thresholds and food webs, we

investigated how food webs can be imperiled when road density surpasses the species-

specific thresholds beyond which local populations may not persist. Specifically, we sought

to evaluate the extent of potential interaction loss across Europe and trophic levels; and

compare, for each species, the amount of distribution area that could be lost or degraded due

to road density exceeding *i*) its specific vulnerability threshold (direct effect) and *ii*) that of

the species that interact with it i.e., prey and/or predators (indirect effect). We expected a

higher loss of interactions and higher degradation of range area with increasing road density

values. However, given the historical presence of roads in Europe and the higher

vulnerability of apex predators to roadkill effects (Rytwinski and Fahrig, 2012), we predicted

a higher direct effect for lower trophic levels and a higher vulnerability of interactions at

higher trophic levels due to indirect effects.

70
Figure 1 – Road-induced simplification of food webs. The diagram illustrates the potential
loss of interactions resulting from road density exceeding species-specific thresholds. In
Panel 1, the road density thresholds for nine species are presented. The garden dormouse (*E.*
quercinus) shows the lowest estimated road density threshold (0.4 km/km^2), while the
Algerian mouse (*M. spretus*) has the highest threshold (3.8 km/km^2). These species
participate in the food web depicted in Panel 2 (network not exhaustive), representing a
scenario without roads. For two regions with lower and higher road densities (Panels 3 and 4;
also indicated in Panel 1), the network may simplify: some species face a risk of local
extinction (shown in grey) when road densities exceed their thresholds, increasing the
likelihood of roadkill and threatening their survival. This leads to a loss of interactions
(indicated by dashed lines). In contrast, species depicted in black are more likely to persist, as
their road density thresholds exceed the levels present in the region. In the example provided,
the Iberian lynx (*L. pardinus*) and garden dormouse may face extinction in region A. In
region B, golden eagle (*A. chrysaetos*) and its prey, including red-legged partridge (*A. rufa*),
southern water vole (*A. sapidus*), and Iberian hare (*L. granatensis*), could also be at risk,
disrupting predator-prey dynamics. Wild cat (*F. silvestris*), common quail (*C. coturnix*), and

Algerian mouse might persist even at the highest road densities, as can their trophic
interactions. Silhouettes from phylopic (www.phylopic.org).

Methods

Species vulnerability to roads

Species-specific thresholds for road density were obtained for birds and mammals (Grilo et

al., 2020). The authors estimated long-term vulnerability to roadkill using the theoretical

demographic model developed by Borda-de-Água et al. (2011), which differentiates between

landscapes with roads and non-roads:

95
$$Max_{road} = \frac{r_1}{r_1 + |r_0|} \text{ (eq. 1)}$$

96
$$r_0 = \frac{-\mu \times RM}{year} \text{ (eq. 2)}$$

Following Grilo et al. (2020), r_1 is the intrinsic population growth rate and r_0 is the

annual rate of population decay on paved roads, where μ is the species' annual natural

mortality rate given its longevity, and *RM* is a scaling factor for road mortality assuming that

roads are unsuitable environments for species to persist, with *RM* = 1000 having been the

value used, in accordance to previously assumed by (Ceia-Hasse et al., 2017). According to

the authors, in this model, non-road areas support positive population growth rates, while

road areas lead to rapid population declines, assuming roads are unsuitable habitats. The

intrinsic population growth rate (r_1) is calculated using a simplified version of the Euler

equation that incorporates species traits such as maturity age and offspring, as well as the

annual natural mortality rate, based on species longevity (Borda-de-Água et al., 2011; Grilo

et al., 2020). Using this approach, the authors calculated species-specific maximum road

density thresholds (eq. 1), beyond which population persistence is unlikely. Importantly, both

the authors and our study consider only paved roads, as these are where roadkill is more

likely to occur due to generally higher traffic volumes [see Borda-de-Água et al. (2011) for

further details on the modelling procedure, and Grilo et al. (2020) for further details on the

estimation of species-specific thresholds]. We regarded the road density threshold as an

indicator that enables a species-specific ranking of regions where species are unlikely to

persist, potentially leading to the loss of their ecological interactions.

Building local potential food webs and risk of interaction disruption

We built local trophic structures using a grid of 50 x 50 km cells overlaid upon the European

region (n=4,743 cells), the same used by Grilo et al. (2020). Building local potential food

webs was guided by the observed interactions outlined in the TETRA-EU metaweb

(Maiorano et al., 2020). This metaweb is based on observed interactions obtained from the

scientific literature and summarizes the potential trophic interactions for tetrapod species

(mammals, breeding birds, reptiles, and amphibians) across Europe and the Northern

Mediterranean basin. If a trophic interaction between two species was documented in the

metaweb, it was considered indicative of their potential interaction within the regional

network (i.e., at the grid cell level). We characterized the trophic level of each species in the

metaweb using a categorical classification, dividing species into three groups: top-level

(species that only have prey), intermediate level (species with both predators and prey), and

basal (species that only have predators). We did this by resorting to the functions

'BasalNodes', 'IntermediateNodes', and 'TopLevelNodes' of the cheddar R package

(Hudson et al., 2013).

We then used the road density thresholds to predict potential points of trophic

interaction disruption in the event of species extinction or severe depletion. For each species,

we assumed that grid cells exceeding their species-specific maximum road density threshold,

i.e., beyond which long-term population persistence becomes improbable (Borda-de-Água et

al., 2011), would exhibit an increased probability of losing trophic interactions associated

with that species. Extinction simulation was conducted using the function 'RemoveNodes' of

the cheddar R package (Hudson et al., 2013).

All the computations were conducted in R (R Core Team, 2021), using the R

packages cheddar (for network analysis) (Hudson et al., 2013), igraph (for network analysis)

(Csárdi et al., 2023), taxize (to combine both datasets, the vulnerability dataset and the

metaweb) (Chamberlain and Szöcs, 2013), and terra (for R based spatial analysis and for

creating the output shapefiles) (Hijmans, 2023). Images were produced in QGIS (2020) and R

(R Development Core Team, 2022). The dataset is available in FIGSHARE. All code is

available on GitHub.

Results

General patterns

Our pool of species, for which a threshold of vulnerability to road density was available and

referenced in the TETRA-EU metaweb, consisted of 551 species (349 birds and 202

mammals). This pool comprised 12 top-level species (2.2% of the total), 107 intermediate-

level species (19.4%), and 432 basal-level species (78.4%) (see Supplementary material S1

for species names). The three trophic levels currently exhibit varying distributions across

Europe (Fig. 2A-C), notably with higher species richness in regions characterized by lower

road density (Fig. 2D), particularly top-level species. This pool of species assembled

altogether a mean (±SD) of 2,057±819 predator-prey interactions per grid cell across Europe.

A greater number of interactions were observed in mountainous regions like the Pyrenees,

Alps, and Carpathians, alongside areas in eastern Europe, regions that exhibit lower road

density (Fig. 2E).

Figure 2 - Richness of top, intermediate and basal level species (A-C), road density in km/km^2 (D), and estimated number of interactions (E) per grid cell, across Europe. All maps use a grid cell resolution of 50x50 km. White lines depict country borders.

Trophic interaction loss

On average, 4.5% (±9.4) of interactions in our regional food webs face potential loss due to road impacts i.e., when road density surpasses the species-specific vulnerability threshold of either the prey or predator engaged in the interaction. However, the simplification of food webs is likely to vary spatially, being particularly intense in areas surrounding large European cities. In these areas, road density can reach 2.5 km/km² (see Fig. 2D), resulting in the loss of over 90% of potential interactions (Fig. 3A). The loss of interactions in the local food webs seems to have a strong relationship with road density, reaching a 30% loss when road density approaches 1 km/km² (Fig. 3B).

Figure 3 - A) Expected loss of interactions due to road impacts; B) Relation between road density and interaction loss (red line depicts the smooth line using a polynomial spline, here used for better visualization of the relation between both variables).

Effects within species range

Among the species examined, 81 (15% of the total) had over 10% of their range with road density surpassing their vulnerability threshold, putting them at significant risk of local extinction due to direct road impacts. These include one top-level species, 12 intermediate species, and 68 basal species (Fig. 4). Notably, the apex predator booted eagle *(Hieraaetus pennatus*) may disappear from 16% of its current range (see Supplementary material S1). The intermediate-level species short-toed snake eagle *(Circaetus gallicus)* and garden dormouse *(Eliomys quercinus)* may experience a substantial loss or degradation within over one-third of their range. For the basal-level species hazel grouse *(Tetrastes bonasia)* and both mole rats *(Spalax graecus* and *S. zemni)*, road densities surpass their vulnerability threshold in nearly half the range (grouse) or the full range (mole rats), leading to potential local extinctions across their entire territories.

On the other hand, 191 species (35% of the total), including 10 top-level, 81 intermediate-level, and 100 basal-level species, exhibit more than 10% of their range at risk of losing trophic interactions due to indirect effects (Fig. 4). Notably, two top-level species,

the golden eagle and the wolf (*Canis lupus*), may experience interaction losses with prey in

over half their territory (Fig. 4). For example, the golden eagle is mostly affected by potential

interaction losses with the hazel grouse (*Bonasa bonasia*; 16% of the total interactions lost),

the European snow vole (*Chionomys nivalis*; 8%), the European mole (*Talpa europaea*; 7%)

and the garden dormouse (7%). As for the wolf, it will potentially lose interactions mostly

with hazel grouse (13% of total interactions), Eurasian pygmy shrew (*Sorex minutus*; 13%)

and the common shrew (*Sorex araneus*; 8%).

Figure 4 – Range area with potential trophic interaction losses due to direct and indirect
effects of roads, by trophic level (top, intermediate and basal). The circle radius is
proportional to the range of the species. Grey diagonal line represents 100% of range areas
affected by combination of direct and indirect effects. Highlighted species are mentioned in
the text.

Likewise, intermediate-level species, such as barn owl (*Tyto alba*) and wild cat (*Felis*

silvestris), may experience losses of trophic interactions in over 70% of their range. The barn

owl may lose most of its interactions as a consumer, namely with the Eurasian pygmy shrew

(8%), the whiskered bat (*Myotis mystacinus*; 5%) and the European mole (5%). Conversely,

as resource, the barn owl may lose interactions with the Eurasian goshawk (*Accipiter gentilis*;

90%) and the tawny owl (*Strix aluco*; 10%). As for the wild cat, it will lose most of its

interactions as a predator with the pygmy shrew (14%), the common shrew (8%) and the

hazel grouse (8%). As a resource, the wild cat will lose interactions with the Eurasian lynx

(*Lynx lynx*; 75%) and the golden eagle (25%). Finally, the basal-level species, Western

Orphean warbler *(Curruca hortensis)* and broom hare (*Lepus castroviejoi*), may lose

interactions as prey in 44% and 91% of their range, respectively (Fig. 4). The Western

Orphean warbler may experience interaction losses mostly from Eurasian scops owl (*Otus*

scops; 39%), booted eagle (26%), woodchat shrike (*Lanius senator*; 17%) or Montagu's

Harrier (*Circus pygargus*; 7%). On the other hand, the broom hare experiences interaction

losses exclusively with the short-toed snake eagle.

These findings strongly imply that the detrimental impacts of road density extend far

beyond what meets the eye. For instance, top-level species currently face a minor risk of

depletion due to roadkill. However, they may suffer from indirect effects across a significant

portion of their range, where at least one prey species goes extinct due to roadkill (Fig. 4).

Discussion

Road-related effects are expected to have a direct impact on species occurrence and

abundance (Andrasi et al., 2021; Ascensão et al., 2022; Borda-de-Água et al., 2011; Forman

and Alexander, 1998). Roadkill, in particular, can be a major driver of population depletion

and reduce the population persistence in road surrounding areas (Ascensão and Desbiez,

2022; Barrientos et al., 2021; Moore et al., 2023). Yet, beyond such population-level effects,

its repercussions are likely to also exert profound consequences on ecological networks. Our

results indicate that increasing road density may significantly reduce the number of trophic

interactions. This simplification of the food web is due not only to the direct depletion of

populations but also to the indirect effects on food chains, such as the elimination of

predators or prey. Furthermore, this loss of interactions occurs differently across space, with a

greater incidence near large cities, and areas coinciding with higher road density, such as

across Belgium, France, Germany, Italy, and the north of the Iberian Peninsula. Such

imbalanced losses may have far-reaching implications for regional food webs, which could

ultimately culminate in local extinctions, disturbing ecosystem functioning, and disrupting

the delivery of essential ecosystem services.

Overall, top and intermediate species seem to follow a pattern dominated by loss of

interactions due to the indirect effects of roads, with intermediate species having higher range

area affected by losses than top-level species. Apart from the booted eagle, which presently

inhabits an extensive region marked by road densities exceeding its estimated vulnerability

threshold, the other top-level species like the golden eagle, brown bear (*Ursus arctos*), or

Eurasian lynx, face minimal risk of being depleted due to direct road-related effects, as their

current ranges have low road densities. This implies that impacts such as roadkill may not

directly affect the persistence of the majority of top-level species. This is not surprising given

that such apex predators are already restricted to habitats with relatively lower human impact

and activity, and consequently with low road density areas (Bleyhl et al., 2021; Cretois et al.,

2021). Conversely, basal species seem to be highly impacted by direct effects, therefore

facing local extinction risk, and potentially eliminating the interactions with their predators.

Yet, several basal species may also benefit from a release of predator interactions in their

range areas.

Research has demonstrated that habitat fragmentation, in general, affects mostly

consumer species, with larger body sizes, at higher trophic levels (Rytwinski and Fahrig,

2013, 2012). These species will generally go extinct faster than species with smaller body

sizes and at lower trophic levels, even if having better dispersal ability (Liao et al., 2017;

Ryser et al., 2019). On the other hand, despite the widespread suitable habitat for large

carnivores, such as the brown bear and the Iberian lynx in Iberian Peninsula, human pressure

is limiting the potential range expansion of these species (Pratzer et al., 2023). Likewise,

wolves and other large carnivores are known to avoid sources of human disturbance,

including high-road-density areas (Bleyhl et al., 2021; Cretois et al., 2021; Rio-Maior et al.,

2019). This might explain why in our dataset top-level species are mostly restricted to lower

road density areas.

Despite the long debate on the role of bottom-up versus top-down control in food

webs (Leroux and Loreau, 2015), some evidence suggests that human activities tend to

amplify bottom-up effects while weakening top-down trophic cascades in terrestrial

ecosystems (Muhly et al., 2013). Our results seem to corroborate this notion, as they indicate

that the impact of roads is particularly strong on prey species, disrupting their populations,

thereby affecting higher trophic levels in the regional networks. Similar to our findings,

previous research has also shown that prey species can benefit from lower exposition to

predation pressure (Liao et al., 2017), and that habitat fragmentation – and probably also high

road density areas – leads to ecological network simplification (Pires et al., 2023). However,

the impact of roads on intermediate and basal-level generalist species may have stronger

implication for food web dynamics, as generalist species play important roles in structuring

the topology of ecological networks (Bastazini et al., 2019; Gaston and Fuller, 2008;

Memmott et al., 2004; Poisot et al., 2013; Richmond et al., 2005; Valiente-Banuet et al.,

2015). These species interact with more species than specialist species, thereby creating a

more cohesive network (Bastazini et al., 2019; Martín González et al., 2010). By linking

otherwise disconnected sub-networks, generalists enhance the overall connectivity, which is

likely to increase their ability to respond to perturbations, increasing their stability

(Bascompte et al., 2003).

We acknowledge that there are three limitations in our study. Firstly, although the

metaweb approach offers valuable insights into species' ecological associations, it overlooks

the spatial and ecological context of these interactions (Chamberlain et al., 2014; Liu and

Gaines, 2022). The strength and distribution of species interactions may differ across

different populations and habitats due to environmental conditions and the presence of other

species. For example, the Iberian lynx may have a few prey species (27 prey species

according to metaweb used). However, its persistence is particularly dependent on the

presence of one key prey, the wild rabbit, which constitutes 75%–99% of its diet (Fedriani et

al., 1999). As such, losing interactions with other prey would probably not have relevant

consequences for the persistence for this top predator. Incorporating interaction strength into

food webs, by using weighted networks instead of unweighted ones, would improve the

realism of our models. However, there is no available information that would allow us to

construct such networks.

On the other hand, the database on trophic interactions builds on species' co-

occurrence. As so, we do not fully account for the spatial structure of these biotic interactions

(e.g., Poisot et al., 2015). The strength of interactions between two species can vary across

different parts of their overlapping ranges, and in some regions, these interactions might even

be absent. Despite the limitations posed by the lack of detailed information on species

interactions, intraspecific variability, and spatial variation in interaction strength, our

approach represents the most viable solution for addressing these research gaps. Furthermore,

undocumented interactions are not included in the metaweb and, as a result, are also missing

from our framework. Nevertheless, to our knowledge, this database remains the most

comprehensive resource available on trophic interactions in Europe.

Finally, following Borda-de-Água et al. (2011) and Grilo et al. (2020), we used road

density as a proxy to measure the impact of roads on wildlife, acknowledging that this metric

may not fully capture the spatial variation in these effects. This choice assumes that road

density generally correlates with traffic intensity across regions, which is reasonable for a

continental-scale study. Incorporating traffic volume into the modeling framework could

potentially provide a more accurate distinction between areas with higher and lower wildlife

impacts. Unfortunately, a significant portion of the road network lacks standardized traffic

volume data, limiting our ability to include this factor. Future research should prioritize

integrating traffic volume data, to refine assessments of areas most at risk from road impacts

on wildlife populations and ecological interactions. Yet, existing studies do not consistently

demonstrate a clear relationship between traffic intensity or speed and roadkill rates

(Denneboom et al., 2021; Grilo et al., 2015). Therefore, using road density might actually be

a more reliable indicator of the potential for roads to disrupt biological communities than

traffic speed or intensity.

Our findings indicate that the impacts of roads hold profound implications for

ecosystem conservation. Roads are likely to prevent or hamper the spread of apex predators

(Maxwell et al., 2016; Ripple et al., 2014). The negative correlation between its distribution

and road density reflects this restrictive relationship. However, while road impacts, such as

roadkill, may be exerting a lower role in directly depleting top-level species in their current

distribution area, roads are likely to be constraining the functional connectivity among their

populations. For example, a major threat to the Iberian and Eurasian lynxes' dispersers is

roadkill (Kramer☐Schadt et al., 2004; Philips, 2020). On the other hand, the loss of prey due

to high road densities may contribute to the failure of top-level species to persist throughout

dispersal areas. Numerous basal species appear to encompass a significant portion of their

ranges with road densities above their estimated threshold of vulnerability (Borda-de-Água et

al., 2011; Grilo et al., 2020). Consequently, roads might be more noticeably depleting prey

species, while generating less visible effects by limiting the presence of predators through

disrupted interactions.

We suggest that roads are contributing to and probably accelerating the simplification

and contraction of ecological networks (Felipe-Lucia et al., 2020; Heleno et al., 2020; Mestre

et al., 2022). With the expected massive expansion of road networks, including the

construction of 3–5 Mio km of new roads worldwide by 2050 (Meijer et al., 2018), these

adverse impacts will likely extend to previously undisturbed ecosystems in developing

regions, while exacerbating them in more developed countries. Therefore, immediate action

is imperative to address the changes in ecosystems that may arise, namely to preserve food

webs, as there is an imminent risk of pushing entire ecosystems beyond their safe limits

(Steffen et al., 2015). This requires improving our understanding of the interconnections

between transport networks and higher levels of ecological organization, such as ecological

communities and their interactions.

This study carries significant implications for road planning and landscape

management strategies, especially concerning protected areas and their connecting corridors.

The focus for conserving top predators should revolve around establishing connectivity

between high-quality habitat areas, strategically steering away from regions having high road

density. However, their conservation must include measures to prevent the depletion of prey

due to high road densities, otherwise top-level species may struggle to persist in the areas

they seek to colonize. On the other hand, the impact of roads on intermediate and basal-level

generalist species warrants a more cautious approach and study. These species have extensive

environmental adaptability and a broad dietary spectrum, often having wide distributions

throughout Europe. As previously mentioned, given that they interact with a

disproportionately large set of coexisting species, they play a fundamental role in shaping

ecological communities and influencing ecosystem functioning (Gaston, 2010). As so, their disappearance, namely due to high roadkill rates, may cause substantial losses in predator-prey interactions. However, the conservation of generalists, and particularly road mitigation targeting those species, is not always a priority. Moreover, while threatened or iconic species receive dedicated research and tailored mitigation measures (e.g., Ascensão et al., 2019b), more common and less appealing species may be disregarded from road management programs. But the prevalence of roadkill among common species is an alarming concern in our ecosystems (Ascensão et al., 2019a; Grilo et al., 2018). Despite their familiarity, these species often face high mortality rates that may tip local populations into extinction, causing a significant decline in their overall abundance and biomass. Such a decline is not just a threat to these species alone but also to all species they interact with, the ecosystem services and contributions from nature they provide, therefore being a pivotal factor contributing to the broader decline in biodiversity.

Overall, road and landscape management should account for the preservation of conditions for species persistence and movement for the different trophic levels. This includes habitat restoration and reducing road fragmentation by implementing tailored measures that bolster the resilience of the different trophic levels within their habitats and road proximity, such as different typologies of road passages (Clevenger et al., 2001; Lesbarrères and Fahrig, 2012). Such targeted species-specific approaches of habitat restoration and functional connectivity could play a pivotal role in mitigating the detrimental impacts of road networks on wildlife and preserving ecological network integrity.

References

Ascensão, F., Desbiez, A.L.J., 2022. Assessing the impact of roadkill on the persistence of wildlife populations: a case study on the giant anteater. bioRxiv. https://doi.org/10.1101/2022.01.18.476626

Ascensão, F., Kindel, A., Teixeira, F.Z., Barrientos, R., D'Amico, M., Borda-de-Água, L., Pereira, H.M., 2019a. Beware that the lack of wildlife mortality records can mask a serious impact of linear infrastructures. Glob. Ecol. Conserv. e00661. https://doi.org/10.1016/j.gecco.2019.e00661

Ascensão, F., Niebuhr, B.B., Moraes, A.M., Alexandre, B.R., Assis, J.C., Alves☐Eigenheer, M.A., Ribeiro, J.W., Morais, M.M., Martins, A.F., Oliveira, A., Moraes, E., Ramos, J.H., Lorini, M.L., Ferraz, L.P., Culot, L., Dietz, J.M., Ruiz☐Miranda, C.R., Ribeiro, M.C., 2019b. End of the line for the golden lion tamarin? A single road threatens 30 years of conservation efforts. Conserv. Sci. Pract. 1. https://doi.org/10.1111/csp2.89

Barbosa, M., Fernandes, G.W., Lewis, O.T., Morris, R.J., 2017. Experimentally reducing species abundance indirectly affects food web structure and robustness. J. Anim. Ecol. 86, 327–336. https://doi.org/10.1111/1365-2656.12626

Barnes, A.D., Jochum, M., Lefcheck, J.S., Eisenhauer, N., Scherber, C., O'Connor, M.I., de Ruiter, P., Brose, U., 2018. Energy Flux: The Link between Multitrophic Biodiversity and Ecosystem Functioning. Trends Ecol. Evol. 33, 186–197. https://doi.org/10.1016/j.tree.2017.12.007

Barrientos, R., Ascensão, F., D'Amico, M., 2020. Inappropriate tourist behavior in protected areas can lead to wildlife road☐kills. Anim. Conserv. 23, 343–344. https://doi.org/10.1111/acv.12547

Barrientos, R., Ascensão, F., D'Amico, M., Grilo, C., Pereira, H.M., 2021. The lost road: do transportation networks imperil wildlife population persistence? Perspect. Ecol. Conserv. https://doi.org/10.1016/j.pecon.2021.07.004

Bascompte, J., Jordano, P., Melián, C.J., Olesen, J.M., 2003. The nested assembly of plant–animal mutualistic networks. Proc. Natl. Acad. Sci. 100, 9383–9387. https://doi.org/10.1073/pnas.1633576100

Bastazini, V.A., Debastiani, V.J., Azambuja, B.O., Jr, P.R.G., Pillar, V.D., 2019. Loss of Generalist Plant Species and Functional Diversity Decreases the Robustness of a Seed Dispersal Network. Environ. Conserv. 46, 52–58. https://doi.org/10.1017/S0376892918000334

Bleyhl, B., Ghoddousi, A., Askerov, E., Bocedi, G., Breitenmoser, U., Manvelyan, K., Palmer, S.C.F., Soofi, M., Weinberg, P., Zazanashvili, N., Shmunk, V., Zurell, D., Kuemmerle, T., 2021. Reducing persecution is more effective for restoring large carnivores than restoring their prey. Ecol. Appl. 31, e02338. https://doi.org/10.1002/eap.2338

Borda-de-Água, L., Navarro, L., Gavinhos, C., Pereira, H.M., 2011. Spatio-temporal impacts of roads on the persistence of populations: analytic and numerical approaches. Landsc. Ecol. 26, 253–265. https://doi.org/10.1007/s10980-010-9546-2

Ceia-Hasse, A., Borda☐de☐Água, L., Grilo, C., Pereira, H.M., 2017. Global exposure of carnivores to roads. Glob. Ecol. Biogeogr. 26, 592–600. https://doi.org/10.1111/geb.12564

Chamberlain, S.A., Bronstein, J.L., Rudgers, J.A., 2014. How context dependent are species interactions? Ecol. Lett. 17, 881–890. https://doi.org/10.1111/ele.12279

Chamberlain, S.A., Szöcs, E., 2013. taxize: taxonomic search and retrieval in R. F1000Research 2, 191. https://doi.org/10.12688/f1000research.2-191.v2

Clevenger, A.P., Chruszcz, B., Gunson, K., 2001. Drainage culverts as habitat linkages and factors affecting passage by mammals. J. Appl. Ecol. 38, 1340–1349. https://doi.org/10.1046/j.0021-8901.2001.00678.x

Cretois, B., Linnell, J.D.C., Van Moorter, B., Kaczensky, P., Nilsen, E.B., Parada, J., Rød, J.K., 2021. Coexistence of large mammals and humans is possible in Europe's

anthropogenic landscapes. iScience 24, 103083.
https://doi.org/10.1016/j.isci.2021.103083

Csárdi, G., Nepusz, T., Müller, K., Horvát, S., Traag, V., Zanini, F., Noom, D., 2023. igraph
for R: R interface of the igraph library for graph theory and network analysis.
https://doi.org/10.5281/ZENODO.7682609

Denneboom, D., Bar-Massada, A., Shwartz, A., 2021. Factors affecting usage of crossing
structures by wildlife – A systematic review and meta-analysis. Sci. Total Environ.
777. https://doi.org/10.1016/j.scitotenv.2021.146061

Dunne, J.A., Williams, R.J., Martinez, N.D., 2002. Network structure and biodiversity loss in
food webs: robustness increases with connectance. Ecol. Lett. 5, 558–567.
https://doi.org/10.1046/j.1461-0248.2002.00354.x

Fedriani, J.M., Palomares, F., Delibes, M., 1999. Niche relations among three sympatric
Mediterranean carnivores. Oecologia 121, 138–148.
https://doi.org/10.1007/s004420050915

Felipe-Lucia, M.R., Soliveres, S., Penone, C., Fischer, M., Ammer, C., Boch, S.,
Boeddinghaus, R.S., Bonkowski, M., Buscot, F., Fiore-Donno, A.M., Frank, K.,
Goldmann, K., Gossner, M.M., Hölzel, N., Jochum, M., Kandeler, E., Klaus, V.H.,
Kleinebecker, T., Leimer, S., Manning, P., Oelmann, Y., Saiz, H., Schall, P., Schloter,
M., Schöning, I., Schrumpf, M., Solly, E.F., Stempfhuber, B., Weisser, W.W.,
Wilcke, W., Wubet, T., Allan, E., 2020. Land-use intensity alters networks between
biodiversity, ecosystem functions, and services. Proc. Natl. Acad. Sci. 117, 28140–
28149. https://doi.org/10.1073/pnas.2016210117

Forman, R., Alexander, L.E., 1998. Roads and Their Major Ecological Effects. Annu. Rev.
Ecol. Syst. 29, 207–231. https://doi.org/10.1146/annurev.ecolsys.29.1.207

Gachassin, M., Najman, B., Raballand, G., 2010. The impact of roads on poverty reduction: a
case study of Cameroon. The World Bank.

Gaston, K.J., 2010. Valuing Common Species. Science 327, 154–155.
https://doi.org/10.1126/science.1182818

Gaston, K.J., Fuller, R.A., 2008. Commonness, population depletion and conservation
biology. Trends Ecol. Evol. 23, 14–19. https://doi.org/10.1016/j.tree.2007.11.001

Grilo, C., Coimbra, M.R., Cerqueira, R.C., Barbosa, P., Dornas, R.A.P., Gonçalves, L.O.,
Teixeira, F.Z., Coelho, I.P., Schmidt, B.R., Pacheco, D.L.K., Schuck, G., Esperando,
I.B., Anza, J.A., Beduschi, J., Oliveira, N.R., Pinheiro, P.F., Bager, A., Secco, H.,
Guerreiro, M., Carvalho, C.F., Veloso, A.C., Custódio, A.E.I., Marçal, O., Ciocheti,
G., Assis, J., Ribeiro, M.C., Francisco, B.S.S., Cherem, J.J., Trigo, T.C., Jardim,
M.M.A., Franceschi, I.C., Espinosa, C., Tirelli, F.P., Rocha, V.J., Sekiama, M.L.,
Barbosa, G.P., Rossi, H.R., Moreira, T.C., Cervini, M., Rosa, C.A., Silva, L.G.,
Ferreira, C.M.M., César, A., Casella, J., Mendes, S.L., Zina, J., Bastos, D.F.O., Souza,
R.A.T., Hartmann, P.A., Deffaci, A.C.G., Mulinari, J., Luzzi, S.C., Rezzadori, T.,
Kolcenti, C., Reis, T.X., Fonseca, V.S.C., Giorgi, C.F., Migliorini, R.P., Kasper, C.B.,
Bueno, C., Sobanski, M., Pereira, A.P.F.G., Andrade, F.A.G., Fernandes, M.E.B.,
Corrêa, L.L.C., Nepomuceno, A., Banhos, A., Hannibal, W., Fonseca, R., Costa, L.A.,
Medici, E.P., Croce, A., Werther, K., Oliveira, J.P., Ribeiro, J.M., Santi, M. de,
Kawanami, A.E., Perles, L., Couto, C. do, Figueiró, D.S., Eizirik, E., Correia, A.A.,
Corrêa, F.M., Queirolo, D., Quagliatto, A.L., Saranholi, B.H., Galetti, P.M.,
Rodriguez☐Castro, K.G., Braz, V.S., França, F.G.R., Buss, G., Rezini, J.A., Lion,
M.B., Cheida, C.C., Lacerda, A.C.R., Freitas, C.H., Venâncio, F., Adania, C.H.,
Batisteli, A.F., Hegel, C.G.Z., Mantovani, J.A., Rodrigues, F.H.G., Bagatini, T., Curi,
N.H.A., Emmert, L., Erdmann, R.H., Costa, R.R.G.F., Martinelli, A., Santos, C.V.F.,

Kindel, A., 2018. BRAZIL ROAD-KILL: a data set of wildlife terrestrial vertebrate road-kills. Ecology 99, 2625–2625. https://doi.org/10.1002/ecy.2464

Grilo, C., Ferreira, F.Z., Revilla, E., 2015. No evidence of a threshold in traffic volume affecting road-kill mortality at a large spatio-temporal scale. Environ. Impact Assess. Rev. 55, 54–58. https://doi.org/10.1016/j.eiar.2015.07.003

Grilo, C., Koroleva, E., Andrášik, R., Bíl, M., González Suárez, M., 2020. Roadkill risk and population vulnerability in European birds and mammals. Front. Ecol. Environ. 18. https://doi.org/10.1002/fee.2216

Heleno, R.H., Ripple, W.J., Traveset, A., 2020. Scientists' warning on endangered food webs. Web Ecol. 20, 1–10. https://doi.org/10.5194/we-20-1-2020

Hijmans, R.J., 2023. terra: Spatial Data Analysis. R package version 1.7-55, <https://CRAN.R-project.org/package=terra>.

Hudson, L.N., Emerson, R., Jenkins, G.B., Layer, K., Ledger, M.E., Pichler, D.E., Thompson, M.S.A., O'Gorman, E.J., Woodward, G., Reuman, D.C., 2013. Cheddar: analysis and visualisation of ecological communities in R. Methods Ecol. Evol. 4, 99–104. https://doi.org/10.1111/2041-210X.12005

Iimi, A., Lancelot, E., Manelici, I., Ogita, S., 2015. Evaluating the Social and Economic Impacts of Rural Road Improvements in the State of Tocantins, Brazil (No. 95574). World Bank Group, Washington, D.C.

Kramer Schadt, S., Revilla, E., Wiegand, T., Breitenmoser, U., 2004. Fragmented landscapes, road mortality and patch connectivity: modelling influences on the dispersal of Eurasian lynx. J. Appl. Ecol. 41, 711–723. https://doi.org/10.1111/j.0021-8901.2004.00933.x

Leighton, G.R.M., Froneman, W., Serieys, L.E.K., Bishop, J.M., 2023. Trophic downgrading of an adaptable carnivore in an urbanising landscape. Sci. Rep. 13, 21582. https://doi.org/10.1038/s41598-023-48868-x

Leroux, S.J., Loreau, M., 2015. Theoretical perspectives on bottom-up and top-down interactions across ecosystems, in: La Pierre, K.J., Hanley, T.C. (Eds.), Trophic Ecology: Bottom-Up and Top-Down Interactions across Aquatic and Terrestrial Systems, Ecological Reviews. Cambridge University Press, Cambridge, pp. 3–28. https://doi.org/10.1017/CBO9781139924856.002

Lesbarrères, D., Fahrig, L., 2012. Measures to reduce population fragmentation by roads: what has worked and how do we know? Trends Ecol. Evol. 27, 374–380. https://doi.org/10.1016/j.tree.2012.01.015

Liao, J., Bearup, D., Blasius, B., 2017. Diverse responses of species to landscape fragmentation in a simple food chain. J. Anim. Ecol. 86, 1169–1178. https://doi.org/10.1111/1365-2656.12702

Liu, O.R., Gaines, S.D., 2022. Environmental context dependency in species interactions. Proc. Natl. Acad. Sci. 119, e2118539119. https://doi.org/10.1073/pnas.2118539119

Maiorano, L., Montemaggiori, A., Ficetola, G.F., O'Connor, L., Thuiller, W., 2020. TETRA-EU 1.0: A species-level trophic metaweb of European tetrapods. Glob. Ecol. Biogeogr. 29, 1452–1457. https://doi.org/10.1111/geb.13138

Martín González, A.M., Dalsgaard, B., Olesen, J.M., 2010. Centrality measures and the importance of generalist species in pollination networks. Ecol. Complex. 7, 36–43. https://doi.org/10.1016/j.ecocom.2009.03.008

Maxwell, S.L., Fuller, R.A., Brooks, T.M., Watson, J.E.M., 2016. Biodiversity: The ravages of guns, nets and bulldozers. Nature 536, 143. https://doi.org/10.1038/536143a

Meijer, J.R., Huijbregts, M.A.J., Schotten, K.C.G.J., Schipper, A.M., 2018. Global patterns of current and future road infrastructure. Environ. Res. Lett. 13, 064006. https://doi.org/10.1088/1748-9326/aabd42

Memmott, J., Waser, N.M., Price, M.V., 2004. Tolerance of pollination networks to species
extinctions. Proc. R. Soc. B Biol. Sci. 271, 2605–2611.
https://doi.org/10.1098/rspb.2004.2909
Mestre, F., Rozenfeld, A., Araújo, M.B., 2022. Human disturbances affect the topology of
food webs. Ecol. Lett. 25, 2476–2488. https://doi.org/10.1111/ele.14107
Moore, L.J., Petrovan, S.O., Bates, A.J., Hicks, H.L., Baker, P.J., Perkins, S.E., Yarnell,
R.W., 2023. Demographic effects of road mortality on mammalian populations: a
systematic review. Biol. Rev. 98, 1033–1050. https://doi.org/10.1111/brv.12942
Morelli, F., Benedetti, Y., Delgado, J.D., 2020. A forecasting map of avian roadkill-risk in
Europe: A tool to identify potential hotspots. Biol. Conserv. 249, 108729.
https://doi.org/10.1016/j.biocon.2020.108729
Muhly, T.B., Hebblewhite, M., Paton, D., Pitt, J.A., Boyce, M.S., Musiani, M., 2013.
Humans Strengthen Bottom-Up Effects and Weaken Trophic Cascades in a Terrestrial
Food Web. PLoS ONE 8, e64311. https://doi.org/10.1371/journal.pone.0064311
Philips, I., 2020. An Agent Based Model to Estimate Lynx Dispersal if Re-Introduced to
Scotland. Appl. Spat. Anal. Policy 13, 161–185. https://doi.org/10.1007/s12061-019-
09297-4
Pires, M.M., Benchimol, M., Cruz, L.R., Peres, C.A., 2023. Terrestrial food web complexity
in Amazonian forests decays with habitat loss. Curr. Biol. 33, 389-396.e3.
https://doi.org/10.1016/j.cub.2022.11.066
Poisot, T., Mouquet, N., Gravel, D., 2013. Trophic complementarity drives the biodiversity–
ecosystem functioning relationship in food webs. Ecol. Lett. 16, 853–861.
https://doi.org/10.1111/ele.12118
Poisot, T., Stouffer, D.B., Gravel, D., 2015. Beyond species: why ecological interaction
networks vary through space and time. Oikos 124, 243–251.
https://doi.org/10.1111/oik.01719
Pratzer, M., Nill, L., Kuemmerle, T., Zurell, D., Fandos, G., 2023. Large carnivore range
expansion in Iberia in relation to different scenarios of permeability of human-
dominated landscapes. Divers. Distrib. 29, 75–88. https://doi.org/10.1111/ddi.13645
QGIS.org, 2020. QGIS Geographic Information System. Open Source Geospatial Foundation
Project. http://qgis.org.
R Development Core Team, 2022. R: A language and environment for statistical computing.
R Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-
project.org/.
Richmond, C.E., Breitburg, D.L., Rose, K.A., 2005. The role of environmental generalist
species in ecosystem function. Ecol. Model. 188, 279–295.
https://doi.org/10.1016/j.ecolmodel.2005.03.002
Rio-Maior, H., Nakamura, M., Álvares, F., Beja, P., 2019. Designing the landscape of
coexistence: Integrating risk avoidance, habitat selection and functional connectivity
to inform large carnivore conservation. Biol. Conserv. 235, 178–188.
https://doi.org/10.1016/j.biocon.2019.04.021
Ripple, W.J., Estes, J.A., Beschta, R.L., Wilmers, C.C., Ritchie, E.G., Hebblewhite, M.,
Berger, J., Elmhagen, B., Letnic, M., Nelson, M.P., Schmitz, O.J., Smith, D.W.,
Wallach, A.D., Wirsing, A.J., 2014. Status and Ecological Effects of the World's
Largest Carnivores. Science 343, 1241484. https://doi.org/10.1126/science.1241484
Ryser, R., Häussler, J., Stark, M., Brose, U., Rall, B.C., Guill, C., 2019. The biggest losers:
habitat isolation deconstructs complex food webs from top to bottom. Proc. R. Soc. B
Biol. Sci. 286, 20191177. https://doi.org/10.1098/rspb.2019.1177

Rytwinski, T., Fahrig, L., 2013. Why are some animal populations unaffected or positively
affected by roads? Oecologia 173, 1143–1156. https://doi.org/10.1007/s00442-013-
2684-x
Rytwinski, T., Fahrig, L., 2012. Do species life history traits explain population responses to
roads? A meta-analysis. Biol. Conserv. 147, 87–98.
https://doi.org/10.1016/j.biocon.2011.11.023
Sewell, S.J., Desai, S.A., Mutsaa, E., Lottering, R.T., 2019. A comparative study of
community perceptions regarding the role of roads as a poverty alleviation strategy in
rural areas. J. Rural Stud. 71, 73–84. https://doi.org/10.1016/j.jrurstud.2019.09.001
Steffen, W., Richardson, K., Rockström, J., Cornell, S.E., Fetzer, I., Bennett, E.M., Biggs, R.,
Carpenter, S.R., Vries, W. de, Wit, C.A. de, Folke, C., Gerten, D., Heinke, J., Mace,
G.M., Persson, L.M., Ramanathan, V., Reyers, B., Sörlin, S., 2015. Planetary
boundaries: Guiding human development on a changing planet. Science 347.
https://doi.org/10.1126/science.1259855
Tylianakis, J.M., Didham, R.K., Bascompte, J., Wardle, D.A., 2008. Global change and
species interactions in terrestrial ecosystems. Ecol. Lett. 11, 1351–1363.
https://doi.org/10.1111/j.1461-0248.2008.01250.x
Valiente-Banuet, A., Aizen, M.A., Alcántara, J.M., Arroyo, J., Cocucci, A., Galetti, M.,
García, M.B., García, D., Gómez, J.M., Jordano, P., Medel, R., Navarro, L., Obeso,
J.R., Oviedo, R., Ramírez, N., Rey, P.J., Traveset, A., Verdú, M., Zamora, R., 2015.
Beyond species loss: The extinction of ecological interactions in a changing world.
Funct. Ecol. 29, 299–307. https://doi.org/10.1111/1365-2435.12356
Van der Ree, R., Smith, D.J., Grilo, C., 2015. Handbook of road ecology. John Wiley &
Sons.
605

 Supplementary Information

 TABLE S1 – Information on species' interaction characteristics.

 Metadata of Table S1

ID	Focal species unique identification number
Species	Focal species
Class	Species class (Aves, Mammalia)
TLevel	Trophic level
Nint_total	Total number of interactions with this species
Nint_pred	Nº of interactions as predator
Nint_prey	Nº of interactions as prey
Mean.int	Mean number of interactions per region
SD.int	Standard deviation of interactions per region
Nint.at.risk_Direct	Nº of total interactions at risk due to direct roadkill (direct effects)
Nint.at.risk_Direct_as.pred	Nº of interactions at risk due to direct roadkill (direct effects) as predator
Nint.at.risk_Direct_as.prey	Nº of interactions at risk due to direct roadkill (direct effects) as prey
Nint.at.risk_Indirect	Nº of total interactions at risk due to indirect effects
Nint.at.risk_Indirect_as.pred	Nº of interactions at risk due to indirect effects as predator
Nint.at.risk_Indirect_as.prey	Nº of interactions at risk due to indirect effects as prey
Range	Range area in nº of cells
Range_area_risk_Direct	Range area in nº of cells at risk due to direct roadkill
Range_area_risk_Indirect	Range area in nº of cells at risk due to indirect effects
Range_area_risk_Direct%	Range area in % of cells at risk due to direct roadkill
Range_area_risk_Indirect%	Range area in % of cells at risk due to indirect effects