Deciphering lentiviral Vpr/Vpx determinants required for HUSH and SAMHD1 antagonism highlights the molecular plasticity of these evolutionary conflicts

Pauline Larrous[1], Cassandre Garnier[1], Marina Morel[1], Michael M. Martin[1], Karima Zarrouk[1], Sarah Maesen[2], Roy Matkovic[1], Andrea Cimarelli[2], Lucie Etienne[2*], Florence Margottin-Goguet[1*]

[1]Institut Cochin, Université Paris Cité, INSERM U1016, CNRS UMR8104, Paris, France
[2]CIRI, Centre International de Recherche en Infectiologie, Inserm U1111, Université Claude Bernard Lyon 1, CNRS UMR5308, Ecole Normale Supérieure de Lyon, Univ Lyon, Lyon, France

* Corresponding authors:

- Florence Margottin-Goguet (22 rue Méchain, 75014 Paris, France; tel: +33 1 40 51 66 16; florence.margottin-goguet@inserm.fr)

- Lucie Etienne (46 Allée d'Italie, 69364 Lyon Cedex 07, France ; +33 (0)4 72 72 86 96 ; lucie.etienne@ens-lyon.fr)

ABSTRACT

SAMHD1 and the HUSH complex constitute two blocks during primate lentivirus infection, the first by limiting reverse transcription and the second by inhibiting proviral expression. Vpr and Vpx of specific lentiviral lineages have evolved to antagonize these antiviral proteins. While the antagonism of SAMHD1 has been well characterized, the evolutionary and molecular determinants of the antagonism against HUSH are unknown. We used chimeric Vpr proteins between SIVagm.Ver and SIVagm.Gri lentiviruses infecting two African green monkey species, to investigate viral determinants involved in HUSH and SAMHD1 antagonisms. We found that different interfaces of closely related Vpr proteins are engaged to degrade different SAMHD1 haplotypes. In addition, we identified distinct viral determinants in SIVagm.Ver Vpr for SAMHD1 and HUSH degradation. The substitution of one residue in SIVagm.Gri Vpr is sufficient to gain the capacity to degrade SAMHD1, while the substitution of α-helix-3 confers HUSH antagonism. We also found that Vpx proteins of HIV-2 from people living with HIV have different abilities to degrade HUSH. These phenotypes rely on small changes in either the N or C terminal part of Vpx, depending on the context. On the host side, we found that HIV-2 and SIVsmm Vpx degrading HUSH from human and vervet monkey cells cannot not degrade HUSH in owl monkey cells, suggesting some host species-specificity.

Altogether, we highlight the molecular plasticity and constraints of viral proteins to adapt to host restrictions. HUSH, like SAMHD1, may have been engaged in ancient and more recent coevolution with lentiviruses and a player in viral fitness.

IMPORTANCE

Antiviral host proteins, the so-called restriction factors, block lentiviruses at different steps of their viral life cycle. In return, primate lentiviruses may counteract these immune proteins to efficiently spread *in vivo*. HIV-2 and some SIVs, but not HIV-1, inactivate SAMHD1 and HUSH, two host antiviral proteins, thanks to their Vpx or Vpr viral proteins. First, we uncovered here viral determinants involved in the function of closely related Vpr proteins from SIVs of African green monkeys and of HIV-2 Vpx alleles from people living with HIV-2. We show how these small viral proteins differently adapted to SAMHD1 polymorphism or to HUSH restriction and highlight their molecular plasticity. Finally, the capacity of divergent lentiviral proteins, including HIV-2 Vpx, to induce the degradation of HUSH depends of the cell/host species. Altogether, our results suggest that HUSH has been engaged in a molecular arms-race along evolution, and therefore is a key player in host-pathogens interaction.

KEYWORDS

HIV, SIV, restriction factors, coevolution, HUSH, SAMHD1, viral antagonism

INTRODUCTION

Host restriction factors are antiviral proteins from the cell autonomous immunity that have been engaged in an evolutionary arms-race with the pathogenic viruses they have been fighting for millions of years (1, 2). They further represent molecular barriers to cross-species transmission of viruses (3-6). When transmitted to humans, the lentiviruses SIVcpz and SIVgor (from chimpanzees and gorillas, respectively) and SIVsmm (from sooty mangabeys) gave rise to HIV-1 and HIV-2, respectively. The different lentiviral lineages share a similar genomic organization, but differ in their set of accessory genes, which produce proteins largely dedicated to the counteraction of restriction factors and which have strongly evolved during lentiviral cross-species transmissions (6, 7). Determining the exact molecular residues/interfaces underlying these conflicts is therefore a major objective to better understand HIV-cell interactions and determinants of virus spillover.

All extant primate lentiviruses, including SIVagm (infecting African green monkeys), encode Vpr, which induces host G2/M cell cycle arrest (8-10). However, despite its importance in the dissemination and pathogenesis of SIVsmm (11), Vpx is found in only two of the eight major lineages of primate lentiviruses, HIV-2/SIVmac/SIVsmm (infecting humans, macaques and sooty mangabeys) and SIVrcm/mnd2 (infecting red-capped mangabeys and mandrills). *Vpr* and *vpx* genes are the results of duplication and recombination events of a precursor gene (reviewed in (12)). The encoded proteins share similarities in size (about 100 amino acids), structure (a N-terminal tail, 3 α-helices and a C-terminal tail) and functions. Nonetheless, they also present highly variable regions: VR1 upstream of helix 1, VR2 across the end of helix 2 and the beginning of helix 3, and VR3, which overlaps the C-terminal tail. In viral lineages that encode both Vpr and Vpx, Vpx induces the proteasomal degradation of the host restriction factor SAMHD1 (SAM and HD domain-containing protein 1), while in some other lineages that do not encode Vpx, such as SIVagm, the Vpr protein performs this function (13-15). Associated phylogenetic analyses showed that an ancestral Vpr protein acquired the anti-SAMHD1 activity prior to the molecular events that gave birth to Vpx (15). SAMHD1 is a 626 amino acid dNTPase that blocks viral DNA synthesis by lowering the pool of nucleotides in macrophages and quiescent CD4+ T cells (16, 17). By degrading SAMHD1, Vpx/Vpr proteins enable the virus to bypass a reverse transcription block. In this process, Vpx/Vpr directly binds SAMHD1 and bridges SAMHD1 to the DCAF1 adaptor of a Cullin4A-based ubiquitin ligase (13, 14, 18-21).

How Vpx interacts with SAMHD1 in a virus-host species-specific manner has been

extensively studied. Strikingly, search for host determinants revealed that HIV-2/SIVsmm

Vpx targets the C-terminus of SAMHD1, while SIVmnd2 and SIVrcm Vpx recognize its N-

terminus (22-24). The divergence in SAMHD1 recognition is further witnessed by the

presence of sites under positive selection during primate evolution in both N- and C-terminal

domains of the protein (15, 22, 25). These site-specific adaptations in SAMHD1 are the

results of host escape from viral antagonism, characteristic of a molecular virus-host arms-

race.

The resolution of crystal structures and functional studies have allowed the identification of

the interfaces between Vpx from different lineages and SAMHD1 (23, 24). In particular, a

cluster of residues in the Vpx α-helix 2 of SIVmnd2 are in contact with SAMHD1, while

these amino-acids are not involved in the case of Vpx from SIVsmm (23, 24).

By studying the coevolution between African green monkeys (AGMs) and their SIVs, Spragg

and Emerman showed that SAMHD1 antagonism is crucial for viral fitness (26). AGMs

comprise at least four closely-related species : *Chlorocebus tantalus*, *sabaeus*, *aethiops*

(Grivet) and *pygerythrus* (Vervet). Each lineage of SIVagm, responsible of the natural

infection of each species, has evolved to antagonize distinct SAMHD1 haplotypes through the

use of Vpr (26). More precisely, among the seven SAMHD1 haplotypes identified in the

AGMs, haplotype IV, but not V, is degraded by SIVagm.Ver Vpr, while the opposite is found

for SIVagm.Gri Vpr; haplotype III is resistant to both Vprs, but is sensitive to SIVagm.Sab

Vpr (26).

In addition to SAMHD1, HIV-2/SIVsmm Vpx and the Vpr from specific lineages can induce

the degradation of the human HUSH complex (27, 28). The HUSH complex is composed of

TASOR, MPP8 and periphilin and contributes to the silencing of cellular genes and

retroelements with the help of MORC2(29). Due to HUSH degradation, Vpx/Vpr proteins

favor viral expression in a model of HIV-1 latency (27, 28). Functional and evolutionary

studies led us to conclude that HUSH antagonism is likely an ancient function of primate

lentiviruses that preceded the birth of Vpx and SAMHD1 antagonism (15). In addition,

HUSH antagonism appeared lentiviral species-specific, with only some Vpx/Vpr proteins

degrading human HUSH. Whether lentiviral species-specificity is accompanied by host

species-specificity, in line with host-virus competition along evolution, has not been

investigated yet in the case of HUSH.

Here, we took advantage of the lentiviral-host specificity within the AGM lineage, both for

SAMHD1 and HUSH antagonism, to identify viral determinants at stake. We found that the

closely related Vpr proteins use different viral determinants to degrade different SAMHD1

haplotypes, highlighting the molecular plasticity and adaptation of the virus to the host. In
addition, viral determinants against HUSH are different from those against SAMHD1. We
further studied Vpx proteins of HIV-2 from people living with HIV-2 (PLWH-2), which were
competent for SAMHD1 degradation (30). We found that some Vpx proteins can induce
HUSH degradation, while others cannot, and this diversity seems independent of the
corresponding status of the viremia. Depending on the Vpx, restoration of the capacity to
induce HUSH degradation was obtained by changing either the N-terminal part or the C-
terminal part of the viral protein.

Finally, we describe one evidence of host-species specificity with HIV-2/SIVsmm Vpx
unable to counteract HUSH in New World monkey cells. Altogether, our results suggest the
existence of a dynamic interplay between HUSH, SAMHD1 and primate lentiviruses along
evolution.

MATERIALS AND METHODS

Plasmids

Vpr SIVagm.ver9063 (KF741096), Vpr SIVagm.gri677 (sequence as the one used in (26)),
Vpr SIVagm.Tan1 (U58991) and Vpr SIVagm.Sab1 (S46351) chimera 1 to 10 (sup Fig 1),
Vpx proteins of HIV-2 from PLWH-2 and Vpx chimeras have been synthesized after codon-
optimisation and subcloned into the pAS1b vector (pAS1b-HA) to get HA-epitope-tagged
proteins (HA at the N-terminus). Vpr SIVagm.Tan1 is also expressed in pCDNA3-3xFlag
vector with a Flag epitope at the N-terminus. The mutants of Vpr were produced by site-
directed mutagenesis according to Phusion polymerase manufacture guide (Thermofisher) or
CloneAmp HiFi polymerase manufacture guide (Takkara) using Vpr SIVagm.ver9063 or
chimera 8 or 9 in the pAS1B vector as templates. Lentiviral proteins Vpx HIV-2 Gh
(P18045.1), Vpx SIVsmm (P19508.1), Vpx SIVmnd2.GAx14 (AAK82846.1), Vpx
SIVrcm.NG411 (AAK69676.1) and Vpx SIVrcm.Gab1 (AAM34564.1) are also expressed
from the pAS1b vector (HA tag at the N-terminus). Human TASOR (NP_001106207.1) and
Owl monkey TASOR (NCBI References Sequences: XP_012316204) are expressed from
vectors pLenti-Flag and pCMV6-Flag respectively, in fusion with the myc-DDK epitope at
the C-terminus. Constructs expressing Haplotypes III, IV and V of SAMHD1 from AGMs are
gifts from M. Emerman and are expressed with a HA epitope at the C-terminus from the
pLPCX vector (KF741043, KF741044 and KF741045).

Cell culture

Cells were tested regularly for mycoplasma contaminations; experiments were only

performed on non-contaminated cells. ATCC-purchased HeLa (CCL-2), VERO (CCL-81),

HEK293T (CLR-3216), HEK293FT (293T cells optimized for VLP production, gift from N.

Manel) and OMK cells were cultivated in media DMEM (Thermofisher) containing 10% fetal

bovine serum (FBS, Eurobio), 1,000 units/mL penicillin, 1,000μg/mL streptomycin (Life

Technologies). J-Lat A1 (gift from E. Verdin) were cultivated in media RPMI (Thermofisher)

supplemented as described for the DMEM medium and, in addition, 2mM glutamine (Life

Technologies). HEK293T stable cell lines expressing the haplotypes III, IV and V of AGM

SAMHD1 were generated by transduction of VLP containing pLPCX-Hap III, pLPCX-Hap

IV and pLPCX-Hap V and cultivated four days before puromycin selection.

siRNA treatment

siRNA transfections were performed with DharmaFECT1 (Dharmacon, GE Lifesciences).

The final concentration for all the siRNA was 40nM. The following siRNA were purchased:

siTASOR: SASI_Hs02_00325516 (Sigma Aldrich); siDCAF1: J-021119-10-0005

(Dharmacon) and the non-targeting control siRNA: MISSION siRNA Universal Negative

Control 1, SIC001 (Sigma Aldrich).

Virus-Like-Particle production, delivery and transduction

VLPs were produced in HEK293FT cells by co-transfection by the calcium-phosphate co-

precipitation method of VSV-G plasmid (3μg), SIV3+ ΔVpr ΔVpx vectors (8μg) and 8μg of

pAS1b-HA-Vpr (or chimeric proteins or point mutants) or pAS1b-HA-Vpx or pAS1b-HA

(empty) and in some experiments with a transfer gene pGAE1.0 (SIVmac-CMV-GFP) used as

a reporter gene. SIV3+ ΔVpr ΔVpx packaging vector is a gift from N. Landau and is

described in Gramberg *et al* (31). VLPs used for the establishment of the HEK293T stable

cell lines expressing HA-SAMHD1 (agm) were produced with VSV-G plasmid (3μg),

pHIT60 MLV packaging vector (8μg) and 8μg of pLPCX vector expressing Hap III or Hap

IV or Hap V. In both cases, 3.10^6 cells were plated in 10cm culture dishes the day prior

transfection. Cell culture medium was collected 48h after transfection and filtered through

0,45μm pores filters. For SAMHD1 VLP, 10mM NaBu were added 24h after transfection and

the cells washed at the end of the day. VLP were concentrated by sucrose gradient and

ultracentrifugation (1h30 at 100,000g). The incorporation of the viral proteins was assessed

by western blot and a quantification of the level of HIV-2 capsid (P27) and HA-Vpr was

performed to deliver the same quantity of viral proteins. J-Lat A1 cells were treated with VLP

for 7h in reduced medium prior to overnight TNFα (1ng/mL) treatment. VERO and OMK

cells were plated in 12-well dishes at a density of 3.10^5 cells and transduced the day after in

reduced medium, cells were harvested the day after transduction.

Flow cytometric analyses

204   J-Lat A1 cells were collected and resuspended in PBS-EDTA (0,5mM). Data were collected

205   and analyzed with BD Accuri C6 cytometer or Attune and software CFlow Plus or FlowJo. At

206   least 10,000 events in P1 were collected, the GFP-positive population was determined using

207   untreated J-Lat A1 cells as their level of GFP expression is low. The same gate was

208   maintained for all conditions and analysis were performed overall GFP-positive population.

SAMHD1 degradation assay

HEK293T Hap III, Hap IV and Hap V were plated in 12-well dishes at $1,5.10^5$ cells and

transfected the following day using the calcium-phosphate co-precipitation method. Different

amounts of the pAS1B vector expressing the different Vpr proteins (1 to 2µg) were

transfected to get the same level of expression of the viral proteins (with adjustment to get the

same level of total DNA per condition). Cells were harvested 48h post-transfection for

western blot analysis. Lysis of the cells was performed in 100µL of RIPAC buffer (50mM

Tris-HCl pH7.5, 150mM NaCl, 10% Glycerol, 2mM EDTA, 0.5% NP40, 0.1% SDS)

containing an anti-protease cocktail (A32965, Thermofisher), lysates were centrifuged at

16,000g for 10min to remove cell debris.

Cell fractionation

All immunoprecipitation experiments are performed in the nuclear fraction of HeLa cells.

Cells grown in 10cm dishes were washed with cold Dulbecco's PBS 1x (ThermoFisher).

After trypsinization (Thermofisher), cells were recovered in 1,5mL tubes and washed once

with ice-cold PBS. After 4 min of centrifugation at 400xg, 1mL of cytoplasmic lysis buffer

(10 mM TRIS-HCl pH7.5, 10 mM NaCl, 3 mM MgCl2, and 0.5% IGEPAL® CA-630 (I8896-

100ML Merck)) was added on the cell pellet and resuspended pellet was incubated on ice for

5 min. Cells were then centrifuged at 300g for 4 min at 4 °C and the supernatant was saved

for cytoplasmic fraction. The pellet was washed with 1mL of cytoplasmic lysis buffer and re-

centrifuged at 300g for 4 min at 4 °C. Finally, the nuclear pellet was lysed with 300µL of

RIPA buffer.

Immunoprecipitation assay, western blot procedure and antibodies

For HA-Vpr (AGM WT or chimeric proteins) and TASOR-Flag immunoprecipitations, HeLa cells were plated at respectively $2,5.10^6$ cells in 10cm dishes and co-transfected by the calcium-phosphate co-precipitation method with pAS1b-HA or pAS1b-HA-Vpx or Vpr (SIVagm WT or chimeric proteins) and plenti-TASOR-FLAG or pCMV6-TASOR-FLAG. Cells were harvested 48h post-transfection for western blot analysis. Lysis of the cells was performed in 700µL of RIPA buffer (50mM Tris-HCl pH7.5, 150mM NaCl, 10% Glycerol, 2mM EDTA, 0.5% NP40) containing an anti-protease cocktail (A32965, Thermofisher) and spun at 16,000g for 10min to remove cell debris. 500µg of cell lysates were incubated with pre-washed EZview Red ANTI-HA or FlagM2 affinity Gel Beads (E6779 and F2426, Merck) at 4 °C under overnight rotation. After three washes in wash buffer (50mM Tris-HCl pH7.5, 150mM NaCl), immunocomplexes were eluted with Laemmli buffer 1X with 20mM DTT and were separated by SDS-PAGE (Bolt Bis-Tris, 4-12%, Life Technologies). Following transfer onto PVDF membranes, proteins were revealed by immunoblot and signals were acquired with Fusion FX (Vilber Lourmat). The following antibodies, with their respective dilution in 5% skimmed milk in PBS-tween 0.1% were used: anti-HA-HRP (3F10) (N°12013819001, Roche) 1/10,000; anti-FLAG-HRP (A-8592, lot 61K9220, Sigma) 1/10,000; anti-HA (HA-7, H3663, lot 066M4837V, Merck) 1/1,000; anti-Flag M2 (F1804-200UG- lot SLCD3990, Merck) 1/1,000; anti-TASOR (HPA006735, lots A106822, C119001, Merck) 1/1,000; anti-DCAF1 (11612-1-AP, ProteinTech) 1/1000; Anti-p27/p55 and anti-P24 were provided by the NIH AIDS research and reference reagent program (ref ARP392/393) 1/1000; anti-βActin (AC40, A3853, Merck) 1/1000; anti-GAPDH (6C5, SC- 32233, Santa Cruz) 1/1,000. All HRP-conjugated secondary antibodies, anti-mouse (31430, lot VF297958, Thermofisher) and anti-rabbit (31460, lots VC297287, UK293475 Thermofisher), were used at a 1/20,000 dilution before reaction with Immobilon Classico (WBLUC0500, Merck Millipore) or Forte (WBLUF0100, Merck Millipore) Western HRP.

Immunofluorescence assay

HeLa cells were cultivated on glass side and transfected as explained before. Cells were fixed with 4% paraformaldehyde for 15 min and permeabilized with 0.1% Triton for 15 min. Blocking step was performed with 2% bovine serum albumin solution for 1h at room temperature. Cells were incubated with antibody against anti-hemagglutinin mouse IgG monoclonal conjugate Alexa Fluor 488 conjugated (Invitrogen) for 1h. The cells were washed and the nuclei counter stained with 40-6-diamidino-2-phenylindole (DAPI, Sigma-Aldrich)

for 20 min. The coverslip cells were mounted with Prolong gold/diamond antifade reagent

(Invitrogen). Immunofluorescence images were captured by using a Leica DMI6000 confocal

microscope at the IMAG'IC core facility.

RESULTS

To identify viral determinants involved in SAMHD1 and HUSH antagonism, we leveraged

the lentiviral-host specificity within the SIV-AGM lineage. Among the seven SAMHD1

haplotypes identified in the AGM population (26), we chose to study three haplotypes (III, IV

and V) that present no more than four amino acid differences and that have been described as:

abundant in the Vervet but very rare in the Grivet AGM species (haplotype IV), or abundant

in the Grivet and absent in the Vervet AGM species (haplotype V), or absent/rare in the

Grivet/Vervet AGM species (haplotype III) (Fig. 1A). The ability of Vpr proteins from

SIVagm viruses to degrade the different SAMHD1 haplotypes was then examined (Fig. 1A

and 1B). To this end, 293T stable cell lines encoding the different HA-tagged SAMHD1

haplotypes were first established and Vpr was then expressed ectopically by transient DNA

transfection (primary sequences of Vpr proteins in Sup Fig. 1). Of note, the sequence of

SIVagm.Gri Vpr (SIVagm.grv677 Vpr) harbors two amino-acid changes (A2T and R102G)

compared to the sequence of the original protein (NCBI RefSeq NP_054371.1), which were

acquired after virus isolation (26, 32). The sequence of SIVagm.Ver Vpr is from

SIVagm.Ver9063 (GenBank KF741096.1). Under these conditions, the Vpr protein derived

from the SIVagm.Ver9063 was able to induce the degradation of SAMHD1 haplotype IV, but

not haplotype V (Fig. 1B, quantification on the right). Conversely, the Vpr protein derived

from the SIVagm.grv677 led to the degradation of SAMHD1 haplotype V but not haplotype

IV (Fig. 1B). As previously shown by Spragg and Emerman (26), neither of the two Vpr

proteins induced the degradation of SAMHD1 haplotype III, which is predominantly found in

Sabaeus AGM species (Fig. 1B). The observed specificities of Vpr proteins could not be

explained by major subcellular localisation differences, because both Vprs exhibited similar

distribution in the nucleus and the cytoplasm, following immunofluorescence and

biochemical fractionation experiments (Fig. 1C and 1D). However, the specificities could be

the result of post-translational modifications, as at least two bands were revealed in different

proportions for SIVagm.Ver Vpr and SIVagm.Gri Vpr, but this was not further investigated

(Fig. 1B and 1D). Overall, phenotypes of SAMHD1 degradation suggest the existence of

specific interactions between SAMHD1 haplotypes present in a given species and the Vpr

protein from the SIVagm infecting the same species.

SIVagm.Ver and SIVagm.Gri Vpr proteins share 69% of identity at the amino acid

level. To decipher the viral determinants involved in SAMHD1 antagonism, we constructed

Vpr protein chimeras by exchanging their different α-helices, N- and C-termini domains (NtD

and CtD) (Fig. 1E, Sup Fig. 1 for sequences). Some chimeras harbored a dimeric form in

addition to the monomeric form, but all were able to bind DCAF1 (Fig. 1F). Chimera 6 was

hardly present as a monomer, possibly impacting its activity (Fig. 1F). All chimeras able to

induce SAMHD1 haplotype IV degradation in human cells contained the CtD of SIVagm.Ver

Vpr (chimeras 1, 4, 5, 7) (Fig. 2A). The replacement of the CtD from SIVagm.Gri by the CtD

from SIVagm.Ver (chimera 9) conferred the SIVagm.Gri Vpr the ability to induce SAMHD1

haplotype IV degradation (gain-of-function, Fig. 2A). These results suggest that the integrity

of the C-terminal domain of SIVagm.Ver Vpr is a critical determinant for SAMHD1

haplotype IV degradation. Furthermore, the CtD of SIVagm.Tan Vpr was also able to confer

the SIVagm.Gri protein (chimera 10) the ability to degrade SAMHD1 haplotype IV (Fig. 2B).

Therefore, SIVagm Vpr proteins that target the C-terminus of SAMHD1 (SIVagm.Tan and

SIVagm.Ver Vpr (26)) rely on C-terminal Vpr determinants to induce SAMHD1 haplotype

IV degradation.

None of the viral chimeras led to consistent degradation of SAMHD1 haplotype V,

suggesting that determinants throughout the viral protein, or the protein's complete

conformation, are essential for this activity (Fig. 2C). Despite all chimeras were able to bind

DCAF1, we cannot exclude that the lack of degradation might result from a global structural

defect. Therefore, only chimeras able to degrade either SAMHD1 haplotype IV (1,4,5,7,9,10)

or TASOR (1,4,5,8, see below) were taken into account to draw conclusions. With this in

mind, our results show that different viral interfaces in SIVagm.Ver Vpr and SIVagm.Gri Vpr

are involved in the degradation of SAMHD1 haplotypes IV and V, respectively.

We then used the same Vpr chimeras to identify the viral determinants involved in the

degradation of TASOR, the core component of the HUSH complex (33). First, we assayed the

degradation of endogenous TASOR in AGM Vervet cells (VERO cells). Vpr proteins were

incorporated into Viral-Like Particles (VLPs) and then delivered in VERO cells. Vpr

incorporation into VLPs was quantified by Western-blot, and VLPs quantities were adjusted

to deliver similar amounts of viral proteins into cells (Fig. 3A). SIVagm.Ver, SIVagm.Sab

and SIVagm.Tan Vpr proteins were able to induce the degradation of endogenous TASOR in

Vervet cells, in contrast to the SIVagm.Gri Vpr (Fig. 3B). Of note, our TASOR antibody detected two bands in VERO cells, but subsequent TASOR siRNA experiment suggested that only the lower band was indeed corresponding to TASOR (Fig. 3B, right). As previously described, similar phenotypes of TASOR degradation were found in human J-Lat A1 cells (Fig. 3C), a Jurkat T-cell line derivative that harbors a latent HIV-1 mini-genome expressing GFP under the control of the proviral LTR promoter (27, 34). In this HIV-1 latency model, TASOR degradation correlated with an increase of the percentage of GFP-positive cells, indicative of the reactivation of the latent provirus (Fig. 3D). Chimeras were further tested in both cell types, VERO and human cells (Fig. 4). All chimeras were well incorporated into VLPs, except chimera 2 (Fig. 4A). Degradation phenotypes were the same in Vero and J-Lat A1 cells: all the Vpr constructs that induced TASOR degradation contained the SIVagm.Ver Vpr α-helix 3 (chimeras 1, 4, 5 and 8) (Fig. 4B for Vero cells and 4C for human J-Lat A1 cells). In particular, chimera 8, which harbors the SIVagm.Ver α-helix 3 in a SIVagm.Gri Vpr background, was competent for TASOR degradation (gain-of-function), indicating that the SIVagm.Ver Vpr α-helix 3 region is a key viral determinant for TASOR degradation. As expected, TASOR degradation correlated with reactivation in the J-Lat-A1 model (Fig. 4D). Altogether, the C-terminal domain of SIVagm.Ver Vpr appears as a critical determinant for SAMHD1 haplotype IV degradation, while α-helix 3 is a key determinant for TASOR degradation.

To further delineate key residues, we performed sequence analyses of the viral proteins in the two regions. We first searched for potential "loss of function" mutations, which would impair SAMHD1 haplotype IV degradation, in the few residues that differ between the CtD of SIVagm.Ver and SIVagm.Gri Vprs proteins (Fig. 5A). We made the corresponding Vpr mutants by changing residues in the CtD of chimera 9, thus producing chimera 9 Q94G, E97S and RANRA-APPP (with RANRA residues at position 110 changed to APPP). We also made the R102G mutant directly in the SIVagm.Ver Vpr. Of note, D104 and D119 were not changed, because they are both present in the phenotypically different Vprs. The resulting proteins were all able to induce SAMHD1 haplotype IV degradation, except SIVagm.Ver Vpr R102G (Fig. 5B). Furthermore, the reciprocal G102R change in SIVagm.Gri Vpr restored its ability to induce SAMHD1 haplotype IV degradation, highlighting R102 as a key residue for this activity (Fig. 5C). Of note, the G102R substitution allowed to recover the original sequence of SIVagm.Gri Vpr; the original protein is now

368   notified with an asterisk throughout the manuscript (i.e., SIVagm.grv677* Vpr). Results are

369   summarized in Figure 5D.

370   Similarly, we searched for potential "loss of function" mutations, which would impair

TASOR degradation, by analyzing differences in α-helix 3 (Fig. 6A). Six residues that were

different between SIVagm.Ver and SIVagm.Gri Vpr proteins in the α-helix 3 were changed

two by two. Incorporation of the corresponding viral proteins into VLPs was checked (Fig.

6B). All the corresponding mutants in the SIVagm.Ver Vpr background retained the capacity

to degrade human TASOR and to reactivate HIV-1 in the J-Lat A1 model (Fig. 6C and 6D).

Because several residues in α-helix 3 of Vpr/Vpx proteins are required for DCAF1 binding

(19, 23, 24), we wondered whether the defect in TASOR degradation could result from a

defect in DCAF1 binding. However, we found that SIVagm.Ver and SIVagm.Gri Vpr

proteins both bound DCAF1 suggesting that differences in TASOR degradation are not linked

to DCAF1 binding (Fig. 1F or 6E). Interestingly, exogenously-expressed Flag-tagged human

TASOR seemed to better interact with SIVagm.Ver Vpr than with SIVagm.Gri Vpr (Fig. 6E).

Vpr differences in TASOR degradation could then result from differences in TASOR binding.

Mutations two by two in α-helix 3 did not impair DCAF1 or TASOR-Flag binding in

agreement with degradation assays (Fig. 6E). Thus, α-helix 3 appears essential for TASOR

degradation, but we could not further narrow-down the specific residues.

386   Overall, recurrent adaptation cycles to antagonize restriction factors during lentiviral

387   evolution have selected different determinants (summarized in Fig. 5D).

389   We next explored the specificity and determinants against SAMHD1 and TASOR

within the HIV-2 clade, which is a single lentiviral lineage as opposed to SIVagms.

Specifically, we tested whether Vpx from different HIV-2 strains could have different

abilities to induce TASOR degradation. We tested a panel of seven HIV-2s from the A or B

pandemic groups (sequences Sup Fig. 1): control Vpx from Ghana-1 (group A), Rod (group

A) and JK (group B) strains and four Vpx alleles of HIV-2 derived from PLWH-2, which

were selected from a study by Yu *et al* as able to induce the degradation of SAMHD1 (30).

Two alleles were from effective controllers (EC: high CD4+T cells and undetectable viral

load for several years) and two from non-controllers (NC: low CD4 count and high viral load)

(30). Vpx from Ghana-1, Rod and JK strains, one NC Vpx (RH2-1-D8, group A) and one EC

Vpx (RH2-135C1, group A) were able to induce TASOR degradation in human J-Lat A1 cells

after VLP delivery and subsequently to reactivate HIV-1 in the J-Lat A1 model (Fig. 7). One

NC Vpx (RH2-7B3, group A) and one EC Vpx (RH2-22-1B4, group B) were impaired in

their ability to induce TASOR degradation and reactivation of the latent virus (Fig. 7, strongly

impaired for B3, poorly for B4). Therefore, the efficiency of Vpx-mediated TASOR

degradation and HIV reactivation was variable within the HIV-2 lineage and did not correlate

with the potency of viral control in PLWH-2. We further took advantage of these phenotypic

differences and constructed chimera to identify Vpx determinants. In particular, we used NC

Vpx RH2-1-D (competent for TASOR degradation) and NC Vpx RH2-7B3 (non competent).

The two share 84.8 % identity, with most differences (highlighted in bold in Fig. 8A) in two

regions at the N-ter (N-ter tail + first part of helix-1) and the C-ter (helix 3 + C-ter tail). We

found that the replacement of the N-ter region (chimera D8B3), but not the C-ter (chimera

B3D8B3), from Vpx RH2-7B3 by RH2-1-D8 led to TASOR degradation and reactivation of

latent HIV-1 (gain-of-function, Fig. 8; C: incorporation, D: degradation, E: reactivation),

showing that RH2-1-D8 Vpx N-ter has key determinants in TASOR antagonism. Yet, in the

context of another chimera (B4D8B4, Fig. 9), we revealed that the C-ter region was this time

important for improving TASOR degradation efficiency (Figure 9). Overall, the capacity to

trigger TASOR degradation is context dependent and relies on N-terminal and C-terminal

regions of the viral protein.

This study and previous work (27, 28) showed differences between Vpr/Vpx proteins

from various lentiviral lineages to counteract human TASOR ("lentiviral lineage-specificity").

To complete the virus-host heterologous assays, we next tested the ability of lentiviral

SIVagm Vpr proteins to degrade TASOR from divergent primate host species, using cells

from New World monkeys (NWMs; owl monkey *Aotus trivirgatus* (aotTri) kidney OMK

cells) in addition to cells used up to now from hominoids (human J-Lat A1 cells) and Old

World monkeys (OWMs; AGM Vervet Vero cells). Of note, in OMK cells, cyclosporin A

(CsA) was added before the delivery of lentiviral proteins to bypass the Trim-CypA block

responsible for capsid destabilization upon entry (36). First, we found that SIVagm Vpr

proteins displayed the same phenotypes in OMK cells as in VERO and J-Lat A1 cells in

presence of CsA (Fig. 10A: incorporation, Fig. 10B: degradation in OMK cells). Therefore,

we extended our panel of lentiviral proteins by testing Vpx proteins from the HIV-2/SIVsmm

lineage (HIV-2 Ghana-1 strain and SIVsmm Vpx). As previously shown by us and others (27,

28, 37), HIV-2 and SIVsmm Vpx induced human TASOR degradation (Fig. 10C); in

addition, we show here that these viral proteins present the same phenotypes in AGM Vervet

cells (VERO) (Fig. 10D). Moreover, although SIVagm.Ver, SIVagm.Tan and SIVagm.Sab

Vpr proteins had similar phenotypes in owl monkey, AGM and human cells, Vpx proteins

from HIV-2 and SIVsmm were unable to induce TASOR degradation in owl monkey cells as

opposed to in human and AGM Vervet cells (Fig. 10E). To further confirm TASOR

degradation phenotypes in owl monkey cells, we incorporated SIVagm.Ver Vpr and HIV-2

Vpx in SIV-derived GFP encoding viruses and found that GFP expression in owl monkey

cells was similar (Fig. 10F). This ruled out the hypothesis that Vpx delivery was impaired in

owl monkey cells, though we cannot exclude the possibility that the stability of the two viral

proteins may differ.

Altogether, on top of lentivirus-lineage specificity, we now have one evidence of host-species

specificity in the interplay between the HUSH complex and the lentiviral Vpr/Vpx proteins

(summary Fig. 10G).

Next, we set out to determine whether the apparent host-species specificity could

result from a defect of binding of Vpx to DCAF1 and/or TASOR. First, both SIVagm.Ver Vpr

and HIV-2 Vpx could interact with owl monkey DCAF1 (Fig. 11A). Unfortunately,

interaction of viral proteins with endogenous TASOR is difficult to detect. Thus, we

generated by gene synthesis a vector encoding a flag-tagged owl monkey TASOR. The

TASOR sequence of the *Aotus trivirgatus* species could not be obtained, we used the

publicly-available sequence from a closely related owl monkey species, *Aotus nancymaae*

(aotNan; XM_012460781.1). We conducted experiments involving the simultaneous

exogenous expression of Flag-tagged owl monkey (aotNan) or human TASOR with, or

without, the HA-tagged HIV-2 Vpx or SIVagm.Ver Vpr. By an anti-Flag co-

immunoprecipitation assay, we found that both HIV-2 Vpx and SIVagm.Ver Vpr exhibited

interaction with human Flag-TASOR but also with owl monkey (aotNan) Flag-TASOR (Fig.

11B). These findings suggest that HIV-2 Vpx is able to interact with owl monkey DCAF1 and

(aotNan) TASOR despite the lack of degradation of TASOR in owl monkey (aotTri) cells.

DISCUSSION

In the molecular arms-race between pathogenic viruses and their hosts, proteins are submitted

to strong selective pressures. Adaptive mutations in the host immune defense may result from

the escape from a viral antagonist, while adaptive changes in viral proteins may maintain the

virus' ability to counteract the host defense. Taking advantage of natural variations in

lentiviruses and host immune defenses from African green monkeys at the inter- and intra-

species levels, we show that closely related Vpr proteins can induce the degradation of

SAMHD1 variants (haplotypes IV and V) through distinct molecular determinants and that

the SIVagm.Ver Vpr protein uses also distinct determinants to trigger SAMHD1 and TASOR

degradation. Lastly, we discovered that HIV-2 Vpx proteins from people living with HIV

present distinct TASOR degradation abilities, which do not correlate with viremia. TASOR-

antagonism determinants are context dependent and are both in the N and C termini. Overall,

our results underline the high plasticity of Vpr and Vpx proteins to hijack ubiquitin ligase

complexes and to eliminate restrictive host proteins. Lastly, we have evidence of host species-

specificity in HUSH antagonism with HIV-2 and SIVsmm Vpx proteins able to induce

TASOR degradation in human and AGM VERO cells, but not in owl monkey OMK cells.

Only four amino acid differences specify the distinct behavior of the SAMHD1 haplotypes IV

and V, with respect to SIVagm Vpr proteins (26). We found that SIVagm.Ver and

SIVagm.Tan Vpr both use their C-ter tails to target SAMHD1 haplotype IV. Interestingly, in

the crystal structures of the ternary complexes of DCAF1, SAMHD1 and Vpx from SIVsmm,

which also targets the C-ter domain of SAMHD1, no residue of the C-terminal tail (also VR3)

of Vpx were found in contact with SAMHD1 (23, 24). Nonetheless, such structural studies

might not be adapted to Vpx C-ter tail, due to its flexible nature. It is also possible that the

recognition mechanism has diverged within the SIVagm lineage or that the C-terminal tail has

only a structural role in maintaining the recognition interface.

Our approach using chimeric proteins did not allow to reveal specific viral determinants of

SIVagm.Gri Vpr involved in haplotype V degradation, suggesting that the integrity of

different parts of the viral protein is required. The use of different interfaces by closely related

viral proteins underlines how antagonism of a given restriction factor results from different

modes of adaptation by the virus. Our results showing that the substitution of only one amino-

acid within a given Vpr protein restores its ability to induce the degradation of SAMHD1

further supports this model of molecular adaptation along evolution.

While the C-terminal tail of SIVagm.Ver Vpr conferred on the SIVagm.Gri protein the ability

to induce SAMHD1 haplotype IV degradation, the α-helix 3 of SIVagm.Ver Vpr conferred on

the SIVagm.Gri protein the ability to degrade the HUSH core protein TASOR. These results

suggest that the C-terminal tail and α-helix 3 are key determinants for SAMHD1 haplotype

IV and TASOR degradation, respectively, but they do not exclude the possibility that the

integrity of other determinants within the viral protein is required for the degradation of each

substrate. Indeed, as the SIVagm.Ver and SIVagm.Gri proteins are very similar in sequence,

chimera may share similar determinants important for the viral protein activity. In any event,

the use of distinct determinants for the degradation of two different substrates fits with a
model of ubiquitin ligase hijacking, in which a viral protein would induce the degradation of
different host factors using distinct viral interfaces for substrate recognition, but hijacking
only one type of ubiquitin ligase. Intriguingly, SIVagm.Gri Vpr could not induce HUSH
degradation, while the integrity of the whole protein seemed required for SAMHD1 haplotype
V degradation. Therefore, the selective pressure imposed on the "entire" virus protein to
counteract SAMHD1 may have limited its ability to adapt and counteract HUSH (27).
Alternatively, it is possible that SIVagm.Gri Vpr cannot degrade human, Vervet AGM or owl
monkey TASORs, but can degrade the TASOR from Grivet AGMs, in a host-species specific
manner. To address this question, all three components of HUSH (TASOR, MPP8 and
periphilin) should be analyzed for polymorphisms and variants within and between AGM
species and more extensively in primates, and tested for their degradation in the presence of
the different SIVagm Vpr proteins. It is also possible that important cofactors are
lacking/different in the cells from other hosts. Altogether, the use of different viral interfaces
between closely related Vpr proteins within the SIVagm lineage highlights the dynamism and
constraints of the molecular interactions between Vpr proteins, SAMHD1 and HUSH, as a
result of a cat-and-mouse game during evolution.

We further highlighted that Vpx proteins of HIV-2 from PLWH-2 harbor different
phenotypes. The ability to induce TASOR degradation can be restored after substitution of
residues either in the N or in the C-ter part (including helix 3) of the viral protein depending
on the Vpx, suggesting that different regions in Vpx could contribute to its interaction with
HUSH.

Our study also opens questions concerning the role of HIV-2 Vpx. By inducing HUSH
degradation, Vpx is able to increase the proviral transcription from the HIV LTR and the
stability of the LTR-driven transcript, thus Vpx counteracts the silencing of the provirus (38).
One could think at first that Vpx confers an advantage in HIV-2 replication by enhancing
LTR-driven RNA expression. However, interestingly, this study of Vpx proteins derived from
both viremic (NCs) or long-term aviremic (EV) PLWH-2 showed that the capacity to induce
TASOR degradation did not correlate with the viremia. The importance of TASOR
degradation over the course of HIV-2 infections remains to be investigated. In addition, we
cannot exclude that some HIV-2 strains may use other non-Vpx determinants in TASOR
antagonism.

Lastly, we determined here that HUSH antagonism presents some features of host-species
specificity in that the HIV-2/SIVsmm Vpx proteins could induce the degradation of human

and AGM vervet, but not of owl monkey TASOR. The underlying causes remain unknown. It is unlikely a defect in the viral protein delivery, because other Vpr/Vpx proteins were well delivered and functional in these cells, but we cannot exclude that HIV-2/SIVsmm Vpx may be particularly unstable in OMK cells. Our coIP results suggest that absence of degradation does not result from a lack of interaction with DCAF1 or TASOR (35). However, one limitation in our coIP experiments is that, due to challenges in sequencing the entire OMK TASOR, we here tested TASOR from a publicly-available sequence of *Aotus nancymaae* TASOR, which could be slightly different from that of OMK cells (*Aotus trivirgatus*).

Virus-host species-specificity is a hallmark of restriction factors. Antagonism of APOBEC3G by Vif occurs in a species-specific manner in Catharine primates and AGM populations, and functional evolutionary studies showed that the specificity of Vif reflects adaptation of the virus to the host including in key cross-species transmission events (3, 39-46). Experimental virus-host heterologous *in vivo* infection of AGMs showed that the lentiviral *vif* gene can adapt to Vif-resistant APOBEC3G haplotypes (43). Although not observed in the same experimental setting with Vpr (26), SIVagm Vpr proteins have acquired distinct interfaces to counteract SAMHD1, suggesting some adaptation of the virus to SAMHD1 selective pressure. The resistance of owl monkey TASOR to HIV-2/SIVsmm Vpx further suggests that HUSH may have evolved in some primate ancestors in response to past lentiviral epidemics. Alternatively, another pathogen, endogenous viral elements or drivers may have shaped primate TASOR resulting in some host species-specificity. More primate sequences in certain lineages and evolutionary analyses combined with functional assays would help to determine the modes and causes of such host evolution. Altogether, on one hand, SAMHD1 and HUSH antagonisms by Vpr/Vpx proteins appear to be critical components of primate lentiviral fitness witnessed by the complex dynamism of the interactions at stake. On the other hand, results from this study and others (30) suggest that Vpx-mediated SAMHD1 or HUSH antagonism does not always correlate with viral control in PLWH-2. Future work will help to resolve this apparent paradox.

ACKNOWLEDGMENTS

We thank all the members of the "Retrovirus, Infection and Latency team" for fruitful discussions. We also thank the members of the LP2L team at CIRI, Lyon, for support. We acknowledge the CYBIO, IMAG'IC core facilities of the Institut Cochin.

This work was supported by grants from the SIDACTION, the French Research Agency on

HIV and Emerging Infectious Diseases ANRS/MIE and Fondation pour la Recherche

Médicale (FRM, EQU202203014684 attributed to F.MG.). P.L. was supported by Université

de Paris Cité, M.M.M. by SIDACTION, C.G. by FRM (EQU202203014684 attributed to

F.MG.), K.Z. by ANRS/MIE, R.M. by SIDACTION and FRM (EQU202203014684

attributed to F.MG.). The work in the laboratory of L.E. is supported by grants from the

ANRS/MIE (#ECTZ118944 to LE) and SIDACTION (n°21-1-AEQ-12972-2 to LE and

FMG).

579 REFERENCES

1.   M. D. Daugherty, H. S. Malik, Rules of engagement: molecular insights from host-
virus arms races. *Annu Rev Genet* **46**, 677-700 (2012).

2.   N. K. Duggal, M. Emerman, Evolutionary conflicts between viruses and restriction
factors shape immunity. *Nat Rev Immunol* **12**, 687-695 (2012).

3.   L. Etienne *et al.*, The Role of the Antiviral APOBEC3 Gene Family in Protecting
Chimpanzees against Lentiviruses from Monkeys. *PLoS Pathog* **11**, e1005149 (2015).

4.   A. Kirmaier *et al.*, TRIM5 suppresses cross-species transmission of a primate
immunodeficiency virus and selects for emergence of resistant variants in the new
species. *PLoS Biol* **8** (2010).

5.   R. M. Pinto *et al.*, BTN3A3 evasion promotes the zoonotic potential of influenza A
viruses. *Nature* 10.1038/s41586-023-06261-8 (2023).

6.   D. Sauter, F. Kirchhoff, Key Viral Adaptations Preceding the AIDS Pandemic. *Cell Host
Microbe* **25**, 27-38 (2019).

7.   L. Etienne, B. H. Hahn, P. M. Sharp, F. A. Matsen, M. Emerman, Gene loss and
adaptation to hominids underlie the ancient origin of HIV-1. *Cell Host Microbe* **14**, 85-
92 (2013).

8.   V. Planelles *et al.*, Vpr-induced cell cycle arrest is conserved among primate
lentiviruses. *J Virol* **70**, 2516-2524 (1996).

9.   G. L. Stivahtis, M. A. Soares, M. A. Vodicka, B. H. Hahn, M. Emerman, Conservation
and host specificity of Vpr-mediated cell cycle arrest suggest a fundamental role in
primate lentivirus evolution and biology. *J Virol* **71**, 4331-4338 (1997).

10.   Y. Zhu *et al.*, Comparison of cell cycle arrest, transactivation, and apoptosis induced
by the simian immunodeficiency virus SIVagm and human immunodeficiency virus
type 1 vpr genes. *J Virol* **75**, 3791-3801 (2001).

11.   V. M. Hirsch *et al.*, Vpx is required for dissemination and pathogenesis of SIV(SM) PBj:
evidence of macrophage-dependent viral amplification. *Nat Med* **4**, 1401-1408
(1998).

12.   T. Schaller, H. Bauby, S. Hue, M. H. Malim, C. Goujon, New insights into an X-
traordinary viral protein. *Front Microbiol* **5**, 126 (2014).

13.   K. Hrecka *et al.*, Vpx relieves inhibition of HIV-1 infection of macrophages mediated
by the SAMHD1 protein. *Nature* **474**, 658-661 (2011).

14.   N. Laguette *et al.*, SAMHD1 is the dendritic- and myeloid-cell-specific HIV-1
restriction factor counteracted by Vpx. *Nature* **474**, 654-657 (2011).
15.   E. S. Lim *et al.*, The ability of primate lentiviruses to degrade the monocyte restriction
factor SAMHD1 preceded the birth of the viral accessory protein Vpx. *Cell Host*
Microbe **11**, 194-204 (2012).
16.   H. M. Baldauf *et al.*, SAMHD1 restricts HIV-1 infection in resting CD4(+) T cells. *Nat*
Med **18**, 1682-1689 (2012).
17.   H. Lahouassa *et al.*, SAMHD1 restricts the replication of human immunodeficiency
virus type 1 by depleting the intracellular pool of deoxynucleoside triphosphates. *Nat*
Immunol **13**, 223-228 (2012).
18.   A. Bergamaschi *et al.*, The human immunodeficiency virus type 2 Vpx protein usurps
the CUL4A-DDB1 DCAF1 ubiquitin ligase to overcome a postentry block in
macrophage infection. *J Virol* **83**, 4854-4860 (2009).
19.   E. Le Rouzic *et al.*, HIV1 Vpr arrests the cell cycle by recruiting DCAF1/VprBP, a
receptor of the Cul4-DDB1 ubiquitin ligase. *Cell Cycle* **6**, 182-188 (2007).
20.   J. Ahn *et al.*, HIV/simian immunodeficiency virus (SIV) accessory virulence factor Vpx
loads the host cell restriction factor SAMHD1 onto the E3 ubiquitin ligase complex
CRL4DCAF1. *J Biol Chem* **287**, 12550-12558 (2012).
21.   S. Srivastava *et al.*, Lentiviral Vpx accessory factor targets VprBP/DCAF1 substrate
adaptor for cullin 4 E3 ubiquitin ligase to enable macrophage infection. *PLoS Pathog*
4, e1000059 (2008).
22.   O. I. Fregoso *et al.*, Evolutionary toggling of Vpx/Vpr specificity results in divergent
recognition of the restriction factor SAMHD1. *PLoS Pathog* **9**, e1003496 (2013).
23.   D. Schwefel *et al.*, Molecular determinants for recognition of divergent SAMHD1
proteins by the lentiviral accessory protein Vpx. *Cell Host Microbe* **17**, 489-499
(2015).
24.   D. Schwefel *et al.*, Structural basis of lentiviral subversion of a cellular protein
degradation pathway. *Nature* **505**, 234-238 (2014).
25.   N. Laguette *et al.*, Evolutionary and functional analyses of the interaction between
the myeloid restriction factor SAMHD1 and the lentiviral Vpx protein. *Cell Host*
Microbe **11**, 205-217 (2012).
26.   C. J. Spragg, M. Emerman, Antagonism of SAMHD1 is actively maintained in natural
infections of simian immunodeficiency virus. *Proc Natl Acad Sci U S A* **110**, 21136-
21141 (2013).
27.   G. Chougui *et al.*, HIV-2/SIV viral protein X counteracts HUSH repressor complex. *Nat*
Microbiol **3**, 891-897 (2018).
28.   L. Yurkovetskiy *et al.*, Primate immunodeficiency virus proteins Vpx and Vpr
counteract transcriptional repression of proviruses by the HUSH complex. *Nat*
Microbiol **3**, 1354-1361 (2018).
29.   I. A. Tchasovnikarova *et al.*, GENE SILENCING. Epigenetic silencing by the HUSH
complex mediates position-effect variegation in human cells. *Science* **348**, 1481-1485
(2015).
30.   H. Yu *et al.*, The efficiency of Vpx-mediated SAMHD1 antagonism does not correlate
with the potency of viral control in HIV-2-infected individuals. *Retrovirology* **10**, 27
(2013).
31.   T. Gramberg, N. Sunseri, N. R. Landau, Evidence for an activation domain at the
amino terminus of simian immunodeficiency virus Vpx. *J Virol* **84**, 1387-1396 (2010).

32. P. R. Johnson *et al.*, Simian immunodeficiency viruses from African green monkeys
display unusual genetic diversity. *J Virol* **64**, 1086-1092 (1990).
33. C. H. Douse *et al.*, TASOR is a pseudo-PARP that directs HUSH complex assembly and
epigenetic transposon control. *Nat Commun* **11**, 4940 (2020).
34. A. Jordan, D. Bisgrove, E. Verdin, HIV reproducibly establishes a latent infection after
acute infection of T cells in vitro. *EMBO J* **22**, 1868-1877 (2003).
35. M. M. Martin *et al.*, Binding to DCAF1 distinguishes TASOR and SAMHD1 degradation
by HIV-2 Vpx. *PLoS Pathog* **17**, e1009609 (2021).
36. D. M. Sayah, E. Sokolskaja, L. Berthoux, J. Luban, Cyclophilin A retrotransposition into
TRIM5 explains owl monkey resistance to HIV-1. *Nature* **430**, 569-573 (2004).
37. E. J. D. Greenwood *et al.*, Promiscuous Targeting of Cellular Proteins by Vpr Drives
Systems-Level Proteomic Remodeling in HIV-1 Infection. *Cell Rep* **27**, 1579-1596
e1577 (2019).
38. R. Matkovic *et al.*, TASOR epigenetic repressor cooperates with a CNOT1 RNA
degradation pathway to repress HIV. *Nat Commun* **13**, 66 (2022).
39. H. P. Bogerd, B. P. Doehle, H. L. Wiegand, B. R. Cullen, A single amino acid difference
in the host APOBEC3G protein controls the primate species specificity of HIV type 1
virion infectivity factor. *Proc Natl Acad Sci U S A* **101**, 3770-3774 (2004).
40. R. Mariani *et al.*, Species-specific exclusion of APOBEC3G from HIV-1 virions by Vif.
Cell **114**, 21-31 (2003).
41. H. Xu *et al.*, A single amino acid substitution in human APOBEC3G antiretroviral
enzyme confers resistance to HIV-1 virion infectivity factor-induced depletion. *Proc*
Natl Acad Sci U S A **101**, 5652-5657 (2004).
42. A. A. Compton, M. Emerman, Convergence and divergence in the evolution of the
APOBEC3G-Vif interaction reveal ancient origins of simian immunodeficiency viruses.
PLoS Pathog **9**, e1003135 (2013).
43. A. A. Compton, V. M. Hirsch, M. Emerman, The host restriction factor APOBEC3G and
retroviral Vif protein coevolve due to ongoing genetic conflict. *Cell Host Microbe* **11**,
91-98 (2012).
44. M. D'Arc *et al.*, Origin of the HIV-1 group O epidemic in western lowland gorillas. *Proc*
Natl Acad Sci U S A **112**, E1343-1352 (2015).
45. M. Letko, T. Booiman, N. Kootstra, V. Simon, M. Ooms, Identification of the HIV-1 Vif
and Human APOBEC3G Protein Interface. *Cell Rep* **13**, 1789-1799 (2015).
46. M. Letko *et al.*, Vif proteins from diverse primate lentiviral lineages use the same
binding site in APOBEC3G. *J Virol* **87**, 11861-11871 (2013).

Figure legends

Figure 1: **Closely related Vpr proteins, which differently antagonize SAMHD1 variants,**

bind DCAF1

(A) Schematic representation of SAMHD1 haplotypes III, IV and V. The phenotype of their

degradation as shown in B is indicated on the right (v: degradation, x: no degradation). (B)

293T cells stably expressing HA-tagged SAMHD1 haplotypes III, IV and V were transfected

with plasmids encoding SIVagm Vpr proteins (ver9063: strain of SIVagm.Ver, grv677: strain

of SIVagm.Gri). (Left) The indicated proteins were revealed by western-blot (one

representative western-blot is shown). (Right) Quantification of SAMHD1 degradation rate

from 3 experiments. A one-sample t-test was performed. Data are presented as mean ±

standard deviation (SD). Statistical significance is indicated as follows: $*p < 0.05$, $**p < 0.01$,

$***p < 0.001$. (C) HA-Vpr proteins were co-expressed in HeLa cells cultured on glass slides.

Cells were fixed and stained with fluorescent probes for DNA (DAPI; blue), HA-Vpr (anti-

HA Alexa Fluor 488-conjugated; green). (Left) Images were acquired using confocal

microscopy, representative images showing the distribution of HA-Vpr are shown. (Right)

Quantification of the mean fluorescence intensity per cell (n = 30 cells) using ImageJ. Error

bars represent the standard deviation. No statistically significant difference in fluorescence

intensity was observed between SIVagm.Ver, and SIVagm.Gri Vpr ($p > 0.05$, Student's t-

test). (D) HeLa cells were transfected with plasmids encoding SIVagm Vpr proteins. Cells

were fractionated before the lysis of the nucleus and the cytoplasm. The indicated proteins

were revealed by western blot. (E) Schematic representation of SIVagm Vpr proteins and

chimeras. The scale is respected for the length of the different domains. (F) Indicated HA-Vpr

constructs were expressed in HeLa cells, then an anti-HA immunoprecipitation was

performed, and proteins were revealed by western-blot.

Figure 2: **Closely related Vpr proteins can induce the degradation of SAMHD1 variants**

(Haplotypes IV and V) through distinct molecular determinants

(A) 293T cells stably expressing HA-tagged SAMHD1 haplotype IV were transfected with

plasmids encoding SIVagm Vpr proteins and chimeras. (Top) A representative western blot

from 3 experiments is shown, (bottom) Quantification of SAMHD1 degradation rate from 3

experiments. A one-sample t-test was performed. Data are presented as mean ± standard

deviation (SD). Statistical significance is indicated as follows: $*p < 0.05$, $**p < 0.01$, $***p <$

0.001. (B) Same as in (A) but with different Vpr constructs as indicated. (C) Same as in (A)

but with cells stably expressing SAMHD1 haplotype V.

Figure 3: **SIV strains of African green monkey species differently degrade TASOR both**

in Vero and human cells

(A) VLPs were produced in 293FT by co-transfection of a packaging vector, an envelope

VSVg vector and a vector encoding HA- or Flag-Vpr as indicated. 48h post transfection,

supernatants were harvested and VLPs, concentrated by ultracentrifugation, were analyzed by

western blot. VLP production was checked with anti-P27 (capsid) antibody and HA-Vpr

incorporation with an anti-HA antibody. (B) (Left) VERO cells were treated with VLPs

containing SIVagm Vpr proteins as indicated and whole-cell extracts analyzed by western

blot. (Middle) Quantification of TASOR degradation rate from 3 experiments. A one-sample

t-test was performed. Data are presented as mean ± standard deviation (SD). Statistical

significance is indicated as follows: $*p < 0.05$, $**p < 0.01$, $***p < 0.001$. (Right) VERO cells

were treated with either siRNA CTRL or siRNA TASOR. (C) and (D) Human J-Lat A1 T

cells were treated with VLPs containing SIVagm Vpr proteins and stimulated overnight with

TNF-α, then cells were analyzed by western-blot and flow cytometry for the percentage of

GFP-positive cells. (C) (Left) Whole-cell extracts were analyzed by western blot, the

immunoblot shown is representative of at least 3 independent VLP productions. (Right)

Quantification of TASOR degradation rate from 3 experiments. A one-sample t-test was

performed. Data are presented as mean ± standard deviation (SD). Statistical significance is

indicated as follows: $*p < 0.05$, $**p < 0.01$, $***p < 0.001$. (D) The reactivation rate

corresponds to the percentage of GFP-positive cells in the presence of one viral protein over
the percentage obtained without viral protein (empty condition). Reactivation rates from 3
independent experiments are shown, along with raw reactivation data from one representative
experiment. A one-sample t-test was performed, with statistical significance denoted as: *p <
0.05, **p < 0.01, ***p < 0.001.

Figure 4: **α-helix 3 of SIVagm.Ver Vpr confers on the SIVagm.Gri protein the ability to**
induce the degradation of the HUSH core protein TASOR
(A) Incorporation of Vpr chimera proteins into VLPs as described in Figure 3A. (B) VERO
cells were treated with VLPs containing SIVagm Vpr proteins. After overnight treatment
cells were lysed. (Top) Whole-cell extracts were analyzed by western blot, the immunoblot
shown is representative of at least 3 independent VLP productions. (Bottom) Quantification
of TASOR degradation rate from 3 experiments. A one-sample t-test was performed. Data are
presented as mean ± standard deviation (SD). Statistical significance is indicated as follows:
*p < 0.05, **p < 0.01, ***p < 0.001. (C) Human J-Lat A1 T cells were treated with VLPs
containing SIVagm Vpr proteins. After overnight treatment with TNF-α, cells were analyzed
by flow cytometry for the percentage of GFP-positive cells. (Top) Whole-cell extracts were
analyzed by western blot, the immunoblot shown is representative of at least 3 independent
VLP productions. (Bottom) Quantification of TASOR degradation rate from 3 experiments. A
one-sample t-test was performed. Data are presented as mean ± standard deviation (SD).
Statistical significance is indicated as follows: *p < 0.05, **p < 0.01, ***p < 0.001. (D) The
reactivation rate corresponds to the percentage of GFP-positive cells in the presence of one
viral protein over the percentage obtained without viral protein (empty condition).
Reactivation rates from 3 independent experiments are shown (left), along with raw
reactivation data from one representative experiment (right). A one-sample t-test was
performed, with statistical significance denoted as: *p < 0.05, **p < 0.01, ***p < 0.001.

Figure 5: **The substitution of only one amino-acid in SIVagm.Gri Vpr restores its ability**
to induce the degradation of SAMHD1
(A) The C-terminal domain sequences from SIVagm.Ver9063, SIVagm.Grv677,
SIVagm.Grv677* (original sequence) and SIVagm.Tan Vpr proteins were aligned to point out
amino acid differences. Substitutions tested in SIVagm.Ver Vpr to assess a functional loss are
highlighted in grey. The key residue identified for haplotype IV SAMHD1 degradation is
shown in red. (B) 293T cells were transfected with plasmids coding for HA-tagged Vpr

proteins, chimera 9 or mutants, as indicated; (top) extracts from these cells were analyzed by
western blot, one representative experiment is shown. (bottom) Quantification of SAMHD1
degradation rate from 3 experiments. A one-sample t-test was performed. Data are presented
as mean ± standard deviation (SD). Statistical significance is indicated as follows: *p < 0.05,
**p < 0.01, ***p < 0.001. (C) SIVagm.Ver Vpr, SIVagm.Gri Vpr and SIVagm.Gri Vpr
G102R were overexpressed by transfection in 293T cells stably expressing the haplotype III
or IV of HA-tagged SAMHD1. (top) Extracts from these cells were analyzed by western blot,
one representative experiment is shown. (bottom) Quantification of SAMHD1 degradation
rate from 3 experiments. A one-sample t-test was performed. Data are presented as mean ±
standard deviation (SD). Statistical significance is indicated as follows: *p < 0.05, **p < 0.01,
***p < 0.001. (D) Schematic representation of key viral determinants involved in SAMHD1
and TASOR degradation.
816

Figure 6: **The whole integrity of SIVagm.Ver Vpr α-helix 3 contributes to the**
degradation of TASOR
(A) α-helix 3 sequences from SIVagm.Ver9063, SIVagm.Grv677, SIVagm.Grv677* (original
sequence) and SIVagm.Tan Vpr proteins were aligned to point out amino acid differences.
Substitutions tested in SIVagm.Ver to assess a functional loss are highlighted in grey. Key
residues identified for TASOR degradation are shown in red. (B) Same as Fig. 3A with
SIVagm.Ver9063 mutants. (C) Human J-Lat A1 T cells were treated with VLPs containing
Vpr AGM proteins. After overnight treatment with TNF-α, (top) whole-cell extracts were
analyzed by western blot (C) and cells were analyzed by flow cytometry for the percentage of
GFP-positive cells (D). (C) The immunoblot shown is representative of at least 3 independent
VLP productions. (bottom) Quantification of TASOR degradation rate from 3 experiments. A
one-sample t-test was performed. Data are presented as mean ± standard deviation (SD).
Statistical significance is indicated as follows: *p < 0.05, **p < 0.01, ***p < 0.001. (D) The
reactivation rate corresponds to the percentage of GFP-positive cells in the presence of one
viral protein over the percentage obtained without viral protein (empty condition).
Reactivation rates from 3 independent experiments are shown (left), along with raw
reactivation data from a representative experiment (right). A one-sample t-test was performed,
with statistical significance denoted as: *p < 0.05, **p < 0.01, ***p < 0.001. (E) Indicated
HA-Vpr constructs were expressed in HeLa cells, then an anti-HA immunoprecipitation was
performed, and proteins were revealed by western-blot.
837

Figure 7: **TASOR degradation by Vpx proteins from different origins, including HIV-2**

from PLWH-2

Human J-Lat A1 T cells were treated with VLPs containing Vpx proteins from different

origins (control Vpx from Ghana-1, Rod and JK strains and four HIV-2 Vpx alleles derived

from PLWH-2, according to Yu *et al.*) and overnight with TNF-α. Whole-cell extracts and

Vpx incorporation into VLPs were analyzed by western blot (A and B) and cells were also

analyzed by flow cytometry for the percentage of GFP-positive cells (C). (A) (Top) the

immunoblot shown is representative of at least 3 independent experiments; (bottom)

Quantification of TASOR degradation rate from 3 experiments. A one-sample t-test was

performed. Data are presented as mean ± standard deviation (SD). Statistical significance is

indicated as follows: *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$. (B) Incorporation of viral proteins

into the VLPs used in 9A is checked by western-blot. (C) The reactivation rate corresponds to

the percentage of GFP-positive cells in the presence of one viral protein over the percentage

obtained without viral protein (empty condition). Reactivation rates from 3 independent

experiments are shown (left), along with raw reactivation data from one representative

experiment (right).

Figure 8: **The N-ter region of Vpx RH2-1D8 confers on Vpx RH2-7B3 the ability to**

induce the degradation of the HUSH core protein TASOR

(A) Sequences from Vpx RH2-1D8 and Vpx RH2-7B3 were aligned to point out amino acid

differences (in red). Cross-referenced sequences for chimera are highlighted in light gray. (B)

Representation of the two indicated Vpx proteins and chimera. (C, D, E) Vpx incorporation

into VLPs, TASOR degradation in J-Lat A1 T cells following Vpx delivery by VLPs and

HIV-1 reactivation were assessed as in Figure 7.

Figure 9: **The C-ter region of Vpx RH2-1D8 confers on Vpx RH2-22-1B4 the ability to**

induce the degradation of the HUSH core protein TASOR

(A) Sequences from Vpx RH2-1D8 and Vpx RH2-22-1B4 were aligned to point out amino

acid differences (in red). The cross-referenced sequence for chimera is highlighted in light

gray. (B) Representation of the two indicated Vpx proteins and chimera. (C, D, E) Vpx

incorporation into VLPs, TASOR degradation in J-Lat A1 T cells following Vpx delivery by

VLPs and HIV-1 reactivation were assessed as in Figure 7.

Figure 10: **HIV-2/SIVsmm Vpx induces TASOR degradation in human and VERO cells,**
but not in Owl monkey cells
(A) Incorporation of Vpr and Vpx proteins into VLPs as described in Figure 3A. (B) OMK
cells were treated with VLPs containing the indicated Vpr/Vpx proteins. After overnight
treatment with cyclosporine CsA (or DMSO in control samples), whole-cell extracts were
analyzed by western blot. (C) Human J-Lat A1 T cells were treated with VLPs containing
Vpx proteins. After overnight treatment with TNF-α, whole-cell extracts were analyzed by
western blot, the immunoblot shown is representative of at least 3 independent VLP
productions. (D) Same as (C), but in VERO cells. (E) same as B, but with a different set of
viral proteins. (F) (Left) SIVagm.Ver Vpr or HIV-2 Vpx were incorporated into SIV-derived
viruses that express GFP following cell transduction. Vpx or Vpr incorporation was checked
by western blot on the VLPs. (Right) OMK cells were treated with the Vpx or Vpr-containing
SIV-derived viruses. After overnight treatment with CsA, whole-cell extracts were analyzed
by western blot. (G) Summary of the degradation phenotypes of the lentiviral proteins. Blue:
efficient degradation of TASOR and gray: no degradation.
886
Figure 11: **HIV-2 Vpx interacts with TASOR and DCAF1 in Owl monkey cells**
(A) Indicated HA-Vpr/Vpx constructs were expressed in OMK cells, then an anti-HA
immunoprecipitation was performed, and proteins were revealed by western-blot. (B)
SIVagm.Ver9063 Vpr or HIV-2.Gh1 Vpx were overexpressed in OMK cells with human or
owl Flag-TASOR, then an anti-Flag immunoprecipitation was performed, and proteins were
revealed by western-blot.

Larrous *et al*, Figure 1

Larrous *et al*, Figure 2

A Incorporation into VLPs

B VERO cells

* aspecific

* aspecific

C J-Lat A1 cells

D Reactivation rate in J-lat A1 cells (n=3)

Raw reactivation data for one sample (n=1)

Larrous *et al*, Figure 3

A *Incorporation into VLPs*

Vpr SIVagm:

30kDa HA-Vpr
P27
15kDa HA-Vpr

35kDa
25kDa
15kDa

B VERO cells

Vpr SIVagm:

TASOR

GAPDH

250 kDa

35kDa

* aspecific

C J-Lat A1 cells

VLP Vpr SIVagm:

TASOR
Actin

250kDa
42kDa

D

Reactivation rate in J-lat A1 cells (n=3)

Raw reactivation data for one sample (n=1)

Larrous *et al*, Figure 4

A

α-helix 2   α-helix 3

70   80

Vpr SIVagm.ver9033   **M** **T** E **R** **A** Y R Y Y R **L** V Q K **A** **L** F V H F R C
Vpr SIVagm.grv677   **W** **N** E **I** **G** Y K Y Y R **I** V Q K **S** **M** F V H F R C
Vpr SIVagm.grv677*   **W** **N** E **I** **G** Y K Y Y R **I** V Q K **S** **M** F V H F R C
Vpr SIVagm.tan1   **M** **I** E **R** **A** Y R Y Y R **L** V Q K **A** **L** F V H F R C

B

Incorporation into VLPs

VLP Vpr: empty   .ver9063   .grv677   .ver9063 M67WT68N   .ver9063 R70IA71G   .ver9063 A81SL82M

P27   25kDa

HA-Vpr/x   15kDa

C

J-lat A1 cells

empty   .ver9063   .grv677   .ver9063 M67WT68N   .ver9063 R70IA71G   .ver9063 A81SL82M

TASOR   250kDa

GAPDH   35kDa

TASOR degradation ratio

1.5

1.0

0.5

0.0

empty   .ver9063   .grv677   .ver9063 M67WT68N   .ver9063 R70IA71G   .ver9063 A81SL82M

****   ***   **   **

D

Reactivation rate in J-lat A1 cells (n=3)

Reactivation rate

1.6
1.4
1.2
1.0
0.8

empty   .ver9063   .grv677   .ver9063 M67WT68N   .ver9063 R70IA71G   .ver9063 A81SL82M

*   *   *   *

Raw reactivation data for one sample (n=1)

% of GFP positive cells

empty   .ver9063   .grv677   .ver9063 M67WT68N   .ver9063 R70IA71G   .ver9063 A81SL82M

E

Input   IP HA-Vpr

Vpr SIVagm: empty   .ver9063   .grv677   .ver9063 M67WT68N   .ver9063 R70IA71G   .ver9063 A81SL82M   empty   .ver9063   .grv677   .ver9063 M67WT68N   .ver9063 R70IA71G   .ver9063 A81SL82M
TASORhu-Flag: +   +   +   +   +   +   +   +   +   +   +   +

TASOR-Flag   250kDa

DCAF-1   250kDa

GAPDH   35kDa

Light Chains

HA-Vpr   15kDa

Larrous *et al*, Figure 6

A J-Lat A1 cells (Hominoïds)

VLP Vpr/x: empty .ver9063 HIV-2.Gh1 HIV-2.JK HIV-2A.Rod HIV-2 RH2-1-D8 (NC) HIV-2 RH2-7B3 (NC) HIV-2 RH2-135C1 (EC) HIV-2 RH2-22-1B4 (EC)

TASOR — 250kDa

GAPDH — 35kDa

B *Incorporation into VLPs*

VLP Vpr/x: empty HIV-2.Gh1 HIV-2.JK HIV-2A.Rod HIV-2 RH2-1-D8 (NC) HIV-2 RH2-7B3 (NC) HIV-2 RH2-135C1 (EC) HIV-2 RH2-22-1B4 (EC) .ver9063

P27 — 25kDa

30kDa Vpx — 30kDa

HA-Vpr/Vpx

15kDa Vpr/x — 15kDa

C *Reactivation rate in J-lat A1 cells (n=3)*

Raw reactivation data for one sample (n=1)

Larrous *et al*, Figure 7

Larrous _et al_, Figure 8

Larrous *et al*, Figure 9

Larrous *et al*, Figure 10

A OMK cells (NWM)

B 293T cells

Vpr SIVagm.ver9063 (GenBank: KF741096.1):
MASGRGPRENRPGEVEIWDLNREPWDEWLRDMLEDLNQEAKLHFGRELLFQVVWNYCQEEGERRGAPMTERAYRYYRLVQKALFVHFRCGCRRRQPFEPYEERRDGQGGGRANRAPPGLD

Vpr SIVagm.GRV677 (same as in Spragg and Emerman 2013):
MTSGRDPREPLPGWLEIWDLDREPWDEWLQDMLRDLNEEARRHFGMNMLIRVWNYCVEEGRRHNTPWNEIGYKYYRIVQKSMFVHFRCGCRRRGPFSPYEEGRNGQGGGAPPPPPGLA

Vpr SIVagm.GRV677 (NCBI Reference Sequence: NP_054371.1):*
MASGRDPREPLPGWLEIWDLDREPWDEWLQDMLRDLNEEARRHFGMNMLIRVWNYCVEEGRRHNTPWNEIGYKYYRIVQKSMFVHFRCGCRRRGPFSPYEERRNGQGGGAPPPPPGLA

Vpr SIVagm.SAB1 (PIR: S46351):
MASGGWLPPVGGDPPKDPPKNPREEIPGWLETWDLPREPFDEWLRDMLQDLNSEAQCHFPRNLLFRLWWNIVEEPAIDHGQTRLEGWYKYCRILQKALFVHMKGRCCKPKTHPAYGPGAGGPPPGLGGASGGAASAAPGL

Vpr SIVagm.Tan1 (GenBank: U58991.1):
MAEGRDSRERRPGWLEIWDLSREPWDEWLRDMVAELNQEAQRHFGRELLFQVVWNFCQEEGERNGAPMIERAYRYYRLVQKALFVHFRCGCRRRTPFEPYEERRNGVGGGRDGREPPGLA

Chimera 1: MTSGRDPREPLPGWLEIWDLDREPWDEWLQDMLRDLNEEARRHFGMNMLIRVWNYCVEEGRRHNTPMTERAYRYYRLVQKALFVHFRCGCRRRQPFEPYEERRDGQGGGRANRAPPGLD
Chimera 2: MASGRGPRENRPGEVEIWDLNREPWDEWLRDMLEDLNQEAKLHFGRELLFQVVWNYCQEEGERRGAPWNEIGYKYYRIVQKSMFVHFRCGCRRRGPFSPYEEGRNGQGGGAPPPPPGLA
Chimera 3: MASGRGPRENRPGEVEIWDLNREPWDEWLRDMLEDLNQEAKLHFGMNMLIRVWNYCVEEGRRHNTPWNEIGYKYYRIVQKSMFVHFRCGCRRRGPFSPYEEGRNGQGGGAPPPPPGLA
Chimera 4: MTSGRDPREPLPGWLEIWDLDREPWDEWLQDMLRDLNEEARRHFGRELLFQVVWNYCQEEGERRGAPMTERAYRYYRLVQKALFVHFRCGCRRRQPFEPYEERRDGQGGGRANRAPPGLD
Chimera 5: MASGRGPRENRPGEVEIWDLNREPWDEWLRDMLEDLNQEAKLHFGMNMLIRVWNYCVEEGRRHNTPMTERAYRYYRLVQKALFVHFRCGCRRRQPFEPYEERRDGQGGGRANRAPPGLD
Chimera 6: MTSGRDPREPLPGWLEIWDLDREPWDEWLQDMLRDLNEEARRHFGRELLFQVVWNYCQEEGERRGAPWNEIGYKYYRIVQKSMFVHFRCGCRRRGPFSPYEEGRNGQGGGAPPPPPGLA
Chimera 7: MASGRGPRENRPGEVEIWDLNREPWDEWLQDMLRDLNEEARRHFGMNMLIRVWNYCVEEGRRHNTPWNEIGYKYYRIVQKSMFVHFRCGCRRRQPFEPYEERRDGQGGGRANRAPPGLD
Chimera 8: MTSGRDPREPLPGWLEIWDLDREPWDEWLQDMLRDLNEEARRHFGMNMLIRVWNYCVEEGRRHNTPMTERAYRYYRLVQKALFVHFRCGCRRRGPFSPYEEGRNGQGGGAPPPPPGLA
Chimera 9: MTSGRDPREPLPGWLEIWDLDREPWDEWLQDMLRDLNEEARRHFGMNMLIRVWNYCVEEGRRHNTPWNEIGYKYYRIVQKSMFVHFRCGCRRRQPFEPYEERRDGQGGGRANRAPPGLD

Vpx HIV-2 Gh1 *UniProtKB/Swiss-Prot: P18045.1:*
MTDPRERVPPGNSGEETIGEAFEWLDRTIEALNREAVNHLPRELIFQVWQRSWRYWHDDQGMSPSYTKYRYLCLMQKAVFIHFKRGCTCLGGGHGPGGWRSGPPPPPPPGLV

Vpx HIV-2 JK *(GenBank: AAL31353.1):*
MDPRERVPPGNSDEETVGEAFAWLERTIVELNREAVNHLPRELIFQVWQRSWAYWRDDQGMSISYTKYRYLLLIQKAMFVHFAKGCTCLRGGHGPGGWRRGPPPPPPPGLA

Vpx HIV-2A Rod *(UniProtKB/Swiss-Prot: P06939.1):*
MTDPRETVPPGNSGEETIGEAFAWLNRTVEAINREAVNHLPRELIFQVWQRSWRYWHDEQGMSESYTKYRYLCIIQKAVYMHVRKGCTCLGRGHGPGGWRPGPPPPPPPGLV

Vpx HIV-2 RH2-13 5C1 (Effective controller):
MADPREAVPPGNSGEETIGEAFAWLERTVEAINREAVNHLPRELIFQVWQKSWRYWHDDQGMSQSYTKYRYLCLIQKALFIHAKKGCTCLGEGHGPGGWRSGPPPPPPPGLV

Vpx HIV-2 RH2-22-1B4 (Effective controller):
MDPRERVPPGNSGEETIREAFEWLERTVAELNREVVNHLPRELIFQVWQRSWAYWHEEQGMSASYTKYRYLCLMQKAMYIHFAKGCTCLREGHGPGGWRPGPPPPPPPGLA

Vpx HIV-2 RH2-1D8 (Non controller):
MADPREIIPPGNSGEETVGEAFAWLERTVEAINREAVNHLPRELIFQVWQRSWRYWHDEQGMSSSYTKYRYLCLMQKAMFIHFTRGCTCLGGGHGPGGWGPGPPPPPPPGLV

Vpx HIV-2 RH2-7B3 (Non controller):
MVDPRETVPPGNSGEETIGEAFAWENRTVEAINREAVNHLPRELIFQVWQRSWRYWHDEQGMSQSYTKYRYLCIIQKALFIHAKQGCTCRGEGHGPGGWRSGPPPPPPPGLV

Chimera D8B3: MADPREIIPPGNSGEETVGEAFAWLERTVEAINREAVNHLPRELIFQVWQRSWRYWHDEQGMSQSYTKYRYLCIIQKALFIHAKQGCTCRGEGHGPGGWRSGPPPPPPPGLV
Chimera B3D8B3: MVDPRETVPPGNSGEETIGEAFAWENRTVEAINREAVNHLPRELIFQVWQRSWRYWHDEQGMSQSYTKYRYLCLMQKAMFIHFTRGCTCLGGGHGPGGWRSGPPPPPPPGLV
Chimera B4D8B4: MDPRERVPPGNSGEETIREAFEWLERTVAELNREVVNHLPRELIFQVWQRSWAYWHEEQGMSASYTKYRYLCLMQKAMFIHFTRGCTCLGGGHGPGGWRPGPPPPPPPGLA

Larrous *et al*, Sup Figure 1