CemR atypical response regulator impacts energy conversion in *Campylobacteria*

Mateusz Noszka[1], Agnieszka Strzałka[2], Jakub Muraszko[1], Dirk Hofreuter[3], Miriam Abele[4], Christina Ludwig[4], Kerstin Stingl[5], Anna Zawilak-Pawlik[1*]

[1] Department of Microbiology, Hirszfeld Institute of Immunology and Experimental Therapy, Polish Academy of Sciences, 53-114 Wrocław, Poland

[2] Department of Molecular Microbiology, Faculty of Biotechnology, University of Wrocław, 50-383 Wrocław, Poland

[3] Department of Biological Safety, Unit of Product Hygiene and Disinfection Strategies, German Federal Institute for Risk Assessment, 12277 Berlin, Germany

[4] Bavarian Center for Biomolecular Mass Spectrometry (BayBioMS), Technical University of Munich (TUM), 85354 Freising, Germany

[5] Department of Biological Safety, National Reference Laboratory for Campylobacter, German Federal Institute for Risk Assessment, 12277 Berlin, Germany

[*] to whom correspondence should be addressed: Anna Pawlik, tel. +48 71 3709949, e-mail: anna.pawlik@hirszfeld.pl

Abstract

Campylobacter jejuni and *Arcobacter butzleri* are microaerobic food-borne human gastrointestinal pathogens that mainly cause diarrhoeal disease. These related species of the *Campylobacteria* class face variable atmospheric environments during infection and transmission, ranging from nearly anaerobic to aerobic conditions. Thus, they need to adjust their metabolism and respiration to the changing oxygen concentrations of the colonisation sites. Our studies revealed that *C. jejuni* and *A. butzleri* lacking a *Campylobacteria*-specific regulatory protein, *C. jejuni* Cj1608 or a homologue *A. butzleri* Abu0127, are unable to reprogram tricarboxylic acid cycle or respiration pathways, respectively, to produce ATP efficiently and, in consequence, adjust growth to changing oxygen supply. Based on these and previous studies on the homologous protein HP1021 in *Helicobacter pylori*, we propose that these regulatory proteins, which we named *Campylobacteria* energy and metabolism regulators (CemR), play an important role in *Campylobacteria* metabolism and energy regulation, allowing bacteria to adapt to variable oxygen availability.

Campylobacter jejuni and *Arcobacter butzleri* are gram-negative, microaerobic bacteria that belong to the *Campylobacteria* class [1]. *C. jejuni* is a commensal bacterium of the gastrointestinal tracts of wildlife and domestic animals. However, in humans, *C. jejuni* is the leading cause of food-borne bacterial diarrhoeal disease[2]. Also, it causes autoimmune neurological diseases such as Guillian Barré and Miller–Fisher syndromes[3]. *A. butzleri* is found in many ecological niches, such as environmental water and animals. Still, in humans, *A. butzleri* causes diarrhoea, enteritis, and bacteremia[2]. Due to the wide distribution of the two species and relatively high resistance to environmental conditions[4] and antibiotics[5,6], both species pose a severe threat to human health.

To adapt to different ecological niches and decide whether conditions are optimal for reproduction and transmission, *C jejuni* and *A. butzleri* encode numerous regulatory proteins. According to the MiST database[7], *C. jejuni* NCTC 11168 (1.64 Mbp; 1643 coding sequences[8]) encodes [7,9]56 signal transduction proteins, which include one-component systems (OCS, 18 proteins) and two-component systems (TCS, 6 histidine kinases (HK) and 10 response regulators (RR)). *A. butzleri* RM4018 (2.34 Mbp, 2259 coding sequences) encodes 218 signal transduction proteins, including 82 OCS, 32 HK and 43 RR of the TCS[7,10]. While *A. butzleri* regulatory systems have been hardly studied[11], the *C. jejuni* signal transduction systems have been partially deciphered[12–16]. Particular interest was directed to studying processes related to the *C. jejuni* colonisation and focused on factors controlling the expression of genes involved in such processes as host adaptation and niche detection (BumSR, DccSR), oxidative stress (e.g., CosR, PerR), metabolism, and respiration (e.g., BumSR, RacSR, CosR, CsrA, LysR (Cj1000), CprSR)[12,14–16] (see also Discussion). It should be noted that *C. jejuni* requires oxygen for growth but, unlike facultatively aerobic *A. butzleri*, is sensitive to aerobic conditions faced during transmission. In addition, *C. jejuni* and *A. butzleri* may encounter conditions deficient in oxygen in the intestine or during intracellular persistence[17]. Thus, these bacteria must adjust their metabolism and respiration as oxygen concentrations change at colonisation sites. However, no orthologs of energy or redox metabolism regulators such as gamma-proteobacterial ArcA, FNR or NarP are found in *C. jejuni*[15,18–20].

The roles of many *C. jejuni* and most *A. butzleri* regulators have still not been explored. One such regulator is the *C. jejuni* NCTC 11168 Cj1608 orphan response regulator, which is homologous to uncharacterised *A. butzleri* RM4018 Abu0127 (39% amino acid (aa) identity, 61% aa similarity) and a relatively well studied *Helicobacter pylori* 26695 HP1021 (32% aa identity, 56% aa similarity). Moreover, this regulator is conserved across most *Campylobacteria* class species, constituting most of the *Campylobacterota* phylum.

The *H. pylori* HP1021 response regulator is one of 28 signal transduction proteins encoded by *H. pylori* 26695[7,21]. HP1021 acts as a redox switch protein, i.e. senses redox imbalance, transmits the signal and triggers the cells' response[22]. The HP1021 regulon, initially determined in *H. pylori* 26695 by microarray analyses[23], has been updated in *H. pylori* N6 using a multi-omics approach[24]. HP1021 controls the transcription of 497 *H. pylori* N6 genes, including 407 genes related to response to oxidative stress. HP1021 directly controls typical reactive oxygen species (ROS) response pathways and less canonical ones, such as central carbohydrate metabolism. The level of ATP and the growth rate of the knock-out *H. pylori* ΔHP1021 are lower than in the wild-type strain, which is possibly due to reduced transcription of many tricarboxylic acid cycle (TCA) genes and/or increased ATP consumption in catabolic processes in ΔHP1021 compared to the wild-type strain. Thus, HP1021 probably controls *H. pylori* metabolic fluxes to maintain the balance between anabolic and catabolic reactions, possibly for efficient oxidative stress response[24]. HP1021 interacts with the origin of chromosome replication region (*oriC*), probably participating in the control of the initiation of *H. pylori* chromosome replication[25]. As for Cj1608, *in vitro,* it interacts with the *C. jejuni oriC* region and the promoter of the *lctP* lactate transporter, probably participating in the control of the initiation of the *C. jejuni* chromosome replication and lactate metabolism[26,27].

In this work, we focused on the two hardly characterised regulatory proteins homologous to *H. pylori* HP1021 of the microaerobic *C. jejuni* NCTC 11168 and the facultatively aerobic *A. butzleri* RM4018, Cj1608 and Abu0127, respectively. To get insight into the function of the proteins, we constructed mutants lacking these regulators and looked at transcriptomic and proteomic changes under optimal growth and oxidative stress conditions. Our results indicate that Cj1608 and Abu0127, which we named <u>*Campylobacteria*</u> energy and <u>m</u>etabolism <u>r</u>egulators (CemR), support energy conservation in bacterial cells by controlling metabolic and respiration pathways in response to oxygen availability.

Results

Influence of the Cj1608 regulator on *C. jejuni* gene and protein expression

To elucidate the role of the Cj1608 regulator in controlling *C. jejuni* gene expression and oxidative stress response, we performed transcriptome analysis (RNA-seq) of the *C. jejuni* NCTC 11168 wild-type (CJ) and deletion mutant strains (ΔCj1608) under microaerobic and oxidative stress conditions. The results of the disc-diffusion assay indicated that from all tested substrates, which induce the formation of reactive oxygen species, paraquat, the superoxide generator agent, inhibited the growth of the ΔCj1608 strain to the greatest extent compared to the growth of the CJ strain (Fig. S1a). Thus, paraquat was used in further analyses to induce oxidative stress in *C. jejuni* (CJS, ΔCj1608S).

A comparison of *C. jejuni* ΔCj1608 and CJ transcriptomes revealed 380 differentially transcribed genes (Fig. 1a, b, Fig. S2a, Supplementary Data 1). The transcription of 232 genes significantly changed in the stressed CJS cells compared to unstressed CJ, while 123 genes were transcribed differently between stressed ΔCj1609S and unstressed ΔCj1608 cells (Fig. 1b, Fig. S2b, c). The transcription of 44 genes was similarly induced or repressed in CJS and ΔCj1609S cells (Fig. S2d, red dots). Thus, these genes responded to oxidative stress, but other or additional factors than Cj1608 controlled them. Using ClusterProfiler[28], we performed a KEGG enrichment analysis to identify processes affected most by the lack of Cj1608 or under stress. In the ΔCj1608 strain, three KEGG groups were activated (nitrogen metabolism, ribosomes, and aminoacyl-tRNA-biosynthesis), and one group was suppressed, namely the tricarboxylic acid cycle (TCA) cycle (Fig.1c). In the CJ strain under stress conditions, six KEGG groups were suppressed (e.g. oxidative phosphorylation) and four KEGG groups were activated (e.g., carbon metabolism).

Since post-transcriptional regulation is common in bacteria, including species closely related to *C. jejuni*[29,30], a proteomics approach (LC-MS/MS) was applied to detect proteins whose levels changed between analysed strains or conditions. A total of 1155 proteins were detected (Supplementary Data 1). In the CJS cells, the expression of ferrohelatase HemH (Cj0503c) was down-regulated while the levels of three proteins were up-regulated (catalase (Cj1385), the chaperon trafficking heme to catalase (Cj1386), and the periplasmic protein Cj0200c) (Fig. S4b, c). The small number of proteins whose expression was altered under oxidative stress confirmed the known phenomenon of ceasing protein synthesis by bacteria under oxidative stress[24,31]. Thus, we focused on analyses concerning differences between proteomes of ΔCj1608 and CJ strains under microaerobic conditions (Fig. S4a). The transcriptomics and proteomics data correlated strongly (Fig. S5a). The expression level of 70 proteins changed in ΔCj1608 at transcription and protein levels, while the expression of 86 proteins changed only at protein level. Up- or down-regulation of gene transcription and protein level correlated for 58 genes, while it was the opposite for 12 genes (Fig. 1d, red and yellow dots, respectively). However, 173 proteins encoded by genes whose transcription changed in ΔCj1608 were not detected by MS (Supplementary Data 1). The ClusterProfiler KEGG enrichment analysis revealed that genes of two KEGG groups, nitrogen metabolism and ribosome, were activated in the ΔCj1608 strain in comparison to CJ, while the genes of three KEGG groups were suppressed (TCA cycle, ABC transporters and carbon metabolism) (Fig. 1e). It should be noted that the nitrogen metabolism group includes proteins such as GlnA and GltBD, which are involved in glutamate metabolism and TCA cycle[32], whose levels increased more than 10-fold, and Nap and Nrf complexes, constituting an electron transport chain (ETC) system (see below).

To conclude, the transcriptomic and proteomic data revealed that Cj1608 impacts the transcription of 585 genes and the expression of 156 proteins in *C. jejuni* NCTC 11168, with the TCA and nitrogen metabolism KEGG groups most affected.

Influence of Abu0127 on *A. butzleri* gene and protein expression

To elucidate the role of the Abu0127 regulator in controlling gene expression and oxidative stress response of *A. butzleri,* we performed RNA-seq and LC-MS/MS analyses similar to those for *C. jejuni*. We analysed gene expression in *A. butzleri* RM4018 wild-type (AB) and deletion mutant strains (ΔAbu0127); paraquat was used to induce oxidative stress (ABS, ΔAbu0127S, Fig. S1b[33]).

Comparison of *A. butzleri* ΔAbu127 and AB transcriptomes revealed 779 differentially transcribed genes (Fig. 2a, b, Fig. S3a, Supplementary Data 2). The transcription of 290 genes significantly changed under stress conditions in ABS compared to AB cells (Fig. 2b, Fig. S3b). The transcription of only 14 genes changed in the ΔAbu0127S compared to ΔAbu0127 cells (Fig. 2b, Fig. S3c); 12 of these differently transcribed genes were also induced in ABS cells (Fig. S3d, red dots). Thus, these genes responded to oxidative stress but were controlled by factors other than or additional to Abu0127. KEGG enrichment analysis revealed that in the ΔAbu0127 strain, two KEGG groups were activated, ABC transporters and sulfur metabolism, while 13 groups were suppressed (e.g., oxidative phosphorylation and TCA cycle) (Fig. 2c). In the AB strain under stress

conditions, seven KEGG groups were suppressed (e.g., oxidative phosphorylation and citrate cycle), while six groups were activated (e.g., sulfur metabolism).

As in the *C. jejuni* studies, a proteomic approach was used to detect proteins whose levels changed between the strains and conditions analysed. 1608 *A butzleri* proteins were detected (Supplementary Data 2). In ABS cells, the level of one protein of unknown function, Abu0530, changed compared to AB cells (Fig. S4e, f). Thus, *A. butzleri*, like *C. jejuni* and other bacteria, ceased protein translation upon oxidative stress. The transcriptomics and proteomics data correlated strongly (Fig. S5b). The comparison of protein abundance between ΔAbu0127 and AB under microaerobic conditions revealed 124 differentially expressed proteins (Fig. 2d, Fig. S4d). The expression level of 64 proteins changed in ΔAbu0127 at transcription and protein levels. Up- or down-regulation of gene transcription and protein level correlated for 54 genes, while it was the opposite for 10 genes (Fig. 2d, red and yellow dots, respectively). However, proteomics did not detect 224 proteins encoded by genes whose transcription changed in ΔAbu0127 (Supplementary Data 2). The ClusterProfiler KEGG enrichment analysis revealed that genes of one KEGG group, sulfur metabolism, were activated in the ΔAbu0127 strain compared to AB. In contrast, the genes of three KEGG groups were suppressed (oxidative phosphorylation, taurine and hypotaurine metabolism, ribosome) (Fig. 2e).

To conclude, the transcriptomics and proteomics data revealed that Abu0127 impacts the transcription of 904 genes and the expression of 124 proteins in *A. butzleri* RM4018, with the oxidative phosphorylation KEGG group most affected.

Cj1608 and Abu0127 are involved in the regulation of energy production and conversion

The transcriptome and proteome analyses of both species revealed that the energy production and conversion processes, such as the TCA cycle and oxidative phosphorylation, are putatively controlled in *C. jejuni* NCTC 11168 by Cj1608 and in *A. butlzeri* RM4018 by Abu0127 (Supplementary Data 1, Supplementary Data 2 and Supplementary Data 3). In *C. jejuni* ΔCj1608, the protein levels of all but two TCA enzymes, fumarate reductase FrdABC and fumarate hydratase FumC, decreased, with the citrate synthase GltA protein level reduced by as much as 70-fold (Fig. 3a, Fig. S6; the level of oxoglutarate-acceptor oxidoreductase Oor was reduced significantly by 1.7-2 fold). Many genes encoding nutrient importers and downstream processing enzymes (e.g., amino acid ABC transporter permease Peb, fumarate importer DcuAB, L-lactate permease LctP and lactate utilisation proteins Lut) were down-regulated in *C. jejuni* ΔCj1608, except for up-regulated DctA succinate/aspartate importer and the enzymes glutamine synthetase GlnA and glutamate synthase small subunit GltB involved in the generation of glutamine and glutamate (Fig. 3a). The levels of many proteins forming ETC complexes were also down-regulated, including subunits of NADH-quinone oxidoreductase Nuo and cytochromes (Fig. 3b). In *A. butzleri* ΔAbu0127, the levels of GltA and isocitrate dehydrogenase Icd proteins were reduced (Fig. 3c, Fig. S7c). As in *C. jejuni*, the levels of many proteins forming ETC complexes decreased, with Nuo complex subunits being the most highly downregulated (Fig. 3d, Fig. S7a, b).

Moreover, we performed a functional analysis annotation with eggNOG-mapper, indicating the Clusters of Orthologous Groups (COG) of differentially expressed genes and proteins. In both species, the most variable groups belonged to two categories, namely energy production and conservation (C) and translation, ribosomal structure, and biogenesis (J) (Fig. 4, the category of unknown genes was excluded from the graph, Supplementary Data 1 and Supplementary Data 2). Many genes and proteins suppressed in both species belong to the energy production and conversion group.

These comprehensive data suggest that mutants of both species produce less energy due to the inhibition of the TCA cycle and oxidative phosphorylation. Therefore, we decided to investigate these processes in more detail and analyse how the lack of Cj1608 and Abu0127 proteins influences energy production and conversion in *C. jejuni* and *A. butzleri*, respectively.

Cj1608 controls *gltA* expression, ATP level and growth in response to O$_2$ supply

To analyse the influence of Cj1608 on *C. jejuni* growth, we compared the growth of wild-type CJ, knock-out ΔCj1608 and complementation COM strains (Fig. 5a). It should be noted that unless an accessory electron donor H$_2$ was supplied in the microaerobic gas mixture (4 % H$_2$, Materials and Methods), we could not obtain the ΔCj1608 mutant strain of *C. jejuni* NCTC 11168. When *C. jejuni* was cultured microaerobically in the presence of H$_2$, the growth of *C. jejuni* ΔCj1608 was slower than the CJ and COM strains (Fig. 5a). The generation time (G) of ΔCj1608 was higher than the generation time of the CJ and complementation (COM) strains under the same conditions (ΔCj1608: 239 ± 17 min, CJ: 204 ± 6 min; COM 201 ± 6 min) (Fig. 5a, b). The ΔCj1608 culture entered the stationary phase at optical density (OD) similar to that of CJ and COM strains (Fig. 5a). Under microaerobic conditions without H$_2$, the CJ and COM strains grew similarly as in the presence

of H_2, albeit reached lower OD_{600} at the stationary phase than in the presence of H_2 (WT G of 210 ± 3 min, COM G of 195 ± 2 min, Fig. 5b). In contrast, ΔCj1608 strain almost did not grow in the absence of H_2, confirming that its growth strictly depended on hydrogen. The ATP analysis indicated that the relative energy levels corresponded to the bacterial growth rates. The ATP level of the CJ strain under microaerobic growth without H_2 was assumed to be 100% (Fig. 5c). The ATP level in CJ and COM strains was constant regardless of the presence or absence of H_2 (85% - 100%). Without H_2, the ATP level in the ΔCj1608 strain reached $25 \pm 4\%$ of the ATP level in the CJ strain. In the presence of H_2, the level of ATP in ΔCj1608 cells increased to 53 ± 8 % compared to that of the CJ strain. These results indicated that the reduced ATP synthesis in ΔCj1608 might be bypassed by using electrons and proton motive force (PMF) generated from hydrogen. Thus, using H_2, the ΔCj1608 cells could produce more energy and multiply more efficiently.

Next, we analysed whether Cj1608 directly affects the TCA cycle efficiency, as suggested by hydrogen boost analysis. We studied the expression of *gltA* since the expression of this gene was severely downregulated in the ΔCj1608 strain at the transcription and translation level (Fig. 5d, 5e, Supplementary Data 3). GltA is the first enzyme in the TCA cycle whose activity impacts the flow of substrates through the TCA cycle and energy production via NADH/FADH/menaquinone cofactors (Fig. 3a). Moreover, in RNA-seq analysis, *gltA* gene transcription was increased under oxidative stress in the wild-type CJ, while it did not change under oxidative stress in the ΔCj1608 strain (Fig. 5d, g, Supplementary Data 3). It was previously shown that *gltA* transcription depends on the O_2 concentration, being lower at 1.88% O_2 and higher at 7.5% O_2, which helps *C. jejuni* optimise energy production and expense during different oxygen availability as the electron acceptor[34]. In this work, we confirmed the interaction of the Cj1608 protein with the *gltA* promoter region by electrophoretic mobility shift assay (EMSA), and we figured that it was specific because Cj1608 did not interact with a control *C. jejuni recJ* region (Fig. 5f). Next, we used RT-qPCR to analyse *gltA* transcription under oxidative stress triggered by paraquat or under diverse O_2 availability: reduced 1% O_2, optimal microaerobic 5% O_2 and increased 10% O_2 (Materials and Methods). Under optimal conditions, *gltA* transcription was lower in ΔCj1608 than in CJ and COM strains (FC of 0.14 ± 0.14), and, unlike in the CJ and COM strain, the transcription of *gltA* in ΔCj1608 was not up-regulated upon paraquat-induced oxidative stress (Fig. 5g). *gltA* was up-regulated in the CJ and COM strains at 5% O_2 compared to 1% O_2 (Fig. 5g). However, in CJ and COM strains, the *gltA* transcription was down-regulated at 10% O_2 compared to 5% O_2 (Fig. 5h), suggesting that the prolonged approx. 5-hour bacterial growth at 10% O_2 possibly caused adaptation to higher oxygen level conditions or stress by reducing TCA cycle activity and possible ROS production during increased aerobiosis (see Materials and Methods). However, *gltA* transcription was constant in ΔCj1608 at all O_2 concentrations, indicating that cells could not adjust *gltA* transcription to changing O_2 conditions. Next, we analysed *C. jejuni* growth and ATP production under various O_2 supplies. Under 1% O_2, the ΔCj1608 strain grew faster than WT and COM strains, suggesting that the metabolic state of the ΔCj1608 mutant strain was optimised for reduced O_2 concentration (Fig. 5i). Under optimal O_2 level, CJ and COM strains grew faster than under 1% O_2 and faster than the ΔCj1608 strain under 5% O_2, indicating that metabolic pathways of CJ and COM strains were adjusted to increased O_2 availability, while the metabolism of ΔCj1608 was not. Under increased O_2 concentration, all three strains grew slower than under optimal conditions, with the ΔCj1608 strain growing slower than the CJ and COM strains. The ATP analyses indicated that the relative ATP levels corresponded to the bacterial growth rates. The ATP level of the CJ strain under optimal microaerobic growth was assumed to be 100%. Compared to that, the level of ATP in the CJ strain under 1% O_2 dropped to $38 \pm 4\%$; at 10% O_2, the level of ATP also decreased to $51 \pm 7\%$ (Fig. 5j, see Discussion). Under optimal conditions in the ΔCj1608 mutant strain, the ATP level reached approximately $70 \pm 8\%$ of that in the CJ strain and was constant regardless of increased or decreased O_2 supply. The ATP levels in the COM strain analysed at different O_2 levels resembled that of the CJ strain. Thus, the results indicated that the ΔCj1608 mutant strain could not adjust pathways responsible for energy conservation to changing O_2 levels.

To summarise *C. jejuni*'s results, Cj1608 helps *C. jejuni* control pathways important for energy conservation dependent on oxygen availability. As we have shown using the *gltA* gene as an example, Cj1608 directly controls gene expression in response to changing O_2 concentrations and oxidative stress.

Abu0127 controls *nuo* expression, ATP level and growth in response to O_2 supply.

Next, we studied the impact of Abu0127 on *A. butzleri's* growth. As for *C. jejuni*, the presence or absence of H_2 did not significantly affect the growth of the wild-type AB strain nor ATP level in AB cells under microaerobic conditions (Fig. 6a-c). In the presence of H_2, the ΔAbu0127 strain grew slower than the AB strain (ΔAbu0127 G of 184 ± 20 min, AB G of 108 ± 16 min) (Fig. 6a, b), and the ATP level of ΔAbu0127 cells reached $77 \pm 3\%$ of that in the WT cells (Fig. 6c). However, the ΔAbu0127 culture entered the stationary phase at OD similar to that of the AB strain. Under microaerobic conditions without H_2, the growth and ATP level of

ΔAbu0127 was further lowered (ΔAbu0127 G of 256 ± 35 min, AB G of 112 ± 6 min; ΔAbu0127 ATP level reached 57 ± 3% of the AB cells), and cells entered stationary phase at lower OD_{600} than the wild-type AB strain (Fig. 6a-c). Despite many attempts, we could not construct an Abu0127 complementation mutant, possibly due to the imperfect molecular biology tools dedicated to *A. butzleri*[35]. Nonetheless, the results indicated that the ΔAbu0127 strain produced less energy than AB cells. However, when additional electron donors and proton motive force derived from hydrogen were available, alternative to that produced by the TCA cycle and used in oxidative phosphorylation, the ΔAbu0127 cells could produce more energy and multiply more efficiently.

Next, we analysed whether Abu0127 directly affects the Nuo complex activity. We studied the expression of *nuoB* since it is the second gene of the *nuoA-N* operon, whose expression was severely downregulated in the ΔAbu0127 strain at the transcription and translation level (Fig. 6d and 6e, Supplementary Data 3). We confirmed the interaction of the Abu0127 protein with the *nuoA-N* promoter region by EMSA, and we figured that it was specific because Abu0127 did not interact with a control *A. butzleri recJ* region (Fig. 6f). Next, we used RT-qPCR to analyse *nuoB* transcription under paraquat-induced oxidative stress and different O_2 supply, as previously in *C. jejuni*. It should be noted, that *A. butzleri* can grow under aerobic conditions[36]; thus, aerobic O_2 concentration is less harmful to *A. butzleri* than to *C. jejuni*. *nuoB* transcription in the wild-type AB strain did not change under paraquat-induced oxidative stress, confirming the trancriptomic results (Fig. 6d, g, and Supplementary Data 3). *nuoB* transcription changed across different O_2 concentrations, being the lowest at 1% O_2 (FC of 0.61 ± 0.06 compared to AB under 5% O_2) and highest at 10% O_2 (FC of 1.7 ± 0.27 compared to AB under 5% O_2) (Fig. 6h). Under optimal O_2 conditions, *nuoB* transcription was lower in the ΔAbu0127 strain than in the AB strain (FC of 0.3 ± 0.05 compared to AB under 5% O_2), and it was stable across different O_2 concentrations (Fig. 6h). Next, we analysed *A. butzleri* growth and ATP production under various O_2 supplies. Under 1% O_2, the AB and ΔAbu0127 strains grew similarly. However, Δabu0127 reached a lower OD_{600} at the stationary growth phase than the AB strain (Fig. 6i). Under optimal and increased O_2 concentrations, the AB strain grew similarly at both concentrations, faster than under 1% O_2, but reached similar OD upon entry to a stationary growth phase as under 1% O_2. The ΔAbu0127 strain grew similarly under 5% and 10% O_2. The growth was faster than under 1% O_2, but slower than AB under the same conditions, and the culture finally reached lower OD than AB (Fig. 6i). The ATP analyses indicated that the relative energy levels corresponded to the bacterial growth rates. The ATP level of the AB strain under microaerobic growth was assumed to be 100%. Compared to that, under 1% O_2, the level of ATP in the AB strain dropped to 44 ± 4%, while the level of ATP did not change in the culture grown under 10% O_2 (Fig. 6j). Under reduced O_2 concentration, the level of ATP in ΔAbu0127 was similar to that of the AB strain under the same conditions (35 ± 4% compared to AB under microaerobic growth). However, under 5% O_2, the ATP level of the ΔAbu0127 mutant strain reached 76 ± 3% of that in the AB strain and did not increase under 10% O_2. Thus, the results indicated that the ΔAbu0127 mutant strain could not efficiently adjust pathways responsible for energy conservation to changing O_2 levels, in which it resembled *C. jejuni* ΔCj1608.

To summarise *A. butzleri*'s results, Abu0127 helps *A. butzleri* control pathways important for energy conservation dependent on oxygen availability. As we have shown using the *nuoB* gene as an example, Abu0127 directly controls gene expression in response to changing O_2 concentrations.

Discussion

One of the challenges for a bacterial cell is to gain energy to grow and reproduce under different environmental conditions. Available energy sources and electron acceptors used for energy conversion are the two major factors defining the activity of metabolic pathways. The regulatory proteins redirect metabolism and ETC pathways, prioritising electron donor or acceptor usage to maximise the energy gain (e.g., aerobic/microaerophilic over anaerobic respiration and fermentation or nitrate respiration over fumarate respiration). The fine-tuning of metabolism and respiration to particular conditions is usually hierarchical and orchestrated by multiple factors[37,38], which often control the same genes. For example, in the model, facultative anaerobe *E. coli*, but also other species of Gammaproteobacteria, three global transcriptional regulators, FNR, ArcA and NarL/NarP mainly control energy conservation processes dependent on electron acceptors availability[20,39]. In the energy reprogramming network, under low oxygen availability, FNR provides the first level of regulation, switching metabolism from aerobic to anaerobic and controlling the expression of other transcriptional factors[40]. ArcA, a response regulator of the ArcAB two-component system[41], possibly controls the switch between anaerobic respiration and fermentation as well as the expression of a set of secondary regulators[42]. NarL/NarP switches on nitrate and nitrite respiration when these electron acceptors are available while repressing genes for less effective anaerobic respiration[43]. Studies have shown that the intensive cross-talk between regulatory proteins, often controlling the same genes hierarchically and controlling genes beyond energy conservation pathways, is essential for efficient bacterial responses or adaptation to diverse conditions[20,44–46].

Till recently, no regulators redirecting energy conversion pathways in response to oxygen as an electron acceptor have been found in *Campylobacterota*[15,47]. It was experimentally shown that *C. jejuni* reprograms its metabolism upon changes to oxygen availability; nonetheless, the regulator remained unknown[34]. It was surprising given that microaerobic species of this phylum must respond or adapt to changing oxygen levels, including harmful oxygen deficiency and excess.

In this and previous works[22,24], we revealed that under microaerobic conditions, three homologous atypical response regulators, HP1021, Cj1608 and Abu0127, which we name <u>*Campylobacteria* energy and metabolism regulators</u> (CemR), control the expression of many genes involved in energy conversion, including TCA and ETC pathways (Fig. 3, Fig. 4, Fig. S8 and Supplementary Data 3 in [24]). Indeed, when *cemR* genes were deleted, the most remarkable gene and protein expression changes occurred in energy production and conversion COG, reaching between 20 to 40% of all the down-regulated genes or proteins (Fig. 4 and Fig. S8, the category of unknown proteins was excluded from the study). Moreover, we showed that in strains lacking CemR, the transcriptional control was lost over the genes whose transcription was dependent on the oxygen level, e.g., *C. jejuni* TCA *gltA*, *A. butzleri* ETC *nuoA-N* operon complex, *H. pylori,* TCA pyruvate:ferredoxin oxidoreductase *PFOR* and ETC cytochrome c oxidase *ccoN-Q* (Fig. 5h, Fig. 6h and [24]). Consequently, under microaerobic conditions in CemR deletion mutants of all three species, the ATP level and the growth rates were lower than in wild-type strains (Fig. 5i, j, Fig. 6i, j and Fig. 3e in [24]). The presence of CemR allowed *C. jejuni* and *A. butlzeri* to optimise energy conservation under microaerobic compared to reduced oxygen concentration, increasing the ATP level and bacterial growth rate (Fig. 5i, j, Fig. 6i, j). The lack of CemR specifically impaired pathways connected with oxygen utilisation as an electron acceptor because in the presence of hydrogen as an alternative electron donor and proton motive force generator, *C. jejuni* and *A. butzleri* gained additional energy, allowing for faster growth (Fig. 5a-c, Fig. 6a-c). All these data indicate that CemR responds to oxygen levels and redirects metabolism towards optimal energy conservation.

An interesting question arises: how many energy conservation regulators are encoded in addition to CemR in each *Campylobacteria* species, and how do they all integrate into the regulatory circuits of a given species? *H. pylori*, *C. jejuni* and *A. butzleri* differ in lifestyles, which shape the complexity of regulatory circuits, as recently illustrated using *Campylobacterota* as a representative phylum[48]. *H. pylori* can be classified as a specialist bacterium[49] - strictly human-associated, using a limited repertoire of electron donors (hydrogen, pyruvate, 2-oxo-glutarate, malate, and succinate) and only oxygen or fumarate as electron acceptors during respiration[19]. Oxygen, being a primary electron acceptor in *H. pylori*, is also toxic for *H. pylori* at higher than microaerobic concentrations[50]. As a consequence of speciation and host adaptation, *H. pylori* lost many regulatory proteins. Nonetheless, the studies on regulatory circuits in *H. pylori* indicated that HP1043[51,52], controlled by unknown stimuli, acid-responsive ArsRS[53], and metal-dependent regulators NikR and Fur[19,54], affect *H. pylori* energy conversion to various extents in response to different stimuli. *C. jejuni* is a host-

associated organism that infects various animals and humans and can survive long periods under aerobic conditions[55]. In contrast to *H. pylori*, *C. jejuni* has a branched electron transport chain with multiple enzymes that utilise more molecules as electron donors and acceptors than *H. pylori*[19]. *C. jejuni* genome size is similar to *H. pylori*; however, the number of *C. jejuni* genes encoding regulatory proteins is higher and reflects the broader spectrum of ecological niches inhabited by this bacterium. Consequently, the number of regulatory proteins controlling energy conversion is higher than in *H. pylori,* with RacRS, LysR (Cj1000), CsrA and CprRS playing the most significant roles (for details, see reviews[12,14–16]). Depending on growth conditions, particularly the availability of electron donors and acceptors, LysR, RacRS, and CprRS regulate the expression of fumarate respiration genes, which is an excellent example of a cross-talk between regulators controlling the same pathways via transcriptional control over the same genes (e.g., *aspA, dcuA, mfr* or *frd*)[47,56,57]. RNA-binding CsrA, a homologue of *E. coli* carbon starvation regulator, is an example of post-transcriptional energy conservation regulation in *C. jejuni,* including control of expression of TCA and ETC genes[58]. *A. butzleri* has the ability to lead anaerobic, microaerophilic and aerobic life styles as an environmental or animal- and human-associated bacterium. As a generalist, it utilises many electron donors and acceptors using an electron transport chain similar to those used by the free-living marine *Campylobacterota*[19]. *A. butzleri*'s genome encodes many signal transduction proteins (see Introduction), including extracytoplasmic sigma factors (ECFσ) to adapt to different conditions. However, hardly anything is known about the regulation of *A. butzleri* energy conversion except that a few species of ECFσimpact electron and carbon metabolism by affecting the transcription of genes from carbon metabolism pathways and electron acceptor complexes[11]. However, one can expect the highest complexity and speciation of *A. butzleri*'s regulatory circuits, including energy conservation, of all three species.

CemR regulators are highly conserved *Campylobacteria* regulators. However, the molecular mechanism of signal perception by CemR is still enigmatic and cannot be predicted by analogy. Transcriptional regulatory proteins may sense respiration status, the concentrations of electron donors or acceptors, or respiration byproducts[59]. We have shown that cysteine residues of HpCemR become oxidised under O_2-triggered oxidative stress in vivo and in vitro, which changes the protein's DNA-binding activity possibly by structural changes of DNA-protein complex rather than on-off mechanism[22,24]. Recent studies on CjCemR interactions with the promoter region of *lctP* suggested a similar mechanism of CjCemR activity control and *lctP* transcription regulation[26]. AbCemR protein also contains cysteine residues, one of which is conserved in all three species (C27 in HpCemR and CjCemR, C31 in AbCemR). The mechanism of activity regulation by the redox state of the cysteine residues resembles that of ArcA-ArcB two-component system regulation. However, in that TCS, the cysteine residues of ArcB sensor kinase are oxidated, triggering autophosphorylation of ArcB, while ArcA response regulator is activated by phosphorylation. CemR proteins are orphan atypical response regulators, in which the same molecule receives and executes the signal. Thus, despite some biochemical similarity to ArcAB system activation, the exact molecular mechanism of CemR activity control waits to be discovered.

In summary, CemR, the global regulatory protein, is a part of cell response to a metabolic redox imbalance, either caused by environmental conditions such as oxygen availability or reactive oxygen species (ROS) or triggered intracellularly due to metabolic changes and increased ROS production. However, the exact pathway of signal transduction by CemR and the levels or hierarchy of regulation are still unknown. Further comprehensive, multi-omic studies are needed to reveal the complex circuits of regulation involving CemR and other regulatory proteins in *Campylobacteria* energy conservation.

Methods

Materials and culture conditions

The strains, plasmids and proteins used in this work are listed in Supplementary Table S1. The primers used in this study are listed in Supplementary Table S2. *C. jejuni* and *A. butzleri* plate cultures were grown on Columbia blood agar base medium supplemented with 5% defibrinated sheep blood (CBA-B). The liquid cultures were prepared in Brain Heart Infusion Broth (BHI) (Oxoid) and incubated with 140 rpm orbital shaking. All *C. jejuni* and *A. butzleri* cultures were supplemented with antibiotic mixes (*C. jejuni*: vancomycin (5 µg/ml), polymyxin B (2.5 U/ml), trimethoprim (5 µµg/ml) and amphotericin B (4 µg/ml)[60]; *A. butzleri*: cefoperazone (8 µg/ml), amphotericin (10 µg/ml), teicoplanin (4 µg/ml)[61]). If necessary for selecting mutants, appropriate antibiotics were used with final concentrations (i) kanamycin 20 µg/ml (*C. jejuni* and *A. butzleri*), (ii) chloramphenicol 8 µg/ml (*C. jejuni*). Routinely, *C. jejuni* and *A. butzleri* were cultivated at 42°C or 30°C, respectively under optimal microaerobic conditions (*C. jejuni*: 5% O_2, 8% CO_2, 4% H_2 and 83 % N_2; *A. butzleri*: 6% O_2, 9% CO_2, and 85% N_2) generated by the jar evacuation-replacement method using Anaerobic Gas System PetriSphere. The gas mixtures were modified by lowering or increasing O_2 and H_2 concentrations for ATP assays and growth curve analyses. Briefly, bacteria were cultured under microaerobic conditions optimal for each species to the late logarithmic growth phase (OD_{600} ~0.8), diluted to OD_{600} ~0.05 and incubated under the desired atmosphere as long as required, up to OD_{600} = 0.2-0.6 for ATP and RT-qPCR or until late stationary phase in growth analyses. The gas mixture with or without H_2 was composed of optimal gas mixtures for *C. jejuni* and *A. butzleri*, respectively. Low oxygen gas mixture was composed of 1% O_2, 10% CO_2, 5% H_2 and 84% N_2, while high oxygen gas mixture was composed of 10% O_2, 5% CO_2, 3% H_2 and 82% N_2. *Escherichia coli* DH5α and BL21 were used for cloning and recombinant protein synthesis. If necessary for selecting *E. coli,* appropriate antibiotics were used with final concentrations (i) kanamycin 50 µg/ml and (ii) ampicillin 100 µg/ml.

Construction of *C. jejuni* mutant strains

C. jejuni mutant strains were constructed using a homologous recombination and natural transformation approach[62]. *C. jejuni* NCTC 11168 was grown in 12 ml BHI to an OD_{600} = 0.2. Next, 150 µl of the culture was centrifuged and resuspended in fresh 150 µl of BHI. 150 ng (1 µg/ml) of EcoRI methylated plasmid was added to the culture and cultivated at 42 °C in microaerobic (3.75% H_2, 5.25% O_2, 7.5% CO_2) conditions with shaking (140 rpm) for 4 h. Next, 100 µl was spread on CBA-B plates with an appropriate antibiotic and incubated for 3 days at 42 °C under microaerobic (3.75% H_2, 5,25% O_2, 7,5% CO_2) conditions.

C. jejuni NCTC 11168 ΔCj1608

The *C. jejuni* *Cj1608* deletion construct (pCR2.1/ΔCj1608) was prepared as follows (Fig. S9a). The upstream and downstream regions of *Cj1608* were amplified by PCR using the P1-P2 and P5-P6 primer pairs, respectively and a *C. jejuni* NCTC 11168 genomic DNA as a template. The *aphA-3* cassette was amplified using the P3-P4 primer pair; pTZ57R/TΔHP1021[25] was used as a template. The resulting fragments were purified on an agarose gel. Subsequently, the fragments were combined into one DNA amplicon with an overlap extension PCR reaction using the P1-P6 primers. The generated amplicon was purified on an agarose gel and cloned to the pCR2.1-TOPO® plasmid (Thermo Fisher Scientific) according to the manufacturer protocol. The DNA fragment cloned in pCR2.1-TOPO® was sequenced. Subsequently, *C. jejuni* NCTC 11168 was transformed with the pCR2.1/ΔCj1608 plasmid, and the transformants were selected by plating on CBA-B plates supplemented with kanamycin. The addition of hydrogen was necessary to obtain the kanamycin-resistant colonies. The allelic exchange was verified by PCR using the P7-P8 primer pair. The lack of *Cj1608* in *C. jejuni* NCTC 11168 ΔCj1608 was confirmed by RNA-seq (Fig. S9b), LC-MS/MS (Supplementary Data 1) and Western blot (Fig. S9c, d).

C. jejuni NCTC 11168 COM

The *C. jejuni* Cj1608 complementation construct (pUC18/COMCj1608) was prepared as follows (Fig. SXa). The regions upstream and downstream of the *Cj1608* gene were amplified by PCR using the P1-P6 primer pair using *C. jejuni* Cj1608 genomic DNA as a template. The resulting fragment was purified on an agarose gel. Subsequently, the generated amplicon and the SmaI digested vector pUC18 were ligated according to the method described by Gibson et al.[63] to give pUC18/COMCj1608. *E. coli* DH5α competent cells were transformed with the construct by heat shock. The insert cloned into pUC18 was sequenced. The pUC18/COMCj1608 was used to complement the lack of *Cj1608* in *C. jejuni* NCTC 11168 ΔCj1608 by an

attempt similar to Multiplex Genome Editing by Natural Transformation (MuGENT)[64]. For the selection, a pSB3021 suicide plasmid was used. The plasmid harbours a *cat* cassette flanked by arms of homology, enabling *cat* cassette integration into the *hsdM* gene. Subsequently, the obtained pUC18/COMCj1608 (1 µg/ml) and pSB3021 (0.25 µg/ml) plasmids were used in the natural transformation of *C. jejuni* NCTC 11168 ΔCj1608. The transformants were selected by plating on CBA-B plates supplemented with chloramphenicol. The bacteria were cultured without hydrogen to increase the selection process. The allelic exchange was verified by PCR using the P7-P9 primers. The presence of *Cj1608* in *C. jejuni* NCTC 11168 COM was confirmed by RNA-seq (Fig. S9b), LC-MS/MS (Supplementary Data 1) and Western blot (Fig. S9c, d).

Construction of *A. butzleri* mutant strain

A. butzleri RM4018 mutant strain was constructed using a homologous recombination and electroporation approach. *A. butzleri* was grown in 25 ml BHI to an $OD_{600} = 0.4$. Next, the culture was transferred on ice and left for 10 min, then centrifuged (4,700 g, 10 min, 4 °C) and washed twice with 20 ml of ice-cold glycerine water (15% glycerol, 7% sucrose, filtrated). Finally, the culture was suspended in 250 µl of ice-cold glycerine water. For electroporation, 50 µl of electrocompetent cells were mixed with 5 µg of appropriate plasmid. Electroporation was performed in 0.1 cm (The Cell Projects) electroporation cuvettes using a Gene Pulser II Electroporator (Bio-Rad) using the following parameters – 12.5 kV/cm, 200 Ω, 25 µF. Cells were regenerated by adding 1 ml of BHI medium to the cuvette, then by transferring the cells to a flask with 2 ml BHI and cultivated with shaking (140 rpm) for 4 hours at 30 °C in microaerobic conditions. Next, the culture was centrifuged (4,700 g, 10 min, 22 °C), resuspended in 100 µl of BHI, spread on CBA-B plates with appropriate antibiotic and incubated for 5 days at 30 °C under microaerobic conditions.

A. butzleri RM4018 ΔAbu0127

The *A. butzleri Abu0127* deletion construct (pUC18/ΔAbu0127) was prepared as follows (Fig. S10a)[65]. The upstream and downstream regions of *Abu0127* were amplified by PCR using the P10-P11 and P14-P15 primer pairs, respectively, and an *A. butzleri* RM4018 genomic DNA as a template. The *aphA-3* cassette was amplified using the P12-P13 primer pair; pTZ57R/TΔHP1021[25] was used as a template. The resulting fragments were purified on an agarose gel. Subsequently, the PCR-amplified fragments and the SmaI digested vector pUC18 were ligated according to the method Gibson et al.[63]. Subsequently, *A. butzleri* RM4018 was transformed via electroporation with the pUC18/ΔAbu0127 plasmid and the transformants were selected by plating on CBA-B plates supplemented with kanamycin. The allelic exchange was verified by PCR using the P16-P17 primer pair. The lack of *Abu0127* in *A. butzleri* RM4018 ΔAbu0127 was confirmed by RNA-seq (Fig. S10b) and LC-MS/MS (Supplementary Data 2).

Disc diffusion assay

Bacteria were cultured in BHI to $OD_{600} = $ ~0.5-0.7 and then diluted to $OD_{600} = 0.1$ in BHI. Each culture was spread by a cotton swab on CBA-B plates. Sterile, glass fibre 6-mm discs were placed on plates, and 5 µl of tested solutions were dropped on discs: 2% H_2O_2 (POCH, 885193111), 2% paraquat dichloride (Acros Organics, 227320010), 5% hydroxyurea (Merck, H8627-1G), 3% sodium nitroprusside (Merck, S0501), 2.5% menadion (Merck, A13593) and 15% sodium hypochlorite (Merck, 1056142500). Sodium hypochlorite solutions for assays were made fresh; the solutions' pH was adjusted by adding HCl to pH 7 and kept in PBS buffer prior to experiments[66]. Menadion solution was prepared in DMSO. The diameter of the inhibition zone around the discs was determined after 3 days of incubation under microaerobic conditions. The experiment was performed using three biological replicates.

RNA isolation

Bacterial cultures (12 ml BHI) of *C. jejuni* and *A. butzleri* strains were grown under microaerobic conditions to OD_{600} of 0.5-0.6. Immediately after opening the jar, 2 ml of the non-stressed culture was added to 2 ml of the RNAprotect Bacteria Reagent (Qiagen), vortexed and incubated for 5 min at room temperature. In parallel, the cultures were treated with 1 mM paraquat (final concentration) for 25 min (*C. jejuni* 42 °C, *A. butzleri* 30 °C, 140 rpm orbital shaking). After oxidative stress, samples were collected similarly to non-stressed cells. After 5 min incubation with RNAprotect Bacteria Reagent, bacteria were collected by centrifugation (4,700 × g, 10 min, RT). RNA was isolated by GeneJET RNA Purification Kit (Thermo Fisher Scientific; K0731) according to the manufacturer's protocol and treated with RNase-free DNase I (Thermo Fisher Scientific). Next, purification by the GeneJET RNA Purification Kit was performed to remove DNase I. A NanoDrop Lite spectrophotometer, agarose gel electrophoresis and Agilent 4200 TapeStation System were used to determine

the RNA quality and quantity. RNA was isolated immediately after bacteria collection, stored at -80 °C for up to one month and used for RNA sequencing. RNA was isolated from three independent cultures.

For analyses of gene transcription dependent on oxygen supply, bacterial cells were collected from cultures under the logarithmic phase of growth (OD$_{600}$ ~0.2-0.5), grown under the desired atmosphere (see Material and culture conditions), and RNA was isolated as described above.

RNA sequencing (RNA-seq)

The prokaryotic directional mRNA library preparation and sequencing were performed at the Novogene Bioinformatics Technology Co. Ltd. (Cambridge, UK). Briefly, the ribosomal RNA was removed from the total RNA, followed by ethanol precipitation. After fragmentation, the first strand of cDNA was synthesised using random hexamer primers. During the second strand cDNA synthesis, dUTPs were replaced with dTTPs in the reaction buffer. The directional library was ready after end repair, A-tailing, adapter ligation, size selection, USER enzyme digestion, amplification, and purification. The library was checked with Qubit, RT-qPCR for quantification and a bioanalyser for size distribution detection. The libraries were sequenced with the NovaSeq 6000 (Illumina), and 150 bp reads were produced.

RNA-seq analysis

The 150 bp paired reads were mapped to the *C. jejuni* NCTC 11168 (NC_002163.1) or *A. butzleri* RM4018 (NC_009850.1) genome depending on the species analysed using Bowtie2 software with local setting (version 2.3.5.1)[67,68] and processed using samtools (version 1.10)[69], achieving more than 10^6 mapped reads on average. Differential analysis was performed using R packages Rsubread (version 2.10), and edgeR (version 3.38)[70,71], following a protocol described in[72]. Genes rarely transcribed were removed from the analysis (less than 10 mapped reads per library). The obtained count data was normalised using the edgeR package, and a quasi-likelihood negative binomial was fitted. Differential expression was tested using the glmTtreat function with a 1.45-fold change threshold. Only genes with a false discovery rate (FDR) less than 0.05 and $|\log_2 FC| \geq 1$ were considered differentially expressed. Data visualisation with volcano plots was done using the EnhancedVolcano and tidyHeatmap R packages (version 1.14 and 1.8.1)[73]. The reproducibility of *C. jejuni* and *A. butzleri* biological replicates was visualized by principal component analysis (PCA) of the normalised RNA-seq CPM data (Fig S11a, b).

Proteomic sample preparation

Bacterial cultures of wild-type *C. jejuni* NCTC 11168 and *A. butzleri* RM4018 strains and their isogenic mutant strains, ΔCj1608 or ΔAbu0127, respectively, were grown in BHI under microaerobic conditions to OD$_{600}$ of 0.5-0.7 and split into 2 sub-cultures of 10 ml each. The first sub-culture was harvested, washed, and lysed immediately after opening the jar; the second sub-culture (only WT strains) was harvested, washed and lysed after incubation with 1 mM paraquate (*C. jejuni*: 42 °C; *A. butzleri*: 30 °C with orbital shaking, 140 rpm) for 30 and 60 minutes. The bacterial proteomes were prepared as described previously by Abele et al.[74]. Briefly, cells were harvested by centrifugation at 10,000 × g for 2 min, media were removed, and cells were washed once with 20 ml of 1 × PBS. The cell pellets were suspended and lysed in 100 μl of 100% trifluoroacetic acid (TFA; Roth)[75] for 5 min at 55 °C. Next, 900 μl of Neutralization Buffer (2 M Tris) was added and vortexed. Protein concentration was measured by Bradford assay (Bio-Rad). 50 μg of protein per sample was reduced (9 mM TCEP) and carbamidomethylated (40 mM CAA) for 5 minutes at 95 °C. The proteins were digested by adding trypsin (proteomics grade, Roche) at a 1/50 enzyme/protein ratio (w/w) and incubation at 37 °C overnight. Digests were acidified by the addition of 3% (v/v) formic acid (FA) and desalted using self-packed StageTips (five disks per micro-column, ø 1.5 mm, C18 material, 3M Empore). The peptide eluates were dried to completeness and stored at -80 °C. Before the LC-MS/MS measurement, all samples were freshly resuspended in 12 μl 0.1% FA in HPLC grade water, and around 25 μg of total peptide amount was injected into the mass spectrometer per measurement. Each experiment was performed using four biological replicates.

Proteomic data acquisition and data analysis

Peptides were analysed on a Vanquish Neo liquid chromatography system (micro-flow configuration) coupled to an Orbitrap Exploris 480 mass spectrometer (Thermo Fisher Scientific). Around 25 µg of peptides were applied onto an Acclaim PepMap 100 C18 column (2 µm particle size, 1 mm ID × 150 mm, 100 Å pore size; Thermo Fisher Scientific) and separated using a two-stepped gradient. In the first step, a 50-minute linear gradient ranging from 3% to 24% solvent B (0.1% FA, 3% DMSO in ACN) in solvent A (0.1% FA, 3% DMSO in HPLC grade water) at a flow rate of 50 µl/min was applied. In the second step, solvent B was further increased from 24% to 31% over a 10-minute linear gradient. The mass spectrometer was operated in data-dependent acquisition (DDA) and positive ionisation mode. MS1 full scans (360 – 1300 m/z) were acquired with a resolution of 60,000, a normalised AGC target value of 100% and a maximum injection time of 50 msec. Peptide precursor selection for fragmentation was carried out using a fixed cycle time of 1.2 sec. Only precursors with charge states from 2 to 6 were selected, and dynamic exclusion of 30 sec was enabled. Peptide fragmentation was performed using higher energy collision-induced dissociation (HCD) and normalised collision energy of 28%. The precursor isolation window width of the quadrupole was set to 1.1 m/z. MS2 spectra were acquired with a resolution of 15,000, a fixed first mass of 100 m/z, a normalised automatic gain control (AGC) target value of 100% and a maximum injection time of 40 msec.

Peptide identification and quantification were performed using MaxQuant (version 1.6.3.4) with its built-in search engine Andromeda[76,77]. MS2 spectra were searched against the *C. jejuni or A. butzleri* proteome database derived from *C. jejuni* NCTC 11168 (NC_002163.1) or *A. butzleri* RM4018 (NC_009850.1), respectively. Trypsin/P was specified as the proteolytic enzyme. The precursor tolerance was set to 4.5 ppm, and fragment ion tolerance to 20 ppm. Results were adjusted to a 1% false discovery rate (FDR) on peptide spectrum match (PSM) level and protein level employing a target-decoy approach using reversed protein sequences. The minimal peptide length was defined as seven amino acids, carbamidomethylated cysteine was set as fixed modification and oxidation of methionine and N-terminal protein acetylation as variable modifications. The match-between-run function was disabled. Protein abundances were calculated using the LFQ algorithm from MaxQuant[78]. Protein intensity values were logarithm transformed (base 2), and a Student t-test was used to identify proteins differentially expressed between conditions. The resulting p-values were adjusted by the Benjamini-Hochberg algorithm[79] to control the false discovery rate (FDR). Since low abundant proteins are more likely to result in missing values, we filled in missing values with a constant of half the lowest detected LFQ intensity per protein. However, if the imputed value was higher than the 20% quantile of all LFQ intensities in that sample, we used the 20% quantile as the imputed value. Only proteins with $|\log_2 FC|$ ≥ 1 were considered differentially expressed[80].

The reproducibility of *C. jejuni* and *A. butzleri* biological replicates was visualized by principal component analysis (PCA) of the normalised LC-MS/MS CPM data (Fig S11c, d).

Omics data analysis

The Kyoto Encyclopedia of Genes and Genomes (KEGG)[81] gene/protein set enrichment analysis of the differentially expressed genes and proteins was performed and visualised based on the clusterProfiler[28] (version 4.8.2) package in R software with P < 0.05. Genome-wide functional annotation was carried out with the eggNOG-mapper (version 5.0)[82] according to the Cluster of Orthologous Genes (COGs) database[83] with an e-value of 0.001 and minimum hit bit-score of 60. Visual representations of RNA expression levels and LC-MS/MS expressed proteins in different COGs were performed with the Circos Table viewer (http://mkweb.bcgsc.ca/tableviewer/, accessed on 05 November 2023). For the analyses, only genes and proteins with $|\log_2 FC|$ ≥ 1 and FDR < 0.05 were considered as significantly changed.

Quantitative polymerase chain reaction

RT-qPCR quantified the mRNA levels of the selected genes. The reverse transcription was conducted using 500 ng of RNA in a 20 µl volume reaction mixture of iScript cDNA Synthesis Kit (Bio-Rad). 2.5 µl of 1:10 diluted cDNA was added to 7.5 µl of Sensi-FAST SYBR No-ROX (Bioline) and 400 nM of forward and reverse primers in a 15 µl final volume. The RT-qPCR program: 95 °C for 3 min, followed by 40 three-step amplification cycles consisting of 5 s at 95 °C, 10 s at 58 °C and 20 s at 72 °C. The following primer pairs were used: P18-P19, *gltA* for *C. jejuni* RT-qPCR and P23-P24, *nuoB* for *A. butzleri* RT-qPCR (Table S2). The relative quantity of mRNA for each gene of *C. jejuni* was determined by referring to the mRNA levels of *recA* (P25-P26 primer pair). The relative quantity of mRNA for each gene of *A. butzleri* was determined by referring to the mRNA levels of *gyrA* (P27-P28 primer pair). The RT-qPCR was performed for three independent

bacterial cultures. The RT-qPCR was performed using the CFX96 Touch Real-Time PCR Detection Systems, and data were analysed with CFX Maestro (Bio-Rad) software.

Construction of plasmids expressing recombinant wild-type Cj1608 and Abu0127 proteins

The *Cj1608* (*Cj1509* in *C. jejuni* 81-116) gene (888 bp) was amplified with primer pairs P29-P30 using *C. jejuni* 81-116 genomic DNA as a template. The *Abu0127* gene (891 bp) was amplified with primer pair P31-P32 using *A. butzleri* RM4018 genomic DNA as a template. The PCR products were digested with BamHI/SalI and cloned into BamHI/SalI sites of pET28Strep[22] to generate pETStrepCj1509 and pETStrepAbu0127, respectively (Table S1).

Protein expression and purification

The recombinant Strep-tagged Cj1608 (Cj1509 of *C. jejuni* 81-116) and Abu0127 proteins were purified according to the Strep-Tactin manufacturer's protocol (IBA Lifesciences). Briefly, *E. coli* BL21 cells (1 l) carrying either the pET28/StrepCj1509 or pET28/StrepAbu0127 vectors were grown at 37 °C. At an OD_{600} of 0.8, protein synthesis was induced with 0.05 mM IPTG for 3 hours at 30 °C for Cj1608 and overnight at 18 °C for Abu0127. The cultures were harvested by centrifugation (10 min, 5,000 × g and 4 °C). The cells were suspended in 20 ml of ice-cold buffer W (100 mM Tris-HCl, pH 8.0; 300 mM NaCl; and 1 mM EDTA) supplemented with cOmplete, EDTA-free protease inhibitor Cocktail (Roche)), disrupted by sonication, and centrifuged (30 min, 31,000 × g and 4 °C). The supernatant was applied onto a Strep-Tactin Superflow high-capacity Sepharose column (1 ml bed volume, IBA). The column was washed with buffer W until Bradford tests yielded a negative result, and then it was washed with 5 ml of buffer W without EDTA. The elution was carried out with approx. 6 × 0.8 ml of buffer E (100 mM Tris-HCl, pH 8.0; 300 mM NaCl; and 5 mM desthiobiotin). Protein purity was analysed by SDS-PAGE electrophoresis using a GelDoc XR+ and ImageLab software (BioRad). The fractions were stored at −20 °C in buffer E diluted with glycerol to a final concentration of 50%.

Electrophoretic mobility shift assay (EMSA)

PCR amplified DNA probes in two steps. DNA fragments were amplified in the first step using unlabeled primer pairs P33-P34 (p*gltA*), P35-P36 (*recJ*) with a *C. jejuni* NCTC 11168 genomic DNA template and P37-P38 (p*nuo*), P39-P40 (*recJ*) with an *A. butzleri* RM4018 genomic DNA template (Table S1). The forward primers were designed with overhangs complementary to P41 (FAM–labelled) for p*nuo* and p*gltA* and P42 (Cy5 labelled) for both *recJ*. The unlabeled fragment was purified and used as a template in the second round of PCR using the appropriate fluorophore-labelled primer and the appropriate reverse primer used in the first step. Both fluorophore-labelled DNA (each 5 nM) were incubated with the Strep-tagged protein at 30°C (Abu0127) or 37°C (Cj1608) for 20 min in Tris buffer (50 mM Tris–HCl, pH 8.0; 100 mM NaCl and 0.2% Triton X-100). The complexes were separated by electrophoresis on a 4% polyacrylamide gel in 0.5 × TBE (1 × TBE: 89 mM Tris, 89 mM borate and 2 mM EDTA) at 10 V/cm in the cold room (approx. 10 °C). The gels were analysed by a Typhoon 9500 FLA Imager and ImageQuant software.

ATP assay

The ATP level was measured with the BacTiter-Glo™ Assay. *C. jejuni and A. butzleri* cultures were grown to the logarithmic growth phase (OD_{600} of 0.5-0.7) in the BHI medium. Cells were diluted to $OD_{600} = 0.1$ with fresh medium. Next, 50 µl of bacteria were mixed with 50 µl of BacTiter-Glo™ (Promega) and incubated at room temperature for 5 min. The luminescence was measured with a CLARIOstar® plate reader on opaque-walled multiwell plates (SPL Life Sciences; 2-200203). Each experiment was performed using three biological replicates.

Statistics and Reproducibility

Statistical analysis was performed using GraphPad Prism (8.4.2) and R (4.3.1) statistical software. All in vivo experiments were repeated at least three times, and data were presented as mean ± SD. The statistical significance between the two conditions was calculated by paired two-tailed Student's t-test. The statistical significance between multiple groups was calculated by one-way ANOVA with Tukey's post hoc test. $P < 0.05$

was considered statistically significant. The proteome and transcriptome correlation were determined with the Pearson correlation coefficient. The EMSA and Western blot experiments were repeated twice with similar results.

Data availability

The *C. jejuni* RNA-seq FASTQ and processed data generated in this study have been deposited in the ArrayExpress database (EMBL-EBI) under accession code E-MTAB-13650.
The *A. butzleri* RNA-seq FASTQ and processed data generated in this study have been deposited in the ArrayExpress database (EMBL-EBI) under accession code E-MTAB-13649.
The raw proteomics data, MaxQuant search results, and the used protein sequence database generated in this study have been deposited in the ProteomeXchange Consortium via the PRIDE partner repository[84] under accession code PXD048711.
Campylobacter jejuni NCTC 11168 reference genome is deposited in the National Center for Biotechnology Information under accession code NC_002163.1 (https://www.ncbi.nlm.nih.gov/nuccore/NC_002163.1).
Arcobacter butzleri RM4018 reference genome is deposited in the National Center for Biotechnology Information under accession code NC_009850.1 (https://www.ncbi.nlm.nih.gov/nuccore/NC_009850).
Source data are provided with this paper.

Code availability

All of the code and data used to generate the figures presented here are deposited in GitHub.

References

1. Waite, D. W., Chuvochina, M. S. & Hugenholtz, P. Road Map of the Phylum *Campylobacterota*. *Bergey's Man. Syst. Archaea Bact.* 1–11 (2019).
2. Chieffi, D., Fanelli, F. & Fusco, V. *Arcobacter butzleri*: Up-to-date taxonomy, ecology, and pathogenicity of an emerging pathogen. *Compr. Rev. food Sci. food Saf.* **19**, 2071–2109 (2020).
3. Burnham, P. M. & Hendrixson, D. R. *Campylobacter jejuni*: collective components promoting a successful enteric lifestyle. *Nat. Rev. Microbiol. 2018 169* **16**, 551–565 (2018).
4. Shange, N., Gouws, P. & Hoffman, L. C. *Campylobacter* and *Arcobacter* species in food-producing animals: prevalence at primary production and during slaughter. *World J. Microbiol. Biotechnol. 2019 359* **35**, 1–16 (2019).
5. Tacconelli, E. *et al.* Discovery, research, and development of new antibiotics: the WHO priority list of antibiotic-resistant bacteria and tuberculosis. *Lancet Infect. Dis.* **18**, 318–327 (2018).
6. Sciortino, S. *et al.* Occurrence and Antimicrobial Resistance of Arcobacter spp. Recovered from Aquatic Environments. *Antibiot. 2021, Vol. 10, Page 288* **10**, 288 (2021).
7. Gumerov, V. M., Ulric, L. E., Or, I. & Zhulin, B. MiST 4.0: a new release of the microbial signal transduction database, now with a metagenomic component. *Nucleic Acids Res.* **2023**, 1–7 (2023).
8. Gundogdu, O. *et al.* Re-annotation and re-analysis of the *Campylobacter jejuni* NCTC11168 genome sequence. *BMC Genomics* **8**, 1–8 (2007).
9. Goodman, K. N., Powers, M. J., Crofts, A. A., Trent, M. S. & Hendrixson, D. R. *Campylobacter jejuni* BumSR directs a response to butyrate via sensor phosphatase activity to impact transcription and colonization. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 11715–11726 (2020).
10. Miller, W. G. *et al.* The Complete Genome Sequence and Analysis of the *Epsilonproteobacterium Arcobacter butzleri*. *PLoS One* **2**, (2007).
11. Martinez-Malaxetxebarria, I. *et al.* Regulation of Energy Metabolism by the Extracytoplasmic Function (ECF) σ Factors of *Arcobacter butzleri*. *PLoS One* **7**, e44796 (2012).
12. Gahamanyi, N. *et al.* Insights into the Virulence of *Campylobacter jejuni* Associated with Two-Component Signal Transduction Systems and Single Regulators. *Microbiol. Res. 2022, Vol. 13, Pages 188-200* **13**, 188–200 (2022).
13. Chandrashekhar, K., Kassem, I. I. & Rajashekara, G. *Campylobacter jejuni* transducer like proteins: Chemotaxis and beyond. *Gut Microbes* **8**, 323 (2017).
14. Casado, J., Lanas, Á. & González, A. Two-component regulatory systems in *Helicobacter pylori* and Campylobacter jejuni: Attractive targets for novel antibacterial drugs. *Front. Cell. Infect. Microbiol.* **12**, 977944 (2022).
15. Taylor, A. J. & Kelly, D. J. The function, biogenesis and regulation of the electron transport chains in *Campylobacter jejuni*: New insights into the bioenergetics of a major food-borne pathogen. *Adv. Microb. Physiol.* **74**, 239–329 (2019).
16. Flint, A., Stintzi, A. & Saraiva, L. M. Oxidative and nitrosative stress defences of *Helicobacter* and *Campylobacter* species that counteract mammalian immunity. *FEMS Microbiol. Rev.* **40**, 938–960 (2016).
17. Espey, M. G. Role of oxygen gradients in shaping redox relationships between the human intestine and its microbiota. *Free Radic. Biol. Med.* **55**, 130–140 (2013).
18. van der Stel, A. X. *et al.* Catabolite repression in *Campylobacter jejuni* correlates with intracellular succinate levels. *Environ. Microbiol.* **20**, 1374–1388 (2018).
19. van der Stel, A.-X. & Wösten, M. M. S. M. Regulation of Respiratory Pathways in *Campylobacterota*: A Review. *Front. Microbiol.* **10**, 424488 (2019).
20. Ravcheev, D. A., Gerasimova, A. V., Mironov, A. A. & Gelfand, M. S. Comparative genomic analysis of regulation of anaerobic respiration in ten genomes from three families of gamma-proteobacteria (Enterobacteriaceae, Pasteurellaceae, Vibrionaceae). *BMC Genomics* **8**, 1–17 (2007).
21. Tomb, J. F. *et al.* The complete genome sequence of the gastric pathogen *Helicobacter pylori*. *Nature* **388**, 539–547 (1997).
22. Szczepanowski, P. *et al.* HP1021 is a redox switch protein identified in *Helicobacter pylori*. *Nucleic Acids Res.* **49**, 6863–6879 (2021).
23. Pflock, M. *et al.* The Orphan Response Regulator HP1021 of *Helicobacter pylori* Regulates Transcription of a Gene Cluster Presumably Involved in Acetone Metabolism. *J. Bacteriol.* **189**, 2339 (2007).
24. Noszka, M. *et al.* Profiling of the *Helicobacter pylori* redox switch HP1021 regulon using a multi-

omics approach. *Nat. Commun. 2023 141* **14**, 1–14 (2023).

25. Donczew, R. *et al.* The atypical response regulator HP1021 controls formation of the *Helicobacter pylori* replication initiation complex. *Mol. Microbiol.* **95**, 297–312 (2015).

26. Sinha, R. *et al.* Gut metabolite L-lactate supports *Campylobacter jejuni* population expansion during acute infection. *Proc. Natl. Acad. Sci.* **121**, e2316540120 (2024).

27. Jaworski, P. *et al.* Structure and function of the *Campylobacter jejuni* chromosome replication origin. *Front. Microbiol.* **9**, 1533 (2018).

28. Wu, T. *et al.* clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. *Innovation* **2**, (2021).

29. Li, S., Lam, J., Souliotis, L., Alam, M. T. & Constantinidou, C. Posttranscriptional Regulation in Response to Different Environmental Stresses in *Campylobacter jejuni*. *Microbiol. Spectr.* **10**, (2022).

30. Sharma, C. M. *et al.* The primary transcriptome of the major human pathogen *Helicobacter pylori*. *Nat. 2010 4647286* **464**, 250–255 (2010).

31. Fasnacht, M. & Polacek, N. Oxidative Stress in Bacteria and the Central Dogma of Molecular Biology. *Front. Mol. Biosci.* **8**, 392 (2021).

32. Stahl, M., Butcher, J. & Stintzi, A. Nutrient Acquisition and Metabolism by *Campylobacter jejuni*. *Front. Cell. Infect. Microbiol.* **2**, 5 (2012).

33. Surmacz, K. Analysis of *Aliarcobacter butzleri* RM4018 sensitivity to oxidative stress. (Wroclaw University of Science and Technology, 2022).

34. Guccione, E. J. *et al.* Transcriptome and proteome dynamics in chemostat culture reveal how *Campylobacter jejuni* modulates metabolism, stress responses and virulence factors upon changes in oxygen availability. *Environ. Microbiol.* **19**, 4326 (2017).

35. Martinez-Malaxetxebarria, I. *et al.* Regulation of Energy Metabolism by the Extracytoplasmic Function (ECF) σ Factors of *Arcobacter butzleri*. *PLoS One* **7**, e44796 (2012).

36. Kiehlbauch, J. A. *et al. Campylobacter butzleri* sp. nov. isolated from humans and animals with diarrheal illness. *J. Clin. Microbiol.* **29**, 376 (1991).

37. Ward, B. Bacterial Energy Metabolism. *Mol. Med. Microbiol.* 201–233 (2015) doi:10.1016/B978-0-12-397169-2.00011-1.

38. Perez, J. C. & Groisman, E. A. Evolution of transcriptional regulatory circuits in bacteria. *Cell* **138**, 233 (2009).

39. Federowicz, S. *et al.* Determining the control circuitry of redox metabolism at the genome-scale. *PLoS Genet.* **10**, (2014).

40. Myers, K. S. *et al.* Genome-scale Analysis of Escherichia coli FNR Reveals Complex Features of Transcription Factor Binding. *PLOS Genet.* **9**, e1003565 (2013).

41. Brown, A. N., Anderson, M. T., Bachman, M. A. & Mobley, H. L. T. The ArcAB Two-Component System: Function in Metabolism, Redox Control, and Infection. *Microbiol. Mol. Biol. Rev.* **86**, (2022).

42. Park, D. M., Akhtar, M. S., Ansari, A. Z., Landick, R. & Kiley, P. J. The bacterial response regulator ArcA uses a diverse binding site architecture to regulate carbon oxidation globally. *PLoS Genet.* **9**, (2013).

43. Stewart, V. Dual interacting two-component regulatory systems mediate nitrate- and nitrite-regulated gene expression in *Escherichia coli*. *Res. Microbiol.* **145**, 450–454 (1994).

44. Iyer, M. S., Pal, A., Srinivasan, S., Somvanshi, P. R. & Venkatesh, K. V. Global Transcriptional Regulators Fine-Tune the Translational and Metabolic Efficiency for Optimal Growth of *Escherichia coli*. *mSystems* **6**, (2021).

45. Overton, T. W. *et al.* Microarray analysis of gene regulation by oxygen, nitrate, nitrite, FNR, NarL and NarP during anaerobic growth of Escherichia coli: new insights into microbial physiology. *Biochem. Soc. Trans.* **34**, 104–107 (2006).

46. Unden, G. & Bongaerts, J. Alternative respiratory pathways of *Escherichia coli*: energetics and transcriptional regulation in response to electron acceptors. *Biochim. Biophys. Acta* **1320**, 217–234 (1997).

47. van der Stel, A. X. *et al.* The *Campylobacter jejuni* RacRS system regulates fumarate utilization in a low oxygen environment. *Environ. Microbiol.* **17**, 1049–1064 (2015).

48. Galperin, M. Y. A census of membrane-bound and intracellular signal transduction proteins in bacteria: Bacterial IQ, extroverts and introverts. *BMC Microbiol.* **5**, 1–19 (2005).

49. Bell, T. H. & Bell, T. Many roads to bacterial generalism. *FEMS Microbiol. Ecol.* **97**, 240 (2021).

50. Bury-Moné, S. *et al.* Is *Helicobacter pylori* a True Microaerophile? *Helicobacter* **11**, 296–303 (2006).

51. Pelliciari, S. *et al.* Insight into the essential role of the *Helicobacter pylori* HP1043 orphan response

regulator: genome-wide identification and characterization of the DNA-binding sites. *Sci. Reports 2017 71* **7**, 1–15 (2017).

52. Olekhnovich, I. N., Vitko, S., Valliere, M. & Hoffmana, P. S. Response to metronidazole and oxidative stress is mediated through homeostatic regulator hsra (hp1043) in *Helicobacter pylori*. *J. Bacteriol.* **196**, 729–739 (2014).

53. Loh, J. T. *et al.* Delineation of the pH-responsive regulon controlled by the *Helicobacter pylori* ArsRS two-component system. *Infect. Immun.* **89**, (2021).

54. Vannini, A. *et al.* Comprehensive mapping of the *Helicobacter pylori* NikR regulon provides new insights in bacterial nickel responses. *Sci. Reports 2017 71* **7**, 1–14 (2017).

55. Gölz, G., Kittler, S., Malakauskas, M. & Alter, T. Survival of Campylobacter in the Food Chain and the Environment. *Curr. Clin. Microbiol. Reports* **5**, 126–134 (2018).

56. Dufour, V. *et al.* Inactivation of the LysR regulator Cj1000 of *Campylobacter jejuni* affects host colonization and respiration. *Microbiol. (United Kingdom)* **159**, 1165–1178 (2013).

57. Svensson, S. L., Hyunh, S., Parker, C. T. & Gaynor, E. C. The *Campylobacter jejuni* CprRS two-component regulatory system regulates aspects of the cell envelope. *Mol. Microbiol.* **96**, 189–209 (2015).

58. Fields, J. A., Li, J., Gulbronson, C. J., Hendrixson, D. R. & Thompson, S. A. *Campylobacter jejuni* CsrA Regulates Metabolic and Virulence Associated Proteins and Is Necessary for Mouse Colonization. *PLoS One* **11**, e0156932 (2016).

59. Price, E. E., Román-Rodríguez, F. & Boyd, J. M. Bacterial approaches to sensing and responding to respiration and respiration metabolites. *Mol. Microbiol.* **116**, 1009–1021 (2021).

60. Contreras, M., Thiberge, J. M., Mandrand-Berthelot, M. A. & Labigne, A. Characterization of the roles of NikR, a nickel-responsive pleiotropic autoregulator of *Helicobacter pylori*. *Mol. Microbiol.* **49**, 947–963 (2003).

61. Atabay, H. I. & Corry, J. E. L. Evaluation of a new arcobacter enrichment medium and comparison with two media developed for enrichment of *Campylobacter spp. Int. J. Food Microbiol.* **41**, 53–58 (1998).

62. Beauchamp, J. M., Leveque, R. M., Dawid, S. & DiRita, V. J. Methylation-dependent DNA discrimination in natural transformation of *Campylobacter jejuni. Proc. Natl. Acad. Sci. U. S. A.* **114**, E8053–E8061 (2017).

63. Gibson, D. G. *et al.* Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nat. Methods 2009 65* **6**, 343–345 (2009).

64. Dalia, A. B., McDonough, E. & Camilli, A. Multiplex genome editing by natural transformation. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 8937–8942 (2014).

65. Cieślak, M. Construction of the plasmid for the analysis of the essentiality of the Abu0127 gene for the survival of *Arcobacter butzleri* RM4018. (Wroclaw University of Science and Technology -, 2021).

66. Perkins, A., Tudorica, D. A., Amieva, M. R., James Remington, S. & Guillemin, K. *Helicobacter pylori* senses bleach (HOCl) as a chemoattractant using a cytosolic chemoreceptor. *PLOS Biol.* **17**, e3000395 (2019).

67. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods 2012 94* **9**, 357–359 (2012).

68. Langmead, B., Wilks, C., Antonescu, V. & Charles, R. Scaling read aligners to hundreds of threads on general-purpose processors. *Bioinformatics* **35**, 421–432 (2019).

69. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079 (2009).

70. Liao, Y., Smyth, G. K. & Shi, W. The R package Rsubread is easier, faster, cheaper and better for alignment and quantification of RNA sequencing reads. *Nucleic Acids Res.* **47**, (2019).

71. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–140 (2010).

72. Chen, Y., Lun, A. T. L. & Smyth, G. K. From reads to genes to pathways: Differential expression analysis of RNA-Seq experiments using Rsubread and the edgeR quasi-likelihood pipeline. *F1000Research* **5**, (2016).

73. Blighe, K, S Rana, and M Lewis. EnhancedVolcano: publication-ready volcano plots with enhanced colouring and labeling. https://github.com/kevinblighe/EnhancedVolcano (2018). 74.   Abele, M. *et al.* Unified workflow for the rapid and in-depth characterization of bacterial proteomes. *Mol. Cell. Proteomics* **22**, 100612 (2023).

75. Doellinger, J., Schneider, A., Hoeller, M. & Lasch, P. Sample Preparation by Easy Extraction and

Digestion (SPEED) - A Universal, Rapid, and Detergent-free Protocol for Proteomics Based on Acid Extraction. *Mol. Cell. Proteomics* **19**, 209–222 (2020).

76. Cox, J. *et al.* Andromeda: a peptide search engine integrated into the MaxQuant environment. *J. Proteome Res.* **10**, 1794–1805 (2011).

77. Tyanova, S., Temu, T. & Cox, J. The MaxQuant computational platform for mass spectrometry-based shotgun proteomics. *Nat. Protoc. 2016 1112* **11**, 2301–2319 (2016).

78. Cox, J. *et al.* Accurate proteome-wide label-free quantification by delayed normalization and maximal peptide ratio extraction, termed MaxLFQ. *Mol. Cell. Proteomics* **13**, 2513–2526 (2014).

79. Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *J. R. Stat. Soc. Ser. B* **57**, 289–300 (1995).

80. GitHub - mengchen18/omicsViewer. https://github.com/mengchen18/omicsViewer.

81. Kanehisa, M., Furumichi, M., Tanabe, M., Sato, Y. & Morishima, K. KEGG: new perspectives on genomes, pathways, diseases and drugs. *Nucleic Acids Res.* **45**, D353–D361 (2017).

82. Huerta-Cepas, J. *et al.* eggNOG 5.0: a hierarchical, functionally and phylogenetically annotated orthology resource based on 5090 organisms and 2502 viruses. *Nucleic Acids Res.* **47**, D309–D314 (2019).

83. Galperin, M. Y., Makarova, K. S., Wolf, Y. I. & Koonin, E. V. Expanded microbial genome coverage and improved protein family annotation in the COG database. *Nucleic Acids Res.* **43**, D261–D269 (2015).

84. Perez-Riverol, Y. *et al.* The PRIDE database resources in 2022: a hub for mass spectrometry-based proteomics evidences. *Nucleic Acids Res.* **50**, D543–D552 (2022).

Acknowledgements

This work has been supported by the OPUS 17, Project Number 2019/33/B/NZ6/01648, founded by the National Science Centre, Poland, to A.Z-P.
This work has been supported by the EPIC-XS, Project Number 823839, funded by the Horizon 2020 Program of the European Union, to C.L.
We thank Kinga Surmacz for help in disc diffusion assays. We thank Maria Cieślak for help in pUC18/ΔAbu0127 plasmid preparation. We thank Verena Breitner and Tayma Midari for technical assistance at BayBioMS.

Author contributions

M.N., K.S. and A.Z-P conceived and designed the experiments. M.N., J.M., M.A. and K.S. performed the experiments. M.N., A.S., D.H., M.A., C.L. and A.Z-P analysed the data. M.N., and A.Z-P wrote the manuscript. M.N., A.S., J.M., D.H., M.A., C.L., K.S. and A.Z-P revised the manuscript. C.L. and A.Z-P provided funding.

Competing interests

The authors declare no competing interests.

Fig. 1: RNA-seq and LC-MS/MS analyses of *C. jejuni* NCTC 11168 gene expression controlled by Cj1608.
a, The comparison of gene transcription in the *C. jejuni* wild-type (CJ), stressed wild-type (CJS) and ΔCj1608 cells revealed by RNA-seq. Genes whose transcription significantly changed ($|\log_2 FC| \geq 1$; FDR ≤ 0.05) are depicted by coloured dots (see the legend in the figure). The genes named on the graph correspond to the citric acid cycle and electron transport chain. **b,** Venn diagram presenting the number of differentially transcribed genes in the analysed strains and conditions. **c,** ClusterProfiler protein enrichment plot showing activation or suppression of KEGG groups in the analysed strains. **d,** The correlation between gene expression in ΔCj1608 and CJ cells at the transcription and proteome levels. Red dots represent homodirectional genes – up-regulated or down-regulated – at the transcriptome and proteome levels. Yellow dots represent opposite changes. Blue dots represent genes that changed only at the transcription level, while green dots represent genes that changed only at the proteome level. Numbers of differentially expressed genes in the indicated strains and conditions are given in parentheses. The proteins named on the graph correspond to the citric acid cycle and electron transport chain. **e,** ClusterProfiler protein enrichment plot showing activation or suppression of KEGG groups in the analysed strains. **a, d,** Values outside the black dashed lines indicate a change in the expression of $|\log_2 FC| \geq 1$. Grey dots correspond to genes whose transcription was not significantly changed (FDR ≤ 0.05).

Fig. 2: RNA-seq and LC-MS/MS analyses of *A. butzleri* RM4018 gene expression controlled by Abu0127.
a, The comparison of gene transcription in the *A. butzleri* wild-type (AB), stressed wild-type (ABS) and ΔAbu0127 cells revealed by RNA-seq. Genes whose transcription significantly changed (|log$_2$FC| ≥ 1; FDR ≤ 0.05) are depicted by coloured dots (see the legend in the figure). The genes named on the graph correspond to the citric acid cycle and electron transport chain. **b,** Venn diagram presenting the number of differentially transcribed genes in the analysed strains and conditions. **c,** ClusterProfiler protein enrichment plot showing activation or suppression of KEGG groups in the analysed strains. **d,** The correlation between gene expression of ΔAbu0127 and AB cells at the transcription and proteome levels. Red dots represent homodirectional genes – up-regulated or down-regulated – at the transcriptome and proteome levels. Yellow dots represent opposite changes. Blue dots represent genes that changed only at the transcription level, while green dots represent genes that changed only at the proteome level. Numbers of differentially expressed genes in the indicated strains and conditions are given in parentheses. The proteins named on the graph correspond to the citric acid cycle and electron transport chain. **e,** ClusterProfiler protein enrichment plot showing activation or suppression of KEGG groups in the analysed strains. **a, d,** Values outside the black dashed lines indicate a change in the expression of |log$_2$FC| ≥ 1. Grey dots correspond to genes whose transcription was not significantly changed (FDR ≤ 0.05).

Fig. 3: Schematic presentation of *C. jejuni* and *A. butzleri* energy conservation factors whose expression changed in ΔCj1608 and ΔAbu0127, respectively. **a,** Regulation of the *C. jejuni* citric acid cycle, the selected associated catabolic pathways, and nutrient transporters. **b,** Regulation of *C. jejuni* electron transport chain enzymes. **c,** Regulation of the *A. butzleri* tricarboxylic acid cycle selected associated catabolic pathways and nutrient transporters. **d,** Regulation of *A. butlzeri* electron transport chain enzymes. **a-d,** The increase or decrease in protein abundance in the deletion mutant strains compared to wild-type strains is shown based on the LC-MS/MS data (Supplementary data 3). When the protein was not detected by MS, but the corresponding transcript level was increased or decreased, the colour code depicts RNA-seq results (denoted as •). For details concerning transcriptomic data or complex transcriptome/proteome interpretation (denoted as *), see the attached Supplementary Data 1-3. Frd, fumarate reductases; Hyd, Hya, Hua, hydrogenases, Lut, Dld, L-lactate utilisation complex, Fdh, formate dehydrogenase; Mqo, malate:quinone oxidoreductase, Nuo, NADH-quinone oxidoreductase; Nrf, nitrate reductase; Tor, SN oxide reductase; Nap, nitrate reductase; Mfr, methylmenaquinol:fumarate reductase (misannotated as succinate dehydrogenase Sdh); SOR, sulfite:cytochrome c oxidoreductase; Cio, cyanide-insensitive cytochrome bd-like quinol oxidase; Cco, cbb3-type cytochrome c oxidase; Pet, ubiquinol-cytochrome C reductase; Gdh, gluconate 2-dehydrogenase; SOX, thiosulfate oxidation by SOX system; Oor, 2-oxoglutarate:acceptor oxidoreductase; POR, pyruvate:acceptor oxidoreductase; PEP, phosphoeneoopyruvate; G-6Pglucose 6-phosphate.; "?" unknown enzyme. All genes and enzyme complexes can be found in Supplementary Data 3.

C - energy production and conversion, **D** - cell cycle control, cell division, chromosome partitioning, **E** - amino acid transport and metabolism, **F** - nucleotide transport and metabolism, **G** - carbohydrate transport and metabolism, **H** - coenzyme transport and metabolism, **I** - lipid transport and metabolism, **J** - translation, ribosomal structure and biogenesis, **K** - transcription, **L** - replication, recombination and repair, **M** - cell wall/membrane/envelope biogenesis, **N** - cell motility, **O** - posttranslational modification, protein turnover, chaperones , **P** - inorganic ion transport and metabolism, **Q** - secondary metabolites biosynthesis, transport and catabolism, **T** - signal transduction mechanisms, **U** - intracellular trafficking, secretion, and vesicular transport, **V** - defense mechanisms

Fig. 4: Impact of Cj1608 and Abu0127 on gene regulation in *C. jejuni* and *A. butzleri*, respectively. **a,** Circos plots presenting the correlation between Cluster of Orthologues Group (COG) and differentially expressed *C. jejuni* genes revealed by RNA-seq in the ΔCj1608 strain compared to the *C. jejuni* wild-type strain. **b,** Circos plots presenting the correlation between COG and differentially expressed *A. butzleri* genes revealed by RNA-seq in the ΔAbu0127 strain compared to the *A. butzleri* wild-type strain. **c,** Circos plots presenting the correlation between COG and differentially expressed *C. jejuni* proteins revealed by LC-MS/MS in the ΔCj1608 strain compared to the *C. jejuni* wild-type strain. **d,** Circos plots presenting the correlation between COG and differentially expressed *A. butzleri* proteins revealed by LC-MS/MS in the ΔAbu0127 strain compared to the *A. butzleri* wild-type strain. **a-d,** Expression changes of log$_2$FC| ≥ 1 and FDR ≥ 0.05 were considered significant and included in the analyses. The category of unknown genes was excluded from the analysis.

Fig. 5: Cj1608-dependent control of *C. jejuni* energy conservation pathways. a, Growth curves of *C. jejuni* wild-type (CJ) and ΔCj1608 strains cultivated in the presence of H₂ or without H₂ (see Materials and Methods). **b,** The generation time of *C. jejuni* CJ and ΔCj1608 strains cultivated as in part a. **c,** ATP production by CJ and ΔCj1608 strains cultivated as in part a. **d,** RNA-seq data profile comparing *gltA* transcription in CJ and ΔCj1608 strains under microaerobic growth and paraquat-induced oxidative stress. Values above the black dashed lines indicate a change in the expression of |log₂FC| ≥ 1; FDR ≤ 0.05. **e,** Logarithmically transformed LFQ intensities of GltA protein in CJ and ΔCj1608 strains. **f,** EMSA analysis of Cj1608 binding to the p*gltA*-FAM promoter region *in vitro*. The *recJ*-Cy5 fragment was used as a negative control. The experiment was repeated twice with similar results. **g,** RT-qPCR analysis of *gltA* transcription in CJ and ΔCj1608 cells cultured under microaerobic or paraquat-induced oxidative stress conditions. **h,** RT-qPCR analysis of *gltA* transcription in CJ and ΔCj1608 cells cultured at different oxygen concentrations: (i) 1% O₂, (ii) 5% O₂, or (iii) 10% O₂ (see Materials and Methods). **i,** Growth curves of CJ and ΔCj1608 strains cultivated as in part h. **j,** ATP production by CJ and ΔCj1608 strains cultivated as in part h. **a-c, e, g-k,** Data presented as the mean values ± SD. Ordinary one-way ANOVA with Tukey's multiple comparison test determined the P value. **e,** Student's unpaired t-test determined the P value. n = 4 biologically independent experiments **a-c, g-k,** n = 3 biologically independent experiments. ND, non-determined.

Fig. 6: Abu0127-dependent control of *A. butzleri* energy conservation pathways. a, Growth curves of *A. butzleri* wild-type (AB) and ΔAbu0127 strains cultivated in the presence of H_2 or without H_2 (see Materials and Methods). **b,** The generation time of AB and ΔAbu0127 strains cultivated as in part a. **c,** ATP production by AB and ΔAbu0127 strains cultivated as in part a. **d,** RNA-seq data profile comparing the transcription of the *nuo* operon in AB and ΔAbu0127 strains under microaerobic growth and paraquat-induced oxidative stress. Values above the black dashed lines indicate a change in the expression of $|\log_2 FC| \geq 1$; FDR ≤ 0.05. **e,** Logarithmically transformed LFQ intensities of Nuo B-F, GltA, NuoG, NuoI proteins in AB and ΔAbu0127 strains. **f,** EMSA analysis of Abu0127 binding to the p*nuoA*-FAM promoter region *in vitro*. The *recJ*-Cy5 fragment was used as a negative control. The experiment was repeated twice with similar results. **g,** RT-qPCR analysis of *nuoB* transcription in *A. butzleri* cultured under microaerobic or paraquat-induced oxidative stress conditions. **h,** RT-qPCR analysis of *nuoB* transcription in AB and ΔAbu0127 strains cultured at different oxygen concentrations: (i) 1% O_2; (ii) 5% O_2 or (iii) 10% O_2 (see Materials and Methods). **i,** Growth curves of AB and ΔAbu0127 strains cultivated as in part h. **j,** ATP production by AB and ΔAbu0127 strains cultivated as in part h. **a-c, e, g-j,** Data presented as the mean values ± SD. Ordinary one-way ANOVA with Tukey's multiple comparison test determined the P value. **e,** n = 4 biologically independent experiments **a-d, g-j, n** = 3 biologically independent experiments.

C - energy production and conversion, **D** - cell cycle control, cell division, chromosome partitioning, **E** - amino acid transport and metabolism, **F** - nucleotide transport and metabolism, **G** - carbohydrate transport and metabolism, **H** - coenzyme transport and metabolism, **I** - lipid transport and metabolism, **J** - translation, ribosomal structure and biogenesis, **K** - transcription, **L** - replication, recombination and repair, **M** - cell wall/membrane/envelope biogenesis, **N** - cell motility, **O** - posttranslational modification, protein turnover, chaperones , **P** - inorganic ion transport and metabolism, **Q** - secondary metabolites biosynthesis, transport and catabolism, **T** - signal transduction mechanisms, **U** - intracellular trafficking, secretion, and vesicular transport, **V** - defense mechanisms