Title: Inhibition of NFAT promotes loss of tissue resident uterine natural killer cells and

attendant pregnancy complications in humans

Authors: Rebecca Asiimwe[1], Brittney Knott[1], Morgan E. Greene[1], Emma Wright[1],

Markayla Bell[1], Daniel Epstein[1], Stefani D. Yates[1], Matthew D. Cheung[1], Michael V.

Gonzalez[2], Samantha Fry[1], Emily Boydston[1], Stephanie Clevenger[3], Jayme E. Locke[1],

James F. George[1], Richard Burney[3], Nitin Arora[4], Virginia E. Duncan[5], Holly E. Richter[3],

Deidre Gunn[3], Aharon G. Freud[6], Shawn C. Little[1], Paige M. Porrett*[1]

Affiliations:

[1]Department of Surgery, Marnix E. Heersink School of Medicine, University of Alabama

at Birmingham, Birmingham, AL, 35233 USA

[2]Center for Cytokine Storm Treatment & Laboratory, Department of Medicine, University

of Pennsylvania, Philadelphia, PA, 19104 USA.

[3]Department of Obstetrics & Gynecology, Marnix E. Heersink School of Medicine,

University of Alabama at Birmingham, Birmingham, AL, 35233 USA

[4]Department of Pediatrics, Marnix E. Heersink School of Medicine, University of Alabama

at Birmingham, Birmingham, AL, 35233 USA

[5]Department of Pathology, Marnix E. Heersink School of Medicine, University of Alabama

at Birmingham, Birmingham, AL, 35233 USA

[6]Department of Pathology, The Ohio State University, Columbus, OH 43210, USA

*Corresponding author: Paige M. Porrett (pmporrett@uabmc.edu)

Abstract: Uterine natural killer cells (uNKs) are a tissue resident lymphocyte population

that are critical for pregnancy success. Although mouse models have demonstrated that

NK deficiency results in abnormal placentation and poor pregnancy outcomes, the

generalizability of this knowledge to humans remains unclear. Here we identify uterus

transplant (UTx) recipients as a human population with reduced endometrial NK cells and

altered pregnancy phenotypes. We further show that the NK reduction in UTx is due to

impaired transcriptional programming of NK tissue residency due to blockade of the

transcription factor nuclear factor of activated T cells (NFAT). NFAT-dependent genes

played a role in multiple molecular circuits governing tissue residency in uNKs, including

early residency programs involving AP-1 transcription factors as well as TGFβ-mediated

upregulation of surface integrins. Collectively, our data identify a previously undescribed

role for NFAT in uterine NK tissue residency and provide novel mechanistic insights into

the biologic basis of pregnancy complications due to alteration of tissue resident NK

subsets in humans.

One Sentence Summary: Role of NFAT in uterine NK cell tissue residency

Main Text:

INTRODUCTION

Successful pregnancy requires the interaction of multiple maternal immune cell types with

uterine parenchymal cells and fetal placental trophoblast at the maternal-fetal interface

(*1, 2*). Among the different immune cells that contribute to maternal and fetal health,

uterine natural killer cells (uNKs) play key roles in multiple physiologic aspects of

pregnancy, including (1) production of molecules to support fetal growth and development

(*3, 4*); (2) trophoblast differentiation and placentation (*5*); and (3) uterine spiral artery

remodeling (*6-22*), a process which transforms the uterine vasculature into a low

pressure, high capacitance system thereby augmenting maternal blood flow to the

placenta (*6, 23*). Importantly, mouse models have established that disruptions of uNK cell

number or function result in pregnancy complications or loss, often due to abnormal

placentation associated with poor spiral artery remodeling (*12, 13, 16, 24*). In humans,

incomplete spiral artery remodeling in decidual biopsies has been similarly associated

with structural placental abnormalities as well as clinical syndromes such as pre-

eclampsia (*25, 26*). However, while a limited number of human studies have suggested

that recurrent pregnancy loss is associated with an abnormal distribution of decidual NK

cell subsets (*27*), pregnancy outcomes have not been well studied in individuals with

known defects in NK cell number or function. Hence, the specific contribution of uNK cells

to pregnancy complications in humans remains unclear.

A related knowledge gap that impairs our ability to understand pregnancy complications

is the mechanism by which uNKs acquire tissue residence. Although a local, tissue-based

precursor pool of hematopoietic cells may give rise to uNKs, a prior study which analyzed

samples from uterus transplant recipients suggested that uNKs have a peripheral origin

(*28*). These findings thus predict that uNKs acquire properties of tissue residency, such

as expression of integrins which promote tissue retention, upon localization into the

uterus. While other work has identified mechanisms of tissue residency in other human

organs (*29-33*), mechanisms governing NK residence in the uterus are largely

unexplored.

Organ transplant recipients represent an immunomodulated population who have

reduced numbers of circulating NK cells (*34, 35*) and rates of pre-eclampsia that are six-

fold greater than the general population (*36-41*). Despite the prevalence of pregnancy

complications in organ transplant recipients, the biologic basis for this increased risk is

unknown (*42*). Studies of uNKs in this at-risk population are notably absent, likely

because of challenges associated with acquisition of samples and the frequency with

which pregnancy is discouraged in this population (*43, 44*). Nevertheless, many of these

barriers may be overcome in the setting of uterus transplantation, where transplantation

occurs for the sole purpose of achieving pregnancy and live birth in patients suffering

from absolute uterine factor infertility (*45, 46*). Notably, pregnancy complications such as

pre-eclampsia are common in uterus transplant recipients (UTx) (*45-47*), suggesting that

these individuals represent an opportunity to better understand the immune pathogenesis

of pregnancy complications.

In this study, complementary single cell and imaging approaches were used to study uNKs derived from endometrial biopsies of healthy control volunteers (HC) and UTx. We found that UTx recipients possessed a reduced frequency of tissue resident natural killer cells (trNKs) that associated with clinical pregnancy complications and histologic placental abnormalities. Transcriptional programs of tissue residency in UTx trNKs were disrupted compared to HC trNKs, with impairment of TGFβ-mediated integrin expression as well as AP-1-based early tissue residency programs that are shared with murine CD8 T resident memory cells. Through comparative analyses of ex vivo samples as well as direct pharmacologic inhibition in vitro, we identified a previously undescribed role for NFAT in tissue residency programming of human uNK cells. Taken together, these studies link impaired tissue residency programming with defects in uNK abundance to improve our understanding of the pathogenesis of pregnancy complications in humans.

RESULTS

Reduced frequency of NK cells among endometrial immune cells in UTx

To begin our study of uNKs in human uterus transplant recipients, we performed serial secretory phase endometrial biopsies on two UTx >3 months after transplantation who had resumed normal menstrual cycling (5 biopsies) (table S1). UTx received standard pharmacologic immunosuppression as previously described (*45-47*); this included perioperative T-cell depletion with anti-thymocyte globulin as well as maintenance immunosuppression with a calcineurin inhibitor (i.e., FK506 / tacrolimus), an anti-proliferative agent (i.e., azathioprine), and an oral steroid (i.e., prednisone). Secretory

phase endometrial biopsies were collected from six HC volunteers (table S1) as this time

point is when NK cells are most abundant (*48, 49*). Given the variable expression of CD56

across NK cell subsets, we assured capture of all NK cells by sorting CD45+ cells from

endometrial biopsies and performing scRNA-seq (Fig. 1A). After data processing and

Harmony integration (*50*), we analyzed 77,481 CD45+ endometrial immune cells from 11

biopsies, which included various lymphoid and myeloid populations such as NK cells, T

cells, group 3 innate lymphocyte cells (ILC3s), B cells, mast cells, plasmacytoid dendritic

cells, and monocytes/macrophages (Fig. 1A and fig. S1A).

We initially identified NK cell clusters among CD45+ endometrial immune cells by

expression of conventional marker genes such as *NCAM1* (CD56) and *FCGR3A* (CD16)

(Fig. 1B and fig. S1B). Clusters 0, 7, 8 and 21 comprised tissue resident NK (trNK) cells

based on high expression of *ITGA1* and *NCAM1* (Fig. 1B), low expression of *FCGR3A*

(Fig.1B), and expression of gene signatures associated with decidual NK cell subsets

(i.e., dNK1, dNK2, dNK3) (*45*) (Fig. 1C). Meanwhile, conventional NK cells (cNK) with

intermediate expression of *NCAM1* and high expression of *FCGR3A* were identified in

cluster 3 (Fig. 1B), which also contained CD8 effector T cells sharing cytotoxic gene

expression (Fig. 1B and fig. S1, C to E). In line with prior work (*46*), NK cells in uterus

transplants were >90% recipient-derived (Fig. 1D), confirming that endometrial NK cells

arise from blood-borne precursors. Expectedly, other immune cell subsets (i.e., T cells

and myeloid cells) were also recipient-derived, while contaminating stromal and epithelial

cells were >70% donor-derived (Fig. 1D). Although NK cells were abundant in HC

endometrial biopsies, NK cells were significantly reduced among sequenced immune

cells in UTx biopsies compared to HCs (Fig. 1E). To validate these results, we used flow

cytometry to further compare the frequency of uNK cells between UTx and HCs. Although

tissue for flow cytometric analysis was available from only three of the five UTx biopsies,

these analyses confirmed the statistically significant reduction in NK cell frequency among

endometrial immune cells in UTx versus HCs (Fig. 1F).

Reduced NK cell density in UTx endometrium

The reduction in NK cell frequency among CD45+ endometrial immune cells in UTx could

result from either a reduction in NK cell density in the tissue or from the presence of

excess CD45+ non-NK cells. To determine whether NK cell density was reduced in UTx

endometrium, we performed immunofluorescence microscopy to evaluate multiple fields

of sections of endometrium from UTx and HCs. Biopsy sections (Fig. 2A) were stained

with DAPI and anti-CD56 and data collected by epifluorescence (Fig. 2, B to F) at high

resolution. Nuclei were segmented by custom semi-automated processing of the DAPI

channel (see Methods) and assigned a status of positive or negative for CD56 expression

(Fig. 2C and D). Multiple independent image fields were collected, each representing an

area of about 0.13 mm^2 and containing 870±128 (mean ± SD) cells per field. The fraction

of CD56+ cells was then calculated for each field (Fig. 2G to I).

CD56+ cells were observed either as isolated individuals or in groups (Fig. 2D). Counts

of cells per field varied widely within the same biopsy (Fig. 2G), suggesting non-

homogenous distribution of spatial cues directing NK localization in the tissue in both HC

and UTx endometrium. By comparing the median values of the fraction CD56+ cells, we

found that NK cell density was reduced by roughly a factor of three in UTx biopsies (2.6%

in HC versus 0.9% in UTx). P-values resulting from pairwise Mann-Whitney U tests

showed that UTx samples tended to differ more from control than from each other (Fig.

2I). In contrast, the same analysis using anti-CD3 staining reveals that the density of

CD3+ cells is reduced by about 40% in UTx compared to control (11% in control versus

7% in UTx; fig. S2). We conclude that NK cell density is reduced in UTx, supporting the

observations from flow cytometry and scRNA-seq analyses.

Maternal vascular malperfusion in UTx placentae

We next considered the potential physiologic consequences of the reduction of NK cells

on UTx recipients. We speculated that placentation would be altered in UTx given that

spiral artery remodeling is impaired in mice with dysfunctional or deficient NK cells (*12,*

13, 16, 24), and inadequate spiral artery remodeling is associated with maternal vascular

malperfusion (MVM) in humans (*8, 23, 26*). To test the hypothesis that placentation would

be abnormal in UTx recipients, we examined placental histology in UTx recipients after

delivery. Both placentas were small for gestational age at 236g (UTxA; 311-464g

expected) and 223g (UTxB; 331-491g expected). H&E examination revealed patchy

regions of increased syncytial knots and slender, sparse villi with widening of the

intervillous space consistent with distal villous hypoplasia and accelerated villous

maturation (Fig. 3A), a histologic phenotype indicative of MVM (*25, 51-53*). Notably, these

histologic findings were identified in both UTx placentae but not HC (Fig. 3B). These

histologic changes were associated with abnormal clinical courses in both UTx recipients.

Both UTx recipients were delivered at ~34 weeks gestation due to pregnancy

complications: UTx "A" developed preterm premature rupture of membranes without

evidence of acute chorioamnionitis, and UTx "B" developed pre-eclampsia with elevated

blood pressure and serum creatinine. Notably, both clinical phenotypes have been

associated with MVM (*51, 52*). Although more data are needed given the small number

of UTx recipients in this study, these data align with reproductive phenotypes in NK

deficient mice and suggest potential physiologic consequences for humans with reduced

NK cell number.

Reduction of tissue resident NK subsets in uterus transplant endometrium

To investigate the mechanism of NK loss in UTx recipients, we first determined whether

the NK cell reduction in UTx affected all uterine NK subsets equally. Consistent with prior

studies (*54-56*), we identified two major populations of endometrial NK cells which could

be distinguished by differential expression of *NCAM1* and *FCGR3A* (Fig. 1A and 4A) as

well as transcriptional programs governing tissue residency and cytotoxic effector function

(Fig. 4B and fig. S3). Tissue resident NK cells (trNK) expressed integrins and transcription

factors frequently associated with tissue resident lymphocytes (i.e. *ITGA1, ITGAE, KLF6*)

(Fig. 4B and fig. S3), whereas cNK expressed a variety of molecules associated with

circulating populations, including *S1PR5* and *KLF2* (Fig. 4B and fig. S3) (*30-33, 57, 58*).

Moreover, the endometrial tissue resident cluster expressed a gene signature associated

with tissue residency in human lung NK cells (Fig. 4C and table S2 (*31*). Whereas cNK

number and frequency were similar in UTx vs. HCs, trNK number and frequency were

significantly reduced in UTx (Fig. 4, D and E). Because uterine trNKs are transcriptionally

and functionally heterogeneous (*56, 59, 60*), we next determined if all trNK subsets were

reduced in UTx or whether the trNK defect was limited to specific trNK subsets. After

identification of canonical trNK subsets by expression of decidual NK signatures (Fig. 1B)

(*60*) and marker genes (Fig. 4F and table S2), we compared the frequency of individual

trNK subsets between UTx and HCs. While HC and UTx had similar frequencies of trNK1

cells, we found statistically significant reductions in trNK2 and trNK3 frequency in UTx

recipients (Fig. 4G). As tissue resident lymphocytes express adhesion molecules that

promote tissue retention, we considered the possibility that tissue resident subsets may

not be retrieved equally from endometrial tissue from HC vs. UTx. Because uterine trNK

subsets in human decidua have been previously defined transcriptionally (*60*), we

validated our scRNA-seq data by performing spatial transcriptomics on one healthy

control biopsy and one uterus transplant biopsy. After deconvolution of the spatial

transcriptomics data using cell2location (*61*) (fig. S4), we found reductions in all trNK

subsets with relative preservation of cNK abundance in the UTx vs. HC (Fig. 4H). Taken

altogether, these data suggest that the NK reduction in UTx endometrium is driven

primarily by a loss of tissue resident NK cells.

Altered transcriptional programming of canonical tissue residency in trNK from

UTx

Having established that trNK cells were reduced in UTx compared to HCs, we aimed to

understand the molecular mechanisms responsible for this reduction. To test the

hypothesis that transcriptional programs of trNK development were altered in UTx trNK,

we performed CITE-seq analysis on CD45+ cells aggregated from three HC and four UTx

endometrial biopsies (table S1). As CITE-seq analysis of uNKs has not yet been reported,

we first characterized uNK subsets with this methodology. NK cells were identified among

CD45+ endometrial immune cells based on their gene expression as well as protein

surface phenotype (i.e., CD56+CD14-CD3-) and were re-clustered for downstream

analyses. As in our prior analyses, NK cells resolved into two major groups of cNK and

trNK subsets which were largely distinguished by differential expression of genes

associated with cytotoxicity (fig. S5A) as well as reciprocal expression of CD49a (*ITGA1*)

and CD16 (*FCGR3A*) (Fig. 5, A and B, and fig. S5, A and B). cNK and trNK subsets were

further distinguished by variable expression of CD56, CD45RA, and KLRG1 among other

genes and proteins (fig. S5, A to C). While CD49a- NK cells were composed mostly of

CD56^{dim}CD16+ cNKs (fig. S5B), CD49a+ cells contained multiple distinct subsets,

including trNK1, trNK2, and trNK3 as well as proliferating cells (Fig. 5C and fig. S5, D and

E). Our analysis leveraging both cell surface protein expression and single-cell

transcriptomics further distinguished multiple trNK2 and trNK3 subsets which shared core

trNK signatures (fig. S5F) yet varied from one another based on expression of effector

molecules (i.e., *GNLY, GZMA*), chemokine ligands (i.e., *CCL3, CCL4, CCL5*) and

integrins (*ITGAX, ITGAD*) (fig. S5D).

Next, we compared endometrial trNK subsets among HC and UTx samples. As in our

prior analyses, we noted that tissue resident subsets were less frequent among UTx

CD56+ NK cells compared to healthy controls (Fig. 5D). Moreover, CD103+ trNK3

subsets (C0&C5) were preferentially and markedly reduced in UTx (Fig. 5D), resulting in

an overall skewing of the trNK compartment toward trNK1 cells (Fig. 5D). Given the critical

role of TGFβ in CD103+ tissue resident lymphocyte development (*30, 58, 62-64*), we

hypothesized that TGFβ target genes were downregulated in trNK3 cells from UTx vs.

HCs. Indeed, the TGFβ gene signature was downregulated in UTx trNK3 subsets (Fig.

5E), including the integrins *ITGA1* and *ITGAE*, as well as *CD9* and *ITGB7* (Fig. 5F).

Accordingly, the surface expression of the corresponding proteins was also reduced in

UTx trNK3 subsets (Fig. 5G), suggesting that cells which possessed trNK3

transcriptomes in UTx had an overall shift in their surface proteome toward trNK2 subsets

(Fig. 5H). Collectively, these data suggested that disruption of TGFβ-mediated tissue

residency programming in UTx samples resulted in an overall reduction of trNK3 subsets

with transcriptional skewing toward trNK1 cells. Notably, changes in cellular

transcriptomes corresponded to changes in surface phenotype, which was independently

confirmed by flow cytometry (Fig. 5I).

Reduced expression of early trNK developmental program in UTx

Although the contribution of TGFβ to tissue residency programming is well established

(*30, 58, 62-64*), less is known about the temporal orchestration of transcriptional

programs of residency as lymphoid cells seed tissues from the blood. Importantly, recent

work in a mouse model of LCMV infection has suggested that an array of immediate early

response genes (i.e., *JUN*, etc.) are not only expressed early after tissue entry but are

required for tissue residence (*65*). Moreover, prior work investigating human uterine NK

development has suggested that a similar transcriptional program of early tissue

adaptation is upregulated in early endometrial immigrants after arrival from the blood (*56*).

We thus tested the hypothesis that trNKs were reduced in UTx due to defective

expression of this early developmental program (EDP) that is conserved between species

and cell types (table S2).

First, we examined expression of the tissue resident EDP in healthy control trNK and cNK

subsets and found highest expression of the EDP in certain trNK clusters (i.e., C5, C6,

C7) (Fig. 6A). Although expression of the tissue resident EDP varied across trNK subsets

in healthy controls (Fig. 6, A&B), we found a statistically significant reduction of

expression of the EDP signature in virtually all UTx trNK clusters (Fig. 6B). Further

examination of the most highly expressed EDP genes revealed two gene modules which

were reciprocally expressed among trNK1 and trNK3 subsets (Fig. 6C). trNK2 subsets

had intermediate expression of both gene modules, suggesting an intermediate

transcriptional state between trNK1 and trNK3 cells, in line with prior results (*56*). Notably,

trNK2 and trNK3 subsets were distinguished from trNK1s in part by increased expression

of the developmental module composed of AP-1 transcription factors (i.e., developmental

module 2) (Fig. 6D), a feature we dissected further after grouping similar trNK subsets

(Fig. 6D, left). These analyses revealed that the developmental module encompassing

AP-1 transcription factors (i.e. developmental module 2) was significantly less expressed

in UTx trNKs than in HCs (Fig. 6D). In contrast, the developmental module which

predominated in trNK1 cells (i.e., developmental module 1) had much more modest

reductions in expression in UTx vs. HC (Fig. 6D). Collectively, these results demonstrate

that the early core developmental programming of tissue residency is impaired in UTx

endometrial NK cells, with larger reductions in trNK2 and trNK3 cells which had higher

expression of AP-1 based developmental tissue residency programs.

Role for NFAT in tissue residency programming in uterine NK cells

We next considered the mechanism by which transcriptional programming of uterine

natural killer cell tissue residency was disrupted in UTx. Because NFAT regulates CD103

gene expression (*66*) and NFAT-dependent genes have been identified within prior

studies of tissue resident lymphocytes (*56, 67*), we hypothesized that tissue residency

programming was reduced in UTx due to exposure of these individuals to

immunosuppression drugs which prevent NFAT nuclear translocation (i.e., calcineurin

inhibitors; tacrolimus). We tested for overlap of a curated gene set of NFAT targets (table

S2) with genes upregulated during mucosal CD8 T_{RM} differentiation (*65*) and found 42

genes governing lymphocyte tissue residency that could be influenced by NFAT (Fig. 7A).

Having identified NFAT-dependent genes involved in the transcriptional programming of

tissue residency (i.e., NFAT^{TR} signature), we then tested for enrichment of the NFAT^{TR}

signature in endometrial trNK from healthy control volunteers and found statistically

significant enrichment of multiple NFAT^{TR} genes in human endometrial trNK compared to

cNK cells (Fig. 7, B and C). We next determined whether the NFAT^{TR} signature was

differentially expressed in endometrial NK cells from uterus transplant recipients versus

healthy controls. We compared UTx with HC NK cells within each major trNK cluster and

then tested for enrichment of the NFAT^{TR} signature among the resulting differentially

expressed genes. These analyses revealed enrichment of the NFAT^{TR} signature among

genes downregulated in UTx compared to HCs in trNK subsets reduced in UTx (i.e., trNK2

and trNK3 subsets) (Fig. 7D). We noted strong conservation of NFAT^{TR} genes that were

downregulated in affected UTx subsets (Fig. 7D). Notably, the NFAT^{TR} genes

downregulated in affected UTx subsets corresponded to genes shared in the

developmental program most expressed in trNK2 and trNK3 subsets (Fig. 7, E and F).

Collectively, these data suggest a role for NFAT in tissue residency programming of

endometrial trNK cells and identify reductions in trNK subsets with highest expression of

transcriptional developmental programs that are enriched for NFAT-dependent genes.

IL-15 promotes integrin expression in endometrial trNK cells through NFAT

Although NFAT has well-established roles conveying signals downstream of antigen

receptors in B and T lymphocytes (*68*), less is known about how innate cells such as NK

cells use NFAT to translate environmental inputs. Nevertheless, a small number of prior

studies have identified NFAT effects downstream of cytokine receptor signals such as IL-

6 (*69*) and IL-15 signals (*70*). Since IL-15 synergizes with TGFβ in NK cells to promote

upregulation of integrins important in tissue residency (*71*), we hypothesized that NFAT

was contributing to tissue residency programming of trNK cells downstream of IL-15

signals. To test this hypothesis, we tested whether NFAT blockade by FK506 would result

in impairment of key integrins on endometrial NK cells in response to IL-15 stimulation.

IL-15 stimulation in vitro for 4-7 days was sufficient to induce significant upregulation of

CD103 and CD49a expression on endometrial NK cells without the addition of exogenous

TGFβ (Fig. 8A). However, the addition of FK506 to these cultures significantly diminished

the frequency of CD103+CD49a+ endometrial trNK cells, with concomitant increases in

the frequencies of CD103-CD49a+ and CD103-CD49a- cells (Fig. 8B). Altogether, these

in vitro studies suggest that cytokine signals which mediate tissue residency programs in

endometrial trNK utilize NFAT.

DISCUSSION

Significant knowledge gaps remain regarding our understanding of the immune

pathogenesis of pregnancy complications in humans. Although mouse models have firmly

established the role played by uNKs in spiral artery remodeling and placentation (*7, 12,*

13, 16, 24), pregnancy outcomes in women with reductions in endometrial NK cells have

not been described. In this work, we identify uterus transplant recipients as an at-risk

cohort who possess a reduced and skewed complement of tissue resident NK cells.

Reductions in trNK number in longitudinal endometrial biopsies collected prior to

pregnancy in UTx recipients were associated with pregnancy complications and histologic

placental abnormalities characteristic of maternal vascular malperfusion. Analysis of trNK

transcriptional and proteomic signatures in UTx samples not only revealed reduced

expression of individual integrins critical for lymphocyte residency in mucosal tissues (*30-*

33, 57, 58, 62-65, 72), but also revealed reductions in NFAT-dependent gene modules

that govern early trNK development (*65*). Collectively, these data suggest that UTx

recipients may recapitulate reproductive phenotypes observed in mice with inadequate

NK cell number or function, thus deepening our understanding of the pathogenesis of

pregnancy complications in humans.

Our work contributes new insights to the molecular events which guide the development

of tissue resident natural killer cells. Although NFAT is known to bind the *ITGAE* enhancer

(*66*) and contributes to CD103 expression in CD8 T cells after TCR stimulation (*73*), a

role for NFAT in regulating CD103 expression in NK cells has not yet been described. We

hypothesized that in the absence of an antigen receptor, NFAT would promote CD103

expression in uterine NK cells downstream of a cytokine signal, as has been described

previously for IL-6 (*69*) and IL-15 (*70*). In this work, we tested the hypothesis that IL-15

could promote CD103 expression through NFAT-dependent mechanisms given that IL-

15 is secreted by decidualizing stromal cells in the secretory phase (*74, 75*). Moreover,

IL-15 is known to synergize with TGFβ to induce CD103 expression on peripheral blood

NK cells through MAPK pathways (*71*). Although NFAT is indeed a target of MAPK in T

cells (*76*), it is unclear what intracellular molecules are promoting NFAT-dependent

CD103 expression in endometrial trNK3 cells, as NFAT may be acting downstream of IL-

15-mediated PI3K activation as well. Additional studies will be needed to elucidate the

molecular details of NFAT signaling on CD103 expression by trNK3 cells, but this work

will likely be highly impactful given the severity of the trNK3 defect we observed in UTx

endometrium.

While the sensitivity of CD103-expressing trNK3 cells to NFAT inhibition may largely be

explained by the role of NFAT in *ITGAE* gene expression, the reduced frequency of

CD103- trNK2 cells in UTx endometrium required further explanation. As our

understanding of tissue resident immune cells has expanded well beyond the adhesome

in recent years, we considered the possibility that other molecular subroutines of tissue

residency were impacted by NFAT. Unexpectedly, we found significant overlap between

genes upregulated early in tissue residency with a curated list of NFAT targets,

suggesting a potential mechanism for loss of CD103- subsets in UTx endometrium.

Nevertheless, not all trNK subsets equally expressed the NFAT-dependent early tissue

residency program, thus providing an explanation for the relative sparing of the trNK1

subset while trNK2 and trNK3 cells were reduced. Precisely how NFAT promotes

expression of the early residency signature in trNK2 and trNK3 cells remains to be

investigated, and in pursuit of this question, the NFAT-independence of trNK1 subsets

may also be deciphered. We anticipate that additional clues will become available as our

understanding of the specific environmental signals which propel trNK1 differentiation is

improved over time.

Reduction of trNK subsets in uterus transplant recipients in our study cohort was

associated with placental pathology and clinical syndromes associated with aberrant

spiral artery remodeling. Taken together, these findings suggest that there are potentially

significant consequences to disruption of tissue residency programming of NK cells on

pregnancy outcomes in humans. Nevertheless, robust conclusions based on these

associations are limited by the small sample size of our UTx cohort and the absence of

decidual studies to assess the magnitude of spiral artery transformation. Such structural

studies on future UTx recipients will be necessary to fully interpret our findings and place

this work in a larger context. Moreover, while we attributed the NK reduction observed in

this work to aberrant residency programming in the setting of NFAT inhibition, UTx

recipients are on a variety of immunosuppression agents whose effects on trNK

recruitment, survival, differentiation, or function have yet to be tested. Additional studies

will be needed to discriminate between these possibilities and determine how these

factors play a role in endometrial immune homeostasis. Finally, our work does not

preclude a role for uterine parenchymal dysfunction or altered decidualization in UTx

recipients due to FK506 effects. Thus, expansion of our study to diverse cell types in other

organ transplant recipients will be necessary to complete our understanding of pre-

eclampsia pathogenesis in this setting. Additional examination of the NFAT axis in trNKs

in other at risk populations will be key to determine the potential benefit of therapeutic

modulation of this pathway.

MATERIALS AND METHODS

Study Design

Research objective:

To test the hypothesis that uterine NK cells are abnormal in a recipient population at

elevated risk for pregnancy complications.

Research location: All aspects of the study (i.e., subject recruitment, sample collection,

and sample analysis) were performed at the University of Alabama at Birmingham (UAB).

Research subjects:

Healthy control (HC) participants were screened for inclusion and exclusion criteria ahead

of their gynecology clinic visits at UAB. Once eligibility was confirmed, participants were

approached for written informed consent for study participation. Inclusion criteria

consisted of reproductive age, uncomplicated pregnancy history, normal menstrual

cycles, and absence of serious medical illness. Exclusion criteria included any history of

the following: current pregnancy, malignancy, diabetes, pharmacologic immune

modulation, systemic autoimmune disease, prior organ or bone marrow transplant, history

of chemotherapy or radiation treatment, chronic or end-stage organ disease, history of

HIV/HBV/HCV infection, instrumentation of the uterine cavity (i.e., D&C) within prior 12

months, use of an intrauterine device in prior 3 months, oral contraceptive use in prior 3

months, pregnancy within prior 12 months, NSAID or aspirin use within the prior 10 days,

and/or medical contraindication to NSAIDs. CMV serostatus was not determined for

healthy control participants. Healthy control participants underwent HCG testing in the

clinical space prior to endometrial biopsy to document a negative pregnancy test.

<u>Uterus transplantation</u> is offered as standard of care treatment outside of a clinical trial at

UAB for XX genotype individuals with absolute uterine factor infertility who meet selection

criteria. Uterus transplant recipients were screened for participation in the endometrial

immunobiology study during clinic visits at UAB. Once eligibility was confirmed,

participants were approached for written informed consent for study participation.

Endometrial biopsies were collected from participants >30 days post-transplant with

normal menstrual cycles and no evidence of infection. Both participants were Caucasian

individuals of reproductive age who met transplant center criteria; specific ages are not

reported given the small nature of the patient cohort. Etiology of uterine factor infertility

included hysterectomy as well as congenital absence of the uterus. PBMCs were collected

one day prior to transplantation for whole exome sequencing. Both participating transplant

recipients received a blood group compatible transplant from a deceased donor of

reproductive age. Transplant recipients received induction immunosuppression with anti-

thymocyte globulin (6mg/kg) and 500mg of intravenous methylprednisolone at the time of

transplantation. Maintenance immunosuppression consisted of tacrolimus, azathioprine,

and prednisone. Target levels of tacrolimus were 5-10 ng/mL and were tested weekly.

Secretory phase endometrial biopsies were obtained while protocol cervical biopsies were

performed for immunologic monitoring of the graft. No endometrial biopsy was included in

the study from a patient treated for rejection based on cervical biopsy findings. Additional

medications taken during the first six months of transplantation included daily

sulfamethoxazole/trimethoprim (for PCP prophylaxis) as well as acyclovir (UTxB) and

valganciclovir (UTxA). Acyclovir was taken for HSV prophylaxis by UTxB given that both

the recipient and the donor were seronegative for cytomegalovirus (CMV). UTxA was

prescribed valganciclovir as she and her donor were both CMV seropositive. All

endometrial biopsies from UTx recipients were obtained >30 days after transplantation

after resumption of normal menstrual cycles but prior to embryo transfer (table S1).

Placentas were examined immediately after cesarean delivery from both participants.

Slides from a <u>healthy control archived placenta</u> that was matched for gestational age (~34

weeks) with UTx deliveries were retrieved from an internal biorepository and evaluated by

a perinatal pathologist (V.E.D.).

<u>Brain-dead deceased donors</u> provided 1) blood for whole exome sequencing, and 2)

endometrium for in vitro co-culture experiments. Deceased donors were identified by

Legacy of Hope personnel for the purpose of organ transplantation and/or research, and

families of the deceased donors provided authorization. PBMCs were collected from

deceased donors (n=2) prior to uterus procurement for the purposes of transplantation in

order to perform donor genotyping through whole exome sequencing. Endometrium for in

vitro co-culture experiments was obtained from deceased donor hysterectomies (n=4) that

were performed during surgical training procurements but were not transplanted. Data on

the menstrual cycle phase for the four deceased donor endometrial samples was

unavailable.

Experimental Design

Design Synopsis: Human endometrium was sampled from 1) healthy control participants

(secretory phase), 2) uterus transplant recipients (secretory phase), and 3) brain-dead

deceased donors (unknown menstrual cycle phase) for this study. Endometrial samples

were analyzed with single cell approaches (i.e., scRNA-seq, CITE-seq, flow cytometry)

as well as imaging (i.e., spatial transcriptomics, immunofluorescence microscopy). Single

cell suspensions of endometrium were cultured with IL-15 and FK506 for 3-7 days.

Rigor & reproducibility considerations

Healthy control and uterus transplant participants used an ovulation predictor kit

(Clearblue® digital ovulation tests (SPD Swiss Precision Diagnostics GmbH; Item number

245-03-0301)) to objectively identify the secretory phase of the menstrual cycle.

Menstrual cycle dating was therefore not dependent on subject reporting based on the

start date of menses. Of note, the digital ovulation predictor kit turns positive when

luteinizing hormone (LH) is detected in the urine. Participants were brought into the clinic

to undergo endometrial biopsy in the days following the LH surge, and biopsy data are

thus reported as "LH+7" to denote menstrual cycle day after the LH surge. Variability in

the uterus transplant data were minimized by collecting biopsies >30 days after

transplantation only from individuals who had resumed normal menstrual cycles.

Use of Biologic and Technical Replicates: For single-cell RNA-seq experiments, six

secretory phase endometrial biopsies were obtained from six independent individuals (for

healthy controls), and five secretory phase endometrial biopsies were obtained from two

independent individuals who had received a uterus transplant (table S1). Endometrial

biopsies were initially analyzed separately and then analyzed together after Harmony

integration to control batch effects. For ex vivo flow cytometric analyses, three secretory

phase endometrial biopsies were obtained from two independent individuals, and 16

secretory phase endometrial biopsies were obtained from healthy control volunteers. For

in vitro co-culture experiments, four secretory phase endometrial biopsies were obtained

from healthy control volunteers and four endometrial biopsies were obtained from

deceased donor hysterectomies (unknown menstrual cycle date). In vitro co-culture

experiments were performed in duplicate or triplicate. For spatial transcriptomics

analyses, two sections were performed for each participant (one healthy control volunteer

and one uterus transplant recipient). Results from one section per participant are reported

in the manuscript but are representative of both sections. For immunofluorescence

microscopy analyses, secretory phase biopsies were obtained from two individual healthy

control volunteers, and three secretory phase biopsies were obtained from two individual

uterus transplant recipients. Although sample number was limited in this study, the

entirety of the tissue was assessed with multiple sections evaluated by an automated

approach with ~10% error. Data from all individual biologic replicates are provided in the

visualizations.

Regulatory considerations:

The University of Alabama at Birmingham Institutional Review Board reviewed and

approved sample collection from healthy control participants under project "Mechanisms

of Uterine NK Cell Differentiation" (IRB-300006859) and sample collection from uterus

transplant recipients under project "Endometrial Immunobiology in Uterus

Transplantation" (IRB-3000007980). All samples were de-identified.

Sample collection & processing

Endometrial biopsies

Secretory phase endometrial biopsies were collected with a Pipelle® suction curette

(Cooper Surgical; Ref. 8200) in the clinical setting and transported to the laboratory in

DPBS with calcium and magnesium (DPBS$^{+/+}$; Gibco™; Cat. 14-040-133) on ice. Multiple

passes of the suction curette were performed to acquire sufficient tissue for multimodal

analysis with processing described below. Endometrial biopsies were then digested to

isolate cells for downstream analysis including sorting, flow cytometry, or in vitro

incubation.

Endometrial biopsies used for imaging analyses, including spatial transcriptomics and

immunofluorescence microscopy, were frozen according to 10x Genomics CG000240

Demonstrated Protocol for Visium Spatial Tissue Preparation Guide. The biopsy pieces

were embedded in O.C.T. mounting media (Tissue-Tek; Cat. No. 4583) in a mold (Tissue-

Tek; Cat. No. 4566). Molds were then placed in liquid nitrogen-cooled isopentane (Fisher

Scientific; Cat. No. O3551-4). Frozen molds were stored at -80°C until sectioned onto

Visium slides or glass slides for immunofluorescence microscopy.

<u>Endometrial biopsies used for single cell isolation</u> (i.e., scRNA-seq, flow cytometric

analysis, in vitro co-culture) were minced with a razor blade on ice and transferred to a 50

mL conical tube on ice. The biopsy had 4 mL of 37°C-warmed digestion buffer added

(DPBS$^{+/+}$ + 40 mL 5 mg/mL Liberase (Sigma Aldrich; SKU 5401127001) + 4 mL 1 mg/mL

DNAse I (Sigma-Aldrich; SKU 11284932001)). Digestion was performed in a 37°C shaking

incubator at 240 RPMs for 20 minutes. The digested sample was filtered through a 70 μm

strainer (Stemcell Technologies; Cat. 27260). The strainer was rinsed 5 times with 1 mL

ice-cold DPBS$^{+/+}$. The digested biopsy was then centrifuged at 300xg for 5 min at 4°C.

The cell pellet was incubated in 3 mL ACK lysis buffer (Quality Biological; Cat. 118-156-

101) on ice for 2 minutes. After ACK lysis, 12 mL ice-cold DPBS without calcium and

magnesium (DPBS$^{-/-}$; Gibco™; Cat. 14-287-080) was added to the cell suspension. The

cells were then centrifuged at 300xg for 5 min at 4°C. The buffer was removed and 0.5-

1mL fresh sort buffe (ice-cold DPBS$^{-/-}$ + 0.04% (w/v) BSA (Jackson ImmunoResearch;

Cat. 001-000-162)), was added. The cells were then suspended and counted. All cells

were counted by the following method: 10 μL of cell suspension from samples was added

to 90 μL 0.04% Trypan blue in PBS (Gibco™; Ref. 15250-061) for a 1:10 dilution, or

serially diluted to make a 1:100 dilution. Cell counts were performed using a disposable

Neubauer improved hemocytometer (INCYTO; DHC-N01-5), following the manufacturer's

instructions for loading.

Placental processing

Placentas were recovered from the operating room at time of delivery and placed in a

sterile plastic container for transport to the clinical surgical pathology laboratory. Placentas

were processed according to the Amsterdam Criteria (Amsterdam Placental Workshop

Group Consensus professional guidelines) and evaluated by an experienced perinatal

pathologist (V.E.D.). In brief, the placental disk was sequentially sectioned and systematic

random samples of the chorionic plate, the mid-zone, and the basal plate were collected

and placed in 10% formalin. Samples were subsequently paraffin embedded, stained with

hematoxylin and eosin, and sectioned at 5μm for histopathologic analysis.

PBMC processing

PBMCs were isolated to perform whole exome sequencing (WES). Blood was collected

in EDTA-K2 tubes vacutainers (BD and Company; Ref. 366643) and blood collection set

(MYCO Medical Supplies, Inc.; Ref. GSBCS23G-7T). PBMC isolation was performed at

room temperature until red blood cell (RBC) lysis. Equal volumes of wash media (DPBS^{-}

$^{/-}$ + 2% (v/v) FBS (Gemini Bio; Ref. 100106)), were added to the whole blood and mixed.

The manufacturer's instructions were followed to isolate PBMCs using Lymproprep^{TM}

(Stemcell Technologies; Ref. 07811) and either a Sepmate™-15 or -50 (Stemcell; Ref.

85420 or 85450), depending on the blood volume. Once the PBMCs were isolated, the

cells were washed with the DPBS$^{-/-}$ and 2% FBS two times, centrifuged once at 300xg for

8 minutes and once at 120xg for 10 minutes with no break. After removing the final wash

buffer, RBCs were lysed using 3 mL of room temperature ACK lysis buffer (Quality

Biological; Cat. 118-156-101) for 2 minutes on ice. Ice-cold DPBS was added to the 14

mL mark and mixed to finish the lysis reaction. Cells were pelleted by centrifugation,

400xg for 5 minutes at 4°C. The buffer was removed and 0.5-1mL DPBS$^{-/-}$ was added;

the cells were suspended and counted. PBMCs were cryopreserved until thawed for DNA

isolation and subsequent whole exome sequencing.

Deceased donor hysterectomies

Surgical hysterectomy for the purposes of research was performed in the Donor Recovery

Center at the University of Alabama at Birmingham from deceased donors after family

authorization for research was obtained. Deceased donor uteri were recovered after

donor cross clamp and removal of other organs for the purposes of transplantation. Uteri

were transported to the laboratory on ice and bivalved to expose the endometrium. A

scalpel blade was used to gently scrape the endometrium into a culture dish, and the

sample was washed and digested as above for flow cytometric and/or in vitro co-culture

experiments.

Flow cytometric analysis & cell sorting

Live, CD45+ immune cells were used for scRNA sequencing and were prepared for

FACS-sorting from endometrium as follows: Cells were first stained using the 150 μL

1:600 LIVE/DEAD™ Fixable Aqua stain kit (Molecular Probes™; Cat. L34957) diluted in

DPBS$^{+/+}$ on ice for 30 minutes. After the incubation, cells were centrifuged at 300xg for 5

minutes at 4°C, and the buffer aspirated. Cells were then washed in staining buffer

(DPBS$^{+/+}$ + 2% FBS + 0.4% (v/v) + 0.5 M EDTA (Fisher Scientific; Cat. No. AAJ15694AP)).

Cells were pelleted and then resuspended in staining buffer and stained for 30 minutes

on ice with 1 mg/mL AF700-mouse-anti-human CD45 (BioLegend; 368508) and PE-

mouse anti-human CD235a in staining buffer. After antibody staining, cells were

centrifuged at 300xg for 5 minutes at 4°C and the staining buffer aspirated. Cells were

washed in 1 mL ice-cold sorting buffer and then centrifuged at 300xg for 5 minutes at 4°C.

The wash buffer was then removed, and the cells were resuspended in ice-cold sorting

buffer at a concentration of 2-7 million cells/mL. Live CD45+ CD235a- cells were then

sorted on a FACSAria I or II (BD Biosciences) at the UAB Flow Cytometry Core Facility

in tubes with 5 mL sorting buffer in the bottom. Cells were counted again to determine cell

number, concentration, and viability.

Cells prepared for flow cytometric analysis were aliquoted to have between 200,000 –

10,000,000 cells per tube. Cells were stained first in 150 µL LIVE/DEAD™ Fixable Aqua

using the staining kit described above. Cells were then resuspended in 50 mL staining

buffer on ice, which included 30 mL of BD Brilliant Stain Buffer (BD Biosciences; Cat. No.

563794) and 1 mL 400 mg/mL Pacific Blue-mouse-anti-human CD45 (BioLegend; Cat.

No. 304029; clone HI30), 1 mL 50 mg/mL PE/Dazzle594-mouse-anti-human CD16

(BioLegend; Cat. No. 302054; clone 3G8), 1 mL 100 mg/mL PE/Cy7-mouse-anti-human

CD39 (BioLegend; Cat. No. 328212; clone A1), 1 mL 50 mg/mL APC-mouse-anti-human

CD56 (BioLegend; Cat. No. 318310; clone HCD56), 0.5 mL 200 mg/mL APC/Cy7-mouse-

anti-human CD3 (BioLegend; Cat. No. 317342; clone OKT3), 2 mL 200 mg/mL BV711-

mouse-anti-human CD49a (BD Biosciences; Cat. No. 742361; clone SR84), and 1 mL 70

mg/mL BV421-mouse-anti-human CD103 (Biolegend; Cat. No. 350214; clone Ber-ACT8).

After the incubation, cells were centrifuged at 300xg for 5 minutes at 4°C and the buffer

aspirated. Cells were then washed in 1mL ice-cold staining buffer. The cells were

centrifuged at 300xg for 5 minutes at 4°C and the buffer aspirated. Cells were

resuspended in DPBS for flow cytometric analysis on a Cytek Northern Lights spectral

cytometer.

In vitro culture

Single-cell suspensions of endometrium were resuspended in complete media and plated

in 24 well plates with 10 ng/mL IL-15 ± 10 ng/mL FK506. 5×10^5 – 1×10^6 cells were plated

in duplicate or triplicate and cultured in a 5% CO_2 incubator for 3-7 days. Culture media

was replaced with fresh IL-15 ± FK506 at day three for 7 day cultures. Upon culture

completion, cells were recovered from the plate and subjected to flow cytometry analysis

as described above.

Whole exome sequencing

PBMCs used for whole exome sequencing (WES) were collected from uterus donors and

recipients prior to uterus transplantation. PBMCs were cryopreserved, as described

above, and then thawed for WES. Cryopreserved cells were partially thawed in a 37°C

water bath while gently swirling. When half of the volume of the cells were thawed, 500

mL of 37°C-warmed thawing media, RPMI with 10% FBS, was added to the cryovials.

Suspended cells were transferred to a 15 mL conical tube with 9 mL of warmed thawing

media. Cells were centrifuged at 400xg for 5 minutes at 4°C. The thaw media was

aspirated, and cells were washed again in thaw media. After centrifugation as before,

thaw media was aspirated, and cells were suspended in DPBS$^{-/-}$ and counted.

Genomic DNA was then isolated from 1-5 million isolated PBMCs using the DNeasy Blood

and Tissue kit (Qiagen; Cat. No. 69504) following manufacturer instructions. WES libraries

were prepared using the SureSelect Human All Exon V8 Kit (Agilent; 5191-6873) and

sequenced on either a NextSeq 2000 (Illumina) or NovaSeq 6000 (Illumina).

scRNA-seq and CITE-seq library preparation

Live CD45+ cells were used for scRNA sequencing after being FACS-enriched as

described above. Cells that were used in CITE-seq experiments were blocked with

TruStain Fc block (BioLegend; Cat. 422301) for 30 minutes. Cells were then incubated

with TotalSeq B™ Universal cocktail (BioLegend; Cat. 399904) in Cell Staining Buffer

(Biolegend; Cat. 420201) and washed following the manufacturer's protocol before being

used for scRNA library preparation.

Cells isolated for scRNA/CITE-seq were suspended in ice-cold cell sorting buffer. scRNA

GEMs and libraries were prepared according to the manufacturer's instructions. 5'

libraries were prepared according to 10x Genomics CG000331 protocol, using Chromium

Next GEM Single Cell 5' Kit v2 (10x Genomics; PN-1000263). 3' libraries were prepared

using 10x Genomics CG000399 protocol, or CG000206 if a CITE-seq antibody panel was

used with the appropriate 3' GEM kit (10x Genomics; PN 1000121). Both 3' and 5' libraries

used Dual Index Kit TT Set A (10x Genomics; PN 3000431). GEMs were created on either

a Chromium Controller or Chromium X (10x Genomics). Either C1000 (Bio-rad) or

SimpliAmp (Applied Biosystems) thermocyclers were used for library prep. Quality and

concentrations of cDNA and final libraries were calculated from 2100 BioAnalyzer

(Agilent) electrograms. Sequencing was performed on a NovaSeq 6000 (Ilumina).

Spatial transcriptomics

Biopsy tissues used for spatial transcriptomics were sectioned on a cryostat at 10 μm

thickness and placed on Visium slides (10x Genomics; CG000239). Visium 3' libraries

were prepared according to 10x Genomics Protocol CG000239 Rev G using a 30-minute

incubation for tissue permeabilization. Either a C1000 (Bio-rad) or SimpliAmp (Applied

Biosystems) thermocycler was used for library preparation. KAPA SYBR FAST qPCR

Master Mix (2X) (Millipore Sigma; KK4600) and a QuantStudio 7 Pro qPCR (Applied

Biosystems) were used to quantify cDNA following manufacturer recommendations. The

Dual Index TT Set A (10x Genomics; PN 3000431) was used for the final library

preparation and a 2100 Bioanalyzer (Agilent) was used to perform library quality control

and concentration quantification. Sequencing was performed on a Novaseq 6000

(Ilumina).

Data analysis

Flow cytometry analysis: All data were analyzed with FlowJo^{TM} software (v.10.8.1)

(Becton, Dickinson, and Company; Ashland, OR, USA). Statistical analysis of flow data

was performed with "stats" package in R (v.4.2.2) after importing into R. Functions used

for statistical testing included wilcox.test() and the additional following arguments

"paired=TRUE" or "alternative = "less"" or "alternative = "greater"" for one-sided testing

for unpaired data.

scRNA-Seq data processing and analysis: Reads from single-cell RNA-Seq experiments

of each HC and UTx library were pre-processed and aligned against the GRCh38 human

reference and quantified using Cell Ranger (v.6.1.1) from 10x Genomics. Downstream

analysis of the generated Unique Molecular Identifier (UMI) count data was completed

using R (v.4.2.1) and Seurat (v.4.4.0), utilizing default parameters unless otherwise

specified. Prior to subsequent analyses, background noise from ambient RNA was

removed using SoupX (v.1.6.2). The overall contamination fraction (rho) was

parameterized using the autoEstCont function to remove > 2% background

contamination in each dataset. The SoupX-corrected count matrixes were then loaded

into R using the *Read10X* function from which Seurat objects were created per dataset

via the *CreateSeuratObject* function. From each dataset, cells with less than 200 features,

and features with less than 3 cells expressing them were filtered. Additionally, respective

count matrixes were filtered based on quality control (QC) metrics and thresholds (table

S3 and fig. S6). Doublets arising from the 10X sample loading procedure were detected

and removed using Scrublet (v.0.2.3). The doublet rate per library was approximated

using estimated rates provided by 10x Genomics.

Filtered count matrices were then merged into one Seurat object, normalized, and

variance stabilized using regularized negative binomial regression via the *SCTransform*

package provided by Seurat (*77*). We also regressed out variation introduced by

mitochondrial expression to prevent any confounding signal. Next, identification of

principal components was performed using (RunPCA). Leveraging identified PCs, and

while considering each individual library as a batch, the normalized dataset was corrected

for batch effects and integrated using Harmony (v1.2.0). The resulting top 50 Harmony-

corrected dimensions were further used to construct a k-nearest neighbor (KNN) graph

using *FindNeighbors().* Cell clustering was subsequently performed

using *FindClusters()* which employs a shared nearest neighbor (SNN) modularity

optimization-based clustering algorithm. Resulting cluster information was used as input

into the uniform manifold approximation and projection algorithm *RunUMAP()* which

further aided the visualization of cell manifolds in a low-dimensional space. We ran

Seurat's implementation of the Wilcoxon rank-sum test *FindMarkers()* to identify

differentially expressed genes in each cluster. The expression of cluster-specific

canonical markers was used to annotate each cell cluster.

CITE-Seq data processing and analysis of CD45+ endometrial immune cells

Three healthy control, and four uterine transplant datasets were further used for cell

surface protein and transcriptomic data analyses. Pre-processing of these data was

conducted as described previously. The RNA gene expression data was normalized using

SCTransform (as earlier described), however, the ADT counts were centered log-ratio

(CLR) normalized. Normalized ADT counts were then scaled and centered using the

ScaleData function. Principal Component Analysis (PCA) was then conducted on both

assays (RNA and ADT) using the *RunPCA* function with default parameters. We

leveraged the ElbowPlot and DimHeatmaps to determine the number of principal

components (PCs) to consider for further downstream analyses. Towards cell clustering,

we found the k-nearest neighbors of each cell using the Seurat *FindNeighbors* function

which then constructed a shared nearest neighbor graph using the top 40 and top 20

dimensions for RNA and ADT assays respectively. As the data constitute multiple

modalities, we leveraged Seurat's Version 4.4.0 to define cellular states based on these

two assays. We applied the weighted-nearest neighbor (WNN) analysis (*78*), an

unsupervised framework to integrate the multiple data types measured within each cell,

and to obtain a unified definition of cellular state based on the multiple modalities. For

each cell, a set of modality weights was learned, which reflects the relative information

content for each data type in that cell. This enabled the generation of a WNN graph that

denotes the most similar cells in the dataset based on a weighted combination of protein

and RNA similarities. To achieve this, we used the Seurat *FindMultiModalNeighbors*

function generating cell neighborhood graphs for further downstream analyses. We then

preformed cell clustering on the generated graphs and identified clusters using the

Louvain algorithm (*FindCluster* function) at the default resolution. We then utilized cluster

information as input into the uniform manifold approximation and projection

(*UMAP*) activated by the *RunUMAP* function which further aided the visualization of cell

manifolds in a two-dimensional space, and as a weighted combination of RNA and protein

data.

To identify differentially expressed (DE) genes between groups of cells, we used the

FindAllMarkers function. Results from differential gene expression analysis were used to

annotate clusters based on the expression of top marker genes in the resulting clusters,

and by leveraging feature plots (*FeaturePlot* function) of canonical cell type specific

marker expression.

Module scores

Module scores were calculated using the *AddModuleScore()* function from Seurat.

Respective feature plots were generated using *FeaturePlot()* with the features set as the

calculated module scores from each gene program.

Identification of donor or recipient cellular origin in uterus transplant samples

To determine the identity of cells from uterus transplant recipients, we used sample-

specific genetic mutations and Demuxlet (v.1.0-5) (*79*). Whole-exome sequencing (WES)

was conducted on DNA isolated from donor and recipient PBMC samples collected prior

to transplantation. Sequenced reads were aligned using STAR. Single nucleotide variants

(SNVs) were detected using Freebayes (v.1.1.0-goolf-1.7.20_39e5e4b) for each donor

and recipient sample. We provided a unique identifier per sample using "bcftools

reheader" and filtered the resulting VCF files to retain only common SNPs with minor

allele frequency (MAF) >= 0.05, and SNPs that overlap exons (defined by GENCODE

release 19). The donor and recipient VCF files were then merged and sorted to match the

recipient .BAM file from cellranger using popscle_helper_tools. Reads were piled up using

the popscle dsc-pileup command using default parameters along with the filtered set of

barcodes from cell ranger. The genetic deconvolution of sample identities in the UTx

single-cell libraries was then conducted using Demuxlet (v.1.0-5) from the popscle

package by taking the merged donor and recipient .vcf file and the recipient .BAM file

from Cell Ranger as inputs. Demuxlet was run at default parameters with the GT filed

used to extract the genotype, likelihood, and posterior, and the offset of the genotype

error rate (geno-error-offset) set to 0.05. Each cell in the final integrated dataset was

annotated as donor or recipient using the BEST.GUESS column from the Demuxlet

output data.

Cell2location

Visium spatial gene expression data processing and analysis using cell2location

i. Cell-type deconvolution of spatial transcriptomics data using cell2location (*60*).

To spatially resolve cell populations identified in healthy controls and uterine transplant

biopsies, we used cell2location, a Bayesian model which estimates cell type abundance

by deconvoluting a spatial expression count matrix into a set of reference cell type

signatures. The model takes a spatial expression matrix with mRNA counts of genes at

spatial locations and a matrix of reference cell type signatures as input. All cell2location

analyses were conducted using Scanpy (v. 1.9) (*80*).

ii. Construction of scRNA-seq reference transcriptomes

Reference transcriptomes input into cell2location were derived from the sorted CD45+

immune cells from the six healthy control endometrial biopsies (scRNA-seq). These data

were pre-processed as described above for scRNA-seq. The data were merged into a

unified object using Seurat's merge function yielding a total of 41,719 cells and 20,636

genes in the reference dataset. The merged dataset was then normalized, scaled, and

variable features identified using the SCTransform() method in Seurat. Identification of

principal components was performed using (RunPCA). Leveraging identified PCs, and

while considering each individual library as a batch, the merged dataset was corrected

for batch effects and integrated using Harmony (v1.2.0). Corrected Harmony embeddings

(n=20) were further used to construct a k-nearest neighbor (KNN) graph

using FindNeighbors. Clustering was subsequently performed using FindClusters which

employs a shared nearest neighbor (SNN) modularity optimization-based clustering

algorithm, resulting in 34 distinct reference cell clusters. Cluster information was used as

input into the uniform manifold approximation and projection algorithm (RunUMAP) which

further aided the visualization of cell manifolds in a low-dimensional space. We ran

Seurat's implementation of the Wilcoxon rank-sum test (FindMarkers) to identify

differentially expressed genes in each cluster. The expression of cluster-specific

canonical markers was used to annotate each identified cell cluster (fig. S4).

iii. Estimating reference cell type signatures

The scRNA-seq reference data and corresponding reference cell type annotations were

loaded into Scanpy using sc.read_10x_mtx. Gene selection and general QC were

conducted as described before. Expression signatures of the 34 reference cell types in

the reference dataset were then estimated using the negative binomial regression model

while accounting for batch effects. The created model was further trained to estimate the

reference cell type signatures on all cells in the dataset (train_size=1), mini-batch size =

2,500 and a training duration of 600 epochs, with GPU acceleration. Model accuracy was

evaluated based on the evidence lower bound (ELBO) loss and reconstruction accuracy

plots.

iv. Spatial mapping of endometrial immune cell-types to spatially barcoded HC and UTx

biopsies using cell2location

We imported the Visium spatial transcriptomics data for each biopsy from 10X Space

Ranger into Scanpy, using scanpy.read_visium. Mitochondrial genes were identified and

removed. Before running cell2location to map our reference cell types to each Visium

dataset, both the estimated reference cell type signatures (gene expression), and spatial

data from each respective biopsy were filtered to select genes shared between the two

data objects. The filtered objects were both used as input to cell2location. We specified

two hyperparameters required to run cell2location: 1) The expected cell abundance per

location (N_cells_per_location), which was estimated by manually counting cells in 10

random capture spots from the respective H&E images and averaging these values to

give us the average cell abundance per tissue sample (HC = 14.5 cells/capture spot; UTx

= 17.7 cells/capture spot); and 2) a parameter which accounted for within-experiment

variation in RNA detection sensitivity (detection_alpha). We set detection alpha = 20 in

order to ensure the greatest accuracy and sensitivity given that strong gradients in mRNA

detection sensitivity are commonly observed in adult human 10x Visium data.

After providing the reference cell type signatures, spatial data, and hyperparameters

specified above as inputs, the cell2location model was trained using max_epochs =

30000 on the full dataset (batch_size = None) while estimating cell abundance at all

locations (train_size = 1). We plotted ELBO loss history during training and assessed

mapping quality by examining reconstruction accuracy plots. We exported estimated

posterior distributions of cell abundance ('num_samples': 1000, 'batch_size':

mod.adata.n_obs, 'use_gpu': True) and added a 5% quantile of the posterior distribution,

representing the value of cell abundance corresponding to high confidence in the model.

We further leveraged Scanpy's plotting function scanpy.pl.spatial to visualize spatial

scatter plots of cell type abundance in spatial coordinates.

<u>Spatially barcoded libraries prepared from one healthy control and one uterine transplant</u>

<u>endometrial biopsy.</u>

i. Data pre-processing

Base call (BCL) files were converted to FASTQ files using the Space Ranger

(v.1.3.1) mkfastq function. Space Ranger count function was used to align the FASTQ

files to the human reference genome GRCh38 to generate count matrices.

Gene set enrichment analysis Gene set enrichment analysis was performed using the

FGSEA package. Inputs included gene sets of interest and ranked differentially

expressed gene lists created with MAST (*81*) and the Seurat *Findmarkers()* function.

Resulting differentially expressed genes were ranked based on the log-fold change (or a

combinatory metric of absolute log-fold change * log transformed p-values) and the

ranked gene list used as input for the fgsea R package (fast preranked GSEA), with a

minimum set size of 15 genes and a maximum of 500 genes. We used the

plotEnrichment() function to visualize the resulting gene set enrichment.

Manual curation of NFAT-dependent genes A set of NFAT dependent genes was manually curated through literature review as well as analysis of publicly available data from NFAT-deficient mice.

Placental Histopathology

H&E-stained slides of the placental disk were evaluated by an experienced perinatal pathologist (V.E.D.). Placental lesions were identified according to the criteria established by the Amsterdam Placental Workshop Group Consensus (*82*) and tissue was processed according to standard laboratory protocols. Placental density was calculated using automated image analysis as previously described (*83*).

Immunofluorescence microscopy

Fresh frozen biopsies in OCT compound (Tissue-Tek) were sectioned on a cryostat at 20 mm thickness and placed on Superfrost Plus slides (Fisher) for processing. Slides were placed in 4% formaldehyde in phosphate buffered saline (PBS), pre-chilled to 4°C, for 30 minutes, then washed in PBS plus 0.3% Tween 20 three times for five minutes per wash at room temperature (RT). Slides were then dried for 15 min at RT, and sections were then treated with blocking buffer (PBS plus 1% Triton X-100, 1% normal goat serum, 1% bovine serum albumin) at RT for 1hr. Primary antibodies (rabbit anti-human NCAM1 (CD56), Cell Signaling Tech. 99746T; hamster anti-human CD3e, BioLegend 100302) were diluted (CD56: 1:50; CD3: 1:200) in blocking buffer and applied overnight at 4°C. Sections were washed in PBST three times (five minutes per wash), then treated with

fluorophore-conjugated secondary antibody (donkey anti-rabbit IgG Alexa Fluor 488,

Invitrogen A32790TR; goat anti-hamster IgG Alexa Fluor 594, BioLegend B388503)

diluted 1:1000 in blocking buffer for 1hr at RT. Sections were then washed three times in

PBST, five min per wash, then treated with DAPI (Invitrogen 62248) diluted 1:1000 in

PBST for 1.5min at RT, washed once in PBST, mounted in Prolong Diamond under

coverglass No. 1.5 and allowed to cure overnight. Images were obtained on ECHO

Revolution in upright mode using a 20x PLAN X Apo 0.8 NA objective and a 5MP 12bit

monochrome CMOS camera. Each imaged field consists of a 2416x2016 pixel array with

pixel physical dimensions of 0.1777x0.1777 mm and each field covering 0.15 mm^2 of

tissue.

Image analysis and error estimation

Raw tagged image files were processed with custom software written in MATLAB. The

analysis consisted of identifying centroids representing the centers of nuclei from the

DAPI channel, then assigning cells of positive or negative CD56 or CD3 expression to

each centroid. This assignment was based on relative signal in the CD56 or CD3 channel

in pixels near each centroid within DAPI-based masks created as described here.

Nucleus centroid positions were estimated by a combination of iterative thresholding,

difference-of-Gaussian (DoG) filtering, and iterative erosion. Single image fields were

initially filtered with a Gaussian of radius 1.25 pixels, then corrected for local background

changes by top-hat filtering with a disk-shaped structuring element with a 50 pixel radius.

The resulting image was filtered with a disk-shaped structuring element of 0.9 pixel radius.

On the resulting image, we performed iterative thresholding to find candidate "connected

component" objects potentially representing single isolated nuclei or multiple overlaying

nuclei. Objects whose area spanned the range of nucleus sizes determined empirically

(covering between 500 and 6000 pixels) and with a circularity (calculated as four times

the Area times pi divided by the perimeter) greater than 0.85 were retained as

representing single nuclei. Noncircular objects were subject to iterative erosion with either

a square structuring element or a diamond structuring element until either the object was

divided into two or more objects, in which case two or more new centroids were taken as

representing single nuclei, or the object disappeared entirely, in which case the original

object was kept as a nucleus. Objects larger than 6000 pixels were subsequently

processed by isolating each object's bounding box from the filtered image and replacing

all pixels outside the object with zeros. This object-specific snippet was then processed

by DoG filtering with Gaussians of 10 and 12 radii. The filtered image was binarized,

morphologically closed, and subjected to the same size and circularity selection criteria

already described. This process was performed iteratively to subdivide all large objects

into candidate nucleus objects with the selected size specificities. Centroids were then

generated from all the candidate objects discovered by this method. A watershed was

then applied to the original connected component mask to assign pixels to the nearest

centroid. To call each centroid as expressing CD56 or CD3, the mean intensity in the

CD56 or CD3 channel was assessed over all pixels assigned to each centroid after

filtering with a Gaussian of radius 15 pixels. Threshold selection was performed by finding

the inflection point in the rate of increase in the fraction of centroids called positive as a

function of candidate threshold level.

For error estimation, we compared the computer performance to that of imaging specialists in a set of ten test images each covering 12,500 mm^2 and containing an average of 130±33 (mean ± s.d.) hand-counted nuclei, for a total of 1,295 nuclei spanning a range of densities. Using hand-selected nuclei as the gold standard, we noted four types of error, two related to nuclear segmentation and two for assigning CD56 expression status: 1) failure to detect a nucleus, usually as a result of incorrectly merging multiple nuclei into a single nucleus; 2) counting excess nuclei, usually as a result of merging of multiple nuclei into a single nucleus; 3) incorrect splitting of single nuclei; 4) failure to call a CD56+ nucleus; 5) incorrectly calling a negative nucleus as positive. To calculate (1) and (2), we generated unique nearest-neighbor pairs of hand-selected and computer-generated centroids while using an exclusion rule prohibiting pairing of centroids separated by more than the average diameter of well-separated nuclei (8 mm). We thereby generated lists of unpaired hand-selected and unpaired computer-generated centroids and calculated the fraction of the total. Errors (3) and (4) were calculated by assessing the fraction of positive (or negative) nuclei that the computer called as negative (or positive) out of the successfully paired set of nuclei. The estimated rate for each error is 0.053±0.029, 0.051±0.043, 0.026±0.013, 0.024±0.015 (mean ± SEM). Assuming these errors arise independently, the total uncertainty in the fraction of CD56+ nuclei may be calculated by finding the root of the sum of the squares of the errors. The estimated uncertainty is thus about 8% and may be as high as 11% (0.082±0.030). Using this upper estimate, the mean fraction of cells that are CD56+ in the healthy control biopsies are

0.0303±0.0033 and 0.0357±0.0039, whereas the transplant biopsies have fractions of

0.0181±0.002, 0.0286±0.0031, 0.0125±0.0014, and 0.0122±0.0013.

LIST OF SUPPLEMENTARY MATERIALS

Figs. S1-S6

Tables S1-S3

REFERENCES AND NOTES

1.   S. E. Ander, M. S. Diamond, C. B. Coyne, Immune responses at the maternal-fetal
interface. *Sci Immunol* **4**, (2019).

2.   M. PrabhuDas *et al.*, Immune mechanisms at the maternal-fetal interface: perspectives
and challenges. *Nat Immunol* **16**, 328-334 (2015).

3.   B. Fu *et al.*, Natural Killer Cells Promote Fetal Development through the Secretion of
Growth-Promoting Factors. *Immunity* **47**, 1100-1113 e1106 (2017).

4.   Y. Zhou *et al.*, PBX1 expression in uterine natural killer cells drives fetal growth. *Sci Transl*
Med **12**, (2020).

5.   Q. Li *et al.*, Human uterine natural killer cells regulate differentiation of extravillous
trophoblast early in pregnancy. *Cell Stem Cell* **31**, 181-195 e189 (2024).

6.   E. D. Albrecht, G. J. Pepe, Regulation of Uterine Spiral Artery Remodeling: a Review.
Reprod Sci **27**, 1932-1942 (2020).

7.   A. A. Ashkar, J. P. Di Santo, B. A. Croy, Interferon gamma contributes to initiation of
uterine vascular modification, decidual integrity, and uterine natural killer cell maturation
during normal murine pregnancy. *J Exp Med* **192**, 259-270 (2000).

8.   J. E. Cartwright, R. Fraser, K. Leslie, A. E. Wallace, J. L. James, Remodelling at the
maternal-fetal interface: relevance to human pregnancy disorders. *Reproduction* **140**,
803-813 (2010).

9.   C. M. Craven, T. Morgan, K. Ward, Decidual spiral artery remodelling begins before
cellular interaction with cytotrophoblasts. *Placenta* **19**, 241-252 (1998).

10.   B. A. Croy *et al.*, Imaging of vascular development in early mouse decidua and its
association with leukocytes and trophoblasts. *Biol Reprod* **87**, 125 (2012).

11.   Y. Cui *et al.*, Role of corin in trophoblast invasion and uterine spiral artery remodelling in
pregnancy. *Nature* **484**, 246-250 (2012).

12.   A. M. Felker, B. A. Croy, Natural cytotoxicity receptor 1 in mouse uNK cell maturation and
function. *Mucosal Immunol* **10**, 1122-1132 (2017).

13.   M. J. Guimond, B. Wang, B. A. Croy, Engraftment of bone marrow from severe combined
immunodeficient (SCID) mice reverses the reproductive deficits in natural killer cell-
deficient tg epsilon 26 mice. *J Exp Med* **187**, 217-223 (1998).

14.   J. Hanna *et al.*, Decidual NK cells regulate key developmental processes at the human
fetal-maternal interface. *Nat Med* **12**, 1065-1074 (2006).

15.   A. D. Hazan *et al.*, Vascular-leukocyte interactions: mechanisms of human decidual spiral
artery remodeling in vitro. *Am J Pathol* **177**, 1017-1030 (2010).

16.   J. Kieckbusch, L. M. Gaynor, A. Moffett, F. Colucci, MHC-dependent inhibition of uterine
NK cells impedes fetal growth and decidual vascular remodelling. *Nat Commun* **5**, 3359
(2014).

17.   G. E. Lash *et al.*, Expression of angiogenic growth factors by uterine natural killer cells
during early pregnancy. *J Leukoc Biol* **80**, 572-580 (2006).

18.   P. D. Lima, J. Zhang, C. Dunk, S. J. Lye, B. A. Croy, Leukocyte driven-decidual
angiogenesis in early pregnancy. *Cell Mol Immunol* **11**, 522-537 (2014).

19.   A. Robson *et al.*, Uterine natural killer cells initiate spiral artery remodeling in human
1026   pregnancy. *FASEB J* **26**, 4876-4885 (2012).

1027   20.   A. Robson *et al.*, Uterine spiral artery muscle dedifferentiation. *Hum Reprod* **34**, 1428-
1028   1438 (2019).

1029   21.   S. D. Smith, C. E. Dunk, J. D. Aplin, L. K. Harris, R. L. Jones, Evidence for immune cell
1030   involvement in decidual spiral arteriole remodeling in early human pregnancy. *Am J Pathol*
1031   **174**, 1959-1971 (2009).

1032   22.   W. Zhang *et al.*, Atrial natriuretic peptide promotes uterine decidualization and a TRAIL-
1033   dependent mechanism in spiral artery remodeling. *J Clin Invest* **131**, (2021).

1034   23.   A. C. Staff *et al.*, Failure of physiological transformation and spiral artery atherosis: their
1035   roles in preeclampsia. *Am J Obstet Gynecol* **226**, S895-S906 (2022).

1036   24.   N. Shreeve *et al.*, The CD94/NKG2A inhibitory receptor educates uterine NK cells to
1037   optimize pregnancy outcomes in humans and mice. *Immunity* **54**, 1231-1244 e1234
1038   (2021).

1039   25.   I. Brosens, R. Pijnenborg, L. Vercruysse, R. Romero, The "Great Obstetrical Syndromes"
1040   are associated with disorders of deep placentation. *Am J Obstet Gynecol* **204**, 193-201
1041   (2011).

1042   26.   C. A. Labarrere *et al.*, Failure of physiologic transformation of spiral arteries, endothelial
1043   and trophoblast cell activation, and acute atherosis in the basal plate of the placenta. *Am*
1044   *J Obstet Gynecol* **216**, 287 e281-287 e216 (2017).

1045   27.   C. Guo *et al.*, Single-cell profiling of the human decidual immune microenvironment in
1046   patients with recurrent pregnancy loss. *Cell Discov* **7**, 1 (2021).

1047   28.   B. Strunz *et al.*, Continuous human uterine NK cell differentiation in response to
1048   endometrial regeneration and pregnancy. *Sci Immunol* **6**, (2021).

29.   C. Dominguez Conde *et al.*, Cross-tissue immune cell analysis reveals tissue-specific

features in humans. *Science* **376**, eabl5197 (2022).

30.   J. I. Gray, D. L. Farber, Tissue-Resident Immune Cells in Humans. *Annu Rev Immunol*

40, 195-220 (2022).

31.   N. Marquardt *et al.*, Unique transcriptional and protein-expression signature in human lung

tissue-resident NK cells. *Nat Commun* **10**, 3841 (2019).

32.   M. M. L. Poon *et al.*, Tissue adaptation and clonal segregation of human memory T cells

in barrier sites. *Nat Immunol* **24**, 309-319 (2023).

33.   P. A. Szabo *et al.*, Single-cell transcriptomics of human T cells reveals tissue and

activation signatures in health and disease. *Nat Commun* **10**, 4706 (2019).

34.   T. J. Kim *et al.*, FK506 causes cellular and functional defects in human natural killer cells.

J Leukoc Biol **88**, 1089-1097 (2010).

35.   H. Wang *et al.*, The unexpected effect of cyclosporin A on CD56+CD16- and

CD56+CD16+ natural killer cell subpopulations. *Blood* **110**, 1530-1539 (2007).

36.   N. A. Deshpande *et al.*, Pregnancy outcomes of liver transplant recipients: a systematic

review and meta-analysis. *Liver Transpl* **18**, 621-629 (2012).

37.   N. A. Deshpande *et al.*, Pregnancy outcomes in kidney transplant recipients: a systematic

review and meta-analysis. *Am J Transplant* **11**, 2388-2404 (2011).

38.   N. Jha, A. K. Jha, S. K. Mishra, S. Parida, Thoracic organ transplantation and pregnancy

outcomes: systematic review and meta-analysis. *Arch Gynecol Obstet* **309**, 385-396

(2024).

39.   A. Lauro, C. S. Matsumoto, I. R. Marino, V. Berghella, A review on pregnancy after

intestinal transplantation. *J Matern Fetal Neonatal Med* **30**, 205-212 (2017).

40.   Z. Stanic *et al.*, Successful spontaneous pregnancy after simultaneous kidney and

pancreas transplant. *Int J Gynaecol Obstet* **162**, 773-774 (2023).

41.   O. Yin *et al.*, Mode of Obstetric Delivery in Kidney and Liver Transplant Recipients and

1075   Associated Maternal, Neonatal, and Graft Morbidity During 5 Decades of Clinical Practice.

1076   *JAMA Netw Open* **4**, e2127378 (2021).

1077   42.   I. Brosens, J. J. Brosens, G. Benagiano, The risk of obstetrical syndromes after solid organ

1078   transplantation. *Best Pract Res Clin Obstet Gynaecol* **28**, 1211-1221 (2014).

1079   43.   D. B. McKay *et al.*, Reproduction and pregnancy in transplant recipients: current practices.

1080   *Prog Transplant* **16**, 127-132 (2006).

1081   44.   D. M. Rupley *et al.*, Preconception counseling, fertility, and pregnancy complications after

1082   abdominal organ transplantation: a survey and cohort study of 532 recipients. *Clin*

1083   *Transplant* **28**, 937-945 (2014).

1084   45.   L. Johannesson *et al.*, The First 5 Years of Uterus Transplant in the US: A Report From

1085   the United States Uterus Transplant Consortium. *JAMA Surg* **157**, 790-797 (2022).

1086   46.   B. P. Jones *et al.*, Human uterine transplantation: a review of outcomes from the first 45

1087   cases. *BJOG* **126**, 1310-1319 (2019).

1088   47.   D. Sawinski *et al.*, A multi-institutional study of renal outcomes and renal-related

1089   pregnancy outcomes in uterus transplant recipients. *Am J Transplant* **22**, 3101-3110

1090   (2022).

1091   48.   A. King, V. Wellings, L. Gardner, Y. W. Loke, Immunocytochemical characterization of the

1092   unusual large granular lymphocytes in human endometrium throughout the menstrual

1093   cycle. *Hum Immunol* **24**, 195-205 (1989).

1094   49.   M. J. van den Heuvel *et al.*, Menstrual cycle hormones induce changes in functional

1095   interactions between lymphocytes and decidual vascular endothelial cells. *J Clin*

1096   *Endocrinol Metab* **90**, 2835-2842 (2005).

1097   50.   I. Korsunsky *et al.*, Fast, sensitive and accurate integration of single-cell data with

1098   Harmony. *Nat Methods* **16**, 1289-1296 (2019).

51. L. M. Ernst, Maternal vascular malperfusion of the placental bed. *APMIS* **126**, 551-560
(2018).

52. Y. M. Kim *et al.*, Failure of physiologic transformation of the spiral arteries in the placental
bed in preterm premature rupture of membranes. *Am J Obstet Gynecol* **187**, 1137-1142
(2002).

53. W. T. Parks, J. M. Catov, The Placenta as a Window to Maternal Vascular Health. *Obstet*
Gynecol Clin North Am **47**, 17-28 (2020).

54. M. Eriksson, S. K. Meadows, C. R. Wira, C. L. Sentman, Unique phenotype of human
uterine NK cells and their regulation by endogenous TGF-beta. *J Leukoc Biol* **76**, 667-675
(2004).

55. E. Giuliani, K. L. Parkin, B. A. Lessey, S. L. Young, A. T. Fazleabas, Characterization of
uterine NK cells in women with infertility or recurrent pregnancy loss and associated
endometriosis. *Am J Reprod Immunol* **72**, 262-269 (2014).

56. M. Greene *et al.*, Sequential transcriptional programs of tissue residency drive human
uterine NK cell development. *bioRxiv* **https://doi.org/10.1101/2024.02.29.582684**,
(2024).

57. P. Dogra *et al.*, Tissue Determinants of Human NK Cell Development, Function, and
Residence. *Cell* **180**, 749-763 e713 (2020).

58. L. Kok, D. Masopust, T. N. Schumacher, The precursors of CD8(+) tissue resident memory
T cells: from lymphoid organs to infected tissues. *Nat Rev Immunol* **22**, 283-293 (2022).

59. O. Huhn *et al.*, Distinctive phenotypes and functions of innate lymphoid cells in human
decidua during early pregnancy. *Nat Commun* **11**, 381 (2020).

60. R. Vento-Tormo *et al.*, Single-cell reconstruction of the early maternal-fetal interface in
humans. *Nature* **563**, 347-353 (2018).

61.   V. Kleshchevnikov *et al.*, Cell2location maps fine-grained cell types in spatial
1124   transcriptomics. *Nat Biotechnol* **40**, 661-671 (2022).

1125   62.   L. K. Mackay *et al.*, The developmental pathway for CD103(+)CD8+ tissue-resident
1126   memory T cells of skin. *Nat Immunol* **14**, 1294-1301 (2013).

1127   63.   L. K. Mackay *et al.*, T-box Transcription Factors Combine with the Cytokines TGF-beta
1128   and IL-15 to Control Tissue-Resident Memory T Cell Fate. *Immunity* **43**, 1101-1111
1129   (2015).

1130   64.   A. P. Nath *et al.*, Comparative analysis reveals a role for TGF-beta in shaping the
1131   residency-related transcriptional signature in tissue-resident memory CD8+ T cells. *PLoS*
1132   *One* **14**, e0210495 (2019).

1133   65.   N. S. Kurd *et al.*, Early precursors and molecular determinants of tissue-resident memory
1134   CD8(+) T lymphocytes revealed by single-cell RNA sequencing. *Sci Immunol* **5**, (2020).

1135   66.   M. Mokrani, J. Klibi, D. Bluteau, G. Bismuth, F. Mami-Chouaib, Smad and NFAT pathways
1136   cooperate to induce CD103 expression in human CD8 T lymphocytes. *J Immunol* **192**,
1137   2471-2479 (2014).

1138   67.   J. J. Milner *et al.*, Runx3 programs CD8(+) T cell residency in non-lymphoid tissues and
1139   tumours. *Nature* **552**, 253-257 (2017).

1140   68.   G. P. Mognol, F. R. Carneiro, B. K. Robbs, D. V. Faget, J. P. Viola, Cell cycle and
1141   apoptosis regulation by NFAT transcription factors: new roles for an old player. *Cell Death*
1142   *Dis* **7**, e2199 (2016).

1143   69.   J. Fric *et al.*, NFAT control of innate immunity. *Blood* **120**, 1380-1389 (2012).

1144   70.   J. Barlic *et al.*, Interleukin (IL)-15 and IL-2 reciprocally regulate expression of the
1145   chemokine receptor CX3CR1 through selective NFAT1- and NFAT2-dependent
1146   mechanisms. *J Biol Chem* **279**, 48520-48534 (2004).

71. L. G. Hawke, B. Z. Mitchell, M. L. Ormiston, TGF-beta and IL-15 Synergize through MAPK

Pathways to Drive the Conversion of Human NK Cells to an Innate Lymphoid Cell 1-like

Phenotype. *J Immunol* **204**, 3171-3181 (2020).

72. M. J. Shannon, E. M. Mace, Natural Killer Cell Integrins and Their Functions in Tissue

Residency. *Front Immunol* **12**, 647358 (2021).

73. S. Klein-Hessling *et al.*, NFATc1 controls the cytotoxicity of CD8(+) T cells. *Nat Commun*

8, 511 (2017).

74. K. Kitaya *et al.*, IL-15 expression at human endometrium and decidua. *Biol Reprod* **63**,

683-687 (2000).

75. W. Wang *et al.*, Single-cell transcriptomic atlas of the human endometrium during the

menstrual cycle. *Nat Med* **26**, 1644-1653 (2020).

76. C. C. Wu, S. C. Hsu, H. M. Shih, M. Z. Lai, Nuclear factor of activated T cells c is a target

of p38 mitogen-activated protein kinase in T cells. *Mol Cell Biol* **23**, 6442-6454 (2003).

77. C. Hafemeister, R. Satija, Normalization and variance stabilization of single-cell RNA-seq

data using regularized negative binomial regression. *Genome Biol* **20**, 296 (2019).

78. Y. Hao *et al.*, Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587

e3529 (2021).

79. H. M. Kang *et al.*, Multiplexed droplet single-cell RNA-sequencing using natural genetic

variation. *Nat Biotechnol* **36**, 89-94 (2018).

80. F. A. Wolf, P. Angerer, F. J. Theis, SCANPY: large-scale single-cell gene expression data

analysis. *Genome Biol* **19**, 15 (2018).

81. G. Finak *et al.*, MAST: a flexible statistical framework for assessing transcriptional

changes and characterizing heterogeneity in single-cell RNA sequencing data. *Genome*

Biol **16**, 278 (2015).

82.   T. Y. Khong *et al.*, Sampling and Definitions of Placental Lesions: Amsterdam Placental

Workshop Group Consensus Statement. *Arch Pathol Lab Med* **140**, 698-713 (2016).

83.   D. Kidron, I. Vainer, Y. Fisher, R. Sharony, Automated image analysis of placental villi and

syncytial knots in histological sections. *Placenta* **53**, 113-118 (2017).

1175

ACKNOWLEDGEMENTS

The authors would like to thank the following individuals and organizations for their

following contributions to this work:

1) Legacy of Hope Organ Procurement Organization

2) Uterus transplant clinical care team at the University of Alabama at Birmingham

3) Staff at the UAB flow cytometry and single cell core facilities

4) Mr. Reid Jones for support of the UAB uterus transplant program

1183

Funding

This work was supported by a pilot grant from the UAB Center for Women's Reproductive

Health (to P.M.P.); UAB AMC21 grants (to J.E.L., J.F.G., & P.M.P.); and start-up funding

from the UAB Heersink School of Medicine (to P.M.P.). Salary support was provided by

R01AI145905 (to P.M.P.); R01CA208353 (to A.G.F. and P.M.P.); American Cancer

Society Research Scholar Grant #RSG-23-1153857-01-IBCD (to A.G.F.).

1190

Author Contributions

R.A.: data curation, analysis, visualization, software.

B.K.: data analysis; writing – original draft; conceptualization.

M.E.G.: investigation; data analysis, conceptualization.

E.W., M.B., D.E., S.D.Y., M.D.C., R.B., N.A., V.E.D., H.E.R., D.G.: investigation,

methodology.

J.E.L.: resources, funding acquisition.

M.V.G.: data analysis, conceptualization.

S.F., E.B., and S.C.: project administration.

J.F.G.: resources, supervision, funding acquisition.

A.G.F.: conceptualization.

S.C.L.: conceptualization, supervision, data analysis, data curation, investigation,

visualizations, writing – original draft, software.

P.M.P.: conceptualization, supervision, data analysis, visualizations, writing – original

draft, funding acquisition.

All authors: writing – review and editing.

1207

1208 **Competing Interests:** The authors have no competing interests to declare.

1209

1210 **Data & Code Availability**

1211 Raw FASTQ files have been deposited to NCBI GEO under accession number

1212 GSEXXXXXX.

1213 Additional data are available upon request from the corresponding author.

1214

1215

Fig. 1

Figure 1. Reduced NK cell frequency among UTx endometrial immune cells. (**A**) UMAP visualization of broad and fine immune clusters among secretory phase endometrial biopsies from healthy control volunteers (n=6) (HC) and uterus transplant recipients (n=5) (UTx). Contaminating stromal and epithelial cells are also shown. (n=77,481 total cells) (**B**) Expression of major cell-type specific marker genes among CD45+ immune cells. (**C**) Module scores of decidual NK cell transcriptomes. (**D**) Frequence of donor-derived (yellow) or recipient-derived (green) cells in each cluster. (**E**) NK cells are reduced in frequency among sequenced CD45+ immune cells in human endometrium. **P<0.01; Mann Whitney U test (one-tailed). ns=not significant. (**F**) Flow cytometric analysis was performed in parallel on three UTx secretory phase biopsies and an expanded cohort of HCs (n=16). Bivariate plots are representative data. Plots are gated on live, CD45+ singlets. **P<0.01; Mann Whitney U test (one-tailed).

	H1	H2	U1	U2	U3
mean	0.0303	0.0357	0.0181	0.0286	0.0125
median	0.0243	0.0285	0.0090	0.0142	0.0054
SD	0.0231	0.0293	0.0294	0.0370	0.0214
# cells	135417	197679	122635	42871	48208
# fields	143	204	149	62	62

Mann-Whitney U test

	H2	U1	U2	U3
H1	0.0136	8.96E-12	0.0984	7.93E-08
H2		3.11E-18	0.0019	1.87E-11
U1			0.0164	0.257
U2				0.0132

Fig. 2

Figure 2. CD56+ cells are reduced in endometrial tissue sections of transplanted uteri. (**A**) Low-magnification DAPI channel from fluorescence microscopy of healthy control endometrial biopsy ("H1" in panels G-I). Box (yellow) indicates area shown in panel B. (**B**) High magnification image showing DAPI (magenta) and anti-CD56 staining (green). (**C**) Computerized rendering of panel B showing CD56+ (green) and CD56- (magenta) nuclei. Boxes indicate areas shown in D. (**D**) Data (left) and computer rendering (right) of the green and orange boxed areas shown in B and C. (**E** and **F**) DAPI (magenta) and anti-CD56 (green) staining in a second control biopsy (E) and a biopsy taken 57 days after uterus transplant (F), designated "H2" and "U1", respectively, in panels G-I. (**G**) Fraction of nuclei designated CD56+ in two healthy control (H1 and H2) and three uterus transplant biopsies (U1-U3). Data points represent individual 0.15 mm2 imaging fields. (**H**) Summary statistics for the five biopsies. (**I**) p-values (Mann-Whitney U test) for all pairwise comparisons of the five biopsies. Scale bars: 500 µm (A), 100 µm (B), 10 µm (D), 200 µm (E,F). See table S1 for additional details regarding samples.

Fig. 3

Figure 3. Maternal vascular malperfusion in UTx placentas. (**A**) UTx "A" and UTx "B" show regions of distal villi that are sparsely distributed, markedly smaller in caliber, show less branching, and have increased numbers of syncytial knots relative to a control placenta (HC) from a similar gestational age. H&E 10X. Gestational age = 34 weeks for all micrographs. (**B**) Density of villi is plotted for each sample as shown. Individual data points represent technical replicates. *P=0.05; one-sided Mann Whitney U test.

Fig. 4

Figure 4. Reduction of trNK cell frequency in UTx. (**A**) Feature plot of trNK and cNK cells in UMAP of aggregated CD45+ immune cells sorted from secretory phase endometrial biopsies from HC (n=6) and UTx (n=5). (**B**) Reciprocal expression of *ITGA1* and *KLF2* in trNK and cNK. (**C**) Expression of human lung tissue resident NK module score in endometrial NK subsets. ****$P<0.0001$; Mann Whitney U test. (**D**) Reduced number of trNK and cNK cells in UTx. (**E**) Frequency of cNK and trNK cells among CD45 immune cells by sample. ns=not significant. **$P<0.01$. Mann Whitney U test. (**F**) Feature plot of trNK subsets and expression of trNK marker genes: *SPINK2* (trNK1), *ITGAX* (trNK2) and *ITGAE* (trNK3). (**G**) Reduced frequency of trNK2 and trNK3 subsets among CD45 endometrial immune cells. **$P<0.01$. Mann Whitney U test. (**H**) Endometrial NK subset abundance in intact biopsies from HC and UTx as measured by spatial transcriptomics.

FIG 5

Figure 5. Impairment of TGFβ-mediated tissue residency programs is associated with loss of trNK subsets in UTx. (**A**) Weighted nearest neighbor UMAP of endometrial NK cells in aggregated secretory phase endometrial biopsies from three HC and 4 UTx (n=7 biopsies). Figure portrays analysis of 8,080 NK cells after re-clustering from CD45+ data. (**B** and **C**) Feature plots of indicated proteins and decidual NK signature scores. (**D**) Loss of trNK3 subsets (clusters 0 & 5; dashed line) in UTx with skewing towards cNK and trNK1 subsets (clusters 1 & 3, respectively). Blue=HC; UTx=Red. *P<0.03, one-sided Mann Whitney U test. (**E**) Enrichment of genes upregulated by TGFβ in healthy control trNK3 subsets. (**F** and **G**) Downregulation of key trNK3 subset genes (F) and proteins (G) by UTx compared to HCs. *P<0.05; **P<0.01; ***P<0.001; ****P<0.0001; ns=not significant; Mann Whitney U test. (**H**) Reduction of frequency of CD103+ CD39- in trNK3 clusters in UTx. Numbers in scatter plots indicate frequency of CD103+CD39- cells within indicated cluster. *P<0.03, one-sided Mann Whitney U test. (**I**) Confirmation of reduction of trNK3 frequency in UTx samples by flow cytometry.

Fig. 6

Figure 6: Reduced expression of early tissue residency developmental program in UTx trNKs. (**A**) Expression of the early developmental program (EDP) of tissue residence in human endometrial NK cells. (**B**) Reduced expression of the EDP in UTx trNKs compared to HCs. (**C**) Reciprocal expression of two developmental modules in trNK1 and trNK3. (**D**) Left: Consolidation of trNK clusters into four groups based on expression of the EDP. Colored bar reflects direction of developmental module expression across the clusters on the UMAP. Right: More pronounced reduction of developmental module 2 (inclusive of AP-1 transcription factors) than developmental module 1 in UTx trNK subsets. ***P<0.001. ****P<0.00001. Mann Whitney U test.

Fig. 7

Figure 7: Reduction of NFAT-dependent tissue residency programs in UTx. (**A**) Identification of 42 NFAT-dependent genes involved in tissue residency programming (NFAT^{TR} signature). (**B**) Enrichment of the NFAT^{TR} signature in endometrial trNKs versus cNKs. (**C**) Variable expression of leading edge genes of the NFAT^{TR} across trNK subsets. (**D**) Enrichment of the endometrial NFAT^{TR} signature in healthy control trNK2 and trNK3 subsets. (**E**) Significant overlap of the endometrial NFAT^{TR} signature with tissue residency developmental module 2. (**F**) Expression of genes in HC and UTx which overlapped between the NFAT^{TR} signature and the tissue residency developmental modules. *P<0.05; **P<0.01; ***P<0.001; ****P<0.0001; ns=not significant; Mann Whitney U test.

Figure 8: NFAT blockade impairs IL-15-induced integrin expression on NK cells. (**A** and **B**) Increase in trNK3 frequency after IL-15 culture in vitro. Secretory phase endometrial biopsies from healthy control volunteers (n=4) (HC) and uteri from deceased donors (DD) (unknown menstrual cycle state) (n=4) were collected and single cell suspension were cultured in vitro under the indicated conditions for 4-7 days. Analysis was performed with flow cytometry. Flow panels indicate representative data. **P<0.01; ns=not significant; paired Wilcoxon signed rank exact test.

Fig. S1

Supplementary Figure 1. Endometrial immune profiling. (**A**) Marker gene expression for major cell clusters of 77,481 CD45+ cells sorted from 11 secretory phase endometrial biopsies (6HC + 5UTx). (**B**) Expression of major marker genes commonly used to define NK cells: *NCAM1* (CD56) and *FCGR3A* (CD16). (**C**) Distinction of CD8 effector T cells from conventional NK cells. Clusters expressing *NCAM1* and *FCGR3A* (C0,3,7,8,12,21) were selected from CD45+ immune cells and re-clustered. (**D**) Expression of T cell-specific genes (i.e., *CD3G*) as well as *NCAM1* identifies cytotoxic CD8 T cell effectors from cNKs.

	H1	H2	U1	U2	U3
mean	0.112	0.108	0.0693	0.0737	0.0794
median	0.105	0.102	0.0531	0.0635	0.0671
SD	0.0584	0.0683	0.0616	0.0644	0.0602
# cells	196188	242124	114970	32510	39106
# fields	219	222	138	48	44

Mann-Whitney U test

	H2	U1	U2	U3
H1	0.202	2.07E-12	1.07E-05	6.70E-05
H2		1.50E-08	6.91E-04	0.00618
U1			0.704	0.153
U2				0.524

Fig. S2

Supplementary Figure 2. CD3-expressing cell density is reduced in transplant endometrium compared to healthy controls. (**A** and **B**) DAPI (magenta) and anti-CD3 (green) staining in health control H1 (A) and transplant U2 (B). Scale bar: 50 µm. (**C** and **D**) Computer rendering of positive (green) and negative (magenta) nuclei in panels A and B. (**E**) Fraction of nuclei designated CD3-positive in healthy control (H1, H2) and uterus transplant (U1-U3) biopsies. (**F**) Summary statistics for the five biopsies. (**G**) p-values (Mann-Whitney U test) for all pairwise comparisons of the five biopsies.

A

Fig. S3

Supplementary Figure 3. Identification of trNK and cNK subsets. (**A**) UMAP embedding of endometrial NK subsets with unique transcriptomes for comparison. NK subsets were selected from CD45+ endometrial immune cells enriched from endometrial biopsies (n=11; 6HC+5UTx). (**B**) Heat map of top 30 DEGs between endometrial trNKs and cNKs. (**C**) Feature plots of gene expression in endometrial trNKs and cNKs which associate with either tissue residency or circulatory programs in lymphocytes.

Fig. S4

Supplementary Figure 4. Cell2location workflow. Spatially barcoded libraries were prepared from one HC secretory phase biopsy and one UTX secretory phase biopsy and input into cell2location along with reference transcriptomes. Reference transcriptomes were derived from aggregated CD45+ endometrial immune cells enriched from HC endometrial biopsies (n=6). Major immune cell clusters are shown for simplicity.

Fig. S5

Supplementary Figure 5. trNK heterogeneity defined by surface proteome and transcriptomes. (**A**) Top 20 DEGs between CD49a- cNK cells and CD49a+ trNK cells. (**B**) Expression of defining marker genes and surface proteins that identify cNK vs. trNKs in human endometrium. (**C**) Differentially expressed proteins on endometrial cNKs. (**D**) Marker genes identify multiple trNK subsets. Only major subsets are shown. (**E**) Differentially expressed proteins identify major trNK subsets: CD69 (trNK3), CD11c (trNK2) and CD39 and KIRs (trNK1). (**F**) Reciprocal expression of trNK1 and trNK3 gene signatures.

Fig. S6

Supplementary Figure 6. scRNA-seq QC Metrics. Figure portrays expression of key QC metrics for scRNA-seq libraries after filtering with the thresholds as described in "Methods".