Differential CheR affinity for chemoreceptor C-terminal pentapeptides biases chemotactic responses

Running title: CheR recognition by chemoreceptors

Félix Velando[#], Elizabet Monteagudo-Cascales[#], Miguel A. Matilla[*] and Tino Krell[*]

Department of Biotechnology and Environmental Protection, Estación Experimental del Zaidín, Consejo Superior de Investigaciones Científicas, Granada, Spain

*Address Correspondence to Tino Krell, tino.krell@eez.csic.es, or Miguel A. Matilla miguel.matilla@eez.csic.es

# Félix Velando and Elizabet Monteagudo-Cascales contributed equally. Author order was determined by a draw.

Summary

The capacity of chemotaxis pathways to respond to signal gradients relies on

adaptation mediated by the coordinated action of CheR methyltransferases and CheB

methylesterases. Many chemoreceptors contain a C-terminal pentapeptide at the end of

a linker. In *Escherichia coli,* this pentapeptide forms a high-affinity binding site for

CheR and phosphorylated CheB, and its removal interferes with adaptation. The

analysis of all available chemoreceptor sequences showed that pentapeptide sequences

vary greatly, and bacteria often possess multiple chemoreceptors that differ in their

pentapeptide sequences. Using the phytopathogen *Pectobacterium atrosepticum*

SCRI1043*,* we assessed whether this sequence variation alters CheR affinity and

chemotaxis. SCRI1043 has 36 chemoreceptors, of which 19 possess a C-terminal

pentapeptide. Using isothermal titration calorimetry, we show that the affinity of CheR

for the different pentapeptides varies up to 11-fold (K_D of 90 nM to 1 μM). The

pentapeptides with the highest and lowest affinities differed only in a single amino acid.

Deletion of the *cheR* gene abolishes chemotaxis. PacC is the sole chemoreceptor for L-

Asp in SCRI1043, and the replacement of its pentapeptide with those having the highest

and lowest affinities significantly interfered with L-Asp chemotaxis. Variable

pentapeptide sequences thus provide a mechanism to bias the responses mediated by

chemoreceptors.

Keywords: signal transduction, chemotaxis, chemoreceptor, CheR, pentapeptide,

molecular recognition

Introduction

Bacterial chemotaxis is the directed swimming of bacteria in chemical gradients. It facilitates the migration of bacteria to sites that are favorable for survival. A major benefit from chemotaxis is access to nutrients (Colin *et al.*, 2021; Matilla *et al.*, 2023). For many human and plant pathogens with different lifestyles and infection mechanisms, chemotaxis is essential for virulence (Matilla and Krell, 2018). Chemotaxis signaling pathways are among the most abundant prokaryotic signal transduction mechanisms (Wuichet and Zhulin, 2010; Gumerov *et al.*, 2023).

Chemotactic responses are typically initiated by a molecule binding to an extracytosolic sensor domain of a chemoreceptor. This binding generates a conformational change that is transmitted across the membrane to modulate the activity of the autokinase CheA, which in turn phosphorylates the CheY response regulator. Only the phosphorylated form of CheY is able to bind to the flagellar motor to control its direction and/or speed of rotation, ultimately resulting in chemotaxis (Parkinson *et al.*, 2015; Bi and Sourjik, 2018).

The capacity of chemotaxis pathways to respond to signal gradients rather than to a constant signal concentration relies on adaptation mechanisms. Canonical adaptation is based on the coordinated action of the CheR methyltransferase and the CheB methylesterase, which catalyze the methylation and demethylation, respectively, of specific glutamyl residues in the chemoreceptor signaling domain (Parkinson *et al.*, 2015; Bi and Sourjik, 2018). The importance of this adaptation mechanism is illustrated by the fact that CheR and CheB are highly conserved proteins that are present in almost all chemosensory pathways (Wuichet and Zhulin, 2010).

In addition to the methylation sites in the chemoreceptor signaling domain, many chemoreceptors contain an additional CheR/CheB-binding site. This consists of a C-terminal pentapeptide fused to the end of a flexible linker (Perez and Stock, 2007; Bartelli and Hazelbauer, 2011; Ortega and Krell, 2020). Pentapeptide-containing chemoreceptors have been identified in 11 different bacterial phyla (Ortega and Krell, 2020). Studies of *Escherichia coli* have shown that the affinity of CheR for the pentapeptide is 50 to 100-fold higher than the affinity for the methylation sites (Wu *et al.*, 1996; Barnakov *et al.*, 1999; Li and Hazelbauer, 2020). Dissociation constants of approximately 2 µM were obtained for CheR binding to the individual pentapeptide and to the pentapeptide-containing chemoreceptor (Wu *et al.*, 1996), indicating that all of the determinants for high-affinity CheR binding are located in the pentapeptide. In

contrast to CheR, unphosphorylated CheB bound to the pentapeptide with low affinity (Barnakov *et al.*, 2002; Velando *et al.*, 2020; Li *et al.*, 2021). However, a stable phosphorylation mimic of *E. coli* CheB had a significantly higher affinity of about 13 µM, so that CheR and phosphorylated CheB both bind to the same high-affinity site (Li *et al.*, 2021). Partial or full truncation of the C-terminal pentapeptide from chemoreceptors greatly decreases both methylation and demethylation *in vivo* and *in vitro,* preventing chemotaxis (Russo and Koshland, 1983; Yamamoto and Imae, 1993; Li *et al.*, 1997; Le Moual *et al.*, 1997; Okumura *et al.*, 1998; Barnakov *et al.*, 1999; Li and Hazelbauer, 2006; Lai *et al.*, 2008; Uchida *et al.*, 2022). However, many bacteria, such as *Pseudomonas putida* KT2440, lack pentapeptide containing chemoreceptors but mediate chemotactic responses that depend on the adaptation enzymes (García-Fontana *et al.*, 2013; García *et al.*, 2015; Martín-Mora *et al.*, 2016), indicating that C-terminal pentapeptides are essential for some receptors, whereas not required for others, which corresponds to an issue that remains poorly understood.

Why are there CheR and CheB-P-binding pentapeptides at the C-terminus of chemoreceptors? It was proposed that high-affinity tethering of CheR and CheB-P to the C-terminal extension increases their local concentration, thereby enhancing their activity and leading to an optimal adaptation (Le Moual *et al.*, 1997; Li and Hazelbauer, 2005; Li *et al.*, 2021). In addition, we have shown previously that pentapeptides also confer specificity in their interaction with CheR and CheB in bacteria that possess multiple pathways. *Pseudomonas aeruginosa* has four chemosensory pathways, each of which contains a CheR and CheB homolog (Matilla *et al.*, 2021). Of the 26 chemoreceptors, McpB (or Aer2) is the only pentapeptide-containing chemoreceptor and the sole chemoreceptor that feeds into the Che2 pathway (Ortega *et al.*, 2017). Only CheR$_2$ and CheB$_2$, the methyltransferase and methylesterase of the Che2 pathway, but not any of the other CheR and CheB homologues bind the McpB/Aer2 pentapeptide (García-Fontana *et al.*, 2014; Velando *et al.*, 2020), permitting the targeting of a particular chemoreceptor with a specific CheR and CheB (García-Fontana *et al.*, 2014; Velando *et al.*, 2020).

All available pentapeptide sequences conserve amino acids with aromatic side chains in positions 2 and 5, whereas a significant variability is observed at the other three positions (Perez and Stock, 2007; Ortega and Krell, 2020). The inspection of the 3D structure of the CheR/pentapeptide complex of *Salmonella enterica* sv. Typhimurium shows that the aromatic side chains at positions 2 and 5 pack into two

hydrophobic pockets, whereas the remaining three side chains establish hydrogen bonds

and a salt bridge with CheR (Djordjevic and Stock, 1998). The conservation of amino

acids at positions 2 and 5 is consistent with studies showing that mutations in these

positions cause severe adaptation defects (Yamamoto and Imae, 1993; Okumura *et al.*,

1998; Shiomi *et al.*, 2000; Lai *et al.*, 2006). Alanine-scanning mutagenesis of residues

at positions 1, 3 and 4 resulted in a reduction of *in vitro* methylation and demethylation

(Lai *et al.*, 2006). Many bacterial strains contain a number of chemoreceptors that differ

in their pentapeptide sequences (Gumerov *et al.*, 2023), and the primary aim of this

study consists in assessing how the naturally occurring variation of pentapeptide

sequences within a bacterium impacts on CheR affinity and magnitude of chemotaxis.

We have addressed this question using *Pectobacterium atrosepticum* as a model.

P. atrosepticum is among the top 10 plant pathogens (Mansfield *et al.*, 2012) and is the

causative agent of soft rot diseases in several agriculturally relevant crops (Toth *et al.*,

2003). The reference strain *P. atrosepticum* SCRI1043 has 36 chemoreceptors, of which

19 possess a C-terminal pentapeptide. We have shown previously that none of these

pentapeptides are recognized by unphosphorylated or the CheB-P mimic beryllium

fluoride-derivatized CheB (Velando *et al.*, 2020). The structure of SCRI1043 CheB can

be closely superimposed onto that of *Salmonella enterica* sv. Typhimurium (Djordjevic

and Stock, 1998; Velando *et al.*, 2020). SCRI1043 has a single chemosensory pathway

(Velando *et al.*, 2020), and the chemoreceptors analyzed thus far respond to amino acids

(PacB, PacC), quaternary amines (PacA), and nitrate (PacN) (Matilla *et al.*, 2022b;

Monteagudo-Cascales *et al.*, 2023; Velando *et al.*, 2023). We show here that there are

significant differences in the affinity of the individual pentapeptides for CheR that are

reflected in differences in the magnitude of the chemotactic responses. The differential

recognition of CheR by pentapeptide may be a mechanism to bias chemotactic

responses.

134 **Results**

135

136 **SCRI1043 chemoreceptors possess 9 different pentapeptides.**

137   A sequence alignment of the C-terminal regions of all chemoreceptors of strain

138 SCRI1043 is shown in Fig. 1A. Nineteen chemoreceptors possess a C-terminal

139 pentapeptide that is tethered to the signaling domain via a linker sequence of 29 to 39

140 amino acids (Fig. 1A). In total, there were 9 different pentapeptide sequences than can

141 be divided into three groups (Table 1). Whereas groups 1 and 2 each contained three

142 pentapeptides that differ in a single amino acid, group 3 contained three pentapeptides

143 with multiple differences. All peptides possess an W and an F at positions 2 and 5,

144 respectively, but share significant variability at the remaining positions. The NWETF

145 pentapeptide that is found in *E. coli* and *S. enterica* sv. Typhimurium receptors is

146 present in eight receptors (Table 1, Fig. 1A). Three pentapeptides, DWTSF, NWTTF

147 and NWEKF, were present in two different chemoreceptors, whereas the remaining

148 pentapeptides were found in only one chemoreceptor (Table 1, Fig. 1A). SCRI1043

149 chemoreceptors possess different sensor domains: PAS, sCache, Cache3_Cache2,

150 dCache, NIT, 4HB and HBM (Velando *et al.*, 2020). Pentapeptides are exclusively

151 found in chemoreceptors that contain a four-helix-bundle sensor domain, either in its

152 monomodular (4HB) or bimodular (HBM) arrangement (Table 1). Pentapeptides

153 NWTTF and DWTSF were found in both 4HB and HBM chemoreceptors.

154

155 **Conserved charge pattern in the linker/pentapeptide sequences.**

156   So far, no conserved feature has been reported for linker sequences. They differ

157 significantly in length, they share no apparent sequence similarities, and they are

158 unstructured (Bartelli and Hazelbauer, 2011; Ortega and Krell, 2020). As expected, the

159 alignment of the 19 linker sequences revealed no apparent sequence similarity (Fig. S1).

160 However, there was a clear pattern of charge distribution (Fig. 1, Table S1). The N-

161 terminal section of the linker frequently had a negative charge, followed by a central

162 positively charged segment. The segment adjacent to the pentapeptide was negatively

163 charged (Fig. 1, Table S1). The average calculated isoelectric point (pI) of the C-

164 terminal 8 amino acids of these 19 linkers was 3.95 ± 0.41, whereas the calculated pI of

165 the remaining linker was 10.82 ± 1.27 (Table S1). The plot of the electrostatic surface

166 charge of Alphafold2 models of linker/pentapeptide segments from three representative

167 chemoreceptors illustrates this pattern (Fig. 1B), which appears to be conserved. It is

present in all pentapeptide-containing chemoreceptors from other chemotaxis model bacteria, including *E. coli*, *S. enterica* sv. Typhimurium, *Sinorhizobium meliloti*, *Azospirillum brasilense*, *Ralstonia solanacearum*, *Comamonas testosteroni*, and *Vibrio cholerae* (Table S2).

SCRI1043 CheR binds to C-terminal pentapeptides with different affinities.

A previous study showed that SCRI1043 CheB and a CheB-P mimic failed to bind all nine C-terminal pentapeptides present in the chemoreceptors of this strain (Velando *et al.*, 2020). To investigate whether these pentapeptides bind CheR, SCRI1043 CheR was overexpressed in *E. coli* and purified from the soluble cell lysate. To verify protein integrity, microcalorimetric titrations with S-adenosylmethionine (SAM) and S-adenosylhomocysteine (SAH), the substrate and product, respectively, of the methylation reaction, were conducted (Fig. 2). Binding of SAM and SAH was characterized by K_D values of 140 ± 20 µM and 480 ± 20 nM, respectively (Table 1), revealing that the reaction product is recognized with strong preference, which agrees with data of CheR from other species (Yi and Weis, 2002; García-Fontana *et al.*, 2013; García-Fontana *et al.*, 2014).

In subsequent studies, CheR was titrated with the nine different pentapeptides of SCRI1043. In all cases, peptides bound, but significant differences in affinity were observed (Fig. 3, Table 1). The titration with the NWETF pentapeptide (Fig. 3), present in *E. coli* and 8 SCRI1043 receptors, revealed a K_D value of 480 nM (Table 1), which is considerably higher affinity than *E. coli* CheR (K_D=2 μM) for this peptide (Wu *et al.*, 1996). NWETF derivatives containing K and Q instead of T bound with similar affinity (Table 1). Of all peptides, GWTTF bound with highest affinity (K_D=90 nM).

Amino acid substitutions at position 1 of the pentapeptide had a pronounced effect on the binding affinity. Substitution of G by N in GWTTF resulted in the pentapeptide that bound with lowest affinity (K_D=1 μM), representing a 11-fold reduction, whereas substitution with D caused a 2.6-fold decrease (Table 1). The three pentapeptides with three changes with respect to NWETF, namely GWQRF, GWEKF and DWTSF, bound with K_D values of 540, 200 and 360 nM, respectively (Table 1). Thus, variation in the residue at the less-conserved positions 1, 3 and 4 still significantly affects CheR binding affinity.

SCRI1043 CheR is essential for chemotaxis.

We have shown previously that SCRI1043 has multiple chemoreceptors for amino acids (Velando *et al.*, 2023). To determine the function of CheR in SCRI1043, we constructed a *cheR* deletion mutant. Quantitative chemotaxis capillary assays showed that this mutant is unable to perform chemotaxis to casamino acids or L-aspartate. Many bacteria are strongly attracted by Krebs cycle intermediates (Matilla *et al.*, 2022a). We observed a strong attraction of SCRI1043 to L-malate (Fig. 4), for which the corresponding chemoreceptor(s) remain(s) unknown. The *cheR* mutant also failed to be attracted to L-malate, indicating that CheR is essential to perform chemotaxis toward amino acids and at least one organic acid.

Alteration of PacC mediated chemotaxis by substitution of pentapeptides

A previous study demonstrated that the strong chemotactic response of SCRI1043 to L-Asp depends entirely on the PacC chemoreceptor (Velando *et al.*, 2023). As in *E. coli* Tar, PacC is encoded in the chemosensory signaling cluster and contains the NWETF pentapeptide. As shown above, CheR recognizes this pentapeptide with a K_D of 480 nM. To investigate the relevance of the CheR affinity for the pentapeptide in the chemotactic response to L-Asp, we generated two chromosomal mutations in which the region encoding the original pentapeptide was altered to encode GWTTF or NWTTF, which bind CheR with either the highest (K_D=90 nM) or lowest (K_D=1 μM) affinity, respectively. Control capillary chemotaxis assays showed that the responses of wt and mutant strains to casamino acids are indistinguishable (Fig. S2).

Chemotaxis experiments with the wt strain showed strong responses towards different L-Asp concentrations, confirming previously published data (Velando *et al.*, 2023). Maximal responses were observed at a concentration of 1 mM. When these experiments were repeated with the mutant strain encoding PacC containing the NWTTF pentapeptide (lowest affinity), a statistically significant drop in the chemotaxis magnitude was observed at a concentrations of 1 mM L-Asp and 10 mM for the receptor containing the GWTTF pentapeptide (Fig. 5). Differences in chemotaxis mediated by the wt and mutant receptors to lower L-Asp concentrations (0.01 and 0.1 Mm) were statistically not significant (Fig. 5). These data demonstrate that the affinity of the CheR-pentapeptide interaction influences the chemotactic response. Either tighter or looser binding of CheR was associated with a decreased chemotaxis response to 1 Mm L-Asp.

Discussion

E. coli and *S. enterica* sv. Typhimurium are the traditional model organisms in

chemotaxis research. Two of their chemoreceptors, Tar and Tsr, contain the NWETF

pentapeptide and most of what we know about pentapeptide function is therefore based

on the study of the NWETF pentapeptide (Wu *et al.*, 1996; Djordjevic and Stock, 1998;

Feng *et al.*, 1999; Barnakov *et al.*, 1999; Shiomi *et al.*, 2000; Lai and Hazelbauer, 2005;

Lai *et al.*, 2006; Lai *et al.*, 2008; Li and Hazelbauer, 2020). Bioinformatic studies of all

available chemoreceptor sequences show that their C-terminal pentapeptides differ in

sequence (Perez and Stock, 2007; Ortega and Krell, 2020), and many strains contain

receptors with pentapeptides having different sequences (Gumerov *et al.*, 2023). Here,

we present the first study evaluating CheR recognition of the complete set of different

pentapeptides within a bacterial strain.

A previous study showed that native CheB or the beryllium fluoride-modified

CheB-P mimic of SCRI1043 did not interact with any of these pentapeptides (Velando

et al., 2020), although it is unknown if an interaction with CheB-P may occur *in vivo*. In

contrast, all 9 pentapeptides were recognized by SCRI1043 CheR. The affinities were

between 90 nM and 1 μM, and thus higher than the affinity of *E. coli* CheR for NWETF

of about 2 μM (Wu *et al.*, 1996; Li and Hazelbauer, 2020), but comparable to the K_D of

about 500 nM derived for the interaction of *P. aeruginosa* CheR$_2$ with the GWEEF

peptide of the McpB/Aer2 chemoreceptor (García-Fontana *et al.*, 2014).

The importance of each of the 5 amino acids of the NWETF pentapeptide in *E.*

coli has been investigated using alanine-scanning mutagenesis (Lai *et al.*, 2006).

Whereas mutation of W and F abolished methylation and demethylation *in vitro*,

mutations of the remaining three amino acids resulted in a partial reduction of both

activities. However, alanine is an amino acid that is rarely found in positions 1, 3 and 4

(Perez and Stock, 2007; Ortega and Krell, 2020). Here, we investigated naturally

occurring pentapeptides and found that they differ in their affinity for CheR by up to 11-

fold. These alterations are consistent with the 3D structure of the *S. enterica*

CheR/pentapeptide complex showing that the N and T of the NWETF pentapeptide

establish hydrogen bonds with the protein, whereas the E forms a salt bridge with a

CheR Arg residue (Djordjevic and Stock, 1998). Altering these interactions by amino

acid replacements might be expected to alter binding affinity.

The notion that residues at positions 1, 3, and 4 do not occur in a random fashion

but are rather the result of evolution is supported by several observations. First, the

sequence logo of all naturally occurring pentapeptides shows clear amino acid

preference. In all three positions aspartate or glutamate residues are predominant

(Ortega and Krell, 2020). Second, the single amino acid substitutions in pentapeptides

of groups 1 and 2 are frequently caused by two nucleotide changes (Table 1) and are

thus not single nucleotide polymorphisms.

In their natural environment, bacteria are exposed to complex mixtures of

compounds, many of which are chemoeffectors. Also, in many bacteria, multiple

chemoreceptors stimulate a single chemotaxis pathway (Gumerov *et al.*, 2023).

However, the mechanisms that permit the generation of specific chemotactic responses

in the presence of complex chemoeffector mixtures is unclear.

How is the weight an individual chemoreceptor in the final response

determined? One contribution is the affinity with which individual chemoeffectors are

recognized by different chemoreceptors; a parameter that is known to determine the

onset of a chemotactic response (Reyes-Darias *et al.*, 2015; Ortega and Krell, 2020).

Other factors are the level of expression and cellular abundance of chemoreceptors

(Feng *et al.*, 1997; Zatakia *et al.*, 2018). Here, we have found that changes in the

pentapeptide sequence can also alter the magnitude of the chemotactic response.

Differences in affinity may result in a differential occupation of chemoreceptors with

CheR. In contrast to *E. coli*, where CheB was found to bind to the pentapeptide (Li *et*

al., 2021), we have failed to observe an interaction of any of the pentapeptides with *P.*

atrosepticum CheB either in its unphosphorylated form or as a CheB-P mimic (Velando

et al., 2020). However, an interaction with CheB-P may occur in the cell, and the failure

to detect this interaction *in vitro* may be due to a technical issue. Considering that

pentapeptides differ in their affinities for CheR, it is reasonable to suggest that such

differences may also occur for binding of CheB-P. Thus, an altered interaction with

CheB-P may also contribute to the dependence of PacC-mediated aspartate taxis seen

with different pentapeptide sequences.

Optimal chemotaxis to L-Asp was observed when the PacC receptor contained

its native NWETF pentapeptide, which has a K_D for CheR of 480 nM. When the

pentapeptide was converted to NWTTF, which binds CheR with a K_D that is ~2-fold

higher (1 µM), or to GWTTF, which binds CheR with a K_D that is ~5-fold lower (90

nM), the chemotactic responses were lower at 1 mM L-Asp. As noted above, the

affinity of different pentapeptides for the other adaptation enzyme, the CheB-P

demethylase, may also vary. Optimal responses were thus observed for the receptor

with the naturally occurring pentapeptide, suggesting that CheR recognition at
chemoreceptors has been finely tuned to achieve optimum chemotaxis.

The linker that leads to the pentapeptide does not have a conserved sequence. It
is considered to be an unstructured, disordered and flexible arm that facilitates the
movement of tethered adaptation enzymes within the receptor array (Bartelli and
Hazelbauer, 2011). We report here that there is a conserved distribution of charge
within SCRI1043 chemoreceptors (Table S1) that is also conserved in the receptors of
bacteria that belong to different phylogenetic categories (Table S2). This pattern is
independent of linker length: it was detected in the 29-residue linker of the *S. enterica*
sv. Typhimurium Tcp chemoreceptor as well as in the 91-residue linker of *R.*
solanacearum RS_RS23910 (Table S2). Our observations should prompt studies to
investigate the functional relevance of this pattern. The signaling domain of
chemoreceptors typically has a significant negative surface charge (Akkaladevi *et al.*,
2018) (Fig. S3). The positive charge of the central linker segment may mediate a charge
attraction with the signaling domain, whereas the negative charge of the C-terminal
portion of the linker that connects to the pentapeptide may cause a repulsion away from
the signaling domain. The notion that this conserved charge pattern is of functional
relevance is supported by a study of *E. coli* Tar (Lai *et al.*, 2008). The authors generated
alanine-substitution mutants of the lysine and three arginine residues present in the Tar
linker (Table S2). Whereas receptors with individual substitutions supported chemotaxis
equivalent to that supported by wild-type Tar, combining all four changes decreased
chemotaxis by about 50%. Importantly, the singly substituted receptors showed a
significant increase in CheA activity that was increased when multiple substitutions
were combined. The authors concluded that the Ala substitutions either stabilize the ON
state or destabilize the OFF state. Whereas the initial 30 amino acids of this linker have
a calculated pI of 10.74 (Table S2), replacing all four positively charged amino acids
with Ala lowered the pI to 3.80. It was suggested that the positively charged linker
either interacts with the negatively charged polar head groups of phospholipids or with
charged residues in the signaling domain of the receptor (Lai *et al.*, 2008). Combined
with the observation that this charge pattern is conserved, this result supports the idea
that this organization of the linker is of fundamental functional relevance.

Collectively, we show here that naturally occurring variations in the
pentapeptide sequence cause important alterations in the binding affinities of CheR.
Maximal responses were obtained for a chemoreceptor containing its naturally

occurring pentapeptide, suggesting that the pentapeptide sequence has been fine-tuned to guarantee maximal responses. A conserved charge pattern was observed in linker sequences that is likely to contribute to the spatial organization of the linker within the chemoreceptor array. Our study forms the basis for investigations on how pentapeptide sequence variation affects the magnitude of chemotaxis in other bacteria and to determine the functional relevance of the observed charge pattern of the linker.

Experimental procedures

Bacterial strains and growth conditions: Bacterial strains used in this study are listed in Table S3. SCRI1043 and its derivative strains were routinely grown at 30 °C in Luria Broth (5 g/l yeast extract, 10 g/l bacto tryptone, 5 g/l NaCl) or minimal medium (0.41 mM $MgSO_4$, 7.56 mM $(NH_4)_2SO_4$, 40 mM K_2HPO_4, 15 mM KH_2PO_4) supplemented with 0.2% (w/v) glucose as carbon source. *E. coli* strains were grown at 37 °C in LB. *Escherichia coli* DH5α was used as a host for gene cloning. Media for propagation of *E. coli* β2163 were supplemented with 300 mM 2,6-diaminopimelic acid. When appropriate, antibiotics were used at the following final concentrations (in $\mu g\ ml^{-1}$): kanamycin, 50; streptomycin, 50; ampicillin, 100. Sucrose was added to a final concentration of 10% (w/v) when required to select derivatives that had undergone a second cross-over event during marker exchange mutagenesis.

In vitro nucleic acid techniques: Plasmid DNA was isolated using the NZY-Tech miniprep kit. For DNA digestion, the manufacturer's instructions were followed (New England Biolabs and Roche). Separated DNA fragments were recovered from agarose using the Qiagen gel extraction kit. Ligation reactions were performed as described in (Sambrook *et al.*, 1989). PCR reactions were purified using the Qiagen PCR Clean-up kit. PCR fragments were verified by DNA sequencing that was carried out at the Institute of Parasitology and Biomedicine Lopez-Neyra (CSIC; Granada, Spain). Phusion® high fidelity DNA polymerase (Thermo Fisher Scientific) was used in the amplification of PCR fragments for cloning.

CheR overexpression and purification: The DNA sequence encoding CheR of SCRI1043 (ECA_RS08375) was amplified by PCR using the oligonucleotides indicated in Table S4 and genomic DNA as template. The PCR product was digested with NheI and SalI and cloned into pET28b(+) linearized with the same enzymes. The resulting plasmid pET28b-CheR was verified by DNA sequencing and transformed into *E. coli* BL21-AI™. The strain was grown under continuous stirring (200 rpm) at 30 ℃ in 2-liter Erlenmeyer flasks containing 500 ml of LB medium supplemented with 50 μg/ml kanamycin. At an OD_{660} of 0.6, protein expression was induced by the addition of 0.1 mM isopropyl β-D-1-thiogalactopyranoside (IPTG) and 0.2 % (w/v) L-arabinose. Growth was continued at 16 °C overnight prior to cell harvest by centrifugation at 10

000 x *g* for 30 min. The cell pellet was resuspended in buffer A (40 mM

KH_2PO_4/K_2HPO_4, 10 mM imidazole, 10% (v/v) glycerol, 1 mM β-mercaptoethanol, pH

7.5) and cells were broken by French press treatment at 62.5 lb/in^2. After centrifugation

at 20 000 x *g* for 30 min, the supernatant was loaded onto a 5-ml HisTrap HP columns

(Amersham Biosciences) equilibrated with buffer A and eluted with an imidazole

gradient of 40–500 mM in the same buffer. Protein-containing fractions were pooled,

and dialyzed overnight against 40 mM KH_2PO_4/K_2HPO_4, 10% (v/v) glycerol, 1 mM β-

mercaptoethanol, pH 7.0. All manipulations were carried out at 4 ºC. Experiments were

conducted with the hexa-histidine tagged protein.

Isothermal titration calorimetry (ITC): All experiments were conducted on a VP-

microcalorimeter (MicroCal, Amherst, MA) at 25 °C. CheR was dialyzed overnight

against 40 mM KH_2PO_4/K_2HPO_4, 10% (v/v) glycerol, 1 mM β-mercaptoethanol, pH

7.0, adjusted to a concentration of 10–20 μM, placed into the sample cell and titrated

with 3.2–4.8 μl aliquots of 0.1–1 mM peptide solutions (purchased from GenScript,

Piscataway, NJ, USA), 2 mM SAM or 250 μM SAH. All solutions were prepared in

dialysis buffer immediately before use. The mean enthalpies measured from the

injection of the peptide into the buffer were subtracted from raw titration data prior to

data analysis with the MicroCal version of ORIGIN. Data were fitted with the 'One

binding site model' of ORIGIN.

Construction of a mutant deficient in cheR. A chromosomal *cheR* mutant of SCRI1043

was constructed by homologous recombination using a derivative plasmid of the suicide

vector pKNG101. The up- and downstream flanking regions of *cheR* were amplified by

PCR using the oligonucleotides listed in Table S4, which were subsequently digested

with EcoRI/BamHI and BamHI/PstI, respectively, and ligated in a three-way ligation

into the EcoRI/PstI sites of pUC18Not to generate pUC18Not_ΔcheR. Subsequently, a

0.95-kb BamHI fragment containing the Km3 cassette of p34S-Km3 was inserted into

the BamHI site of pUC18Not_ΔcheR, giving rise to pUC18Not_ΔcheR_km3. Lastly, a

~2.5-kb NotI fragment of pUC18Not_ΔcheR_km3 was cloned at the same site into

pKNG101, resulting in pKNG101_ΔcheR. This plasmid was transferred to SCRI1043

by biparental conjugation using *E. coli* β2163. Then, cells were spread onto LB plates

containing kanamycin at 50 μg/ml. Selected merodiploid colonies were spread onto LB

plates containing 10 % (w/v) sucrose to select derivatives that had undergone a second

cross-over event during marker exchange mutagenesis. All plasmids and the final
mutant were confirmed by PCR and sequencing.

Construction of SCRI1043 mutant strains pacC-*NWTTF and* pacC-*GWTTF*.
Chromosomal mutants of SCRI1043 encoding PacC variants with altered pentapeptides
were constructed by homologous recombination using derivative plasmids of the suicide
vector pKNG101. Briefly, an overlapping PCR mutagenesis approach was employed to
construct *pacC* gene variants in which the region encoding the original NWETF
pentapeptide sequence was altered to encode GWTTF or NWTTF, using
oligonucleotides listed in Table S4. The corresponding ~1-kb PCR products were then
digested with EcoRI/PstI and cloned into the same sites of pUC18Not to generate
pUC18Not-pacC-NWTTF and pUC18Not-pacC-GWTTF. Subsequently, ~1.1-kb NotI
fragments of these plasmids were cloned at the same site in pKNG101 to generate
pKNG-pacC-NWTTF and pKNG-pacC-GWTTF, which were transferred to *P.*
atrosepticum SCRI1043 by biparental conjugation using *E. coli* β2163. After overnight
incubation at 30 ℃, cells were spread onto LB plates containing streptomycin at 50
µg/ml. Selected merodiploid colonies were spread onto LB plates containing 10 % (w/v)
sucrose to select derivatives that had undergone a second cross-over event during
marker exchange mutagenesis. All plasmids and final mutants were confirmed by PCR
and sequencing.

Quantitative capillarity chemotaxis assays. Overnight bacterial cultures of SCRI1043
were grown at 30 ℃ in minimal medium. At an OD_{660} of 0.3-0.4, the cultures were
washed twice by centrifugation ($1,667 \times g$ for 5 min at room temperature) and
subsequent resuspension in chemotaxis buffer (50 mM K_2HPO_4/KH_2PO_4, 20 µM EDTA
and 0.05% (v/v) glycerol, pH 7.0). Cells were diluted to an OD_{660} of 0.1 in the same
buffer. Subsequently, 230 µl aliquots of the resulting bacterial suspension were place
into 96-well plates. One-microliter capillary tubes (P1424, Microcaps; Drummond
Scientific) were heat-sealed at one end and filled with either the chemotaxis buffer
(negative control) or chemotaxis buffer containing the chemoeffectors to test. The
capillaries were immersed into the bacterial suspensions at its open end. After 30 min at
room temperature, the capillaries were removed from the bacterial suspensions, rinsed
with sterile water and the content expelled into tubes containing 1 ml of minimal
medium salts. Serial dilutions were plated onto minimal medium supplemented with 15

mM glucose as carbon source. The number of colony forming units was determined

after 36 h incubation at 30 °C. In all cases, data were corrected with the number of cells

that swam into buffer containing capillaries. Data are the means and standard deviations

from at least three biological replicates conducted in triplicate.

Acknowledgements: We are indebted to Dr. Michael Manson for his constructive

scientific criticism and editing the English of this manuscript. We would also like to

thank Dr. Gerald Hazelbauer for his advice. This work was supported by the Spanish

Ministry for Science and Innovation/*Agencia Estatal de Investigación*

10.13039/501100011033 through grants PID2020-112612GB-I00 to TK and PID2019-

103972GA-I00 to MAM, CSIC grant 2023AEP002 to MAM, and the Junta de

Andalucía (grant P18-FR-1621 to TK).

Author contributions: FV: investigation, formal analysis, visualization, writing –

review & editing; EMC: investigation, formal analysis, visualization, writing – review

& editing; MAV: Conceptualization, funding acquisition, project administration,

supervision, writing – review & editing; TK: Conceptualization, funding acquisition,

project administration, supervision, writing – original draft preparation, writing –

review & editing

Abbreviated Summary: The nine different pentapeptides present in *P. atrosepticum*

SCRI1043 differ in their affinity for the CheR methyltransferase up to 11-fold.

Replacement of the naturally occurring NWETF pentapeptide at the PacC

chemoreceptor with pentapeptides the bind with lower and higher affinity to CheR

reduces the magnitude of chemotaxis to 1 mM L-Asp. The linker sequences showed a

charge pattern that was conserved in all pentapeptide containing chemoreceptors of

other chemotaxis model strains.

Abbreviations: 4HB, four-helix bundle domain; HBM, helical bimodular domain; ITC,

isothermal titration calorimetry; LBD, ligand binding domain; SCRI1043,

Pectobacterium atrosepticum SCRI1043; SAM, S-adenosylmethionine; SAH, S-

adenosylhomocysteine.

Figure legends

Fig. 1) C-terminal pentapeptides at *P. atrosepticum* SCRI1043 chemoreceptors. A) The C-terminal section of the sequence alignment of SCRI1043 chemoreceptors is shown. Pentapeptides are shaded in grey. The linker sequences are underlined. Green: conserved residues in the signaling domain; red: Asp and Glu; blue: Lys and Arg. B) The electrostatic surface potential of the linker + pentapeptide sequences from three representative SCRI1043 chemoreceptors. Fragments from molecular models of the entire receptor generated by Alphafold2 (Jumper *et al.*, 2021) are shown. Calculations were made using the "APBS Electrostatics" plugin of PyMOL (Schrodinger, 2010). Blue: positive charge; red: negative charge.

Fig. 2) Binding of S-adenosylmethionine (SAM) and S-adenosylhomocysteine (SAH) to *P. atrosepticum* CheR. Microcalorimetric titrations of 10 µM CheR with 16 µl aliquots of 2 mM SAM (A) and of 14 µM CheR with 4.8 µl aliquots of 250 µM SAH (B). Upper panel: Raw titration data. Lower panel: Data corrected for dilution heat and concentration-normalized integrated peak areas of raw titration data. Data were fitted with the "One binding site model" of the MicroCal version of ORIGIN.

Fig. 3) Interaction between *P. atrosepticum* pentapeptides and CheR. Upper panel: Microcalorimetric titrations of 16-20 µM CheR with 1 mM solutions of pentapeptides NWTTF, NWETF and GWTTF. Lower panel: Data corrected for dilution heat and concentration-normalized integrated peak areas of raw titration data. Data were fitted with the "One binding site model" of the MicroCal version of ORIGIN. Derived binding parameters are provided in Table 1.

Fig. 4) Quantitative capillary chemotaxis assays of wild-type *P. atrosepticum* SCRI1043 and a Δ*cheR* mutant towards different chemoattractants. Data have been corrected for the number of cells that swam into buffer containing capillaries (113 ± 19, 370 ± 64, and 305 ± 49 cells/capillary for L-malate, casamino acids (CAA), and L-aspartate, respectively). The means and standard deviations from three independent experiments conducted in triplicate are shown.

Fig. 5) Quantitative L-Asp capillary chemotaxis assays of wild type P. atrosepticum SCRI1043 and chromosomal mutants containing variations in the PacC pentapeptide sequence. Data have been corrected for the number of bacteria that swam into buffer-containing capillaries (872 ± 248 for wt, 975 ± 517 for the PacC-NWTTF mutant, and 445 ± 109 for the PacC-GWTTF mutant). Data are the means and standard deviations from at least three biological replicates conducted in triplicate. * $P<0.05$ in Student's t-test, ** $P<0.01$.

Table 1) Binding parameters for the interaction of *P. atrosepticum* CheR with S-adenosylmethionine (SAM), S-adenosylhomocysteine (SAH) and pentapeptides. Amino acid changes and the corresponding alterations in the nucleotide sequence in pentapeptides from groups 1, 2 and 3 with respect to the reference pentapeptide in each group are highlighted in red color.

Pentapeptide group[a]	Ligand	Nucleotide sequence in corresponding gene	Chemoreceptor (sensor domain type)[b]	K_D[c] (nM)
-	SAM	-	-	$144,000 \pm 20$
	SAH	-	-	480 ± 20
Group 1	NWETF	AACTGGGAAACCTTT	ECA_RS21450 (4HB)	480 ± 40
		AACTGGGAAACCTTT	ECA_RS21455 (4HB)	
		AATTGGGAAACCTTC	ECA_RS08370 (4HB)	
		AATTGGGAAACGTTT	ECA_RS06345 (4HB)	
		AACTGGGAAACGTTC	ECA_RS08330 (4HB)	
		AATTGGGAAACGTTC	ECA_RS19280 (4HB)	
		AACTGGGAAACTTTC	ECA_RS13300 (4HB)	
		AACTGGGAAACCTTT	ECA_RS21445 (4HB)	
	NWE**K**F	AACTGGGAAA**AA**TTC	ECA_RS20365 (4HB)	790 ± 150
		AACTGGGAAA**AA**TTC	ECA_RS06625 (4HB)	
	NWE**Q**F	AACTGGGAA**CAG**TTC	ECA_RS21440 (4HB)	430 ± 20
Group 2	NWTTF	AATTGGACGACGTTC	ECA_RS00900 (4HB)	1000 ± 0.20
		AACTGGACAACGTTC	ECA_RS00400 (HBM)	
	GWTTF	**GG**CTGGACGACATTC	ECA_RS12635 (HBM)	90 ± 10
	DWTTF	**G**ACTGGACCACGTTC	ECA_RS08780 (4HB)	230 ± 40
Group 3	GWQRF	GGCTGGCAGCGTTTT	ECA_RS15955 (4HB)	540 ± 110
	GW**EK**F	GGATGG**GA**AA**AG**TTC	ECA_RS07510 (4HB)	200 ± 40
	DW**TS**F	**G**ACTGG**ACATCG**TTC	ECA_RS18000 (4HB)	360 ± 60
		GACTGG**ACGTCG**TTT	ECA_RS12640 (HBM)	

[a]The pentapeptides were classified into 3 groups depending on the number of changes in the amino acid residues with respect to the reference pentapeptide in the group, with groups 1 and 2 presenting one change, whereas pentapeptides in group 3 present two or three amino acid changes. Residues that differ from NWETF (group 1), NWTTF (group 2) or GWQRF (group 3) are shown in red font.

[b]4HB: 4-helix bundle domain; HBM: helical bimodular domain.

[c]Data were derived from microcalorimetric titrations (Fig. 3).

References

Akkaladevi, N., Bunyak, F., Stalla, D., White, T.A., and Hazelbauer, G.L. (2018) Flexible Hinges in Bacterial Chemoreceptors. *J Bacteriol* **200**: e00593-17.

Barnakov, A.N., Barnakova, L.A., and Hazelbauer, G.L. (1999) Efficient adaptational demethylation of chemoreceptors requires the same enzyme-docking site as efficient methylation. *Proc Natl Acad Sci U S A* **96**: 10667–72.

Barnakov, A.N., Barnakova, L.A., and Hazelbauer, G.L. (2002) Allosteric enhancement of adaptational demethylation by a carboxyl-terminal sequence on chemoreceptors. *J Biol Chem* **277**: 42151–6.

Bartelli, N.L., and Hazelbauer, G.L. (2011) Direct evidence that the carboxyl-terminal sequence of a bacterial chemoreceptor is an unstructured linker and enzyme tether. *Protein Sci* **20**: 1856–66.

Bi, S., and Sourjik, V. (2018) Stimulus sensing and signal processing in bacterial chemotaxis. *Curr Opin Microbiol* **45**: 22–29.

Colin, R., Ni, B., Laganenka, L., and Sourjik, V. (2021) Multiple functions of flagellar motility and chemotaxis in bacterial physiology. *FEMS Microbiol Rev* **45**: fuab038.

Djordjevic, S., and Stock, A.M. (1998) Chemotaxis receptor recognition by protein methyltransferase CheR. *Nat Struct Biol* **5**: 446–50.

Feng, X., Baumgartner, J.W., and Hazelbauer, G.L. (1997) High- and low-abundance chemoreceptors in *Escherichia coli*: differential activities associated with closely related cytoplasmic domains. *J Bacteriol* **179**: 6714–20.

Feng, X., Lilly, A.A., and Hazelbauer, G.L. (1999) Enhanced function conferred on low-abundance chemoreceptor Trg by a methyltransferase-docking site. *J Bacteriol* **181**: 3164–71.

García, V., Reyes-Darias, J.-A., Martín-Mora, D., Morel, B., Matilla, M.A., and Krell, T. (2015) Identification of a Chemoreceptor for C2 and C3 Carboxylic Acids. *Appl Environ Microbiol* **81**: 5449–5457.

García-Fontana, C., Corral Lugo, A., and Krell, T. (2014) Specificity of the CheR2 methyltransferase in *Pseudomonas aeruginosa* is directed by a C-terminal pentapeptide in the McpB chemoreceptor. *Sci Signal* **7**: ra34.

García-Fontana, C., Reyes-Darias, J.A., Muñoz-Martínez, F., Alfonso, C., Morel, B., Ramos, J.L., and Krell, T. (2013) High specificity in CheR methyltransferase function: CheR2 of *Pseudomonas putida* is essential for chemotaxis, whereas CheR1 is involved in biofilm formation. *J Biol Chem* **288**: 18987–18999.

Gumerov, V.M., Ulrich, L.E., and Zhulin, I.B. (2023) MiST 4.0: a new release of the microbial signal transduction database, now with a metagenomic component. *Nucleic Acids Res* gkad847.

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., *et al.* (2021) Highly accurate protein structure prediction with AlphaFold. *Nature* **596**: 583–589.

Lai, R.-Z., Bormans, A.F., Draheim, R.R., Wright, G.A., and Manson, M.D. (2008) The region preceding the C-terminal NWETF pentapeptide modulates baseline activity and aspartate inhibition of *Escherichia coli* Tar. *Biochemistry* **47**: 13287–13295.

Lai, W.-C., Barnakova, L.A., Barnakov, A.N., and Hazelbauer, G.L. (2006) Similarities and differences in interactions of the activity-enhancing chemoreceptor pentapeptide with the two enzymes of adaptational modification. *J Bacteriol* **188**: 5646–5649.

Lai, W.-C., and Hazelbauer, G.L. (2005) Carboxyl-terminal extensions beyond the conserved pentapeptide reduce rates of chemoreceptor adaptational modification. *J Bacteriol* **187**: 5115–5121.

Le Moual, H., Quang, T., and Koshland, D.E., Jr. (1997) Methylation of the *Escherichia coli* chemotaxis receptors: intra- and interdimer mechanisms. *Biochemistry* **36**: 13441–8.

Li, J., Li, G., and Weis, R.M. (1997) The serine chemoreceptor from *Escherichia coli* is methylated through an inter-dimer process. *Biochemistry* **36**: 11851–7.

Li, M., and Hazelbauer, G.L. (2005) Adaptational assistance in clusters of bacterial chemoreceptors. *Mol Microbiol* **56**: 1617–26.

Li, M., and Hazelbauer, G.L. (2006) The carboxyl-terminal linker is important for chemoreceptor function. *Mol Microbiol* **60**: 469–79.

Li, M., and Hazelbauer, G.L. (2020) Methyltransferase CheR binds to its chemoreceptor substrates independent of their signaling conformation yet modifies them differentially. *Protein Sci* **29**: 443–454.

Li, M., Xu, X., Zou, X., and Hazelbauer, G.L. (2021) A Selective Tether Recruits Activated Response Regulator CheB to Its Chemoreceptor Substrate. *mBio* **12**: e0310621.

Mansfield, J., Genin, S., Magori, S., Citovsky, V., Sriariyanum, M., Ronald, P., *et al.* (2012) Top 10 plant pathogenic bacteria in molecular plant pathology. *Mol Plant Pathol* **13**: 614–29.

Martín-Mora, D., Reyes-Darias, J.-A., Ortega, Á., Corral-Lugo, A., Matilla, M.A., and Krell, T. (2016) McpQ is a specific citrate chemoreceptor that responds preferentially to citrate/metal ion complexes. *Environ Microbiol* **18**: 3284–3295.

Matilla, M.A., Gavira, J.A., and Krell, T. (2023) Accessing nutrients as the primary benefit arising from chemotaxis. *Curr Opin Microbiol* **75**: 102358.

Matilla, M.A., and Krell, T. (2018) The effect of bacterial chemotaxis on host infection and pathogenicity. *FEMS Microbiol Rev* **42**: fux052.

Matilla, M.A., Martín-Mora, D., Gavira, J.A., and Krell, T. (2021) *Pseudomonas*
aeruginosa as a Model To Study Chemosensory Pathway Signaling. *Microbiol Mol*
Biol Rev **85**: e00151-20.

Matilla, M.A., Velando, F., Martín-Mora, D., Monteagudo-Cascales, E., and Krell, T.
(2022a) A catalogue of signal molecules that interact with sensor kinases,
chemoreceptors and transcriptional regulators. *FEMS Microbiol Rev* **46**: fuab043.

Matilla, M.A., Velando, F., Tajuelo, A., Martín-Mora, D., Xu, W., Sourjik, V., *et al.*
(2022b) Chemotaxis of the Human Pathogen *Pseudomonas aeruginosa* to the
Neurotransmitter Acetylcholine. *mBio* **13**: e0345821.

Monteagudo-Cascales, E., Ortega, Á., Velando, F., Morel, B., Matilla, M.A., and Krell,
T. (2023) Study of NIT domain-containing chemoreceptors from two global
phytopathogens and identification of NIT domains in eukaryotes. *Mol Microbiol* **119**:
739–751.

Okumura, H., Nishiyama, S., Sasaki, A., Homma, M., and Kawagishi, I. (1998)
Chemotactic adaptation is altered by changes in the carboxy-terminal sequence
conserved among the major methyl-accepting chemoreceptors. *J Bacteriol* **180**:
1862–8.

Ortega, Á., and Krell, T. (2020) Chemoreceptors with C-terminal pentapeptides for
CheR and CheB binding are abundant in bacteria that maintain host interactions.
Comput Struct Biotechnol J **18**: 1947–1955.

Ortega, D.R., Fleetwood, A.D., Krell, T., Harwood, C.S., Jensen, G.J., and Zhulin, I.B.
(2017) Assigning chemoreceptors to chemosensory pathways in *Pseudomonas*
aeruginosa. *Proc Natl Acad Sci USA* **114**: 12809–12814.

Parkinson, J.S., Hazelbauer, G.L., and Falke, J.J. (2015) Signaling and sensory
adaptation in Escherichia coli chemoreceptors: 2015 update. *Trends Microbiol* **23**:
257–66.

Perez, E., and Stock, A.M. (2007) Characterization of the *Thermotoga maritima*
chemotaxis methylation system that lacks pentapeptide-dependent methyltransferase
CheR:MCP tethering. *Mol Microbiol* **63**: 363–78.

Reyes-Darias, J.A., Yang, Y., Sourjik, V., and Krell, T. (2015) Correlation between
signal input and output in PctA and PctB amino acid chemoreceptor of *Pseudomonas*
aeruginosa. *Mol Microbiol* **96**: 513–525.

Russo, A.F., and Koshland, D.E. (1983) Separation of signal transduction and
adaptation functions of the aspartate receptor in bacterial sensing. *Science* **220**:
1016–1020.

Sambrook, J., Fritsch, E.F., and Maniatis, T. (1989) *Molecular Cloning: A Laboratory*
Manual, 2nd edn. Cold Spring Harbor Laboratory Press, New York, NY, USA, .

Schrodinger, L. (2010) The PyMOL Molecular Graphics System, Version 1.3r1. *New*
York: Schrödinger, LLC .

Shiomi, D., Okumura, H., Homma, M., and Kawagishi, I. (2000) The aspartate
chemoreceptor Tar is effectively methylated by binding to the methyltransferase
mainly through hydrophobic interaction. *Mol Microbiol* **36**: 132–40.

Toth, I.K., Bell, K.S., Holeva, M.C., and Birch, P.R. (2003) Soft rot erwiniae: from
genes to genomes. *Mol Plant Pathol* **4**: 17–30.

Uchida, Y., Hamamoto, T., Che, Y.-S., Takahashi, H., Parkinson, J.S., Ishijima, A., and
Fukuoka, H. (2022) The Chemoreceptor Sensory Adaptation System Produces
Coordinated Reversals of the Flagellar Motors on an Escherichia coli Cell. *J
Bacteriol* **204**: e0027822.

Velando, F., Gavira, J.A., Rico-Jimenez, M., Matilla, M.A., and Krell, T. (2020)
Evidence for Pentapeptide-Dependent and Independent CheB Methylesterases. *Int J
Mol Sci* **21**: E8459.

Velando, F., Matilla, M.A., Zhulin, I.B., and Krell, T. (2023) Three unrelated
chemoreceptors provide *Pectobacterium atrosepticum* with a broad-spectrum amino
acid sensing capability. *Microb Biotechnol* **16**: 1548–1560.

Wu, J., Li, J., Li, G., Long, D.G., and Weis, R.M. (1996) The receptor binding site for
the methyltransferase of bacterial chemotaxis is distinct from the sites of
methylation. *Biochemistry* **35**: 4984–93.

Wuichet, K., and Zhulin, I.B. (2010) Origins and diversification of a complex signal
transduction system in prokaryotes. *Sci Signal* **3**: ra50.

Yamamoto, K., and Imae, Y. (1993) Cloning and characterization of the *Salmonella
typhimurium*-specific chemoreceptor Tcp for taxis to citrate and from phenol. *Proc
Natl Acad Sci U S A* **90**: 217–21.

Yi, X., and Weis, R.M. (2002) The receptor docking segment and S-adenosyl-L-
homocysteine bind independently to the methyltransferase of bacterial chemotaxis.
Biochim Biophys Acta **1596**: 28–35.

Zatakia, H.M., Arapov, T.D., Meier, V.M., and Scharf, B.E. (2018) Cellular
Stoichiometry of Methyl-Accepting Chemotaxis Proteins in *Sinorhizobium meliloti*. *J
Bacteriol* **200**: e00614-17.

Fig. 1

A

```
ECA_RS00895  ...LADAVSAFKIPSYSHGNAGSYESVPMSTPTLSLALARKE
ECA_RS00900  ...LAEAVSTFKLLSYGNGKTASYASAPTRTPTLSLAPAAAKNQSNNDNWTTF
ECA_RS08780  ...LARTVSVFNLGASYKSAALNRKTETPALAAPKNNRAEKTSAKGELADWTTF
ECA_RS10160  ...LSHAVAAFRI
ECA_RS18000  ...LAETVSQFRLGNGHQIARTPAAAASLTLRPALAAPGKSGISAGEGDWTSF
ECA_RS21440  ...LVELMKVFIVEGGSSQRIAPPLKRPSSAKFSLANPKGSAGSNNQNWEQF
ECA_RS21445  ...LVELMKVFIVESGSSQRTTPELKRPSSAKLSLASPKGRTKSDSQNWETF
ECA_RS20365  ...LVTLMNHFHLRGTPAARPAPMAKKAQTARLALAPVGNTQDNWEKF
ECA_RS21450  ...LLELMGVFKLNGIQTKAPRLTSQVKQPAAPRLALASKSGHTSSDNWETF
ECA_RS21455  ...LVELMGVFKIDGTQSQRAVPQVTTLSRPKLALAGNSSNTNWETF
ECA_RS12390  ...LNEQAQELSRTVEQFRVDESAGSYLALGAR
ECA_RS02220  ...LRDAVRFFKVNQDHLLRIH
ECA_RS06345  ...LQNAVEVFKINQAVAQEHRAASASSLAALPKSLLPKPTSAGSSNANWETF
ECA_RS08370  ...LNQAVAVFRLSEDTGSFRRTTQATAGQKPVLLAPSVNGGKKAKEGSSTDNWETF
ECA_RS08330  ...LTRAVATFKLSSHLSSGHSAPARPNALAAKGRSSLALPRQANTENGNWETF
ECA_RS19280  ...LTQAVAVFKLSGIVQQVRSSLPKSAPQPRLAPAMAIAGSSKGNSNQNWETF
ECA_RS18955  ...LNLGVSRFHLM
ECA_RS06625  ...LNQTVSLFQLSDTQSALQVAAKPVRKAQAIAPRAGKALPTSSDNWEKF
ECA_RS07510  ...LSAVVDVFNLRDSDSDQQTAFSRPAIAAPVHRAVAQSTTPLLSVHGRHGEGWEKF
ECA_RS13300  ...LTEAVSVFQLSAAEAPRRPQQRLAEKAPAAQKPMLLAAAGGKKGNANDNWETF
ECA_RS12635  ...LESMVANFRLSENEGRKPKANISGLPPQQKYLPPAAKQTQDSGWTTF
ECA_RS12640  ...LEKLLEHFRVSQSDNRVASRASSSIPRHTLPKSVSAKAASSESDWTSF
ECA_RS00400  ...LATLMSVFRISDKDVARLQGSNTGNPNSGNKATARLPTLASRDNGNDNWTTF
ECA_RS00455  ...LKQAVSVFRLANAQHDDTPAGIAFNNQPHRLHAPR
ECA_RS10935  ...LAQTIEHFRLEQQHALPHALLR
ECA_RS17750  ...LAESMVQFKVQSQEFAAIGRF
ECA_RS17910  ...LMRSMALFQVEPRLS
ECA_RS02005  ...LQQSVSRFQIARENREMDNVLPGLRQNITLADAR
ECA_RS15955  ...LSQLVGQFIVGQIASSSLIPALASVPSGLSAPRLASAKNKNALAQDEAGWQRF
ECA_RS05475  ...LNSVVGAFRV
ECA_RS02210  ...MESLVSHFKVDDSAPQQPLQHALLSR
ECA_RS09870  ...MSEAVSVFSIPR
ECA_RS09875  ...MVQAASVFSLSR
ECA_RS17685  ...LNSAINVYGS
ECA_RS17860  ...LSELVSVFRI
ECA_RS11380  ...LNTSVSLFILPSTEADVNPMIDQREMTQRIPMMG
```

B

ECA_RS12635 ECA_RS21440 ECA_RS08370

N609 E521 S518

F561

F559

F644

Fig. 2

Fig. 3

Fig. 4

Fig. 5