Phage resistance is associated with density increases in the soil bacterium

***Variovorax* sp. and results in changes to community structure**

Authors: Meaghan Castledine[1], Daniel Padfield[1], Rai Lewis[1] & Angus Buckling[1]

[1]College of Life and Environmental Sciences, Environment and Sustainability

Institute, University of Exeter, Penryn, Cornwall, TR10 9EZ, U.K.

Corresponding author: Meaghan Castledine (mcastledine96@gmail.com)

Author contributions: MC and AB conceived and designed the study. Experiments

conducted by MC. DNA extractions for population sequencing conducted by RL.

Bioinformatics conducted by DP and MC. Data analysis conducted by MC. All

authors contributed to the writing of the manuscript.

Competing Interests: We have no competing interests

Acknowledgements: We thank Stineke van Houte and Michael Brockhurst for

helpful feedback on a manuscript draft. This work was funded by grant no.

MR/N0137941/1 for the Great West 4 BIOMED Medical Research Council Doctoral

Training Partnership, awarded to the Universities of Bath, Bristol, Cardiff, and Exeter

from the Medical Research Council/UK Research and Innovation, awarded to MC.

This work was supported by NERC awards NE/V012347/1 and NE/S000771/1

awarded to AB. DP is funded by a NERC independent research fellowship

(NE/W008890/1).

29
Data availability: All data and R code used in the analysis are available on GitHub
(https://github.com/mcastledine96/Variovorax_phage_resistance_increases_density_
2024). Assemblies and sequencing data are available from ENA (Project Accession
PRJEB74093).

Abstract

Lytic bacteriophages ('phages') can limit bacterial densities, either directly through

lysis or indirectly through costs to resistance. However, phages have also been

reported to have no, and in some cases even positive, effects on host densities.

Here, we investigate the mechanisms behind an increase in host density in

Variovorax sp. populations following a fixation of resistance that was maintained

after phage extinction. Our results demonstrate that the density increase was a

genetic trait coinciding with resistance emergence. Growth curves showed that

phage resistance shifted population growth curves such that density was higher in

the death-phase. This density-increasing effect of resistance had important

implications for community structure with phage resistant *Variovorax* decreasing the

density of a conspecific. That resistance to lytic phage can increase host densities

has implications for wider ecology and phage therapy where lytic phages are

presumed to have negative effects on their hosts.

Introduction

Bacteriophages are hypothesised to have major impacts on structuring microbial communities, and in regulation of wider nutrient cycles (1–3). Part of this assumption comes from the ubiquity of bacteriophages (4). By lysing their hosts, phages can suppress fast-growing bacteria densities and give a relative advantage to slow-growing or rarer microbial taxa (3). Phages are shown to lyse algal blooms, resulting in crashes in algae abundances and nutrient release which benefits the surrounding community (1, 5, 6). Furthermore, a common assumption of many models is that phage resistance is costly (3), which leads to suppression of dominant or previously-fast growing bacteria populations (3, 7).

However, the effects of phage resistance on host phenotypes, and the associated cost, is highly system specific. If evolving phage resistance typically involves modification of a receptor essential to host growth, resistance is likely to be associated with a growth rate cost (8). For example, to become resistant to OMKO1, *Pseudomonas aeruginosa* mutates an efflux pump required for drug resistance (9). As a result, while antibiotics are present, resistance to OMKO1 imposes a cost to *P. aeruginosa* (9). Similarly, phage resistance involving mutations to flagella or pili are further shown to slow bacteria growth rates (10, 11). However, there are numerous examples of phage resistance being non costly to growth including in *Pseudomonas syringae* (12), *Synechococcus* (13), and *P. aeruginosa* (14).

Phage resistance may also indirectly increase host fitness (beyond the direct effect of resisting phage). For instance, increased biofilm production is a phage resistance mechanism which physically blocks phage access to bacteria cells (15–17). Biofilm

production has consequential fitness benefits in resisting wider environmental stress

such as antibiotics (18), and can be advantageous in competition (19). Phages can

also accelerate the molecular evolution of their hosts and select for hypermutator

strains, therefore giving increased adaptive potential to secondary stressors (20, 21).

P. fluorescens phage has been shown to increase host densities in soil, although

only when grown in monoculture (in the absence of a soil community) (22). Here,

phages are hypothesised to have increased bacterial densities by reducing

intraspecific competition through slowing of growth rates or facilitating diversification

into separate ecotypes (22), although the exact mechanism is unknown.

Understanding how phages can increase bacteria densities is important for phage

therapy where increasing bacteria densities would be undesirable (15). Furthermore,

investigating this effect has important implications for our understanding of how

phages shape microbial communities where, in nature, phages are presumed to

have predominantly negative effects their host densities (1, 3). In a previous study,

we noted that densities of *Variovorax* sp. increased following phage exposure and

fixation of phage resistance (23) even when phage had been driven extinct. Here, we

investigate the mechanism underpinning phage-mediated increases in *Variovorax*

densities, and the consequence this has for the composition of a three species

synthetic bacteria community. We show that phage resistance is associated with

density increases resulting from a shift in growth curves that led to higher densities in

the death-phase. Phage resistance had ramifications for community structure with

decreases in the proportion of another community member which is parasitised by

Variovorax.

Results

Phage resistant clones have higher densities than phage susceptible clones

In a previous study (23), we evolved *Variovorax* and its lytic phage in the presence and absence of two other bacteria-phage pairs (Figure 1). We observed an interesting interaction with *Variovorax* densities increasing after two weeks of phage exposure (phage present: $\bar{x}$ = 10$^{7.79}$ CFU/mL, 95%CI = 10$^{7.74}$ - 10$^{7.85}$; phage absent: $\bar{x}$ = 10$^{7.69}$ CFU/mL, 95%CI = 10$^{7.63}$ - 10$^{7.74}$; ANOVA comparing models with and without phage: χ^2_1= 7.21, p = 0.007; Tukey HSD: p = 0.013). This coincided with a fixation of phage resistance in polyculture (90.3%; SE ±6.6) and monoculture (100%), and extinction of phage after two weeks. As such, we investigated the hypothesis that phage resistance has led to increased *Variovorax* densities.

111 Figure 1. Overview of experimental design in which *Variovorax* and its phage were

112 evolved in monoculture and polyculture, and assays performed to understand the

mechanism of this density increase. Sequencing analyses and phenotypic assays aimed to determine the mechanism by which phages led to increased *Variovorax* densities across experimental evolution. Created in BioRender. Castledine, M. (2024)

BioRender.com/c73o264.

To assess whether higher densities in phage-exposed populations was likely a result of a genetic change (as opposed to epigenetic or physiological), we measured the densities of individual clones (isolated from weeks two (pre phage loss) and four (post phage loss) from monoculture treatments) grown in isolation in common garden experiments (no phage or other clones present so any density effects are purely heritable). All clones isolated from phage treatments were phage resistant as phage resistance reached fixation in monoculture, while all clones from the no-phage treatment were susceptible. Consistent with the population dynamics during experimental evolution, clones isolated from phage-exposed replicates had 66% higher densities ($\bar{x} = 10^{7.51}$ CFU/mL, 95%CI = $10^{5.60}$ - $10^{9.41}$) than phage-unexposed clones ($\bar{x} = 10^{7.29}$ CFU/mL, 95%CI = $10^{5.38}$ - $10^{9.19}$; ANOVA comparing models with and without phage: $\chi^2_1 = 6.60$, p = 0.102; Tukey HSD: p = 0.018; Figure 2). Additionally, there was a significant independent effect of time with clones from week four ($\bar{x} = 10^{7.49}$ CFU/mL, 95%CI = $10^{5.30}$ - $10^{9.68}$) reaching higher densities than clones from week two ($\bar{x} = 10^{7.31}$ CFU/mL, 95%CI = $10^{5.12}$ - $10^{9.49}$; Tukey HSD: p < 0.001; Figure 2). Overall, these results suggest effects imposed by phage were hereditary as phage did not need to be actively present for effects to be observed.

Figure 2. Common garden experiments showing the density of individual clones (points) which have been isolated from evolution lines where phage have been absent or present. Tops and bottoms of the bars represent the 75th and 25th percentiles of the data, the middle lines are the medians, and the whiskers extend from their respective hinge to the smallest or largest value no further than 1.5* interquartile range. Points represent individual clones.

Fitness advantage is associated with resistance mutations

Given the consistent higher growth rate that phage exposed populations and phage resistant clones had over phage absent populations and phage susceptible clones,

we looked at whether there were underlying genetic changes that could explain this.

We sequenced 3 resistant and 3 sensitive clones coexisting within 3 polyculture

treatment replicates. Of the phage resistant clones, 2/3 clones had a missense

mutation at the same locus (gene ID: 01575) and 1 had a substitute mutation at a

nearby locus (gene ID: 01584; 10.4k base pairs downstream; this was also the only

mutation in this clone) that was not found in phage susceptible isolates. No other

mutations were found at the same loci across clones. While identified in a

hypothetical protein, the predicted protein based on amino acid sequence of 01575

suggests that the mutation occurred in a gene responsible for metabolism or a

membrane protein. Of matches related to microorganisms (excluding comparisons to

plant and human proteins), glycosyltransferase (97% coverage to *Saccharomyces*

cerevisiae with 99.4% model confidence; 83% coverage to *Staphylococcus aureus*

with 100% model confidence) was identified. 17/20 of the top protein predictions

were to transferase enzymes, suggesting that this mutated protein likely belongs to

this family of proteins. Glycosyltransferases are a broad group of enzymes

responsible for transferring sugars to acceptor molecules which, in bacteria, is

important for signalling, cell wall biosynthesis and adhesion (24). In *Klebsiella*

pneumoniae, mutation in the glycosyltransferase enzyme responsible for

exopolysaccharide synthesis resulted in phage resistance (25). Therefore, these

results would suggest phage resistance occurred via mutation at genes which

influence host metabolism and potentially cell structure. Single point mutations and

receptor loss are consistent with surface modification forms of resistance (8, 26).

Generally, few mutations were found, with one resistant clone having four mutations

while the other two had only one mutation each. Similarly, one phage susceptible

clone had two mutations while the other two had one mutation. 01584 sequence

(mutation found in 1/3 resistant clones) had no comparable genes or protein

structures in BLAST or Phyre2, however it's proximity to the other genes could imply

a similar function (27, 28).

We sequenced whole populations taken from seven weeks of evolution in

monoculture (*Pseudomonas* and *Ochrobactrum* absent), with the aim of identifying

further genetic changes linked to phage resistance and increased density. In total,

we found 470 genetic variants, with 239 shared between resistant and susceptible

populations. Phage resistant populations had 117 unique variants (within 60 genes)

while phage susceptible populations had 114 unique variants (within 64 genes). Most

variants were found at low frequency (96.8% of variants at <50% frequency) with

resistant and susceptible populations having on average 147 variants. 5/6 phage

resistant populations had at least one unique genetic variant at high frequency

(>50%) compared to only 2/6 of the phage susceptible populations, suggesting

specific mutations were selected by phage. While we did not observe mutations in

01575 (mutation found in 2/3 resistant clones above), mutations were found in 01584

(mutation found in 1/3 resistant clones) in 4/6 phage resistant populations (92.7%,

68.2%, 12.3%, 9.6%). Mutations selected to high frequency were also found in *prpR*

(69.3% in 1 population; transcription regulator gene) and in five other hypothetical

proteins or intergenic regions (Figure 3). The lack of convergence suggests phage

resistance may be maintained by several different mechanisms that have the same

phenotypic effects (resistance and higher population density). As a result, we did not

find phage presence significantly differed in genetic composition (PERMANOVA:

$F_{1,10} = 0.889$, p = 0.81, Figure S1) or distance from the ancestral genome (ANOVA:

$F_{1,10} = 0.547$, p = 0.477). Overall, evidence from fixation of resistance (100% clones

tested being resistant to phage), unique mutations and common garden experiments

suggest a link between resistance and phenotypic increases in population density;

however, mechanistic links cannot be made between genotypes and phenotypes

owing to the lack of convergent evolution.

Figure 3. The frequency of genetic variants found in phage resistant but not

phage susceptible populations. Where multiple variants are found in the same

gene, the average has been presented. Each replicate population is identified on the

x-axis. Genes of known function are named on the y-axis while hypothetical proteins

are given a numerical denomination. Gene names separated by a forward slash

indicate mutations in intergenic regions. Samples indicate each of the six replicate

populations that evolved in the presence of phage.

Phage resistant clones do not have a fitness advantage

We next determined whether phage resistance conferred a fitness advantage in the

absence of phage. In competition assays with a marked ancestral strain, relative

fitness was not significantly affected by phage resistance ($F_{2,15}$ = 0.460, p = 0.640;

Figure 4). The greater density associated with resistance did not therefore result in a

competitive fitness advantage.

Figure 4. Relative fitness of ancestral, phage susceptible and resistant isolates against a marked ancestral strain of *Variovorax*. Points represent independent replicates. Tops and bottoms of the bars represent the 75th and 25th percentiles of the data, the middle lines are the medians, and the whiskers extend from their respective hinge to the smallest or largest value no further than 1.5* interquartile range.

Phage resistance affects growth dynamics

We next investigated if density benefits resulting from resistance arose from changes in growth dynamics. This may have arisen if, for example, logistic growth rates of resistant isolates were slowed, resulting in resources becoming less exhausted when cultures were transferred to fresh media on day seven. As such, we measured the

growth curves of *Variovorax* populations isolated from week seven of experimental

evolution over one week. Here, there was a significant interaction between treatment

and day of measurement (ANOVA comparing models with and without the

interaction: χ^2_1 = 7.25, p = 0.007), suggesting phage resistance had altered

population growth dynamics (Figure 5). Consistent with our previous observation

based on CFU counts (Figure 2), optical density after one week was higher in phage

resistant ($\bar{x}$ = 0.067, 95%CI = 0.062 - 0.072) than susceptible ($\bar{x}$ = 0.057, 95%CI =

0.053 - 0.062) populations (Figure 5). Consistent with an absence of fitness

differences between susceptible and resistant isolates (Figure 4), there was no

significant difference in exponential growth rate measured in the 24 hr growth rate

assay ($F_{1,9}$ = 0.622, p = 0.451) or carrying capacity in the one week growth rate

assay (phage resistant: $\bar{x}$ = 0.08, 95%CI = 0.075 − 0.085; phage susceptible: $\bar{x}$ =

0.076, 95%CI = 0.072 − 0.081; Figure 5). Phage resistant isolates show similar

growth dynamics to susceptible isolates, except they maintain higher densities

during the death phase at day 7.

Figure 5. The seven-day growth curves of (a) phage susceptible and (b) phage resistant *Variovorax* populations that had evolved for seven weeks. Points represent individual replicate populations measured repeatedly through time. Solid lines represent the mean predictions, and shaded bands represent the 95% confidence intervals of the best fitted model.

The supernatant of phage resistant populations is not less toxic

If bacterial populations can maintain a higher density during the stationary/death phase, this suggests that bacteria are potentially producing fewer toxic metabolites and/or are less susceptible to them. We investigated these possibilities by growing week seven populations in spent supernatants from either their own treatment or the opposite treatment. However, population density was non-significantly different between populations irrespective of prior phage exposure (= 1.48, p = 0.224) or

supernatant type (= 2.12, p = 0.145; interaction between effects: = 0.148, p =

0.7004; Figure 6).

Figure 6. Density changes when populations which are (a) phage susceptible

and (b) phage resistant, are grown in supernatant isolated from either

population. E.g. phage susceptible growing in supernatant from phage susceptible

populations (same evolutionary history) or phage susceptible growing in supernatant

from phage resistant populations (different evolutionary history). Points represent

densities of individual replicates while lines indicate the change when that same

population is grown in a different supernatant.

Indirect benefit of phage resistance impacts community structure

Considering that phage resistance increased *Variovorax* densities, we considered

whether this had implications for the structure of a simple microbial community. We

had previously cultured *Variovorax* with two other species, *Ochrobactrum* and

Pseudomonas, from which it derives a fitness benefit to the other species' detriment

(29). As such, we predicted that phage resistance increasing *Variovorax*'s density

may have negative implications for the other two species. To this end, we re-

assembled communities with different *Variovorax* isolates with ancestral

Ochrobactrum and *Pseudomonas*. Consistent with this prediction, the relative

proportion of *Pseudomonas* in the community was significantly lower with phage

resistant *Variovorax* ($\bar{x}$ = 0.092, 95%CI = 0.08 − 0.11) compared to ancestral ($\bar{x}$ =

0.143, 95%CI = 0.128 − 0.158; Tukey HSD comparing *Pseudomonas* proportion with

ancestral vs resistant *Variovorax*: estimate = 0.495, z-ratio = 5.02, p < 0.001) or

phage susceptible *Variovorax* ($\bar{x}$ = 0.128, 95%CI = 0.114 − 0.143; ANOVA

comparing models with and without interaction between *Variovorax* phage resistance

and species identity: χ_4^2 = 34.1, p < 0.001; Tukey HSD comparing *Pseudomonas*

proportion with resistant vs susceptible *Variovorax*: estimate = -0.367, z-ratio = -3.64,

p < 0.001; Figure 7). *Pseudomonas*'s relative proportion was non-significantly

different when cultured with ancestral or phage susceptible *Variovorax* (Tukey HSD:

estimate = 0.129, z-ratio = 1.403, p = 0.339). Neither *Ochrobactrum*'s or *Variovorax*'s

relative proportion was significantly affected by phage resistance in *Variovorax*

(Tukey HSDs > 0.05, Table S1; Figure 7).

Figure 7. The relative proportion of each species in replicate communities when cultured with ancestral, phage resistant or phage susceptible *Variovorax* isolates. Points represent individual treatment replicates. Tops and bottoms of the bars represent the 75th and 25th percentiles of the data, white lines indicate the medians, and whiskers extend from their respective hinge to the smallest or largest value no further than 1.5× the interquartile range.

Discussion

Bacteriophages have been previously shown to positively influence bacteria densities, but only when propagated in monoculture and the mechanism was not determined (22). Here, we showed that evolved resistance to phage changed the

growth curves of *Variovorax* populations resulting in higher densities during the

death phase. High levels of resistance, and the associated positive effect on density,

was observed for several weeks past phage extinction within monocultures (no other

bacterial species present) and in a community context. That this effect was

consistent in both the presence and absence of other species is particularly

interesting as the growth of this species is enhanced by the presence of the other

two species (30). As such, irrespective of the metabolic state the bacteria are in

(feeding on the media or on supernatant), the phage had a positive effect on

Variovorax density. This effect had consequences for community structure with

phage resistance in *Variovorax* suppressing *Pseudomonas* in communities. These

results demonstrate that phage resistance can give some species a relative

advantage against others in community contexts, contrary to the prevailing wisdom

of phage community ecology (3).

Phage resistance did not have a significant effect on exponential growth rate but did

affect the death phase, with phage resistant populations maintaining a higher

density. Although we did not find evidence that this was linked to a change in

metabolites produced after one week, it may reflect a broader ability of the bacteria

being able to cope in stressful conditions. *Variovorax* is a metabolically plastic

species and is capable of being lithoautotrophic or chemoorganotrophic, including

the metabolism of toxic or complex compounds (31). Benefits from phage-exposure

were not linked directly to a single phage resistance mutation, however resistance

was found to likely emerge via metabolism for a surface protein (glycosyltransferase

related (25)), often with missense mutations resulting in a loss of function. Loss of

this receptor or surface structure could have shifted *Variovorax* metabolism to give

the effects shown. However, confirming this is challenging due to the complexity of this nutrient medium. Additionally, population sequencing revealed multiple mutations emerging and several reached high frequencies (>50%) in 5/6 phage treatment populations, although with a lack of convergent evolution. All high frequency genetic variants were in hypothetical proteins, making mechanistic inferences impossible within this project. Interestingly, only two no-phage culture had a unique mutation that reached high frequency (>50%) compared to 5/6 phage cultures, suggesting phage presence increased the likelihood of variants reaching a high frequency.

While resistance was clearly beneficial in the presence and absence of phage, reaching and being maintained at very high frequencies, we did not find any phage resistance emerge in phage-free controls across eight weeks. This was presumably because there was no selective benefit of phage resistant mutants competing against phage sensitive clones in the absence of phage, and hence resistant mutations would not have increased in frequency. Nevertheless, we may expect this resistance-linked growth phenotype to be favoured in highly structured metapopulations, where limited dispersal can result in different patches seeded by single clones with the most productive patches (i.e. those seeded with resistant clones) producing more individuals to colonise new patches (32). The resistance mutations may also allow populations to subsequently explore alternative evolutionary trajectories in the longer term (17).

Importantly, phage resistance increasing *Variovorax* densities shifted community structure, resulting in lower proportions of *Pseudomonas. Variovorax* has been

shown in previous work to benefit from the presence of *Pseudomonas* to

Pseudomonas' detriment (29). Increasing densities of phage resistant *Variovorax*

therefore likely resulted in increased costs experienced by *Pseudomonas* in

coculture. Phage lysis and phage resistance are typically assumed to be most costly

to dominant community members (3). However, many examples exist of non-costly

phage resistance and species dominating communities with high levels of phage

resistance (33, 34). Consequently, our results in-part support the king-of-the-

mountain' hypothesis in which dominant community members are more able to

evolve resistance owing to greater population sizes (33, 35) – albeit in this case

likely by mutation than horizontal-gene-transfer (starting population isogenic).

Synonymously, the 'king-of-the-mountain' hypothesis also relies on a low trade-off in

defence and competition (36), evident in *Variovorax*.

Few examples exist of phages improving bacteria densities (22), suggesting this

observation may not be ubiquitous. However, much research has heavily focussed

on model strains of bacteria and phage which are unlikely to be generalisable to all

bacteria and phages (23, 34). Furthermore different mechanisms of phage

resistance have the potential to improve bacteria densities in specific contexts. For

example, phages can select for increased biofilm production (14, 37) and persister

populations (38) which can further confer resistance to stressors such as antibiotics

and immune cells (18, 39). If phages can improve the density of soil bacteria, such

as by driving host diversification (40), this may have positive impacts on carbon

cycling and crop yield (41). *Variovorax* species, for instance, are symbionts to plants

and are found in the rhizosphere where they can improve plant growth (31).

Alternatively, if phages are used therapeutically (42, 43) then density enhancement

of target bacterial pathogens is clearly detrimental. Work considering wider bacteria

and phage pairs, beyond model systems, will allow greater generalisations to be

made and understanding of how phages can affect their hosts in the short and long-

term.

The influence of phages on bacterial populations are highly diverse, making

predictions and generalisations challenging from model systems to phages in nature.

Our work highlights the surprising outcomes that can arise from bacteria-phage

interactions, and that the simple assumption that lytic phages negatively affect host

densities may not always be true. Wider research analysing the effects of different

phages on host bacteria, both during and after phage extinction would give important

insight into how phage therapy may affect pathogens during infection and how

phages operate in nature.

Materials and Methods

Experimental evolution

In a previous study, species *Ochrobactrum* sp., *Pseudomonas* sp., and *Variovorax*

sp. were grown as monocultures and as a three-species polyculture with and without

their respective phage (*Ochrobactrum* phage ORM_20, *Pseudomonas* phage

CH7FMC, *Variovorax* phage VAC_51) (23). Experimental treatments included:

monoculture, monoculture with phage (each bacterial species individually with only

its species-specific bacteriophage), polyculture (all three bacteria present) and

polyculture with phage (all three bacteria with all three phages present) (Figure 1).

Each treatment was replicated six times. Experimental set-up is described in (23).

Briefly, cultures were established using isogenic strains of each bacteria and phage,

with phage added at an MOI of 0.01 (to phage present treatments). Cultures were grown in 1/64 TSB (tryptic soy broth diluted in demineralised H_2O), at 28 °C and transferred weekly to fresh medium. Bacteria and phage densities were estimated every two weeks. For *Variovorax* cultures, we estimated phage resistance in week two and eight monocultures (phage present and absent). Phage resistance was analysed using spot assays with ancestral phage.

Clonal density measures

We first wanted to assess whether higher densities conferred to phage-exposed populations was a heritable trait observable in single clones picked and regrown in the absence of phage (common garden experiments). From week two and four cultures (the weeks preceding and following higher densities), twelve colonies were isolated from each replicate from monoculture treatments. Colonies were individually grown in 1/64 TSB and density normalised as before. To fresh 6 mL 1/64 TSB, ~10^6 CFUs of each clone were added to individual vials. Cultures were grown for 1 week, frozen, and plated from frozen onto KB agar as previous.

Clonal and population sequencing and bioinformatics

To gain insight into the mechanism of resistance and whether this could be linked to changes in bacterial density and growth rate, isolates were sent for whole genome sequencing (WGS). Three phage resistant and three phage susceptible clones from three independent week two polyculture lines were selected for WGS. Polyculture lines were selected as these were the only treatment which had phage susceptible and resistant clones coexisting (100% resistance was found in monoculture lines). This allowed for direct comparison between different clones that have evolved from

the same starting population but only differed in their resistance profile. DNA

extractions were done on each isolate by MicrobesNG using in-house protocols

(MicrobesNG Sequencing Service Methods, 2021). WGS was performed by

MicrobesNG using an Illumina Novaseq 6000 to create 250 base pair (bp) paired-

end reads. In-house processing by MicrobesNG trimmed the adapters and removed

reads with a quality <15.

To further understand genetic changes that resulted in increased growth rates of

phage resistant *Variovorax*, cultures from week 7 monoculture lines of evolution

were sent for population sequencing. Overnight cultures (120 µL frozen stock in 1/64

TSB grown shaking at 28 ºC. 1.8 mL was spun at 16000 x g for 5 mins (Progen

GenFuge 24D centrifuge). The supernatant was removed and another 1.8 mL of

fresh sample was added to each tube and centrifuged for a further 5 mins and

supernatant removed. The Qiagen DNeasy Ultra Clean DNA Extraction Kit reagents

and protocol were used to extract the gDNA from the cultures. The concentration of

gDNA in each sample was determined using the QuBit dsDNA HS Assay kit regents

and protocol. Sequencing was done by the University of Liverpool's Centre for

Genomic Research (CGR) using an Illumina Novaseq to create 150bp paired end

reads. The CGR trimmed for the presence of Illumina adapter sequences using

Cutadapt (v1.2.1) and removed reads with a minimum quality score of 20 and which

were shorter 15bp in length.

For both clonal and population sequencing, we further filtered the trimmed reads to

remove Illumina adapter sequences and for reads with a quality score below 10

using TrimGalore (v0.6.4_dev). We used the breseq pipeline (v0.36) – a

computational tool for analysing short-read DNA data - to map the trimmed reads to

a high quality reference genome of *Variovorax* sp. (GenBank accession:

GCA_037482385.1) and predict genetic variants (44). As our reference genome has

two contigs, we added the parameters "-c" and "—contig-reference" so that all the

sequences are assumed to have the same average coverage and coverage

distribution. In the clonal sequencing, 97.6% of reads mapped to the reference

genome (minimum of 95.4%), giving on average ~876,989 reads per replicate

(minimum 444,292) with an average coverage of 62.21 (minimum of 31.84). We only

investigated genetic variants that were at a frequency of 1 (we do not expect

polymorphisms in any of the predicted genetic variants). For the population

sequencing, we ran breseq in polymorphism mode (-p). Variants with <5% frequency

in the population samples were removed.

When mutations unique to resistant clones appeared in a hypothetical protein, we

aimed to understand the potential role of this protein by simulating its structure. The

amino acid sequence of the protein was run through Phyre2 (Protein

Homology/analogY Recognition Engine V 2.0) (45). Results were assessed based

on whether proteins were homologous to microbial proteins, and the most common

matches up to the first 20 modelled proteins.

Phage resistance and competition assay

We determined if phage resistance conferred a growth rate benefit to *Variovorax*,

thereby leading to higher bacterial densities. Twelve clones per replicate from week

two monoculture-lines were isolated and grown in 96-well plates with 150 µL TSB.

Phage resistance was determined by streaking isolated colonies against the

ancestral phage (streak assay). Week two isolates were chosen as this time-point

preceded changes in bacterial density from phage exposure and phage extinction.

We elected for phage resistant clones generated during experimental evolution

(instead of selecting spontaneous mutants) in the hope that resistance

mutations/mechanisms (and therefore phenotypic effects) were consistent with those

from the evolution experiments. One phage resistant clone from the phage

resistance assays was chosen for each replicate (n = 6), and one clone from each

no-phage treatment replicate was isolated as a phage susceptible control (n = 6).

Each colony was grown individually in 1/64 TSB individually as described previously.

In addition, marked (ancestor-cp; chloramphenicol resistance (46)) and unmarked

ancestral *Variovorax* sp. were grown. Cultures were diluted to normalise densities as

described previously. The unmarked ancestral strain and each phage resistant and

susceptible colony was competed against the marked ancestral strain at equal

starting densities (~3 x 10^6 CFUs of each competitor inoculated (30 μL diluted

culture)). Diluted cultures used in inoculation were frozen and plated on KB agar as

above for starting density estimation. Cultures were grown for one week as previous

and plated from frozen onto KB and KB-chloramphenicol (25 μg/mL) agar. Relative

fitness was calculated from the ratio of the estimated Malthusian parameters,

$m_{resistant}$, $m_{susceptible}$ or $m_{ancestor}$:$m_{ancestor-cp}$, which were calculated as $m = \ln(N_1/N_0)$,

where N_1 is the final density and N_0 is the starting density (47).

Growth curves

Next, we were interested if phage resistance had shifted the growth curve of

Variovorax populations thereby leading to higher densities at transfer. 120 μL from

each monoculture line from week seven were revived in 6 mL growth medium and

grown shaking for two days at 28 ⁰C. Densities were normalised as previous and 10

µL was inoculated into fresh growth medium. Every 24 hrs for one week, cultures

were vortexed and optical density (OD_{600}; Multiskan Sky) was measured from a 200

µL sample.

To characterise the log/exponential growth phase of phage resistant and susceptible

cultures at a higher resolution, week seven cultures were revived as previous and 20

µL of diluted culture (~10^6 CFUs) were inoculated into 180 µL growth medium.

Optical density reads were measured every 45 mins for 24 hrs.

Supernatant assay

We hypothesised that higher densities and shifts in growth curve may be caused by

production of less-toxic metabolites. This hypothesis was tested using a supernatant

assay. Week seven cultures were revived and normalised as previous using five

replicates of resistant and susceptible populations (one replicate each of susceptible

and resistant cultures became contaminated and were not used in the experiment).

Two replicates of each week seven culture (two technical replicates for each

biological replicate) were established to ensure enough supernatant was available

for the reciprocal transplant assay. Cultures were grown for one week. On day five,

fresh week seven cultures were revived and grown for two days. Week-old cultures

were sterilised by passing medium through a 0.22 µm filter syringe. Replicate

supernatants from phage resistant and susceptible populations were pooled into

separate supernatant stocks. To separate vials, 3 mL of phage resistant and

susceptible supernatant was added to 25 mL vials. Two-day (fresh) culture densities

were normalised and each culture was inoculated into both supernatant types. This

established two controls in which cultures were grown in the same supernatant (e.g.

phage resistant in phage resistant supernatant) and treatments in which cultures

were transplanted into the opposite supernatant (e.g. phage resistant in phage

susceptible supernatant and vice versa). Cultures were grown for two days and

plated as previous.

Phage resistance and community structure

Next we assayed whether resistance to phage gives *Variovorax* an advantage over

Ochrobactrum and *Pseudomonas* when phages are absent. Six independent phage

resistant clones and six independent phage susceptible clones were isolated from

phage and no phage week two monoculture evolution lines respectively. 'No phage'

lines were selected for isolation of phage susceptible clones as phage resistance

was too high in 'phage' lines for enough individual susceptible clones to be isolated.

Comparison between 'no phage' and 'phage' lines accounts for any change in

general laboratory adaptation. Each clone and ancestral *Variovorax* were individually

grown for two days at 28 °C, shaking (180 r.p.m.) in 6 mL 1/64 TSB. Ancestral

Pseudomonas and *Ochrobactrum* isolates were also grown in the same conditions.

Isolate densities were normalised to 10^5 CFUs/mL and 10 μL added to relevant vials.

Each *Variovorax* clone (ancestral, resistant, susceptible) was grown as a polyculture

with *Pseudomonas* and *Ochrobactrum*, with six replicates per treatment. Cultures

were grown static for 1 week at 28 °C, then frozen as previous and plated from

frozen.

Statistical analyses

All data were analysed using R (v.4.2.1) in RStudio (48) and all plots were made

using the package '*ggplot2*' (49). Model simplification was conducted using likelihood

ratio tests and Tukey's post hoc multiple comparison tests were done using the R

package '*emmeans*' (50).

Variovorax densities were analysed in a linear mixed effects model with density

(log10 CFU/mL) tested against interacting fixed effects of phage exposure, culture

type (polyculture or monoculture) and time (weeks two, four, six, eight) with a

random effect of treatment replicate. Clonal densities (log10 CFU/mL) from common

garden experiments were analysed against interacting fixed effects of phage

exposure and time (week two or four) with random effects of treatment replicate and

block (experiment conducted in two blocks for feasibility). In competition assays,

relative fitness was analysed in a linear model with a fixed effect of competitor

(ancestor, phage resistant or susceptible).

One week growth curves were analysed in a mixed effects model with optical density

tested against a fixed term of day and interacting fixed effects of treatment (phage

resistant / susceptible) and a quadratic term of day to account for the non-linear

relationship. A random effect of treatment replicate was included. 24-hour assays of

bacterial growth were used to estimate exponential growth rate. One replicate was

removed from the no-phage treatment for failing to grow. Per replicate, exponential

growth was estimated using rolling regression, taking the steepest slope of the linear

regression between OD_{600} and time in hours in a shifting window of every nine time

points (~6.75 hr). A linear model tested growth rate against treatment.

The supernatant assay was analysed in a linear mixed effects model with density

(log10 CFU/mL) analysed against interacting fixed effects of supernatant type and

original treatment with a random effect of treatment replicate.

To test whether consistent genetic differences occur within treatments, we performed

nonmetric multidimensional scaling on the Euclidean distance matrix of SNPs/indels

and their proportions in each population using '*metaMDS*' in the R package '*vegan*'

(51). Nonmetric multidimensional scaling aims to collapse information from multiple

dimensions (i.e., from different populations and multiple SNPs/indels per population)

into just a few, allowing differences between samples to be visualized and

interpreted. Permutational ANOVA tests were run using '*vegan::adonis*', with

Euclidean distance as the response term and treatment as the predictor variable.

Additionally, we analysed whether evolution with phage affected the genetic

difference of each population (phage, no phage) from the ancestral population. The

genetic distance from the ancestral population was calculated as the sum of the

difference of the proportion of each SNP/indel in each population from the ancestral

proportion. Genetic distance was then analysed in a linear model analysing distance

against phage presence/absence.

To assess whether *Variovorax* phage resistance influences community structure in

the absence of phage, we analysed the relative proportion of each species in a

generalised linear mixed effects model. Here, each species proportion was analysed

against interacting fixed effects of *Variovorax* phage resistance and species identity,

with a random effect of treatment replicate and a binomial error structure.

References

1. Breitbart M, Bonnain C, Malki K, Sawaya NA. 2018. Phage puppet masters of the marine microbial realm. Nat Microbiol 3:754–766.

2. Breitbart M. 2012. Marine Viruses: Truth or Dare. Annu Rev Mar Sci 4:425–448.

3. Winter C, Bouvier T, Weinbauer MG, Thingstad TF. 2010. Trade-Offs between Competition and Defense Specialists among Unicellular Planktonic Organisms: the "Killing the Winner" Hypothesis Revisited. Microbiol Mol Biol Rev 74:42–57.

4. Clokie MRJ, Millard AD, Letarov AV, Heaphy S. 2011. Phages in nature. Bacteriophage 1:31–45.

5. Flynn KJ, Mitra A, Wilson WH, Kimmance SA, Clark DR, Pelusi A, Polimene L. 2022. 'Boom☐and☐busted' dynamics of phytoplankton–virus interactions explain the paradox of the plankton. New Phytologist 234:990–1002.

6. Liao H, Liu C, Ai C, Gao T, Yang Q-E, Yu Z, Gao S, Zhou S, Friman V-P. 2023. Mesophilic and thermophilic viruses are associated with nutrient cycling during hyperthermophilic composting. ISME J 17:916–930.

7. Gandon S, Buckling A, Decaestecker E, Day T. 2008. Host-parasite coevolution and patterns of adaptation across time and space. J Evol Biol 21:1861–1866.

8. van Houte S, Buckling A, Westra ER. 2016. Evolutionary Ecology of Prokaryotic Immune Mechanisms. Microbiol Mol Biol Rev 80:745–763.

9. Chan BK, Sistrom M, Wertz JE, Kortright KE, Narayan D, Turner PE. 2016. Phage selection restores antibiotic sensitivity in MDR Pseudomonas aeruginosa. Sci Rep 6:26717.

10. Buckling A, Wei Y, Massey RC, Brockhurst MA, Hochberg ME. 2006. Antagonistic

coevolution with parasites increases the cost of host deleterious mutations. Proc Royal

Soc B 273:45–49.

11. Li N, Zeng Y, Wang M, Bao R, Chen Y, Li X, Pan J, Zhu T, Hu B, Tan D. 2022.

Characterization of Phage Resistance and Their Impacts on Bacterial Fitness in

Pseudomonas aeruginosa. Microbiol Spectr 10:e02072-22.

12. Koskella B, Lin DM, Buckling A, Thompson JN. 2012. The costs of evolving resistance

in heterogeneous parasite environments. Proc R Soc B 279:1896–1903.

13. Lennon JT, Khatana SAM, Marston MF, Martiny JBH. 2007. Is there a cost of virus

resistance in marine cyanobacteria? The ISME Journal 1:300–312.

14. Castledine M, Padfield D, Sierocinski P, Soria Pascual J, Hughes A, Mäkinen L, Friman

V-P, Pirnay J-P, Merabishvili M, de Vos D, Buckling A. 2022. Parallel evolution of

Pseudomonas aeruginosa phage resistance and virulence loss in response to phage

treatment in vivo and in vitro. eLife 11:e73679.

15. Abedon ST. 2023. Ecology and Evolutionary Biology of Hindering Phage Therapy: The

Phage Tolerance vs. Phage Resistance of Bacterial Biofilms. Antibiotics 12:245.

16. Harper D, Parracho H, Walker J, Sharp R, Hughes G, Werthén M, Lehman S, Morales

S. 2014. Bacteriophages and Biofilms. Antibiotics 3:270–284.

17. Salathé M, Soyer OS. 2008. Parasites lead to evolution of robustness against gene loss

in host signaling networks. Mol Syst Biol 4:202.

18. Crabbé A, Jensen PØ, Bjarnsholt T, Coenye T. 2019. Antimicrobial Tolerance and

Metabolic Adaptations in Microbial Biofilms. Trends in Microbiology 27:850–863.

19. Oliveira NM, Martinez-Garcia E, Xavier J, Durham WM, Kolter R, Kim W, Foster KR.

2015. Biofilm Formation As a Response to Ecological Competition. PLoS Biol

13:e1002191.

20. Jayaraman R. 2011. Hypermutation and stress adaptation in bacteria. J Genet 90:383–

391.

21. Pal C, Maciá MD, Oliver A, Schachar I, Buckling A. 2007. Coevolution with viruses

drives the evolution of bacterial mutation rates. Nature 450:1079–1081.

22. Gómez P, Buckling A. 2011. Bacteria-Phage Antagonistic Coevolution in Soil. Science

332:106–109.

23. Castledine M, Padfield D, Schoeman M, Berry A, Buckling A. 2024. Bacteria-phage

(co)evolution is constrained in a synthetic community across multiple bacteria-phage

pairs https://doi.org/10.1101/2024.10.14.618343.

24. Zhu F, Wu R, Zhang H, Wu H. 2013. Structural and Biochemical Analysis of a

Bacterial Glycosyltransferase, p. 29–39. *In* Brockhausen, I (ed.), Glycosyltransferases.

Humana Press, Totowa, NJ.

25. Fang Q, Feng Y, McNally A, Zong Z. 2022. Characterization of phage resistance and

phages capable of intestinal decolonization of carbapenem-resistant Klebsiella

pneumoniae in mice. Commun Biol 5:48.

26. Beckett SJ, Williams HTP. 2013. Coevolutionary diversification creates nested-modular

structure in phage–bacteria interaction networks. Interface Focus 3:20130033.

27. Demerec M, Hartman PE. 1959. Complex Loci in Microorganisms. Annu Rev

Microbiol 13:377–406.

28.   Ballouz S, Francis AR, Lan R, Tanaka MM. 2010. Conditions for the Evolution of Gene

Clusters in Bacterial Genomes. PLoS Comput Biol 6:e1000672.

29.   Castledine M, Pennycook J, Newbury A, Lear L, Erdos Z, Lewis R, Kay S, Sanders D,

Sünderhauf D, Buckling A, Hesse E, Padfield D. 2024. Characterizing a stable five-

species microbial community for use in experimental evolution and ecology.

Microbiology 170.

30.   Castledine M, Padfield D, Buckling A. 2020. Experimental (co)evolution in a

multi☐species microbial community results in local maladaptation. Ecol Lett 23:1673–

1681.

31.   Satola B, Wübbeler JH, Steinbüchel A. 2013. Metabolic characteristics of the species

Variovorax paradoxus. Appl Microbiol Biotechnol 97:541–560.

32.   Griffin AS, West SA, Buckling A. 2004. Cooperation and competition in pathogenic

bacteria. Nature 430:1024–1027.

33.   Zhao Y, Temperton B, Thrash JC, Schwalbach MS, Vergin KL, Landry ZC, Ellisman

M, Deerinck T, Sullivan MB, Giovannoni SJ. 2013. Abundant SAR11 viruses in the

ocean. Nature 494:357–360.

34.   Castledine M, Buckling A. 2024. Critically evaluating the relative importance of phage

in shaping microbial community composition. Trends in Microbiology

S0966842X2400057X.

35.   Giovannoni SJ. 2017. SAR11 Bacteria: The Most Abundant Plankton in the Oceans.

Annu Rev Mar Sci 9:231–255.

36.   Thingstad TF, Våge S, Storesund JE, Sandaa R-A, Giske J. 2014. A theoretical analysis

of how strain-specific viruses can control microbial species diversity. Proc Natl Acad

Sci USA 111:7813–7818.

37.   Scanlan PD, Buckling A. 2012. Co-evolution with lytic phage selects for the mucoid

phenotype of Pseudomonas fluorescens SBW25. ISME J 6:1148–1158.

38.   Fernández-García L, Kirigo J, Huelgas-Méndez D, Tomás M, García-Contreras R,

Wood TK. 2023. Phages Produce Persisters https://doi.org/10.1101/2023.10.17.562728.

39.   Niu H, Gu J, Zhang Y. 2024. Bacterial persisters: molecular mechanisms and

therapeutic development. Sig Transduct Target Ther 9:174.

40.   Brockhurst MA, Rainey PB, Buckling A. 2004. The effect of spatial heterogeneity and

parasites on the evolution of host diversity. Proc R Soc Lond B 271:107–111.

41.   Dy RL, Rigano LA, Fineran PC. 2018. Phage-based biocontrol strategies and their

application in agriculture and aquaculture. Biochemical Society Transactions 46:1605–

1613.

42.   Oechslin F. 2018. Resistance Development to Bacteriophages Occurring during

Bacteriophage Therapy. Viruses 10:351.

43.   Abedon ST. 2019. Use of phage therapy to treat long-standing, persistent, or chronic

bacterial infections. Adv Drug Deliv Rev 145:18–39.

44.   Deatherage DE, Barrick JE. 2014. Identification of mutations in laboratory-evolved

microbes from next-generation sequencing data using breseq. Methods Mol Biol

1151:165–188.

45.   Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJE. 2015. The Phyre2 web

portal for protein modeling, prediction and analysis. Nat Protoc 10:845–858.

46.   Sünderhauf D, Klümper U, Gaze WH, Westra ER, Van Houte S. 2023. Interspecific

competition can drive plasmid loss from a focal species in a microbial community. The

ISME Journal 17:1765–1773.

47.   Lenski RE, Rose MR, Simpson SC, Tadler SC. 1991. Long-Term Experimental

Evolution in Escherichia coli. I. Adaptation and Divergence During 2,000 Generations.

Am Nat 138:1315–1341.

48.   Team RC. 2013. R: a language and environment for statistical computing.

49.   Wickham H. 2016. ggplot2: elegant graphics for data analysis. Springer, New York.

50.   Lenth R. 2018. Emmeans: Estimated marginal means, aka least☐squares means. R

Package Version.

51.   Oksanen J, Simpson GL, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'Hara RB,

Solymos P, Stevens MHH, Szoecs E, Wagner H, Barbour M, Bedward M, Bolker B,

Borcard D, Carvalho G, Chirico M, De Caceres M, Durand S, Furneaux B, Hannigan G,

Hill MO, Lahti L, McGlinn D, Ouellette M-H, Ribeiro Cunha E, Smith T, Stier A, Ter

Braak CJF, Weedon J. 2019. vegan: Community Ecology Package.

 Supplementary Information

Figure S1. The divergence of each treatment group as indicated by Euclidean distances of each gene (arrows) from the ancestral genome. Points represent individual replicates sequenced from phage resistant or susceptible treatment groups.

741 Table S1. Tukey HSD comparisons comparing the relative proportion of

742 *Ochrobactrum* (O), *Pseudomonas* (P) and *Variovorax* (V) when the *Variovorax* strain

743 is the ancestral type, phage resistant or susceptible. P-values are adjusted by the

744 tukey method for comparing a family of three estimates.

Species	Contrast	Estimate	z-ratio	p-value
O	Ancestral - Resistant	-0.146	-2.252	0.063
O	Ancestral - Susceptible	-0.049	-0.763	0.726
O	Resistant - Susceptible	0.096	1.483	0.299
P	Ancestral - Resistant	0.495	5.018	<0.001
P	Ancestral - Susceptible	0.129	1.403	0.339
P	Resistant - Susceptible	-0.367	-3.637	0.001
V	Ancestral - Resistant	-0.091	-1.257	0.42
V	Ancestral - Susceptible	-0.019	-0.259	0.964
V	Resistant - Susceptible	0.072	0.995	0.58

745
746