Loss of Bbs8 leads to cystic kidney disease in mice with reduced acetylation of ciliary alpha-tubulin through HDAC2

Emilia Kieckhöfer[1,2], Julia Günzler[1,2,#], Peter A. Matthiessen[3,#], Lena K. Ebert[1,2], Christina Klausen[4], Dagmar Wachten[4], Metin Cetiner[5], Max C. Liebau[6], Thomas Benzing[1,2], Helen May-Simera[3,*], Bernhard Schermer[1,2,*,§]

[1] Department II of Internal Medicine and Center for Molecular Medicine Cologne, University of Cologne, Faculty of Medicine and University Hospital Cologne, Cologne, Germany.

[2] Cologne Excellence Cluster on Cellular Stress Responses in Aging-Associated Diseases (CECAD), University of Cologne, Faculty of Medicine and University Hospital Cologne, Cologne, Germany.

[3] Faculty of Biology, Ciliary Cell Biology, Institute of Molecular Physiology, Johannes Gutenberg-University, Mainz, Germany.

[4] Institute of Innate Immunity, Biophysical Imaging, Medical Faculty, University of Bonn, Bonn, Germany.

[5] Department of Pediatric Nephrology, Pediatrics II, University of Duisburg-Essen, Essen, Germany.

[6] Department of Pediatrics and Center for Molecular Medicine Cologne, University Hospital Cologne and Faculty of Medicine, University of Cologne, Cologne, Germany.

# These authors contributed equally.

* These authors contributed equally.

§corresponding author: bernhard.schermer@uk-koeln.de

Keywords: BBS, primary cilium, kidney, ciliopathies, cysts, HDAC2

Abstract

Primary cilia dysfunction underlies a group of severe disorders known as ciliopathies. These include

Bardet-Biedl syndrome (BBS), which is caused by mutations in BBS genes encoding for components of

ciliary protein complexes essential for the assembly and maintenance of primary cilia. As in most

ciliopathies, a hallmark feature of BBS is the development of cystic kidney disease. However, the

molecular mechanisms linking ciliary dysfunction to cystogenesis remain incompletely understood. Here,

we show that $Bbs8^{-/-}$ mice develop late-onset cystic kidney disease accompanied by increased regulated

cell death and fibrosis. While the number and length of cilia are not affected, loss of BBS8 reduces K40

acetylation of α-tubulin within primary cilia, compromising ciliary stability. Notably, proteomic analysis

revealed a significant upregulation of histone deacetylase HDAC2 in $Bbs8^{-/-}$ kidneys, which we confirmed

in $Bbs8^{-/-}$ mouse embryonic fibroblasts (MEFs) and in urine-derived renal epithelial cells (URECs) from a

BBS8 patient. We further demonstrate the protein interaction between BBS8 and HDAC2, implicating a

disrupted BBS8–HDAC2 regulatory axis in disease pathogenesis. Consistent with a role of excessive

HDAC2 activity in the BBS8 deficient cells, pharmacological inhibition of HDAC2 restored tubulin

acetylation in BBS8 urine-derived cells. Thus, modulation of HDAC2 activity may represent a strategy to

alter ciliary stability *in vivo* which could explain positive effects of class I specific HDAC inhibitors in

models of cystic kidney disease.

Introduction

The primary cilium is a specialized microtubule-based sensory organelle extending from most

mammalian cells' surfaces (Pazour and Bloodgood, 2008). It consists of the axoneme, ensheathed by the

ciliary membrane, the transition zone, and the basal body, which anchors the cilium to the cell body. The

transition zone between the basal body and the ciliary shaft is critical for protein trafficking to and from

the ciliary compartment (Reiter, Blacque and Leroux, 2012; Garcia-Gonzalo and Reiter, 2017). Tight

regulation of the protein composition is required for cilia to function as a hub for chemo- and mechano-

transduction, as well as ciliary signaling (Reiter, Blacque and Leroux, 2012). As such, they play a pivotal

role in modulating various signaling pathways, e.g. Hedgehog, Wnt, and TGF-β signaling (Fan and Tessier-

Lavigne, 1994; Munsterberg *et al.*, 1995; Pourquié *et al.*, 1996). Unlike any other organelle, cilia

periodically disappear during the mitotic cell cycle: When cells re-enter mitosis, cilia are typically

disassembled to subsequently release the basal body, a modified mother centriole, to serve as

microtubule organizing centers at the spindle poles. Disassembly of cilia mechanistically involves the

deacetylation of ciliary α-tubulin by HDAC6 and, thus, destabilization of microtubules (Pugacheva *et al.*,

2007; Gopalakrishnan *et al.*, 2023). It is well established that pathogenetic variants in proteins, which

alter the structure or function of cilia, lead to a heterogeneous group of genetic diseases and syndromes

referred to as ciliopathies (Hildebrandt, Benzing and Katsanis, 2011). Ciliopathies comprise a diverse

group of genetic disorders with overlapping phenotypes in different organs and tissues, such as the

brain, eye, skeleton, liver, vasculature, and kidney (Forsythe and Beales, 2013; Reiter and Leroux, 2017).

Thereby, cystic and fibrotic kidney diseases are a common occurrence among many ciliopathies, leading

to the frequent classification of a significant subgroup, namely: renal ciliopathies (Hildebrandt and Zhou,

2007; McConnachie, Stow and Mallett, 2021).

The Bardet-Biedl Syndrome is an archetypical autosomal-recessive ciliopathy characterized by obesity,

retinopathy, polydactyly, and kidney disease (Forsythe and Beales, 2013). This syndrome is caused by

pathogenic variants of BBS genes encoding for BBS proteins. Most BBS proteins can be classified

according to the particular protein complexes which they form. The BBSome complex (Nachury *et al.*,

2007; Loktev *et al.*, 2008) (BBS1, BBS2, BBS4, BBS5, BBS7, BBS8, BBS9, BBS18) is required for trafficking of

ciliary membrane proteins whereas, the chaperonin-like complex (BBS6, BBS10, and BBS12), catalyzes

the assembly of the BBSome (Seo *et al.*, 2010). The remaining BBS proteins have varying specific

functions but are predicted to influence ciliary trafficking. The kidney phenotype in BBS patients is

variable with kidneys showing parenchymal, medullary, and most often corticomedullary cysts, but also

renal fibrosis, unilateral agenesis, or dysplastic kidneys (Beales *et al.*, 1999; Putoux *et al.*, 2012; Forsythe

et al., 2017; Elawad *et al.*, 2022). However, these phenotypes are highly variable among families and

could be dependent on mutation type (Putoux *et al.*, 2012; Forsythe *et al.*, 2017).

Studies on various *Bbs* mutant mice, including *Bbs1^{M390R}* transgenic, *Bbs2$^{-/-}$*, *Bbs4$^{-/-}$*, *Bbs5$^{-/-}$*, *Bbs6$^{-/-}$*, *Bbs8$^{-/-}$*

and *Bbs10$^{-/-}$*, revealed significant parallels to human disease. These mice exhibit a broad spectrum of

vision impairment, obesity, male infertility, and neurological deficits (Kulaga *et al.*, 2004; Mykytyn *et al.*,

2004; Nishimura *et al.*, 2004; Fath *et al.*, 2005; Davis *et al.*, 2007; Rahmouni *et al.*, 2008; Seo *et al.*, 2009;

Cognard *et al.*, 2015; Kretschmer *et al.*, 2019; Bentley-Ford *et al.*, 2021; Rödig *et al.*, 2022; Sieckmann *et*

al., 2024). For BBS10, complete knockout resulted in renal abnormalities primarily affecting glomeruli

and podocytes without cyst formation, while a tubular epithelial cell-specific knockout did not show any

overt phenotype (Cognard *et al.*, 2015). Cyst formation has been observed in *Bbs2$^{-/-}$* and *Bbs4$^{-/-}$* mice, but

again primarily affecting glomeruli (Nishimura *et al.*, 2004; Guo *et al.*, 2011). Loss of *Bbs8* has so far

shown to cause the most severe retinal phenotype in mice (Tadenev *et al.*, 2011; Dilan *et al.*, 2018;
Kretschmer *et al.*, 2019; Schneider *et al.*, 2021), which is not only caused by ciliary defects of
photoreceptor cells but also due to ciliary defects in the retinal pigment epithelium (RPE) (Kretschmer *et*
al., 2023; Schneider *et al.*, 2021). BBS8 is one of the key components of the BBSome, which might explain
the severity of the *Bbs8*$^{-/-}$ phenotype. BBS8 (TTC8) is a tetratricopeptide repeat (TPR) protein with a
critical role in planar cell polarity and laterality (May-Simera *et al.*, 2010, 2015; Patnaik *et al.*, 2019). Loss
of *Bbs8* leads to changes in the composition of the octameric BBSome complex, where BBS8 is a direct
interactor of the scaffold protein BBS9 (Zhang *et al.*, 2012).

Until now, research into BBS8 function has predominantly focused on embryonic development and
ocular health. A subtle renal phenotype, characterized by mild tubular dilation at around 8 months of
age, has been reported (Tadenev *et al.*, 2011), but specific roles of BBS8 in maintaining renal cilia and
preserving kidney architecture remain largely unexplored. In this study, we report the occurrence of
tubular cystic kidney disease in *Bbs8*-deficient mice at 46 weeks of age, accompanied by renal fibrosis
and increased rate of regulated cell death. Interestingly, neither cilia number, nor length differed
significantly from wild-type littermates; however, the levels of acetylated tubulin within primary cilia
were markedly reduced. We demonstrate that this reduction, which destabilizes cilia, is driven by
increased expression of histone deacetylase HDAC2. This effect is confirmed in isolated mouse
embryonic fibroblasts (MEFs) and urine-derived renal epithelial cells (URECs) from a BBS8 patient. Thus,
beyond potential therapeutic implications of HDAC2 inhibition, our data highlight the importance of not
relying solely on acetylated tubulin as a marker for cilia in functional studies, particularly in disease
contexts where ciliary stability is affected.

Methods

Mouse lines

Bbs8$^{-/-}$ mice have been previously described (Tadenev *et al.*, 2011). All animals were housed, handled,
and animal studies conducted, in accordance with approved institutional animal care and use committee
procedures. All experiments had ethical approval from the Landesuntersuchungsamt Rheinland-Pfalz and
were performed in accordance with institutional guidelines for animal welfare, German animal
protection law, and the guidelines given by the ARVO Statement for the Use of Animals in Ophthalmic
and Vision Research. Animal maintenance and handling were performed in line with the Federation for
Laboratory Animal Science Associations (FELASA) recommendations. Animals were housed in a 12 h
light/dark cycle with food and water available *ad libitum*. For the preparation, mice were weighted,

followed by cervical dislocation; blood was collected prior to the perfusion of the kidney with PBS

through the aorta. Further, other organs and fat tissue were taken and, as the kidney, processed by

either fixation in 4 % formaldehyde and embedded in paraffin or snap-frozen for further tissue analysis.

The blood was incubated for 2 h at RT, followed by centrifugation. Serum creatinine levels were

measured by the Institute of Clinical Chemistry, University Hospital of Cologne, Germany.

Immunohistology

For histological analysis, tissue was cut into 2-μm-thick sections and deparaffinized by xylene treatment

and rehydration in graded ethanol. For PAS staining, the sections were stained with 0,9 % periodic acid

(cat# 3257.1, Roth) and Schiffsches Reagent (cat# 1.09033, Merck) both for 10 min embedded into

washing steps with H_2O. Finally, to visualize nuclei in blue, the samples were stained with Mayer's

Haematoxylin for 20 s. For the Masson staining the Masson-Goldner's trichrome staining kit (cat# 3459,

Roth) was used and performed according to the manufacturer's instructions. After dehydration of the

sections, they were embedded with Histomount (HS-103, National Diagnostics).

Cyst index analysis

The cyst index was calculated for whole slide images using open-source software for bioimage analysis

QuPath (v0.4.0) (Bankhead *et al.*, 2017). Cysts were detected using an Artificial Neural Network-based

pixel classifier. Initially detected cysts were filtered for the minimal area of 400 μm² and a minimum

circularity value of 0.35. Plots were generated using the Plots of Data web app (Postma and Goedhartid,

2019). For statistical analysis, all results were normalized to the control followed by a two-tailed

Student's *t*-test (p<0.05).

Proteome and phosphoproteome analysis

For each biological replicate a quarter of kidney tissue was used and dounced with a Wheaton Dounce

tissue grinder in urea buffer (8 M Urea, 50 mM ammonium bicarbonate) supplemented with Halt

protease-phosphatase-inhibitor cocktail (Thermo Scientific™). After clearing of the sample (16.000 xg, 1

h at 4°C), the lysates were reduced (10 mM dithiothreitol), alkylated (50 mM chloroacetamide) and

digested (LysC; 1:75). Samples (800 μg) were diluted to 2 M urea and subjected to tryptic digestion

(1:50). After overnight incubation, phosphoenrichment was performed in the CECAD proteomics facility

using the Thermo Scientific™ Kit High Select TiO2 Kit (#A32993). All samples were analyzed as well by the

CECAD proteomics facility on a Q Exactive Plus Orbitrap mass spectrometer that was coupled to an EASY

nLC (both Thermo Scientific™). Peptides were loaded with solvent A (0.1% formic acid in water) onto an

in-house packed analytical column (50 cm, 75 µm inner diameter, filled with 2.7 µm Poroshell EC120 C18,

Agilent). Peptides were chromatographically separated at a constant flow rate of 250 nL/min using the

following gradient: 3-5% solvent B (0.1% formic acid in 80 % acetonitrile) within 1.0 min, 5-30% solvent B

within 121.0 min, 30-40% solvent B within 19.0 min, 40-95% solvent B within 1.0 min, followed by

washing and column equilibration. The mass spectrometer was operated in data-dependent acquisition

mode. The MS1 survey scan was acquired from 300-1750 m/z at a resolution of 70,000. The top 10 most

abundant peptides were isolated within a 1.8 Th window and subjected to HCD fragmentation at a

normalized collision energy of 27%. The AGC target was set to 5e5 charges, allowing a maximum

injection time of 55 ms. Product ions were detected in the Orbitrap at a resolution of 17,500. Precursors

were dynamically excluded for 25.0 s. All mass spectrometric raw data were processed with MaxQuant

(Tyanova, Temu and Cox, 2016) (version 2.2.0.0) using default parameters against the UniProt canonical

murine database (UP10090, downloaded 20.01.2023) with the match-between-runs option enabled

between replicates. Samples were sorted into two parameter groups, either containing the enriched or

non-enriched samples. Enriched samples had the phosphorylation (STY) variable modification added,

whereas non-enriched samples were quantified by LFQ. A follow-up analysis was done in Perseus 1.6.15

(Tyanova *et al.*, 2016). Results were cleaned up by removing hits from the decoy database, the

contaminant list and, in case of non-enriched fractions, those only identified by modified peptides were

removed. Afterwards, results were filtered for data completeness in at least one condition and LFQ

values (WP) or intensities (PP), imputed using sigma downshift with standard settings. Finally, FDR-

controlled two-sided t-tests between sample groups were performed (S0=0, FDR≤0.05) as well as a 1D

annotation enrichment using Perseus (version 1.6.15.0).

Isolation and immortalization of mouse embryonic fibroblasts (MEFs)

For the isolation of mouse embryonic fibroblasts (MEFs), timed matings were set up with one male and

two females of the desired genotype. At day 13, the pregnant mouse was anesthetized using isoflurane

(Piramal Healthcare) followed by a cervical dislocation. The lower abdomen was opened by an abdominal

incision to extract the two uterine horns. Embryos were isolated and transferred into a 24-well plate

filled with PBS. The head and the red organs (heart and liver) were removed. The rest of the embryo was

placed into a 12-well plate filled with 2 ml ice-cold 0.25 % Trypsin/PBS (diluted from 2.5 % Trypsin,

Gibco). The embryos were chopped into small pieces and incubated overnight at 4 °C. The trypsin

solution was discarded and the remaining Trypsin/tissue mixture was incubated for 30 min in a 37 °C

water bath. Afterward, medium (composition: DMEM/Glutamax, 10 % FCS, 1 % sodium pyruvate (100x),

1 % Pen Strep), was added, and the cell suspension was pipetted vigorously up and down to break up the

digested tissue into a single cell suspension. After 1 min, the cell suspension was collected, to remove

the sedimentation of the remaining tissue. This step was repeated and the cell suspension was filtered

through a 100 µm cell-strainer (Corning). Cells were plated and after 24 h, the medium was changed.

Immortalization of MEFs was performed as described previously (Todaro and Green, 1963). Briefly, cells

are split every three days and seeded with the same cell density. From passage three onwards, cells were

seeded on at least two 10 cm culture dishes. After around 15 passages, cells started to regrow. When

MEFs were immortalized, frozen back-ups were made.

Isolation, immortalization, and BBS8 re-expression of urine renal epithelial cells (URECS)

BBS8[c.915delG] patient urine (male; age 14) was collected at the University Hospital Essen, Germany, in

cooperation with the Network for early onset cystic kidney disease (Neocyst). . Urine sample was

centrifuged (10 min, 400 x g), and the pellet was resuspended in 10 ml washing buffer (sterile 1x PBS,

100 U Penicillin/Streptomycin (Gibco™), 500 ng/ml Amphotericin B (Sigma)). After centrifugation (10

min, 200 x g) the supernatant was removed leaving ~0.2 – 0.5 ml in the tube. The pellet was resuspended

in primary medium (PUC; DMEM-F12 (Sigma), REGM SingleQuot kit (Lonza, cat# CC-4127), 10% FBS

(Gibco™), 2 mM GlutaMAX (Gibco™), 1.0% Penicillin/ Streptomycin (Gibco™), 500 ng/ml Amphotericin B

(Sigma)) and plated in a single well of a galantine-coated 12-well plate. The plate was incubated for 48 h

at 37 °C, before the first PUC medium change. With the first visible colonies, PUC medium was changed

to normal growth medium (DMEM-F12 (Sigma), 10% FBS (Gibco™), 2 mM GlutaMAX (Gibco™), 1.0%

Penicillin/ Streptomycin (Gibco™)). Reaching a confluency of 60-80 %, cells were split in two wells of a 6-

well plate using TrypLE Select Express (Gibco™). One well was expanded, and stored as primary cells, and

one well was used for immortalization. Immortalization was started by adding 1 ml of previously

produced virus supplemented with 1 µl of Polybrene 8mg/ml (Sigma) to 40 % confluent cells and

incubated for 24 h at 37°C. The virus was removed and replaced by normal growth medium. Cells were

split thrice, while expansion in multiple 10 cm culture dishes before frozen back-ups were made.

Immortalized urine-derived renal epithelial cells (URECS) were additionally expressed exogenous with

viruses produced with pLenti6.3 F.hBBS8 and pLenti6.3 F.GFP in HEK293T cells. Sterilized harvest virus

supplemented with Polybrene was added to 40 % confluent cells and incubated for 24 h at 37°C. After

removing the virus, the cells were selected by adding normal growth medium supplemented with 0,001

mg/ml Blasticidin (Invivogen). Cells were split three times before selection was stopped and cells were

used for experiments.

Cell culture and treatment

Human embryonic kidney 293T cells (HEK293T, ATCC®), mIMCD3 cells, MEFs and URECs were cultivated

at 37°C and 5% CO_2. Thereby, HEK293T cells were maintained in DMEM + GlutaMAX™ medium (Gibco™)

supplemented with 10% fetal bovine serum (FBS, Gibco™), MEFs in DMEM/F-12 + GlutaMAX™ medium

(Gibco™) complemented with 10% FBS and 1% penicillin-streptomycin.URECs and mIMCD3 cells were

cultured in normal growth medium (DMEM-F12 (Sigma), 10% FBS (Gibco™), 2 mM GlutaMAX (Gibco™).

For URECs, 1.0% Penicillin/ Streptomycin (Gibco™) was added. All cell lines were tested negative for

mycoplasma (PCR Mycoplasma Test Kit I/C, PromoKine). To induce a higher amount of ciliogenesis,

mIMCD3 cells were grown in serum-free media for 24 h at 37°C. HEK293T cells were transiently

transfected with5 µg (IP) or 6 µg (Interactome) of F.hBBS8 in pcDNA6 using the calcium phosphate

method. As negative control, equal amounts of F.EPS[1-225] in pcDNA6 were used. The HDAC2 inhibitor

Tucidinostat (TUC, 160 nM; Biozol, cat# HY-109015) treatment was performed in serum-free media at

37°C for 24 h. Control treatment was performed with DMSO (AppliChem). For IF stainings, confluent

mIMCD3 cells were transfected with 5 µg of either F.EPS or F.HDAC2 using Lipofectamine®3000

(Invitrogen) in starvation medium and fixed 24 h after transfection. Every plasmid used for this study

added an N-terminal F9-tag to the protein of interest, which refers to a FLAG-tag followed by 9 histidine

residues. Only mHDAC2 was expressed with a single FLAG-tag.

Harvest and fixation

Cells were seeded in dishes or on glass coverslips in 12 well plates and incubated for 24 h, prior to

starvation or TUC treatment; both for an additional 24 h at 37°C. To harvest cells, they were washed with

PBS (1x), scraped in fresh PBS (1x), and centrifuged for 10 min, 10.000 rpm at 4°C. Cell pellets could be

used for western blot or RNA isolation. Harvest of whole cell lysate samples was performed by

resuspending the cells in cold 1x PBS followed by centrifugation for 5 min, 4.000 rpm at 4°C. Afterwards

cells were resuspended in 1x Laemmli with DTT and boiled for 5 min at 95°C. For Immunofluorescence

staining cells were fixed by adding 4% PFA (Walter CMP GmBH & Co) for 5 min, followed by methanol

(Roth) for 4 min. Fixed cells were washed thrice with PBS (1x) before staining.

Cilia stability assay

211 Depolymerization of microtubules was induced by incubating starved cells for 30 min on ice. The

212 subsequent fixation was continuously performed on ice. The untreated control cells were fixed at room

213 temperature. Fixed cells were stored at 4°C in PBS until staining and quantifying cilia.

Quantitative real-time PCR

RNA isolation was performed from kidney tissue samples, thereby using one-quarter of a kidney. Tissue

was ground with BeadBeater (Roth) using a Precelly24 with 5.000 rpm two times for 30 s in Tri-Reagent.

RNA isolation from cells was performed by resuspending snap-frozen cell pellets in Phenol. RNA

extraction was performed with the Direct-zol RNA Miniprep kit (Zymo Research) following the

manufacturer's instructions, including a DNase1 treatment step. Before the reverse transcription using

the High-Capacity cDNA Reverse Transcription kit (Applied Biosystems), RNA concentration and sample

quality were assessed on a Nanodrop spectrophotometer (Peqlab). mRNA was assessed by qPCR with

SYBR Green (Thermo Scientific™) using mHprt1 or hHPRT1 as endogenous control. The qPCR experiments

were performed on a QuantStudio Q5 Real-time PCR System (ThermoFisher Scientific). For data analysis,

all results were normalized to the housekeeping gene *Hprt1* using the delta-delta CT followed by a two-

tailed Student's *t*-test (p<0.05).

Co-Immunoprecipitation

Co-immunoprecipitation (Co-IP) was performed as previously described (Habbig *et al.*, 2011) using the IP

Buffer (20 mM Tris, 1% (v/v) TritonX-100, 50 mM NaCl, 15 mM $Na_4P_2O_7$, 50 mM NaF, pH 7.5)

supplemented with inhibitors (44 µg/µl PMSF, 2 mM Na_3VO_4) in four replicates. Input samples (lysates)

were collected, and the remaining samples were incubated with α-FLAG M2 Beads (A2220, Sigma-

Aldrich) for 2 h. Samples were denatured and reduced in 2x Laemmli before SDS-PAGE. For proteomic

analysis, the IP Buffer was supplemented with cOmplete™ Protease Inhibitor Cocktail (Roche). After

sonication, the lysates were first centrifuged 30 min at 14.000 g at 4°C followed by ultracentrifugation

for 45 min at 100.000 g and 4°C. After three washing steps the beads were incubated with 5% SDS in PBS

for 5 min at 95 °C, followed by reduction with 10 mM dithiothreitol and alkylation with 40 mM

chloroacetamide.

Interactome

Samples were analyzed by the CECAD Proteomics Facility on an Orbitrap Exploris 480 (Thermo Scientific,

granted by the German Research Foundation under INST 1856/71-1 FUGG) mass spectrometer equipped

with a FAIMSpro differential ion mobility device that was coupled to an Vanquish neo in trap-and-elute

setup (Thermo Scientific). Samples were loaded onto a precolumn (Acclaim 5µm PepMap 300 µ

Cartridge) with a flow of 60 µl/min before reverse-flushed onto an in-house packed analytical column (30

cm length, 75 µm inner diameter, filled with 2.7 µm Poroshell EC120 C18, Agilent). Peptides were

chromatographically separated with an initial flow rate of 400 nL/min and the following gradient: initial

2% B (0.1% formic acid in 80 % acetonitrile), up to 6 % in 3 min. Then, flow was reduced to 300 nl/min

and B increased to 20% B in 26 min, up to 35% B within 15 min and up to 98% solvent B within 1.0 min

while again increasing the flow to 400 nl/min, followed by column wash with 98% solvent B and re-

equilibration to initial condition. The FAIMS pro was operated at -50V compensation voltage and

electrode temperatures of 99.5 °C for the inner and 85 °C for the outer electrode. The mass

spectrometer was operated in data-dependent acquisition top 24 mode with MS1 scans acquired from

350 m/z to 1400 m/z at 60k resolution and an AGC target of 300%. MS2 scans were acquired at 15 k

resolution with a maximum injection time of 22 ms and an AGC target of 300% in a 1.4 Th window and a

fixed first mass of 110 m/z. All MS1 scans were stored as profile, all MS2 scans as centroid. All mass

spectrometric raw data were processed with Maxquant (version 2.4) (Tyanova, Temu and Cox, 2016)

using default parameters against the Uniprot HUMAN canonical database (UP5640) with the match-

between-runs option enabled between replicates. Follow-up analysis was done in Perseus 1.6.15

(Tyanova *et al.*, 2016). Protein groups were filtered for potential contaminants and insecure

identifications. Remaining IDs were filtered for data completeness in at least one group and missing

values imputed by sigma downshift (0.3 σ width, 1.8 σ downshift). Afterwards, FDR-controlled two-sided

t-tests were performed (S0=0, FDR≤0.05). Protein structure prediction was done using the Alphafold

server (Abramson *et al.*, 2024), while ChimeraX has been used for visualization (Pettersen *et al.*, 2021;

Meng *et al.*, 2023).

Immunoblotting

Confluent grown , and starved MEF cells were harvest with RIPA buffer (50 mM Tris/HCl, 150 mM NaCl,

1% (v/v) NP-4O, 0.5% (w/v) Sodium deoxycholate, 0.1% (w/v) SDS), supplemented with 1% Halt Protease

and Phosphatase Inhibitor Cocktail (Thermo Scientific™) (Brücker *et al.*, 2023). Cells were lysed on ice

and sonicated for 2 s. Protein concentration of lysates was determined and loaded onto a 10% SDS-PAGE

and subsequently transferred to a PVDF-FL membrane (Millipore). URECSs cell pellets were lysed in

adjusted inhibitor-supplemented RIPA buffer (50 mM Tris/HCL pH 7.5, 150 mM NaCl, 0.1 % NP-40, 0.5 %

Na-Deoxycholat, 0.1 % SDS) supplemented with Benzonase® (70746-3 Millipore), cOmplete™

(4693159001, Roche) and PhosSTOP™ (4906845001, Roche), on ice for 30 min. This is followed by

centrifugation for 1 h, 14.000 rpm at 4°C, determination of protein concentration, and supplementing

and boiling with 5x Laemmli. Samples were loaded onto a gel as before and transferred to a PVDF

membrane (Millipore), including whole cell lysates of mIMCD3 cells in 1x Laemmli. After blocking,

membranes were incubated overnight at 4°C in primary antibodies (FLAG, F7425, Sigma-Aldrich, 1:1000;

HDAC2, ab32117 Abcam, 1:1000; acetylated Tubulin, T6793 Sigma, 1:1000; Calnexin, 10427-2 AP

ProteinTech, 1:1000). Secondary antibodies were incubated for 1 h at RT either fluorescent (Li-COR

Biosciences IRDye680 and IRDye800 1:10.000: rb680, 925-68073; mm680, 925-68072; rb800, 925-32213)

or horseradish peroxidase-conjugated antibodies. Fluorescent signals were visualized using the Odyssey

Infrared Imaging System 2800 (Li-COR), for peroxidase-conjugated ECL detection, and images were

acquired with a Fusion Solo S (Vilber Lourmat Germany GmbH, Eberhardzell, Germany).

Immunofluorescence, Immunochemistry and TUNEL staining

The kidney tissue staining of 4 μm (cilia staining) and 2 μm (other stainings) fixed sections were

performed as previously described (Dafinger *et al.*, 2021). The primary antibodies (NF-kB, #8242 Cell

Signaling, 1:1000; CD3, MCA-1477 Biorad, 1:100; acetylated Tubulin, T6793 Sigma, 1:1000; Slc12a3,

HPA028748, Sigma Aldrich, 1:500; ARL13B, 17711-1-AP ProteinTech, 1:500) were incubated overnight at

4°C, followed by incubation with secondary antibodies. For immunochemistry, the DBA kit (K3468 DAKO,

30 min for NF-kB; SK-4105 Vector, 5 min for CD3) was used and samples counterstained with

Hematoxylin. For immunofluorescence, fluorophore-coupled antibodies (Jackson ImmunoResearch,

1:500: anti-mouse-Cy5, # 715-175-150; anti-rabbit A647, 711-605-152; anti-mouse-Cy3, 715-165-150;

and FITC-Lotus Tetragonolobus Lectin (LTL), FL-1321-2; Vector laboratories, 1:500) were used for 1 h at

RT. Samples were mounted after a short incubation with Hoechst 33342 (ThermoFisher Scientific,

1:5000) with ProLong™ Diamond (ThermoFisher Scientific). The DeadEnd™ Fluorometric TUNEL System

(Promega) was performed following the manufacturer's instructions, with the exception that the

samples were mounted, with a pre-incubation of Hoechst with ProLong™ Diamond. Fixed MEFs were

quenched with 50 mM NH_4Cl for 10 min, before permeabilisation with 0,3% PBS-TritonX-100 for 20 min.

Antibodies were diluted in Fish-Block (0.1 % (w/v) ovalbumin, 0.5 % (w/v) fish gelatine, in PBS),

supplemented with 0,3% TritonX-100. The primary antibody (ARL13B 1:800 cat# ab136648, Abcam;

acetylated Tubulin, 1:800 cat# T6793, Sigma) were incubated overnight at 4°C followed by the secondary

antibodies (anti-rabbit 488, A11034 Invitrogen, 1:400; anti-mouse 555, A31570, Invitrogen, 1:400; DAPI,

Roth, 1:400). Finally, coverslips were mounted with Fluoromount-G (ThermoFisher, 00-4958-02) and

imaged with the Leica microscope CTR6000, with DM6000B Laser and DFC360FX camera. MEF cell

images were deconvoluted with the Leica imaging software LASX. Cilia number and length were

determined with the open-source Fiji software (Schindelin *et al.*, 2012). Fixed BBS8[c.915delG] URECs were

blocked for 1 h in PBS containing 0.1 % triton-X-100 and 5 % normal donkey serum. Fixed mIMCD3 cells

were permeabilized for 10 min with 0.1 % triton X-100 diluted in PBS followed by a 1 h blocking step with

10 % NDS in PBS supplement with Ca^{2+} and Mg^{2+}. The primary antibody (ARL13B 1:1000 cat# ab136648,

Abcam; ARL13b 1:400, cat# 17711-1-AP, Proteintech; phospho-HDAC6 1:500 cat# PA5-105035,

Invitrogen; acetylated Tubulin 1:1000, Sigma; beta-Tubulin 1:50, cat# E7, DSHB) were incubated at 4°C.

Secondary Antibodies (donkey anti-mouse Cy3, cat# 715-165-150, Jackson ImmunoResearch; donkey

anti-rabbit A647, cat# 711-605-152, Jackson ImmunoResearch; DYKDDDDK Tag (L5), Alexa Fluor™ 488,

1:500; Invitrogen) were incubated at RT. Nuclei staining was performed with Hoechst 33342 Solution

(1:5000, cat# H3570 Invitrogen). Finally, coverslips were mounted with ProLong™ Diamond w/o DAPI

(cat# P36965, Thermo Fisher Scientific). URECs and IMCD3 cells were imaged with the AxioObserver

microscope with an axioCam ICc1, Axiocam 702 mono, Apoptome system (Carl Zeiss MicroImaging, Jena,

Germany). Images were exported through Zen Blue Software by Zeiss and aligned together using the free

design tool Inkscape.

Quantification and statistical analysis

Data are expressed as mean ± standard deviation (SD). All experiments were performed in at least 3

independent biological replicates. The data were statistically analyzed with GraphPad Prism version 9.5.1

unless otherwise mentioned. Statistical testing was performed as indicated in the figure legends.

Results and discussion

Cystic kidney disease in Bbs8 knockout mice is accompanied by inflammation and fibrosis

To investigate the function of BBS8 in the kidney, we analyzed the kidneys of a conventional $Bbs8^{-/-}$

mouse line (Tadenev *et al.*, 2011) where a mild renal phenotype with moderately dilated tubules in the

deep cortical region has previously been reported at 8 months of age. Consistent with this, no apparent

renal abnormalities were observed at 24 weeks. Therefore, we extended the observation period to 46

weeks. At this age, $Bbs8^{-/-}$ mice exhibited marked obesity, with a body weight approximately 30% higher

than that of control animals **(Fig. 1A)**, consistent with a typical BBS phenotype. In addition, signs of renal

involvement began to emerge, as indicated by a trend toward elevated serum urea levels **(Fig. 1B)** and a

slight increase in kidney weight **(Supp. Fig. 1A)**, although these changes did not reach statistical

significance. PAS staining of kidney sections revealed cyst formation and tubular dilatation **(Fig. 1C)**, with

a cyst index approximately six-fold higher in $Bbs8^{-/-}$ compared to control mice **(Fig. 1D)**. Further analysis

revealed that although the size of individual cysts detected in $Bbs8^{-/-}$ was similar to the size of dilated

tubules and cysts occasionally found in age-matched control animals **(Supp. Fig. 1B)**, the number of

dilated tubules and cysts was significantly increased **(Supp. Fig. 1C)**. Staining for proximal and distal

tubules in the kidney revealed that the majority of cysts originated from distal tubules, as they were

positive for Slc12a3 **(Fig. 1E)**. Furthermore, $Bbs8^{-/-}$ kidneys showed a significant increase in inflammation

and fibrosis in PAS and Masson's trichrome stained sections **(Fig. 1F)**. In line with inflammation and

fibrosis, we observed an accumulation of T-cells (CD3[+]), as well as an increased nuclear expression of the

NF-κB subunit RelA/p65, in particular in inflammation-rich areas (arrowheads) but also in kidney tubules

(arrows), in Bbs8[-/-] kidney tissue **(Fig. 1G)**. Increased TUNEL-positive cells indicated ongoing cell death in

Bbs8[-/-] kidneys **(Fig. 1H)**. To unravel the role of potential players involved in these processes of cell death,

we found the mRNA transcription of *RelA/p65*, as well as, of the pyroptosis markers *Nlrp3* and *GsdmD* to

be significantly increased **(Fig. 1I)**. Taken together, Bbs8[-/-] mice, in addition to obesity, fatty liver disease,

retinal degeneration, and occasional polydactyly, develop a relatively late-onset cystic fibrotic kidney

disease accurately reflecting the patient renal manifestation. Cyst formation predominantly originates

from the distal tubular segments and increased renal expression of genes associated with pyroptosis and

the inflammasome, key factors of renal fibrosis and inflammation.

The loss of Bbs8 affects ciliary tubulin acetylation and primary cilia stability

Since BBS8 is a component of the BBSome and the role of cilia in Bardet-Biedl syndrome (BBS) is well

established, we next examined primary cilia in kidney tissue from Bbs8[-/-] mice. In contrast to mouse

olfactory tissue, where a dramatic loss of cilia had been described (Tadenev *et al.*, 2011), we did not

observe any significant difference in the number and length of cilia in kidney sections of Bbs8[-/-] mice

compared to wildtype littermates **(Fig. 2A)**. Since quantification of cilia in tissue sections has technical

limitations, we also used mouse embryonic fibroblasts (MEFs) generated from Bbs8[-/-] and control mice. A

high percent of MEFs typically form cilia, and quantification of cilia number and length via ARL13B

staining revealed no significant difference in serum-starved MEFs **(Fig. 2B)**, confirming our finding in the

kidney sections. As an additional model, we isolated URECs from a BBS8 patient with a homozygous BBS8

variant (BBS8[c.915delG]) that results in a frameshift and premature protein termination (p.Met305Ilefs*15).

In these cells we re-expressed F.hBBS8 or F.GFP via lentiviral gene transfer. Expression was validated by

immunoblotting **(Suppl Fig. 2A)**. Consistently, cilia numbers as determined by ARL13B staining were not

significantly different in these cell lines **(Fig. 2C)**. Surprisingly, however, we observed a discrepancy

between ARL13B staining and acetylated tubulin, another classical ciliary marker: in kidney sections **(Fig.**

2E), as well as in MEFs **(Fig. 2F)** and URECs **(Fig. 2G)**, a significant number of cilia in the BBS8-deficient

group identified by ARL13B staining were negative for acetylated tubulin staining. Taking into account

that tubulin K40 acetylation is important for microtubule and, thus, ciliary stability, we analyzed this in

the BBS8[c.915delG] URECs with and without re-expression of F.hBBS8 by incubating the cells on ice to

depolymerize microtubules and destabilize cilia. This assay revealed that upon treatment, the amount of

cilia was significantly reduced in BBS8[c.915delG] F.GFP URECs as compared to the rescued cells **(Fig. 2H/I)**.

Thus, the loss of BBS8 does not affect cilia number but leads to a reduction of acetylated tubulin within

the cilium, which in turn impacts ciliary stability.

Proteomic analysis of Bbs8-deficient kidneys

To gain more insights into the kidney pathology and to decipher the mechanism contributing to the loss

of tubulin acetylation, we performed unbiased proteomic expression analysis of kidney lysates from

Bbs8$^{-/-}$ and control mice**.** The principal component analysis clearly separated the two genotypes in both

datasets **(Fig. 3A)**. After quality control (QC), we found a total of 2926 proteins in the whole proteome

(WP) **(Fig. 3B; Suppl. Table 1)**. Based on the Student's *t*-test (S0=0; FDR≤0.05), only 10 proteins were

significantly up- and 16 significantly downregulated in *Bbs8*$^{-/-}$ kidney lysates compared to control **(Fig.**

3C). Among those differentially expressed proteins were several proteins previously associated with

ciliopathies, cystogenesis, and kidney disease progression: For example, dynein cytoplasmic 2 heavy

chain 1 (DYNC2H1) and dystrobrevin binding protein 1 (DTNBP1) were found to be downregulated in the

knockout. DYNC2H1, a subunit of the IFT-dynein motor, drives retrograde IFT-rafts and plays a role in the

formation of the ciliary transition zone. Mutations lead to a skeletal ciliopathies, namely Jeune

asphyxiating thoracic dystrophy, short-rib polydactyly, and also to non-syndromic retinal degeneration

(Jensen *et al.*, 2018; Vig *et al.*, 2020; Hammarsjö *et al.*, 2021; Piceci-Sparascio *et al.*, 2023). Mutations in

DTNBP1, encoding a subunit of the biogenesis of lysosome-related organelles complex-1 (BLOC-1), cause

late-onset cystic kidney disease in mice (Monis, Faundez and Pazour, 2017). Proteins with increased

expression compared to control included retinol-binding protein 4 (RBP4), uromodulin (UMOD),

nucleoporin 98 (NUP98), and histone deacetylase 2 (HDAC2), each of which has previously been

associated with kidney diseases or primary cilia (Zaucke *et al.*, 2010; Kobayashi *et al.*, 2017; Endicott and

Brueckner, 2018; Xun *et al.*, 2018). Given our finding of decreased acetylated ciliary tubulin, we were

surprised to find neither the key acetylation enzyme αTAT1 nor the deacetylase HDAC6 to be

significantly altered. Previous studies have shown that HDAC6 upon phosphorylation by Aurora A kinase

(AurA) promotes tubulin deacetylation in cilia and ciliary disassembly (Pugacheva *et al.*, 2007). This

process can be prevented by BBS proteins through the recruitment of Inversin/NPHP2 to the ciliary base

(Patnaik *et al.*, 2019). However, AurA levels were not significantly changed, and phosphoproteomic (PP)

analyses performed in parallel **(Suppl. Fig. 3A/B; Suppl. Table 1)** also showed no alteration in HDAC6 or

AurA phosphorylation. Similarly, localization and expression levels of pHDAC6 in urine-derived

BBS8$^{c.915delG}$ URECs at the ciliary base was not altered **(Suppl. Fig. 3C)**. However, our data revealed a

significantly increased expression of another histone-deacetylase, HDAC2, in *Bbs8*$^{-/-}$ kidneys. This finding

could be further confirmed by Western blot experiments in both MEFs **(Fig. 3D)** and BBS8-deficient

URECs **(Fig. 3E)**.

BBS8 co-precipitates with HDAC2

Intrigued by upregulation of HDAC2 upon loss of BBS8, we questioned whether there might be a

connection between BBS8 and HDAC2 in the context of a shared protein complex. For an unbiased

interactomic analysis, we exogenously expressed F.hBBS8 or an equally FLAG-tagged control protein in

HEK293T cells and performed an anti-FLAG immunoprecipitation followed by MS/MS. The principal

component analysis clearly separated the F.hBBS8 pulldown from the control **(Fig. 4A)**. We identified

1151 proteins of which 986 were significantly enriched upon precipitation with BBS8 **(Fig. 4B, Suppl.**

Table 2). Notably, among those were many known members of the BBSome (BBS1, BBS2, BBS4, BBS5,

BBS7, BBS9) and the BBS chaperonin complex (CCT6A, CCT5, CCT3, CCT2, TCP1, CTT8, CCT4 and CCT7).

We again found HDAC2 as the only enriched histone-deacetylase. Additional protein groups shown to

potentially interact with BBS8 include heat shock binding proteins and proteins related to the Wnt

signaling pathway **(Supp. Fig. 4A)**. To confirm this potential interaction, we performed a similar co-

immunoprecipitation experiment and were able to detect endogenous HDAC2 upon pull down with

BBS8. **(Fig. 4C)**. Co-immunoprecipitation in the reverse direction confirmed BBS8 co-precipitating with

F9.HDAC2 **(Fig. 4D)**.

HDAC2 induces the deacetylation of alpha tubulin

HDAC2 has previously been associated with reduced cilia number in pancreatic ductal adenocarcinoma, a

phenomenon attributed to increased transcription of AurA and consequently enhanced ciliary

disassembly by HDAC6 (Kobayashi *et al.*, 2017). However, our data from *Bbs8*[-/-] tissue provided no

evidence for elevated AurA expression or HDAC6 activity. Therefore, we aimed to demonstrate that

HDAC2 directly influences tubulin acetylation. To this end, we transiently transfected HEK293 cells with

F.mHDAC2 or F.GFP and included F.hHDAC6 as a positive control. Immunoblot analysis of cell lysates

revealed that HDAC2 expression reduced of overall cellular levels of K40-acetylated α-tubulin, albeit to a

slightly lesser extent than HDAC6 **(Fig. 5A).** This functional similarity between HDAC2 and HDAC6 is

further supported by structural predictions using AlphaFold. The acetylated K40 residue of tubulin is

predicted to interact with the HDAC domain of HDAC2 in a manner similar to its interaction with the

second HDAC domain of HDAC6 **(Suppl. Fig. 5)**. To further substantiate that the loss of ciliary tubulin

acetylation in BBS8-deficient cells is attributable to HDAC2 activity, we employed the selective HDAC

inhibitor Tucidinostat (TUC). TUC is a specific low nanomolar inhibitor of the class I HDAC enzymes

(HDAC1/2/3; EC50: 100-200 nM) and class IIb HDAC enzymes (HDAC10; EC50: 100 nM) with almost no

effect on HDAC6 (EC50 >10 µM) and has already been used in a clinical trial for patients relapsed or

refractory peripheral T-cell lymphomas (Ho, Chan and Ganesan, 2020; Liu *et al.*, 2021; Utsunomiya *et al.*,

2022). We treated the urine-derived BBS8[c.915delG] URECs expressing F.GFP or F.BBS8 with TUC (160 nM) or

DMSO for 24 h in serum-free medium and stained for primary cilia with antibodies directed against

ARL13B and acetylated-tubulin. As before, we observed in the control group (DMSO) a significantly

decreased percentage of acetylated-tubulin positive cilia in the urine-derived BBS8[c.915delG] URECS as

compared to BBS8 re-expression. However, HDAC2-inhibition significantly increased the percentage of

acetylated-tubulin positive cilia in BBS8[c.915delG] cells, to a similar level as BBS8 re-expression did **(Fig. 5B)**.

In summary, we conclude that loss of BBS8 results in a pathological upregulation of HDAC2 activity,

which targets acetylated α-tubulin within the cilium, thereby perturbing ciliary dynamics and stability.

Consistent with the relatively mild and late-onset kidney phenotype, these changes do not result in

major alterations of cilia and have no impact on cilia length or number. Nevertheless, these subtle

modifications in primary cilia architecture may affect ciliary dynamics and, over time, negatively

influence the homeostasis of renal tissue. Thus, HDAC2 may indeed represent a promising therapeutic

target. Supporting this notion, a chemical screen in two zebrafish models of polycystic kidney disease

(PKD) identified both a pan-HDAC inhibitor and a class I HDAC inhibitor as suppressors of cyst formation

(Cao *et al.*, 2009). Our study provides a mechanistic basis for this positive effect of HDAC inhibitors by

identifying the HDAC2-induced acetylation-defect of cilia which can be reversed by a class 1 inhibitor as

potential driver of cystogenesis in BBS8, linked to subtle ciliary defects.

Data Availability

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via

the PRIDE (Perez-Riverol *et al.*, 2024) partner repository with the dataset identifier PXD064988

(interactome) and PXD064990 (proteomeBBS8-/- kidneys). Login details are available from the authors

upon request.

Figure legends:

Figure 1. Bbs8[-/-] mice develop late-onset cystic kidney disease with fibrosis and inflammation.

46-week-old *Bbs8[-/-]* compared to control (*Bbs8[+/+]*) mice showed a significantly increased **(A)** body weight

(n=4) and an increasing trend **(B)** of measured urea level of blood serum (n=4). **(C)** PAS staining of the

whole kidney in control and *Bbs8[-/-]* animals, showing cyst formation in the knockout; Scale bar: 1 mm. **(D)**

Cyst index described in a cyst-to-tissue area ratio, normalized to control animals (n=4). **(E)** Staining of

distal (Slc12a3, magenta) and proximal tubules (LTL, green) in the kidney showing cyst formation in the

distal convoluted tubules; Scale bar: 500 µm. **(F)** Representative microscopic images of the PAS staining

(higher magnified area), showing clusters of inflammation (Arrow; Scale bar: 200 µm). The additional

Masson's trichrome staining, displayed fibrosis in *Bbs8*$^{-/-}$ animals (Arrowhead; Scale bar: 300 µm). **(G)**

Representative microscopic images of CD3$^+$ and NF-kB (RelA/p65) expression revealed inflammation in

Bbs8$^{-/-}$ animals; Scale bar: 300 µm. **(H)** TUNEL staining of control and Bbs8$^{-/-}$ kidney tissue; Scale bar: 100

µm. **(I)** Quantitative real-time PCR of *RelA/p65* and inflammasome genes in control and *Bbs8*$^{-/-}$ kidney

samples (n=4). All of the statistical analyses were performed using a two-sided Student's *t*-test (*p*-value:

<0.001***; 0.002**; 0.033*; ns⬛=⬛0.12).

Figure 2. BBS8 deficiency results in reduced ciliary tubulin acetylation and stability without significantly

affecting the number or length of cilia.

(A) Control and *Bbs8*$^{-/-}$ kidney tissue stained for cilia using ARL13B antiserum (Scale bar: 50 µm). **(B)** *Bbs8*$^-$

$^{/-}$ and control mouse embryonic fibroblasts (MEF) quantified for the number and length of cilia by

ARL13B staining (n=4), statistically analyzed using an unpaired Student's *t*-test (*p*-value: <0.001***;

0.002**; 0.033*; ns⬛=⬛0.12). **(C)** BBS8$^{c.915delG}$ and BBS8 re-expressing URECs stained for cilia using ARL13B

antiserum (n=7), statistically analyzed using an unpaired Student's *t*-test (*p*-value: <0.001***; 0.002**;

0.033*; ns⬛=⬛0.12). **(D-F)** Representative images of primary cilia co-stained for ARL13B (green) and

acetylated-tubulin (ac-Tub, magenta; scale bar: 10 µm); graphs showing the percentage of cells with a ac-

Tub positive cilium from the total amount of cilia, as identified by ARL13B staining. Arrowhead indicates

ac-Tub negative cilia. Statistically analyzed using an unpaired Student's *t*-test (*p*-value: <0.001***;

0.002**; 0.033*; ns⬛=⬛0.12). Staining and analysis were performed in **(D)** control and *Bbs8*$^{-/-}$ kidney

tissue (n=3), **(E)** cultured *Bbs8*$^{-/-}$ and control MEFs (n=4), **(F)** BBS8$^{c.915delG}$ patient-isolated URECs stably

expressing either F.hBBS8 or F9.GFP (n=4). **(G)** Cilia stabilization assay was performed in BBS8$^{c.915delG}$

patient-isolated URECs stably expressing either F.hBBS8 or F.GFP. Destabilization was induced by keeping

the cells for 30 min on ice. Representative images of primary cilia stained for ARL13B (green; n=3; Scale

bar: 10 µm). Statistical analysis was performed by using a one-way ANOVA followed by a two-sided

Student's *t*-test (*p*-value: >0.001***; 0.002**; 0.033*; ns⬛=⬛0.12).

Figure 3. HDAC2 is one of the few proteins significantly upregulated in BBS8$^{-/-}$ *kidney tissue*

(A) Principal component analysis (PCA) plot of the whole proteome (WP) of control and *Bbs8*$^{-/-}$ kidney

samples. The axes represent the percentages of variation explained by the principal components, three

biological replicates per group. **(B/C)** Scatter blot of the proteome. Significantly (FDR≤0.05) up-regulated

(yellow, 1-10) or downregulated (cyan, 1-16) proteins are marked and additionally listed in the table. **(D)**

Lysates from control and *Bbs8*[-/-] mouse embryonic fibroblasts (MEFs) were immunoblotted against

HDAC2, with Calnexin as loading control. **(E)** Urine-derived BBS8[c.915delG] URECs stably expressing either

F.hBBS8 or F.GFP were immunoblotted against HDAC2 with Calnexin as a loading control.

Figure 4. BBS8 and HDAC2 are components of a common protein complex

(A) Principal component analysis (PCA) plot of the interactome data of F.hEPS and F.hBBS8 in HEK293T

cells. The axes represent the percentages of variation explained by the principal components. Four

individual replicates were performed. **(B)** Scatter plot of the interactome, marked with potential

interactors of BBS8 (FDR≤0.05) in magenta. **(C)** Co-IP from HEK293T cells, expressing F.hBBS8 or F.hEPS

as control. Immuno-precipitation from lysates was done with α-FLAG (IP), immunoblot was performed

with α-HDAC2 and with α-FLAG. **(D)** Co-IP from HEK293T cells expressing F.mHDAC2 or F.hEPS together

with eGFP.hBBS8. Immuno-precipitation from lysates was performed with α-FLAG (IP), immunoblot with

α-eGFP and α-FLAG.

Figure 5. HDAC2 promotes tubulin deacetylation, and its inhibition increases tubulin acetylation in BBS8-

deficient cells.

(A) Lysates of HEK293T cells transfected with F.hEPS, F.mHDAC2, F.hHDAC6 and co-transfected with

F.GFP were immunoblotted with α-acetylated tubulin and α-FLAG. F.GFP expression was used as control

for quantitative comparison (n=3) with the statistical analysis of a one-way ANOVA followed by a two-

sided Student's *t*-test (*p*-value: >0.001***; 0.002**; 0.033*; ns=0.12). Data were normalized to the

acetylated tubulin amount in F.hEPS expressing cells. **(B)** BBS8[c.915delG] patient-isolated URECs stably

(re)expressing either F.hBBS8 or F.GFP were treated with the HDAC2 inhibitor Tucidinostat (TUC, 160

nM) or DMSO as control. Representative images of primary cilia co-stained with the cilia marker ARL13B

(green) and acetylated-tubulin (ac-Tub, magenta; scale bar: 10 µm) displaying the restoration of

acetylation within primary cilia in F.GFP BBS8[c.915delG] URECs upon HDAC2 inhibition (n=4; arrowheads: ac-

Tub positive cilia; arrows: ac-Tub negative cilia); graph showing the percentage of acetylated-tubulin

positive cilia of the total amount of cilia determined by ARL13B staining. Statistical analysis was

performed by using a one-way ANOVA followed by a two-sided Student's *t*-test (*p*-value: >0.001***;

0.002**; 0.033*; ns = 0.12).

Supplementary Figure 1. Small-cyst formation in Bbs8$^{-/-}$ kidney and systemic inflammation.

(A) Kidney weight of 46-week-old control and *Bbs8*$^{-/-}$ mice did not significantly alter (n=4). **(B)** The

measured cyst area is depicted as jittered dots with the mean as a horizontal line. The values are plotted

on a log$_{10}$ scale. Single individuals are plotted separately (n=4). **(C)** The overall number of cysts is

significantly increased in *Bbs8*$^{-/-}$ animals (n=4). For statistical analysis, a two-sided Student's *t*-test *was*

used (*p*-value: <0.001***; 0.002**; 0.033*; ns = 0.12).

Supplementary Figure 2. Re-expression of BBS8 in BBS8$^{c.915delG}$ URECs

(A) Urine-derived BBS8$^{c.915delG}$ URECS stably expressing either F.hBBS8 or F.GFP, were immunoblotted

against FLAG with Calnexin as control.

Supplementary Figure 3. Additional analyses of the phosphoproteome data

(A) Principal component analysis (PCA) plot of the phosphoproteome (PP) data of control and *Bbs8*$^{-/-}$

kidney samples. The axes represent the percentages of variation explained by the principal components.

Three individual replicates were performed. **(B)** Analysis of the PP identified 6223 phosphosites, within a

total of 2188 individual proteins from which 151 were significantly up- and 1041 were significantly down-

regulated. Among these were multiple HDAC2 phospho-peptides containing 3 serine (Ser) residues:

Ser394, Ser422 and Ser424. Notably, HDAC6 was not found to be changed in its phosphorylation. **(C)** Co-

Staining for the cilia (ARL13B, green) and phopho-HDAC6 (orange) in urine-derived BBS8$^{c.915delG}$ URECs re-

expressing F.hBBS8 or F.GFP (Scale bar: 10 μm). Ciliary base is indicated by arrowheads.

Supplementary Figure 4. BBS8 interactors involved in cilia signaling pathways

(A) Scatter plot of different protein clusters expressed in the BBS8 interactome for different GO-terms:

BBSome, chaperonin-containing T-complex, heat shock protein binding and Wnt signaling pathway.

Supplementary Figure 5. Modeling the interaction of HDAC2 and HDAC6 with K40-acetylated alpha

Tubulin.

(A) Human TBA1A (UniProt ID: Q71U36) with the PTM "N6-acetyl-L-lysine" added to K40 was co-

predicted in AlphaFold with the HDAC domain of HDAC2 (Q92769; residues 9–322) or the second HDAC

domain of HDAC6 (Q9UBN7; residues 482–800), a well-characterized tubulin deacetylase. The ipTM and

pTM values were obtained from the AlphaFold server. The acetylated residue K40 is highlighted in red.

Supplementary Figure 6. Original data: full-sized immunoblots.

Original western blots only cropped to gel size **(A)** of Supp. Figure 2 A, **(B)** of Figure 3 D, **(C)** of Figure 3 E,

(D) of Figure 4 C, **(E)** of Figure 4 D and **(F)** of Figure 5 A. To visualize the expression of F.hBBS8, the

membrane was captured with a short and longer exposure time.

References:

Abramson, J. *et al.* (2024) 'Accurate structure prediction of biomolecular interactions with AlphaFold 3',

Nature 2024 630:8016, 630(8016), pp. 493–500. doi: 10.1038/s41586-024-07487-w.

Bankhead, P. *et al.* (2017) 'QuPath: Open source software for digital pathology image analysis', *Scientific*

Reports 2017 7:1, 7(1), pp. 1–7. doi: 10.1038/s41598-017-17204-5.

Beales, P. L. *et al.* (1999) 'New criteria for improved diagnosis of Bardet-Biedl syndrome: results of a

population survey', *Journal of Medical Genetics*, 36(6), p. 437. doi: 10.1136/jmg.36.6.437.

Bentley-Ford, M. R. *et al.* (2021) 'A mouse model of BBS identifies developmental and homeostatic

effects of BBS5 mutation and identifies novel pituitary abnormalities', *Human molecular genetics*, 30(3–

4), pp. 234–246. doi: 10.1093/HMG/DDAB039.

Brücker, L. *et al.* (2023) 'The actin-bundling protein Fascin-1 modulates ciliary signalling', *Journal of*

Molecular Cell Biology, 15(4), p. 22. doi: 10.1093/JMCB/MJAD022.

Cao, Y. *et al.* (2009) 'Chemical modifier screen identifies HDAC inhibitors as suppressors of PKD models',

Proceedings of the National Academy of Sciences of the United States of America, 106(51), pp. 21819–

21824. doi: 10.1073/PNAS.0911987106.

Cognard, N. *et al.* (2015) 'Comparing the Bbs10 complete knockout phenotype with a specific renal

epithelial knockout one highlights the link between renal defects and systemic inactivation in mice', *Cilia*,

4. Available at: https://ciliajournal.biomedcentral.com/articles/10.1186/s13630-015-0019-8 (Accessed:

30 March 2021).

Dafinger, C. *et al.* (2021) 'Targeted deletion of Ruvbl1 results in severe defects of epidermal development

and perinatal mortality', *Molecular and Cellular Pediatrics*, 8(1). doi: 10.1186/S40348-021-00111-1.

Davis, R. E. *et al.* (2007) 'A knockin mouse model of the Bardet-Biedl syndrome 1 M390R mutation has

cilia defects, ventriculomegaly, retinopathy, and obesity', *Proceedings of the National Academy of*

Sciences of the United States of America, 104(49), pp. 19422–19427. doi: 10.1073/PNAS.0708571104.

Dilan, T. L. *et al.* (2018) 'Bardet-Biedl syndrome-8 (BBS8) protein is crucial for the development of outer

segments in photoreceptor neurons.', *Human molecular genetics*, 27(2), pp. 283–294. doi:

10.1093/hmg/ddx399.

Elawad, O. A. M. A. *et al.* (2022) 'Bardet–Biedl syndrome: a case series', *Journal of Medical Case Reports*,

16(1), pp. 1–9. doi: 10.1186/S13256-022-03396-6/FIGURES/4.

Endicott, S. J. and Brueckner, M. (2018) 'NUP98 Sets the Size-Exclusion Diffusion Limit through the Ciliary

Base', *Current biology : CB*, 28(10), pp. 1643-1650.e3. doi: 10.1016/J.CUB.2018.04.014.

Fan, C.-M. and Tessier-Lavigne, M. (1994) 'Patterning of mammalian somites by surface ectoderm and

notochord: Evidence for sclerotome induction by a hedgehog homolog', *Cell*, 79(7), pp. 1175–1186. doi:

10.1016/0092-8674(94)90009-4.

Fath, M. A. *et al.* (2005) 'Mkks-null mice have a phenotype resembling Bardet-Biedl syndrome', *Human*

molecular genetics, 14(9), pp. 1109–1118. doi: 10.1093/HMG/DDI123.

Forsythe, E. *et al.* (2017) 'Risk factors for severe renal disease in bardet-biedl syndrome', *Journal of the*

American Society of Nephrology, 28(3), pp. 963–970. doi: 10.1681/ASN.2015091029.

Forsythe, E. and Beales, P. L. (2013) 'Bardet–Biedl syndrome', *European Journal of Human Genetics*,

21(1), pp. 8–13. doi: 10.1038/ejhg.2012.115.

Garcia-Gonzalo, F. R. and Reiter, J. F. (2017) 'Open Sesame: How Transition Fibers and the Transition

Zone Control Ciliary Composition', *Cold Spring Harbor perspectives in biology*, 9(2). doi:

10.1101/CSHPERSPECT.A028134.

Gopalakrishnan, J. *et al.* (2023) 'Emerging principles of primary cilia dynamics in controlling tissue

organization and function', *The EMBO Journal*, p. e113891. doi: 10.15252/EMBJ.2023113891.

Guo, D. F. *et al.* (2011) 'Inactivation of Bardet-Biedl syndrome genes causes kidney defects', *American*

Journal of Physiology - Renal Physiology, 300(2), p. F574. doi: 10.1152/AJPRENAL.00150.2010.

Habbig, S. *et al.* (2011) 'NPHP4, a cilia-associated protein, negatively regulates the Hippo pathway', *The*

Journal of cell biology, 193(4), pp. 633–642. doi: 10.1083/JCB.201009069.

Hammarsjö, A. *et al.* (2021) 'High diagnostic yield in skeletal ciliopathies using massively parallel genome

sequencing, structural variant screening and RNA analyses', *Journal of Human Genetics 2021 66:10*,

66(10), pp. 995–1008. doi: 10.1038/s10038-021-00925-x.

Hildebrandt, F., Benzing, T. and Katsanis, N. (2011) 'Ciliopathies', *The New England journal of medicine*,

364(16), pp. 1533–1543. doi: 10.1056/NEJMRA1010172.

Hildebrandt, F. and Zhou, W. (2007) 'Nephronophthisis-associated ciliopathies.', *Journal of the American*

Society of Nephrology : JASN, 18(6), pp. 1855–71. doi: 10.1681/ASN.2006121344.

Ho, T. C. S., Chan, A. H. Y. and Ganesan, A. (2020) 'Thirty Years of HDAC Inhibitors: 2020 Insight and
Hindsight', *Journal of Medicinal Chemistry*, 63(21), pp. 12460–12484. doi:
10.1021/ACS.JMEDCHEM.0C00830/ASSET/IMAGES/LARGE/JM0C00830_0023.JPEG.

Jensen, V. L. *et al.* (2018) 'Role for intraflagellar transport in building a functional transition zone', *EMBO*
reports, 19(12). doi: 10.15252/EMBR.201845862.

Kobayashi, T. *et al.* (2017) 'HDAC2 promotes loss of primary cilia in pancreatic ductal adenocarcinoma',
EMBO reports, 18(2), pp. 334–343. doi: 10.15252/EMBR.201541922.

Kretschmer, V. *et al.* (2019) 'Progressive Characterization of Visual Phenotype in Bardet-Biedl Syndrome
Mutant Mice', *Investigative ophthalmology & visual science*, 60(4), pp. 1132–1143. doi: 10.1167/IOVS.18-
25210.

Kretschmer, V. *et al.* (2023) 'Deletion of IFT20 exclusively in the RPE ablates primary cilia and leads to
retinal degeneration', *PLoS biology*, 21(12). doi: 10.1371/JOURNAL.PBIO.3002402.

Kulaga, H. M. *et al.* (2004) 'Loss of BBS proteins causes anosmia in humans and defects in olfactory cilia
structure and function in the mouse', *Nature genetics*, 36(9), pp. 994–998. doi: 10.1038/NG1418.

Liu, W. *et al.* (2021) 'A Multi-Center, Real-World Study of Chidamide for Patients With Relapsed or
Refractory Peripheral T-Cell Lymphomas in China', *Frontiers in Oncology*, 11, p. 750323. doi:
10.3389/FONC.2021.750323/BIBTEX.

Loktev, A. V. *et al.* (2008) 'A BBSome Subunit Links Ciliogenesis, Microtubule Stability, and Acetylation',
Developmental Cell, 15(6), pp. 854–865. doi: 10.1016/j.devcel.2008.11.001.

May-Simera, H. L. *et al.* (2010) 'Bbs8, together with the planar cell polarity protein Vangl2, is required to
establish left-right asymmetry in zebrafish', *Developmental biology*, 345(2), pp. 215–225. doi:
10.1016/J.YDBIO.2010.07.013.

May-Simera, H. L. *et al.* (2015) 'Ciliary proteins Bbs8 and Ift20 promote planar cell polarity in the
cochlea', *Development (Cambridge, England)*, 142(3), pp. 555–566. doi: 10.1242/DEV.113696.

McConnachie, D. J., Stow, J. L. and Mallett, A. J. (2021) 'Ciliopathies and the Kidney: A Review', *American*
journal of kidney diseases : the official journal of the National Kidney Foundation, 77(3), pp. 410–419.
doi: 10.1053/J.AJKD.2020.08.012.

Meng, E. C. *et al.* (2023) 'UCSF ChimeraX: Tools for structure building and analysis', *Protein Science*,

32(11), p. e4792. doi: 10.1002/PRO.4792.

Monis, W. J., Faundez, V. and Pazour, G. J. (2017) 'BLOC-1 is required for selective membrane protein
trafficking from endosomes to primary cilia', *The Journal of cell biology*, 216(7), pp. 2131–2150. doi:
10.1083/JCB.201611138.

Munsterberg, A. E. *et al.* (1995) 'Combinatorial signaling by Sonic hedgehog and Wnt family members
induces myogenic bHLH gene expression in the somite.', *Genes & Development*, 9(23), pp. 2911–2922.
doi: 10.1101/gad.9.23.2911.

Mykytyn, K. *et al.* (2004) 'Bardet-Biedl syndrome type 4 (BBS4)-null mice implicate Bbs4 in flagella
formation but not global cilia assembly', *Proceedings of the National Academy of Sciences of the United*
States of America, 101(23), pp. 8664–8669. doi: 10.1073/PNAS.0402354101.

Nachury, M. V. *et al.* (2007) 'A Core Complex of BBS Proteins Cooperates with the GTPase Rab8 to
Promote Ciliary Membrane Biogenesis', *Cell*, 129(6), pp. 1201–1213. doi: 10.1016/j.cell.2007.03.053.

Nishimura, D. Y. *et al.* (2004) 'Bbs2-null mice have neurosensory deficits, a defect in social dominance,
and retinopathy associated with mislocalization of rhodopsin', *Proceedings of the National Academy of*
Sciences of the United States of America, 101(47), pp. 16588–16593. doi: 10.1073/PNAS.0405496101.

Patnaik, S. R. *et al.* (2019) 'Bardet–Biedl Syndrome proteins regulate cilia disassembly during tissue
maturation', *Cellular and Molecular Life Sciences*, 76(4), pp. 757–775. doi: 10.1007/s00018-018-2966-x.

Pazour, G. J. and Bloodgood, R. A. (2008) 'Targeting Proteins to the Ciliary Membrane', *Current Topics in*
Developmental Biology, 85, pp. 115–149. doi: 10.1016/S0070-2153(08)00805-3.

Pettersen, E. F. *et al.* (2021) 'UCSF ChimeraX: Structure visualization for researchers, educators, and
developers', *Protein science : a publication of the Protein Society*, 30(1), pp. 70–82. doi:
10.1002/PRO.3943.

Perez-Riverol, J. *et al.* (2024) 'The PRIDE database at 20 years: 2025 update', *Nucleic Acids Res*. doi:
10.1093/nar/gkae1011

Piceci-Sparascio, F. *et al.* (2023) 'Clinical variability in DYNC2H1-related skeletal ciliopathies includes Ellis-
van Creveld syndrome', *European Journal of Human Genetics 2023 31:4*, 31(4), pp. 479–484. doi:
10.1038/s41431-022-01276-7.

Postma, M. and Goedhartid, J. (2019) 'PlotsOfData-A web app for visualizing data together with their

summaries'. doi: 10.1371/journal.pbio.3000202.

Pourquié, O. *et al.* (1996) 'Lateral and Axial Signals Involved in Avian Somite Patterning: A Role for
BMP4', *Cell*, 84(3), pp. 461–471. doi: 10.1016/S0092-8674(00)81291-X.

Pugacheva, E. N. *et al.* (2007) 'HEF1-dependent Aurora A activation induces disassembly of the primary
cilium', *Cell*, 129(7), pp. 1351–1363. doi: 10.1016/J.CELL.2007.04.035.

Putoux, A. *et al.* (2012) 'Phenotypic variability of Bardet-Biedl syndrome: focusing on the kidney',
Pediatric Nephrology, 27(1), pp. 7–15. doi: 10.1007/s00467-010-1751-3.

Rahmouni, K. *et al.* (2008) 'Leptin resistance contributes to obesity and hypertension in mouse models of
Bardet-Biedl syndrome', *The Journal of clinical investigation*, 118(4), pp. 1458–1467. doi:
10.1172/JCI32357.

Reiter, J. F., Blacque, O. E. and Leroux, M. R. (2012) 'The base of the cilium: roles for transition fibres and
the transition zone in ciliary formation, maintenance and compartmentalization', *EMBO Reports*, 13(7),
p. 608. doi: 10.1038/EMBOR.2012.73.

Reiter, J. F. and Leroux, M. R. (2017) 'Genes and molecular pathways underpinning ciliopathies', *Nature*
Reviews Molecular Cell Biology 2017 18:9, 18(9), pp. 533–547. doi: 10.1038/nrm.2017.60.

Rödig, N. *et al.* (2022) 'Behavioral Phenotyping of Bbs6 and Bbs8 Knockout Mice Reveals Major
Alterations in Communication and Anxiety', *International Journal of Molecular Sciences*, 23(23). doi:
10.3390/IJMS232314506/S1.

Schindelin, J. *et al.* (2012) 'Fiji: an open-source platform for biological-image analysis', *Nature Methods*
2012 9:7, 9(7), pp. 676–682. doi: 10.1038/nmeth.2019.

Schneider, S. *et al.* (2021) 'Loss of Ciliary Gene Bbs8 Results in Physiological Defects in the Retinal
Pigment Epithelium', *Frontiers in Cell and Developmental Biology*, 9, p. 607121. doi:
10.3389/FCELL.2021.607121/FULL.

Seo, S. *et al.* (2009) 'Requirement of Bardet-Biedl syndrome proteins for leptin receptor signaling',
Human molecular genetics, 18(7), pp. 1323–1331. doi: 10.1093/HMG/DDP031.

Seo, S. *et al.* (2010) 'BBS6, BBS10, and BBS12 form a complex with CCT/TRiC family chaperonins and
mediate BBSome assembly.', *Proceedings of the National Academy of Sciences of the United States of*
America, 107(4), pp. 1488–93. doi: 10.1073/pnas.0910268107.

Sieckmann, K. *et al.* (2024) 'BBSome-dependent ciliary Hedgehog signaling governs cell fate in the white
adipose tissue', *bioRxiv*, p. 2024.01.25.576526. doi: 10.1101/2024.01.25.576526.

Tadenev, A. L. D. *et al.* (2011) 'Loss of Bardet-Biedl syndrome protein-8 (BBS8) perturbs olfactory
function, protein localization, and axon targeting', *Proceedings of the National Academy of Sciences of*
the United States of America, 108(25), pp. 10320–10325. doi: 10.1073/PNAS.1016531108.

Todaro, G. J. and Green, H. (1963) 'Quantitative studies of the growth of mouse embryo cells in culture
and their development into established lines', *The Journal of cell biology*, 17(2), pp. 299–313. doi:
10.1083/JCB.17.2.299.

Tyanova, S. *et al.* (2016) 'The Perseus computational platform for comprehensive analysis of
(prote)omics data', *Nature Methods 2016 13:9*, 13(9), pp. 731–740. doi: 10.1038/nmeth.3901.

Tyanova, S., Temu, T. and Cox, J. (2016) 'The MaxQuant computational platform for mass
spectrometry–based shotgun proteomics', *Nature Protocols*, 11. doi: 10.1038/nprot.2016.136.

Utsunomiya, A. *et al.* (2022) 'Oral histone deacetylase inhibitor tucidinostat (HBI-8000) in patients with
relapsed or refractory adult T-cell leukemia/lymphoma: Phase IIb results', *Cancer Science*, 113(8), pp.
2778–2787. doi: 10.1111/CAS.15431.

Vig, A. *et al.* (2020) 'DYNC2H1 hypomorphic or retina-predominant variants cause nonsyndromic retinal
degeneration', *Genetics in Medicine*, 22(12), p. 2041. doi: 10.1038/S41436-020-0915-1.

Xun, C. *et al.* (2018) 'Circulating RBP4 Increase and Its Diagnosis of Chronic Kidney Disease', *Annals of*
clinical and laboratory science, 48(2), pp. 205–207. Available at:
https://pubmed.ncbi.nlm.nih.gov/29678848/ (Accessed: 20 February 2024).

Zaucke, F. *et al.* (2010) 'Uromodulin is expressed in renal primary cilia and UMOD mutations result in
decreased ciliary uromodulin expression', *Human Molecular Genetics*, 19(10), p. 1985. doi:
10.1093/HMG/DDQ077.

Zhang, Q. *et al.* (2012) 'Intrinsic protein-protein interaction-mediated and chaperonin-assisted sequential
assembly of stable bardet-biedl syndrome protein complex, the BBSome.', *The Journal of biological*
chemistry, 287(24), pp. 20625–35. doi: 10.1074/jbc.M112.341487.

Acknowledgements:

We would like to express our sincere gratitude to Angelika Köser, Serena Greco-Torres, and Stefanie Keller for their exceptional technical assistance. We also deeply appreciate the outstanding support provided by the CECAD Imaging Facility and the CECAD Proteomics Facility. Our heartfelt thanks go to all members of our research group. Lastly, we extend our appreciation to the patient and their advocate for their generous contributions and commitment to advancing research. We would also like to thank Edward Seto for kindly providing us with the HDAC2 Plasmid. Molecular graphics and analyses were performed with UCSF ChimeraX, developed by the Resource for Biocomputing, Visualization, and Informatics at the University of California, San Francisco, with support from National Institutes of Health R01-GM129325 and the Office of Cyber Infrastructure and Computational Biology, National Institute of Allergy and Infectious Diseases (Meng *et al.*, 2023). Funding: This study was supported by the German Research Foundation (DFG; SFB1403, project number 414786233, A09 to B.S. and T.B.; and FOR5547 to HMS, DW, and BS, project number 503306912, under Germany's Excellence Strategy – EXC2151 – Project-ID 390873048 (to D.W.), and WA 3382/8-1 – Project-ID 513767027 (to D.W.). The NEOCYST consortium (Network for Early Onset Cystic Kidney Diseases) supported the project, with funding provided by the German Ministry of Education and Research (BMBF) under grant agreement number 01GM1515D. This funding (to B.S., M.C.L. and M.C.) had no impact on the selection of the topic or the content of the position paper. B.S. and M.C.L.. were also supported by the European Union (TheraCil). P.M. was supported by the Studienstiftung des Deutschen Volkes. The CECAD proteomics facility was supported by the large instrument grant INST 1856/71-1 FUGG by the German Research Foundation (DFG Großgeräteantrag).

Author contributions:

EK, HMS and BS designed the study. EK and JG performed most of the experiments. PAM prepared the experiments for MEFs. LKE performed and analysed the BBS8 interactome MS/MS experiments. HMS, CK, DW, MC, MCL, TB provided materials and support for the interpretation of results. EK assembled the final figures. EK and BS wrote the original draft. All of the authors discussed the project and contributed to the final version of the manuscript.

Figure 1

Figure 2

Figure 3

Figure 4

A

HEKs

B

URECs

Figure 5

Supp. Figure 1

A

URECs

A

control ■ *Bbs8⁻ᐟ⁻* ●

PP

B

n: 6223
up: 151
down: 1041

HDAC2_Ser424

HDAC2_Ser422/424

HDAC6

HDAC2_Ser422

HDAC2_Ser394

HDAC6

Student's *t*-test difference *Bbs8⁻ᐟ⁻* vs control

HEKs

C

URECs

A

A

HDAC domain of
HDAC2 (9-322)

acetylated K40

Tubulin alpha-1A
(TBA1A; K40 acetylated)

ipTM=0,78; pTM=0,64)

2nd HDAC domain of
HDAC6 (482-800)

acetylated K40

Tubulin alpha-1A
(TBA1A; K40 acetylated)

ipTM=0,85; pTM=0,71)

Supp. Figure 5

Supp. Figure 6