HPV18 E7 inhibits LATS1 kinase and activates YAP1 by degrading PTPN14

William J. Blakely[1], Joshua Hatterschide[1,2], Elizabeth A. White[1*]

[1]Department of Otorhinolaryngology: Head and Neck Surgery, University of Pennsylvania

Perelman School of Medicine, Philadelphia, PA, USA

[2]Current address: Department of Integrative Immunobiology, Duke University School of

Medicine, Durham, NC, USA

*Correspondence: Elizabeth A. White, eawhite@pennmedicine.upenn.edu

Classification: Biological Sciences, Microbiology

Keywords: Human papillomavirus, differentiation, tumor suppressor, keratinocyte, YAP1, Hippo,

PTPN14

17
18

ABSTRACT

High-risk human papillomavirus (HPV) oncoproteins inactivate cellular tumor

suppressors to reprogram host cell signaling pathways. HPV E7 proteins bind and degrade the

tumor suppressor PTPN14, thereby promoting the nuclear localization of the YAP1 oncoprotein

and inhibiting keratinocyte differentiation. YAP1 is a transcriptional coactivator that drives

epithelial cell stemness and self-renewal. YAP1 activity is inhibited by the highly conserved

Hippo pathway, which is frequently inactivated in human cancers. MST1/2 and LATS1/2 kinases

form the core of the Hippo kinase cascade. Active LATS1 kinase is phosphorylated on threonine

1079 and inhibits YAP1 by phosphorylating it on amino acids including serine 127. Here, we

tested the effect of high-risk (carcinogenic) HPV18 E7 on Hippo pathway activity. We found that

either PTPN14 knockout or PTPN14 degradation by HPV18 E7 decreased phosphorylation of

LATS1 T1079 and YAP1 S127 in human keratinocytes and inhibited keratinocyte differentiation.

Conversely, PTPN14-dependent differentiation required LATS kinases and certain PPxY motifs

in PTPN14. Neither MST1/2 kinases nor the putative PTPN14 phosphatase active site were

required for PTPN14 to promote differentiation. Taken together, these data support that

degradation of PTPN14 by HPV18 E7 reduces LATS1 activity, promoting active YAP1 and

inhibiting keratinocyte differentiation.

SIGNIFICANCE

The Hippo kinase cascade inhibits YAP1, an oncoprotein and driver of cell stemness and

self-renewal. There is mounting evidence that the Hippo pathway is targeted by tumor viruses

including human papillomavirus. The high-risk HPV E7 oncoprotein promotes YAP1 nuclear

localization and the carcinogenic activity of high-risk HPV E7 requires YAP1 activity. Blocking

HPV E7-dependent YAP1 activation could inhibit HPV-mediated carcinogenesis, but the

mechanism by which HPV E7 activates YAP1 has not been elucidated. Here we report that by

degrading the tumor suppressor PTPN14, HPV18 E7 inhibits LATS1 kinase, reducing inhibitory

phosphorylation on YAP1. These data support that an HPV oncoprotein can inhibit Hippo

signaling to activate YAP1 and strengthen the link between PTPN14 and Hippo signaling in

human epithelial cells.

INTRODUCTION

Human papillomaviruses are double-stranded DNA viruses with a tropism for

keratinocytes in stratified squamous epithelia (1). The human papillomavirus (HPV) E6 and E7

proteins target host cellular proteins to reprogram cell signaling pathways, thereby enabling

HPV replication and persistence. A subset of HPVs are the 'high-risk' genotypes that cause

many anogenital and oropharyngeal cancers (2). High-risk HPV E6 and E7 proteins have

biological activities that are not shared among the E6 and E7 proteins encoded by other HPVs,

and high-risk HPV E6 and E7 expression can drive the immortalization and oncogenic

transformation of human keratinocytes (1, 3).

One well-documented activity of many high-risk and low-risk (non-oncogenic) HPV E7

proteins is to bind and inactivate the retinoblastoma tumor suppressor (RB1) (4). RB1

inactivation drives the transcription of E2F target genes, thereby promoting the cell cycle

progression that is required for HPV DNA replication (5, 6). In addition to binding RB1, high-risk

HPV E7 proteins also target it for proteasome-mediated degradation (7). However, evidence

from keratinocyte immortalization assays, other cell-based assays of HPV E7 transforming

activity, and from studies of HPV E7 activity in transgenic mouse models supports that RB1

inactivation is necessary, but insufficient for the carcinogenic activity of high-risk HPV E7 (8-17).

We and others have determined that many HPV E7 proteins also target a second tumor

suppressor, Protein Tyrosine Phosphatase Non-receptor Type 14 (PTPN14) (18-20). Several

amino acids, including a highly conserved arginine in the HPV E7 C-terminus, enable E7 to bind

directly to PTPN14 (21, 22). A conserved acidic amino acid in the HPV E7 N-terminus binds to

the ubiquitin ligase UBR4, which is required for PTPN14 degradation and for the transforming

activity of high-risk HPV E7 (13, 18). PTPN14 degradation by high-risk HPV E7 inhibits

keratinocyte differentiation and extends primary epithelial cell lifespan, contributing to the

immortalization of keratinocytes in culture (18, 21).

PTPN14 is mutated or its expression is reduced in several cancer types, and data from

human cancer studies and cell-based assays provide evidence for the tumor suppressive

activity of PTPN14 (23-34). The phosphatase domain of PTPN14 likely lacks enzymatic function

and the putative catalytic site does not appear important for tumor suppressive activity (29, 30,

35). The best-characterized mechanism by which PTPN14 acts as a tumor suppressor is by

inactivating the YAP1 oncoprotein, a potent driver of cell stemness and self-renewal (29, 31-34,

36). YAP1 is a transcriptional coregulator that is active when localized to the nucleus and bound

to transcription factors including Transcriptional Enhanced Associate Domain (TEAD) proteins

(37).

YAP1 and its paralog TAZ are normally repressed by the Hippo signaling pathway and

Hippo pathway components are frequently mutated in human cancers (36, 38). The MST1/2 and

LATS1/2 kinases form the core of the conserved Hippo kinase cascade (39, 40). MST1/2

activate LATS kinases at sites including threonine 1079 on LATS1 (41). Active LATS1/2 then

phosphorylate YAP1 at serine 127 (42-45). LATS-dependent phosphorylation causes YAP1 to

interact with 14-3-3 proteins and be retained in the cytoplasm where it is transcriptionally

inactive and targeted for degradation (36, 38, 40, 45). Many additional protein-protein

interactions regulate Hippo kinase activity and influence the assembly of Hippo components at

the plasma membrane (36, 39). SAV1 promotes MST1 binding to LATS1 and MOB1A/B

promote LATS1 binding to YAP1 (41, 46-49). Other proteins including NF2 and WWC1/2/3 likely

participate in and influence Hippo protein interactions (50, 51). The NF2 protein localizes to the

plasma membrane in proximity to cytoskeletal scaffold elements and interacts with LATS1/2 (52-

54). NF2 recruits LATS to the plasma membrane and promotes LATS kinase activity and Hippo

signaling, but without affecting the MST kinase activity (51). WWC1 (KIBRA) and its paralogs

WWC2 and WWC3 have been proposed to bind and regulate LATS kinase activity, perhaps by

helping recruit LATS1/2 to the MST1/2-containing complex (55-57). Other data support that

WWC proteins interact with NF2 (58-61). Of the three WWC paralogs, WWC1 has been studied

most extensively (57). Many of the interactions between Hippo components are mediated by

binding of PPxY motifs in one protein to WW domains in another. Initial overexpression studies

suggested that PPxY motifs in PTPN14 bound to WW domains in YAP1 (29, 31, 32). However,

later biochemical experiments revealed that a separate set of PPxY motifs in PTPN14 make an

exceptionally high-affinity interaction with WW motifs in WWC1, suggesting that the primary

effect of PTPN14 does not occur through direct binding to YAP1 (62). The tumor suppressive

activity of WWC1 requires PTPN14 (63).

Several observations highlight the potential role of YAP1, PTPN14, and the Hippo

pathway in HPV carcinogenesis. We previously found that high-risk HPV E7 proteins promote

the nuclear localization of YAP1 in the basal layer of stratified epithelial cell cultures, dependent

on the ability of E7 to degrade PTPN14 (64). We also found that YAP1/TEAD-dependent

transcriptional activity was required for high-risk HPV E7 proteins to extend the lifespan of

primary human keratinocytes (64). Somatic mutation data in human cancers also argues for the

influence of an HPV oncoprotein on Hippo signaling and YAP1. The three mitogenic pathways

most differentially altered with high mutation rates in HPV-negative and low mutation rates in

HPV-positive head and neck squamous cell carcinoma (HNSCC) are p53 (targeted by E6), cell

cycle/RB1 (targeted by E7), and Hippo/YAP1 (64, 65).

Our findings support that RB1 inactivation and PTPN14 degradation are independent

activities of high-risk HPV E7 that are each required for E7 transforming activity. The primary

transcriptional readout in keratinocytes downstream of HPV E7-mediated PTPN14 degradation

and YAP1 activation is a repression of epithelial differentiation. However, it was not known

whether or how E7 and PTPN14 act on the Hippo pathway to regulate YAP1. Here we report

that HPV18 E7 degrades PTPN14 to inhibit Hippo signaling and activate YAP1, thereby

inhibiting the commitment to keratinocyte differentiation.

RESULTS

PTPN14 knockout reduces phosphorylation on YAP1 S127 and LATS1 T1079 in human

keratinocytes. HPVs infect human keratinocytes and PTPN14 knockout or HPV E7-mediated

PTPN14 degradation promotes YAP1 nuclear localization in the basal layer of stratified epithelia

(64). However, the role of PTPN14 in Hippo signaling has not been tested in human

keratinocytes. To test the hypothesis that PTPN14 degradation by E7 inhibits Hippo signaling,

we monitored Hippo pathway activity in the presence and absence of PTPN14. First, we directly

manipulated PTPN14. We transduced *hTERT* immortalized human keratinocytes (N/Tert-1) with

lentiviral vectors to deliver Cas9 and one of two different sgRNAs targeting PTPN14, or non-

targeting control sgRNAs. Disruption of the actin cytoskeleton activates Hippo kinases and we

treated cells with the actin polymerization inhibitor cytochalasin D to induce Hippo signaling.

YAP1 S127 phosphorylation (YAP1 pS127), which reflects decreased YAP1 activity, increased

relative to total YAP1 during 90 minutes of cytochalasin D treatment in each nontargeting control

cell line (Figure 1). In contrast, YAP1 pS127 did not increase over time in cytochalasin D-treated

PTPN14 knockout cell lines.

Having observed that YAP1 pS127 was inhibited in PTPN14 knockout keratinocytes, we

sought to determine whether PTPN14 knockout reduced phosphorylation on LATS, which is

well-established as a marker of LATS kinase activity. To complement the cytochalasin D

treatment experiments, we induced Hippo kinase activity by detachment of cells from their

growth substrate (66-68). Two independent sets of nontargeting control or PTPN14 knockout

N/Tert-1 cells were cultured at low density, then detached by trypsinization and incubated in

suspension for 10 minutes. YAP1 pS127 levels increased upon suspension in control cells, but

the increase in YAP1 pS127 was low or negligible in PTPN14 knockout cells (Figure 2A, B).

Similarly, phosphorylation on LATS1 T1079 (LATS1 pT1079) showed a greater increase upon

detachment in nontargeting control cells than in PTPN14 knockout cells. LATS2 protein could

not be detected by Western blot in N/Tert-1 cells.

NF2 binds to LATS kinases and tethers them to the plasma membrane (53, 54, 57, 69).

Phosphorylation on NF2 S518 is reduced when Hippo signaling is active (70, 71).

Dephosphorylated S518 indicates active NF2 and active Hippo signaling. Culturing cells in

suspension decreased NF2 pS518 levels in both control and PTPN14 knockout N/Tert-1 cells

(Figure 2A, B), indicating that PTPN14 knockout alters phosphorylation on LATS1 and YAP1,

but not NF2.

HPV18 E7-mediated PTPN14 degradation reduced YAP1 S127 and LATS1 T1079

phosphorylation in human keratinocytes. To test whether PTPN14 degradation by a high-risk

HPV E7 oncoprotein also reduces YAP1 and LATS1 phosphorylation, we used N/Tert-1 cells

that express HPV18 E7, HPV18 E7 R84S, or a vector control. HPV18 E7 R84S cannot bind or

degrade PTPN14 but can destabilize RB1 (21). Again, cytochalasin D treatment or culture in

suspension were used to stimulate Hippo activity. YAP1 pS127 increased relative to total YAP1

over time in cytochalasin D-treated vector control cells and in HPV18 E7 R84S cells, but not in

wild type HPV18 E7 cells (Figure 3A). Stimulating Hippo kinase activity using cell detachment

had a similar effect. YAP1 pS127 and LATS1 pT1079 increased after cell detachment in vector

control and HPV18 E7 R84S cells, but the detachment-dependent increase was smaller in the

wild type HPV18 E7 cells (Figure 3B). Overall, these data support that HPV18 E7 can dampen

Hippo-dependent phosphorylation on LATS1 T1079 and YAP1 S127, dependent on its ability to

bind and degrade PTPN14.

PTPN14 promotes differentiation and Hippo pathway activity. PTPN14 knockout reduces

the expression of epithelial differentiation genes and dampens the expression of differentiation

genes induced by cell detachment (18, 21). To test whether the effects of PTPN14 on

differentiation correlate with its effects on YAP1 phosphorylation, we first tested the effect of

PTPN14 knockout on differentiation in N/Tert-1 cells. Calcium treatment induces differentiation

gene expression in keratinocytes and we confirmed that transcript levels for the differentiation

marker genes Keratin 10 (*KRT10*) and involucrin (*IVL*) were reduced in calcium-treated PTPN14

knockout cells compared to matched controls (Figure 4).

The observation that PTPN14 limits the expression of differentiation genes prompted us

to test whether the converse was also true: that elevated PTPN14 levels could increase

differentiation gene expression. N/Tert-1 keratinocytes were transduced with a lentivirus

encoding HA-tagged doxycycline-inducible PTPN14 (pLIX-PTPN14). *IVL* and *KRT10* transcript

levels increased in doxycycline-treated N/Tert-pLIX-PTPN14 cells (Figure 5A), a finding

consistent with our previous report that PTPN14 overexpression increased the expression of

differentiation marker genes *KRT1* and *IVL* in HFK (64). Having established that increased

PTPN14 could promote differentiation, we tested whether it altered phosphorylation on Hippo

proteins. Doxycycline treatment induced PTPN14, increased YAP1 pS127 and LATS1 pT1079,

but did not affect NF2 pS518 (Figure 5B). Overall, PTPN14 overexpression increased

phosphorylation on the same sites and promoted differentiation gene expression, and

conversely PTPN14 knockout inhibited YAP1 pS127 and LATS1 pT1079 and reduced

differentiation gene expression.

The PTPN14 PY1/2 motifs are required to induce keratinocyte differentiation. Next we

used PTPN14 overexpression to determine which region(s) of PTPN14 upregulate

differentiation. Several protein components of the Hippo pathway interact with one another via

complementary WW domains and PPxY (PY) motifs (62, 72). PTPN14 contains two sets of PY

motifs and has a C-terminal protein tyrosine phosphatase domain. However, the amino acid

sequence of the PTPN14 phosphatase domain diverges from the consensus and it is likely to

have little or no catalytic activity (35). We designed two mutants, each lacking a PY motif pair,

and a third containing a serine point mutant C1121S at the conserved reactive cysteine of the

putative phosphatase active site (Figure 6A) (73, 74). We expressed wild-type (WT) and mutant

PTPN14 in primary human foreskin keratinocytes (HFK), then measured *KRT1* differentiation

marker transcript levels (Figure 6B). *KRT1* levels increased in WT PTPN14, PTPN14 ΔPY3/4,

and PTPN14 C1121S cells, but not in PTPN14 ΔPY1/2 cells. The PTPN14 PY1/2 motifs bind

with high affinity to the WW domain region of WWC1 (KIBRA) (62). We therefore tested if WT

and PY mutant forms of PTPN14 bound to WWC1 in human keratinocytes. N/Tert-Cas9 cells

were transfected with a PTPN14 sgRNA to eliminate endogenous PTPN14, then transduced

with pLIX vectors encoding WT or mutant HA-tagged PTPN14. PTPN14 and KRT10 protein

levels increased upon induction of WT PTPN14 (Figure S1). Cells were treated with doxycycline

and lysates subjected to anti-HA immunoprecipitation. Wild-type PTPN14 and PTPN14 ΔPY3/4,

but not PTPN14 ΔPY1/2, co-immunoprecipitated WWC1 (Figure 6C). We conclude that the

ability of PTPN14 to induce keratinocyte differentiation requires the PY motifs that bind to

WWC1, but not the PTPN14 C1121 residue.

LATS kinases are required for PTPN14 to induce keratinocyte differentiation. We

previously used siRNA knockdown combined with PTPN14 overexpression to demonstrate that

YAP1 is required for PTPN14 to promote differentiation gene expression (64). The finding was

based on the observation that although knockdown of YAP1 and its paralog TAZ increased

differentiation gene expression, PTPN14 overexpression could not further increase the levels of

differentiation genes in YAP1/TAZ-depleted cells. Our data further supported that the

predominant effect of PTPN14 on differentiation is mediated by YAP1, not TAZ. We also

observed that knockdown of either LATS1 or LATS2 in HFK reduced basal levels of

differentiation gene expression (64).

Here we used the same knockdown plus overexpression strategy to determine which of

several additional components of the Hippo pathway are required for PTPN14 to induce

keratinocyte differentiation. We transfected HFKs with siRNA targeting YAP1 and TAZ, LATS1

and LATS2, or a non-targeting control. Twenty-four hours post transfection, cells were

transduced with lentiviruses encoding PTPN14 or GFP, then total cellular RNA was harvested

72h post transfection. In siControl treated cells, PTPN14 expression increased the levels of

KRT10, *KRT1*, and *IVL* (Figure 7, Figure S2). Consistent with our previous finding, YAP1 and

TAZ knockdown in HFK increased the expression of differentiation genes, but PTPN14 could not

further increase differentiation in cells depleted of YAP1/TAZ. Although PTPN14 retained some

ability to increase *KRT10, KRT1, and IVL* RNA in cells treated with two different sets of siRNAs

targeting LATS1 and LATS2 (Figure 7A, Figure S2), levels of all three differentiation marker

genes remained low in LATS knockdown cells throughout the experiment. In the absence of

LATS kinases, PTPN14 did not increase the levels of *KRT10, KRT1, or IVL* beyond the amount

observed in untreated control cells. Knockdowns and PTPN14 overexpression were validated

by qRT-PCR (Figure S3).

Next we tested whether PTPN14 required MST1 and MST2, NF2, or WWC1 and its

paralogs WWC2 and WWC3 to induce differentiation. We transfected HFK with siRNA targeting

MST1 and MST2, NF2, or all three WWC genes, transduced the cells with lentiviruses encoding

GFP or PTPN14, and measured *KRT10* transcript levels. Depletion of MST1 and 2 did not

interfere with the ability of PTPN14 to induce *KRT10* (Figure 7B). Similar to the knockdown of

LATS1 and LATS2, knockdown of NF2 limited, although did not completely eliminate, the ability

of PTPN14 to induce *KRT10* (Figure 7C). Depletion of WWC genes had an intermediate effect,

but PTPN14 was still able to induce *KRT10* several-fold in the absence of WWC1/2/3 (Figure

7D). These data support that LATS kinases and NF2 are required for full induction of

differentiation by PTPN14.

PTPN14 knockout promotes anchorage independent growth and reduces YAP1 pS127 in

HEK TER cells. Finally, we tested the effect of PTPN14 knockout in a human cell

transformation assay. HEK TER cells are human embryonic kidney (HEK) cells that express the

catalytic subunit of human telomerase (*hTERT*), Simian Virus 40 (SV40) Large T antigen (LT),

and an oncogenic allele of *ras* (H-*ras* V12) (75, 76). The cells are immortalized but do not grow

without attachment to a substrate. HEK TER cells acquire the capacity for anchorage-

independent growth upon expression of SV40 Small T (ST) antigen. ST activates YAP1 and

transforms HEK TER cells by interacting with a PP2A subunit of the STRIPAK complex

upstream of the Hippo pathway (77). We previously reported that HPV16 E7 could promote the

anchorage-independent growth of HEK TER cells (11). The HPV16 E7 transforming activity in

HEK TER cells mapped to the regions required to degrade PTPN14 (13, 19).

To test whether PTPN14 knockout promotes the anchorage-independent growth of HEK

TER cells, we generated CRISPR/Cas9 edited HEK TER cells that do not express PTPN14.

LATS1/2 knockout HEK TER cells were included to verify that Hippo pathway inhibition

promoted HEK TER growth and HEK TER cells stably expressing SV40 ST (11) were included

as a positive control. HEK TER cells depleted of PTPN14, LATS1/2, or that expressed SV40 ST

readily formed colonies in soft agar assays (Figure 8A), whereas HEK TER cells transfected

with non-targeting sgRNA exhibited minimal anchorage-independent growth. Consistent with our

findings in human keratinocytes and with the previous report that SV40 ST reduces YAP1

phosphorylation (77), cytochalasin D treatment induced YAP1 pS127 more strongly in HEK TER

sgNT cells than in cells depleted of PTPN14, LATS1 and LATS2, or that expressed SV40 ST

(Figure 8B). Using a second set of sgRNAs directed against PTPN14 and LATS1 and LATS2

resulted in a similar trend, albeit with less efficient reduction in LATS1 protein levels (Figure S4).

DISCUSSION

YAP1 is a transcriptional coactivator that promotes epithelial cell stemness and self-

renewal, limiting differentiation (42, 50, 78-85). We previously reported that oncogenic high-risk

HPV E7 proteins bind and target PTPN14 for proteasome-mediated degradation (19, 86). The

HPV18 E7-mediated degradation of PTPN14 results in increased nuclear YAP1 localization and

the maintenance of basal epithelial identity in three-dimensional models of stratified epithelial

tissue (64). However, we had not tested whether HPV E7-mediated degradation of PTPN14

alters Hippo pathway activity, which controls YAP1, in keratinocytes.

Several reports indicate that PTPN14 can inhibit YAP1 activity, although the mechanistic

basis of such inhibition is incompletely understood (24, 29-34, 55, 64). Initial studies suggested

that PTPN14 bound directly to YAP1 via the second set of PTPN14 PY motifs (PY3/4 in this

paper), but more recent analysis of PY-WW interactions among Hippo components determined

that the PTPN14 PY1/2 tandem motif has nanomolar affinity for WW domains in Hippo

components including WWC1 (31-33, 62). These and other data support that PTPN14, WWC1,

and LATS1/2 work together to inhibit YAP1 (24, 55, 56), but not by binding YAP1 directly.

Certain assays of tumor suppression show a mutual requirement for PTPN14 and WWC1

(KIBRA) (55, 62, 63, 87, 88), but potential contributions of WWC2 and WWC3 in the same

assays have not been thoroughly tested. Other data indicate that PTPN14 and/or WWC proteins

act through NF2, which interacts with LATS1/2 and recruits LATS1/2 to a plasma membrane-

associated complex that also contains MST1/2 and its cofactor SAV1 (51, 60, 61, 70, 89-91).

Our data strengthen the model that PTPN14 and WWC proteins act together to regulate

LATS kinases and that NF2 is required for the complex to assemble, likely at the plasma

membrane. In support of this model, we found that either PTPN14 knockout or HPV18 E7-

mediated PTPN14 degradation reduced the levels of LATS1 pT1079 and YAP1 pS127,

indicating that PTPN14 degradation reduces Hippo activity and activates YAP1 (Figure 1, Figure

2, Figure 3). The effect of HPV18 E7 on YAP1 was independent of its ability to degrade RB1

(Figure 3). Conversely, PTPN14 overexpression increased phosphorylation on LATS1 T1079

and YAP1 S127 (Figure 5).

We established the connection between PTPN14, YAP1 activity, and keratinocyte

differentiation (Figure 4, Figure 5, (18)) and used keratinocyte differentiation as a readout to

probe the effect of PTPN14 on Hippo signaling. We found that the induction of differentiation by

PTPN14 required its PY1/2 motifs, which bind WWC1, but not the catalytic cysteine of the

putative active site (Figure 6). PTPN14 required LATS1/2 kinases and NF2 to promote

differentiation (Figure 7), further supporting that PTPN14 acts on LATS kinases only in the

presence of NF2. Our finding that PTPN14 could still induce differentiation in cells transfected

with MST1/2 siRNAs is consistent with the finding that MST1/2 are not required for PTPN14

overexpression to induce LATS1 phosphorylation (55). It is possible that one of several other

kinases can phosphorylate LATS in the absence of MST1/2 (59, 92-96). Finally, PTPN14 could

still promote differentiation in cells treated with WWC1/2/3 siRNA, even though the WWC1

binding motif was required for PTPN14 to promote differentiation in a separate assay (Figure

6B, Figure 7). Incomplete siRNA-based depletion of WWC1/2/3 may have resulted in a weaker

phenotype than that observed with the PTPN14 ΔPY1/2 deletion mutant. Finally, our finding that

in anchorage independent growth assays PTPN14 knockout promoted colony formation and

reduced the levels of phosphorylation on YAP1 S127 (Figure 8) further strengthens the

connection between YAP1, PTPN14, and oncogenic transformation.

Consistent with the central role of YAP1 in epithelial homeostasis and the epithelial tropic

nature of HPV infection, there are additional links between HPV biology and YAP1/Hippo

signaling. Cutaneous HPV8 E6 can bind TEAD and activate YAP1/TEAD-dependent

transcription (97), and HPV8 E6 can also reduce LATS phosphorylation during failed cytokinesis

(98). HPV E6/E7 expression in cancer cell lines downregulates MST1 by increasing miR-18A

levels (99). Both high- and low-risk mucosal HPV E6 proteins have some ability to promote

YAP1 nuclear localization in basal epithelial cells (64, 100). There is growing recognition that

YAP1 and Hippo signaling are affected by other DNA tumor viruses. SV40 ST antigen promotes

the interaction between STRIPAK and MAP4K4, thereby decreasing MAP4K4 activity and

activating YAP1 (77). The Epstein-Barr Virus (EBV) protein LMP1 reduces phosphorylation on

YAP1 S127, and the analogous TAZ S89 site, by increasing Src kinase activity (101).

Our findings provide the first evidence of a viral protein, HPV18 E7, targeting a Hippo

pathway component to suppress LATS kinase activity, the central regulatory step immediately

upstream of YAP1. PTPN14 is expressed specifically in the basal layer of stratified epithelia (64,

102), and so our finding that PTPN14 can control YAP1 and LATS phosphorylation in human

keratinocytes supports that PTPN14 is a basal keratinocyte-specific regulator of YAP1 and the

Hippo pathway. Our previous work supports that PTPN14 degradation and YAP1/TEAD

transcriptional activity are required for the transforming activity of high-risk HPV E7 (21, 64).

Other direct interactions between HPV oncoproteins and cellular targets, such as E7/RB1 and

E6/p53, are essential for HPV carcinogenesis but so far have not been tractable targets for

therapeutic intervention. Understanding the mechanism by which E7 degrades PTPN14 to

activate YAP1 has the potential both to provide a new therapeutic opportunity in HPV disease

and to elucidate details of how HPV E7 proteins promote basal epithelial identity. Our future

studies will also use E7 and PTPN14 as tools to better understand the complex interactions that

control Hippo signaling and YAP1 activity in human epithelial cells.

MATERIALS AND METHODS

Plasmids and cloning

MSCV-puro C-terminal FlagHA retroviral expression vectors encoding HPV18 E7, HPV18 E7

R84S, or empty vector controls were previously described (21, 86). Sequences for sgRNA

selected from the Broad Institute Brunello library (103) were ligated into LentiCRISPRv2 vectors

following standard cloning protocols. A lentiviral vector (pHAGE) was used for PTPN14

overexpression in HFK as previously described (64). A doxycycline-inducible, HA-tagged

PTPN14 overexpression vector was developed by cloning the wild-type PTPN14 coding

sequence into a lentiviral vector pLIX_402, a gift from David Root (Addgene #41394), using

Gateway recombination. More information for all plasmids is listed in Table S1. Mutant versions

of PHAGE-PTPN14 and pLIX-PTPN14 were generated by site-directed mutagenesis and

Gateway recombination.

Cell culture

Discarded, deidentified primary human foreskin keratinocytes (HFKs) were obtained from the

Skin Biology and Diseases Resource-based Center (SBDRC) at the University of Pennsylvania.

N/Tert-1 keratinocytes were the gift of James Rheinwald (104). All keratinocytes were cultured

as previously described (64). N/Tert-1 cell lines were established by previously described

methods for lentiviral or retroviral vector expression (18, 86). Gene knockouts in N/Tert-1 cells

were accomplished by one of two methods. In Figures 1 and 2, N/Tert-1 LentiCRISPR knockout

cell lines were established by lentiviral transduction of LentiCRISPRv2 constructs encoding

spCas9 and a sgRNA sequence selected from the Broad Institute Brunello Library for targeting

PTPN14 or nontargeting, then selecting with G418. In Figures 4 and 6C, N/Tert-1 Cas9

knockout cells were generated by transduction with a spCas9 lentiviral expression vector

(Addgene # 52962) and selecting with blasticidin, after which cells were transiently transfected

with sgRNA targeting PTPN14, a non-targeting sgRNA (Synthego), or mock transfected (18).

Doxycycline inducible PTPN14 expressing N/Tert-1 keratinocytes (Figure 5) were generated by

transducing N/Tert-1 cells with the lentiviral expression vector pLIX-PTPN14 and selecting with

puroymcin. In Figure 5, the inducible PTPN14 cells were established using parental N/Tert-1

cells. In Figure 6C, the inducible PTPN14 cells were established using N/Tert-1

Cas9:sgPTPN14 knockout cells as the starting material. N/Tert-1 Cas9:sgPTPN14 knockout

cells were transduced with pLIX-PTPN14 lentiviral constructs containing the wild-type PTPN14

gene or PTPN14 ΔPY1/2, ΔPY3/4 or C1121S mutants and selected with puromycin.

Keratinocytes expressing HPV18 E7 were generated by transducing N/Tert-1 cells with MSCV-

Puro retroviral vectors encoding either wild-type HPV18 E7, HPV18 E7 R84S, or empty control,

then selecting with puromycin (Figure 3).

HEK TER cells were described previously and cultured in Minimum Essential Media (MEM,

Gibco) supplemented with 10% fetal bovine serum (FBS), 2mM L-glutamine and penicillin-

streptomycin (76). HEK TER cells expressing spCas9 were generated by transduction with a

Cas9 lentiviral vector followed by blasticidin selection. HEK TER Cas9 cells were transfected

with sgRNA targeting PTPN14, LATS1, LATS2, or nontargeting control (Synthego) (18). HEK

TER cells stably expressing SV40 ST were described previously (11, 76).

Immunoblots

Immunoblots were performed using Mini-PROTEAN or Criterion (Bio-Rad) Tris/Glycine 4-20%

polyacrylamide gradient SDS-PAGE gels. Proteins were transferred to polyvinylidene difluoride

(PVDF) membranes as previously described (19). Membranes were blocked in 5% nonfat dried

milk (NFDM) in Tris-buffered saline, pH 7.4, with 0.05% Tween 20 (TBS-T) unless otherwise

specified. Blots were then incubated in 5% NFDM/TBS-T with primary antibodies listed in Table

S2. Blots were incubated with anti-mouse or anti-rabbit secondary antibody conjugated to

horseradish peroxidase, washed with TBS-T, and proteins detected by chemiluminescent

substrate. Blots were imaged using a Chemidoc MP imaging system (BioRad). Membranes

probed for YAP1 or with one of several phospho-specific antibodies (YAP1 pS127, LATS1

pT1079, NF2 pS518) were blocked with TBS-based Intercept Blocking Buffer (LI-COR) and

incubated with antibodies diluted in TBS-based Intercept Antibody Diluent (LI-COR). Blots

probed for LATS1 pT1079, YAP1 pS127, YAP1 and beta-actin were incubated with secondary

antibody-fluorophore conjugates anti-mouse IRDye 680LT (LI-COR) or anti-rabbit IRDye 800CW

(LI-COR) diluted in Intercept Antibody Diluent (LI-COR). Blots were washed with TBS-T and

fluorescence was detected using a LI-COR imager. Blots probed for NF2 pS518 were incubated

with anti-rabbit secondary antibody conjugated to horseradish peroxidase, and diluted in

Intercept Antibody Diluent (LI-COR), then washed with TBS-T and detected by

chemiluminescent substrate using a Chemidoc MP imaging system (BioRad).

Immunoprecipitation

Immunoprecipitation with anti-HA conjugated agarose beads was described previously (19).

Whole cell protein lysates were incubated with anti-HA agarose beads, then washed with lysis

buffer. Immunoprecipitates were eluted by boiling in protein sample buffer. Elution and input

samples were analyzed by SDS-PAGE and immunoblot.

Hippo activation assays

For Cytochalasin D treatment, 2.5×10^5 cells were plated in 6 cm dishes and grown overnight

before treatment with 5 µM Cytochalasin D in appropriate media (MEM for HEK TER cells, K-

SFM for keratinocytes). Plates were harvested every 30 minutes starting at T_0 by washing once

with TBS and scraping into Radioimmunoprecipitation Assay (RIPA) lysis buffer containing

protease and phosphatase inhibitors. For suspension assays, 5×10^5 cells were plated in 10 cm

dishes and grown for 24 hours before treatment with 0.05% Trypsin/EDTA for 10 minutes, then

quenched with DMEM/F12 containing 10% FBS. Suspended cells were pelleted and washed

with TBS before being pelleted again and resuspended in RIPA lysis buffer containing protease

and phosphatase inhibitors. Control cells were washed once with TBS in plates and scraped

into RIPA with protease and phosphatase inhibitors. All cell lysates were incubated on ice at

least 30 minutes to complete lysis then protein concentrations were normalized by Bradford

assay.

Differentiation induction with calcium

N/Tert-1 mock or sgPTPN14 cells were seeded in 6 well plates at a density of 10,000 cells per

well and cultured for four days. Cells were then maintained in K-SFM (untreated) or switched to

culture in Keratinocyte Basal Medium (KBM, Lonza) with $CaCl_2$ added to a final concentration of

1.5 mM. Treated and untreated cells were harvested for RNA analysis and qRT-PCR at times

indicated.

Differentiation induction with PTPN14 mutants

HFK were seeded in 12 well plates with 5.0×10^4 cells per well and transduced with pHAGE

lentiviral constructs encoding GFP, wild-type PTPN14, PTPN14 ΔPY1/2, PTPN14 ΔPY3/4 or

PTPN14 C1121S. Total RNA was collected 48 hours post-transduction and analyzed by qRT-

PCR for KRT1 transcripts normalized to GAPDH.

qRT-PCR

Total RNA from cell lysates was isolated with a NucleoSpin RNA purification kit (Macherey-

Nagel). Purified whole cell RNA was converted to cDNA using the high-capacity cDNA reverse

transcription kit (Applied Biosystems). Specific genes were quantified by qRT-PCR with Fast

SYBR green master mix (Applied Biosystems) using a Quant-Studio 3 system (ThermoFisher

Scientific). KiCqStart qRT-PCR primers (MilliporeSigma) were used to quantify the following

transcripts: KRT1, KRT10, IVL, PTPN14, YAP1, LATS1, STK3 (MST2), STK4 (MST1), NF2,

WWC1, WWC2, WWC3, and GAPDH.

PTPN14 overexpression and siRNA knockdown assay

Primary HFK were reverse transfected with 40nM siRNAs (Dharmacon) using Dharmafect 1

transfection reagent. Transfected HFK were transduced at 24 hours post-transfection with

pHAGE-PTPN14 lentivirus. At 48 hours post-transfection media was exchanged for 1:1 DF-K/K-

SFM (86). Cells were lysed and total RNA collected 72 hours post-transfection for qRT-PCR

analysis. The siRNAs used were: nontargeting siControl #1, siYAP1-06, siWWTR-06, siLATS1-

05, siLATS1-08, siLATS2-09, siLATS2-10, siSTK3-06, siSTK3-08, siSTK4-06, siSTK4-08,

siNF2-05, siNF2-07, siWWC1-09, siWWC1-10, siWWC2-17, siWWC2-20, siWWC3-17,

siWWC3-18.

Anchorage independent growth assay

HEK TER cell lines were suspended in Dulbecco's Modified Eagle Medium (DMEM, Gibco) and

mixed with 0.4% Noble agar, then seeded onto 6 cm plates coated with DMEM/0.6% Noble

agar, allowed to solidify at room temperature, then incubated at $37^{\circ}C/5\%$ CO_2 (11). DMEM was

supplemented with 10% FBS and antibiotic-antimycotic. Cells were seeded in triplicate plates

(5.0×10^4 cells/plate) for each condition and colonies allowed to form for 18 days, then

photographed with a GelDoc XR+ imager (BioRad). Colonies were counted using ImageJ

software.

ACKNOWLEDGEMENTS

We are grateful to Drs. Karl Munger and Molly Patterson for helpful comments on the

manuscript and to Dr. Pushkal Ramesh for help with primer validation. We thank the members

of our laboratory for helpful discussions. This work was supported by American Cancer Society

grant 131661-RSG-18-048-01-MPC, NIH/NIAID R01 AI148431, and NIH/NIAID R21 AI176035

to EAW. WJB was supported by NIH/NCI T32 CA115299 and NIH/NIDCR F32 DE032573. JH

was supported by NIH/NIAID T32 AI007324 and NIH/NIDCR F31 DE030365. The University of

Pennsylvania SBDRC was funded by NIH/NIAMS P30 AR069589.

AUTHOR CONTRIBUTIONS

Conception and design: WJB, JH, EAW. Acquisition of data: WJB, JH, EAW. Analysis and

interpretation of data: WJB, JH, EAW. Drafting or revising the article: WJB, JH, EAW.

REFERENCES

1. A. A. McBride, Human papillomaviruses: diversity, infection and host interactions. *Nat Rev Microbiol* **20**, 95-108 (2022).
2. C. de Martel, M. Plummer, J. Vignat, S. Franceschi, Worldwide burden of cancer attributable to HPV by site, country and HPV type. *Int J Cancer* **141**, 664-670 (2017).
3. C. A. Moody, L. A. Laimins, Human papillomavirus oncoproteins: pathways to transformation. *Nat Rev Cancer* **10**, 550-560 (2010).
4. N. Dyson, P. M. Howley, K. Munger, E. Harlow, The human papilloma virus-16 E7 oncoprotein is able to bind to the retinoblastoma gene product. *Science* **243**, 934-937 (1989).
5. J. DeGregori, D. G. Johnson, Distinct and Overlapping Roles for E2F Family Members in Transcription, Proliferation and Apoptosis. *Curr Mol Med* **6**, 739-748 (2006).
6. N. Dyson, The regulation of E2F by pRB-family proteins. *Genes Dev* **12**, 2245-2262 (1998).
7. S. N. Boyer, D. E. Wazer, V. Band, E7 protein of human papilloma virus-16 induces degradation of retinoblastoma protein through the ubiquitin-proteasome pathway. *Cancer Res* **56**, 4620-4624 (1996).
8. F. Ciccolini, G. Di Pasquale, F. Carlotti, L. Crawford, M. Tommasino, Functional studies of E7 proteins from different HPV types. *Oncogene* **9**, 2633-2638 (1994).
9. T. Ibaraki, M. Satake, N. Kurai, M. Ichijo, Y. Ito, Transacting activities of the E7 genes of several types of human papillomavirus. *Virus Genes* **7**, 187-196 (1993).
10. R. J. Jewers, P. Hildebrandt, J. W. Ludlow, B. Kell, D. J. McCance, Regions of human papillomavirus type 16 E7 oncoprotein required for immortalization of human keratinocytes. *J Virol* **66**, 1329-1335 (1992).
11. E. A. White *et al.*, Papillomavirus E7 oncoproteins share functions with polyomavirus small T antigens. *J Virol* **89**, 2857-2865 (2015).
12. L. Banks, C. Edmonds, K. H. Vousden, Ability of the HPV16 E7 protein to bind RB and induce DNA synthesis is not sufficient for efficient transforming activity in NIH3T3 cells. *Oncogene* **5**, 1383-1389 (1990).
13. K. W. Huh *et al.*, Association of the human papillomavirus type 16 E7 oncoprotein with the 600-kDa retinoblastoma protein-associated factor, p600. *Proc Natl Acad Sci U S A* **102**, 11492-11497 (2005).
14. A. M. Helt, D. A. Galloway, Destabilization of the retinoblastoma tumor suppressor by human papillomavirus type 16 E7 is not sufficient to overcome cell cycle arrest in human keratinocytes. *J Virol* **75**, 6737-6747 (2001).
15. K. Strati, P. F. Lambert, Role of Rb-dependent and Rb-independent functions of papillomavirus E7 oncogene in head and neck cancer. *Cancer Res* **67**, 11585-11593 (2007).
16. S. Balsitis, F. Dick, N. Dyson, P. F. Lambert, Critical roles for non-pRb targets of human papillomavirus type 16 E7 in cervical carcinogenesis. *Cancer Res* **66**, 9393-9400 (2006).

526   17.   S. Balsitis *et al.*, Examination of the pRb-dependent and pRb-independent functions of
527   E7 in vivo. *J Virol* **79**, 11392-11402 (2005).
528   18.   J. Hatterschide *et al.*, PTPN14 degradation by high-risk human papillomavirus E7 limits
529   keratinocyte differentiation and contributes to HPV-mediated oncogenesis. *Proc Natl*
530   *Acad Sci U S A* **116**, 7033-7042 (2019).
531   19.   E. A. White, K. Munger, P. M. Howley, High-Risk Human Papillomavirus E7 Proteins
532   Target PTPN14 for Degradation. *mBio* **7** (2016).
533   20.   A. Szalmas *et al.*, The PTPN14 Tumor Suppressor Is a Degradation Target of Human
534   Papillomavirus E7. *J Virol* **91** (2017).
535   21.   J. Hatterschide, A. C. Brantly, M. Grace, K. Munger, E. A. White, A Conserved Amino
536   Acid in the C Terminus of Human Papillomavirus E7 Mediates Binding to PTPN14 and
537   Repression of Epithelial Differentiation. *J Virol* **94** (2020).
538   22.   H. Y. Yun *et al.*, Structural basis for recognition of the tumor suppressor protein PTPN14
539   by the oncoprotein E7 of human papillomavirus. *PLoS Biol* **17**, e3000367 (2019).
540   23.   Z. Wang *et al.*, Mutational analysis of the tyrosine phosphatome in colorectal cancers.
541   *Science* **304**, 1164-1166 (2004).
542   24.   R. Wang, Y. Du, J. Shang, X. Dang, G. Niu, PTPN14 acts as a candidate tumor
543   suppressor in prostate cancer and inhibits cell proliferation and invasion through
544   modulating LATS1/YAP signaling. *Mol Cell Probes* **53**, 101642 (2020).
545   25.   R. Mehra *et al.*, Biallelic Alteration and Dysregulation of the Hippo Pathway in Mucinous
546   Tubular and Spindle Cell Carcinoma of the Kidney. *Cancer Discov* **6**, 1258-1266 (2016).
547   26.   X. Bonilla *et al.*, Genomic analysis identifies new drivers and progression pathways in
548   skin basal cell carcinoma. *Nat Genet* **48**, 398-406 (2016).
549   27.   A. Schramm *et al.*, Mutational dynamics between primary and relapse neuroblastomas.
550   *Nat Genet* **47**, 872-877 (2015).
551   28.   T. Olafsdottir *et al.*, Loss-of-Function Variants in the Tumor-Suppressor Gene PTPN14
552   Confer Increased Cancer Risk. *Cancer Res* **81**, 1954-1964 (2021).
553   29.   W. Wang *et al.*, PTPN14 is required for the density-dependent control of YAP1. *Genes*
554   *Dev* **26**, 1959-1971 (2012).
555   30.   S. S. Mello *et al.*, A p53 Super-tumor Suppressor Reveals a Tumor Suppressive p53-
556   Ptpn14-Yap Axis in Pancreatic Cancer. *Cancer Cell* **32**, 460-473 e466 (2017).
557   31.   X. Liu *et al.*, PTPN14 interacts with and negatively regulates the oncogenic function of
558   YAP. *Oncogene* **32**, 1266-1273 (2013).
559   32.   J. M. Huang *et al.*, YAP modifies cancer cell sensitivity to EGFR and survivin inhibitors
560   and is negatively regulated by the non-receptor type protein tyrosine phosphatase 14.
561   *Oncogene* **32**, 2220-2229 (2013).
562   33.   C. Michaloglou *et al.*, The tyrosine phosphatase PTPN14 is a negative regulator of YAP
563   activity. *PLoS One* **8**, e61916 (2013).
564   34.   X. Han *et al.*, Nonreceptor tyrosine phosphatase 14 promotes proliferation and migration
565   through regulating phosphorylation of YAP of Hippo signaling pathway in gastric cancer
566   cells. *J Cell Biochem* **120**, 17723-17730 (2019).
567   35.   V. Reiterer *et al.*, The dead phosphatases society: a review of the emerging roles of
568   pseudophosphatases. *FEBS J* **287**, 4198-4220 (2020).
569   36.   K. F. Harvey, X. Zhang, D. M. Thomas, The Hippo pathway and human cancer. *Nat Rev*
570   *Cancer* **13**, 246-257 (2013).
571   37.   B. Zhao *et al.*, TEAD mediates YAP-dependent gene induction and growth control.
572   *Genes Dev* **22**, 1962-1971 (2008).
573   38.   P. C. Calses, J. J. Crawford, J. R. Lill, A. Dey, Hippo Pathway in Cancer: Aberrant
574   Regulation and Therapeutic Opportunities. *Trends Cancer* **5**, 297-307 (2019).
575   39.   S. Ma, Z. Meng, R. Chen, K. L. Guan, The Hippo Pathway: Biology and Pathophysiology.
576   *Annu Rev Biochem* **88**, 577-604 (2019).

40. B. A. Edgar, From cell structure to transcription: Hippo forges a new path. *Cell* **124**, 267-
273 (2006).
41. E. H. Chan *et al.*, The Ste20-like kinase Mst2 activates the human large tumor
suppressor kinase Lats1. *Oncogene* **24**, 2076-2086 (2005).
42. B. Zhao *et al.*, Inactivation of YAP oncoprotein by the Hippo pathway is involved in cell
contact inhibition and tissue growth control. *Genes Dev* **21**, 2747-2761 (2007).
43. T. Oka, V. Mazack, M. Sudol, Mst2 and Lats kinases regulate apoptotic function of Yes
kinase-associated protein (YAP). *J Biol Chem* **283**, 27534-27546 (2008).
44. Y. Hao, A. Chun, K. Cheung, B. Rashidi, X. Yang, Tumor suppressor LATS1 is a negative
regulator of oncogene YAP. *J Biol Chem* **283**, 5496-5509 (2008).
45. B. Zhao, L. Li, K. Tumaneng, C. Y. Wang, K. L. Guan, A coordinated phosphorylation by
Lats and CK1 regulates YAP stability through SCF(beta-TRCP). *Genes Dev* **24**, 72-85
(2010).
46. M. Praskova, A. Khoklatchev, S. Ortiz-Vega, J. Avruch, Regulation of the MST1 kinase
by autophosphorylation, by the growth inhibitory proteins, RASSF1 and NORE1, and by
Ras. *Biochem J* **381**, 453-462 (2004).
47. B. A. Callus, A. M. Verhagen, D. L. Vaux, Association of mammalian sterile twenty
kinases, Mst1 and Mst2, with hSalvador via C-terminal coiled-coil domains, leads to its
stabilization and phosphorylation. *FEBS J* **273**, 4264-4276 (2006).
48. Z. Lin, R. Xie, K. Guan, M. Zhang, A WW Tandem-Mediated Dimerization Mode of SAV1
Essential for Hippo Signaling. *Cell Rep* **32**, 108118 (2020).
49. J. Bothos, R. L. Tuttle, M. Ottey, F. C. Luca, T. D. Halazonetis, Human LATS1 is a mitotic
exit network kinase. *Cancer Res* **65**, 6568-6575 (2005).
50. Z. Meng, T. Moroishi, K. L. Guan, Mechanisms of Hippo pathway regulation. *Genes Dev*
30, 1-17 (2016).
51. F. Yin *et al.*, Spatial organization of Hippo signaling at the plasma membrane mediated
by the tumor suppressor Merlin/NF2. *Cell* **154**, 1342-1355 (2013).
52. H. Han *et al.*, Regulation of the Hippo Pathway by Phosphatidic Acid-Mediated Lipid-
Protein Interaction. *Mol Cell* **72**, 328-340 e328 (2018).
53. A. I. McClatchey, M. Giovannini, Membrane organization and tumorigenesis--the NF2
tumor suppressor, Merlin. *Genes Dev* **19**, 2265-2277 (2005).
54. D. Lallemand, M. Curto, I. Saotome, M. Giovannini, A. I. McClatchey, NF2 deficiency
promotes tumorigenesis and metastasis by destabilizing adherens junctions. *Genes Dev*
17, 1090-1100 (2003).
55. K. E. Wilson *et al.*, PTPN14 forms a complex with Kibra and LATS1 proteins and
negatively regulates the YAP oncogenic function. *J Biol Chem* **289**, 23693-23700 (2014).
56. F. Uliana *et al.*, Phosphorylation-linked complex profiling identifies assemblies required
for Hippo signal integration. *Mol Syst Biol* **19**, e11024 (2023).
57. D. O. Wennmann *et al.*, Evolutionary and molecular facts link the WWC protein family to
Hippo signaling. *Mol Biol Evol* **31**, 1710-1723 (2014).
58. S. Qi *et al.*, WWC proteins mediate LATS1/2 activation by Hippo kinases and imply a
tumor suppression strategy. *Mol Cell* **82**, 1850-1864 e1857 (2022).
59. R. Chen, R. Xie, Z. Meng, S. Ma, K. L. Guan, STRIPAK integrates upstream signals to
initiate the Hippo kinase cascade. *Nat Cell Biol* **21**, 1565-1577 (2019).
60. I. Poernbacher, R. Baumgartner, S. K. Marada, K. Edwards, H. Stocker, Drosophila Pez
acts in Hippo signaling to restrict intestinal stem cell proliferation. *Curr Biol* **22**, 389-396
(2012).
61. A. Elbediwy *et al.*, Integrin signalling regulates YAP and TAZ to control skin homeostasis.
Development **143**, 1674-1687 (2016).
62. Z. Lin *et al.*, Decoding WW domain tandem-mediated target recognitions in tissue growth
and cell polarity. *Elife* **8** (2019).

63.   J. F. Knight *et al.*, KIBRA (WWC1) Is a Metastasis Suppressor Gene Affected by
Chromosome 5q Loss in Triple-Negative Breast Cancer. *Cell Rep* **22**, 3191-3205 (2018).
64.   J. Hatterschide *et al.*, YAP1 activation by human papillomavirus E7 promotes basal cell
identity in squamous epithelia. *Elife* **11** (2022).
65.   F. Sanchez-Vega *et al.*, Oncogenic Signaling Pathways in The Cancer Genome Atlas.
Cell **173**, 321-337 e310 (2018).
66.   B. Zhao *et al.*, Cell detachment activates the Hippo pathway via cytoskeleton
reorganization to induce anoikis. *Genes Dev* **26**, 54-68 (2012).
67.   Y. Si *et al.*, Src Inhibits the Hippo Tumor Suppressor Pathway through Tyrosine
Phosphorylation of Lats1. *Cancer Res* **77**, 4868-4880 (2017).
68.   A. Barzegari *et al.*, The role of Hippo signaling pathway and mechanotransduction in
tuning embryoid body formation and differentiation. *J Cell Physiol* **235**, 5072-5083
(2020).
69.   R. F. Hennigan, J. S. Fletcher, S. Guard, N. Ratner, Proximity biotinylation identifies a set
of conformation-specific interactions between Merlin and cell junction proteins. *Sci
Signal* **12** (2019).
70.   M. C. Primi, E. S. Rangarajan, D. N. Patil, T. Izard, Conformational flexibility determines
the Nf2/merlin tumor suppressor functions. *Matrix Biol Plus* **12**, 100074 (2021).
71.   E. I. Surace, C. A. Haipek, D. H. Gutmann, Effect of merlin phosphorylation on
neurofibromatosis 2 (NF2) gene function. *Oncogene* **23**, 580-587 (2004).
72.   Z. Salah, A. Alian, R. I. Aqeilan, WW domain-containing proteins: retrospectives and the
future. *Front Biosci (Landmark Ed)* **17**, 331-348 (2012).
73.   A. J. Barr, J. E. Debreczeni, J. Eswaran, S. Knapp, Crystal structure of human protein
tyrosine phosphatase 14 (PTPN14) at 1.65-A resolution. *Proteins* **63**, 1132-1136 (2006).
74.   K. E. Chen *et al.*, Substrate specificity and plasticity of FERM-containing protein tyrosine
phosphatases. *Structure* **23**, 653-664 (2015).
75.   W. C. Hahn *et al.*, Creation of human tumour cells with defined genetic elements. *Nature*
400, 464-468 (1999).
76.   W. C. Hahn *et al.*, Enumeration of the simian virus 40 early region elements necessary
for human cell transformation. *Mol Cell Biol* **22**, 2111-2123 (2002).
77.   J. W. Kim *et al.*, STRIPAK directs PP2A activity toward MAP4K4 to promote oncogenic
transformation of human cells. *Elife* **9** (2020).
78.   J. Huang, S. Wu, J. Barrera, K. Matthews, D. Pan, The Hippo signaling pathway
coordinately regulates cell proliferation and apoptosis by inactivating Yorkie, the
Drosophila Homolog of YAP. *Cell* **122**, 421-434 (2005).
79.   A. Pocaterra, P. Romani, S. Dupont, YAP/TAZ functions and their regulation at a glance.
J Cell Sci **133** (2020).
80.   A. Beverdam *et al.*, Yap controls stem/progenitor cell proliferation in the mouse postnatal
epidermis. *J Invest Dermatol* **133**, 1497-1505 (2013).
81.   B. C. Heng *et al.*, Role of YAP/TAZ in Cell Lineage Fate Determination and Related
Signaling Pathways. *Front Cell Dev Biol* **8**, 735 (2020).
82.   A. D. Szymaniak, J. E. Mahoney, W. V. Cardoso, X. Varelas, Crumbs3-Mediated Polarity
Directs Airway Epithelial Cell Fate through the Hippo Pathway Effector Yap. *Dev Cell* **34**,
283-296 (2015).
83.   J. Hicks-Berthet *et al.*, Yap/Taz inhibit goblet cell fate to maintain lung epithelial
homeostasis. *Cell Rep* **36**, 109347 (2021).
84.   D. Yimlamai *et al.*, Hippo pathway activity influences liver cell fate. *Cell* **157**, 1324-1338
(2014).
85.   R. Zhao *et al.*, Yap tunes airway epithelial size and architecture by regulating the identity,
maintenance, and self-renewal of stem cells. *Dev Cell* **30**, 151-165 (2014).

86. E. A. White *et al.*, Systematic identification of interactions between host cell proteins and
E7 oncoproteins from diverse human papillomaviruses. *Proc Natl Acad Sci U S A* **109**,
E260-267 (2012).
87. K. E. Wilson, N. Yang, A. L. Mussell, J. Zhang, The Regulatory Role of KIBRA and
PTPN14 in Hippo Signaling and Beyond. *Genes (Basel)* **7** (2016).
88. G. Singh, S. Mishra, H. Chander, KIBRA Team Up with Partners to Promote Breast
Cancer Metastasis. *Pathol Oncol Res* **26**, 627-634 (2020).
89. R. F. Hennigan, C. S. Thomson, K. Stachowski, N. Nassar, N. Ratner, Merlin tumor
suppressor function is regulated by PIP2-mediated dimerization. *PLoS One* **18**,
e0281876 (2023).
90. A. W. Hong *et al.*, Critical roles of phosphoinositides and NF2 in Hippo pathway
regulation. *Genes Dev* **34**, 511-525 (2020).
91. A. V. Anzalone *et al.*, Search-and-replace genome editing without double-strand breaks
or donor DNA. *Nature* **576**, 149-157 (2019).
92. G. Seo *et al.*, MAP4K Interactome Reveals STRN4 as a Key STRIPAK Complex
Component in Hippo Pathway Regulation. *Cell Rep* **32**, 107860 (2020).
93. Y. Tang *et al.*, Architecture, substructures, and dynamic assembly of STRIPAK
complexes in Hippo signaling. *Cell Discov* **5**, 3 (2019).
94. M. Kim *et al.*, cAMP/PKA signalling reinforces the LATS-YAP pathway to fully suppress
YAP in response to actin cytoskeletal changes. *EMBO J* **32**, 1543-1555 (2013).
95. Z. Meng *et al.*, MAP4K family kinases act in parallel to MST1/2 to activate LATS1/2 in
the Hippo pathway. *Nat Commun* **6**, 8357 (2015).
96. Y. Zheng *et al.*, Identification of Happyhour/MAP4K as Alternative Hpo/Mst-like Kinases
in the Hippo Kinase Cascade. *Dev Cell* **34**, 642-655 (2015).
97. S. C. Wu, M. Grace, K. Munger, The HPV8 E6 protein targets the Hippo and Wnt
signaling pathways as part of its arsenal to restrain keratinocyte differentiation. *mBio* **14**,
e0155623 (2023).
98. D. Dacus, C. Cotton, T. X. McCallister, N. A. Wallace, Beta Human Papillomavirus 8E6
Attenuates LATS Phosphorylation after Failed Cytokinesis. *J Virol* **94** (2020).
99. E. L. Morgan *et al.*, MicroRNA-18a targeting of the STK4/MST1 tumour suppressor is
necessary for transformation in HPV positive cervical cancer. *PLoS Pathog* **16**,
e1008624 (2020).
100. W. Yin *et al.*, HPV E6 inhibits E6AP to regulate epithelial homeostasis by modulating
keratinocyte differentiation commitment and YAP1 activation. *PLoS Pathog* **19**,
e1011464 (2023).
101. D. R. Singh *et al.*, Epstein-Barr virus LMP1 protein promotes proliferation and inhibits
differentiation of epithelial cells via activation of YAP and TAZ. *Proc Natl Acad Sci U S A*
120, e2219755120 (2023).
102. S. Wang *et al.*, Single cell transcriptomics of human epidermis identifies basal stem cell
transition states. *Nat Commun* **11**, 4239 (2020).
103. J. G. Doench *et al.*, Optimized sgRNA design to maximize activity and minimize off-
target effects of CRISPR-Cas9. *Nat Biotechnol* **34**, 184-191 (2016).
104. M. A. Dickson *et al.*, Human keratinocytes that express hTERT and also bypass a
p16(INK4a)-enforced mechanism that limits life span become immortal yet retain normal
growth and differentiation characteristics. *Mol Cell Biol* **20**, 1436-1447 (2000).
723
724

FIGURES AND TABLES

Figure 1. Phosphorylation on YAP1 S127 is reduced in PTPN14 knockout keratinocytes.

N/Tert-1 keratinocytes were transduced with LentiCRISPRv2 vectors encoding spCas9 and an

sgRNA sequence targeting PTPN14 or a nontargeting control sequence. sgRNA sequences

were chosen from the Broad Institute Brunello Library. Four cell lines were generated using two

different non-targeting control (sgNT-1, sgNT-2) and two different PTPN14 (sgPTPN14-1,

sgPTPN14-3) sgRNAs. Cells were treated with cytochalasin D for up to 90 minutes and whole

cell lysates were harvested at 30 minute intervals. Whole cell protein lysates were separated by

SDS-PAGE and proteins were detected by immunoblotting for YAP1, YAP1 pS127, PTPN14,

and actin.

Figure 2. Phosphorylation on LATS1 T1079 is reduced in PTPN14 knockout

keratinocytes. Control and PTPN14 knockout N/Tert-1 cells generated using LentiCRISPRv2

vectors were trypsinized and kept in suspension for 10 minutes. Whole cell protein lysates were

separated by SDS-PAGE and proteins were detected by immunoblotting for YAP1, YAP1 pS127,

LATS1, LATS1 pT1079, NF2, NF2 pS518, PTPN14, and actin. Panels (A, B) show replicate

experiments using the same four distinct cell lines as in Figure 1.

Figure 3. HPV18 E7-mediated PTPN14 degradation reduces phosphorylation on YAP1

S127 and LATS1 T1079. N/Tert-1 keratinocytes that stably express HA-tagged HPV18 E7 or

HA-tagged HPV18 E7 R84S, which is unable to bind or degrade PTPN14, were used in assays

of Hippo pathway activity. Cells transduced with empty vector were included as a control. (A)

Cells were treated with cytochalasin D for up to 90 minutes and whole cell lysates were

harvested at 30 minute intervals. Whole cell protein lysates were separated by SDS-PAGE and

proteins were detected by immunoblotting for YAP1, YAP1 pS127, PTPN14, RB1, HA, and

actin. (B) Cells were trypsinized and kept in suspension for 10 minutes. Whole cell protein

lysates were separated by SDS-PAGE and proteins were detected by immunoblotting for YAP1,

YAP1 pS127, LATS1, LATS1 pT1079, HA, PTPN14, and actin.

Figure 4. PTPN14 knockout impairs keratinocyte differentiation. (A) N/Tert-1 cells stably

expressing Cas9 were transfected with sgRNA targeting PTPN14 (Synthego) or mock

transfected and PTPN14 knockout was validated by immunoblot. (B) Mock transfected and

PTPN14 knockout N/Tert-1 cells were cultured in media containing 1.5 mM calcium and

harvested for RNA analysis at indicated times. KRT10 and IVL RNA levels were analyzed by

qRT-PCR and normalized to G6PD. Graphs show values for technical duplicate experiments.

Error bars display mean ± range.

Figure 5. PTPN14 promotes keratinocyte differentiation and Hippo pathway activity.

N/Tert-1 keratinocytes were transduced with a lentiviral vector encoding doxycycline-inducible

PTPN14 (pLIX-PTPN14). Cells were treated with 1 μg/mL doxycycline for 24 hours or left

untreated. (A) KRT10 and IVL RNA levels were measured by qRT-PCR and normalized to

GAPDH. Graphs show data points for two technical replicate experiments. Error bars display

mean ± range. (B) Whole cell protein lysates were separated by SDS-PAGE and proteins

analyzed by immunoblotting for YAP1, YAP1 pS127, LATS1, LATS1 pT1079, NF2, NF2 pS518,

PTPN14, and actin.

Figure 6. PTPN14 PY1/2 motifs are required to induce differentiation. (A) Schematic of

PTPN14 domains and location of PPxY motifs. (B) HFK were transfected with lentiviral

constructs encoding GFP, wild-type (WT) PTPN14, or PTPN14 mutants as indicated. Total

cellular RNA was collected and KRT1 transcript levels were analyzed by qRT-PCR and

normalized to GAPDH. Graph shows individual data points for three independent experiments.

Data is plotted as mean ± standard deviation. Significance was determined by ANOVA with

Holm-Šidak's multiple comparisons test. (ns, not significant; ****, P < 0.0001). (C) N/Tert-1

immortalized keratinocytes transduced with pLIX-PTPN14 were treated with 1 µg/mL

doxycycline for 48 hours to induce HA-tagged PTPN14 expression. Whole cell protein lysates

were collected and PTPN14 immunoprecipitated with anti-HA agarose beads. Protein lysates

from input and elution fractions were separated by SDS-PAGE and proteins were analyzed by

immunoblotting for WWC1, PTPN14, and actin.

Figure 7. LATS kinases and NF2 are required for PTPN14 to induce KRT10 expression.

HFK were transfected with siRNA then transduced with lentiviruses encoding GFP or PTPN14

at 24h post transfection. Total cellular RNA was collected 72h post-knockdown and 48h post-

transduction. RNA transcripts for KRT10 were measured by qRT-PCR and normalized to

GAPDH. Six individual experiments were conducted, each in technical duplicate. Each

experiment included an siControl condition, a YAP1 and TAZ siRNA-treated condition, and

siRNAs targeting additional component(s) of the Hippo pathway. Each component of the Hippo

pathway was targeted with two different siRNAs per gene, denoted A and B. Panels display data

from knockdowns as follows: (A) LATS1 and LATS2, (B) MST1 and MST2, (C) NF2, or (D)

WWC1, WWC2, and WWC3. Data are graphed as mean ± standard deviation of combined

replicate data. PTPN14/GFP denotes the ratio of KRT10 level in PTPN14 transduced cells vs

GFP transduced cells in each siRNA-treated condition. Since Experiment 3 included several

siRNAs (Control, YAP1/TAZ, MST1/2, NF2, WWC1/2/3), the same siControl and siYAP1/TAZ

data from experiment 3 is included in panels B, C, and D.

Figure 8. PTPN14 knockout promotes anchorage independent growth and reduces YAP1

phosphorylation in HEK TER cells. (A) HEK TER cells expressing spCas9 were transfected

with sgRNA targeting PTPN14 or LATS1 and LATS2. HEK TER cells expressing SV40 ST were

used as a positive control for colony formation. Cells were plated in soft agar in technical

triplicate and incubated at 37°C for 18 days, then photographed. Colonies were counted and

quantified using ImageJ software. Graphs show individual data points for each plate and

indicate mean ± standard deviation. Statistical significance of the nontargeting control condition

compared to experimental conditions was determined by ANOVA with Dunnett's multiple

comparisons test (*, P < 0.05; ***, P < 0.001). (B) Cells were treated with cytochalasin D for up

to 90 minutes and whole cell lysates were harvested at 30 minute intervals. Whole cell protein

lysates were separated by SDS-PAGE and proteins were detected by immunoblotting for YAP1,

YAP1 pS127, PTPN14, LATS1, and actin.

Figure S1. PTPN14 increases KRT10 protein levels. N/Tert-1 Cas9:sgPTPN14 keratinocytes

transduced with the pLIX-PTPN14 overexpression construct were incubated with 0, 1 or 2 µg/ml

doxycycline for 48 hours or 72 hours. Whole cell protein lysates were separated by SDS-PAGE

and proteins were detected by immunoblotting for HA, PTPN14, KRT10, and actin.

Figure S2. PTPN14 increases KRT1 and IVL transcript levels. HFK transfected with siRNA

for 72 hours were also transduced for 48 hours with lentiviral vectors for GFP or PTPN14

overexpression. Cells were transfected with siRNA: nontargeting control, YAP1, TAZ, and two

separate pairs for LATS1 and LATS2 denoted (A) and (B). Total cellular RNA was measured by

qRT-PCR for KRT1 and IVL for two biological replicate experiments, each performed in

technical duplicate. Transcript levels were normalized to GAPDH. Graphs show mean ±

standard deviation. Samples are the same as those analyzed in Figure 7.

Figure S3. Validation of siRNA knockdowns. HFK were transfected with siRNA then

transduced with lentiviruses encoding GFP or PTPN14 at 24h post transfection. Total cellular

RNA was collected 72h post-knockdown and 48h post-transduction. RNA transcripts for KRT10

were measured by qRT-PCR and normalized to GAPDH. Six individual experiments were

conducted, each in technical duplicate, and samples are those that are analyzed in Figure 7. (A)

Total cellular RNA was analyzed by qRT-PCR with primer sets as indicated to validate

knockdown efficiency. Transcript levels are normalized to GAPDH. Graphs show mean RNA

levels ± range. TAZ and LATS2 transcript levels were below the limit of detection and are not

shown. (B) PTPN14 transcripts were measured in all six experiments by qRT-PCR, normalized

to GAPDH, and plotted as mean ± range.

Figure S4. PTPN14 knockout promotes anchorage independent growth and suppresses

YAP1 phosphorylation in HEK TER cells. (A) HEK TER cells expressing Cas9 were

transfected with a second set of sgRNA compared to those used in Figure 8. HEK TER cells

expressing SV40 ST were used as a positive control for colony formation. Cells were plated in

soft agar in technical triplicates and incubated at 37°C for 18 days, then photographed. Colonies

were counted and quantified using ImageJ software. Graphs show individual data points for

each plate and indicate mean ± standard deviation. Statistical significance of the nontargeting

control condition compared to experimental conditions was determined by ANOVA with

Dunnett's multiple comparisons test (ns, not significant; ****, P < 0.0001). (B) Cells were treated

with cytochalasin D for up to 90 minutes and whole cell lysates were harvested at 30 minute

intervals. Whole cell protein lysates were separated by SDS-PAGE and proteins were detected

by immunoblotting for YAP1, YAP1 pS127, PTPN14, LATS1 and actin.

848
849   **Supplementary Table 1. List of plasmids used in this study.**

851   **Supplementary Table 2. List of antibodies used in this study.**

A

Untreated | Suspension

sgNT-1 | sgPTPN14-1 | sgNT-1 | sgPTPN14-1

YAP1 — pS127 — 75 kDa
Total — 75 kDa
LATS1 — pT1079 — 150 kDa
Total — 150 kDa
NF2 — pS518 — 75 kDa
Total — 75 kDa
PTPN14 — 150 kDa
Actin — 50 kDa

B

Untreated | Suspension

sgNT-2 | sgPTPN14-3 | sgNT-2 | sgPTPN14-3

YAP1 — pS127 — 75 kDa
Total — 75 kDa
LATS1 — pT1079 — 150 kDa
Total — 150 kDa
NF2 — pS518 — 75 kDa
Total — 75 kDa
PTPN14 — 150 kDa
Actin — 50 kDa

A

N/Tert-1 + Cas9

sgRNA
Mock
PTPN14-1

PTPN14 — 150 kDa
Actin — 50 kDa

B

KRT10

IVL

☐ Mock
● sgPTPN14-1

Fold change RNA

Time post CaCl$_2$ (days)

Time post CaCl$_2$ (days)

A

Fold change RNA

B

N/Tert-1
pLIX-PTPN14

A

1   FERM   341   900   PTP   1187

PY1/2   PY3   PY4   C1121

B

KRT1/GAPDH

20 ┤

15 ┤

10 ┤

5 ┤

0 ┤

GFP   WT ΔPY1/2 ΔPY3l4 C1121S

PTPN14

▲ Exp 1
● Exp 2
■ Exp 3

**** ns **** ****

C

pLIX-PTPN14

WT   WT   ΔPY1/2 ΔPY3/4

Dox: -   +   +   +

IP: HA
WWC1 ——— 150 kDa
PTPN14 ——— 150 kDa

Input
WWC1 ——— 150 kDa
PTPN14 ——— 150 kDa
Actin ——— 50 kDa

A

B

HEK TER

LentiCas9

| | sgNT-1 | sgPTPN14-1 | sgLATS1/2-1 | SV40 ST |
Cyto D (min): 0 30 60 90 | 0 30 60 90 | 0 30 60 90 | 0 30 60 90

YAP1 ⎰ pS127 — 75 kDa
⎱ Total — 75 kDa

PTPN14 — 150 kDa

LATS1 — 150 kDa

Actin — 50 kDa