Urban *Cepaea nemoralis* snails are less likely to have nematodes trapped within their shells

Maxime Dahirel*, Hannah Reyné, Katrien De Wolf, Dries Bonte

Ghent University, Department of Biology, B-9000 Gent

ORCID:

MD: 0000-0001-8077-7765

KDW: 0000-0002-9255-913X

DB: 0000-0002-3320-7505

*corresponding author: maxime.dahirel@yahoo.fr

Abstract

Urbanization is a major human-induced environmental change which can impact not only individual species, but also the way these species interact with each other. As a group, terrestrial molluscs interact frequently with a wide diversity of parasites, yet the way these interactions vary across space and in response to environmental pressures is poorly documented. In this study we leveraged a recently discovered defence mechanism, by which snails trap parasitic nematodes in their shells, to explore how snail-nematodes interactions may vary in response to city life. We examined shells from the generalist snail *Cepaea nemoralis* sampled in three urban areas in Belgium for trapped nematodes, and attempted to link this to urbanization and shell phenotypic traits. We found that even a small degree of urbanization led to large decreases in the rates of shell encapsulation, and that larger snails were more likely to contain trapped nematodes. However, we found no evidence that shell colour, which had been previously linked to immune function, was correlated to encapsulation rates. We discuss how between-population variation in encapsulation rates can result from urbanization-induced changes on the nematodes side, the snail side, or both, and suggest potential tests for future studies aiming to disentangle these mechanisms.

Key words: Biotic interactions, Gastropoda, immunity, parasites.

Introduction

Urbanization is a major and all-encompassing human-induced environmental change, leading to changes in land use, local climate, soil imperviousness, light and chemical pollution… (Parris, 2016). The (often negative) impacts of these multivariate changes on biodiversity are increasingly well-documented: many species decline in cities, while some become successful "urban adapters", leading to major restructuring of biological communities along urbanization gradients (e.g. McKinney, 2008; Piano et al., 2020; Fenoglio et al., 2020; Liang et al., 2023). In parallel, urbanization can also cause within-species phenotypic and genetic changes (Alberti et al., 2017; Szulkin et al., 2020; Diamond & Martin, 2021). Urbanization may also have second-order impacts by reshaping ecological interactions, if tightly connected species respond to environmental change in different ways (Theodorou, 2022). Such urbanization-induced changes in ecological interactions, in both positive and negative directions, have been recorded for plant-pollinator interactions (Liang et al., 2023), plant-herbivore and prey-predator interactions (Eötvös et al., 2018; Valdés-Correcher et al., 2022; Gámez et al., 2022; Korányi et al., 2022), as well as host-parasite interactions (Murray et al., 2019; Korányi et al., 2022).

Terrestrial molluscs (snails and slugs) are potentially valuable models in urban ecology and evolution, in part because of their limited movement abilities, which means they often cannot move to escape environmental changes. Like in many other taxa, urbanization can reshape molluscan communities (Lososová et al., 2011; Horsák et al., 2013; Barbato et al., 2017; Hodges & McKinney, 2018), and drive evolutionary responses in urban populations (Kerstes et al., 2019). Interestingly, in a comparative cross-taxon study of urbanization impacts, snail species richness were less negatively affected, compared to other more mobile groups (Piano et al., 2020). Land molluscs are hosts to a diverse array of metazoan parasites, including nematodes, flies, mites or trematodes (Barker, 2004; Segade et al., 2013; Żbikowska et al., 2020). How urbanization reshapes these interactions remains understudied, despite some of these parasites being of increasing veterinary interest (Aziz et al., 2016; Giannelli et al., 2016).

Snails and slugs can defend themselves against metazoan parasites through a variety of behavioural (Wilson et al., 1999; Wynne et al., 2016; Rae, 2023) or immune responses (Furuta & Yamaguchi, 2001; Scheil et al., 2014; Coaglio et al., 2018). Among the latter, it has been discovered that land molluscs can use their shells to trap parasitic nematodes, killing them and fusing them to the inner shell surface (Rae et al., 2008; Williams & Rae, 2015; Rae, 2017). This ability seems phylogenetically widespread, even present in slugs with vestigial shells (Rae et al., 2008; Rae, 2017), and could therefore provide a relatively easy to access record of ecological interactions. Following anecdotal records of mites and trematodes encapsulated in shells, it has further been suggested that this shell encapsulation might extend to other metazoan parasites (Dahirel et al., 2022; Gérard et al., 2023). However, given how rare these non-nematode records are, they may be merely by-products of a defence mechanism targeted towards nematodes, rather than evidence of a more generalized defence response (Gérard et al., 2023). The few snail species in which this phenomenon has been studied across multiple populations show that the prevalence of individuals trapping nematodes can vary widely between sites (Rae, 2017; Rae, 2018; Cowlishaw et al., 2019), but there has been no attempt, to our knowledge, to assess whether this variation could be non-random with respect to environmental context.

To that end, we combine here publicly available and standardized urbanization metrics with
observations of field-collected snails across three cities in Belgium, using the grove snail *Cepaea*
nemoralis (Gastropoda, family Helicidae) as a model. Like other helicids, *C. nemoralis* can
encapsulate and trap parasitic nematodes in its shell (Williams & Rae, 2016; Rae, 2017; Dahirel et
al., 2022; Gérard et al., 2023). This snail is also common both outside and within cities (Kerstes et
al., 2019), and therefore a very suitable model to study variation in encapsulation rates, whether
it is due to urbanization or to spatial (between-cities) differences. Furthermore, the shell colour
variation that made *Cepaea* species iconic models in evolutionary biology (Jones et al., 1977;
Ożgo, 2009) may also influence their immune response, with some evidence that darker morphs
mount better defences against nematodes (Dahirel et al., 2022; but see Scheil et al., 2014). On
the other hand, this morph variation in resistance might not translate to shell encapsulation
(Williams & Rae, 2016; Dahirel et al., 2022). However, existing comparisons were either limited
to one type of colour variation (banding pattern only, Dahirel et al., 2022), or analysed
experimental infections by one model nematode (Williams & Rae, 2016); we here test whether
this remains true when analysing naturally occurring snail-nematode interactions and accounting
for more dimensions of shell colour variation.

Methods

Site selection and sampling

We searched for *Cepaea nemoralis* snails from early October to mid-November 2022 in and
around the urban areas of Brussels, Ghent and Leuven in Belgium (**Fig. 1**). Potential sites were
selected based on pre-existing online crowdsourced records (iNaturalist contributors &
iNaturalist, 2024) combined with personal observations and virtual fieldwork using Google
StreetView to identify suitable habitats (based on Falkner et al., 2001). We visited 36 sites chosen
to be roughly balanced between the three cities (including their surrounding areas; Brussels: 13
sites, Ghent: 13 sites, Leuven: 10 sites). In each site, we sampled living snails by hand during visual
search, in a radius of up to 50 m around a designated site centroid (though search was *de facto*
mostly concentrated within a 20 m radius). Individuals were mainly searched in known favourable
micro-habitats, i.e. on tall herbs and shrubs, under piled wood and cardboard or loose rocks, or
on fences, walls, and tree trunks (Falkner et al., 2001). Field identification of *Cepaea nemoralis*
snails is easy based on shell shape, size and colour (Cameron, 2008). We only collected adults,
which can easily be separated from subadults by the presence of a reflected shell lip marking the
end of shell growth (Cameron, 2008). Each site was visited by 1 to 3 people (mean: 2.03) for a
duration of 5 to 30 person-minutes (mean: 15). We collected a total of 298 snails from 28 of the
36 sites visited (Brussels: 9 sites, Ghent: 10 sites, Leuven: 9 sites). However, 2 shells were lost
before examination for parasites due to handling errors, and another shell was accidentally
broken for parasite examination before photographs or size measurements could be done; this
led to a final complete dataset of 295 snails in 28 sites. For each of these 28 sites, the nearest
neighbouring site with snails found was between 153 and 1516 m away (mean: 768 m), which is
in any case farther than the maximal dispersal distances (Kramarenko, 2014), indicating that even
nearby sites could be considered separate populations.

Figure 1. Location of study sites within western Europe and central Belgium. The Functional Urban Areas (roughly corresponding to commuter zones, Schiavina et al., 2019; Moreno-Monroy et al., 2021) that were used to link each site to a city are also displayed as solid black lines, while the corresponding core urban areas (Urban Centres *sensu* Eurostat (European Commission), 2021; Schiavina, Melchiorri, et al., 2023) are displayed with dashed lines.

Urbanization metrics

It is well-known that urban environmental changes are complex and multivariate (e.g. Parris, 2016); however, given our relatively low number of sites, and the risk of collinearity between urban metrics, we decided to use simple overarching metrics focused on building presence and human population density. We assessed urbanization at each site using raster layers from the Global Human Settlement Layer project for the year 2020 (https://ghsl.jrc.ec.europa.eu/, Joint Research Centre (European Commission), 2023). We first used built-up surface (GHS-BUILT-S) and population density (GHS-POP) at 100 m and 1000 m resolutions (Pesaresi & Politis, 2023; Schiavina, Freire, et al., 2023). The former spatial scale matches the scale of maximal dispersal movements over timespans of up to a couple years in helicid snails, while the latter is closer to the scale of longer term (over several decades) population spread (Kramarenko, 2014). As an additional categorical metric, we also used the Degree of Urbanization as recorded in the Settlement Model layer (GHS-SMOD, available only at 1000 m resolution, Eurostat (European Commission), 2021; Schiavina, Melchiorri, et al., 2023). At the highest level of classification, the standardized Degree of Urbanization methodology mainly uses population density and contiguity rules to classify grid cells as either part of a continuous high-density Urban Centre, as low-density rural cells or as intermediate peri-urban/suburban cells. For each site and urbanization metric, we recorded the value of the corresponding grid cell. Interestingly, Degree of Urbanization classes, primarily based on population density, divide our sites in almost the same non-linear way as another, independent, three-level classification based on built-up surfaces used in previous urban ecology studies in the study region (e.g. Piano et al., 2020) (**Supplementary Material S1**).

Snail shell analysis

Snail size was measured using a caliper as the shell greatest diameter (to the nearest 0.1 mm). Snail shell colour morphs were scored following e.g. Cain (1988) for background colour (from lighter to darker: yellow, pink or brown), number of dark bands (0 to 5 bands) and on the presence or absence of band fusions (which increase the proportion of the shell covered by dark bands). Snails were killed by first inducing dormancy at 6°C, then by freezing at -20°C. We removed bodies from shells with forceps and lightly cleaned shells with water (bodies were stored in ethanol for separately planned studies). We then broke each shell into fragments using forceps, examined fragments under a binocular microscope, and recorded all animals found encapsulated within the shell as in e.g. Gérard et al. (2023). A total of 606 nematodes were found in 104 shells (**Fig. 2**); we found no mites, trematodes or other parasites in any of the shells. Shells with nematodes contained 5.83 nematodes on average (SD: 9.95, range: 1-58). As this method is destructive, we took standardised photographs of the shells beforehand (dorsal and apertural views following Callomon, 2019) for archival and potential future studies.

Figure 2. Fragment of a *Cepaea nemoralis* shell (A) containing encapsulated nematodes (B, C). The arrow in (B) points to the nematode shown in (C).

 Statistical analysis

All analyses were done in R version 4.3.2 (R Core Team, 2023), with the help of the *tidyverse*
(Wickham et al., 2019) and *sf* (Pebesma, 2018) packages for data processing, as well as additional
packages detailed below for model fitting and exploration.

We analysed the probability a shell contained nematodes as a binary yes/no response at the
individual level, using Generalized Linear Mixed Models (GLMMs) (binomial family, logit link). We
ran six models; the first five all included shell size, shell morph traits (background colour, band
number and fusion), urbanization and city identity (Brussels, Ghent or Leuven) as fixed effects,
only differing by which urbanization metric they used (among the five described above in
Urbanization metrics). Given our sample size, we did not include interactions between our
explanatory variables, especially as we had no *a priori* hypotheses regarding these (but see
Discussion). Numeric predictors were centred and scaled to unit 1 SD. Sampling site was included
as a random intercept. The sixth model was a "null" model, identical to the other ones except
that it did not include an urbanization metric. We ran our models using the *glmmTMB* package
(Brooks et al., 2017), and then used AICc to compare them. As one model largely outperformed
the others (see **Results**), we did all further analyses on that best model.

We checked for residual spatial autocorrelation using a spline correlogram (*ncf* package,
Bjornstad, 2022), and found no evidence of spatial structure in the best model. We then used the
car (Fox & Weisberg, 2019) and *emmeans* (Lenth, 2023) packages to test for overall effects of our
variables in the best model and to run (Tukey-corrected) pairwise comparisons, respectively.
Finally, we estimated the marginal and conditional R^2 (Nakagawa & Schielzeth, 2013) as measures
of the proportion of variation explained by fixed effects (R_m^2) and both fixed and random effects
(R_c^2) respectively (using the delta method, Nakagawa et al., 2017).

Results

The model using the categorical Degree of Urbanization (GHS-SMOD) as an urbanization metric
outperformed all other models based on AICc (**Table 1**). Fixed effects and random effects
explained similar amounts of variance ($R_m^2 = 0.19$; $R_c^2 = 0.38$). The probability that a shell had
trapped nematodes was dependent on urbanization level ($\chi^2 = 15.97$, df $= 2, p = 3.40 \times 10^{-4}$)
but did not vary significantly between cities ($\chi^2 = 3.89$, df $= 2, p = 0.14$). Snails from rural sites
were more likely to contain nematodes than snails from intermediate and Urban Centre
populations (**Fig. 3**; rural - intermediate difference on the logit scale $\pm$ SE: 3.71 ± 0.95 ; rural -
Urban Centre difference: 2.34 ± 0.77). Larger shells were more likely to contain nematodes ($\chi^2 =$
4.17, df $= 1, p = 0.04$; standardised coefficient $\beta = 0.35 \pm 0.17$). There was no clear evidence
that any of the shell colour traits affected encapsulation rates (background colour: $\chi^2 = 2.17$,
df $= 2, p = 0.34$; band number: $\chi^2 = 1.90$, df $= 1, p = 0.17$; fusion: $\chi^2 = 0.17$, df $= 1, p =$
0.68).

Table 1. Model selection table for the effect of urbanization on shell encapsulation rates. All models otherwise include effects of city identity, shell size and shell morph (background colour, number of bands and band fusion).

Urbanization variable in model	df	log-likelihood	AICc	Δ	AICc weight
Degree of Urbanization categories (SMOD, 1000 m resolution grid)	11	-149.5	322.0	0.00	0.95
None ("null" model)	9	-155.9	330.4	8.43	0.01
Population density (1000 m resolution grid)	10	-155.2	331.1	9.11	0.01
Built-up surface (1000 m resolution grid)	10	-155.4	331.5	9.49	0.01
Population density (100 m resolution grid)	10	-155.4	331.6	9.61	0.01
Built-up surface (100 m resolution grid)	10	-155.6	331.9	9.91	0.01

Figure 3. Effect of the Degree of Urbanization in 1000 m grid cells on the probability a snail shell contained encapsulated nematodes. Grey dots are observed proportions per population, with the size of the dot proportional to the number of snails; white dots (and error bars) are estimated marginal means from the best model (and their 95% confidence intervals), with the effects of the other predictors averaged out.

Discussion

We found that the prevalence of *Cepaea nemoralis* snails encapsulating nematodes in their shell as a defence mechanism were partly driven by environmental conditions, with lower trapping rates in more urban sites (**Fig. 3**). This effect was better explained by a categorical classification of the Degree of Urbanization, rather than by linear effects of continuous urbanization variables. This indicates that the response to urbanization is non-linear, as the increases in population density/built-up rates needed to go from a rural to an intermediate area are much smaller than those needed to go from intermediate to Urban Centre, and most of the variation in density/built-up is within Urban Centres rather than between categories (**Supplementary Material S1**).

A difficulty for interpreting our results is that nematodes trapped in shells accumulate with time (Williams & Rae, 2015; Rae, 2017), meaning that as they may have endured more infections, older snails may be more likely to have them. If the urban heat island influences snail survival rates (Wolda, 1967, Manoli et al., 2019), then our urban-rural differences in nematodes trapped might merely reflect differences in average snail age/survival. Age estimation in terrestrial snails is challenging (Pollard et al., 1977; Williamson, 1979), and any age proxy is likely to be influenced by local conditions, making it useless to compare age between populations without thorough calibration studies. However, while there is substantial population variation, the number of nematodes found in infected shells does not decrease in more urbanized sites (**Supplementary Material S2**), contrary to what we would expect if variation in shell encapsulation was primarily explained by variation in time available to accumulate nematodes.

If we assume that our results reflect differences in snail-nematodes interactions between urban and non-urban areas, several mutually non-exclusive mechanisms may explain why urban *Cepaea nemoralis* shells are less likely to trap nematodes. Each of these mechanisms directly suggests potential tests for future studies:

- First, snail parasitic nematodes infecting *C. nemoralis* may be less abundant in cities. Many nematodes known to infect land snails have at least one free-living life stage in the soil, and some are facultative parasites (Morand et al., 2004; Pieterse et al., 2017). Increasingly impervious substrates in cities (Parris, 2016) may deprive these of habitat critical for their life cycle. Where habitat is available, soil nematode communities are profoundly altered by urbanization, like other taxa (Li et al., 2022; Gong et al., 2024). However, this does not lead to overarching declines in nematode abundance; rather, some trophic groups decline while others thrive (Li et al., 2022; Gong et al., 2024). Unfortunately, detailed information on nematodes parasitizing animals is typically lacking from these analyses; soil sampling specifically targeting parasitic nematodes (Jaffuel et al., 2019) would be here particularly useful. The few studies available are mixed on the effects of urbanization on the infection of land molluscs by parasitic nematodes. In Wales, urban and suburban slugs are more, not less, likely to be infected by *Angiostrongylus vasorum* compared to rural ones (Aziz et al., 2016). By contrast, data from Andrus et al. (2022) spanning urban and non-urban sites suggest that the prevalence of nematode infection may be slightly lower in urban molluscs, although they did not themselves analyze the effect of urbanization. Both

studies however analyzed nematode prevalence in molluscs, not their
abundance/availability in the urban environment.

• Second, individual differences in behaviour, especially space-related behaviour, may lead
to differences in the risk of encountering and then being infected by parasites (Barber &
Dingemanse, 2010). Habitat loss and fragmentation associated with urbanization are
expected to exert strong selection pressures on movement and space use (Cote et al.,
2017). If this results in lower movement in urban snail populations, this might then reduce
their encounter rates with parasites. In the snail *Cornu aspersum*, urbanization does not
lead to reduced habitat boundary-crossing behaviour (Dahirel et al., 2016), although that
is only one component of mobility. Urbanization-induced increases in temperature may
also alter the frequency at which snails hide into shelters or climb above the substrate
(Rosin et al., 2018), and potentially again the risk of encountering parasites. The picture is
complicated by behaviour-parasite feedbacks, where while host behaviour shapes
infection risk, infection can then alter host behaviour in turn (Ezenwa et al., 2016). In
Cepaea nemoralis, nematode infection itself might lead to reduced movement propensity,
but only in some morphs (Dahirel et al., 2022). More studies of movement behaviour
across urbanization gradients are here needed.

• Third, shell encapsulation rates are not direct records of snail-nematode interactions, but
rather informative on the host's ability to mount a defence in such interactions. This
defence is not always effective, as field-caught snails sometimes show active infections
but zero shell-trapped nematodes (see e.g. data in Dahirel et al., 2022). If immune
response declines with urbanization, then this alone could explain our results even in the
absence of changes in nematode communities. In vertebrates, urban living can lead to
both depressed or stimulated immune function, depending on taxon and context,
especially food availability (Murray et al., 2019; Minias, 2023). In terrestrial molluscs,
chemical pollutants seem to negatively impact many, but not all, physiological
components of immune defence (Radwan et al., 2020). The exact physiological pathways
involved in shell encapsulation in land molluscs remain however unstudied, to the best of
our knowledge.

Interestingly, nematode encapsulation prevalence was seemingly more variable between Urban
Centre populations than between populations in the other urban categories, with a few sites
having observed prevalences largely above the predicted mean (**Fig. 3**). While this may simply be
sampling variability as these populations have very small sample sizes, this suggests that there
could be non-random within-city variability in snail-nematode interactions. As a first post-hoc
exploration, we have re-run the models with continuous urbanization variables as predictors,
using only the Urban Centre subset of sites (see **Data and code availability**). After accounting for
phenotype and city of origin, we found no indication that built-up or population density levels
influenced prevalence within the Urban Centre category. Nonetheless, cities remain highly
heterogeneous environments even beyond built-up and population density; for instance, within-
city variation in vegetation, mediated in part by neighborhood-level socio-economic differences,
may shape biodiversity, including species interactions (e.g. Martin et al., 2024). While we are not
able to identify the causes of this heterogeneity in our current dataset, as we are hampered by

our small number of sites per Urban Centre, future studies designed to target this within-city
heterogeneity may uncover more on the fine-scale drivers of snail responses to parasites.

On the individual phenotype side, larger shells were more likely to contain trapped nematodes. If
shell size also varied in response to urbanization, then this could open an indirect pathway linking
urbanization to encapsulation mediated by snail size, potentially accentuating or dampening the
direct effect we describe above. However, we found no clear effect of urbanization on *C.*
nemoralis shell size (**Supplementary Material S3**). In addition and as a post-hoc exploration, we
re-ran our model set adding size × urbanization interactions, and found no significant interaction,
and no evidence that the urbanization effect changed in response (**Supplementary Material S4**).
The relationship between size and nematode encapsulation could be the result of survivor bias
alone, if larger snails are more likely to survive infection. However, and although we cannot
exclude that other nematodes have larger effects, experimental nematode infections by
Phasmarhabditis are almost never lethal in adult *Cepaea nemoralis*, contrary to other snail
species (Wilson et al., 2000; Williams & Rae, 2016). Other potential explanations for this result
can be sorted along three non-exclusive lines, similar to the mechanisms suggested above to
explain the effect of urbanization:

• Larger snails might harbour larger parasite infections (e.g. Daniels et al., 2013), which
would increase the likelihood that some nematodes are trapped. However, there is no link
between nematode abundance in active infections and snail size in *C. nemoralis* (Dahirel
et al., 2022), and no clear effect of shell size on the number of nematodes trapped in the
present study (**Supplementary Material S2**).

• If large and small snails differ in their space use, they might also differ in their parasite
exposure risk. Evidence for a link between shell size and space use is mixed in *Cepaea*
nemoralis, and this may depend on the scale of the movements in question (short-term
routine vs. dispersal movements; Oosterhoff, 1977; Dahirel et al., 2022).

• Finally, small and large snails may differ in their immune defence abilities. Comparative
studies suggest that large and small snail species and subspecies differ in their immune
strategies at the physiological level (Russo & Madec, 2011, 2013). However, the range of
body size and life history variation is much larger in these scenarios than among adults of
C. nemoralis, limiting the transferability of these results. More physiological studies
focused on within-, rather than among-species variation may help understand better this
link between body size and encapsulation rates.

In contrast to shell size, we found no relationship between any of the shell colour traits and
nematode trapping rate. This confirms experimental results from Williams & Rae (2016) using
infections by *Phasmarhabditis hermaphrodita*. However, colour morphs do differ in active
infection rates or other aspects of immune response in *C. nemoralis* (Dahirel et al., 2022) and
other polymorphic snails (Scheil et al., 2013, 2014). This discrepancy may indicate that shell
encapsulation is driven by different physiological pathways than other components of snail
immune defence.

Beyond the effects of phenotype or environment, whether and how the prevalence of nematodes trapped in shells is correlated with rates of active parasite infections remains an open and complex question (which we could not tackle here as snail bodies were reserved for other investigations). If variation in snail-nematode interactions is driven by e.g. variation in nematode density in the environment, we may expect a positive correlation, as higher nematode densities should drive up rates of both shell encapsulation (Rae, 2018) and active infection (although if encapsulation is highly effective, it may end up suppressing dose-dependent effects on active infection, Williams & Rae, 2015). On the other hand, if variation is mostly driven by snail immune response, we may expect a negative correlation: snails with more effective immune systems may be more likely to successfully trap nematodes in shells while being less likely to harbour active infections. While this would need to be validated, the strength and direction of between- and within-sites correlations between active infections and shell-trapped nematodes may provide useful indicators of the main drivers of snail-nematodes interactions in response to city life.

We acknowledge that the relatively small size of our sample does not allow us to draw firm causal conclusions. Nonetheless, we hope our results may encourage larger studies regarding host-parasite interactions in land molluscs in the context of environmental change. As new technical developments such as micro-CT imaging allow non-destructive analyses of snail shells (Falkingham & Rae, 2021), these may extend to using museum and other natural history collections to understand how interactions vary in space and time (Cowlishaw et al., 2019), reaffirming their value for urban ecology and evolution (Shultz et al., 2020).

Author contributions

Initial study idea: MD, DB. Site selection and fieldwork: MD, HR, KDW. Shell data collection: HR, after initial training by MD. Data analysis: MD, after preliminary analyses by HR. Initial manuscript draft: MD. All authors contributed critically to edits and gave final approval for publication.

Funding

This study has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 101022802 (HELICITY). KDW and DB received funding from the Fund for Scientific Research Flanders (grant G080221N).

Conflict of interest disclosure

The authors declare they have no financial conflict of interest in relation with the content of this article. DB is a recommender for PCI Ecology and PCI Evolutionary Biology.

Data and code availability

Data and R scripts to reproduce all analyses presented in this article, as well as a copy of the Supplementary Materials, are available on Github (https://github.com/mdahirel/HELICITY-2022_shell-nematodes) and archived in Zenodo (DOI: https://doi.org/10.5281/zenodo.10794928).

References

Alberti M, Correa C, Marzluff JM, Hendry AP, Palkovacs EP, Gotanda KM, Hunt VM, Apgar TM, Zhou Y (2017) Global urban signatures of phenotypic change in animal and plant populations. *Proceedings of the National Academy of Sciences*, 201606034. https://doi.org/10.1073/pnas.1606034114

Andrus PS, Rae R, Wade CM (2022). Nematodes and trematodes associated with terrestrial gastropods in Nottingham, England. *Journal of Helminthology*, **96**, e81. https://doi.org/10.1017/S0022149X22000645

Aziz NAA, Daly E, Allen S, Rowson B, Greig C, Forman D, Morgan ER (2016) Distribution of *Angiostrongylus vasorum* and its gastropod intermediate hosts along the rural–urban gradient in two cities in the United Kingdom, using real time PCR. *Parasites & Vectors*, **9**, 56. https://doi.org/10.1186/s13071-016-1338-3

Barbato D, Benocci A, Caruso T, Manganelli G (2017) The role of dispersal and local environment in urban land snail assemblages: an example of three cities in Central Italy. *Urban Ecosystems*, **20**, 919–931. https://doi.org/10.1007/s11252-017-0643-8

Barber I, Dingemanse NJ (2010) Parasitism and the evolutionary ecology of animal personality. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **365**, 4077–4088. https://doi.org/10.1098/rstb.2010.0182

Barker GM (Ed.) (2004) *Natural enemies of terrestrial molluscs*. CABI, Wallingford, UK.

Bjornstad ON (2022) R package ncf: Spatial covariance functions.

Brooks M E., Kristensen K, Benthem K J.,van, Magnusson A, Berg C W., Nielsen A, Skaug H J., Mächler M, Bolker B M. (2017) GlmmTMB balances speed and flexibility among packages for zero-inflated generalized linear mixed modeling. *The R Journal*, **9**, 378. https://doi.org/10.32614/RJ-2017-066

Cain AJ (1988) The scoring of polymorphic colour and pattern variation and its genetic basis in molluscan shells. *Malacologia*, **28**, 1–15.

Callomon P (2019) *Standard views for imaging mollusk shells*. American Malacological Society.

Cameron R (2008) *Land Snails in the British Isles*. FSC Publications, Telford, UK.

Coaglio AL, Ferreira MAND, dos Santos Lima W, de Jesus Pereira CA (2018) Identification of a phenoloxidase- and melanin-dependent defence mechanism in *Achatina fulica* infected with *Angiostrongylus vasorum*. *Parasites & Vectors*, **11**, 113. https://doi.org/10.1186/s13071-018-2710-2

Cote J, Bestion E, Jacob S, Travis J, Legrand D, Baguette M (2017) Evolution of dispersal strategies and dispersal syndromes in fragmented landscapes. *Ecography*, **40**, 56–73. https://doi.org/10.1111/ecog.02538

Cowlishaw RM, Andrus P, Rae R (2019) An investigation into nematodes encapsulated in shells of
wild, farmed and museum specimens of *Cornu aspersum* and *Helix pomatia*. *Journal of*
Conchology, **43**, 385–392.

Dahirel M, Proux M, Gérard C, Ansart A (2022) Morph-dependent nematode infection and its
association with host movement in the land snail *Cepaea nemoralis* (Mollusca, Gastropoda).
Journal of Zoology, **318**, 181–192. https://doi.org/10.1111/jzo.13012

Dahirel M, Séguret A, Ansart A, Madec L (2016) Dispersal-related traits of the snail *Cornu*
aspersum along an urbanisation gradient: maintenance of mobility across life stages despite high
costs. *Urban Ecosystems*, **19**, 1847–1859. https://doi.org/10.1007/s11252-016-0564-y

Daniels RR, Beltran S, Poulin R, Lagrue C (2013) Do parasites adopt different strategies in different
intermediate hosts? Host size, not host species, influences *Coitocaecum parvum* (Trematoda) life
history   strategy,   size   and   egg   production. *Parasitology*,   **140**,   275–283.
https://doi.org/10.1017/S0031182012001564

Diamond SE, Martin RA (2021) Evolution in cities. *Annual Review of Ecology, Evolution, and*
Systematics, **52**, 519–540. https://doi.org/10.1146/annurev-ecolsys-012021-021402

Eötvös CB, Magura T, Lövei GL (2018) A meta-analysis indicates reduced predation pressure with
increasing   urbanization. *Landscape   and   Urban   Planning*,   **180**,   54–59.
https://doi.org/10.1016/j.landurbplan.2018.08.010

Eurostat (European Commission) (2021) *Applying the degree of urbanisation: a methodological*
manual to define cities, towns and rural areas for international comparisons : 2021 edition.
Publications Office of the European Union, LU.

Ezenwa VO, Archie EA, Craft ME, Hawley DM, Martin LB, Moore J, White L (2016) Host behaviour–
parasite feedback: an essential link between animal behaviour and disease ecology. *Proceedings*
of   the   Royal   Society   B: Biological   Sciences,   **283**,   20153078.
https://doi.org/10.1098/rspb.2015.3078

Falkingham P, Rae R (2021) 3D morphology of nematode encapsulation in snail shells, revealed
by micro-CT imaging. *Scientific Reports*, **11**, 2523. https://doi.org/10.1038/s41598-021-82106-6

Falkner G, Obrdlik P, Castella E, Speight MCD (2001) *Shelled Gastropoda of western Europe*.
Friedrich Held Gesellschaft, München, Germany.

Fenoglio MS, Rossetti MR, Videla M (2020) Negative effects of urbanization on terrestrial
arthropod communities: a meta-analysis. *Global Ecology and Biogeography*, **29**, 1412–1429.
https://doi.org/10.1111/geb.13107

Fox J, Weisberg S (2019) *An R Companion to Applied Regression*. Sage, Thousand Oaks, CA.

Furuta E, Yamaguchi K (2001) Haemolymph: blood cell morphology and function. In: *The biology*
of terrestrial molluscs (ed Barker GM), pp. 289–306. CABI, Wallingford, UK.

Gámez S, Potts A, Mills KL, Allen AA, Holman A, Randon PM, Linson O, Harris NC (2022) Downtown
diet: a global meta-analysis of increased urbanization on the diets of vertebrate predators.
Proceedings of the Royal Society B: Biological Sciences, **289**, 20212487.
https://doi.org/10.1098/rspb.2021.2487

Gérard C, De Tombeur Y, Dahirel M, Ansart A (2023) Land snails can trap trematode cercariae in
their shell: Encapsulation as a general response against parasites? *Parasite*, **30**, 1.
https://doi.org/10.1051/parasite/2023001

Giannelli A, Cantacessi C, Colella V, Dantas-Torres F, Otranto D (2016) Gastropod-borne
helminths: a look at the snail–parasite interplay. *Trends in Parasitology*, **32**, 255–264.
https://doi.org/10.1016/j.pt.2015.12.002

Gong X, Qiao Z, Yao H, Zhao D, Eisenhauer N, Scheu S, Liang C, Liu M, Zhu Y-G, Sun X (2024)
Urbanization simplifies soil nematode communities and coincides with decreased ecosystem
stability. *Soil Biology and Biochemistry*, **190**, 109297.
https://doi.org/10.1016/j.soilbio.2023.109297

Hodges MN, McKinney ML (2018) Urbanization impacts on land snail community composition.
Urban Ecosystems, **21**, 721–735. https://doi.org/10.1007/s11252-018-0746-x

Horsák M, Lososová Z, Čejka T, Juřičková L, Chytrý M (2013) Diversity and biotic homogenization
of urban land-snail faunas in relation to habitat types and macroclimate in 32 Central European
cities. *PLOS ONE*, **8**, e71783. https://doi.org/10.1371/journal.pone.0071783

iNaturalist contributors, iNaturalist (2024) iNaturalist Research-grade Observations.
iNaturalist.org. Occurrence dataset. https://doi.org/10.15468/ab3s5x

Jaffuel G, Půža V, Hug A-S, Meuli RG, Nermuť J, Turlings TCJ, Desurmont GA, Campos-Herrera R
(2019) Molecular detection and quantification of slug parasitic nematodes from the soil and their
hosts. *Journal of Invertebrate Pathology*, **160**, 18–25. https://doi.org/10.1016/j.jip.2018.11.005

Joint Research Centre (European Commission) (2023) *GHSL data package 2023*. Publications
Office of the European Union, LU.

Jones JS, Leith BH, Rawlings P (1977) Polymorphism in *Cepaea*: a problem with too many
solutions? *Annual Review of Ecology and Systematics*, **8**, 109–143.

Kerstes NAG, Breeschoten T, Kalkman VJ, Schilthuizen M (2019) Snail shell colour evolution in
urban heat islands detected via citizen science. *Communications Biology*, **2**, 264.
https://doi.org/10.1038/s42003-019-0511-6

Korányi D, Egerer M, Rusch A, Szabó B, Batáry P (2022) Urbanization hampers biological control
of insect pests: A global meta-analysis. *Science of The Total Environment*, **834**, 155396.
https://doi.org/10.1016/j.scitotenv.2022.155396

Kramarenko S (2014) Active and passive dispersal of terrestrial mollusks: a review. *Ruthenica*, **24**,
1–14.

Lenth R (2023) emmeans: estimated marginal means, aka least-squares means (R package).

Liang H, He Y-D, Theodorou P, Yang C-F (2023) The effects of urbanization on pollinators and
pollination: A meta-analysis. *Ecology Letters*, **26**, 1629–1642. https://doi.org/10.1111/ele.14277

Li X, Liu T, Li H, Geisen S, Hu F, Liu M (2022) Management effects on soil nematode abundance
differ among functional groups and land-use types at a global scale. *Journal of Animal Ecology*,
91, 1770–1780. https://doi.org/10.1111/1365-2656.13744

Lososová Z, Horsák M, Chytrý M, Čejka T, Danihelka J, Fajmon K, Hájek O, Juřičková L, Kintrová K,
Láníková D, Otýpková Z, Řehořek V, Tichý L (2011) Diversity of Central European urban biota:
effects of human-made habitat types on plants and land snails. *Journal of Biogeography*, **38**,
1152–1163. https://doi.org/10.1111/j.1365-2699.2011.02475.x

Manoli G, Fatichi S, Schläpfer M, Yu K, Crowther TW, Meili N, Burlando P, Katul GG, Bou-Zeid E
(2019) Magnitude of urban heat islands largely explained by climate and population. *Nature*, **573**,
55–60. https://doi.org/10.1038/s41586-019-1512-9

Martin E, El-Galmady S, Johnson MTJ (2024) Urban socioeconomic variation influences the
ecology and evolution of trophic interactions. *Ecology Letters*, **27**, e14407.
https://doi.org/10.1111/ele.14407

McKinney ML (2008) Effects of urbanization on species richness: a review of plants and animals.
Urban Ecosystems, **11**, 161–176. https://doi.org/10.1007/s11252-007-0045-4

Minias P (2023) The effects of urban life on animal immunity: Adaptations and constraints. *Science*
of The Total Environment, **895**, 165085. https://doi.org/10.1016/j.scitotenv.2023.165085

Morand S, Wilson MJ, Glen DM (2004) Nematodes (Nematoda) parasitic in terrestrial molluscs.
In: *Natural enemies of terrestrial molluscs* (ed Barker GM), pp. 525–557. CABI, Wallingford, UK.

Moreno-Monroy AI, Schiavina M, Veneri P (2021) Metropolitan areas in the world. Delineation
and population trends. *Journal of Urban Economics*, **125**, 103242.
https://doi.org/10.1016/j.jue.2020.103242

Murray MH, Sánchez CA, Becker DJ, Byers KA, Worsley-Tonks KE, Craft ME (2019) City sicker? A
meta-analysis of wildlife health and urbanization. *Frontiers in Ecology and the Environment*, **17**,
575–583. https://doi.org/10.1002/fee.2126

Nakagawa S, Johnson PCD, Schielzeth H (2017) The coefficient of determination R^2 and intra-class
correlation coefficient from generalized linear mixed-effects models revisited and expanded.
Journal of The Royal Society Interface, **14**, 20170213. https://doi.org/10.1098/rsif.2017.0213

Nakagawa S, Schielzeth H (2013) A general and simple method for obtaining R^2 from generalized
linear mixed-effects models. *Methods in Ecology and Evolution*, **4**, 133–142.
https://doi.org/10.1111/j.2041-210x.2012.00261.x

Oosterhoff LM (1977) Variation in growth rate as an ecological factor in the landsnail *Cepaea*
nemoralis (L.). *Netherlands Journal of Zoology*, **27**, 1–132.
https://doi.org/10.1163/002829677X00072

Ożgo M (2009) Current problems in the research of *Cepaea* polymorphism. *Folia Malacologica*,
16, 55–60. https://doi.org/10.12657/folmal.016.009

Parris KM (2016) *Ecology of urban environments*. Wiley-Blackwell, Chichester, UK.

Pebesma E (2018) Simple features for R: standardized support for spatial vector data. *The R*
Journal, **10**, 439. https://doi.org/10.32614/RJ-2018-009

Pesaresi M, Politis P (2023) GHS-BUILT-S R2023A - GHS built-up surface grid, derived from
Sentinel2 composite and Landsat, multitemporal (1975-2030).
https://doi.org/10.2905/9F06F36F-4B11-47EC-ABB0-4F8B7B1D72EA

Piano E, Souffreau C, Merckx T, Baardsen LF, Backeljau T, Bonte D, Brans KI, Cours M, Dahirel M,
Debortoli N, Decaestecker E, Wolf KD, Engelen JMT, Fontaneto D, Gianuca AT, Govaert L,
Hanashiro FTT, Higuti J, Lens L, Martens K, Matheve H, Matthysen E, Pinseel E, Sablon R, Schön I,
Stoks R, Doninck KV, Dyck HV, Vanormelingen P, Wichelen JV, Vyverman W, Meester LD,
Hendrickx F (2020) Urbanization drives cross-taxon declines in abundance and diversity at
multiple spatial scales. *Global Change Biology*, **26**, 1196–1211.
https://doi.org/10.1111/gcb.14934

Pieterse A, Malan AP, Ross JL (2017) Nematodes that associate with terrestrial molluscs as
definitive hosts, including *Phasmarhabditis hermaphrodita* (Rhabditida: Rhabditidae) and its
development as a biological molluscicide. *Journal of Helminthology*, **91**, 517–527.
https://doi.org/10.1017/S0022149X16000572

Pollard E, Cooke AS, Welch JM (1977) The use of shell features in age determination of juvenile
and adult Roman snails *Helix pomatia*. *Journal of Zoology*, **183**, 269–279.
https://doi.org/10.1111/j.1469-7998.1977.tb04186.x

R Core Team (2023) R: a language and environment for statistical computing.

Radwan MA, El-Gendy KS, Gad AF (2020) Biomarker responses in terrestrial gastropods exposed
to pollutants: A comprehensive review. *Chemosphere*, **257**, 127218.
https://doi.org/10.1016/j.chemosphere.2020.127218

Rae R (2017) The gastropod shell has been co-opted to kill parasitic nematodes. *Scientific Reports*,
7, 4745. https://doi.org/10.1038/s41598-017-04695-5

Rae R (2018) Shell encapsulation of parasitic nematodes by *Arianta arbustorum* (Linnaeus, 1758)
in the laboratory and in field collections. *Journal of Molluscan Studies*, **84**, 92–95.
https://doi.org/10.1093/mollus/eyx045

Rae R (2023) Avoidance and attraction behaviour of slugs exposed to parasitic nematodes. *Journal*
of Invertebrate Pathology, **197**, 107896. https://doi.org/10.1016/j.jip.2023.107896

Rae RG, Robertson JF, Wilson MJ (2008) Susceptibility and immune response of *Deroceras*
reticulatum, *Milax gagates* and *Limax pseudoflavus* exposed to the slug parasitic nematode
Phasmarhabditis hermaphrodita. *Journal of Invertebrate Pathology*, **97**, 61–69.
https://doi.org/10.1016/j.jip.2007.07.004

Rosin ZM, Kwieciński Z, Lesicki A, Skórka P, Kobak J, Szymańska A, Osiejuk TS, Kałuski T, Jaskulska
M, Tryjanowski P (2018) Shell colour, temperature, (micro)habitat structure and predator
pressure affect the behaviour of *Cepaea nemoralis*. *The Science of Nature*, **105**, 35.
https://doi.org/10.1007/s00114-018-1560-2

Russo J, Madec L (2011) Dual strategy for immune defense in the land snail *Cornu aspersum*
(Gastropoda, Pulmonata). *Physiological and Biochemical Zoology: Ecological and Evolutionary*
Approaches, **84**, 212–221. https://doi.org/10.1086/659123

Russo J, Madec L (2013) Linking immune patterns and life history shows two distinct defense
strategies in land snails (Gastropoda, Pulmonata). *Physiological and Biochemical Zoology*, **86**,
193–204. https://doi.org/10.1086/669482

Scheil AE, Hilsmann S, Triebskorn R, Köhler H-R (2013) Shell colour polymorphism, injuries and
immune defense in three helicid snail species, *Cepaea hortensis*, *Theba pisana* and *Cornu*
aspersum maximum. *Results in Immunology*, **3**, 73–78.
https://doi.org/10.1016/j.rinim.2013.06.002

Scheil AE, Hilsmann S, Triebskorn R, Köhler H-R (2014) Shell colouration and parasite tolerance in
two helicoid snail species. *Journal of Invertebrate Pathology*, **117**, 1–8.
https://doi.org/10.1016/j.jip.2014.01.003

Schiavina M, Freire S, Carioli A, MacManus K (2023) GHS-POP R2023A - GHS population grid
multitemporal (1975-2030). https://doi.org/10.2905/2FF68A52-5B5B-4A22-8F40-
C41DA8332CFE

Schiavina M, Melchiorri M, Pesaresi M (2023) GHS-SMOD R2023A - GHS settlement layers,
application of the Degree of Urbanisation methodology (stage I) to GHS-POP R2023A and GHS-
BUILT-S R2023A, multitemporal (1975-2030). https://doi.org/10.2905/A0DF7A6F-49DE-46EA-
9BDE-563437A6E2BA

Schiavina M, Moreno-Monroy A, Maffenini L, Veneri P (2019) GHS-FUA R2019A - GHS functional
urban areas, derived from GHS-UCDB R2019A (2015). https://doi.org/10.2905/347F0337-F2DA-
4592-87B3-E25975EC2C95

Segade P, García-Estévez J, Arias C, Iglesias R (2013) Parasitic infections in mixed system-based
heliciculture farms: dynamics and key epidemiological factors. *Parasitology*, **140**, 482–497.
https://doi.org/10.1017/S0031182012001795

Shultz AJ, Adams BJ, Bell KC, Ludt WB, Pauly GB, Vendetti JE (2020) Natural history collections are
critical resources for contemporary and future studies of urban evolution. *Evolutionary*
Applications, **14**, 233–247. https://doi.org/10.1111/eva.13045

Szulkin M, Munshi-South J, Charmantier A (Eds.) (2020) *Urban Evolutionary Biology*. Oxford University Press. https://doi.org/10.1093/oso/9780198836841.001.0001

Theodorou P (2022) The effects of urbanisation on ecological interactions. *Current Opinion in Insect Science*, **52**, 100922. https://doi.org/10.1016/j.cois.2022.100922

Valdés-Correcher E, Popova A, Galmán A, Prinzing A, Selikhovkin AV, Howe AG, Mrazova A, Dulaurent A-M, Hampe A, Tack AJM, Bouget C, Lupaștean D, Harvey D, Musolin DL, Lövei GL, Centenaro G, Halder IV, Hagge J, Dobrosavljević J, Pitkänen J-M, Koricheva J, Sam K, Barbaro L, Branco M, Ferrante M, Faticov M, Tahadlová M, Gossner M, Cauchoix M, Bogdziewicz M, Duduman M-L, Kozlov MV, Bjoern MC, Mamaev NA, Fernandez-Conradi P, Thomas RL, Wetherbee R, Green S, Milanović S, Moreira X, Mellerin Y, Kadiri Y, Castagneyrol B (2022) Herbivory on the pedunculate oak along an urbanization gradient in Europe: Effects of impervious surface, local tree cover, and insect feeding guild. *Ecology and Evolution*, **12**, e8709. https://doi.org/10.1002/ece3.8709

Wickham H, Averick M, Bryan J, Chang W, McGowan L, François R, Grolemund G, Hayes A, Henry L, Hester J, Kuhn M, Pedersen T, Miller E, Bache S, Müller K, Ooms J, Robinson D, Seidel D, Spinu V, Takahashi K, Vaughan D, Wilke C, Woo K, Yutani H (2019) Welcome to the Tidyverse. *Journal of Open Source Software*, **4**, 1686. https://doi.org/10.21105/joss.01686

Williamson P (1979) Age determination of juvenile and adult *Cepaea*. *Journal of Molluscan Studies*, **45**, 52–60.

Williams AJ, Rae R (2015) Susceptibility of the Giant African snail (*Achatina fulica*) exposed to the gastropod parasitic nematode *Phasmarhabditis hermaphrodita*. *Journal of Invertebrate Pathology*, **127**, 122–126. https://doi.org/10.1016/j.jip.2015.03.012

Williams A, Rae R (2016) *Cepaea nemoralis* (Linnaeus, 1758) uses its shell as a defence mechanism to trap and kill parasitic nematodes. *Journal of Molluscan Studies*, **82**, 349–350. https://doi.org/10.1093/mollus/eyv064

Wilson MJ, Hughes LA, Hamacher GM, Glen DM (2000) Effects of *Phasmarhabditis hermaphrodita* on non-target molluscs. *Pest Manag Sci*, **56**, 711–716. https://doi.org/10.1002/1526-4998(200008)56:8%3C711::AID-PS185%3E3.0.CO;2-O

Wilson MJ, Hughes LA, Jefferies D, Glen DM (1999) Slugs (*Deroceras reticulatum* and *Arion ater* agg.) avoid soil treated with the rhabditid nematode *Phasmarhabditis hermaphrodita*. *Biological Control*, **16**, 170–176. https://doi.org/10.1006/bcon.1999.0757

Wolda H (1967) The effect of temperature on reproduction in some morphs of the landsnail *Cepaea nemoralis* (L.). *Evolution*, **21**, 117–129. https://doi.org/10.1111/j.1558-5646.1967.tb00135.x

Wynne R, Morris A, Rae R (2016) Behavioural avoidance by slugs and snails of the parasitic nematode *Phasmarhabditis hermaphrodita*. *Biocontrol Science and Technology*, **26**, 1129–1138. https://doi.org/10.1080/09583157.2016.1185513

607   Żbikowska E, Marszewska A, Cichy A, Templin J, Smorąg A, Strzała T (2020) *Cepaea* spp. as a source
608   of *Brachylaima mesostoma* (Digenea: Brachylaimidae) and *Brachylecithum* sp. (Digenea:
609   Dicrocoeliidae) larvae in Poland. *Parasitology Research*, **119**, 145–152.
610   https://doi.org/10.1007/s00436-019-06516-2