Urban *Cepaea nemoralis* snails are less likely to have nematodes trapped within their shells

Maxime Dahirel*, Hannah Reyné, Katrien De Wolf, Dries Bonte

Ghent University, Department of Biology, B-9000 Gent

ORCID:

MD: 0000-0001-8077-7765

KDW: 0000-0002-9255-913X

DB: 0000-0002-3320-7505

*corresponding author: maxime.dahirel@yahoo.fr

Abstract

Urbanisation is a major human-induced environmental change which can impact not only individual species, but also the way these species interact with each other. As a group, terrestrial molluscs interact frequently with a wide diversity of parasites, yet the way these interactions vary across space and in response to environmental pressures is poorly documented. In this study we leveraged a recently discovered defence mechanism, by which snails trap parasitic nematodes in their shells, to explore how snail-nematodes interactions may vary in response to city life. We examined shells from the generalist snail *Cepaea nemoralis* sampled in three urban areas in Belgium for trapped nematodes, and attempted to link this to urbanisation and shell phenotypic traits. We found that even a small degree of urbanisation led to large decreases in the rates of shell encapsulation, and that larger snails were more likely to contain trapped nematodes. However, we found no evidence that shell colour, which had been previously linked to immune function, was correlated to encapsulation rates. We discuss how between-population variation in encapsulation rates can result from urbanisation-induced changes on the nematodes side, the snail side, or both, and suggest potential tests for future studies aiming to disentangle these mechanisms.

Key words: Biotic interactions, Gastropoda, immunity, parasites.

Introduction

Urbanisation is a major and all-encompassing human-induced environmental change, leading to changes in land use, local climate, soil imperviousness, light and chemical pollution… (Parris, 2016). The (often negative) impacts of these multivariate changes on biodiversity are increasingly well-documented: many species decline in cities, while some become successful "urban adapters", leading to major restructuring of biological communities along urbanisation gradients (e.g. McKinney, 2008; Piano et al., 2020; Fenoglio et al., 2020; Liang et al., 2023). In parallel, urbanisation can also cause within-species phenotypic and genetic changes (Alberti et al., 2017; Szulkin et al., 2020; Diamond & Martin, 2021). Urbanisation may also have second-order impacts by reshaping ecological interactions, if tightly connected species respond to environmental change in different ways (Theodorou, 2022). Such urbanisation-induced changes in ecological interactions, in both positive and negative directions, have been recorded for plant-pollinator interactions (Liang et al., 2023), plant-herbivore and prey-predator interactions (Eötvös et al., 2018; Valdés-Correcher et al., 2022; Gámez et al., 2022; Korányi et al., 2022), as well as host-parasite interactions (Murray et al., 2019; Korányi et al., 2022).

Terrestrial molluscs (snails and slugs) are potentially valuable models in urban ecology and evolution, in part because of their limited movement abilities, which means they often cannot move to escape environmental changes. Like in many other taxa, urbanisation can reshape molluscan communities (Lososová et al., 2011; Horsák et al., 2013; Barbato et al., 2017; Hodges & McKinney, 2018), and drive evolutionary responses in urban populations (Kerstes et al., 2019). Interestingly, in a comparative cross-taxon study or urbanisation impacts, snail species richness seemed less negatively affected, compared to other more mobile groups (Piano et al., 2020). This may be because their locally high abundances and small home ranges allow their persistence in even small habitat fragments. Land molluscs are hosts to a diverse array of metazoan parasites, including nematodes, flies, mites or trematodes (Barker, 2004; Segade et al., 2013; Żbikowska et al., 2020). How urbanisation reshapes these interactions remains understudied, despite some of these parasites being of increasing veterinary interest (Aziz et al., 2016; Giannelli et al., 2016).

Snails and slugs can defend themselves against metazoan parasites through a variety of behavioural (Wilson et al., 1999; Wynne et al., 2016; Rae, 2023) or immune responses (Furuta & Yamaguchi, 2001; Scheil et al., 2014; Coaglio et al., 2018). Among the latter, it has been discovered that land molluscs can use their shells to trap parasitic nematodes, killing them and fusing them to the inner shell surface (Rae et al., 2008; Williams & Rae, 2015; Rae, 2017). This ability seems phylogenetically widespread, even present in slugs with vestigial shells (Rae et al., 2008; Rae, 2017), and could therefore provide a relatively easy to access record of ecological interactions. Following anecdotal records of mites and trematodes encapsulated in shells, it has further been suggested that this shell encapsulation might extend to other metazoan parasites (Dahirel et al., 2022; Gérard et al., 2023). However, given how rare these non-nematode records are, they may be merely by-products of a defence mechanism targeted towards nematodes, rather than evidence of a more generalized defence response (Gérard et al., 2023). The few snail species in which this phenomenon has been studied across multiple populations show that the prevalence of individuals trapping nematodes can vary widely between sites (Rae, 2017;

Rae, 2018; Cowlishaw et al., 2019), but there has been no attempt, to our knowledge, to assess
whether this variation could be non-random with respect to environmental context.

To that end, we combine here publicly available and standardized urbanisation metrics with
observations of field-collected snails across three cities in Belgium, using the grove snail *Cepaea*
nemoralis (Gastropoda, family Helicidae) as a model. Like other helicids, *C. nemoralis* can
encapsulate and trap parasitic nematodes in its shell (Williams & Rae, 2016; Rae, 2017; Dahirel
et al., 2022; Gérard et al., 2023). This snail is also common both outside and within cities
(Kerstes et al., 2019), and therefore a very suitable model to study variation in encapsulation
rates, whether it is due to urbanisation or to spatial (between-cities) differences. Furthermore,
the shell colour variation that made *Cepaea* species iconic models in evolutionary biology (Jones
et al., 1977; Ożgo, 2009) may also influence their immune response, with some evidence that
darker morphs mount better defences against nematodes (Dahirel et al., 2022; but see Scheil et
al., 2014). On the other hand, this morph variation in resistance might not translate to shell
encapsulation (Williams & Rae, 2016; Dahirel et al., 2022). However, existing comparisons were
either limited to one type of colour variation (banding pattern only, Dahirel et al., 2022), or
analysed experimental infections by one model nematode (Williams & Rae, 2016); we here test
whether this remains true when analysing naturally occurring snail-nematode interactions and
accounting for more dimensions of shell colour variation.

Methods

Site selection and sampling

We searched for *Cepaea nemoralis* snails from early October to mid-November 2022 in and
around the urban areas of Brussels, Ghent and Leuven in Belgium (**Fig. 1**). Potential sites were
selected based on pre-existing online crowdsourced records (iNaturalist contributors &
iNaturalist, 2024) combined with personal observations and virtual fieldwork using Google
StreetView to identify suitable habitats (based on Falkner et al., 2001). We visited 36 sites
chosen to be roughly balanced between the three cities (including their surrounding areas;
Brussels: 13 sites, Ghent: 13 sites, Leuven: 10 sites). In each site, we sampled living snails by
hand during visual search, in a radius of up to 50 m around a designated site centroid (though
search was *de facto* mostly concentrated within a 20 m radius). Individuals were mainly
searched in known favourable micro-habitats, i.e. on tall herbs and shrubs, under piled wood
and cardboard or loose rocks, or on fences, walls, and tree trunks (Falkner et al., 2001). Field
identification of *Cepaea nemoralis* snails is easy based on shell shape, size and colour (Cameron,
2008). We only collected adults, which can easily be separated from subadults by the presence
of a reflected shell lip marking the end of shell growth (Cameron, 2008). Each site was visited by
1 to 3 people (mean: 2.03) for a duration of 5 to 30 person-minutes (mean: 15). We collected a
total of 298 snails from 28 of the 36 sites visited (Brussels: 9 sites, Ghent: 10 sites, Leuven: 9
sites). However, 2 shells were lost before examination for parasites due to handling errors, and
another shell was accidentally broken for parasite examination before photographs or size
measurements could be done; this led to a final dataset of 295 snails still in 28 sites. For each of
these 28 sites, the nearest neighbouring site with snails found was between 153 and 1516 m
away (mean: 768 m), which is in any case farther than the maximal dispersal distances

(Kramarenko, 2014), indicating that even nearby sites could be considered separate
populations.

Urbanisation metrics

It is well-known that urban environmental changes are complex and multivariate (e.g. Parris,
2016); however, given our relatively low number of sites, and the risk of collinearity between
urban metrics, we decided to use simple overarching metrics focused on building presence and
human population density. We assessed urbanisation at each site using raster layers from the
Global Human Settlement Layer project for the year 2020 (https://ghsl.jrc.ec.europa.eu/, Joint
Research Centre (European Commission), 2023). We first used built-up surface (GHS-BUILT-S)
and population density (GHS-POP) at 100 m and 1000 m resolutions (Pesaresi & Politis, 2023;
Schiavina, Freire, et al., 2023). The former spatial scale matches the scale of maximal dispersal
movements over timespans of up to a couple years in helicid snails, while the latter is closer to
the scale of longer term (over several decades) population spread (Kramarenko, 2014). As an
additional categorical metric, we also used the Degree of Urbanisation as recorded in the
Settlement Model layer (GHS-SMOD, available only at 1000 m resolution, Eurostat (European
Commission), 2021; Schiavina, Melchiorri, et al., 2023). At the highest level of classification, the
standardized Degree of Urbanisation methodology mainly uses population density and
contiguity rules to classify grid cells as either part of a continuous high-density Urban Centre, as
low-density rural cells or as intermediate peri-urban/suburban cells. For each site and
urbanisation metric, we recorded the value of the corresponding grid cell. Interestingly, Degree
of Urbanisation classes, which are primarily based on population density, divide our sites in
almost the same non-linear way as another, independent, three-level classification based on

built-up surfaces used in previous urban ecology studies in the study region (e.g. Piano et al.,
2020) (**Supplementary Material S1**).

Snail shell analysis

Snail size was measured using a caliper as the shell greatest diameter (to the nearest 0.1 mm).
Snail shell colour morphs were scored following e.g. Cain (1988) for background colour (from
lighter to darker: yellow, pink or brown), number of dark bands (0 to 5 bands) and on the
presence or absence of band fusions (which increase the proportion of the shell covered by dark
bands). Snails were killed by first inducing dormancy at 6°C, then by freezing at -20°C. We
removed bodies from shells with forceps and lightly cleaned shells with water (bodies were
stored in ethanol for separately planned studies). We then broke each shell into fragments using
forceps, examined fragments under a binocular microscope, and recorded all animals found
encapsulated within the shell as in e.g. Gérard et al. (2023). A total of 606 nematodes were
found in 104 shells; no other metazoan parasites were recorded. As this method is destructive,
we took standardised photographs of the shells beforehand (dorsal and apertural views
following Callomon, 2019) for archival and potential future studies.

Statistical analysis

All analyses were done in R version 4.3.2 (R Core Team, 2023), with the help of the *tidyverse*
(Wickham et al., 2019) and *sf* (Pebesma, 2018) packages for data processing, as well as
additional packages detailed below for model fitting and exploration.

We analysed the probability a shell contained nematodes as a binary yes/no response at the
individual level, using Generalized Linear Mixed Models (GLMMs) (binomial family, logit link).
We ran six models; the first five all included shell size, shell morph traits (background colour,
band number and fusion), urbanisation and city identity (Brussels, Ghent or Leuven) as fixed
effects, only differing by which urbanisation metric they used (among the five described above
in **Urbanisation metrics**). Numeric predictors were centred and scaled to unit 1 SD. Sampling
site was included as a random intercept. The sixth model was a "null" model, identical to the
other ones except that it did not include an urbanisation metric. We ran our models using the
glmmTMB package (Brooks et al., 2017), and then used AICc to compare them. As one model
largely outperformed the others (see **Results**), we did all further analyses on that best model.

We checked for residual spatial autocorrelation using a spline correlogram (*ncf* package,
Bjornstad, 2022), and found no evidence of spatial structure in the best model. We then used
the *car* (Fox & Weisberg, 2019) and *emmeans* (Lenth, 2023) packages to test for overall effects
of our variables in the best model and to run (Tukey-corrected) pairwise comparisons,
respectively. Finally, we estimated the marginal and conditional R^2 (Nakagawa & Schielzeth,
2013) as measures of the proportion of variation explained by fixed effects (R_m^2) and both fixed
and random effects (R_c^2) respectively (using the delta method, Nakagawa et al., 2017).

Results

The model using the categorical Degree of Urbanisation (GHS-SMOD) as an urbanisation metric
outperformed all other models based on AICc (**Table 1**). Fixed effects and random effects

explained similar amounts of variance ($R_m^2 = 0.19$; $R_c^2 = 0.38$). The probability that a shell had
trapped nematodes was dependent on urbanisation level ($\chi^2 = 15.97$, df $= 2$, $p = 3.40 \times$
10^{-4}) but did not vary significantly between cities ($\chi^2 = 3.89$, df $= 2$, $p = 0.14$). Snails from
rural sites were more likely to contain nematodes than snails from intermediate and Urban
Centre populations (**Fig. 2**; rural - intermediate difference on the logit scale $\pm$ SE: 3.71 ± 0.95 ;
rural - Urban Centre difference: 2.34 ± 0.77). Larger shells were more likely to contain
nematodes ($\chi^2 = 4.17$, df $= 1$, $p = 0.04$; standardised coefficient $\beta = 0.35 \pm 0.17$). There was
no clear evidence that any of the shell colour traits affected encapsulation rates (background
colour: $\chi^2 = 2.17$, df $= 2$, $p = 0.34$; band number: $\chi^2 = 1.90$, df $= 1$, $p = 0.17$; fusion:
$\chi^2 = 0.17$, df $= 1$, $p = 0.68$).

Table 1. Model selection table for the effect of urbanisation on shell encapsulation rates. All models otherwise include effects of city identity, shell size and shell morph (background colour, number of bands and band fusion).

Urbanisation variable in model	df	log-likelihood	AICc		AICc weight
Degree of Urbanisation categories (SMOD, 1000 m resolution grid)	11	-149.5	322.0	0.00	0.95
None ("null" model)	9	-155.9	330.4	8.43	0.01
Population density (1000 m resolution grid)	10	-155.2	331.1	9.11	0.01
Built-up surface (1000 m resolution grid)	10	-155.4	331.5	9.49	0.01
Population density (100 m resolution grid)	10	-155.4	331.6	9.61	0.01
Built-up surface (100 m resolution grid)	10	-155.6	331.9	9.91	0.01

Figure 2. Effect of the Degree of Urbanisation in 1000 m grid cells on the probability a snail shell contained encapsulated nematodes. Grey dots are observed proportions per population, with the size of the dot proportional to the number of snails; white dots (and error bars) are estimated marginal means from the best model (and their 95% confidence intervals), with the effects of the other predictors averaged out.

Discussion

We found that population differences in the prevalence of *Cepaea nemoralis* snails encapsulating nematodes in their shell as a defence mechanism were partly driven by environmental conditions, with trapping rates decreasing with increasing urbanisation (**Fig. 2**). This effect was better explained by a categorical classification of the Degree of Urbanisation, rather than by linear effects of continuous urbanisation variables. This indicates that the response to urbanisation is here non-linear, as the increases in population density/built-up rates needed to go from a rural to an intermediate area are much smaller than those needed to go from intermediate to Urban Centre, and most of the variation in density/built-up is within Urban Centres rather than between categories (**Supplementary Material S1**).

A difficulty for interpreting our results is that nematodes trapped in shells will accumulate with time (Williams & Rae, 2015; Rae, 2017), meaning that as they may have endured more infections, older snails may be more likely to have them. If urbanisation reduces snail survival rates, a plausible proposal because snail survival is strongly temperature-dependent (e.g. Wolda, 1967) and cities are generally hotter environments (Manoli et al., 2019), then our urban-rural differences in nematodes trapped might merely reflect differences in average snail age/survival. Age estimation in shelled gastropods, especially in terrestrial snails with determinate growth like *Cepaea*, is notoriously challenging. There are documented proxies of time to maturity, such as the number of dormancy-induced growth stops visible on the shell, or of adult age, such as shell wear or shell layers in cross-sections (Pollard et al., 1977; Williamson, 1979). However, they are imprecise, and we may expect them to vary in response to urbanisation, as spatial variation in local climate can lead for instance to spatial variation in dormancy regimes (Staikou et al., 2024), and so potentially in the relationship between age and growth stops. This could make these proxies useless to compare age between populations, as would be needed here, unless thorough calibration studies are undertaken. We note though, reassuringly, that the number of nematodes found in infected shells does not seem to decrease in more urbanised sites (**Supplementary Material S2**). This is contrary to what we would expect if variation in shell encapsulation was primarily explained by variation in age/time available to accumulate nematodes and urban snails were on average younger.

If we assume that our results reflect differences in snail-nematodes interactions between urban and non-urban areas, several mutually non-exclusive mechanisms may explain why urban *Cepaea nemoralis* shells are less likely to trap nematodes. Each of these mechanisms directly suggests potential tests for future studies:

- First, snail parasitic nematodes infecting *C. nemoralis* may be less abundant in cities. Many of the nematodes known to infect land snails have at least one free-living life stage in the soil, and some are facultative parasites (Morand et al., 2004; Pieterse et al., 2017). Increasingly impervious substrates in cities (Parris, 2016) may deprive these of habitat critical for their life cycle. Where habitat is available, soil nematode communities are profoundly altered by urbanisation, like other taxa (Li et al., 2022; Gong et al., 2024). However, this does not lead to overarching declines in nematode abundance; rather, some trophic groups decline while others thrive (Li et al., 2022; Gong et al., 2024).

Unfortunately, detailed information on nematodes parasitizing animals is typically
lacking from these analyses; soil sampling specifically targeting parasitic nematodes
(Jaffuel et al., 2019) would here be particularly useful.

• Second, individual differences in behaviour, especially space-related behaviour, may lead
to differences in the risk of encountering and then being infected by parasites (Barber &
Dingemanse, 2010). In chipmunks (*Tamias sibiricus*), some individuals are consistently
more explorative than others, which affects their risk of encountering ticks (Boyer et al.,
2010). A similar result was also found in firebugs (*Pyrrhocoris apterus*) (Gyuris et al.,
2016). Habitat loss and fragmentation associated with urbanisation are expected to
exert strong selection pressures on movement and space use (Cote et al., 2017). If this
results in lower movement in urban snail populations, this might then reduce their
encounter rates with parasites. In the snail *Cornu aspersum*, another related species
commonly found in cities, urbanisation does not seem to lead to reduced boundary-
crossing behaviour in that species however (Dahirel et al., 2016), although that is only
one component of mobility. Urbanisation-induced increases in temperature may also
alter the frequency at which snails hide into shelters or climb above the substrate (Rosin
et al., 2018), and potentially again the risk of encountering parasites. The picture is
however complicated by behaviour-parasite feedbacks, where while host behaviour
shapes infection risk, infection can then alter host behaviour in turn (Ezenwa et al.,
2016). In *Cepaea nemoralis*, nematode infection itself might lead to reduced movement
propensity, but only in some morphs (Dahirel et al., 2022). More studies of movement
behaviour across urbanisation gradients are here needed.

• Third, shell encapsulation rates are not direct records of snail-nematode interactions,
but rather informative on the host's ability to mount a defence in such interactions. This
defence is not always effective, as demonstrated by cases where field-caught snails
showed active infections but zero shell-trapped nematodes (see e.g. data in Dahirel et
al., 2022). If immune response in general and the pathways involved with shell
encapsulation specifically decline with urbanisation, then this alone could explain our
results even in the absence of changes in nematode communities. In vertebrates,
available evidence suggests that urban living can lead to both depressed or stimulated
immune function, depending on taxon and context, especially food availability (Murray
et al., 2019; Minias, 2023). In terrestrial molluscs, chemical pollutants seem to negatively
impact many, but not all, physiological components of immune defence (Radwan et al.,
2020). In the freshwater snail *Lymnaea stagnalis*, higher temperatures led to declines in
immune function (Leicht et al., 2017). The exact physiological pathways involved in shell
encapsulation in land molluscs remain however unstudied, to the best of our knowledge.

On the phenotype side, larger shells were more likely to contain trapped nematodes. This could
be the result of survivor bias alone, if larger snails are more likely to survive infection. Although
we cannot exclude that other nematodes have larger effects, experimental nematode infections
by *Phasmarhabditis* are almost never lethal in adult *Cepaea nemoralis*, contrary to other snail
species (Wilson et al., 2000; Williams & Rae, 2016). Other potential explanations for this result

can be sorted along three non-exclusive lines, similar to the mechanisms suggested above to
explain the effect of urbanisation:

• Larger snails might harbour larger parasite infections (e.g. Daniels et al., 2013), which
would increase the likelihood that some nematodes are trapped. However, there is no
link between nematode abundance in active infections and snail size in *C. nemoralis*
(Dahirel et al., 2022). Furthermore, although our sample size is small, we find no clear
effect of body size on the number of nematodes trapped in the present study
(**Supplementary Material S2**).

• If large and small snails differ in their space use, they might also differ in their parasite
exposure risk. On the one hand, while only one dimension of space use was tested, there
was no link between short-term movement and size in adult *C. nemoralis* (Dahirel et al.,
2022). On the other hand, there is a positive correlation between adult size and
emigration propensity (Oosterhoff, 1977).

• Finally, small and large snails may differ in their immune defence abilities. Comparative
studies suggest that large and small snail species and subspecies differ in their immune
strategies at the physiological level (Russo & Madec, 2011, 2013). However, the range of
body size and life history variation is much larger in these scenarios than among adults
of *C. nemoralis*, limiting the transferability of these results. Larger adults might also be
larger because they were able to access more or better resources during development,
which may make them able to defend against parasites more easily. However, the link
between resource quality or access and adult size (as opposed to growth rate or adult
mass) is actually ambiguous in helicid snails (Wacker & Baur, 2004; Nicolai et al., 2012).
More physiological studies focused on within-, rather than among-species variation may
help understand better this link between body size and encapsulation rates.

In contrast to shell size, we found no relationship between any of the shell colour traits and
nematode trapping rate. This confirms experimental results from Williams & Rae (2016) using
infections by *Phasmarhabditis elegans*. However, colour morphs do differ in active infection
rates or other aspects of immune response in *C. nemoralis* (Dahirel et al., 2022) and other
polymorphic snails (Scheil et al., 2013, 2014). This discrepancy may indicate that shell
encapsulation is driven by different physiological pathways than other components of snail
immune defence.

Beyond the effects of phenotype or environment, how to relate the prevalence of nematodes
trapped in shells with rates of active parasite infections remains an open and complex question.
We were not able to compare the two here as snail bodies were reserved for other
investigations. Nematodes trapped in shells reflect the cumulative history of infection, which
may or may not be meaningfully correlated with active infection at a given time. For instance, if
variation in snail-nematode interactions is driven by e.g. variation in nematode density in the
environment, we may expect a positive correlation, as higher nematode densities should drive
up rates of both shell encapsulation (Rae, 2018) and active infection (although if encapsulation
is highly effective, it may end up suppressing dose-dependent effects on active infection,
Williams & Rae, 2015). On the other hand, if variation is mostly driven by snail immune

response, we may expect a negative correlation instead: nematode density being equal, snails with more effective immune systems may be more likely to successfully trap nematodes in shells while being less likely to harbour active infections. Additionally, we do not know much about whether some nematode species are more likely than others to end up trapped in shells (identification by sequencing of encapsulated nematodes may be promising here, Cowlishaw et al., 2019). While this would need to be validated, the strength and direction of between- and within-sites correlation between active infections and shell-trapped nematodes may provide useful indicators of the main drivers of snail-nematodes interactions in response to city life.

We acknowledge that the relatively small size of our sample does not allow us to draw firm causal conclusions. Nonetheless, we hope our results may encourage larger studies regarding host-parasite interactions in land molluscs in the context of environmental change. As new technical developments such as micro-CT imaging allow non-destructive analyses of snail shells (Falkingham & Rae, 2021), these may extend to using museum and other natural history collections to understand how interactions vary in space and time (Cowlishaw et al., 2019), reaffirming their value for urban ecology and evolution (Shultz et al., 2020).

Author contributions

Initial study idea: MD, DB. Site selection and fieldwork: MD, HR, KDW. Shell data collection: HR, after initial training by MD. Data analysis: MD, after preliminary analyses by HR. Initial manuscript draft: MD. All authors contributed critically to edits and gave final approval for publication.

Funding

This study has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 101022802 (HELICITY). KDW and DB received funding from the Fund for Scientific Research Flanders (grant G080221N).

Conflict of interest disclosure

The authors declare they have no financial conflict of interest in relation with the content of this article. DB is a recommender for PCI Ecology and PCI Evolutionary Biology.

Data availability

Data and R scripts to reproduce all analyses presented in this article, as well as a copy of the Supplementary Materials, are available on Github (https://github.com/mdahirel/HELICITY-2022_shell-nematodes) and archived in Zenodo (DOI: https://doi.org/10.5281/zenodo.10794928).

References

Alberti M, Correa C, Marzluff JM, Hendry AP, Palkovacs EP, Gotanda KM, Hunt VM, Apgar TM, Zhou Y (2017) Global urban signatures of phenotypic change in animal and plant populations.

Proceedings of the National Academy of Sciences, 201606034.
https://doi.org/10.1073/pnas.1606034114

Aziz NAA, Daly E, Allen S, Rowson B, Greig C, Forman D, Morgan ER (2016) Distribution of
Angiostrongylus vasorum and its gastropod intermediate hosts along the rural–urban gradient
in two cities in the United Kingdom, using real time PCR. *Parasites & Vectors*, **9**, 56.
https://doi.org/10.1186/s13071-016-1338-3

Barbato D, Benocci A, Caruso T, Manganelli G (2017) The role of dispersal and local environment
in urban land snail assemblages: an example of three cities in Central Italy. *Urban Ecosystems*,
20, 919–931. https://doi.org/10.1007/s11252-017-0643-8

Barber I, Dingemanse NJ (2010) Parasitism and the evolutionary ecology of animal personality.
Philosophical Transactions of the Royal Society B: Biological Sciences, **365**, 4077–4088.
https://doi.org/10.1098/rstb.2010.0182

Barker GM (Ed.) (2004) *Natural enemies of terrestrial molluscs*. CABI, Wallingford, UK.

Bjornstad ON (2022) R package ncf: Spatial covariance functions.

Boyer N, Réale D, Marmet J, Pisanu B, Chapuis J-L (2010) Personality, space use and tick load in
an introduced population of Siberian chipmunks *Tamias sibiricus*. *Journal of Animal Ecology*, **79**,
538–547. https://doi.org/10.1111/j.1365-2656.2010.01659.x

Brooks M E., Kristensen K, Benthem K J.,van, Magnusson A, Berg C W., Nielsen A, Skaug H J.,
Mächler M, Bolker B M. (2017) GlmmTMB balances speed and flexibility among packages for
zero-inflated generalized linear mixed modeling. *The R Journal*, **9**, 378.
https://doi.org/10.32614/RJ-2017-066

Cain AJ (1988) The scoring of polymorphic colour and pattern variation and its genetic basis in
molluscan shells. *Malacologia*, **28**, 1–15.

Callomon P (2019) *Standard views for imaging mollusk shells*. American Malacological Society.

Cameron R (2008) *Land Snails in the British Isles*. FSC Publications, Telford, UK.

Coaglio AL, Ferreira MAND, dos Santos Lima W, de Jesus Pereira CA (2018) Identification of a
phenoloxidase- and melanin-dependent defence mechanism in *Achatina fulica* infected with
Angiostrongylus vasorum. *Parasites & Vectors*, **11**, 113. https://doi.org/10.1186/s13071-018-
2710-2

Cote J, Bestion E, Jacob S, Travis J, Legrand D, Baguette M (2017) Evolution of dispersal
strategies and dispersal syndromes in fragmented landscapes. *Ecography*, **40**, 56–73.
https://doi.org/10.1111/ecog.02538

Cowlishaw RM, Andrus P, Rae R (2019) An investigation into nematodes encapsulated in shells
of wild, farmed and museum specimens of *Cornu aspersum* and *Helix pomatia*. *Journal of*
Conchology, **43**, 385–392.

Dahirel M, Proux M, Gérard C, Ansart A (2022) Morph-dependent nematode infection and its
association with host movement in the land snail *Cepaea nemoralis* (Mollusca, Gastropoda).
Journal of Zoology, **318**, 181–192. https://doi.org/10.1111/jzo.13012

Dahirel M, Séguret A, Ansart A, Madec L (2016) Dispersal-related traits of the snail *Cornu*
aspersum along an urbanisation gradient: maintenance of mobility across life stages despite
high costs. *Urban Ecosystems*, **19**, 1847–1859. https://doi.org/10.1007/s11252-016-0564-y

Daniels RR, Beltran S, Poulin R, Lagrue C (2013) Do parasites adopt different strategies in
different intermediate hosts? Host size, not host species, influences *Coitocaecum parvum*
(Trematoda) life history strategy, size and egg production. *Parasitology*, **140**, 275–283.
https://doi.org/10.1017/S0031182012001564

Diamond SE, Martin RA (2021) Evolution in cities. *Annual Review of Ecology, Evolution, and*
Systematics, **52**, 519–540. https://doi.org/10.1146/annurev-ecolsys-012021-021402

Eötvös CB, Magura T, Lövei GL (2018) A meta-analysis indicates reduced predation pressure with
increasing urbanization. *Landscape and Urban Planning*, **180**, 54–59.
https://doi.org/10.1016/j.landurbplan.2018.08.010

Eurostat (European Commission) (2021) *Applying the degree of urbanisation: a methodological*
manual to define cities, towns and rural areas for international comparisons : 2021 edition.
Publications Office of the European Union, LU.

Ezenwa VO, Archie EA, Craft ME, Hawley DM, Martin LB, Moore J, White L (2016) Host
behaviour–parasite feedback: an essential link between animal behaviour and disease ecology.
Proceedings of the Royal Society B: Biological Sciences, **283**, 20153078.
https://doi.org/10.1098/rspb.2015.3078

Falkingham P, Rae R (2021) 3D morphology of nematode encapsulation in snail shells, revealed
by micro-CT imaging. *Scientific Reports*, **11**, 2523. https://doi.org/10.1038/s41598-021-82106-6

Falkner G, Obrdlik P, Castella E, Speight MCD (2001) *Shelled Gastropoda of western Europe*.
Friedrich Held Gesellschaft, München, Germany.

Fenoglio MS, Rossetti MR, Videla M (2020) Negative effects of urbanization on terrestrial
arthropod communities: a meta-analysis. *Global Ecology and Biogeography*, **29**, 1412–1429.
https://doi.org/10.1111/geb.13107

Florczyk A, Corbane C, Schiavina M, Pesaresi M, Maffenini L, Melchiorri M, Politis P, Sabo F,
Freire S, Ehrlich D, Kemper T, Tommasi P, Airaghi D, Zanchetta L (2019) GHS-UCDB R2019A - GHS
Urban Centre Database 2015, multitemporal and multidimensional attributes.

Fox J, Weisberg S (2019) *An R Companion to Applied Regression*. Sage, Thousand Oaks, CA.

Furuta E, Yamaguchi K (2001) Haemolymph: blood cell morphology and function. In: *The biology*
of terrestrial molluscs (ed Barker GM), pp. 289–306. CABI, Wallingford, UK.

Gámez S, Potts A, Mills KL, Allen AA, Holman A, Randon PM, Linson O, Harris NC (2022)
Downtown diet: a global meta-analysis of increased urbanization on the diets of vertebrate
predators. *Proceedings of the Royal Society B: Biological Sciences*, **289**, 20212487.
https://doi.org/10.1098/rspb.2021.2487

Gérard C, De Tombeur Y, Dahirel M, Ansart A (2023) Land snails can trap trematode cercariae in
their shell: Encapsulation as a general response against parasites? *Parasite*, **30**, 1.
https://doi.org/10.1051/parasite/2023001

Giannelli A, Cantacessi C, Colella V, Dantas-Torres F, Otranto D (2016) Gastropod-borne
helminths: a look at the snail–parasite interplay. *Trends in Parasitology*, **32**, 255–264.
https://doi.org/10.1016/j.pt.2015.12.002

Gong X, Qiao Z, Yao H, Zhao D, Eisenhauer N, Scheu S, Liang C, Liu M, Zhu Y-G, Sun X (2024)
Urbanization simplifies soil nematode communities and coincides with decreased ecosystem
stability. *Soil Biology and Biochemistry*, **190**, 109297.
https://doi.org/10.1016/j.soilbio.2023.109297

Gyuris E, Hankó JF, Feró O, Barta Z (2016) Personality and ectoparasitic mites (*Hemipteroseius*
adleri) in firebugs (*Pyrrhocoris apterus*). *Behavioural Processes*, **122**, 67–74.
https://doi.org/10.1016/j.beproc.2015.11.011

Hodges MN, McKinney ML (2018) Urbanization impacts on land snail community composition.
Urban Ecosystems, **21**, 721–735. https://doi.org/10.1007/s11252-018-0746-x

Horsák M, Lososová Z, Čejka T, Juřičková L, Chytrý M (2013) Diversity and biotic homogenization
of urban land-snail faunas in relation to habitat types and macroclimate in 32 Central European
cities. *PLOS ONE*, **8**, e71783. https://doi.org/10.1371/journal.pone.0071783

iNaturalist contributors, iNaturalist (2024) iNaturalist Research-grade Observations.
iNaturalist.org. Occurrence dataset. https://doi.org/10.15468/ab3s5x

Jaffuel G, Půža V, Hug A-S, Meuli RG, Nermuť J, Turlings TCJ, Desurmont GA, Campos-Herrera R
(2019) Molecular detection and quantification of slug parasitic nematodes from the soil and
their hosts. *Journal of Invertebrate Pathology*, **160**, 18–25.
https://doi.org/10.1016/j.jip.2018.11.005

Joint Research Centre (European Commission) (2023) *GHSL data package 2023*. Publications
Office of the European Union, LU.

Jones JS, Leith BH, Rawlings P (1977) Polymorphism in *Cepaea*: a problem with too many
solutions? *Annual Review of Ecology and Systematics*, **8**, 109–143.

Kerstes NAG, Breeschoten T, Kalkman VJ, Schilthuizen M (2019) Snail shell colour evolution in
urban heat islands detected via citizen science. *Communications Biology*, **2**, 264.
https://doi.org/10.1038/s42003-019-0511-6

Korányi D, Egerer M, Rusch A, Szabó B, Batáry P (2022) Urbanization hampers biological control
of insect pests: A global meta-analysis. *Science of The Total Environment*, **834**, 155396.
https://doi.org/10.1016/j.scitotenv.2022.155396

Kramarenko S (2014) Active and passive dispersal of terrestrial mollusks: a review. *Ruthenica*,
24, 1–14.

Leicht K, Seppälä K, Seppälä O (2017) Potential for adaptation to climate change: family-level
variation in fitness-related traits and their responses to heat waves in a snail population. *BMC*
Evolutionary Biology, **17**, 140. https://doi.org/10.1186/s12862-017-0988-x

Lenth R (2023) emmeans: estimated marginal means, aka least-squares means (R package).

Liang H, He Y-D, Theodorou P, Yang C-F (2023) The effects of urbanization on pollinators and
pollination: A meta-analysis. *Ecology Letters*, **26**, 1629–1642. https://doi.org/10.1111/ele.14277

Li X, Liu T, Li H, Geisen S, Hu F, Liu M (2022) Management effects on soil nematode abundance
differ among functional groups and land-use types at a global scale. *Journal of Animal Ecology*,
91, 1770–1780. https://doi.org/10.1111/1365-2656.13744

Lososová Z, Horsák M, Chytrý M, Čejka T, Danihelka J, Fajmon K, Hájek O, Juřičková L, Kintrová K,
Láníková D, Otýpková Z, Řehořek V, Tichý L (2011) Diversity of Central European urban biota:
effects of human-made habitat types on plants and land snails. *Journal of Biogeography*, **38**,
1152–1163. https://doi.org/10.1111/j.1365-2699.2011.02475.x

Manoli G, Fatichi S, Schläpfer M, Yu K, Crowther TW, Meili N, Burlando P, Katul GG, Bou-Zeid E
(2019) Magnitude of urban heat islands largely explained by climate and population. *Nature*,
573, 55–60. https://doi.org/10.1038/s41586-019-1512-9

McKinney ML (2008) Effects of urbanization on species richness: a review of plants and animals.
Urban Ecosystems, **11**, 161–176. https://doi.org/10.1007/s11252-007-0045-4

Minias P (2023) The effects of urban life on animal immunity: Adaptations and constraints.
Science of The Total Environment, **895**, 165085.
https://doi.org/10.1016/j.scitotenv.2023.165085

Morand S, Wilson MJ, Glen DM (2004) Nematodes (Nematoda) parasitic in terrestrial molluscs.
In: *Natural enemies of terrestrial molluscs* (ed Barker GM), pp. 525–557. CABI, Wallingford, UK.

Moreno-Monroy AI, Schiavina M, Veneri P (2021) Metropolitan areas in the world. Delineation
and population trends. *Journal of Urban Economics*, **125**, 103242.
https://doi.org/10.1016/j.jue.2020.103242

Murray MH, Sánchez CA, Becker DJ, Byers KA, Worsley-Tonks KE, Craft ME (2019) City sicker? A
meta-analysis of wildlife health and urbanization. *Frontiers in Ecology and the Environment*, **17**,
575–583. https://doi.org/10.1002/fee.2126

Nakagawa S, Johnson PCD, Schielzeth H (2017) The coefficient of determination R^2 and intra-
class correlation coefficient from generalized linear mixed-effects models revisited and
expanded. *Journal of The Royal Society Interface*, **14**, 20170213.
https://doi.org/10.1098/rsif.2017.0213

Nakagawa S, Schielzeth H (2013) A general and simple method for obtaining R^2 from generalized
linear mixed-effects models. *Methods in Ecology and Evolution*, **4**, 133–142.
https://doi.org/10.1111/j.2041-210x.2012.00261.x

Nicolai A, Filser J, Lenz R, Valérie B, Charrier M (2012) Composition of body storage compounds
influences egg quality and reproductive investment in the land snail *Cornu aspersum*. *Canadian
Journal of Zoology*, **90**, 1161–1170. https://doi.org/10.1139/z2012-081

Oosterhoff LM (1977) Variation in growth rate as an ecological factor in the landsnail *Cepaea
nemoralis* (L.). *Netherlands Journal of Zoology*, **27**, 1–132.
https://doi.org/10.1163/002829677X00072

Ożgo M (2009) Current problems in the research of *Cepaea* polymorphism. *Folia Malacologica*,
16, 55–60. https://doi.org/10.12657/folmal.016.009

Parris KM (2016) *Ecology of urban environments*. Wiley-Blackwell, Chichester, UK.

Pebesma E (2018) Simple features for R: standardized support for spatial vector data. *The R
Journal*, **10**, 439. https://doi.org/10.32614/RJ-2018-009

Pesaresi M, Politis P (2023) GHS-BUILT-S R2023A - GHS built-up surface grid, derived from
Sentinel2 composite and Landsat, multitemporal (1975-2030).
https://doi.org/10.2905/9F06F36F-4B11-47EC-ABB0-4F8B7B1D72EA

Piano E, Souffreau C, Merckx T, Baardsen LF, Backeljau T, Bonte D, Brans KI, Cours M, Dahirel M,
Debortoli N, Decaestecker E, Wolf KD, Engelen JMT, Fontaneto D, Gianuca AT, Govaert L,
Hanashiro FTT, Higuti J, Lens L, Martens K, Matheve H, Matthysen E, Pinseel E, Sablon R, Schön I,
Stoks R, Doninck KV, Dyck HV, Vanormelingen P, Wichelen JV, Vyverman W, Meester LD,
Hendrickx F (2020) Urbanization drives cross-taxon declines in abundance and diversity at
multiple spatial scales. *Global Change Biology*, **26**, 1196–1211.
https://doi.org/10.1111/gcb.14934

Pieterse A, Malan AP, Ross JL (2017) Nematodes that associate with terrestrial molluscs as
definitive hosts, including *Phasmarhabditis hermaphrodita* (Rhabditida: Rhabditidae) and its
development as a biological molluscicide. *Journal of Helminthology*, **91**, 517–527.
https://doi.org/10.1017/S0022149X16000572

Pollard E, Cooke AS, Welch JM (1977) The use of shell features in age determination of juvenile
and adult Roman snails *Helix pomatia*. *Journal of Zoology*, **183**, 269–279.
https://doi.org/10.1111/j.1469-7998.1977.tb04186.x

R Core Team (2023) R: a language and environment for statistical computing.

Radwan MA, El-Gendy KS, Gad AF (2020) Biomarker responses in terrestrial gastropods exposed
to pollutants: A comprehensive review. *Chemosphere*, **257**, 127218.
https://doi.org/10.1016/j.chemosphere.2020.127218

Rae R (2017) The gastropod shell has been co-opted to kill parasitic nematodes. *Scientific*
Reports, **7**, 4745. https://doi.org/10.1038/s41598-017-04695-5

Rae R (2018) Shell encapsulation of parasitic nematodes by *Arianta arbustorum* (Linnaeus, 1758)
in the laboratory and in field collections. *Journal of Molluscan Studies*, **84**, 92–95.
https://doi.org/10.1093/mollus/eyx045

Rae R (2023) Avoidance and attraction behaviour of slugs exposed to parasitic nematodes.
Journal of Invertebrate Pathology, **197**, 107896. https://doi.org/10.1016/j.jip.2023.107896

Rae RG, Robertson JF, Wilson MJ (2008) Susceptibility and immune response of *Deroceras*
reticulatum, *Milax gagates* and *Limax pseudoflavus* exposed to the slug parasitic nematode
Phasmarhabditis hermaphrodita. *Journal of Invertebrate Pathology*, **97**, 61–69.
https://doi.org/10.1016/j.jip.2007.07.004

Rosin ZM, Kwieciński Z, Lesicki A, Skórka P, Kobak J, Szymańska A, Osiejuk TS, Kałuski T, Jaskulska
M, Tryjanowski P (2018) Shell colour, temperature, (micro)habitat structure and predator
pressure affect the behaviour of *Cepaea nemoralis*. *The Science of Nature*, **105**, 35.
https://doi.org/10.1007/s00114-018-1560-2

Russo J, Madec L (2011) Dual strategy for immune defense in the land snail *Cornu aspersum*
(Gastropoda, Pulmonata). *Physiological and Biochemical Zoology: Ecological and Evolutionary*
Approaches, **84**, 212–221. https://doi.org/10.1086/659123

Russo J, Madec L (2013) Linking immune patterns and life history shows two distinct defense
strategies in land snails (Gastropoda, Pulmonata). *Physiological and Biochemical Zoology*, **86**,
193–204. https://doi.org/10.1086/669482

Scheil AE, Hilsmann S, Triebskorn R, Köhler H-R (2013) Shell colour polymorphism, injuries and
immune defense in three helicid snail species, *Cepaea hortensis*, *Theba pisana* and *Cornu*
aspersum maximum. *Results in Immunology*, **3**, 73–78.
https://doi.org/10.1016/j.rinim.2013.06.002

Scheil AE, Hilsmann S, Triebskorn R, Köhler H-R (2014) Shell colouration and parasite tolerance
in two helicoid snail species. *Journal of Invertebrate Pathology*, **117**, 1–8.
https://doi.org/10.1016/j.jip.2014.01.003

Schiavina M, Freire S, Carioli A, MacManus K (2023) GHS-POP R2023A - GHS population grid
multitemporal (1975-2030). https://doi.org/10.2905/2FF68A52-5B5B-4A22-8F40-
C41DA8332CFE

Schiavina M, Melchiorri M, Pesaresi M (2023) GHS-SMOD R2023A - GHS settlement layers,
application of the Degree of Urbanisation methodology (stage I) to GHS-POP R2023A and GHS-

BUILT-S R2023A, multitemporal (1975-2030). https://doi.org/10.2905/A0DF7A6F-49DE-46EA-
9BDE-563437A6E2BA

Schiavina M, Moreno-Monroy A, Maffenini L, Veneri P (2019) GHS-FUA R2019A - GHS functional
urban areas, derived from GHS-UCDB R2019A (2015). https://doi.org/10.2905/347F0337-F2DA-
4592-87B3-E25975EC2C95

Segade P, García-Estévez J, Arias C, Iglesias R (2013) Parasitic infections in mixed system-based
heliciculture farms: dynamics and key epidemiological factors. *Parasitology*, **140**, 482–497.
https://doi.org/10.1017/S0031182012001795

Shultz AJ, Adams BJ, Bell KC, Ludt WB, Pauly GB, Vendetti JE (2020) Natural history collections
are critical resources for contemporary and future studies of urban evolution. *Evolutionary*
Applications, **14**, 233–247. https://doi.org/10.1111/eva.13045

Staikou A, Sagonas K, Spanoudi O, Savvidou K, Nazli Z, Feidantsis K, Michaelidis B (2024)
Activities of antioxidant enzymes and Hsp levels in response to elevated temperature in land
snail species with varied latitudinal distribution. *Comparative Biochemistry and Physiology Part*
B: Biochemistry and Molecular Biology, **269**, 110908.
https://doi.org/10.1016/j.cbpb.2023.110908

Szulkin M, Munshi-South J, Charmantier A (Eds.) (2020) *Urban Evolutionary Biology*. Oxford
University Press. https://doi.org/10.1093/oso/9780198836841.001.0001

Theodorou P (2022) The effects of urbanisation on ecological interactions. *Current Opinion in*
Insect Science, **52**, 100922. https://doi.org/10.1016/j.cois.2022.100922

Valdés-Correcher E, Popova A, Galmán A, Prinzing A, Selikhovkin AV, Howe AG, Mrazova A,
Dulaurent A-M, Hampe A, Tack AJM, Bouget C, Lupaştean D, Harvey D, Musolin DL, Lövei GL,
Centenaro G, Halder IV, Hagge J, Dobrosavljević J, Pitkänen J-M, Koricheva J, Sam K, Barbaro L,
Branco M, Ferrante M, Faticov M, Tahadlová M, Gossner M, Cauchoix M, Bogdziewicz M,
Duduman M-L, Kozlov MV, Bjoern MC, Mamaev NA, Fernandez-Conradi P, Thomas RL,
Wetherbee R, Green S, Milanović S, Moreira X, Mellerin Y, Kadiri Y, Castagneyrol B (2022)
Herbivory on the pedunculate oak along an urbanization gradient in Europe: Effects of
impervious surface, local tree cover, and insect feeding guild. *Ecology and Evolution*, **12**, e8709.
https://doi.org/10.1002/ece3.8709

Wacker A, Baur B (2004) Effects of protein and calcium concentrations of artificial diets on the
growth and survival of the land snail *Arianta arbustorum*. *Invertebrate Reproduction &*
Development, **46**, 47–53. https://doi.org/10.1080/07924259.2004.9652605

Wickham H, Averick M, Bryan J, Chang W, McGowan L, François R, Grolemund G, Hayes A, Henry
L, Hester J, Kuhn M, Pedersen T, Miller E, Bache S, Müller K, Ooms J, Robinson D, Seidel D, Spinu
V, Takahashi K, Vaughan D, Wilke C, Woo K, Yutani H (2019) Welcome to the Tidyverse. *Journal*
of Open Source Software, **4**, 1686. https://doi.org/10.21105/joss.01686

Williamson P (1979) Age determination of juvenile and adult *Cepaea*. *Journal of Molluscan*
Studies, **45**, 52–60.

Williams AJ, Rae R (2015) Susceptibility of the Giant African snail (*Achatina fulica*) exposed to
the gastropod parasitic nematode *Phasmarhabditis hermaphrodita*. *Journal of Invertebrate*
Pathology, **127**, 122–126. https://doi.org/10.1016/j.jip.2015.03.012

Williams A, Rae R (2016) *Cepaea nemoralis* (Linnaeus, 1758) uses its shell as a defence
mechanism to trap and kill parasitic nematodes. *Journal of Molluscan Studies*, **82**, 349–350.
https://doi.org/10.1093/mollus/eyv064

Wilson MJ, Hughes LA, Hamacher GM, Glen DM (2000) Effects of *Phasmarhabditis*
hermaphrodita on non-target molluscs. *Pest Manag Sci*, **56**, 711–716.
https://doi.org/10.1002/1526-4998(200008)56:8%3C711::AID-PS185%3E3.0.CO;2-O

Wilson MJ, Hughes LA, Jefferies D, Glen DM (1999) Slugs (*Deroceras reticulatum* and *Arion ater*
agg.) avoid soil treated with the rhabditid nematode *Phasmarhabditis hermaphrodita*. *Biological*
Control, **16**, 170–176. https://doi.org/10.1006/bcon.1999.0757

Wolda H (1967) The effect of temperature on reproduction in some morphs of the landsnail
Cepaea nemoralis (L.). *Evolution*, **21**, 117–129. https://doi.org/10.1111/j.1558-
5646.1967.tb00135.x

Wynne R, Morris A, Rae R (2016) Behavioural avoidance by slugs and snails of the parasitic
nematode *Phasmarhabditis hermaphrodita*. *Biocontrol Science and Technology*, **26**, 1129–1138.
https://doi.org/10.1080/09583157.2016.1185513

Żbikowska E, Marszewska A, Cichy A, Templin J, Smorąg A, Strzała T (2020) *Cepaea* spp. as a
source of *Brachylaima mesostoma* (Digenea: Brachylaimidae) and *Brachylecithum* sp. (Digenea:
Dicrocoeliidae) larvae in Poland. *Parasitology Research*, **119**, 145–152.
https://doi.org/10.1007/s00436-019-06516-2