A reactant-dependent algebraic framework to define, construct, characterize and assess the redundant
forms of a biochemical network

Siddhartha Kundu[1, a]

[1] Department of Biochemistry,
All India Institute of Medical Sciences,
Ansari Nagar,
New Delhi 110029,
INDIA
[a] Correspondence: siddhartha_kundu@yahoo.co.in, siddhartha_kundu@aiims.edu

Abstract

The intracellular milieu, replete with concomitantly occurring biochemical networks, constitutes a

complex physicochemical environment wherein macromolecules interact and whose activities can be

modified by small molecules. These metabolism-contributing, signal-transducing and response-regulating

networks, and the redundant forms that result, thereof, are responsible for complex biochemical-,

physiological- and clinical-outcomes. It is unclear whether a unique mathematical structure, which is

concurrently resolvable into redundant, unequal and functionally relevant substructures, can be ascribed

to a biochemical network. Here, we develop a reactant-dependent algebraic framework to define,

construct, characterize and assess the redundant forms of a biochemical network. We deploy a

constraint-based approach to populate and select plausible reaction vectors in accordance with known

paradigms of biochemically relevant outcomes. There is a strict lower bound for the number of

participating reactants with integral changes, across all molecules of a reactant, which are Boolean

(zero, signed non-zero) and combinatorially distributed across linearly independent plausible reaction

vectors. The reactants and the selected plausible reaction vectors form a set of unique stoichiometry

number matrices, each of which represents a distinct biochemical network which when multiplied by a

set of positive and non-unitary rational numbers will generate a library of equivalent rational number

matrices. The matrices that comprise this library are unequal, disjoint, form a semigroup with respect

to addition, share the same null space and represent the redundant forms of a biochemical network.

The presented framework is theoretically sound, mathematically rigorous, readily implementable, easily

parameterized and can be used to assess the biomedical relevance for the redundant forms of a

biochemical network. We conclude our thesis with plausible explanations into the genesis of

formaldehyde resistance and thence nosocomial infections by *Klebsiella spp.*

Keywords: Reactant-dependent algebraic framework to model, characterize and assess the redundant

forms of a biochemical network; Constraint-based, mathematically rigorous and biochemically relevant

construction; Reactant-specific plausible reaction vectors; Assembly of preselected reactant-specific

plausible reaction vectors into unique stoichiometry number matrices; Set of strictly positive and non-

unitary rational number multipliers; Library of stoichiometry number matrix specific equivalent rational

number matrices; Dissociation constant-based non-local and local parameterization;

Introduction

Biochemical networks of enzyme-mediated transformations, exchange- and diffusion-reactions, and formation/disassembly of molecular complexes) are regulated by short (feedback mechanisms, post-translational modifications)- and long (*de novo* synthesis, degradation)-term mechanisms [1, 2]. A biochemical network is either studied by optimizing the biomass of a reactant of interest and enumerating elementary vectors or is data-driven, inverse and inferential. Algorithms which optimize one or more reactants are likely to be biased in choice, type and usage of constraints, and will result in studies with reduced biomedical relevance despite being numerically sound [1, 3-6]. Enumeration-based algorithms on the other hand, involve extensive computations that are NP-hard and will become intractable for large biochemical networks [2, 6-8]. The availability of- and accessibility to- empirical- and clinical-datasets has the potential to recreate, without bias, relevant data-driven biochemical networks, infer directionality for a given biochemical network (polynomial equation) or represent an arbitrary state of a modelled network (continuous-time Lyapunov equation) [9-12]. In these approaches perturbations are modelled as stochastic fluctuations and are used to derive the covariance (weak)- or correlation (strong)-interaction matrices for the set of nodes that comprise the modelled biochemical network [13-15].. This reliance on empirical data both, *a priori* and *a posteriori*, along with the computational complexity in computing the Jacobian are limitations in the inferential study of biochemical networks [13-15].

These limitations notwithstanding, inferred biochemical networks have a role in precision medicine and data mining [16-20]. For example, the use of guidance-based radio- and immune-therapeutics in managing chronic inflammatory diseases, neurodegenerative pathologies and recalcitrant and recurring malignancies is being used to assess dose response and predict outcome such as the propensity, progress and resolution of an adverse drug reaction [16-20]. In particular, these investigations combine the expression/abundance profiles of system-specific panels of biomarkers (genes, proteins, small molecules) into one of several machine learning (ML)-algorithms and predict patient response and/or

compliance [18-20]. However, despite these advancements the primary or raw data (time-series kinetic data, "omics"-based datasets) is dependent on the platform used, experimental design and will require several numerical assumptions and extensive- and varied-normalization protocols [9-14]. Additionally, the use of expensive equipment, reagents and manpower can potentially restrict the utility of these investigations. In contrast to the aforementioned discussion on large scale modelling and analysis, causal networks and the inferences that result thereof, examine the effects of variable stimuli on several equally plausible and well-defined outcomes of a few nodes. Although causal network modelling has had several proven successes in investigating discrete real world events, the requirement for reference data, simple architecture and simpler rules, precludes consideration of complex biochemical systems other than for mining large biological datasets for lists of statistically significant genes/proteins of well characterized networks and in response to defined perturbations [21-25].

As opposed to the data-driven approach discussed vide supra, algebraic models for biochemical networks are usually formulated as population models [26, 27]. In particular, chemical reaction networks, with regards to the partaking reactants, are modelled as either self-sustaining/persistent once initialized or limiting (zero-occurring) [28, 29]. Although these investigations have enhanced our repertoire of modelling strategies for functional biochemical networks, novel data-analytics and parameter-free frameworks, the absence of biochemically-relevant parameterization may mitigate the potential biomedical impact and relevance of these studies [30, 31]. The dissociation (Kd)-, association (Ka)- and Michaelis-Menten (Km)-constants are important parameters in comprehending the outcome of a biochemical reaction (non-enzyme, enzyme) and are empirically determined (dual polarization interferometry, surface plasmon resonance, stopped-flow spectroscopy), inferred (regularized least squares pKa spectrum, static binding isotherms) and/or theoretically derived [32-38]. The probable dissociation constant, for the i^{th}-reaction, is an unbiased numerical measure that is computed from a null space-generated subspace of combinatorially summed non-redundant and non-trivial vectors of the stoichiometry number matrix for a biochemical network [7, 8]. The probable dissociation constant is a strictly positive real number which can unambiguously compute the forward-, reverse- or equivalent-outcomes for a reaction [7, 8].

A biochemical network is modelled as an undirected graph with a finite number of nodes and edges, as opposed to a specific pathway which is, in contrast, rule-based and a directed version of the same.

However, modelling the multitude of genome-scale derived data points mandates the use of vertex-

bounded polytopes such as an n-dimensional simplicial complex with $n + 1$ affinely-independent

vertices/nodes and a finite number ($\sum_{m=0}^{m=n} \binom{n+1}{m+1}$) of faces. Here too, the absence of a biochemical

rationale for the all- or none-manner in which the architecture of a network is perturbed whence

node(s) are added/subtracted, choice of polytope, computational intractability at higher dimensions and

the number of steady-state mapping constraints present unique challenges [39-41]. Redundant forms of

a biochemical network (molecular, modular, network-level), are not uncommon, contribute significantly

to physiological-, cellular- and biochemical-homeostasis (complement activation, blood clotting pathways,

circadian rhythms) and are well-characterized (KEGG, MetaCyc) [42-46]. Relevance-studies, for these

forms, include node-deletion (knock-out) or -inclusion (knock-in) with sensitivity analysis (single

nucleotide polymorphism (SNP)-mapping) and last common ancestor (LCA)-tracing of molecular variants

for epidemiological studies (source of an infecting pathogen, pedigree analysis, phylogenetic analysis)

[3-7, 47, 48]. It is however, unclear whether a unique mathematical structure, which is concurrently

resolvable into redundant, unequal and functionally relevant substructures, can be ascribed to a whole

biochemical network. This paucity of mathematical descriptors notwithstanding, network-level redundant

forms of a biochemical network occur in groups of organisms with significant taxonomic relatedness

(species, strains, genotypes), are likely to facilitate herd immunity and may confer-/transfer-resistance to

infectious- and chemotherapeutic-agents [49].

Here, we seek network-level redundant forms of a biochemical network, which are unequal, have a

sound theoretical basis, mathematically rigorous, biochemically relevant and amenable to minimal

perturbation(s). In order to accomplish this, we will develop a reactant-dependent algebraic framework

to define and construct the redundant forms of a biochemical network, study its properties via suitable

metrics (mathematical, computational) and assess biomedical relevance of the same (known biochemical

pathways). In section 2, pertinent definitions and concepts in the form of biochemically relevant

Constraints will be introduced and discussed. In "Methods" (section 3), we will describe the

mathematical and computational tools needed to complement the theoretical assertions that have been

outlined. This section will also describe the derivation of network-specific metrics which can be used

to parameterize and establish relevance of the developed algebraic framework. In section 4 ("Results"),

a detailed examination of the mathematical properties of the developed framework will be presented

along with in-depth computational studies of known biochemical pathways. The "Discussion" (section

5) will highlight the rationale, motivation and biomedical relevance of this framework. The study

concludes (section 6) by summarizing the salient features, major findings and limitations of this study.

Detailed proofs, in order of appearance, for the mathematical formalism deployed, i.e., Constraints (Cs),

definitions-environment (D), lemmas (L), theorems (T), corollaries (C) and propositions (P), are

included in section 7 as supplementary material (STx) and cited wherever applicable

2　Definitions, concepts and preliminary results for a reactant-specific algebraic framework

Biochemical networks comprise interactions between finite numbers of molecular species of distinct

reactants. The "reactions" that ensue are parameterized using enzyme-mediated catalysis (Km, Kcat,

turnover number, catalytic efficiency), membrane-limited transport and macromolecular complex

formation with disassembly (Kd, Ka).

2.1 Dissociation constants as units of change for reactants of a biochemical network

Consider a bimolecular non-enzymatic reaction,

$$a.\,A + c.\,C \underset{k_b}{\overset{k_f}{\rightleftarrows}} ac.\,AC \qquad\qquad RXN(1)$$

where,

$A, C := \text{Reactants undergoing forward reaction with rate constant } k_f \left(Rf = k_f.[A]^{\theta_A}.[C]^{\theta_C}\right)$ (1)

$AC := \text{Reactant undergoing reverse reaction with rate constant } k_b \left(Rb = k_b.[AC]^{\theta_{AC}}\right)$ (2)

$(-a, -c, ac) := \text{Stoichiometry numbers for reactants and products of the forward reaction}$ (3)

$(a, c, -ac) := \text{Stoichiometry numbers for reactants and products of the reverse reaction}$ (4)

$(\theta_A, \theta_C, \theta_{AC}) := \text{Stoichiometry coefficients for reactants of a reaction (forward, reverse)}$ (5)

$Kd := \text{Dissociation constant for the reversible reaction RXN (1)}$ (6)

14　Let us assume that this reaction takes place at a steady state,

$$Rf = Rb \qquad\qquad (6)$$

$$\Rightarrow \frac{[AC]^{\theta_{AC}}}{[A]^{\theta_A}.[C]^{\theta_C}} = \frac{k_f}{k_b} \in \mathbb{R} \cap (0, +\infty) \text{ and } k_b \neq 0 \quad (6.1)$$

$$= Kd(AC) \qquad\qquad (6.2)$$

$$= \frac{1}{Ka(AC)} \qquad\qquad (6.3)$$

16　We can extend the annotation viz. **Eqs. (1-6)** to define the forward and reverse reactions,

$$\{Kd(AC), Ka(AC)^{-1}\} \in \mathbb{R}_+ \text{ where } k_f, k_b \neq 0 \quad \textbf{(D1)}$$

18　Let us describe a stoichiometry number for a reactant(s) from the dissociation constant with the linear

19　map (h),

$$h: Kd(.) \in \mathbb{R} \cap (0, +\infty) \mapsto \Delta m_j = \sum_{u=1}^{u=U} m_{j,u,t+\delta t} - m_{j,u,t} \in \mathbb{Z} \text{ where } m_j = U. m_{j,u} \text{ species} \quad \textbf{(D2)}, (7), (8)$$

$$\Delta m_j \propto \lfloor \ln(Kd) \rfloor = \left| \ln\left(\frac{[AC]^{\theta_{AC}}}{[A]^{\theta_A}.[C]^{\theta_C}} \right) \right| \quad (9), (10)$$

$$\textbf{Case1}: (Kd > 1) \Rightarrow \frac{[AC]^{\theta_{AC}}}{[A]^{\theta_A}.[C]^{\theta_C}} > 1 \Rightarrow \left| \ln\left(\frac{[AC]^{\theta_{AC}}}{[A]^{\theta_A}.[C]^{\theta_C}} \right) \right| \geq 1 > 0 \Rightarrow \Delta m_j > 0 \quad (11-14)$$

$$\textbf{Case2}: (Kd \in (0,1)) \Rightarrow \frac{[AC]^{\theta_{AC}}}{[A]^{\theta_A}.[C]^{\theta_C}} \in (0,1) \Rightarrow \left| \ln\left(\frac{[AC]^{\theta_{AC}}}{[A]^{\theta_A}.[C]^{\theta_C}} \right) \right| < 0 \Rightarrow \Delta m_j < 0 \quad (15-17)$$

$$\textbf{Case3}: (Kd \approx 1) \Rightarrow \frac{[AC]^{\theta_{AC}}}{[A]^{\theta_A}.[C]^{\theta_C}} \approx 1 \Rightarrow \left| \ln\left(\frac{[AC]^{\theta_{AC}}}{[A]^{\theta_A}.[C]^{\theta_C}} \right) \right| = 0 \Rightarrow \Delta m_j = 0 \quad (18-20)$$

2   This map is onto and many-one since,

$$\text{if}(Kd(.) \in \mathbb{R} \cap (0,1)) \text{ then } \Delta m_j \in \mathbb{Z}_- \quad (21)$$
$$\text{if}(Kd(.) \in \mathbb{R} \cap \{1\}) \text{ then } \Delta m_j = 0 \quad (22)$$
$$\text{if}(Kd(.) \in \mathbb{R} \cap (1, +\infty)) \text{ then } \Delta m_j \in \mathbb{Z}_+ \quad (23)$$

4   We now describe a biochemical network of j-indexed $(j = 1, \ldots, J)$ m-reactants $(m_1, \ldots, m_J)$ where each

5   m_j is comprised of $U. m_{j,u}$ species,

$$\mathbb{m} = [m_1, \ldots, m_J]^T \text{ where each } m_j \sim (m_{j,u=1}, \ldots, m_{j,u=U}) = U. m_{j,u} \quad \textbf{(D3)}, (24), (25)$$

7   The reactions that comprise this network are i-indexed $(i = 1, \ldots, I)$ r-reactions $(r_1, \ldots, r_I)$ each of

8   which is represented by a dissociation constant which we can rewrite as,

$$r_i \equiv Kd_i \twoheadrightarrow \Delta m_j \in \mathbb{Z} \quad [\text{From D2, Eqs.} (21-23)] \quad \textbf{(D4)}$$

10   We will express the stoichiometry number for each reactant in terms of a reaction-specific dissociation

11   constant in accordance with **D2-D4**. Since we are concerned with biochemical networks within a cell

12   we will subsume a reactant's contribution to each and every reaction, i.e., a reaction vector (RV),

$$Kd_i(\mathbb{m}) = [Kd_i(m_1), \ldots, Kd_i(m_j)]^T = [\Delta m_{1i}, \ldots, \Delta m_{Ji}]^T = [\Delta m_1, \ldots, \Delta m_J]_i^T \overset{\text{def}}{=} r_i \in \mathbb{Z}^J \quad (26-28), \textbf{(D5)}$$

2.2 Constraints to construct a reactant-dependent algebraic framework

The intracellular milieu is an interplay of physico- and bio-chemical processes working in concert to

achieve complex function. In our study we define and classify known instances of redundancies that

are observed for a biochemical network in terms of the number of reactants. In order to assign a

numerical measure to this constraint we will populate each RV in accordance with known biochemical

paradigms.

1   *2.2.1   Constraints to define, describe and populate a plausible reaction vector*

2   The basic unit of our algebraic framework is the plausible reaction vector (PRV) which unlike an RV

3   will be, once populated, numerically tenable and representative of a true biochemical reaction.

4   Constraint 1 **(Cs1):** The contribution to the i^{th}-PRV by J-reactants is Boolean (non-zero, zero),

$$\exists\left(\Delta m_{ji} \neq 0, \Delta m_{ki} = 0\right) \in \mathbf{r}_i \qquad \textbf{(D6)}$$
where
$$\Delta m_{ji}, \Delta m_{ki} \in \mathbf{r}_i \in \mathbf{R} - \{\mathbf{0}\} \subset \mathbb{Z}^J \qquad (29)$$
$$j, k = 1, \dots, J; j \neq k; i = 1, \dots, \hat{I} \in \mathbb{N}; J \geq 1$$

For example,
$$\mathbf{r}_i \in \left\{[\,2 - 2\ \ 0\ 0]_i^T, [-1\ 0\ 1\ 0]_i^T, [0 - 3\ 0\ 3]_i^T, [0\ 3\ 0 - 3\ ]_i^T\right\} \quad (30)$$
$$\mathbf{r}_i \notin \left\{[\,2 - 2\ -1 - 1]_i^T, [-11\ 1 - 1]_i^T\right\} \qquad (31)$$

6   Constraint 2 **(Cs2)**: The cardinality of a set of i-indexed $\hat{I}$-PRVs $\left(\#[\mathbf{r}_1, \dots, \mathbf{r}_{\bar{i}}]^T\right)$, for a biochemical

7   network with j-indexed J-reactants $(\Delta m_1, \dots, \Delta m_J)$, is a function of the number of unique reactants that

8   partake in the network.

$$\#\mathbf{R} \overset{\text{def}}{=} \hat{I} = 2^J \qquad \textbf{(D7)}, (32), (33)$$
$$\text{where } \mathbf{R} = [\mathbf{r}_1, \dots, \mathbf{r}_{\bar{i}}]^T \subset \mathbb{Z}^J; i = 1, \dots, \hat{I}; J \geq 1 \qquad (34 - 36)$$

10   In order to uphold the premise of this study, i.e., a framework to model and characterize the

11   redundant forms of a biochemical network, we will exclude PRVs that contribute to molecular

12   (isoforms, isoenzymes, alternate splicing, pseudogenes, duplicate genes, truncated proteins)- and modular

13   (futile cycles, sequestrations) –redundancy, for a biochemical network.

14   Constraint 3 **(Cs3)**: The PRVs used to construct the algebraic framework must be linearly independent,

$$\text{if}(\lambda_i. \mathbf{r}_i, \lambda_k. \mathbf{r}_k \in \mathbf{R} \subset \mathbb{Z}^J \text{ s.t. } \lambda_i. \mathbf{r}_i + \lambda_k. \mathbf{r}_k = \mathbf{0} \text{ where } \lambda_i, \lambda_k \in \mathbb{R} \text{ and } \mathbf{r}_i, \mathbf{r}_k \neq \mathbf{0})$$
$$\text{then}(\lambda_i = \lambda_k = 0) \qquad \textbf{(D8)}$$
$$\text{where } i \neq k \text{ and } i, k \in \left[1, \hat{I}\right]$$

Consider the contradiction,
$$\mathbf{r}_i = (-1). \mathbf{r}_k \qquad (37)$$
$$\mathbf{r}_i + \mathbf{r}_k = (-1). \mathbf{r}_k + \mathbf{r}_k = \mathbf{0} \qquad (38)$$

16   We will ensure that **Cs3** remains relevant at all stages of the modelling process by using a

17   combinatorial approach to select- and populate-PRVs and identify reactant classes. The latter are

18   combinations of subsets of reactants taken simultaneously. Additionally, we will preclude PRVs that are

unlikely to be relevant from a biochemical perspective. For example, a zero vector means that every

reactant is non-contributing, which is improbable given the complexity of the intracellular milieu

$\left(\Delta m_{ji} = 0 \; \forall j; \mathbf{D9}, \mathbf{Eq.(39)}\right)$. It is equally improbable that every reactant will participate in every

reaction (futile cycles, sequestration by membrane-bound compartments, feedback loops, partial reactions,

competing metabolites) $\left(\Delta m_{ji} \neq 0 \; \forall j; \mathbf{D10}, \mathbf{Eq.(40)}\right)$. On the other hand, a singly signed non-zero entry

-populated PRV of the form $\{\Delta m_{ji}, \Delta m_{ki}\}$ where,

$\Delta m_{ji} = 0, \Delta m_{ki} = \begin{cases} \wedge_{k=1}^{k=J}(+)\Delta m_{ki} \\ \wedge_{k=1}^{k=J}(-)\Delta m_{ki} \end{cases}$ s.t. $k \neq j$ and $k = 1, \dots, J; \mathbf{D11}, \mathbf{Eq.(41)}$ will indicate partial- or

incomplete-reactions for a reactant or a product.

Constraint 4 (**Cs4**): The cardinality for the subset of excluded PRVs $\left(\#\widehat{\mathbf{R}}\right)$, for J-reactants, is

$$\#\widehat{\mathbf{R}} = \begin{pmatrix} \binom{J}{0} + \binom{J}{1} + \binom{J}{J-1} + \binom{J}{J} = \underbrace{\binom{J}{0} + \binom{J}{J}}_{[\text{From } \mathbf{D9,D10}]} + \underbrace{\binom{J}{1} + \binom{J}{J-1}}_{[\text{From } \mathbf{D11}]} \\ = 1 + 1 + J + J = 2 + 2.J \end{pmatrix} \overset{\text{def}}{=} 2.(1+J) \quad (\mathbf{D12}), (42-45)$$

Constraint 5 (**A5**): The revised cardinality for the set of J-dependent PRVs,

$$\hat{I} = \#\left(\mathbf{R}\backslash\widehat{\mathbf{R}}\right) = \#\mathbf{R} - \#\widehat{\mathbf{R}} \overset{\text{def}}{=} 2^J - 2.(1+J) \quad [\text{From } \mathbf{A6, A8}] \quad (\mathbf{D13}), (46-48)$$
For example,
$J = 4: \hat{I} = 6$ $\qquad\qquad\qquad\qquad\qquad\qquad (49), (50)$
$J = 5: \hat{I} = 20$ $\qquad\qquad\qquad\qquad\qquad\qquad (51), (52)$

We will utilize signed non-zero reactants $\left((+)\Delta m_{ji}, (-)\Delta m_{ki} \text{ where } k \neq j, k = 1, \dots, J\right)$ to populate a

PRV in terms of their role in a reaction. A molecular species that is utilized/converted by a

biochemical reaction is a reactant and is annotated as $\left((-)\Delta m_{ji}\right)$ (**D14**). A product is defined as a

molecular species which is formed by a biochemical reaction $\left((+)\Delta m_{ji}\right)$ (**D15**). Clearly, **D14** and **D15**

are only possible if the generating reaction is strictly forward or reverse $(Kd_i \in \mathbb{R}_+ - \{1\})$. In contrast,

a reactant without a net contribution $\left(\Delta m_{ji} = 0\right)$ is likely to be due to uncertainty in the raw data or

even an equivalent reaction $(Kd_i \in \mathbb{R}_+ \cap \{1\})$ [7, 8].

Constraint 6 (**A6**): The stoichiometry contribution by any j-indexed m-reactant $\left(\Delta m_j\right)$ to a PRV is

[**From D2, D3, D14, D15**],

$$\Delta m_{ji} = \begin{cases} (-)\Delta m_{ji} \overset{\underline{def}}{=} \text{Integral change across all U} - \text{molecular species as a reactant} \\ (+)\Delta m_{ji} \overset{\underline{def}}{=} \text{Integral change across all U} - \text{molecular species as a product} \\ 0 \overset{\underline{def}}{=} \text{Integral change across all U} - \text{molecular species without a net contribution} \end{cases} \quad \textbf{(D16)}$$

where,

$$\Delta m_{ji} \in \mathbf{r}_i \in \mathbf{R} - \{\mathbf{0}\} \subset \mathbb{Z}^J; i = 1, \dots, \hat{I}; J \geq 1 \tag{53}$$

For RXN(1),

	Forward (f)	Reverse (b)	
A	$-a$	a	(54)
AB	$\mathfrak{a}$	$-\mathfrak{a}$	
C	0	0	

Constraint 7 (Cs7): The i^{th}-PRV must have at least one pair of reactants that are differently/oppositely signed,

$$\exists \left((-)m_{ji}, (+)m_{ki} \right) \in \mathbf{r}_i \tag{D17}$$

where

$$m_{ji}, m_{ki} \in \mathbf{r}_i \in \mathbf{R} - \{\mathbf{0}\} \subset \mathbb{Z}^J; j, k \in [1, J]; j \neq k \tag{55 - 57}$$
$$i = 1, \dots, \hat{I}; J \geq 1$$

For example

$$\mathbf{r}_i = [\, 2\ -2\ 0\,]_i^T \vee [-1\ 0\ 1]_i^T \vee [-3\ 0\ 3]_i^T \vee [1\ 1\ -1\,]_i^T \vee [0\ 3\ -3\,]_i^T \dots \tag{58}$$

Constraint 8 (Cs8): The i^{th}-PRV must comprise two non-zero and two zero entries/reactants,

$$\#\mathcal{Z}, \#\bar{\mathcal{Z}} \geq 2 \qquad [\text{From } \mathbf{Cs2} - \mathbf{Cs5}] \quad (59), (60)$$

where

$$\mathcal{Z} \ni m_{ji} \subset \mathbf{r}_i | m_{ji} = 0 \tag{D18}$$
$$\bar{\mathcal{Z}} \ni m_{ji} \subset \mathbf{r}_i | m_{ji} \neq 0 \tag{D19}$$
$$\mathbf{r}_i \in \mathbf{R} \backslash \widehat{\mathbf{R}} \subset \mathbb{Z}^J \tag{D20}$$

and

$$i = 1, \dots, \hat{I}$$

Constraint 9 (Cs9): The lower bound for the cardinality of an i^{th}-PRV is,

$$\dim(\mathbf{r}_i) = J \geq 4 \qquad [\text{From } \mathbf{Cs8}] \quad (D21)$$
$$\text{where } \mathbf{r}_i \in \mathbf{R} \backslash \widehat{\mathbf{R}} \subset \mathbb{Z}^J \text{ and } i = 1, \dots, \hat{I}$$

On the basis of **Cs1- Cs9** we will now refine our definition for a PRV as a non-zero, linearly independent vector with differently signed stoichiometry numbers for contributing reactants and zero values for non-contributing members both of which will not be less than 2 entries. We can now formally define the lower bound for the number of reactants $(\min(J) \geq 4; \mathbf{D22}$ [From **Cs9**]) that will result in network-level redundant forms of a biochemical network thereby distinguishing it from molecular $(J = 1)$- and modular $(J = 2,3)$-versions of the same.

2.2.3   Constraints to assemble a reactant-dependent stoichiometry number matrix

We have defined the criteria to populate and select subsets from the PRVs. We now describe

Constraints that will combine selected PRVs into unique reactant-specific stoichiometry number

matrices.

Constraint 10 (**Cs10**): The number of distinct subset(s) that J-reactants is divided into, where each

subset comprises $J \in [s + 1, J - 2]$ reactants that are concurrently considered, is computed from an s-

indexed $\mathbb{J}$-membered ordered list,

$$(.) \left(\hat{\mathbf{J}}(J) \right) \stackrel{\text{def}}{=} (.)(\hat{\mathbf{J}}_J) = (.)_{\hat{J}_J} = \left((.)_{\hat{J}_{J,1}}, \dots, (.)_{\hat{J}_{J,\mathbb{J}}} \right)^{\text{T}} \subset \mathbb{N}^{\mathbb{J}} \quad [\text{From } \mathbf{Cs9}] \quad (\mathbf{D23}), (61), (62)$$

$$\text{where } \mathbb{J} \stackrel{\text{def}}{=} \#\hat{J}_J = J - 3 = J + 1 - 4 = (J + 1) - (\#\hat{\mathbf{R}}) \qquad (\mathbf{D24}), (63 - 65)$$

$$\text{and } J \geq 4, \#\hat{\mathbf{R}} = 4 \text{ and } s = 1, \dots, \mathbb{J}$$

Examples:

$$J = 4: \#\hat{J}_4 = 1; (.)_{\hat{J}_4} = \left[(.)_{\hat{J}_{4,1}} \right]^{\text{T}} \subset \mathbb{N} \qquad\qquad (66 - 68)$$

$$J = 5: \#\hat{J}_5 = 2; (.)_{\hat{J}_5} = \left[(.)_{\hat{J}_{5,1}}, (.)_{\hat{J}_{5,2}} \right]^{\text{T}} \subset \mathbb{N}^2 \qquad\qquad (69 - 71)$$

9   Constraint 11 (**Cs11**): We denote $\bar{\theta}_{\hat{J}_{J,s}} \in \bar{\boldsymbol{\theta}}_{\hat{J}_J} \subset \mathbb{N}^{\mathbb{J}}$ as the subset of a J-reactant dependent biochemical

10   network where $\bar{\theta}_{\hat{J}_{J,s}}$ is the number of reactants considered simultaneously,

$$\bar{\theta}_{\hat{J}_{J,s}} \stackrel{\text{def}}{=} s + 1 \text{ where } J \geq 4, s \geq 1 \text{ and } \mathbb{J} = \#\hat{J}_J \qquad (\mathbf{D25})$$

$$\bar{\boldsymbol{\theta}}(\hat{\mathbf{J}}_J) = \bar{\boldsymbol{\theta}}_{\hat{J}_J} = \left(\bar{\theta}_{\hat{J}_{J,1}}, \dots, \bar{\theta}_{\hat{J}_{J,\mathbb{J}}} \right)^{\text{T}} \subset \mathbb{N}^{\mathbb{J}} \text{ [From } \mathbf{Cs10}] \quad (72), (73)$$

Examples:

$$J = 4: \#\hat{J}_4 = 1; \bar{\boldsymbol{\theta}}_{\hat{J}_4} = \left[\bar{\theta}_{\hat{J}_{4,1}} = 2 \right]^{\text{T}} \qquad\qquad (74 - 76)$$

$$J = 5: \#\hat{J}_5 = 2; \bar{\boldsymbol{\theta}}_{\hat{J}_5} = \left[\bar{\theta}_{\hat{J}_{5,1}} = 2, \bar{\theta}_{\hat{J}_{5,2}} = 3 \right]^{\text{T}} \qquad (77 - 79)$$

12   Constraint 12 **(Cs12):** We denote $\hat{I}_{\hat{J}_{J,s}} \in \hat{\mathbf{I}}_{\hat{J}_J} \subset \mathbb{N}^{\#\hat{J}_J}$ as the subset of a J-reactant dependent biochemical

13   network where $\hat{I}_{\hat{J}_{J,s}}$ is the number of selected PRVs,

$$\left(\hat{I}_{\hat{j}_{J,s}} \stackrel{def}{=} \binom{J}{\overline{\theta}_{\hat{j}_{J,s}}} = \frac{J!}{\overline{\theta}_{\hat{j}_{J,s}}! \left(J - \overline{\theta}_{\hat{j}_{J,s}} \right)!} = \frac{J \cdot (J-1) \dots \left(J - \overline{\theta}_{\hat{j}_{J,s}} + 1 \right) \cdot \left(J - \overline{\theta}_{\hat{j}_{J,s}} \right)!}{\overline{\theta}_{\hat{j}_{J,s}}! \left(J - \overline{\theta}_{\hat{j}_{J,s}} \right)!} = \right.$$
$$\left. = \frac{J \cdot (J-1) \dots \left(J - \overline{\theta}_{\hat{j}_{J,s}} + 1 \right)}{\overline{\theta}_{\hat{j}_{J,s}}!} = \left(\frac{\prod_{s=1}^{s=\#\overline{\theta}_{\hat{j}_J}} \left(J - \left(\overline{\theta}_{\hat{j}_{J,s}} - s \right) \right)}{\overline{\theta}_{\hat{j}_{J,s}}!} \right) \right)$$

(D26), (80 − 83)

where $J \geq 4, s \geq 1, \mathbb{J} = \#\hat{j}_J$ and $\hat{I}(\hat{j}_J) = \hat{I}_{\hat{j}_J} = \left(\hat{I}_{\hat{j}_{J,1}}, \dots, \hat{I}_{\hat{j}_{J,\mathbb{J}}} \right)^T \subset \mathbb{N}^{\mathbb{J}}$ [From **Cs10**] (84)

Examples:

$$J = 4; \#\hat{j}_4 = 1; \hat{I}_{\hat{j}_4} = \left[\hat{I}_{\hat{j}_{4,1}} = 6 \right]^T$$ (85 − 87)

$$J = 5; \#\hat{j}_5 = 2; \hat{I}_{\hat{j}_5} = \left(\hat{I}_{\hat{j}_{5,1}} = 10, \hat{I}_{\hat{j}_{5,2}} = 10 \right)$$ (88 − 90)

Constraint 13 (**Cs13**): A subset of i-indexed **r**-PRVs $\left(\mathbf{r}_{i=1}, \dots, \mathbf{r}_{i=\#\hat{I}_{\hat{j}_{J,s}}} \right)$ will form a stoichiometry number

matrix $\tilde{\mathcal{S}}(.)_{\hat{j}_{J,s}}$,

$$\left(\left[\mathbf{r}_{i=1}, \dots, \mathbf{r}_{i=\#\hat{I}_{\hat{j}_{J,s}}} \right]^T \in \mathbb{Z}^{J \cdot \left(\hat{I}_{\hat{j}_{J,s}} \right) \times 1} = \left[\left[m_{1,1}, \dots, m_{\overline{\theta}_{\hat{j}_{J,s}},1} \right]^T, \dots, \left[m_{1,\hat{I}_s}, \dots, m_{\overline{\theta}_{\hat{j}_{J,s}},\hat{I}_{\hat{j}_{J,s}}} \right]^T \right]^T \right.$$
$$\left. = \mathbb{Z}^J \otimes \mathbb{Z}^{\hat{I}_{\hat{j}_{J,s}}} = \begin{bmatrix} m_{1,1} & \cdots & m_{1,\hat{I}_{\hat{j}_{J,s}}} \\ \vdots & \ddots & \vdots \\ m_{J,1} & \cdots & m_{J,\hat{I}_{\hat{j}_{J,s}}} \end{bmatrix}_{J \times \hat{I}_{\hat{j}_{J,s}}} \stackrel{def}{=} \tilde{\mathcal{S}}(.)_{\hat{j}_{J,s}} \in \mathbb{Z}^{J \times \hat{I}_{\hat{j}_{J,s}}} \right)$$

(D27), (91 − 94)

where $J \geq 4, s \geq 1, \mathbb{J} = \#\hat{j}_J$ and $\left\{ \overline{\theta}_{\hat{j}_{J,s}} \in \overline{\theta}_{\hat{j}_J}, \hat{I}_{\hat{j}_{J,s}} \in \hat{I}_{\hat{j}_J} \right\} \subset \mathbb{N}^{\mathbb{J}}$ (95)

Constraint 14 (**Cs14**): The rank of $\tilde{\mathcal{S}}(.)_{\hat{j}_{J,s}}$ corresponds to the number of $\hat{I}_{\hat{j}_{J,s}}$-PRVs,

$$\text{Rank } \tilde{\mathcal{S}}(.)_{\hat{j}_{J,s}} \stackrel{def}{=} \hat{I}_{\hat{j}_{J,s}} \qquad \text{[From Cs2, Cs3]} \quad (D28)$$
$$\text{where } J \geq 4, s \geq 1 \text{ and } \tilde{\mathcal{S}}(.)_{\hat{j}_{J,s}} \in \mathbb{Z}^{J \times \hat{I}_{\hat{j}_{J,s}}}$$

Whilst **Cs10**- **Cs14** result in a generic matrix $\tilde{\mathcal{S}}(.)_{\hat{j}_{J,s}} \in \mathbb{Z}^{J \times \hat{I}_{\hat{j}_{J,s}}}$ where $s \geq 1$, the choice of reactants $(s +$

1, non-zero, $\overline{\theta}_{\hat{j}_{J,s}}$; combinatorial, without replacement) in tandem with the corresponding $\hat{I}_{\hat{j}_{J,s}}$-PRVs will

result in different outcomes or stoichiometry number matrices.

Constraint 15 (**Cs15**): We denote $\mathfrak{S}_{\hat{j}_{J,s}}$ as a class-specific subset of x-indexed $\tilde{\mathcal{S}}(.)_{\hat{j}_{J,s,1}}, \dots, \tilde{\mathcal{S}}(.)_{\hat{j}_{J,s,\#\mathfrak{S}_{\hat{j}_{J,s}}}}$-

distinct stoichiometry number matrices for a J-reactant dependent biochemical network with $\overline{\theta}_{\hat{j}_{J,s}}$-

reactants taken concurrently, without replacement and in tandem with $\hat{I}_{\hat{j}_{J,s}}$-PRVs,

$$\mathfrak{S}_{\hat{\jmath}_{J,s}} \stackrel{\text{def}}{=} \left(\tilde{\mathcal{S}}(.)_{\hat{\jmath}_{J,s,1}}, \ldots, \tilde{\mathcal{S}}(.)_{\hat{\jmath}_{J,s,\#\mathfrak{S}_{\hat{\jmath}_{J,s}}}} \right) \subseteq \mathbb{Z}^{J \times \hat{I}_{J,s}} \tag{D29}$$

$$\text{where } \#\mathfrak{S}_{\hat{\jmath}_{J,s}} \stackrel{\text{def}}{=} X_{\hat{\jmath}_{J,s}} = \begin{pmatrix} \hat{I}_{\hat{\jmath}_{J,s}} \\ \bar{\theta}_{\hat{\jmath}_{J,s}} \end{pmatrix} = \left(\frac{\prod_{s=1}^{s=\#\bar{\theta}_{\hat{\jmath}_J}} \left(\hat{I}_{\hat{\jmath}_{J,s}} - \left(\bar{\theta}_{\hat{\jmath}_{J,s}} - s \right) \right)}{\bar{\theta}_{\hat{\jmath}_{J,s}}!} \right) \text{ [From Cs13, Cs14]} \tag{D30}, (96)$$

$$\text{and } J \geq 4, s \geq 1, \mathbb{J} = \#\hat{\jmath}_J, \left\{ \bar{\theta}_{\hat{\jmath}_{J,s}} \in \bar{\boldsymbol{\theta}}_{\hat{\jmath}_J}, \hat{I}_{\hat{\jmath}_{J,s}} \in \hat{\mathbf{I}}_{\hat{\jmath}_J} \right\} \subset \mathbb{N}^{\mathbb{J}}$$

2   In order to explore the definitions discussed thus far, we present several numerical case studies for

3   greater clarity (**Table 1**). We now present some preliminary results on the basis of **Cs1- Cs15.**

4   Theorem 1 (**T1; STx1: Eqs. (1-7)**): There exist x-indexed $X_{\hat{\jmath}_{J,s,x}}$-membered subset(s) of stoichiometry

5   number matrices for a J-dependent biochemical network $\left(\tilde{\mathcal{S}}(.)_{\hat{\jmath}_{J,s,x}} \in \mathfrak{X}_J \right)$ and for $\bar{\theta}_{\hat{\jmath}_{J,s}}$-reactants

6   (concurrent, without replacement) and in tandem with $\hat{I}_{\hat{\jmath}_{J,s}}$-PRVs,

7
$$\mathfrak{X}_J \stackrel{\text{def}}{=} \left\{ \left(\bigcup_{s=1}^{s=\mathbb{J}} \bigcup_{x=1}^{x=X_{\hat{\jmath}_{J,s}}} \tilde{\mathcal{S}}(.)_{\hat{\jmath}_{J,s,x}} \right) - \left(\bigcap_{s=1}^{s=\mathbb{J}} \bigcap_{x=1}^{x=X_{\hat{\jmath}_{J,s}}} \tilde{\mathcal{S}}(.)_{\hat{\jmath}_{J,s,x}} \right) \right\} \tag{D31}$$
$$\text{where } J \geq 4, s \geq 1 \text{ and } \mathbb{J} = \#\hat{\jmath}_J$$

8   Corollary 1 (**C1; without proof**): The stoichiometry number matrices that comprise the superset of J-

9   dependent biochemical networks $\left(\tilde{\mathcal{S}}(.)_{\hat{\jmath}_{J,s,x}} \in \mathfrak{X}_J \right)$ are non-overlapping, unique and unequal (From **T1**),

10
$$\mathfrak{X}_J \stackrel{\text{def}}{=} \left\{ \mathfrak{S}_{\hat{\jmath}_{J,s=1}}, \ldots, \mathfrak{S}_{\hat{\jmath}_{J,s=\mathbb{J}}} \right\} = \left\{ \left(\bigcup_{s=1}^{s=\mathbb{J}} \bigcup_{x=1}^{x=X_{\hat{\jmath}_{J,s}}} \tilde{\mathcal{S}}(.)_{\hat{\jmath}_{J,s,x}} \right) - \emptyset \right\} \tag{D32}, (97)$$
$$= \left\{ \left(\tilde{\mathcal{S}}(.)_{\hat{\jmath}_{J,1,1}} \neq \cdots \neq \tilde{\mathcal{S}}(.)_{\hat{\jmath}_{J,1,X_{\hat{\jmath}_{J,s}}}} \right), \ldots, \left(\tilde{\mathcal{S}}(.)_{\hat{\jmath}_{J,\mathbb{J},1}} \neq \cdots \neq \tilde{\mathcal{S}}(.)_{\hat{\jmath}_{J,\mathbb{J},X_{\hat{\jmath}_{J,s}}}} \right) \right\}$$
$$\text{where } J \geq 4, s \geq 1 \text{ and } \mathbb{J} = \#\hat{\jmath}_J$$

11   Corollary 2 (**C2; without proof**): A J-reactant dependent superset $\left(\mathfrak{X}_J \right)$ can be expressed as a vector of

12   the independent classes of stoichiometry number matrices that comprises it $\left(\mathbf{z}_{\mathfrak{X}_J} \right)$,

13
$$\mathfrak{X}_{\hat{\jmath}} \times \mathbb{I}_{\mathbb{J}} \stackrel{\text{def}}{=} \mathbf{z}_{\mathfrak{X}_J} = \left(\sum_{s=1}^{s=\mathbb{J}} X_s . \widehat{s1} \ldots \sum_{s=1}^{s=\mathbb{J}} X_s . \widehat{s\mathbb{J}} \right)^T \in \mathbb{N}^{1 \times \mathbb{J}} \tag{D33}, (98)$$
$$\left(\begin{array}{c} \Rightarrow \left\| \mathbf{z}_{\mathfrak{X}_J} \right\|_1 \stackrel{\text{def}}{=} \#\mathfrak{X}_J = \left(\sum_{s=1}^{s=\mathbb{J}} X_s . \widehat{s1} + \cdots + \sum_{s=1}^{s=\mathbb{J}} X_s . \widehat{s\mathbb{J}} \right)^T \\ = \left(X_{\hat{\jmath}_{J,1}} + \cdots + X_{\hat{\jmath}_{J,\mathbb{J}}} \right)^T = \sum_{s=1}^{s=\mathbb{J}} X_{\hat{\jmath}_{J,s}} \end{array} \right) \tag{D34}, (99)$$
$$\text{where } \mathfrak{X}_{\hat{\jmath}} \stackrel{\text{def}}{=} \left(X_{\hat{\jmath}_{J,1}}, \ldots, X_{\hat{\jmath}_{J,\mathbb{J}}} \right)^T \in \mathbb{N}^{1 \times \mathbb{J}} \text{ and } \mathbb{I}_{\mathbb{J}} \subset \mathbb{N}^{\mathbb{J} \times \mathbb{J}} \tag{100}, (101)$$
$$\text{Here } \widehat{s1} = [1 \ 0 \ldots 0]^T, \ldots, \widehat{s\mathbb{J}} = [0 \ 0 \ldots 1]^T \text{ and } \mathbb{J} = \#\hat{\jmath}_J$$

1   Theorem 2 (**T2**; **without proof**): The J-reactant dependent superset $\mathfrak{X}\left(\mathbf{z}_{\mathfrak{X}_J}\right)$ of stoichiometry number

2   matrices can be expressed as a unit vector $\left(\hat{\mathbf{z}}_{\mathfrak{X}_J}\right)$,

$$\hat{\mathbf{z}}_{\mathfrak{X}_J} \stackrel{\text{def}}{=} \frac{\mathbf{z}_{\mathfrak{X}_J}}{\left\|\mathbf{z}_{\mathfrak{X}_J}\right\|_2} \tag{D35}$$

3   Here, $\left\|\hat{\mathbf{z}}_{\mathfrak{X}_J}\right\|_2 = \left\|\dfrac{\mathbf{z}_{\mathfrak{X}_J}}{\left\|\mathbf{z}_{\mathfrak{X}_J}\right\|_2}\right\|_2 = \left\|\dfrac{X_{s=1}}{\left\|\mathbf{z}_{\mathfrak{X}_J}\right\|_2} \cdot \widehat{s1} + \cdots + \dfrac{X_{s=\mathbb{J}}}{\left\|\mathbf{z}_{\mathfrak{X}_J}\right\|_2} \cdot \widehat{s\mathbb{J}}\right\|_2 \approx 1$ (102), (103)

and $\left\|\mathbf{z}_{\mathfrak{X}_J}\right\|_2 = \left(\sum_{s=1}^{s=\mathbb{J}} X_s^{\,2}\right)^{\frac{1}{2}}$ (104)

where $\mathbf{z}_{\mathfrak{X}_J} \stackrel{\text{def}}{=} X_{\hat{J}_{J,1}} \cdot \widehat{s1} + \cdots + X_{\hat{J}_{J,\mathbb{J}}} \cdot \widehat{s\mathbb{J}}, \widehat{s1} = [1\ 0\ \dots 0]^T, \widehat{s2} = [0\ 1\ \dots 0]^T, \widehat{s\mathbb{J}} = [0\ 0\ \dots 1]^T, \mathbb{J} = \#\hat{J}_J$ (105)

4   The algebraic model presented is J-reactant dependent and considers reactants as "siphons" where the

5   contributions of the participating reactants are described as intermittent (**Table 1**) [28-30].

6   Constraint 16 (**Cs16**): The set of J-reactant dependent stoichiometry number matrices is an $\mathbb{x}$-indexed

7    $(\mathbb{x} = 1, \dots, \mathbb{X})$ $\mathbb{X}$-membered $\left(\tilde{\mathcal{S}}(.)_{J,\mathbb{x}=1}, \dots, \tilde{\mathcal{S}}(.)_{J,\mathbb{x}=\mathbb{X}}\right)$ ordered set of constrained, unique and unequal

8   stoichiometry number matrices [**From Cs15, T1, C1, C2**],

$$\mathfrak{X}_J \stackrel{\text{def}}{=} \left(\tilde{\mathcal{S}}(.)_{J,\mathbb{x}=1}, \dots, \tilde{\mathcal{S}}(.)_{J,\mathbb{x}=\mathbb{X}}\right) \subset \mathbb{Z}^{J\times \max(\hat{I}_{\mathbb{x}})} \text{ where } \tilde{\mathcal{S}}(.)_{J,\mathbb{x}=1} \neq \cdots \neq \tilde{\mathcal{S}}(.)_{J,\mathbb{x}=\mathbb{X}} \tag{D36}$$

$$\mathbb{x} = 1, \dots, \mathbb{X}; \; \mathbb{X} \equiv \#\mathfrak{X}_J = \sum_{x=1}^{x=J-3} \begin{pmatrix} \hat{I}_x = \binom{J}{x+1} \\ \hat{J}_x = x+1 \end{pmatrix} \text{and } J \geq 4 \tag{106 - 108}$$

We simplify the notation without any loss of generality,

$$\text{If}\left(\tilde{\mathcal{S}}(.)_{J,\mathbb{x}} \in \mathfrak{X}_J \subset \mathbb{Z}^{J\times\max(\hat{I}_{\mathbb{x}})}\right) \text{ then } \left(\tilde{\mathcal{S}}(.)_{J,\mathbb{x}} \equiv \tilde{\mathcal{S}}_{\wp_{J,\mathbb{x}}}\right) \quad \textbf{(D37)}$$

2.2.3   Constraints to describe and incorporate indices of biochemical function

Whilst the Constraints described vide supra will ensure that our algebraic framework retains

biochemical relevance, we will, nonetheless require direct indices of biochemical function by which we

can model and characterize network-redundancy.

Constraint 17 (**Cs17**): The null space for any $\mathbb{x}$-stoichiometry number matrix $N\left(\tilde{\mathcal{S}}_{\wp_{J,\mathbb{x}}}\right)$ represents a

biochemical network at equilibrium,

$$\left\{ \mathbf{v}_k \in \mathrm{N}\left(\tilde{\mathcal{S}}_{\wp_{J,\mathbb{x}}}\right) \subset \mathbb{R}^{\hat{\mathbb{I}}_\mathbb{x}} | \mathbf{v}_k{}^{\mathbf{T}} \times \tilde{\mathcal{S}}_{\wp_{J,\mathbb{x}}} = \vec{\mathbf{0}}^{\mathbf{T}} \right\} \qquad \textbf{(D38)}$$

$$\text{where } \mathbf{v}_k \neq \vec{\mathbf{0}}, \mathbf{v}_k \neq \mathbf{v}_l | \mathbf{v}_k, \mathbf{v}_l \in \mathrm{N}\left(\tilde{\mathcal{S}}_{\wp_{J,\mathbb{x}}}\right) ; k, l \in \mathbb{N} \quad (109-112)$$

$$\tilde{\mathcal{S}}_{\wp_{J,\mathbb{x}}} \in \mathfrak{X}_J \subset \mathbb{Z}^{J \times \max(\hat{\mathbb{I}}_\mathbb{x})}; \mathbb{x} \in [1, \mathbb{X}]; \mathbb{X} = \#\mathfrak{X}_J; J \geq 4$$

Biochemical networks can be "trivially"-perturbed as in sequestrations (complex formation, membrane-

bound compartmentalization), post-translational modifications (acetylation, glycosylation, phosphorylation,

etc.) and single nucleotide polymorphism (SNPs) variants of a reactant. Our rationale to preclude the

"non-trivial" subset (deleterious mutations) is that these events are extremely rare and will alter the

biochemical network irrevocably disallowing any meaningful comparison [7, 43].

Constraint 18 (**Cs18**): The $\mathbb{x}^{\text{th}}$-stoichiometry number matrix $\left(\tilde{\mathcal{S}}_{\wp_{J,\mathbb{x}}}\right)$ is perturbed $\left(\tilde{\mathcal{S}}_{\wp_{J,\mathbb{x}}} \to \tilde{\mathcal{S}}_{\tilde{\wp}_{J,\mathbb{x}}}\right)$ with ρ-

PRVs and be studied by comparing the number and proportion of shared pairs of PRVs,

$$\text{If} \left(\begin{array}{l} \tilde{\mathbb{I}}_\mathbb{x} := \text{PRVs specific to } \tilde{\mathcal{S}}_{\tilde{\wp}_{J,\mathbb{x}}} \\ \text{where } \tilde{\mathbb{I}}_\mathbb{x} \stackrel{\text{def}}{=} \hat{\mathbb{I}}_\mathbb{x} + \rho; \rho \geq 1; \rho \in \mathbb{N}, \mathbb{x} \in [1, \mathbb{X}] \text{ and } \mathbb{X} = \#\mathfrak{X}_J \end{array} \right)$$

$$\text{then } \left(\left(\hat{\mathbf{r}}_{\mathbb{x}_{i=1}}, \tilde{\mathbf{r}}_{\mathbb{x}_{i=1}}\right), \dots, \left(\hat{\mathbf{r}}_{\mathbb{x}_{i=\hat{\mathbb{I}}_\mathbb{x}}}, \tilde{\mathbf{r}}_{\mathbb{x}_{i=\tilde{\mathbb{I}}_\mathbb{x}-\rho}}\right) \text{ where } \hat{\mathbf{r}}_{\mathbb{x}_i} \in \tilde{\mathcal{S}}_{\wp_{J,\mathbb{x}}}{}^{J \times 1}, \tilde{\mathbf{r}}_{\mathbb{x}_i} \in \tilde{\mathcal{S}}_{\tilde{\wp}_{J,\mathbb{x}}}{}^{J \times 1} \right) \quad \textbf{(D39)}, (113-116)$$

10   Constraint 19 (**Cs19**): There exists a bijection between the $\mathbb{x}^{\text{th}}$-stoichiometry number matrix (J-reactants,

11    $\hat{\mathbb{I}}_\mathbb{x}$-PRVs) and the $\mathbb{x}^{\text{th}}$-biochemical network (J-reactants, $\hat{\mathbb{I}}_\mathbb{x}$-reactions),

$$\text{If} \left(\mathbf{r}_{\mathbb{x}_i} \in \tilde{\mathcal{S}}_{\wp_{J,\mathbb{x}}}{}^{J \times 1} \subset \mathbb{N}^J \; \exists i \in \wp_{J,\mathbb{x}} \text{ s.t. } \mathbf{r}_{\mathbb{x}_i} \leftrightarrow i \, \forall i \right) \text{ then } \left(\tilde{\mathcal{S}}_{\wp_{J,\mathbb{x}}} \sim \wp_{J,\mathbb{x}} \right) \quad [\text{From } \textbf{Cs18}]$$

$$\text{where } \tilde{\mathcal{S}}_{\wp_{J,\mathbb{x}}} \in \mathfrak{X}_J \subset \mathbb{Z}^{J \times \max(\hat{\mathbb{I}}_\mathbb{x})}; \mathbb{x} \in [1, \mathbb{X}]; \mathbb{X} = \#\mathfrak{X}_J; J \geq 4 \qquad \textbf{(D40)}, (117)$$

13   Constraint 20 (**Cs20**): The probable dissociation constant for the i^{th}-reaction of an $\mathbb{x}$-specific

14   biochemical network $\left(\eta_i\left(\wp_{J,\mathbb{x}}\right)\right)$ can be classified as Boolean,

$$\eta_i\left(\wp_{J,\mathbb{x}}\right) \stackrel{\text{def}}{=} Kd_i \in \mathbb{R} \cap (0, +\infty) = ((b \vee e) \vee f) = ((b \vee f) \vee e) = (ne \vee e) \quad \textbf{(D41)}, (118-120)$$

$$\text{where } i \in [1, \hat{\mathbb{I}}_\mathbb{x}], \mathbb{x} \in [1, \mathbb{X}], \mathbb{X} = \#\mathfrak{X}_J \text{ and } J \geq 4$$

3   Methods

Constraints (**Cs1- Cs20**) and definitions (**D1-D41**), for a J-reactant dependent algebraic framework, will

result in a set of $\mathbb{x}$-indexed ($\mathbb{x} = 1, \dots, \mathbb{X}$) $\mathbb{X}$-unique stoichiometry number matrices $\left(\mathfrak{X}_J\right)$ where each $\mathbb{x}$-

stoichiometry number matrix corresponds to an $\mathbb{x}$-specific biochemical network $\left(\wp_{J,\mathbb{x}}\right)$ (**Cs19**). Since we

wish to model and characterize the redundant forms of a biochemical network, we seek $\mathbb{x}$-indexed

($\mathbb{x} = 1, \dots, \mathbb{X}$) $\mathbb{X}$-libraries of equivalent matrices where the $\mathbb{x}^{th}$-library can be mapped uniquely to the

$\mathbb{x}^{th}$-stoichiometry number matrix and studied.

3.1 Strictly positive non-unitary rational number scalar multipliers as generators of a library of equivalent matrices for the $\mathbb{x}^{th}$-biochemical network

Stoichiometry coefficients are numerical representations of the reaction conditions and can be used to

approximate the predicted/theoretical/assumed stoichiometry numbers for a set of reactants (**D1** and **D2**

(**Eqs. (1-5)**)). These can therefore, be fractions, negative numbers and zero, wherever applicable. Here,

we use strictly positive $\mathbb{Q}_+$ (stoichiometry preserving) and non-unitary rational numbers as scalar

multipliers for any $\tilde{\delta}_{p_{J,\mathbb{x}}} \in \mathfrak{X}_J \subset \mathbb{Z}^{J \times \max(\hat{I}_{\mathbb{x}})}$ to generate an $\mathbb{x}$-network specific library of equivalent

matrices. Although the set of rational numbers ($\mathbb{Q}$) is uncountable, we will restrict our consideration to

a countable infinite subset with dimensions $W \times Y$ where $W, Y \in \mathbb{N}$ (**Eqs. (121-155)**; **D42-D59**).

Theorem 3 (**T3**; **without proof**): The set of non-zero and strictly positive rational number multipliers

is modelled as the square matrix,

$$\boldsymbol{\Phi} \stackrel{\text{def}}{=} W \otimes Y = \phi(w, y) = (\phi)_{w,y} = \begin{cases} \left(\dfrac{w}{y}\right) \in LT(\boldsymbol{\Phi}) | w > y \\ \left(\dfrac{y}{w}\right) \in UT(\boldsymbol{\Phi}) | w < y \\ \left(\left(\dfrac{w}{w}\right) \vee \left(\dfrac{y}{y}\right)\right) \in D(\boldsymbol{\Phi}) | w = y \end{cases} \subset \left(\mathbb{Q}_+ \backslash \{1\}\right)^{W \times Y} \quad (\textbf{D42}), (121 - 123)$$

where,
rows($\boldsymbol{\Phi}$) = 1, ... , W   (124)
cols($\boldsymbol{\Phi}$) = 1, ... , Y   (125)
UT($\boldsymbol{\Phi}$) := Upper triangular matrix   (126)
LT($\boldsymbol{\Phi}$) := Lower triangular matrix   (127)
D($\boldsymbol{\Phi}$) := Diagonal elements   (128)

15   Corollary 3 (**C3; STx1: Eq. (8-20)**): The matrix of non-zero non-unitary rational number multipliers

16    $\left(\boldsymbol{\Phi} \subset \left(\mathbb{Q}_+ \backslash \{1\}\right)^{W \times Y}\right)$ is characterized by,

$$\#D(\boldsymbol{\Phi}) = (W) \vee (Y) \quad\quad\quad (\textbf{D43}), (129)$$

$$\#LT(\boldsymbol{\Phi}) = \left(\frac{W.(W-1)}{2}\right) \quad\quad (\textbf{D44}), (130)$$

$$\#UT(\boldsymbol{\Phi}) = \left(\frac{Y.(Y-1)}{2}\right) \quad\quad (\textbf{D45}), (131)$$

$$\#Vec(\boldsymbol{\Phi}) = (W^2) \vee (Y^2) \vee (W.Y) \quad (\textbf{D46}), (132)$$

$$\text{where } \#rows(\boldsymbol{\Phi}) = W, \#cols(\boldsymbol{\Phi}) = Y \quad (133), (134)$$

18   Corollaries 4-8 (**C4-C8; STx1: Eqs. (21-61)**): The following general results hold true for $\boldsymbol{\Phi}$,

$$\mathbf{C4}: (\mathbf{\Phi}^T)^T = \mathbf{\Phi} \qquad\qquad (\mathbf{D47}), (135)$$
$$\mathbf{C4}: q.\,\mathbf{\Phi}^T = (q.\,\mathbf{\Phi})^T \text{ s.t. } q \in \mathbb{R} - \{0\} \qquad (\mathbf{D48}), (136)$$
$$\mathbf{C6}: (\mathbf{\Phi}^p)^T = (\mathbf{\Phi}^T)^p \text{ s.t. } p \in \mathbb{R} \qquad (\mathbf{D49}), (137)$$
$$\mathbf{C7}: \mathbf{\Phi} \times \mathbf{\Phi}^T \ni \bar{\bar{\phi}}_{\widehat{w},\widehat{y}} = \bar{\bar{\phi}}_{\widehat{y},\widehat{w}} \text{ where } \widehat{w} \neq \widehat{y} \quad (\mathbf{D50}), (138)$$
$$\mathbf{C8}: \det(\mathbf{\Phi}) = \det(\mathbf{\Phi}^T) \qquad\qquad (\mathbf{D51}), (139)$$

2   Lemma 1 (**L1**; **without proof**): The scalar product of a transposable pair of elements of $\mathbf{\Phi}$ (off-

3   diagonal pair) is unity,

$$\left(\begin{array}{c} \text{if } \left(\phi_{w,y}, \phi_{y,w} \in \mathbf{\Phi} \subset \left(\mathbb{Q}_+ \backslash \{1\} \right)^{W \times Y} \text{ where } w \neq y \right) \\ \text{then } \phi_{w,y}.\,\phi_{y,w} = \phi_{w,y}.\,\phi_{w,y}^{-1} = 1 \end{array} \right) \quad \text{[From } \mathbf{D42}\text{]} \quad (\mathbf{D52}), (140), (141)$$

where,
$$w = 1, \dots, W \in \mathbb{N}, y = 1, \dots, Y \in \mathbb{N} \qquad\qquad (142), (143)$$

5   Theorem 4 (**T4; STx1: Eq. (62-71)**): The scalar product of all the off-diagonal elements of $\mathbf{\Phi}$ is

6   numerically equal to unity,

$$\mathrm{LT}(\mathbf{\Phi}).\,\mathrm{UT}(\mathbf{\Phi}) = 1 \qquad\qquad (144)$$
where
$$\mathrm{LT}(\mathbf{\Phi}) = \prod_{w=2}^{w=W} \prod_{y=1}^{y=w-1} \phi_{w,y} \qquad (145)$$

$$\mathrm{UT}(\mathbf{\Phi}) = \prod_{y=2}^{y=Y} \prod_{w=1}^{w=y-1} \phi_{w,y} \qquad (146)$$

$$\text{and } \phi_{w,y} \in \mathbf{\Phi} \subset \left(\mathbb{Q}_+ \backslash \{1\} \right)^{W \times Y} \text{ s.t. } w \neq y$$

8   If we choose only off-diagonal elements of $\left(\mathbb{Q}_+ \backslash \{1\} \right)^{W \times Y}$ as rational number multipliers for $\tilde{\delta}_{\wp_{J,x}} \in$

9     $\mathfrak{X}_J \subset \mathbb{Z}^{J \times \max(\hat{I}_x)}$, we can ensure the bijection with an x-biochemical network-specific a-indexed $\mathbb{A}$-

10   membered library of unique and equivalent rational number matrices $\mathcal{S}_{\wp_{J,x,a}} \in \mathfrak{C}_{J,x} \subset \mathbb{Q}\backslash\{-1,1\}^{J \times \max(\bar{I}_x)}$.

11   The rational number reaction vectors (RRVs; $\bar{I}_x$) for $\mathcal{S}_{\wp_{x_a}}$ are then simply, transformed PRVs ($\hat{I}_x$) of

12     $\tilde{\delta}_{\wp_{J,x}}$,

$$\#\hat{I}_x = \#\bar{I}_x \text{ s.t. } \hat{I}_x \in \mathbb{Z}^J, \bar{I}_x \in \left\{ \mathbb{Q}\backslash\{-1,1\} \right\}^J \quad (147), (\mathbf{D53}), (\mathbf{D54})$$
$$\text{and } J \geq 4$$

14   Theorem 5 (**T5**; **without proof**): A strictly positive and non-unitary rational number multiplier of

15     $\tilde{\delta}_{\wp_{J,x}} \in \mathfrak{X}_J \subset \mathbb{Z}^{J \times \max(\hat{I}_x)}$ will generate an equivalent pair of rational number matrices,

$$\text{if} \left(\begin{array}{l} \mathcal{S}_{\wp_{J,x,a}} = \left(\phi_{w,y} \right) . \tilde{\mathcal{S}}_{\wp_{J,x}} \text{ and } \mathcal{S}_{\wp_{J,x,b}} = \left(\phi_{y,w} \right) . \tilde{\mathcal{S}}_{\wp_{J,x}} \\ \text{where } \mathcal{S}_{\wp_{J,x,a}}, \mathcal{S}_{\wp_{J,x,b}} \in \mathfrak{C}_{J,x} \subset \mathbb{Q}\backslash\{-1,1\}^{J\times\max(\hat{I}_x)} \end{array} \right)$$

$$\text{then } \mathcal{S}_{\wp_{J,x,a}} \neq \mathcal{S}_{\wp_{J,x,b}} \text{ iff } \left(\phi_{w,y} \right) \neq 1 \text{ and } a \neq b \qquad \text{[From Cs17, T3, C3]} \quad (\textbf{D55}), (148-152)$$

$$\text{where } \phi_{w,y}, \phi_{y,w} \in \boldsymbol{\Phi} \subset \left(\mathbb{Q}_+\backslash\{1\} \right)^{W\times Y} \text{s. t. } w \neq y$$

$$\tilde{\mathcal{S}}_{\wp_{J,x}} \in \mathfrak{X}_J \subset \mathbb{Z}^{J\times\max(\hat{I}_x)} \text{ and } w, y, x, a, b \in \mathbb{N}$$

2 Theorem 6 (**T6; without proof**): Any $\tilde{\mathcal{S}}_{\wp_{J,x}} \in \mathfrak{X}_J \subset \mathbb{Z}^{J\times\max(\hat{I}_x)}$ where $x \in [1, \mathbb{X}]$ and $\mathbb{X} = \#\mathfrak{X}_J$ whence

3 multiplied by strictly positive and non-unitary rational number multipliers will result in a library of

4 equivalent rational number matrices for the x^{th}-biochemical network,

$$\left(\mathcal{S}_{\wp_{J,x,a=1}}, \dots, \mathcal{S}_{\wp_{J,x,a=\mathbb{A}}} \right) = \mathfrak{C}_{J,x} \subset \left(\mathbb{Q}\backslash\{-1,1\} \right)^{J\times\max(\bar{I}_x)} | \mathcal{S}_{\wp_{J,x,a}} = \left(\phi_{w,y} \right) . \tilde{\mathcal{S}}_{\wp_{J,x}} \qquad (\textbf{D56}), (153)$$

$$\Rightarrow \left(\phi_{w,y} \right) . \tilde{\mathcal{S}}_{\wp_{J,x}}, \dots, \left(\phi_{W,Y} \right) . \tilde{\mathcal{S}}_{\wp_{J,x}} \forall W. Y \qquad (154)$$

$$\overset{\text{def}}{=} \left(\mathcal{S}_{\wp_{J,x,a=1}} \neq \dots \neq \mathcal{S}_{\wp_{J,x,a=\mathbb{A}}} \right) \qquad \text{[From T5]} \qquad (\textbf{D57})$$

5 where,

$$a = \begin{cases} 2(|w - y|) \text{ iff } w - y > 0 \\ 2(|w - y|) - 1 \text{ iff } w - y < 0 \end{cases} \qquad (\textbf{D58})$$

$$\phi_{w,y} \in \left(\mathbb{Q}_+\backslash\{1\} \right)^{W\times Y}; w \neq y$$

$$w = 1, \dots, W; y = 1, \dots, Y; \mathbb{A} = W. Y \text{ and } x, w, y, \mathbb{A}, W, Y \in \mathbb{N} \qquad (155)$$

6 Corollary 9 (**C9; without proof**): For any $\tilde{\mathcal{S}}_{\wp_{J,x}} \in \mathfrak{X}_J \subset \mathbb{Z}^{J\times\max(\hat{I}_x)}$ we have a J-reactant dependent x-

7 library of a-indexed $\mathbb{A}$-redundant forms for the x^{th}-biochemical network,

$$\left(\mathcal{S}_{\wp_{J,x,a=1}} \neq \dots \neq \mathcal{S}_{\wp_{J,x,a=\mathbb{A}}} \right) = \mathfrak{C}_{J,x} \subset \left(\mathbb{Q}\backslash\{-1,1\} \right)^{J\times\max(\bar{I}_x)} \qquad \text{[From T5]}$$

8 $$\Rightarrow \wp_{J,x,a=1} \sim \dots \sim \wp_{J,x,a=\mathbb{A}} \qquad \text{[From Cs18, Cs19]} \quad (\textbf{D59})$$

$$\text{where } \tilde{\mathcal{S}}_{\wp_{J,x}} \in \mathfrak{X}_J \subset \mathbb{Z}^{J\times\max(\hat{I}_x)} \text{ and } x \in [1, \mathbb{X}]; \mathbb{X} = \#\mathfrak{X}_J; \mathbb{A} = W. Y; J \geq 4$$

3.2 Probable dissociation constants, as parameters, to interrogate specific biochemical pathways

We utilize "ReDirection" to conduct detailed computational studies on known and well-characterized

biochemical pathways [7]. "ReDirection" is an R-package that deploys a mathematically rigorous

algorithm to compute the probable dissociation constant and assign an outcome to every reaction of a

user-defined biochemical pathway [7, 8]. Briefly, "ReDirection" exhaustively checks a network-specific

stoichiometry number matrix for linear dependency (rows, columns), lower bounds (reactants, PRVs)

and reaction vector orientation (rows, columns) [7]. "ReDirection" then serially generates null space-

generated subspaces by removing all redundant and trivial vectors, populating a reaction-specific

sequence vector with numerical values drawn from each row and across all columns of this subspace

and thence summing ($\|.\|_1$) for the same [7, 8]. These steps are done iteratively and recursively until the following conditions are met,

$$\|.\|_1 \sim \begin{cases} \|+(.)\|_1 > 1 \\ \|-(.)\|_1 < -1 \\ \|+(.)\|_1 \approx \|-(.)\|_1 \cong 0 \end{cases} \quad (156)$$

For each iteration thereafter, "ReDirection" will compute reaction-specific sequence vector descriptors such as the arithmetic mean, standard deviation, etc., screen each term and bin them to outcome-specific subsets whose terms are, in turn, summed and mapped to a positive real number [7]. These constitute the i^{th}-reaction-specific output vector and its p1-norm is the probable dissociation constant $\left(\eta_i \in \mathbb{R} \cap (0, +\infty)\right)$ [7, 8]. The aforementioned steps are continued until every reaction ($\forall i$) has been assigned an unambiguous outcome in accordance with **Eq. (156)**.

3.3 Evaluating the consequences of introducing ρ-reactions into a specific biochemical pathway

The introduction of ρ-(reactions/RRVs) into an a-indexed redundant form of the x^{th}-biochemical network $\left(\mathscr{p}_{J,x,a}, \tilde{\mathscr{p}}_{J,x,a}\right)$ will be used as input to "ReDirection" since $\left(\mathscr{p}_{J,x,a}, \tilde{\mathscr{p}}_{J,x,a}\right) \leftrightarrow \left(S_{\mathscr{p}_{J,x,a}}, S_{\tilde{\mathscr{p}}_{J,x,a}}\right)$ (**A19, C9**) (**Figure 2**). The probable dissociation constants will be computed and used to unambiguously assign outcomes to every reaction. In order to ascertain the effects of $\rho \geq 1$-reactions, we will need to compute the number of shared reactions ($\tilde{I}_{J,x,a} - \rho$; **A19**, **Figure 2**). From this subset, we compute the number, distribution and proportion of matched (m)-reactions which are non-equivalent (mne) and equivalent (me) (**Eqs. (157-160))** (**Table 1; D60-D69**),

$$\#\text{ne}_{\left(\mathscr{p}_{J,x,a}, \tilde{\mathscr{p}}_{J,x,a}\right)} = \#\text{f}_{\left(\mathscr{p}_{J,x,a}, \tilde{\mathscr{p}}_{J,x,a}\right)} + \#\text{b}_{\left(\mathscr{p}_{J,x,a}, \tilde{\mathscr{p}}_{J,x,a}\right)} \quad (157)$$

$$\#\text{mne}_{\left(\mathscr{p}_{J,x,a}, \tilde{\mathscr{p}}_{J,x,a}\right)} = \#\text{mf}_{\left(\mathscr{p}_{J,x,a}, \tilde{\mathscr{p}}_{J,x,a}\right)} + \#\text{mb}_{\left(\mathscr{p}_{J,x,a}, \tilde{\mathscr{p}}_{J,x,a}\right)} \quad (158)$$

Combining (157) and (158) to derive the metrics,

$$\omega_{\left(\mathscr{p}_{J,x,a}, \tilde{\mathscr{p}}_{J,x,a}\right)} = \left(\left(\frac{\#\text{mf}_{\left(\mathscr{p}_{J,x,a}, \tilde{\mathscr{p}}_{J,x,a}\right)} + \#\text{mb}_{\left(\mathscr{p}_{J,x,a}, \tilde{\mathscr{p}}_{J,x,a}\right)}}{\#\text{f}_{\left(\mathscr{p}_{J,x,a}, \tilde{\mathscr{p}}_{J,x,a}\right)} + \#\text{b}_{\left(\mathscr{p}_{J,x,a}, \tilde{\mathscr{p}}_{J,x,a}\right)}}\right).(100)\right) = \left(\frac{\#\text{mne}_{\left(\mathscr{p}_{J,x,a}, \tilde{\mathscr{p}}_{J,x,a}\right)}}{\#\text{ne}_{\left(\mathscr{p}_{J,x,a}, \tilde{\mathscr{p}}_{J,x,a}\right)}}\right).(100) \quad (159)$$

$$\psi_{\left(\mathscr{p}_{J,x,a}, \tilde{\mathscr{p}}_{J,x,a}\right)} = \left(\frac{\#\text{me}_{\left(\mathscr{p}_{J,x,a}, \tilde{\mathscr{p}}_{J,x,a}\right)}}{\#\text{e}_{\left(\mathscr{p}_{J,x,a}, \tilde{\mathscr{p}}_{J,x,a}\right)}}\right).(100) \quad (160)$$

3.4 Computational studies to evaluate the consequences of introducing ρ-reactions into known biochemical pathways

Biochemical pathways for the urea cycle and folate metabolism will be analysed with "ReDirection" [7, 8]. These are well characterized, conserved across taxa and make significant contributions to the intracellular milieu and thereby to the physiological function of a cell.

The human urea cycle is an important pathway in the utilization and removal of non-protein nitrogen (ammonia) and occurs, partly, in the mitochondria and the cytosol of the liver and kidneys. The inability to excrete urea over a prolonged period of time predisposes an individual to varying grades of uremia and encephalopathy. The clinical relevance of the urea cycle notwithstanding, Ornithine, a critical intermediate of the urea cycle is a significant contributor to polyamine biosynthesis (Putrescine, Spermidine, Spermine) [50, 51]. Polyamines (PAs) contribute non-trivially to recombination-mediated DNA repair, inwardly-rectifying potassium channels (K_{ir}), cardiac conduction, cell proliferation and macrophage polarization [52-55]. The biochemical relevance of PAs is offset by their propensity to form free radicals *in vivo* and a non-trivial contribution to dysregulated cell growth [52-55]. Endogenous production of PAs is stringently regulated by the rate limiting enzymes Ornithine (EC 4.1.1.17)- and S-adenosylmethionine (EC 4.1.1.50)- decarboxylases [55, 56]. Here, we examine the effects of a multi-step and indirect reaction(s) (N-Acetylglutamate-, L-Glutamate-, 4-hydroxyglutamate-semialdehydes; 1-pyrroline-5- and 1-pyrroline-3-hydroxy-5-carboxylates) from Arginine, Proline and Glutamate (r_{12}) on an integrated biochemical network of urea metabolism and polyamine biosynthesis. Folate metabolism connects disparate reactants (macromolecular, micronutrients) such as proteins, carbohydrates, lipids, DNA/RNA and vitamins. The "Folate trap" is a clinicopathological condition that is characterized by relative folate deficiency in the presence of vitamin B12 deficiency [57, 58]. Since folate is an important contributor to 1C-metabolism and DNA synthesis, its deficiency will result in macrocytic anemia (large and immature RBCs). In its role as a cofactor, folate is required by enzymes involved in the metabolism of several amino acids (Serine, Glycine, Histidine, Cysteine) [57, 58]. The direct conversion of dUMP to dTMP (r_{18}) is likely to alter the flux of various reactants in an integrated biochemical network for folate and methionine metabolism and will be investigated in this study. Thymidine is required for DNA synthesis and the enzyme responsible for this conversion is thymidylate synthase (TS; EC 2.1.1.45) [58, 59]. Defects of this enzyme will result in the accumulation of dUMP and other deoxynucleotides which can damage DNA. Conversely, TS is a known contributor in the pathogenesis of hepatocellular carcinoma and development of chemoresistance [57-59].

4 Results

Here, we will investigate algebraic properties of the $\mathbb{x}^{th}$-biochemical network-specific library of

equivalent rational number matrices, discuss the implications of introducing and thence perturbing

known biochemical networks and evaluating parameter selection strategies to characterize network-level

redundant forms of a biochemical network.

4.1 Defining and elucidating the properties of a library of equivalent rational number matrices for
the $\mathbb{x}^{th}$-biochemical network

Corollaries -10 (**C10**) and -11 (**C11**) follow readily and is presented without formal proofs,

9
$$\textbf{C10:} \bigcap_{a=1}^{a=\mathbb{A}} \mathcal{S}_{\wp_{J,\mathbb{x},a}} = \emptyset \qquad\qquad \text{[From \textbf{T6}]} \quad (161)$$

$$\textbf{C11:} \mathfrak{C}_{J,\mathbb{x}} = \bigcup_{a=1}^{a=\mathbb{A}} \mathcal{S}_{\wp_{J,\mathbb{x},a}} - \bigcap_{a=1}^{a=\mathbb{A}} \mathcal{S}_{\wp_{J,\mathbb{x},a}} = \bigcup_{a=1}^{a=\mathbb{A}} \mathcal{S}_{\wp_{J,\mathbb{x},a}} \qquad \text{[From \textbf{C10}]} \quad (162)$$

$$\text{where } \left(\mathcal{S}_{\wp_{J,\mathbb{x},a=1}} \neq \cdots \neq \mathcal{S}_{\wp_{J,\mathbb{x},a=\mathbb{A}}} \right) = \mathfrak{C}_{J,\mathbb{x}} \subset (\mathbb{Q}\backslash\{-1,1\})^{J \times \max(\bar{I}_{\mathbb{x}})}$$

$$\text{and } \mathbb{x} \in [1, \mathbb{X}]; \mathbb{X} = \#\mathfrak{X}_J; \mathbb{A} = W.Y; J \geq 4$$

10 The null space is a numerical construct which is used to model a system of biochemical reactions at

11 equilibrium (**A15**). The null space-generated subspace has also been shown to be biochemically relevant

12 with the computation of the probable dissociation constant for every reaction of a user-defined

13 biochemical network [7, 8].

14 Theorem 7 (**T7; STx1: Eqs. (72-84)**): The null spaces generated by members of $\mathfrak{C}_{J,\mathbb{x}}$ are equal,

15
$$N\left(\mathcal{S}_{\wp_{J,\mathbb{x},a=1}}\right) = \cdots = N\left(\mathcal{S}_{\wp_{J,\mathbb{x},a=\mathbb{A}}}\right) \qquad\qquad (\textbf{D70})$$

$$\text{where } \left(\mathcal{S}_{\wp_{J,\mathbb{x},a=1}} \neq \cdots \neq \mathcal{S}_{\wp_{J,\mathbb{x},a=\mathbb{A}}} \right) = \mathfrak{C}_{J,\mathbb{x}} \subset (\mathbb{Q}\backslash\{-1,1\})^{J \times \max(\bar{I}_{\mathbb{x}})}$$

$$\text{and } \mathbb{x} \in [1, \mathbb{X}]; \mathbb{X} = \#\mathfrak{X}_J; \mathbb{A} = W.Y; J \geq 4$$

16 Theorem 8 (**T8; STx1: Eqs. (85-100)**): The linear map between $\mathfrak{C}_{J,\mathbb{x}}$ and the corresponding null space

17 is many-to-one,

18
$$f: \mathfrak{C}_{J,\mathbb{x}} \twoheadrightarrow N\left(\mathfrak{C}_{J,\mathbb{x}}\right) \quad (\textbf{D71})$$

19 In accordance with **T7** and **T8** we refine our definition of $\mathfrak{C}_{J,\mathbb{x}}$ to include this degeneracy (D) (**Figure**

20 **1**). In other words,

1 $$\text{if}\left(\mathfrak{C}_{J,x} = \left(\mathcal{S}_{\wp_{J,x,a=1}}, \dots, \mathcal{S}_{\wp_{J,x,a=\mathbb{A}}}\right) \subset (\mathbb{Q}\backslash\{-1,1\})^{J\times\max(\bar{I}_x)} \text{ s.t. } \mathcal{S}_{\wp_{J,x,a=1}} \neq \dots \neq \mathcal{S}_{\wp_{J,x,a=\mathbb{A}}}\right)$$
$$\text{then} \left(N\left(\mathcal{S}_{\wp_{J,x,a=1}}\right) = \dots = N\left(\mathcal{S}_{\wp_{J,x,a=\mathbb{A}}}\right)\right) \qquad \textbf{(D72)}, (163), (164)$$
$$\text{where } x \in [1,\mathbb{X}], \mathbb{X} = \#\mathfrak{X}_J, \mathbb{A} = W.Y \text{ and } J \geq 4$$

2 We will now present a refined definition for an a-indexed $\mathbb{A}$-membered library of constrained, rational

3 number, equivalent and redundant matrices for the x^{th}-biochemical network (CRaTERx),

4
$$\left(\mathcal{S}_{\wp_{J,x,a=1}}, \dots, \mathcal{S}_{\wp_{J,x,a=\mathbb{A}}}\right) = \mathfrak{C}_{J,x} \subset (\mathbb{Q}\backslash\{-1,1\})^{J\times\max(\bar{I}_x)} \,|\, \mathcal{S}_{\wp_{J,x,a}} = \left(\phi_{w,y}\right).\tilde{S}_{\wp_{J,x}} \quad \textbf{(D73)}$$
$$\text{where,}$$
$$\mathcal{S}_{\wp_{J,x,a=1}} \neq \dots \neq \mathcal{S}_{\wp_{J,x,a=\mathbb{A}}}$$
$$N\left(\mathcal{S}_{\wp_{J,x,a=1}}\right) = \dots = N\left(\mathcal{S}_{\wp_{J,x,a=\mathbb{A}}}\right)$$
$$\tilde{S}_{\wp_{J,x}} \in \mathfrak{X}_J \subset \mathbb{Z}^{J\times\max(\hat{I}_x)}; x \in [1,\mathbb{X}], \mathbb{X} = \#\mathfrak{X}_J \text{ and } J \geq 4$$
$$\phi_{w,y} \in \left(\mathbb{Q}_+\backslash\{1\}\right)^{W\times Y}; w \neq y$$
$$\text{and}$$
$$a = \begin{cases} 2(|w-y|) \text{ iff } w-y > 0 \\ 2(|w-y|)-1 \text{ iff } w-y < 0 \end{cases}$$
$$w = 1, \dots, W; y = 1, \dots, Y; \mathbb{A} = W.Y; x, w, y, \mathbb{A}, W, Y \in \mathbb{N}$$

5 Generic mathematical properties for $\mathfrak{C}_{J,x}$ include being rectangular $\left(4 \leq J < \max(\hat{I}_x); \textbf{Eq.}(\textbf{165})\right)$ and

6 comprising reaction vectors that are linearly dependent.

7 Lemma 2 (**L2; without proof**): The set $\mathfrak{C}_{J,x} \subset (\mathbb{Q}\backslash\{-1,1\})^{J\times\max(\bar{I}_x)}$ is commutative or Abelian with

8 regards to addition,

9
$$\mathcal{S}_{\wp_{J,x,a}} + \mathcal{S}_{\wp_{J,x,b}} = \mathbf{S}_{\wp_{J,x,(a+b)}} = \mathbf{S}_{\wp_{J,x,(b+a)}} = \mathcal{S}_{\wp_{J,x,b}} + \mathcal{S}_{\wp_{J,x,a}} \quad (166-168)$$
$$\text{where } \mathcal{S}_{\wp_{J,x,a}}, \mathcal{S}_{\wp_{J,x,b}} \in \mathfrak{C}_{J,x}; \left\{\mathbf{S}_{\wp_{J,x,a}}, \mathbf{S}_{\wp_{J,x,b}}\right\} \neq \mathbf{0}^{J\times\bar{I}_x} \qquad (169), (170)$$
$$\text{and } a, b \in [1, \mathbb{A}]; a \neq b; \mathbb{A} = W.Y; J \geq 4$$

10 Theorem 9 (**T9; STx1: Eqs. (101-108)**): The set $\mathfrak{C}_{J,x} \subset (\mathbb{Q}\backslash\{-1,1\})^{J\times\max(\bar{I}_x)}$ is a semigroup with

11 respect to addition and is conditionally Abelian,

12
$$g: \left(\mathfrak{C}_{J,x}, +\right) \to \mathfrak{C}_{J,x} \qquad (171)$$
$$\text{where } \mathcal{S}_{\wp_{J,x,a}}, \mathcal{S}_{\wp_{J,x,b}}, \mathcal{S}_{\wp_{J,x,c}} \in \mathfrak{C}_{J,x}; \left\{\mathbf{S}_{\wp_{J,x,a}}, \mathbf{S}_{\wp_{J,x,b}}, \mathcal{S}_{\wp_{J,x,c}}\right\} \neq \mathbf{0}^{J\times\max(\bar{I}_x)} \quad (172), (173)$$
$$\text{and } a, b, c \in [1, \mathbb{A}]; a \neq b \neq c; \mathbb{A} = W.Y; J \geq 4 \qquad (174-176)$$

13 **4.2 "ReDirection"-based analysis of known biochemical pathways**

14 We now complement our theoretical assertions with detailed computational studies on pathways for

15 urea synthesis and folate metabolism. Perturbations are introduced as single extraneous reactions and

the probable dissociation constants for each pair of shared reactions (unperturbed, perturbed) is

computed $\left(\eta_{i(\wp_{J,x,a})}, \tilde{\eta}_{i(\tilde{\wp}_{J,x,a})}\right)$ [7, 8].

The data, for the urea cycle and thence polyamine biosynthesis suggests that there is a decrease in the

absolute number of equivalent reactions (**Figures 3a and 3b, Table 3**). The increase in Glutamate

synthesis from Arginine ($\tilde{\eta}_{12} > 1$) will stimulate the synthesis of Ornithine ($\tilde{\eta}_8 \gg 1$) which will

condense with urea and form Arginine ($\tilde{\eta}_5 < 1$) (**Figures 3a and 3b, Table 2**). This, in turn, will

drive the urea cycle via the formation of Citrulline, Aspartate and Fumarate ($\tilde{\eta}_3, \tilde{\eta}_6, \tilde{\eta}_1, \tilde{\eta}_2 > 1$)

(**Figures 3a and 3b, Table 3).** These modifications are compounded by reductions in the probable

dissociation constants for the syntheses of Glutamate from Fumarate and thence Aspartate ($\tilde{\eta}_{11} <$

$\eta_{11}, \tilde{\eta}_7 < \eta_7$) (**Figures 3a and 3b, Table 3).** Our results also demonstrate a heightened synthesis of

Ornithine via Glutamate ($\tilde{\eta}_8 \approx 32$). This, we combined with the futile utilization via Citrulline

($\{\eta_9, \tilde{\eta}_9\} \approx 1.000$) and synthesis to Arginine ($\tilde{\eta}_5 \approx 0.032$) directly will ensure increased availability of

Ornithine and thence PAs (**Figures 3a and 3b, Table 3).**

We also observe and study the relevance of interconversion dUMP and dTMP ($\tilde{\eta}_{18} \approx 1.000$), in folate

metabolism, is equivalent and likely to be a key regulatory step. We also note that although the

proportion of equivalent and non-equivalent reactions remains unchanged, the magnitudes of the

probable dissociation constants differ significantly (**Figures 3c and 3d, Table 3**). Amongst the

reactions that exhibit a complete reversal are the reactants Cysteine, Alanine, Pyruvate and Glucose

($\tilde{\eta}_{13}, \tilde{\eta}_{10} < 1; \tilde{\eta}_{14}, \tilde{\eta}_{11} > 1$) (**Figures 3c and 3d, Table 3**). On the other hand, alterations in the

magnitude of the probable dissociation constants are also observed for reactions that involve

methionine, S-adenosylmethionine, Homocysteine, S-adenosylhomocysteine and the tetrahydrofolates

($\tilde{\eta}_1 > \eta_1, \tilde{\eta}_2 > \eta_2, \tilde{\eta}_3 > \eta_3, \tilde{\eta}_4 > \eta_4, \tilde{\eta}_5 > \eta_5, \tilde{\eta}_6 > \eta_6, \tilde{\eta}_7 > \eta_7$) (**Figures 3c and 3d, Table 3**). Since $1 >$

$\tilde{\eta}_4 > \eta_4$, a greater quantity of Homocysteine may be available for its reaction with 5-

methylenetetrahydrofolate ($\tilde{\eta}_1 \approx 31$) as opposed to the simple reversal to Methionine ($\eta_4 \approx 0.044$). The

ensuing reactions converts tetrahydrofolate to 5,10-methylenetetrahydrofolate ($\tilde{\eta}_1 \approx 31$) and thence to

dTMP via methylation of dUMP. A critical mediator favouring this step is the availability of Ribose

via Glucose, Alanine and Pyruvate (Glucose/Alanine/Cysteine, Cysteine/Glucose/Pyruvate) (**Figures 3c**

and 3d, Table 3). Since $1 > \tilde{\eta}_{13} > \eta_{14}$, the conversion of Glucose to Alanine ($\tilde{\eta}_{13} \approx 0.137$) is much

slower as compared to the same via Pyruvate ($\eta_{14} \approx 0.024$). This results a greater availability of

Glucose for Ribose synthesis and subsequent ribosylation of the de novo synthesized Thymine base

(Thymidine). This reactant is critical in the synthesis of the Uridine-Thymidine transversion and thence

DNA synthesis and a perturbation will shift the equilibrium towards compensatory synthesis (**Figures**

3c and 3d, Table 3; ST2, SD2).

4.3 Perturbation-based metrics to characterize the redundant forms of a biochemical network

Although the a^{th}-redundant form of an $\mathbb{x}^{\text{th}}$-biochemical network is easily evaluated (unperturbed,

perturbed) $\left\{ (.)(\mathbb{J}, \mathbb{x}, a) \sim (.)_{\left(\wp_{\mathbb{J},\mathbb{x},a}, \tilde{\wp}_{\mathbb{J},\mathbb{x},a} \right)} | \wp_{\mathbb{J},\mathbb{x},a} \sim \mathcal{S}_{\wp_{\mathbb{J},\mathbb{x},a}}, \tilde{\wp}_{\mathbb{J},\mathbb{x},a} \sim \mathcal{S}_{\tilde{\wp}_{\mathbb{J},\mathbb{x},a}}, a \in [1, \mathbb{A}] \right\}$ (**Eq. (177)**), we will need a

metric that compares $\mathbb{A}$-redundant forms of a biochemical network concurrently. Here, we will discuss

the derivation of metric which be used to compare any pair of redundant forms

$\left\{ (.)(\mathbb{J}, \mathbb{x}, (a,b)) \sim (.)_{\left(\wp_{\mathbb{J},\mathbb{x},a}, \wp_{\mathbb{J},\mathbb{x},b} \right)} | \wp_{\mathbb{J},\mathbb{x},a} \sim \mathcal{S}_{\wp_{\mathbb{J},\mathbb{x},a}}, \wp_{\mathbb{J},\mathbb{x},b} \sim \mathcal{S}_{\wp_{\mathbb{J},\mathbb{x},b}}, a, b \in [1, \mathbb{A}], a \neq b \right\}$ (**Eq. (178)**) (**Cs19**, **C9**). In

order to accomplish this, we will reformulate previously derived metrics into non-local (less sensitive,

numerically robust, specific) and local (greater sensitivity, reduced specificity, robustness)-versions (**D74**,

D75).

Let us compute confusion matrices for the final outcomes between pairs of shared reactions of the

unperturbed- and perturbed-forms of a biochemical network, which we will evaluate as parameters

$\left(\mathbb{T}^{1}_{\mathbb{J},\mathbb{x},a} := m^{\mathbb{T}1}e_{\left(\wp_{\mathbb{J},\mathbb{x},a}, \tilde{\wp}_{\mathbb{J},\mathbb{x},a} \right)} \sim \text{TP}; \mathbb{T}^{2}_{\mathbb{J},\mathbb{x},a} := m^{\mathbb{T}2}ne_{\left(\wp_{\mathbb{J},\mathbb{x},a}, \tilde{\wp}_{\mathbb{J},\mathbb{x},a} \right)} \sim \text{TP} \right)$ (**Eqs. (179-182)**) (**Table 1; D76-D83**),

$$\text{Case1:} \quad \text{Positive outcome} \quad := \text{Number of shared reactions with equivalent outcomes} \tag{179}$$

$$\text{Confusion matrix} \quad \overset{\text{def}}{=} \begin{bmatrix} m^{\mathbb{T}1}e_{\left(\wp_{\mathbb{J},\mathbb{x},a}, \tilde{\wp}_{\mathbb{J},\mathbb{x},a} \right)} = \text{TP} & u^{\mathbb{T}1}ne_{\left(\wp_{\mathbb{J},\mathbb{x},a}, \tilde{\wp}_{\mathbb{J},\mathbb{x},a} \right)} = \text{FN} \\ u^{\mathbb{T}1}e_{\left(\wp_{\mathbb{J},\mathbb{x},a}, \tilde{\wp}_{\mathbb{J},\mathbb{x},a} \right)} = \text{FP} & m^{\mathbb{T}1}ne_{\left(\wp_{\mathbb{J},\mathbb{x},a}, \tilde{\wp}_{\mathbb{J},\mathbb{x},a} \right)} = \text{TN} \end{bmatrix} = \mathbb{T}^{1}_{\mathbb{J},\mathbb{x},a} \subset \mathbb{Z}_{+}^{2 \times 2} \tag{180}$$

$$\text{Case2:} \quad \text{Positive outcome} \quad := \text{Number of shared reactions with non} - \text{equivalent outcomes} \tag{181}$$

$$\text{Confusion matrix} \quad \overset{\text{def}}{=} \begin{bmatrix} m^{\mathbb{T}2}ne_{\left(\wp_{\mathbb{J},\mathbb{x},a}, \tilde{\wp}_{\mathbb{J},\mathbb{x},a} \right)} = \text{TP} & u^{\mathbb{T}2}e_{\left(\wp_{\mathbb{J},\mathbb{x},a}, \tilde{\wp}_{\mathbb{J},\mathbb{x},a} \right)} = \text{FN} \\ u^{\mathbb{T}2}ne_{\left(\wp_{\mathbb{J},\mathbb{x},a}, \tilde{\wp}_{\mathbb{J},\mathbb{x},a} \right)} = \text{FP} & m^{\mathbb{T}2}e_{\left(\wp_{\mathbb{J},\mathbb{x},a}, \tilde{\wp}_{\mathbb{J},\mathbb{x},a} \right)} = \text{TN} \end{bmatrix} = \mathbb{T}^{2}_{\mathbb{J},\mathbb{x},a} \subset \mathbb{Z}_{+}^{2 \times 2} \tag{182}$$

18 The indices of accuracy (Ac), precision (Pr), recall (Re), false positive rate (FPr) and specificity

19 (SPy) will be computed (**Eqs. (183-190)**) (**Table 4**),

$$\text{Pr}_{(\wp_{J,\mathbb{x},a},\tilde{\wp}_{J,\mathbb{x},a})} \overset{\text{def}}{=} \left(\frac{\text{TP}}{\text{TP}+\text{FP}}\right).100 \tag{183}$$

$$\text{Re}_{(\wp_{J,\mathbb{x},a},\tilde{\wp}_{J,\mathbb{x},a})} \overset{\text{def}}{=} \left(\frac{\text{TP}}{\text{TP}+\text{FN}}\right).100 \tag{184}$$

$$\text{Ac}_{(\wp_{J,\mathbb{x},a},\tilde{\wp}_{J,\mathbb{x},a})} \overset{\text{def}}{=} \left(\frac{\text{TP}+\text{TN}}{\text{TP}+\text{FP}+\text{TN}+\text{FN}}\right).100 \tag{185}$$

$$\text{FPr}_{(\wp_{J,\mathbb{x},a},\tilde{\wp}_{J,\mathbb{x},a})} \overset{\text{def}}{=} \left(\frac{\text{FP}}{\text{FP}+\text{TN}}\right).100 \tag{186}$$

$$\text{SPy}_{(\wp_{J,\mathbb{x},a},\tilde{\wp}_{J,\mathbb{x},a})} \overset{\text{def}}{=} \left(1-\text{FPr}_{(\wp_{J,\mathbb{x},a},\tilde{\wp}_{J,\mathbb{x},a})}\right).100 = \left(1-\left(\frac{\text{FP}}{\text{FP}+\text{TN}}\right)\right).100 \tag{187}$$

where $\wp_{J,\mathbb{x},a}\sim\mathcal{S}_{\wp_{J,\mathbb{x},a}}; \tilde{\wp}_{J,\mathbb{x},a}\sim\mathcal{S}_{\tilde{\wp}_{J,\mathbb{x},a}}; \mathcal{S}_{\wp_{J,\mathbb{x},a}} \in \mathfrak{C}_{J,\mathbb{x}} \subset \mathbb{Q}\backslash\{-1,1\}^{J\times\max(\bar{I}_{\mathbb{x}})}$ $\quad(188-190)$
and $\mathbb{x} \in [1,\mathbb{X}]; \mathbb{X} = \#\mathfrak{X}_J; a \in [1,\mathbb{A}]; \mathbb{A} = \text{W.Y}$

We can rewrite $\mathbb{T}^1_{J,\mathbb{x},a}$ and $\mathbb{T}^2_{J,\mathbb{x},a}$ (**Eqs. (180), (182)**) in terms of the previously derived network-specific (**Eqs. (183-190)**) and generalize this as the bijection (g),

$$\text{g}: \mathbb{T}^{(.)}_{J,\mathbb{x},a}\sim\text{g}^{(.)}(.) \xleftarrow{\begin{array}{c}\left([\text{Pr Re Ac FPr SPy}]^T_{(\wp_{J,\mathbb{x},a},\tilde{\wp}_{J,\mathbb{x},a})}=\frac{1}{100}.[\text{Pr Re Ac FPr SPy}]^T_{(\wp_{J,\mathbb{x},a},\tilde{\wp}_{J,\mathbb{x},a})}\right)\\=(100^{-1}.[\text{Pr Re Ac FPr SPy}]^T)_{(\wp_{J,\mathbb{x},a},\tilde{\wp}_{J,\mathbb{x},a})}=\text{g}^{(.)}(\wp_{J,\mathbb{x},a},\tilde{\wp}_{J,\mathbb{x},a})\end{array}} \text{g}^{(.)}(J,\mathbb{x},a) \subset \mathbb{R}^5 \text{ (D84)}, (191-194)$$
where $\text{g}^{(.)} = 100^{-1}.[\text{Pr Re Ac FPr SPy}]^T$ $\tag{195}$

Let us now partition into non-local $\left(\text{g}^1(.) \in \mathcal{G}(\mathfrak{C}_{J,\mathbb{x}})\right)$- and local $\left(\text{g}^2(.) \in \mathcal{H}(\mathfrak{C}_{J,\mathbb{x}})\right)$-parameters to

evaluate and characterize the redundant forms of a biochemical network after a perturbation. Whilst the

former will evaluate the shift or lack thereof, for reactions with equivalent outcomes between pairs of

concurrently occurring redundant forms, the latter will characterize the alteration in flux of

reactants/products through reactions with non-equivalent outcomes (forward, reverse) for any single

redundant form. Paraphrasing the aforementioned assertions we can define,

$$\text{g}^{(.)}(.) \overset{\text{def}}{=} \begin{cases} \text{g}^1(.) \in \mathcal{G}(\mathfrak{C}_{J,\mathbb{x}}) \equiv \Delta e\{(\wp_{J,\mathbb{x},a},\tilde{\wp}_{J,\mathbb{x},a}),(\wp_{J,\mathbb{x},b},\tilde{\wp}_{J,\mathbb{x},b})\} \sim \text{Non}-\text{local} \\ \text{g}^2(.) \in \mathcal{H}(\mathfrak{C}_{J,\mathbb{x}}) \equiv \Delta ne(\wp_{J,\mathbb{x},a},\tilde{\wp}_{J,\mathbb{x},a}) \sim \text{Local} \end{cases} \text{ (D85)}, (196), (197)$$
where $\wp_{J,\mathbb{x},a}\sim\mathcal{S}_{\wp_{J,\mathbb{x},a}}, \wp_{J,\mathbb{x},b}\sim\mathcal{S}_{\wp_{J,\mathbb{x},b}}, \{\mathcal{S}_{\wp_{J,\mathbb{x},a}}, \mathcal{S}_{\wp_{J,\mathbb{x},b}}\} \in \mathfrak{C}_{J,\mathbb{x}} \subset \mathbb{Q}\backslash\{-1,1\}^{J\times\max(\bar{I}_{\mathbb{x}})}$
with $\mathbb{x} \in [1,\mathbb{X}]; \mathbb{X} = \#\mathfrak{X}_J; a,b \in [1,\mathbb{A}]; a \neq b; \mathbb{A} = \text{W.Y}$

Expanding each of these and rewriting **Eqs.**-(**198**) and -(**199**) in accordance with **D84**,

$$\text{g}^1_{(\wp_{J,\mathbb{x},a},\wp_{J,\mathbb{x},b})} = \text{g}^1_{\{(\wp_{J,\mathbb{x},a},\tilde{\wp}_{J,\mathbb{x},a}),(\wp_{J,\mathbb{x},b},\tilde{\wp}_{J,\mathbb{x},b})\}} = \text{g}^1(J,\mathbb{x},(a,b))\sim\text{g}^1\left(\mathcal{S}_{\wp_{J,\mathbb{x},a}},\mathcal{S}_{\wp_{J,\mathbb{x},b}}\right) \quad (198),(199),(\textbf{D86})$$

$$\text{g}^2_{(\wp_{J,\mathbb{x},a},\wp_{J,\mathbb{x},b})} = \text{g}^2_{\{(\wp_{J,\mathbb{x},a},\tilde{\wp}_{J,\mathbb{x},a}),(\wp_{J,\mathbb{x},b},\tilde{\wp}_{J,\mathbb{x},b})\}} = \text{g}^2(J,\mathbb{x},a),\text{g}^2(J,\mathbb{x},b)\sim\text{g}^2\left(\mathcal{S}_{\wp_{J,\mathbb{x},a}}\right),\text{g}^2\left(\mathcal{S}_{\wp_{J,\mathbb{x},b}}\right) \quad (200),(201),(\textbf{D87})$$

where $\wp_{J,\mathbb{x},a}\sim\mathcal{S}_{\wp_{J,\mathbb{x},a}}, \wp_{J,\mathbb{x},b}\sim\mathcal{S}_{\wp_{J,\mathbb{x},b}}, \{\mathcal{S}_{\wp_{J,\mathbb{x},a}}, \mathcal{S}_{\wp_{J,\mathbb{x},b}}\} \in \mathfrak{C}_{J,\mathbb{x}} \subset \mathbb{Q}\backslash\{-1,1\}^{J\times\max(\bar{I}_{\mathbb{x}})}$
with $\mathbb{x} \in [1,\mathbb{X}]; \mathbb{X} = \#\mathfrak{X}_J; a,b \in [1,\mathbb{A}]; a \neq b; \mathbb{A} = \text{W.Y}$

5 Discussion

The intracellular milieu is complex and any model must incorporate contributors to this complexity.

Our algebraic framework can be used to model and characterize the redundant forms of a biochemical

network with parameters that are both, theoretically sound and biochemically relevant.

5.1 Parameter selection to analyse the redundant-forms of a biochemical network

Although the outcome of reaction equivalence suggests data ambiguity for a reaction, inferring a

regulatory role for the same is equally plausible. Conversely, reactions with non-equivalent (forward,

reverse) outcomes are likely to represent the prevailing physicochemical conditions and may represent

the response elements of a biochemical network. These hypotheses were examined, numerically, *post*

hoc by the evaluation of $\mathbb{g}^{(.)}(.)$ for the unperturbed- and perturbed-forms of the urea cycle and folate

metabolism pathways (**D85**; **Table 4**).

The recall of a metric is a measure of how sensitive a pathway is to perturbation. Since these are

almost identical $\left(Re_{(Urea,Folate)} \approx 52\%\right)$ and of moderately low magnitude, the utilization of reaction

equivalence as the relevant non-local parameter is easily justified $\left(SPy_{(Urea,Folate)} \approx\right.$

$\left.75\%, Ac_{(Urea,Folate)} \approx 62\%\right)$ (**Table 4**). The precision or positive predictive value of a metric is an

indicator of robustness upon change. This, too, for an outcome of reaction equivalence is a useful

parameter to comprehend and compare biochemical pathways pre- and post-perturbation

$\left(Pr_{(Urea,Folate)} \approx 80\%\right)$ (**Table 4**). Interestingly, if the original confusion matrix was redefined with a

reaction outcome of non-equivalence as TP and reaction equivalence as TN $\left(\mathbb{T}^1_{J,\mathbb{x},a} \rightarrow \mathbb{T}^2_{J,\mathbb{x},a}\right)$ we get

different results (**Table 4**). Since reactions with non-equivalent outcomes are likely to be influenced by

the local intracellular milieu, this measure will be a poor choice whence comparing biochemical

networks $\left(Re_{(Urea,Folate)} \approx 75\%\right)$ (**Table 4**). However, since forward- and reverse-reactions regulate the

flux of the reactants/products through a biochemical network, outcomes of non-equivalence for the

majority of the reactions will result in a net direction for a pathway $\left(SPy_{(Urea.Folate)} \approx 52\%\right)$ rather

than robustness $\left(Pr_{(Urea,Folate)} \approx 49\%\right)$ (**Table 4**).

We conclude, from these studies, that reactions with an equivalent outcome may constitute a useful

metric of comparison between the redundant forms of a biochemical network whence taken

concurrently. On the other hand, since the flux of reactants/products through reactions with a non-

equivalent outcome (forward, reverse) is dependent on the prevailing physicochemical conditions, this

metric is better suited to characterizing an instance of a redundant form of a biochemical network.

Extending these results to $\mathbb{A}$-redundant forms of an J-reactant dependent $\mathbb{x}^{th}$-biochemical network

(CRaTER$\mathbb{x}$),

$$\mathcal{G}(\mathfrak{C}_{J,\mathbb{x}}) \stackrel{\text{def}}{=} \mathcal{G}\left(\mathcal{S}_{\wp_{J,\mathbb{x},a=1}}, \dots, \mathcal{S}_{\wp_{J,\mathbb{x},a=\mathbb{A}}}\right) \ni \left\{\mathbb{g}^1\left(\mathcal{S}_{\wp_{J,\mathbb{x},a}}, \mathcal{S}_{\wp_{J,\mathbb{x},b}}\right), \dots\right\} \tag{202}$$

$$\mathcal{H}(\mathfrak{C}_{J,\mathbb{x}}) \stackrel{\text{def}}{=} \mathcal{H}\left(\mathcal{S}_{\wp_{J,\mathbb{x},a=1}}, \dots, \mathcal{S}_{\wp_{J,\mathbb{x},a=\mathbb{A}}}\right) \ni \left\{\left(\mathbb{g}^2\left(\mathcal{S}_{\wp_{J,\mathbb{x},a}}\right), \dots, \mathbb{g}^2\left(\mathcal{S}_{\wp_{J,\mathbb{x},\mathbb{A}}}\right)\right), \dots\right\} \tag{203}$$

where $\wp_{J,\mathbb{x},a} \sim \mathcal{S}_{\wp_{J,\mathbb{x},a}}, \wp_{J,\mathbb{x},b} \sim \mathcal{S}_{\wp_{J,\mathbb{x},b}}, \left\{\mathcal{S}_{\wp_{J,\mathbb{x},a}}, \mathcal{S}_{\wp_{J,\mathbb{x},b}}\right\} \in \mathfrak{C}_{J,\mathbb{x}} \subset \mathbb{Q}\backslash\{-1,1\}^{J \times \max(\bar{I}_{\mathbb{x}})}$
with $\mathbb{x} \in [1, \mathbb{X}]; \mathbb{X} = \#\mathfrak{X}_J; a, b \in [1, \mathbb{A}]; a \neq b; \mathbb{A} = W.Y$

5.2 Biomedical relevance of the redundant forms of a biochemical network for the HMP-shunt pathway in the genesis of Klebsiella spp.-mediated formaldehyde resistance

Nosocomial (healthcare-associated) infections are usually observed in patients, in-residence, at a

healthcare facility for a significant period of time [61]. Formaldehyde (HCHO) is a surface-disinfectant,

a fixative and is also used to sterilize instruments such as endoscopes which cannot be autoclaved or

heated. HCHO is also generated intracellularly as part of the respiratory burst (macrophages,

neutrophils) in response to injury/infection, enzyme demethylation and tetrahydrofolate degradation [62-

64]. The electrophilic reactivity of HCHO will result in adduct-formation and crosslinking

(DNA/RNA/proteins) which is toxic to cells and non-methylotrophic bacteria [62, 63]. Interestingly,

several studies have demonstrated that the absence of HCHO-assimilating genes does not preclude non-

methylotrophic bacteria from utilizing and detoxifying the same [62-65]. In fact, gram negative bacteria

possess several molecular mechanisms to sense, detoxify and thrive in the presence of HCHO (*NmlR-*

adhC-estD, *FrmR-frmAB*) [62-65]. For example, a comparative analysis of several *Neisseria spp.*

suggests that the greater virulence of *N. meningitidis* could be its rapid dissemination. This, in turn,

could be on account of possessing both, *adhC* (alcohol dehydrogenase, EC1.1.2.84) and *estD*

thioesterase, EC3.1.2.12) genes as opposed to *N. gonorrhoea* (*estD*)..[62, 63]. Similar mechanisms are

operative for *Bacillus* spp. (*HxlR-hxlAB*) and *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*,

Escherichia coli (*FrmR-frmAB*) [63]. The biomedical and clinical relevance for these horizontally

acquired genes was appreciated when it was discovered that certain multi-drug resistant (MDR) strains

of *Klebsiella pneumoniae* (sequence type 4523; ST4523) and *Pseudomonas aeruginosa* were found to

possess the FrmRAB operon and could thrive in an environment of HCHO along with other biocides

[66, 67].

The molecular mechanisms to activate and thence assimilate HCHO depends on it condensing with reduced Glutathione (GSH) which in turn, is oxidized (GSSG) and will need to be reduced. The HMP-shunt pathway (oxidative, non-oxidative) is a conserved (eukaryotes, bacteria) pathway that produces reduced Nicotinamide adenine dinucleotide phosphate ($NADPH + H^+$), phosphoribosyl pyrophosphate (PRPP) and many other potentially anaplerotic intermediates for Kreb's cycle [68-71]. Although $NADPH + H^+$ partakes in several important pathways, it is the major source of reducing equivalents in the reduction and subsequent regeneration of GSH [62-65]. Clearly the integrity and functionality of the HMP-shunt pathway is important for the growth and continual survival of bacteria, a factor facilitated by the concomitant formation of PRPP. The Ribulose-5-phosphate (RuMP)-pathway is either the bona fide oxidative phase of the pathway, as in methylotrophic bacteria (*M. aminofaciens*, *B. bravis*, *Bacillus* spp.) and some archaea (methanogens, methylotrophic), or complements an existent HMP-shunt pathway (non-methylotrophic organisms; *S. cerevisae*, *E. coli*) [64-67, 72, 73]. RuMP/XuMP condenses with formaldehyde to give a hexulose-6-phosphate which isomerizes into fructose-6-phosphate via the reversible reactions that are catalysed by hexulose-6-phosphate synthase (EC4.1.2.43) and phosphohexose isomerase (EC5.3.1.9) respectively [64-67].

Horizontal gene transmission (HGT) may contribute to HCHO-mediated resistance in nosocomial infections via the HMP-pathway. Although, the inter-conversion between β-D fructose 6-phosphate and D-ribulose 6-phosphate (r_{19}) on a redundant form of the HMP-shunt pathway, in a multi-drug variant of *K. pneumoniae* (ST4523), does not alter the outcome of the shared equivalent reactions ($\eta_8 = \tilde{\eta}_8, \eta_{10} = \tilde{\eta}_{10}, \eta_{11} = \tilde{\eta}_{11}, \eta_{12} = \tilde{\eta}_{12,}$), the same does not hold true for the shared non-equivalent reactions (**Figures 4a and 4b, Table 2; ST3, SD3**). These exhibit a reduction in magnitude ($\eta_4 > \tilde{\eta}_4, \eta_{14} > \tilde{\eta}_{14}$) as well as reversed outcomes ($\{\eta_7, \eta_9\} > 1 \rightarrow \{\tilde{\eta}_7, \tilde{\eta}_9\} \in (0,1); \{\eta_{10}, \eta_{11}, \eta_{12}\} \in (0,1) \rightarrow \{\tilde{\eta}_{10}, \tilde{\eta}_{11}, \tilde{\eta}_{12}\} > 1$) (**Figures 4a and 4b, Table 2; ST3, SD3**). From this data it is clear that any existing and host-specific Ribulose-5-phosphate is utilized ($r_{7,9}$) and/or redirected (r_{10-12}). This apparent loss of D-Ribulose-5-phosphate will, by the law of mass action and in concert with r_4 and r_{14}, result in additional D-Ribulose-5-phosphate being formed from D-Fructose-6-phosphate via r_{19} ($\tilde{\eta}_{19} \approx 18.0$) (**Figures 4a and 4b, Tables 3** and **4; ST3, SD3**). The subsequent condensation of HCHO by D-Ribulose-5-phosphate and thence assimilation/utilization into D-Fructose-6-phosphate will allow *K. pneumoniae* (ST4523) to grow, thrive and disseminate (**Figure 4c**).

6   Concluding remarks

This study develops a reactant-dependent algebraic framework to study and characterize the redundant forms of a biochemical network with a library of constrained, rational number, equivalent and redundant/degenerate matrices for the $\mathfrak{x}^{\text{th}}$-biochemical network (CRaTER$\mathfrak{x}$). These matrices form a many-one map with the stoichiometry number matrix that they are derived from, have distinct mathematical properties, are easily parameterized, can be perturbed and can simulate a biochemical network under differing intracellular conditions. The theoretical results are complemented with detailed computational studies of known biochemical pathways. Future studies will endeavour to search for and identify additional parameters to characterize network-redundancy, identify bounds for the numbers of perturbing PRVs and incorporate/import/analyse data from publicly available datasets.

7   Detailed proofs (STx)

Detailed proofs to construct and analyse a reactant-dependent algebraic framework

3.4.2   Proofs to define and construct a reactant-dependent algebraic framework

Proof (T1):

$$\mathfrak{X}(J) = \mathfrak{X}_J \tag{1}$$

$$= \left\{ \left(\bar{\theta}_{\hat{\jmath}_{J,1}}, \hat{I}_{\hat{\jmath}_{J,1}} \right), \dots, \left(\bar{\theta}_{\hat{\jmath}_{J,\mathbb{J}}}, \hat{I}_{\hat{\jmath}_{J,\mathbb{J}}} \right) \right\} \tag{2}$$

$$= \left\{ \left(\frac{\hat{I}_{\hat{\jmath}_{J,1}}}{\bar{\theta}_{\hat{\jmath}_{J,1}}} \right), \dots, \left(\frac{\hat{I}_{\hat{\jmath}_{J,\mathbb{J}}}}{\bar{\theta}_{\hat{\jmath}_{J,\mathbb{J}}}} \right) \right\} \tag{3}$$

$$= \#\mathfrak{S}_{\hat{\jmath}_J} \tag{4}$$

$$= \left\{ \#\mathfrak{S}_{\hat{\jmath}_{J,1}}, \dots, \#\mathfrak{S}_{\hat{\jmath}_{J,\mathbb{J}}} \right\} \tag{5}$$

$$\Rightarrow \left\{ \left(\tilde{\mathcal{S}}(.)_{\hat{\jmath}_{J,s=1,x=1}}, \dots, \tilde{\mathcal{S}}(.)_{\hat{\jmath}_{J,s=1,x=X_{\hat{\jmath}_{J,s}}}} \right), \dots, \left(\tilde{\mathcal{S}}(.)_{\hat{\jmath}_{J,s=\mathbb{J},x=1}}, \dots, \tilde{\mathcal{S}}(.)_{\hat{\jmath}_{J,s=\mathbb{J},x=X_{\hat{\jmath}_{J,s}}}} \right) \right\} \tag{6}$$

$$= \left\{ \left(\bigcup_{s=1}^{s=\mathbb{J}} \tilde{\mathcal{S}}(.)_{\hat{\jmath}_{J,s,x}} - \bigcap_{x=1}^{x=X_{\hat{\jmath}_{J,s}}} \tilde{\mathcal{S}}(.)_{\hat{\jmath}_{J,s,x}} \right), \dots, \left(\bigcup_{s=1}^{s=\mathbb{J}} \tilde{\mathcal{S}}(.)_{\hat{\jmath}_{J,s,x}} - \bigcap_{x=1}^{x=X_{\hat{\jmath}_{J,s}}} \tilde{\mathcal{S}}(.)_{\hat{\jmath}_{J,s,x}} \right) \right\} \tag{7}$$

$$= \left\{ \left(\bigcup_{s=1}^{s=\mathbb{J}} \bigcup_{x=1}^{x=X_{\hat{\jmath}_{J,s}}} \tilde{\mathcal{S}}(.)_{\hat{\jmath}_{J,s,x}} \right) - \left(\bigcap_{s=1}^{s=\mathbb{J}} \bigcap_{x=1}^{x=X_{\hat{\jmath}_{J,s}}} \tilde{\mathcal{S}}(.)_{\hat{\jmath}_{J,s,x}} \right) \right\} \tag{7.1}$$

∎

Proof (C3):

$$\#D(\mathbf{\Phi}) = \binom{\#LT(\mathbf{\Phi}) \vee \#UT(\mathbf{\Phi})}{1} \tag{8}$$

$$= \binom{\#rows(\mathbf{\Phi}) \vee \#cols(\mathbf{\Phi})}{1} \tag{9}$$

$$= \left(\frac{W!}{1!\,(W-1)!}\right) \vee \left(\frac{Y!}{1!\,(Y-1)!}\right) \tag{10}$$

$$= \left(\frac{W.(W-1)!}{1!\,(W-1)!}\right) \vee \left(\frac{Y.(Y-1)!}{1!\,(Y-1)!}\right) \tag{10.1}$$

$$= (W) \vee (Y) \tag{11}$$

$$\#LT(\mathbf{\Phi}) = \binom{\#rows(\mathbf{\Phi})}{2} \tag{12}$$

$$= \left(\frac{W!}{2!\,(W-2)!}\right) \tag{13}$$

$$= \left(\frac{W.(W-1).(W-2)!}{2!\,(W-2)!}\right) \tag{13.1}$$

$$= \left(\frac{W.(W-1)}{2!}\right) \tag{13.2}$$

$$\#UT(\mathbf{\Phi}) = \binom{\#cols(\mathbf{\Phi})}{2} \tag{14}$$

$$= \left(\frac{Y!}{2!\,(Y-2)!}\right) \tag{15}$$

$$= \left(\frac{Y.(Y-1).(Y-2)!}{2!\,(Y-2)!}\right) \tag{15.1}$$

$$= \left(\frac{Y.(Y-1)}{2!}\right) \tag{15.2}$$

$$\#Vec(\mathbf{\Phi}) = \#D(\mathbf{\Phi}) + \#LT(\mathbf{\Phi}) + \#UT(\mathbf{\Phi}) \tag{16}$$

$$= \big((W) \vee (Y)\big) + \left(\frac{W.(W-1)}{2!}\right) + \left(\frac{Y.(Y-1)}{2!}\right) \tag{17}$$

$$\Rightarrow \big((W) \vee (Y)\big) + \left(2.\left(\frac{W.(W-1)}{2!}\right)\right) \vee \left(2.\left(\frac{Y.(Y-1)}{2!}\right)\right) \quad [\text{Since } W = Y] \tag{18}$$

$$= \left(W + 2.\left(\frac{W.(W-1)}{2!}\right)\right) \vee \left(Y + 2.\left(\frac{Y.(Y-1)}{2!}\right)\right) \tag{19}$$

$$= \big(W + W.(W-1)\big) \vee \big(Y + Y.(Y-1)\big) \tag{19.1}$$

$$= (W + W^2 - W) \vee (Y + Y^2 - Y) \vee (W.Y) \tag{19.2}$$

$$= (W^2) \vee (Y^2) \vee (W.Y) \quad [\text{Since } W = Y] \tag{20}$$

∎

Proof (C4-C8):

C4:
$$(\mathbf{\Phi}^T)^T = \left(\left(\phi\right)_{w,y}{}^T\right)^T = \left(\left(\phi\right)_{y,w}\right)^T = \left(\phi\right)_{y,w}{}^T = \left(\phi\right)_{w,y} = \mathbf{\Phi} \tag{21-25}$$
∎

C5:
$$(\mathbf{\Phi}^p)^T = \left(\left(\phi\right)_{w,y}{}^p\right)^T = \left(\left(\phi\right)_{w,y}{}^T\right)^p = \left(\left(\phi\right)_{y,w}\right)^p = \left(\left(\phi\right)_{w,y}{}^T\right)^p = (\mathbf{\Phi}^T)^p \tag{26-31}$$
∎

C6:
$$q.\,(\mathbf{\Phi})^T = q.\,(\phi)_{w,y}{}^T = q.\,(\phi)_{y,w} = (q.\,\phi)_{y,w} = (q.\,\phi)_{w,y}{}^T = (q.\,\mathbf{\Phi})^T \tag{32-36}$$
∎

C7:
For an off $-$ diagonal element $\bar{\bar{\phi}}_{\widehat{w},\widehat{y}} \in \mathbf{\Phi} \times \mathbf{\Phi}^T$ where $\phi_{w,y} \in \mathbf{\Phi}, \widehat{\phi}_{w,y} \in \mathbf{\Phi}^T$ and $\widehat{w} \neq \widehat{y}$ $\quad$ (37 $-$ 39)
$\widehat{w} := \text{row } (1, \dots, W \vee Y)$ and $\widehat{y} := \text{column } (1, \dots, W \vee Y)$ $\qquad\qquad\quad$ (40), (41)
$n := \text{Number of terms } (W \vee Y)$ $\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\quad$ (42)

$$\bar{\bar{\phi}}_{\widehat{w},\widehat{y}} = \sum_{n=1}^{n=W\vee Y}\left(\phi_{\widehat{w},n}\right).\left(\widehat{\phi}_{n,\widehat{y}}\right) = \left(\phi_{\widehat{w},1}.\,\widehat{\phi}_{1,\widehat{y}}\right) + \cdots + \left(\phi_{\widehat{w},W}.\,\widehat{\phi}_{W,\widehat{y}}\right) \tag{43}$$

Swapping the row and column indices (transpose) and rewriting Eq. (43)
$$\bar{\bar{\phi}}_{\widehat{w},\widehat{y}} = \left(\phi_{\widehat{y},1}.\,\widehat{\phi}_{1,\widehat{w}}\right) + \cdots + \left(\phi_{\widehat{y},W}.\,\widehat{\phi}_{W,\widehat{w}}\right) = \sum_{n=1}^{n=W\vee Y}\left(\phi_{\widehat{y},n}\right).\left(\widehat{\phi}_{n,\widehat{w}}\right) = \bar{\bar{\phi}}_{\widehat{y},\widehat{w}} \tag{44}$$
∎

C8:
Let $k_1, k_2 \in \mathbb{R} - \{0\}$ $\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\quad$ (45), (46)
Let $A_1, A_2 \in \mathbb{R}$ where $A_1 = \det(\mathbf{\Phi}), A_2 = \det(\mathbf{\Phi}^T)$ $\qquad\qquad\qquad\qquad$ (47), (48)
$k_1.\,A_1 = \det(k_1.\,\mathbf{\Phi}) = \det\left(k_1.\,(\phi)_{w,y}\right) = \det\left((\phi)_{y,w}\right) = \det(\mathbf{\Phi}^T) = A_2$ $\quad$ (49)
$k_2.\,A_2 = \det(k_2.\,\mathbf{\Phi}^T) = \det\left(k_2.\,(\phi)_{y,w}\right) = \det\left((\phi)_{w,y}\right) = \det(\mathbf{\Phi}) = A_1$ $\quad$ (50)

Solving these simultaneously,
$$k_1 = \frac{A_1}{A_1.\,k_2} = k_2{}^{-1} \tag{51}$$

$$k_2 = \frac{A_2}{A_2.\,k_1} = k_1{}^{-1} \tag{52}$$

Clearly, $k_1.\,k_2 = k_1.\,k_1{}^{-1} = k_2{}^{-1}.\,k_2 = 1 \overset{\text{def}}{=} k$ $\qquad\qquad\qquad\qquad\qquad$ (53 $-$ 56)
$k_1.\,A_1 = A_2 = k.\,A_1$ $\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\quad$ (57), (58)
$k_2.\,A_2 = A_1 = k.\,A_2$ $\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\quad$ (59), (60)

$A_1 = A_2$ $\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad$ (61)
∎

Proof (T4):

$$\prod_{w=1}^{w=W} \prod_{y=1}^{y=Y} \phi_{w,y} \tag{62}$$

$$\left(\prod_{w=2}^{w=W} \prod_{y=1}^{y=w-1} \phi_{w,y}\right) \cdot \left(\prod_{y=2}^{y=Y} \prod_{w=1}^{w=y-1} \phi_{y,w}\right) \tag{63}$$

$$\Rightarrow LT(\mathbf{\Phi}).\,UT(\mathbf{\Phi}) \tag{64}$$

Since $\phi_{y,w} = \phi_{w,y}{}^{-1}$ for $w \in [1, W]$ and $y \in [1, Y]$ **[From T3, L1]**

$$(\phi_{w,y}).\,(\phi_{y,w}) = (\phi_{w,y}).\,(\phi_{w,y}{}^{-1}) = \left(\frac{w}{y}\right).\left(\frac{y}{w}\right) = 1 \; \forall w \forall y \tag{65 - 67}$$

Conversely since $\phi_{w,y} = \phi_{y,w}{}^{-1}$ we can rewrite Eq. (65) as: **[From T3, L1]**

$$(\phi_{w,y}).\,(\phi_{y,w}) = (\phi_{y,w}{}^{-1}).\,(\phi_{y,w}) = \left(\frac{w}{y}\right).\left(\frac{y}{w}\right) = 1 \; \forall w \forall y \tag{68 - 70}$$

$$\prod_{w=1}^{w=W} \prod_{y=1}^{y=Y} \phi_{w,y} = 1 \qquad\qquad \textbf{[From Eqs. (67) and (70)]} \tag{71}$$

∎

3.4.2 Proofs of describe the properties of the redundant forms or equivalent rational number matrices for the $\mathbb{x}^{th}$-biochemical network via the computed null space

Proof (T7):

$$\mathcal{S}_{\wp_{J,\mathbb{x},a}} = \left((\phi_{w,y}) \vee (\phi_{y,w}) \right) . \tilde{\mathcal{S}}_{\wp_{J,\mathbb{x}}} \tag{72}$$

$$= \left(\phi_{w,y} \vee \phi_{y,w} \right) . \tilde{\mathcal{S}}_{\wp_{J,\mathbb{x}}} \tag{73}$$

$$N\left(\mathcal{S}_{\wp_{J,\mathbb{x},a}} \right) = N\left(\left(\phi_{w,y} \vee \phi_{y,w} \right) . \tilde{\mathcal{S}}_{\wp_{J,\mathbb{x}}} \right) \tag{74}$$

$$= N\left(\left(\phi_{w,y} . \tilde{\mathcal{S}}_{\wp_{J,\mathbb{x}}} \vee \phi_{y,w} . \tilde{\mathcal{S}}_{\wp_{J,\mathbb{x}}} \right) \right) \tag{74.1}$$

$$= N\left(\left(\phi_{w,y} . \tilde{\mathcal{S}}_{\wp_{J,\mathbb{x}}} \right) \vee \left(\phi_{y,w} . \tilde{\mathcal{S}}_{\wp_{J,\mathbb{x}}} \right) \right) \tag{74.2}$$

$$= N\left(\left(\phi_{w,y} \right) . \tilde{\mathcal{S}}_{\wp_{J,\mathbb{x}}} \right) \vee N\left(\left(\phi_{y,w} \right) . \tilde{\mathcal{S}}_{\wp_{J,\mathbb{x}}} \right) \tag{74.3}$$

$$= N\left(\mathcal{S}_{\wp_{J,\mathbb{x},b}} \right) \vee N\left(\mathcal{S}_{\wp_{J,\mathbb{x},c}} \right) \tag{74.4}$$

Rewriting (74.4)

$$\left(N\left(\mathcal{S}_{\wp_{J,\mathbb{x},a}} \right) = N\left(\mathcal{S}_{\wp_{J,\mathbb{x},b}} \right) \right) \vee \left(N\left(\mathcal{S}_{\wp_{J,\mathbb{x},a}} \right) = N\left(\mathcal{S}_{\wp_{J,\mathbb{x},c}} \right) \right) \tag{75}$$

$$\Rightarrow \left(N\left(\mathcal{S}_{\wp_{J,\mathbb{x},b}} \right) = N\left(\mathcal{S}_{\wp_{J,\mathbb{x},c}} \right) \right) \tag{76}$$

From (75), (76)

$$N\left(\boldsymbol{\mathcal{S}}_{\wp_{J,\mathbb{x},a}} \right) = N\left(\boldsymbol{\mathcal{S}}_{\wp_{J,\mathbb{x},b}} \right) = N\left(\boldsymbol{\mathcal{S}}_{\wp_{J,\mathbb{x},c}} \right) \tag{77}$$

where $\left\{ \boldsymbol{\mathcal{S}}_{\wp_{J,\mathbb{x},a}}, \boldsymbol{\mathcal{S}}_{\wp_{J,\mathbb{x},b}}, \boldsymbol{\mathcal{S}}_{\wp_{J,\mathbb{x},c}} \right\} \in \mathfrak{C}_{J,\mathbb{x}}$ and $\boldsymbol{\mathcal{S}}_{\wp_{J,\mathbb{x},a}} \neq \boldsymbol{\mathcal{S}}_{\wp_{J,\mathbb{x},b}} \neq \boldsymbol{\mathcal{S}}_{\wp_{J,\mathbb{x},c}}$ (78), (79)

and $a, b, c \in [1, \mathbb{A}]; a \neq b \neq c$ (80), (81)

Eq. (77) will hold for any ordered pair (w, y) where $w = 1, \dots, W$ and $y = 1, \dots, Y$ (82), (83)

$$\Rightarrow N\left(\boldsymbol{\mathcal{S}}_{\wp_{J,\mathbb{x},a=1}} \right) = \cdots = N\left(\boldsymbol{\mathcal{S}}_{\wp_{J,\mathbb{x},a=\mathbb{A}}} \right) \text{ for } \tilde{\mathcal{S}}_{\wp_{J,\mathbb{x}}} \in \mathfrak{X}_J \subset \mathbb{Z}^{J \times \max(\hat{l}_\mathbb{x})} \tag{84}$$

where $\mathbb{x} \in [1, \mathbb{X}], \mathbb{X} = \#\mathfrak{X}_J, \mathbb{A} = W.Y$ and $J \geq 4$

∎

Proof (T8): By contradiction

Since $f: \mathfrak{C}_{J,x} \twoheadrightarrow N(\mathfrak{C}_{J,x})$

$$\Rightarrow \left(\begin{array}{c} \text{for each } a = 1, \dots, \mathbb{A} \; \exists \mathcal{S}_{\wp_{J,x,a}} \in \mathfrak{C}_x \text{ s.t. } f\left(\boldsymbol{S}_{\wp_{J,x,a}}\right) \in N(\mathfrak{C}_{J,x}) \\ \text{Clearly } f\left(\boldsymbol{S}_{\wp_{J,x,a=1}}\right) = \cdots = f\left(\boldsymbol{S}_{\wp_{J,x,a=\mathbb{A}}}\right) \equiv N\left(\boldsymbol{S}_{\wp_{J,x,a=1}}\right) = \cdots = N\left(\boldsymbol{S}_{\wp_{J,x,a=\mathbb{A}}}\right) \text{ and } f^{-1}\left(N(\mathfrak{C}_{J,x})\right) \in \mathfrak{C}_{J,x} \end{array} \right) \quad (85-87)$$

Since $f^{-1}\left(N(\mathfrak{C}_x)\right) \in \mathfrak{C}_{J,x}$

$$\Rightarrow f^{-1} \circ f(\mathfrak{C}_{J,x}) \in \mathfrak{C}_{j,x} \tag{88}$$

$$= f^{-1} \circ \left(f\left(\boldsymbol{S}_{\wp_{J,x,a=1}}\right) \vee \dots \vee f\left(\boldsymbol{S}_{\wp_{J,x,a=\mathbb{A}}}\right) \right) \in \mathfrak{C}_{J,x} \tag{89}$$

$$= f^{-1} \circ \left(\bigvee_{a=1}^{a=\mathbb{A}} f\left(\boldsymbol{S}_{\wp_{J,x,a}}\right) \right) \in \mathfrak{C}_{J,x} \tag{90}$$

$$= f^{-1} \circ f\left(\boldsymbol{S}_{\wp_{J,x,a=1}}\right) \vee \dots \vee f^{-1} \circ f\left(\boldsymbol{S}_{\wp_{J,x,a=\mathbb{A}}}\right) \in \mathfrak{C}_{J,x} \text{ where } f^{-1} \circ f\left(\boldsymbol{S}_{\wp_{J,x,a=1}}\right) \neq \cdots \neq f^{-1} \circ f\left(\boldsymbol{S}_{\wp_{J,x,a=\mathbb{A}}}\right) \quad (90.1),(91)$$

Let $\mathcal{B} \in N(\mathfrak{C}_{J,x})$ s.t. $\mathcal{B} \neq \left\{ f\left(\boldsymbol{S}_{\wp_{J,x,a=1}}\right), \dots, f\left(\boldsymbol{S}_{\wp_{J,x,a=\mathbb{A}}}\right) \right\}$ $\qquad(92)$

$$\Rightarrow f^{-1}(\mathcal{B}) \in \mathfrak{C}_{J,x} \text{ s.t. } f^{-1}(\mathcal{B}) \neq \left\{ f^{-1} \circ f\left(\boldsymbol{S}_{\wp_{J,x,a=1}}\right) \neq \cdots \neq f^{-1} \circ f\left(\boldsymbol{S}_{\wp_{J,x,a=\mathbb{A}}}\right) \right\} \quad (93),(94)$$

$$\Rightarrow f^{-1}(\mathcal{B}) \neq \left\{ (f^{-1} \circ f)\left(\boldsymbol{S}_{\wp_{x_{a=1}}}\right) \neq \cdots \neq (f^{-1} \circ f)\left(\boldsymbol{S}_{\wp_{J,x,a=\mathbb{A}}}\right) \right\} \neq \left\{ \boldsymbol{S}_{\wp_{J,x,a=1}}, \dots, \boldsymbol{S}_{\wp_{J,x,a=\mathbb{A}}} \right\} \quad (95),(96)$$

But $f \circ f^{-1}(\mathcal{B}) \in N(\mathfrak{C}_{J,x})$ $\qquad(97)$

where $f \circ f^{-1}(\mathcal{B}) = (f \circ f^{-1})(\mathcal{B}) = \mathcal{B} = \left\{ f\left(\boldsymbol{S}_{\wp_{J,x,a=1}}\right), \dots, f\left(\boldsymbol{S}_{\wp_{J,x,a=\mathbb{A}}}\right) \right\}$ $\qquad(98-100)$

Clearly Eq. (100) contradicts Eq. (92)

∎

3.4.2 Proofs to describe the algebraic properties of the redundant forms or equivalent rational number matrices for the x^{th}-biochemical network

Proof (T9):

No Additive identity:

if $\boldsymbol{S}_{\wp_{J,x,a}} \in \mathfrak{C}_{J,x} \Rightarrow \boldsymbol{S}_{\wp_{J,x,a}} \neq \boldsymbol{0}^{J \times \max(\bar{I}_x)}$ [By definition] $\qquad(101)$

Closure: From (**L2**),
if $\boldsymbol{S}_{\wp_{J,x,a}}, \boldsymbol{S}_{\wp_{J,x,b}} \in \mathfrak{C}_{J,x}$ where $\boldsymbol{S}_{\wp_{J,x,a}} \neq \boldsymbol{S}_{\wp_{J,x,b}}; a,b \in [1,\mathbb{A}]; a \neq b$
then $\boldsymbol{S}_{\wp_{J,x,a}} + \boldsymbol{S}_{\wp_{J,x,b}} = \boldsymbol{S}_{\wp_{J,x,(a+b)}} \in \mathfrak{C}_{J,x}$ $\qquad(102)$

Association:
if $\boldsymbol{S}_{\wp_{J,x,a}}, \boldsymbol{S}_{\wp_{J,x,b}}, \boldsymbol{S}_{\wp_{J,x,c}} \in \mathfrak{C}_{J,x}$ where $\boldsymbol{S}_{\wp_{J,x,a}} \neq \boldsymbol{S}_{\wp_{J,x,b}} \neq \boldsymbol{S}_{\wp_{J,x,c}}; a,b,c \in [1,\mathbb{A}]; a \neq b \neq c$

then $\left(\boldsymbol{S}_{\wp_{J,x,a}} + \boldsymbol{S}_{\wp_{J,x,b}}\right) + \boldsymbol{S}_{\wp_{J,x,c}} = \boldsymbol{S}_{\wp_{J,x,(a+b)}} + \boldsymbol{S}_{\wp_{J,x,c}} = \boldsymbol{S}_{\wp_{J,x,(a+b+c)}} \in \mathfrak{C}_{J,x}$ $\qquad(103)$

and $\boldsymbol{S}_{\wp_{J,x,a}} + \left(\boldsymbol{S}_{\wp_{J,x,b}} + \boldsymbol{S}_{\wp_{J,x,c}}\right) = \boldsymbol{S}_{\wp_{J,x,a}} + \boldsymbol{S}_{\wp_{J,x,(b+c)}} = \boldsymbol{S}_{\wp_{J,x,(a+b+c)}} \in \mathfrak{C}_{J,x}$ $\qquad(104)$

From (103), (104)
$$\left(\boldsymbol{S}_{\wp_{J,x,a}} + \boldsymbol{S}_{\wp_{J,x,b}}\right) + \boldsymbol{S}_{\wp_{J,x,c}} = \boldsymbol{S}_{\wp_{J,x,a}} + \left(\boldsymbol{S}_{\wp_{J,x,b}} + \boldsymbol{S}_{\wp_{J,x,c}}\right) \tag{105}$$

Generalizing,
$$\left(\boldsymbol{S}_{\wp_{J,x,a}} + \boldsymbol{S}_{\wp_{J,x,b}}\right) \dots + \boldsymbol{S}_{\wp_{J,x,\mathbb{A}}} = \boldsymbol{S}_{\wp_{J,x,a}} + \left(\boldsymbol{S}_{\wp_{J,x,b}} \dots + \boldsymbol{S}_{\wp_{J,x,\mathbb{A}}}\right) = \boldsymbol{S}_{\wp_{J,x,(a+\dots+\mathbb{A})}} = \boldsymbol{S}_{\wp_{J,x,\left(\frac{(\mathbb{A}).(\mathbb{A}+1)}{2}\right)}} \in \mathfrak{C}_x \quad (106-108)$$

∎

Acknowledgements

This study was supported, in part by an extramural grant from the Science and Engineering Research Board (SERB), Department of Science and Technology, Government of India under the Mathematical Research Impact-Centric Support (MATRICS) scheme to SK (MTR/2021/000290).

Author Contributions

SK, designed the study; defined, formulated, developed and wrote proofs for the theorems, lemmas and corollaries; developed the assessment metrics and algorithms; conducted numerical studies and collated data; wrote all necessary code and manuscript.

Competing Interests

The author declares no conflict of interest.

References

1.   Kim OD, Rocha M, Maia P. A Review of Dynamic Modeling Approaches and Their Application in Computational Strain Optimization for Metabolic Engineering. Front Microbiol. 2018;9:1690. Epub 2018/08/16. doi: 10.3389/fmicb.2018.01690. PubMed PMID: 30108559; PubMed Central PMCID: PMCPMC6079213.

2.   Maarleveld TR, Khandelwal RA, Olivier BG, Teusink B, Bruggeman FJ. Basic concepts and principles of stoichiometry modeling of metabolic networks. Biotechnol J. 2013;8(9):997-1008. Epub 2013/07/31. doi: 10.1002/biot.201200291. PubMed PMID: 23893965; PubMed Central PMCID: PMCPMC4671265.

3.   Klamt S, Muller S, Regensburger G, Zanghellini J. A mathematical framework for yield (vs. rate) optimization in constraint-based modeling and applications in metabolic engineering. Metab Eng. 2018;47:153-69. Epub 2018/02/11. doi: 10.1016/j.ymben.2018.02.001. PubMed PMID: 29427605; PubMed Central PMCID: PMCPMC5992331.

4.   Klamt S, Regensburger G, Gerstl MP, Jungreuthmayer C, Schuster S, Mahadevan R, et al. From elementary flux modes to elementary flux vectors: Metabolic pathway analysis with arbitrary linear flux constraints. PLoS Comput Biol. 2017;13(4):e1005409. Epub 2017/04/14. doi: 10.1371/journal.pcbi.1005409. PubMed PMID: 28406903; PubMed Central PMCID: PMCPMC5390976.

5.   Orth JD, Thiele I, Palsson BO. What is flux balance analysis? Nat Biotechnol. 2010;28(3):245-8. Epub 2010/03/10. doi: 10.1038/nbt.1614. PubMed PMID: 20212490; PubMed Central PMCID: PMCPMC3108565.

6.   Muller S, Regensburger G. Elementary Vectors and Conformal Sums in Polyhedral Geometry and their Relevance for Metabolic Pathway Analysis. Front Genet. 2016;7:90. Epub 2016/06/03. doi: 10.3389/fgene.2016.00090. PubMed PMID: 27252734; PubMed Central PMCID: PMCPMC4877377.

7.   Kundu, S. ReDirection: an R-package to compute the probable dissociation constant for every reaction of a user-defined biochemical network. *Frontiers in Molecular Biosciences* **10**, doi:10.3389/fmolb.2023.1206502 (2023).

8.   Kundu, S. A mathematically rigorous algorithm to define, compute and assess relevance of the probable dissociation constants in characterizing a biochemical network. Sci Rep 14, 3507, doi:10.1038/s41598-024-53231-9 (2024).

9.   Srividhya J, Crampin EJ, McSharry PE, Schnell S. Reconstructing biochemical pathways from time course data. Proteomics. 2007;7(6):828-38. Epub 2007/03/21. doi: 10.1002/pmic.200600428. PubMed PMID: 17370261.

10.   Oksuz M, Sadikoglu H, Cakir T. Sparsity as cellular objective to infer directed metabolic networks from steady-state metabolome data: a theoretical analysis. PLoS One. 2013;8(12):e84505. Epub 2014/01/07. doi: 10.1371/journal.pone.0084505. PubMed PMID: 24391961; PubMed Central PMCID: PMCPMC3877278.

11.   Nagele T. Linking metabolomics data to underlying metabolic regulation. Front Mol Biosci. 2014;1:22. Epub 2014/01/01. doi: 10.3389/fmolb.2014.00022. PubMed PMID: 25988163; PubMed Central PMCID: PMCPMC4428386.

12.   Cakir T, Hendriks MM, Westerhuis JA, Smilde AK. Metabolic network discovery through reverse engineering of metabolome data. Metabolomics. 2009;5(3):318-29. Epub 2009/09/01. doi: 10.1007/s11306-009-0156-4. PubMed PMID: 19718266; PubMed Central PMCID: PMCPMC2731157.

13.   Khatibipour MJ, Kurtoglu F, Cakir T. JacLy: a Jacobian-based method for the inference of metabolic interactions from the covariance of steady-state metabolome data. PeerJ. 2018;6:e6034. Epub 2018/12/20. doi: 10.7717/peerj.6034. PubMed PMID: 30564518; PubMed Central PMCID: PMCPMC6286809.

14.   Wang T, Ren Z, Ding Y, Fang Z, Sun Z, MacDonald ML, et al. FastGGM: An Efficient Algorithm for the Inference of Gaussian Graphical Model in Biological Networks. PLoS Comput Biol. 2016;12(2):e1004755. Epub 2016/02/13. doi: 10.1371/journal.pcbi.1004755. PubMed PMID: 26872036; PubMed Central PMCID: PMCPMC4752261.

15.   Nagele T, Mair A, Sun X, Fragner L, Teige M, Weckwerth W. Solving the differential biochemical Jacobian from metabolomics covariance data. PLoS One. 2014;9(4):e92299. Epub 2014/04/04. doi: 10.1371/journal.pone.0092299. PubMed PMID: 24695071; PubMed Central PMCID: PMCPMC3977476.

16.   Rottman BM, Hastie R. Reasoning about causal relationships: Inferences on causal networks. Psychol Bull. 2014;140(1):109-39. Epub 2013/04/03. doi: 10.1037/a0031903. PubMed PMID: 23544658; PubMed Central PMCID: PMCPMC3988659.

17.   Goto T, Fernandes AFA, Tsudzuki M, Rosa GJM. Causal phenotypic networks for egg traits in an F2 chicken population. Mol Genet Genomics. 2019;294(6):1455-62. Epub 2019/06/27. doi: 10.1007/s00438-019-01588-2. PubMed PMID: 31240383.

18.   Seyhan AA, Carini C. Are innovation and new technologies in precision medicine paving a new era in patients centric care? J Transl Med. 2019;17(1):114. Epub 2019/04/07. doi: 10.1186/s12967-019-1864-9. PubMed PMID: 30953518; PubMed Central PMCID: PMCPMC6451233.

19.   Biane C, Delaplace F. Causal Reasoning on Boolean Control Networks Based on Abduction: Theory and Application to Cancer Drug Discovery. IEEE/ACM Trans Comput Biol Bioinform. 2019;16(5):1574-85. Epub 2018/12/26. doi: 10.1109/TCBB.2018.2889102. PubMed PMID: 30582550.

20.   Pai S, Bader GD (2018) Patient Similarity Networks for Precision Medicine. *J Mol Biol* 430: 2924-2938.

21.   Du Q, Gong C, Wang Q, Zhou D, Yang H, Pan W, et al. Genetic architecture of growth traits in Populus revealed by integrated quantitative trait locus (QTL) analysis and association studies. New Phytol. 2016;209(3):1067-82. Epub 2015/10/27. doi: 10.1111/nph.13695. PubMed PMID: 26499329.

22.   Gove RP, Chen W, Zweber NB, Erwin R, Rychtar J, Remington DL. Effects of causal networks on the structure and stability of resource allocation trait correlations. J Theor Biol. 2012;293:1-14. Epub 2011/10/19. doi: 10.1016/j.jtbi.2011.09.034. PubMed PMID: 22004994.

23.   Salvador AC, Arends D, Barrington WT, Elsaadi AM, Brockmann GA, Threadgill DW. Sex-specific genetic architecture in response to American and ketogenic diets. Int J Obes (Lond). 2021;45(6):1284-97. Epub 2021/03/17. doi: 10.1038/s41366-021-00785-7. PubMed PMID: 33723359; PubMed Central PMCID: PMCPMC8159743.

24.   Gopalan L, Sebastian A, Praul CA, Albert I, Ramachandran R. Metformin Affects the Transcriptomic Profile of Chicken Ovarian Cancer Cells. Genes (Basel). 2021;13(1). Epub 2022/01/22. doi: 10.3390/genes13010030. PubMed PMID: 35052372; PubMed Central PMCID: PMCPMC8774788.

25.   Lu J, Dumitrascu B, McDowell IC, Jo B, Barrera A, Hong LK, et al. Causal network inference from gene transcriptional time-series response to glucocorticoids. PLoS Comput Biol. 2021;17(1):e1008223. Epub 2021/01/30. doi: 10.1371/journal.pcbi.1008223. PubMed PMID: 33513136; PubMed Central PMCID: PMCPMC7875426

26.   Gross, E., Harrington, H. A., Rosen, Z. & Sturmfels, B. Algebraic Systems Biology: A Case Study for the Wnt Pathway. Bull Math Biol 78, 21-51, doi:10.1007/s11538-015-0125-1 (2016).

27.   Loutchko, D. Semigroup models for biochemical reaction networks. J Math Biol 86, 78, doi:10.1007/s00285-023-01898-5 (2023).

28.   Angeli, D., De Leenheer, P. & Sontag, E. D. A Petri net approach to the study of persistence in chemical reaction networks. Math Biosci 210, 598-618, doi:10.1016/j.mbs.2007.07.003 (2007).

29.   Shiu, A. & Sturmfels, B. Siphons in chemical reaction networks. Bull Math Biol 72, 1448-1463, doi:10.1007/s11538-010-9502-y (2010).

30.   Harrington, H. A., Ho, K. L., Thorne, T. & Stumpf, M. P. Parameter-free model discrimination criterion based on steady-state coplanarity. Proc Natl Acad Sci U S A 109, 15746-15751, doi:10.1073/pnas.1117073109 (2012).

31.   Taylor, D. et al. Topological data analysis of contagion maps for examining spreading processes on networks. Nat Commun 6, 7723, doi:10.1038/ncomms8723 (2015).

32.   Xue Y, Wang Y, Wang S, Yan M, Huang J, Yang X. Label-Free and Regenerable Aptasensor for Real-Time Detection of Cadmium(II) by Dual Polarization Interferometry. Anal Chem. 2020;92(14):10007-15. Epub 2020/07/04. doi: 10.1021/acs.analchem.0c01710. PubMed PMID: 32618180.

33.   Da P, Li W, Lin X, Wang Y, Tang J, Zheng G. Surface plasmon resonance enhanced real-time photoelectrochemical protein sensing by gold nanoparticle-decorated TiO(2) nanowires. Anal Chem. 2014;86(13):6633-9. Epub 2014/06/11. doi: 10.1021/ac501406x. PubMed PMID: 24915128.

34.   Nagai H, Ebisu S, Abe R, Goto T, Takahashi N, Hosaka T, et al. Development of a novel PPARgamma ligand screening system using pinpoint fluorescence-probed protein. Biosci Biotechnol Biochem. 2011;75(2):337-41. Epub 2011/02/11. doi: 10.1271/bbb.100810. PubMed PMID: 21307572.

35.   Smith DS, Ferris FG. Specific surface chemical interactions between hydrous ferric oxide and iron-reducing bacteria determined using pK(a) spectra. J Colloid Interface Sci. 2003;266(1):60-7. Epub 2003/09/06. doi: 10.1016/s0021-9797(03)00667-2. PubMed PMID: 12957582.

36.   Shindo Y, Kondo Y, Sako Y (2018) Inferring a nonlinear biochemical network model from a heterogeneous single-cell time course data. *Sci Rep* 8: 6790.

37.   Koutrouli M, Karatzas E, Paez-Espino D, et al. (2020) A Guide to Conquer the Biological Network Era Using Graph Theory. Front Bioeng Biotechnol 8: 34.

38.   Kundu S. Modeling ligand-macromolecular interactions as eigenvalue-based transition-state dissociation constants may offer insights into biochemical function of the resulting complexes. Mathematical Biosciences and Engineering. 2022;19:13252-13275.

39.   De Martino, D., Mori, M. & Parisi, V. Uniform sampling of steady states in metabolic networks: heterogeneous scales and rounding. PLoS One 10, e0122670, doi:10.1371/journal.pone.0122670 (2015).

40.   Samal, S. S., Grigoriev, D., Frohlich, H., Weber, A. & Radulescu, O. A Geometric Method for Model Reduction of Biochemical Networks with Polynomial Rate Functions. Bull Math Biol 77, 2180-2211, doi:10.1007/s11538-015-0118-0 (2015).

41.   Theorell, A., Jadebeck, J. F., Noh, K. & Stelling, J. PolyRound: polytope rounding for random sampling in metabolic networks. Bioinformatics 38, 566-567, doi:10.1093/bioinformatics/btab552 (2022).

42.   Caspi, R. et al. The MetaCyc database of metabolic pathways and enzymes - a 2019 update. Nucleic Acids Res 48, D445-D453, doi:10.1093/nar/gkz862 (2020).

43.   Altman, T., Travers, M., Kothari, A., Caspi, R. & Karp, P. D. A systematic comparison of the MetaCyc and KEGG pathway databases. BMC Bioinformatics 14, 112, doi:10.1186/1471-2105-14-112 (2013).

44.   Noris, M. & Remuzzi, G. Overview of complement activation and regulation. Semin Nephrol 33, 479-492, doi:10.1016/j.semnephrol.2013.08.001 (2013).

45.   Stanton, D., Justin, H. S. & Reitzel, A. M. Step in Time: Conservation of Circadian Clock Genes in Animal Evolution. Integr Comp Biol 62, 1503-1518, doi:10.1093/icb/icac140 (2022).

46.   Hoffman, M. & Monroe, D. M. Coagulation 2006: a modern view of hemostasis. Hematol Oncol Clin North Am 21, 1-11, doi:10.1016/j.hoc.2006.11.004 (2007).

47.   Kakita, T. et al. Novel genotypes of Leptospira interrogans serogroup Sejroe isolated from human patients in Okinawa Prefecture, Japan. J Med Microbiol 69, 587-590, doi:10.1099/jmm.0.001169 (2020).

48.   Kundu, S. & Sharma, R. Origin, evolution, and divergence of plant class C GH9 endoglucanases. BMC Evol Biol 18, 79, doi:10.1186/s12862-018-1185-2 (2018).

49.   Bakkeren, E. et al. Impact of horizontal gene transfer on emergence and stability of cooperative virulence in Salmonella Typhimurium. Nat Commun 13, 1939, doi:10.1038/s41467-022-29597-7 (2022).

50.   Li, Z. et al. Arginase: shedding light on the mechanisms and opportunities in cardiovascular diseases. *Cell Death Discov* **8**, 413, doi:10.1038/s41420-022-01200-4 (2022).

51.   Caldwell, R. W., Rodriguez, P. C., Toque, H. A., Narayanan, S. P. & Caldwell, R. B. Arginase: A Multifaceted Enzyme Important in Health and Disease. *Physiol Rev* **98**, 641-665, doi:10.1152/physrev.00037.2016 (2018).

52.   Yao Y, Xu XH, Jin L. Macrophage Polarization in Physiological and Pathological Pregnancy. Front Immunol. 2019;10:792. Epub 2019/05/01. doi: 10.3389/fimmu.2019.00792. PubMed PMID: 31037072; PubMed Central PMCID: PMCPMC6476302.

53.   Latour, Y. L., Gobert, A. P. & Wilson, K. T. The role of polyamines in the regulation of macrophage polarization and function. *Amino Acids* **52**, 151-160, doi:10.1007/s00726-019-02719-0 (2020).

54.   Sagar, N. A., Tarafdar, S., Agarwal, S., Tarafdar, A. & Sharma, S. Polyamines: Functions, Metabolism, and Role in Human Disease Management. *Med Sci (Basel)* **9**, doi:10.3390/medsci9020044 (2021).

55.   Moinard, C., Cynober, L. & de Bandt, J. P. Polyamines: metabolism and implications in human diseases. *Clin Nutr* **24**, 184-197, doi:10.1016/j.clnu.2004.11.001 (2005).

56.   Perez-Leal, O. & Merali, S. Regulation of polyamine metabolism by translational control. Amino Acids 42, 611-617, doi:10.1007/s00726-011-1036-6 (2012).

57.   Banerjee, D. et al. Novel aspects of resistance to drugs targeted to dihydrofolate reductase and thymidylate synthase. *Biochim Biophys Acta* **1587**, 164-173, doi:10.1016/s0925-4439(02)00079-0 (2002).

58.   Peters, G. J. et al. Induction of thymidylate synthase as a 5-fluorouracil resistance mechanism. *Biochim Biophys Acta* **1587**, 194-205, doi:10.1016/s0925-4439(02)00082-0 (2002).

59.   Carreras, C. W. & Santi, D. V. The catalytic mechanism and structure of thymidylate synthase. *Annu Rev Biochem* **64**, 721-762, doi:10.1146/annurev.bi.64.070195.003445 (1995).

60.   Santhekadur, P. K. et al. The transcription factor LSF: a novel oncogene for hepatocellular carcinoma. *Am J Cancer Res* **2**, 269-285 (2012).

61.   Porter, L. et al. How long do nosocomial pathogens persist on inanimate surfaces? A scoping review. *J Hosp Infect* **147**, 25-31, doi:10.1016/j.jhin.2024.01.023 (2024).

62.   Chen, N. H. et al. A glutathione-dependent detoxification system is required for formaldehyde resistance and optimal survival of Neisseria meningitidis in biofilms. *Antioxid Redox Signal* **18**, 743-755, doi:10.1089/ars.2012.4749 (2013).

63.   Chen, N. H., Djoko, K. Y., Veyrier, F. J. & McEwan, A. G. Formaldehyde Stress Responses in Bacterial Pathogens. *Front Microbiol* **7**, 257, doi:10.3389/fmicb.2016.00257 (2016).

64.   Orita, I. et al. The ribulose monophosphate pathway substitutes for the missing pentose phosphate pathway in the archaeon Thermococcus kodakaraensis. *J Bacteriol* **188**, 4698-4704, doi:10.1128/JB.00492-06 (2006).

65.   Ferenci T, Strom T, Quayle JR. Purification and properties of 3-hexulose phosphate synthase and phospho-3-hexuloisomerase from Methylococcus capsulatus. Biochem J. 1974;144(3):477-86. Epub 1974/12/01. doi: 10.1042/bj1440477. PubMed PMID: 4219834; PubMed Central PMCID: PMCPMC1168525.

66.   Jia, P. et al. Emergence of a Novel NDM-5-Producing Sequence Type 4523 Klebsiella pneumoniae Strain Causing Bloodstream Infection in China. *Microbiol Spectr* **10**, e0084222, doi:10.1128/spectrum.00842-22 (2022).

67.   Namaki, M. et al. Prevalence of resistance genes to biocides in antibiotic-resistant Pseudomonas aeruginosa clinical isolates. *Mol Biol Rep* **49**, 2149-2155, doi:10.1007/s11033-021-07032-2 (2022).

68.   Alfarouk, K. O. *et al.* The Pentose Phosphate Pathway Dynamics in Cancer and Its Dependency on Intracellular pH. *Metabolites* **10**, doi:10.3390/metabo10070285 (2020).

69.   Horecker, B. L. The pentose phosphate pathway. *J Biol Chem* **277**, 47965-47971, doi:10.1074/jbc.X200007200 (2002).

70.   Kruger, N. J. & von Schaewen, A. The oxidative pentose phosphate pathway: structure and organisation. *Curr Opin Plant Biol* **6**, 236-246, doi:10.1016/s1369-5266(03)00039-6 (2003).

71.   Kawai, S. & Murata, K. Structure and function of NAD kinase and NADP phosphatase: key enzymes that regulate the intracellular balance of NAD(H) and NADP(H). *Biosci Biotechnol Biochem* **72**, 919-930, doi:10.1271/bbb.70738 (2008).

72.   Kelso, P. A., Chow, L. K. M., Carpenter, A. C., Paulsen, I. T. & Williams, T. C. Toward Methanol-Based Biomanufacturing: Emerging Strategies for Engineering Synthetic Methylotrophy in Saccharomyces cerevisiae. *ACS Synth Biol* **11**, 2548-2563, doi:10.1021/acssynbio.2c00110 (2022).

73.   Wang, Y., Fan, L., Tuyishime, P., Zheng, P. & Sun, J. Synthetic Methylotrophy: A Practical Solution for Methanol-Based Biomanufacturing. *Trends Biotechnol* **38**, 650-666, doi:10.1016/j.tibtech.2019.12.013 (2020).

Figure 1: Construction and analysis of a reactant-dependent algebraic framework to model and characterize the redundant forms of a biochemical network: An algebraic framework to model and characterize redundant forms of a reactant-dependent biochemical network is constructed and populated in accordance with biochemically relevant paradigms. The framework is mathematical rigorous and comprises a set of unique stoichiometry number matrices. Each matrix is constrained by a lower bound for the number of reactants and thence the selected PRVs, is populated by integral changes which are Boolean (signed non-zero, zero) and combinatorially distributed across selected PRVs and represents a biochemical network. In order to study the redundant forms of a biochemical network, the dot product of a stoichiometry number matrix with non-zero non-unitary rational number multipliers is taken. The rational number matrices that result from these computations are similarly constrained, equivalent, degenerate with respect to a shared null space and comprise the redundant forms of a modelled biochemical network (CRaTER𝕩). The matrices have well defined algebraic properties and the pathways that they represent are easily parameterized in terms of probable dissociation constants and trivially perturbed. **Abbreviations**: **CRaTER𝕩**, Library of 𝕩-stoichiometry number matrix $(\mathcal{S}_{p_{J,𝕩}} \sim p_{J,𝕩})$ specific a-indexed $\mathbb{A}$-membered rational number matrices $(\mathcal{S}_{p_{J,𝕩,a}} \sim p_{J,𝕩,a})$ which are similarly constrained, equivalent, unequal and null-space redundant; $p_{(.)}, \tilde{p}_{(.)}$, unperturbed- and perturbed-forms of a biochemical network; **PRV**, Plausible reaction vector;

Figure 2: Perturbation studies for parameter selection and assessment of relevance of the redundant forms of a biochemical network: A perturbation is modelled by the inclusion of one or more extraneous reactions and its effect on the modelled biochemical network is studied by the pairwise comparison of the probable dissociation constants between shared reactions. This is possible since there is a one-one correspondence between a pathway (reactants, reactions) and its encoding rational number matrix (reactants, RRVs). The probable dissociation constant is a purely numerical measure that is computed by the R-package "ReDirection" for every reaction of a user-defined biochemical pathway. The metrics for these analyses are the numbers and proportions (matched, unmatched) of equivalent and non-equivalent reactions between the unperturbed- and perturbed-versions of a redundant form of an x^{th}-biochemical network (a-indexed CRaTERx-matrix). These network-derived characteristics are reformulated as metrics to compare the redundant forms of a biochemical network. These include $\Delta e\{(\wp_{J,x,a}, \tilde{\wp}_{J,x,a}), (\wp_{J,x,b}, \tilde{\wp}_{J,x,b})\}$ for pairs of concurrently considered redundant forms and $\Delta ne(\wp_{J,x,a}, \tilde{\wp}_{J,x,a})$ for single instances of the same. Both of these are computed by populating the appropriate confusion matrices with reaction-outcome data, after perturbing the system(s) and computing the indices (precision, accuracy, recall) which are then assessed. **Abbreviations**: **CRaTERx**, Library of x-stoichiometry number matrix ($\mathcal{S}_{\wp_{J,x}} \sim \wp_{J,x}$) specific a, b indexed $\mathbb{A}$-membered rational number matrices ($\mathcal{S}_{\wp_{J,x,a}} \sim \wp_{J,x,a}; \mathcal{S}_{\wp_{J,x,b}} \sim \wp_{J,x,b}$) which are similarly constrained, equivalent, unequal ($a \neq b$) and null-space redundant; $\wp_{(.)}, \tilde{\wp}_{(.)}$, unperturbed- and perturbed-forms of a biochemical network; $\Delta e, \Delta ne$, Change in fraction of reactions with equivalent (e)- or non-equivalent (ne)-outcomes of a biochemical network after being perturbed; **PRV**, Plausible reaction vector; **RRV**, Rational number reaction vector;

Figure 3: Computational studies to model, evaluate and establish biochemical and physiological relevance for a reactant-dependent algebraic framework: a), **b)** The urea cycle is characterized by several shunt pathways via Aspartate (Kreb's cycle, r_6) and Carbamoyl phosphate/Glutamate (Kreb's cycle, r_7; D-amino acids, r_5; Arginine-Proline, r_8). The synthesis of Citrulline (r_3) with Nitric oxide synthase (NOS, EC1.14.13.39) utilizes NADPH and nitric oxide which is an important member of reactive oxygen (RO)- and reactive nitrogen (RN)-species (ROS, RNS). In this study, we investigate the effects of converting Arginine to Glutamate (r_{12}) and thence to urea and Ornithine. Ornithine, in turn, is a critical precursor in the biosynthetic pathway of the polyamines (Putrescine, Spermidine, Spermine). These compounds are involved in DNA synthesis, cell proliferation and growth and the regulation of a subclass of potassium receptors and **c), d)** The folate cycle (THF, 5-methylTHF and 5,10-methyleneTHF) is responsible for the movement and exchange of several small molecules (methyl groups, vitamins, amino acids) between DNA/RNA, proteins and lipids. We examine effects of a direct and equally probable interconversion between dTMP to dUMP (r_{18}) on a biochemical network of folate metabolism. **Abbreviations**: **CRaTER$\mathbb{x}$**, Library of $\mathbb{x}$-stoichiometry number matrix ($\boldsymbol{S}_{\boldsymbol{p}_{J,\mathbb{x}}} \sim \boldsymbol{p}_{J,\mathbb{x}}$) specific a-indexed $\mathbb{A}$-membered rational number matrices ($\boldsymbol{S}_{\boldsymbol{p}_{J,\mathbb{x},a}} \sim \boldsymbol{p}_{J,\mathbb{x},a}$) which are similarly constrained, equivalent and redundant; $\eta_i(\boldsymbol{p}_{J,\mathbb{x},a}), \tilde{\eta}_i(\tilde{\boldsymbol{p}}_{J,\mathbb{x},a})$, probable dissociation constant for a shared reaction between pathways or redundant forms of a biochemical networks; **mne,** shared and matched non-equivalent reactions; **me,** shared and matched equivalent reactions; **THF**, Tetrahydrofolate;

Figure 4: Biomedical relevance of the redundant forms of a biochemical network for the HMP-shunt pathway in the genesis of formaldehyde resistance: a), b) Analysis of a biochemical network of the hexose monophosphate shunt pathway in *K. pneumoniae* (ST4523) before and after the introduction of r_{19}. In this study the direct conversion of β-D fructose 6-phosphate to D-ribulose-5-phosphate (r_{19}) is investigated in terms of the magnitude and distribution of probable dissociation constants for the shared equivalent- and non-equivalent reactions of redundant forms of the biochemical network for the HMP-pathway (*Klebsiella pneumoniae*, *Klebsiella pneumoniae* ST4523). Although this conversion is absent in humans, the reaction is a significant contributor to formaldehyde fixation in bacteria and some archaea. The reactions occur via D-arabino hex-3-ulose 6-phosphate and includes the enzymes (6-phospho-3-hexuloisomerase, EC5.3.1.27; 3-hexulose-6-phosphate synthase, EC4.1.2.43) as independent polypeptides (archaea) or bifunctional protein (methylotrophic bacteria) and **c)** Schematic diagram for a model of enhanced nosocomial infection. Here, an increased virulence and resistance in *Klebsiella pneumoniae* and other non-methylotrophic bacteria such as *Pseudomonas aeruginosa* and *Escherichia coli* may occur if genes to support the RuMP pathway were horizontally acquired. These bacteria would then possess the RuMP-pathway in addition to the conserved HMP-shunt pathway and formaldehyde detoxifying genes (*adhC*, *estD*). This may result in bacteria that could thrive on formaldehyde exposure in a hospital setting (host-pathogen interface, formaldehyde-sterilized endoscopic instruments). **Abbreviations**: **EC**, enzyme commission; **F6P**, Fructose-6-phosphate; **G6P**, Glucose-6-phosphate; **NADPH**, reduced Nicotinamide adenine dinucleotide phosphate; **mM**, Millimolar; (**Pa⁺, Kp⁺**), *Pseudomonas aeruginosa* and *Klebsiella pneumoniae* with horizontally acquired genes to support the RuMP pathway; **PRPP**, phosphoribosyl pyrophosphate; **RuMP**, Ribulose-5-phosphate;

Table 1: Glossary to define, construct, describe and analyse an algebraic framework to model and characterize redundant forms of a biochemical network

	Glossary	Rmks
1.	Non-equivalent forward reaction $\left(\eta \in \mathbb{R} \cap (1, +\infty)\right)$ Non-equivalent reverse reaction $\left(\eta \in \mathbb{R} \cap (0,1)\right)$ Equivalent reaction $(\eta \in \mathbb{R} \cap \{1\})$ j^{th}-reactant of a biochemical network Number of reactants for a biochemical network Number of reactant classes considered concomitantly Class specific stoichiometry number matrices i^{th}-reaction vector of a biochemical network Number of reaction vectors for a biochemical network Subset of selected PRVs for a biochemical network Collection of unique reactant-dependent stoichiometry number matrices RRVs for a biochemical network derived random number matrix Library of unique equivalent biochemical network-specific random number matrices Redundant form of biochemical network (unperturbed) Redundant form of biochemical network (perturbed)	$\left.\begin{array}{l} f \\ b \\ r \end{array}\right\}$ Dissociation constant based outcomes $\left.\begin{array}{l} m_j \\ J \\ \mathbb{J} \\ \mathfrak{S} \end{array}\right\}$ Reactant relevant details $\left.\begin{array}{l} r_i \\ I \\ \hat{I} \\ \mathfrak{x} \end{array}\right\}$ Plausible reaction vector based $\left.\begin{array}{l} \bar{\bar{I}} \\ \mathbb{C} \\ \wp \\ \tilde{\wp} \end{array}\right\}$ Rational number based reaction vec
2.	Generic example of a bimolecular reaction (RXN) Stoichiometry numbers as linear maps from reaction-specific dissociation constants Reactants as a finite tuple of identical molecules or species Integral contribution of reactant species to a reaction vector (RV)	$(\textbf{Eqs. 1} - \textbf{6})$ **D1** $(\textbf{Eqs. 7} - \textbf{23})$ **D2** $(\textbf{Eqs. 24, 25})$ **D3** $(\textbf{Eqs. 26} - \textbf{28})$ **D4, D5**
3.	Constraints to define and populate a PRV Constraints to describe an algebraic model for a biological networks Constraints to define indices of biochemical function Rational number multipliers of the stoichiometry number matrices	$\text{Cs1} - \text{Cs9}\,(\textbf{Eqs. 29} - \textbf{60})$ **D6** − **D21** $\text{Cs10} - \text{Cs16}\,(\textbf{Eqs. 61} - \textbf{108})$ **D23** − **D36** $\text{Cs17} - \text{Cs20}\,(\textbf{Eqs. 109} - \textbf{120})$ **D38** − **D41** $(\textbf{Eqs. 121} - \textbf{155})$ **D42** − **D59**

	Glossary	Rmks	
4.	Number of forward reactions for an x^{th}-biochemical network (unperturbed) Number of matched (m) forward reactions after perturbation Number of reverse reactions for an x^{th}-biochemical network (unperturbed) Number of matched (m) reverse reactions after perturbation Number of non-equivalent (ne) reactions for an x^{th}-biochemical network (unperturbed) Number of matched (m) non-equivalent (ne) reactions after perturbation Number of unmatched (u) non-equivalent (ne) reactions after perturbation Number of equivalent (e) reactions for an x^{th}-unperturbed biochemical network Number of matched (m) equivalent (e) reactions after perturbation Number of unmatched (u) equivalent (e) reactions after perturbation	$f_{\wp_{J,x,a}}$ $mf_{\wp_{J,x,a}}$ $b_{\wp_{J,x,a}}$ $mb_{\wp_{J,x,a}}$ $ne_{\wp_{J,x,a}}$ $mne_{\wp_{J,x,a}}$ $une_{\wp_{J,x,a}}$ $e_{\wp_{J,x,a}}$ $me_{\wp_{J,x,a}}$ $ue_{\wp_{J,x,a}}$	**D60** **D61** **D62** **D63** **D64** **D65** **D66** **D67** **D68** **D69**
5.	Number of equivalent reactions with are unaltered (e→e) (TP; $\mathbb{T}^1_{J,x,a}$) Number of altered equivalent reactions (e→{f,b}=ne) (FP; $\mathbb{T}^1_{J,x,a}$) Number of reactions which are reversed (f⇌b) or altered (ne={f,b}→e) (FN; $\mathbb{T}^1_{J,x,a}$) Number of non-equivalent reactions which are unaltered (ne→ne) (TN; $\mathbb{T}^1_{J,x,a}$) Number of non-equivalent reactions which are unaltered (ne→ne) (TP; $\mathbb{T}^2_{J,x,a}$) Number of reactions which are reversed (f⇌b) or altered (ne={f,b}→e) (FP; $\mathbb{T}^2_{J,x,a}$) Number of altered equivalent reactions (e→{f,b}=ne) (FN; $\mathbb{T}^2_{J,x,a}$) Number of equivalent reactions which are unaltered (e→e) (TN; $\mathbb{T}^2_{J,x,a}$)	$m^{\mathbb{T}1}e_{\wp_{J,x,a}}$ $u^{\mathbb{T}1}e_{\wp_{J,x,a}}$ $u^{\mathbb{T}1}ne_{\wp_{J,x,a}}$ $m^{\mathbb{T}1}ne_{\wp_{J,x,a}}$ $m^{\mathbb{T}2}ne_{\wp_{J,x,a}}$ $u^{\mathbb{T}2}ne_{\wp_{J,x,a}}$ $u^{\mathbb{T}2}e_{\wp_{J,x,a}}$ $m^{\mathbb{T}2}e_{\wp_{J,x,a}}$	**D76** **D77** **D78** **D79** **D80** **D81** **D82** **D83**

Abbreviations:

      D: Definition

      A: Constraint

    Eqs.: Equations

      η: Probable dissociation constant for a reaction

      e: Equivalent reaction

ne:	Non-equivalent reaction {f, b}
$\mathcal{S}$:	Stoichiometry number matrix
$\mathbb{T}^1_{J,\mathbb{x},a}$:	Confusion matrix with equivalence as the positive outcome
$\mathbb{T}^2_{J,\mathbb{x},a}$:	Confusion matrix with non-equivalence as the positive outcome
TP:	True positive
FP:	False positive
FN:	False negative
TN:	True negative

Table 2: Numerical case studies for a J-reactant dependent algebraic model to model and characterize the redundant forms of a biochemical network

Reactants (J)	RV (2^J)	PRVs $(2^J - 2(1+J))$	Classes $(J - 3\ (\mathbb{J}))$	Set(s) of reactants $\left(\bar{\theta}_{\hat{\jmath}_J}\right)$	Set(s) of PRVs $\left(\hat{I}_{\hat{\jmath}_J}\right)$	Set(s) of SNMs $(\#\mathfrak{X}_J = \mathbb{X})$	CRaTERx $\left(\begin{matrix}\#\mathfrak{C}_{J,\mathbb{x}} = \mathbb{A}\\ \mathbb{x} \in [1, \mathbb{X}]\end{matrix}\right)$
4	16	6	1	$\bar{\theta}_{\hat{\jmath}_4} = \left\{\bar{\theta}_{\hat{\jmath}_{4,1}} = 2\right\}$	$\hat{I}_{\hat{\jmath}_4} = \left(\hat{I}_{\hat{\jmath}_{4,1}} = 6\right)$	$\#\mathfrak{X}_4 = \left(\mathfrak{S}_{\hat{\jmath}_{4,1}}(2,6)\colon \binom{6}{2} = 15\right)$	$\mathfrak{C}_{J,\mathbb{x},a}$ $a = 1, \ldots, \mathbb{A}$ $\mathbb{x} \in [1,15]$
5	32	20	2	$\bar{\theta}_{\hat{\jmath}_5} = \begin{pmatrix}\bar{\theta}_{\hat{\jmath}_{5,1}} = 2\\ \bar{\theta}_{\hat{\jmath}_{5,2}} = 3\end{pmatrix}$	$\hat{I}_{\hat{\jmath}_5} = \begin{pmatrix}\hat{I}_{\hat{\jmath}_{5,1}} = 10\\ \hat{I}_{\hat{\jmath}_{5,2}} = 10\end{pmatrix}$	$\#\mathfrak{X}_5 = \begin{pmatrix}\mathfrak{S}_{\hat{\jmath}_{5,1}}(2,10)\colon \binom{10}{2} = 45\\ \mathfrak{S}_{\hat{\jmath}_{5,2}}(3,10)\colon \binom{10}{3} = 120\end{pmatrix} = 165$	$\mathfrak{C}_{J,\mathbb{x},a}$ $a = 1, \ldots, \mathbb{A}$ $\mathbb{x} \in [1,165]$
6	64	50	3	$\bar{\theta}_{\hat{\jmath}_6} = \begin{pmatrix}\bar{\theta}_{\hat{\jmath}_{6,1}} = 2\\ \bar{\theta}_{\hat{\jmath}_{6,2}} = 3\\ \bar{\theta}_{\hat{\jmath}_{6,3}} = 4\end{pmatrix}$	$\hat{I}_{\hat{\jmath}_6} = \begin{pmatrix}\hat{I}_{\hat{\jmath}_{6,1}} = 15\\ \hat{I}_{\hat{\jmath}_{6,2}} = 20\\ \hat{I}_{\hat{\jmath}_{6,3}} = 15\end{pmatrix}$	$\#\mathfrak{X}_6 = \begin{pmatrix}\mathfrak{S}_{\hat{\jmath}_{6,1}}(2,15)\colon \binom{15}{2} = 105\\ \mathfrak{S}_{\hat{\jmath}_{6,2}}(3,20)\colon \binom{20}{3} = 180\\ \mathfrak{S}_{\hat{\jmath}_{6,3}}(4,15)\colon \binom{15}{4} = 1365\end{pmatrix} = 1650$	$\mathfrak{C}_{J,\mathbb{x},a}$ $a = 1, \ldots, \mathbb{A}$ $\mathbb{x} \in [1,1650]$
7	128	112	4	$\bar{\theta}_{\hat{\jmath}_7} = \begin{pmatrix}\bar{\theta}_{\hat{\jmath}_{7,1}} = 2\\ \bar{\theta}_{\hat{\jmath}_{7,2}} = 3\\ \bar{\theta}_{\hat{\jmath}_{7,3}} = 4\\ \bar{\theta}_{\hat{\jmath}_{7,4}} = 5\end{pmatrix}$	$\hat{I}_{\hat{\jmath}_7} = \begin{pmatrix}\hat{I}_{\hat{\jmath}_{7,1}} = 21\\ \hat{I}_{\hat{\jmath}_{7,2}} = 35\\ \hat{I}_{\hat{\jmath}_{7,3}} = 35\\ \hat{I}_{\hat{\jmath}_{7,4}} = 21\end{pmatrix}$	$\#\mathfrak{X}_7 = \begin{pmatrix}\mathfrak{S}_{\hat{\jmath}_{7,1}}(2,21)\colon \binom{21}{2} = 210\\ \mathfrak{S}_{\hat{\jmath}_{7,2}}(3,35)\colon \binom{35}{3} = 6545\\ \mathfrak{S}_{\hat{\jmath}_{7,3}}(4,35)\colon \binom{35}{4} = 52{,}360\\ \mathfrak{S}_{\hat{\jmath}_{7,4}}(5,21)\colon \binom{21}{5} = 20{,}349\end{pmatrix} = 79464$	$\mathfrak{C}_{J,\mathbb{x},a}$ $a = 1, \ldots, \mathbb{A}$ $\mathbb{x} \in [1,79464]$

Abbreviations:

J:	Reactants
RV	Reaction vector
PRV:	Plausible reaction vector
SNM:	Stoichiometry number matrix
CRaTERx:	Constrained, rational number, equivalent and redundant matrices for the $\mathbb{x}^{\text{th}}$-biochemical network

Table 3: "ReDirection"-based analysis for a redundant form (unperturbed, perturbed) of known biochemical networks

	Biochemical network	Type	$\hat{I}_x$	J	$\mathcal{V}$	#F	#B	#E	Time (s^{-1})
1.	Urea cycle	$\wp_{J,x,a}$	11	9	2	2	4	5	0.09
		$\tilde{\wp}_{J,x,a}$	12	9	3	8	1	3	1.25
2.	Folate metabolism	$\wp_{J,x,a}$	17	15	3	8	3	6	0.05
		$\tilde{\wp}_{J,x,a}$	18	15	3	8	3	7	0.06
3.	HMP shunt pathway (*K. pneumoniae* spp.)	$\wp_{J,x,a}$	18	16	3	6	8	4	25.68
		$\tilde{\wp}_{J,x,a}$	19	16	4	6	9	4	25.07

Abbreviations:

$\hat{I}_x$:   Number of PRVs for a biochemical network

PRV: Plausible reaction vector for stoichiometry number matrix

J: Number of reactant for a biochemical network

$\mathcal{V}$:   Initial nullity of null space-generated subspace of stoichiometry number matrix for a biochemical network

$\wp_{J,x,a}$:   Unperturbed form of a biochemical network

$\tilde{\wp}_{J,x,a}$:   Perturbed form of a biochemical network

#F: Number of forward reactions for a biochemical network

#B: Number of reverse reactions for a biochemical network

#E: Number of equivalent reactions for a biochemical network

Table 4: Evaluating the confusion matrices for a redundant form (unperturbed, perturbed) of known biochemical networks

Confusion matrix	Network	TP	FP	FN	TN	Pr	Re	Ac	SPy
$\mathbb{T}^1_{J,x,a}$	Urea	3	2	4	2	60%	$\approx 43\%$	$\approx 46\%$	50%
	Folate	6	0	4	7	100%	60%	$\approx 77\%$	100%
	HMP (K.pa$^+$)	4	0	5	9	100%	$\approx 44\%$	$\approx 72\%$	100%
$\mathbb{T}^2_{J,x,a}$	Urea	2	4	2	3	$\approx 33\%$	50%	$\approx 46\%$	$\approx 43\%$
	Folate	7	4	0	6	$\approx 64\%$	100%	$\approx 77\%$	60%
	HMP (K.pa^{++})	9	5	0	4	$\approx 64\%$	100%	$\approx 72\%$	$\approx 44\%$

Abbreviations

$\mathbb{T}^1_{J,x,a}$: Confusion matrix with equivalence as the positive outcome

$\mathbb{T}^2_{J,x,a}$: Confusion matrix with non-equivalence as the positive outcome

TP: True positive

FP: False positive

FN: False negative

TN: True negative

Pr: Precision or positive predictive value

Re: Recall or sensitivity of test

Ac: Accuracy of test

SPy: Specificity of test

K.pa$^+$: *K. pneumoniae*

K.pa^{++}: *K. pneumoniae (ST4523)*