The sulfur-related metabolic status of *Aspergillus fumigatus* during infection reveals cytosolic serine
hydroxymethyltransferase as a promising antifungal target.

In vivo transcriptomics to identify new antifungal targets and validation of SHMT

Reem Alharthi[1], Monica Sueiro-Olivares[1], Isabelle Storer[1], Hajer Alshammri[1], Jennifer Scott[1], Reem Al-
Shidhani[2], Rachael Fortune-Grant[1], Elaine Bignell[3], Lydia Tabernero[2], Michael Bromley[1], Can Zhao[4], and
Jorge Amich[1,5]*

1. Manchester Fungal Infection Group (MFIG), Division of Evolution, Infection, and Genomics,
Faculty of Biology, Medicine and Health, University of Manchester, Manchester, M13 9PT UK.
2. School of Biological Sciences, Faculty of Biology Medicine and Health, Manchester Academic
Health Science Centre, University of Manchester, Manchester M13 9PT, UK
3. MRC Centre for Medical Mycology, University of Exeter, Exeter, United Kingdom.
4. Department of Life Sciences, Manchester Metropolitan University, Manchester, M1 5GD,
United Kingdom.
5. Mycology Reference Laboratory (Laboratorio de Referencia e Investigación en Micología
LRIM), National Centre for Microbiology, Instituto de Salud Carlos III (ISCIII), Majadahonda,
28220 Madrid, Spain.

*Correspondence to Jorge Amich: jamich@isciii.es

ABSTRACT

Sulfur metabolism is an essential aspect of fungal physiology and is known to be crucial for

pathogenicity. Fungal sulfur metabolism comprises anabolic and catabolic routes that are not well-

conserved in mammals, and therefore can be considered a promising source of prospective novel

antifungal targets. To gain insight into the status of the *Aspergillus fumigatus* sulfur-related

metabolism during infection we used a NanoString custom nCounter TagSet and compared the

expression of 68 key metabolic genes in different murine models of invasive pulmonary aspergillosis,

at three different time-points, and a variety of *in vitro* conditions. We identified a set of 15 genes that

are consistently expressed at higher levels *in vivo* than *in vitro*, suggesting that they may be particularly

relevant for intrapulmonary growth and therefore constitute promising drug targets. Indeed, the role

of five of the fifteen genes had previously been empirically validated, supporting the likelihood that

the remaining candidates are relevant. In addition, the analysis of the dynamics of gene expression at

the early (16h), mid (24h) and late (72h) time-points uncovered potential disease initiation and

progression factors. We further characterised one of the identified genes, encoding the cytosolic

serine hydroxymethyltransferase ShmB, and demonstrated that it is an essential gene of *A. fumigatus*

and that it is also required for virulence in a murine model of established pulmonary infection. We

further show that the structure of the ligand binding pocket of the fungal enzyme differs significantly

from its human counterpart, suggesting that specific inhibitors can be designed. Therefore, *in vivo*

transcriptomics is a powerful tool to identify genes crucial for fungal pathogenicity that might encode

promising antifungal target candidates.

AUTHOR SUMMARY

Aspergillus fumigatus is an opportunistic human fungal pathogen that causes devastating chronic and

invasive infections in immunocompromised patients. Our arsenal of antifungal drugs to fight this and

other fungal pathogens is very limited, partly because of the high similarity between eukaryotic fungal

and human cells makes the identification of suitable drug targets a challenging task. Furthermore,

targets identified *in vitro* are often not effective *in vivo*, as their action is not relevant for fungal

virulence. To address this challenge, we compared the expression profiles of a set of genes involved

in sulfur metabolism, a promising source of potential drug targets, in numerous *in vitro* and *in vivo*

conditions to identify favourable antifungal candidates. Subsequently, we validated one of the

highlighted genes, demonstrating that it is essential for *A. fumigatus* viability and virulence, and that

it can likely be targeted by specific inhibitors. Hence, we show the potential of using *in vivo*

transcriptomics to identify targets that contribute to virulence, propose various candidates for future

studies and present a novel target validated for further antifungal drug development.

INTRODUCTION

Aspergillus fumigatus is a ubiquitous mould that normally lives in the soil feeding on decaying organic matter [1]. During its natural life-cycle *A. fumigatus* produces thousands of spores, also known as conidia, that are disseminated in the air. Because of their high abundance and prevalence, it has been estimated that every human breathes in several hundred of these conidia on a daily basis [2] and, due to their small size (2-3 µm), they have the potential to penetrate deep into the respiratory tract and even reach the lung alveoli [3]. In immunocompetent individuals, this has no consequence as epithelial mucocilliarity and a proper immune response eliminate the spores very efficiently [4]. However, patients with an imbalanced immune response are susceptible to *A. fumigatus*, which can cause a wide range of diseases collectively termed aspergilloses [5]. Chronic and invasive aspergilloses are life-threatening infections with high mortality rates, even in patients under antifungal treatment [6, 7]. First line therapy against aspergilloses is based on the use of azoles, the only class of antifungals that can be orally administered. Worryingly, the number of clinical *A. fumigatus* isolates that are resistant to triazole antifungals is increasing worldwide, which correlates with higher mortality rates [6, 8, 9]. Consequently, new agents to fight this fungal pathogen are urgently required.

Metabolism is central to the virulence of fungal pathogens, and therefore its targeting is considered a promising strategy for the development of novel antifungals [10, 11]. Fungal sulfur metabolism is particularly interesting as it comprises routes and enzymes that are often not conserved in humans, and thus it is considered a propitious source of novel antifungal targets [12]. Indeed, we have previously shown that a proper regulation of sulfur metabolism is crucial for *A. fumigatus* infective capacity [13], demonstrated that biosynthesis of the S-containing amino acids is required for virulence [14], and comprehensively characterised an enzyme in the trans-sulfuration pathway, methionine synthase, as a promising antifungal target [15]. Others have also shown that sulfur is involved in the biosynthesis of essential fungal metabolites [16] and proposed sulfur assimilation and the trans-sulfuration pathway as sources of antifungal targets [17, 18]. Here, we aimed to expand on that knowledge by using *in vivo* transcriptomics to better characterise the fungal sulfur metabolic status during infection of the murine lung, and to use the expression profile of genes to identify those that may be relevant for infection, proposing them as promising candidates for further investigations. However, detection and analysis of the fungal transcriptome during infection is a persistent challenge, as fungal RNA usually represents only between 0.05 and 0.5% of the total RNA isolated from infected organs [19]. To overcome this obstacle, we have used NanoString nCounter, a novel probe-based technology with excellent reproducibility, sensitivity, and specificity when compared to other transcriptomic methods such as microarray hybridisation or RT-qPCR [20, 21]. Its high sensitivity, being capable of detecting mRNA concentrations in the femtomolar range [22], makes this technology ideal

for detecting and quantifying fungal RNA during infection. Furthermore, its incomparable specificity is

fundamental to prevent any crosstalk with the abundant host RNA. Using this technology, we have

assessed the transcription profile of a selection of metabolic genes during infection of the murine lung

in two models of immunosuppression and at three time-points post infection, and compared them

with the expression profiles under a variety of *in vitro* conditions. By these means we have uncovered

interesting aspects of the *in vivo* sulfur metabolic status and identified genes that can be proposed as

relevant for growth in the tissues. We have further characterised a promising candidate, the cytosolic

serine hydroxymethyltransferase (*shmB*) encoding gene, demonstrating that it is essential for *A.*

fumigatus viability as well as for virulence and that it is promising to aim for the development of

fungal-specific inhibitors against this target, which can serve to address the emerging global threat of

antifungal resistance [23, 24].

RESULTS

Linear amplification of Aspergillus fumigatus mRNAs allows gene expression profiling of low input in

vivo samples.

To determine the transcriptional status of genes related to sulfur metabolism during fungal infection

of the murine lung, we developed a custom NanoString nCounter Elements TagSet comprising 68

metabolic genes (51 directly related to S-metabolism and 17 metabolic genes of interest indirectly

related with S-metabolism) and 4 housekeeping controls (Table S1). We isolated RNA from *A.*

fumigatus propagated under a variety of *in vitro* conditions (Table S2) and from the lungs of infected

mice immunosuppressed with two different regimens, the leukopenic and the corticosteroid model

(Table S2 and see material and methods). We isolated the lungs of infected mice at three different

time-points to investigate gene expression at an early (16h), medium (24h) and late (72h) time point

of infection (3 mice per time-point). We further assayed a chronic model of pulmonary aspergillosis

[25] to determine if this model provides a significantly different S-environment (Table S2). A

preliminary experiment revealed that fungal RNA could not be reliable detected at 16h and 24h post-

infection, which is expected as the fungal burden at those early times is very low. To circumvent this

problem, we applied the Nanostring Single Cell Gene Expression protocol (see Material and Methods)

to perform linear amplification using Multiplexed Target Enrichment (MTE) primers. To verify that the

MTE enrichment does not affect the relative abundance of transcripts, we compared the fold changes

between diluted-amplified and non-amplified (undiluted) *in vitro* (-S limiting conditions) and *in vivo*

samples (lungs extracted from two leukopenic mice 72h after infection) (Fig 1A). This direct

comparison demonstrated that the correlation of the fold changes detected with non-amplified and

amplified samples was very high ($R^2=0.95$ for mouse 1 and $R^2=0.84$ for mouse 2). Therefore, MTE

enrichment of low input fungal RNA samples does not affect the relative abundance of transcripts and

can be used to investigate gene expression in the mammalian lung at early time-points after infection.

Fig. 1

A

B

Figure 1. Identification of genes potentially relevant *for Aspergillus fumigatus* virulence using NanoString *in vivo* transcriptomics.

A) Validation of the linear amplification performed with the Nanostring Single Cell Gene Expression protocol with Multiplexed Target Enrichment (MTE) primers. The fold changes between diluted-amplified and non-amplified (undiluted) *in vitro* (-S limiting conditions) and *in vivo* samples (lungs extracted from two leukopenic mice 72h after infection) were compared. For both mice, there was a linear relationship between the FCs of non-amplified and amplified samples, demonstrating that amplification did not affect the relative abundance of transcripts. **B)** Hierarchical clustering of all samples formed two distinct branches, one of *in vitro* samples and one with the *in vivo* samples. There were 15 genes (red line) that were always expressed at higher levels in the *in vivo* samples, compared with all the *in vitro* samples.

The fungal S-related metabolic status in vivo is different from that of in vitro conditions, revealing genes potentially relevant for infection

To analyse the S-related fungal metabolic status *in vivo* and *in vitro*, we hybridized and read our custom NanoString codeset with isolated RNAs or amplified cDNAs (Table S2). The obtained raw data was subjected to quality control, which revealed that the samples obtained from two leukopenic mice 72h had not run properly: they were borderline for the recommended threshold of overall number of counts (10,000) and did not reach the minimum values for the positive counts reads (Fig. S1A and S1B). Consequently, these samples were excluded from all analyses. In addition, inspection of the normalized data (see Material and Methods for details) revealed that five genes (AFUA_3G06492, AFUA_1G06940, AFUA_5G08600, AFUA_4G03950 and AFUA_6G00760) had a very low number of counts (<20) in all samples. This could be due to the genes not being expressed at all or at very low levels in the tested conditions, or to failure of the NanoString designed probes to detect these mRNAs. Accordingly, these genes were also excluded from all downstream analyses.

We first performed a hierarchical clustering of all samples analysed (Fig. 1B). By this stratification, the *in vivo* samples clearly clustered separately from the *in vitro* samples, indicating that the fungal S-related metabolic status during infection is significantly different from that during growth in diverse experimental conditions This suggests that the lung provides a unique environment in terms of sulfur metabolism that is distinct from defined culture media. Based on this, we hypothesised that genes which are consistently more highly expressed *in vivo*, compared to in the great variety of tested *in vitro* conditions, may be particularly relevant for the fungal capacity to grow in the lung tissues, and

therefore for fungal virulence. We detected 15 genes which were always expressed more *in vivo* than *in vitro* (Fig. 1B, red box, Table 1). Interestingly, we and others have already demonstrated the importance of at least five of those genes for *A. fumigatus* virulence, and a further six genes had already been proposed to be potentially relevant (Table 1 and see discussion), supporting the rationale that the uncharacterized genes might be valid candidates for further analyses.

GENE ID	DESCRIPTION	VIRULENCE
AFUA_1G11980	Glutamine amidotransferase (HisfF/H)	Suspected [26]
AFUA_8G02500	Glutathione S-transferase	Suspected [27]
AFUA_3G06530	ATP sulfurylase (sC)	Unknown
AFUA_5G02180	Cysteine synthase (CysB)	Proven [14]
AFUA_3G09320	Serine hydroxymethyltransferase (SmhB)	This work
AFUA_1G17170	Taurine dioxygenase	Unknown
AFUA_3G13900	Glutamate-cysteine ligase (Gcs1)	Suspected [27]
AFUA_5G03920	bZIP transcription factor (HapX)	Proven [28]
AFUA_2G07680	L-ornithine N5-oxygenase (SidA)	Proven [29]
AFUA_4G12470	bZIP transcription factor (CpcA)	Proven [30]
AFUA_6G08790	C6 transcription factor (PrnA)	Unknown
AFUA_5G06060	SCF-complex subunit (SkpA)	Suspected [31, 32]
AFUA_8G04340	cystathionine gamma-lyase (MecB)	Proven [33]
AFUA_7G06810	L-amino acid oxidase (LaoA)	Suspected [34]
AFUA_2G07620	Cystathionine beta-synthase (mecA)	Suspected [33]

Table 1. Genes that consistently showed higher expression *in vivo* compared to any of the tested *in vitro* conditions.

Fungal sulfur related in vivo metabolism is influenced by the model of immunosuppression and the course of infection.

To investigate if the fungal S-related metabolic status is influenced by the model of immunosuppression, we performed hierarchical clustering of all the *in vivo* samples (Fig. 2A). This clustering showed that the fungal S-related status is quite variable *in vivo* (notice that the dynamic range of expression are much smaller than when comparing *in vitro* and *in vivo* samples in Fig. 1B), and is partially dependent on the model of immunosuppression, as 5 out of the 8 leukopenic mice aggregated in a separated cluster. Interestingly, the chronic model clustered in this separated branch, suggesting that the fungal S-related metabolic status in this type of infection is more similar to the status in the leukopenic model than in the steroid-induced one. Nevertheless, some samples from leukopenic mice interleaved with the samples from steroid-treated mice, demonstrating that the model of immunosuppression is not the only factor that determines the fungal metabolic status.

To investigate to what extent the fungal S-related metabolic status is influenced by the course of
infection, we performed hierarchical clustering of the samples from leukopenic and steroid-treated
mice separately (Fig 2B). In both models of suppression there was a partial aggregation of samples
according to the time point post infection, suggesting that the course of infection is mirrored by the
S-related metabolic status of the fungal pathogen.

In conclusion, the model of immunosuppression as well as the time of duration of infection appear to
influence the status of the fungal S-related metabolism. However, these variables alone or combined
cannot explain the variances among samples, indicating that other factors must be involved. For
instance, it is to be expected that the exact anatomic location of the foci of fungal growth would
account for different microenvironments that in turn affect the fungal S-related metabolism.

Fig. 2

Figure 2. The fungal S-related sulfur metabolism *in vivo* is influenced by the model of suppression and the time of infection.

A) Hierarchical clustering of all *in vivo* samples created two major branches, one almost exclusive for leukopenic samples, which only included the chronic model sample. The second major branch was divided again in two branches, one exclusive for steroid samples and the other mixed with leukopenic and steroid samples. Therefore, the model of suppression influences but is not the only factor that determines the fungal metabolic status. **B)** Separated hierarchical clustering of leukopenic or steroid samples partially aggregated by the time of infection. Therefore, the fungal S-related metabolism is influenced *in vivo* by the status of infection.

Temporal dynamics of gene expression in vivo versus in vitro reveals the existence of potential disease initiation and progression factors

Since we found that the S-related metabolic status correlates with the time period post infection, we reasoned that some genes may be expressed at different levels as the fungal infection progresses. To investigate this hypothesis, we first examined the fold change of the (geometric) mean expression of all genes in each time-point with respect to the corresponding 16h condition (Fig. 3A). This analysis revealed that indeed many genes change the level of expression during the course of infection: transcription of several genes seemed to consistently increase, a few decreased expression as infection progressed, and some appeared to increase or decrease expression at mid-infection (24h) before returning to initial levels in late infection (72h). To explore whether any of these genes that show temporal dynamic expression could be particularly important for *in vivo* growth, we analysed the fold change of all genes in each *in vitro* condition versus each of the leukopenic (Fig. 3B) or steroid-treated (Fig. 3C) samples grouped by the time after infection. As expected, we identified some genes which showed a temporal variance *in vivo* compared with *in vitro*. In detail, in leukopenic mice we observed that genes *metAT* (AFUA_2G13630), *cdoB* (AFUA_5G14410), *cysAT* (AFUA_1G09470) and *isa1* (AFUA_4G10690) had higher transcript levels *in vivo* at early (16h) than at mid (24h) or late (72h) time points after infection (Fig. 3B orange box), suggesting that these genes may be particularly important for the initiation of infection, but possibly not as relevant for progression and maintenance of infection. In contrast, *cdoA* (AFUA_1G05570) had higher *in vivo* transcription during advanced infection (Fig. 3B, dark blue box), suggesting that this gene may be relevant for growth in progressively invaded tissue. The genes *hisB* (AFUA_6G04700), *sD* (AFUA_1G10820), *metR* (AFUA_4G06530) and *hapX* (AFUA_5G03920) displayed higher expression at early (16h) and late (72h) time after infection (Fig. 3B, purple box), suggesting that they may be particularly important to initiate infection and during

established infection, but that they are not crucial for infection to progress. Similarly, in steroid treated mice, we observed that genes *sFα* (AFUA_6G08920), *sB* (AFUA_1G05020), *sA* (AFUA_3G06540), *mupA* (AFUA_4G09040), *mupB* (AFUA_7G00730), *sfnB* (AFUA_8G01010), two taurine dioxygenase encoding genes (AFUA_3G0796 and AFUA_3G01010) and one arylsulfatase encoding gene (AFUA_8G02520) had higher expression *in vivo* at mid-infection time (24h) (Fig. 3C, orange boxes), suggesting that these genes may play a role in this model for the progression of infection. Finally, genes *metAT* (AFUA_2G13630), *cysAT* (AFUA_1G09470), *isa1* (AFUA_4G10690) and *ssuA* (AFUA_7G01790) had higher expression at mid- (24h) and late (72h) times after infection (Fig. 3C, purple box), indicating that they may be more relevant for progression and maintenance of infection than for its initiation.

We therefore propose that the temporal dynamics of *in vivo* transcript levels may be useful to identify and differentiate fungal factors of disease initiation and progression, defined as products that are essential for the initiation of infection or that facilitate persistence and continued disease progression [35]. Although this hypothesis needs to be validated further by confirming the role of some of our identified candidates as true disease initiation or progression factors in appropriate models of infection, we emphasise the potential of *in vivo* transcriptomics at distinct stages of infection to detect such initiation and progression factors, as has already been proposed [35].

Fig. 3

Figure 3. Temporal dynamics of gene expression reveals the existence of potential initiation and disease progression factors.

A) Fold change of expression of genes in each time-point with respect to the correspondent 16h condition revealed that most genes changed expression during the course of infection The geometric mean value for each time-point was used. **B)** Fold change of all genes in each *in vitro* condition versus each of the leukopenic samples ordered by the time after infection identified temporal dynamics in the expression of certain genes. Four genes showed higher expression *in vivo* at 16h than at 24h and 72h after infection (orange box). One gene had had higher expression *in vivo* at 72h time (red arrow). Four genes had higher expression at 16h and 72h time after infection (purple box). **C)** Fold change of all genes in each *in vitro* condition versus each of the steroid-treated samples ordered by the time after infection identified temporal dynamics in the expression of certain genes. Nine genes had higher expression *in vivo* at 24h (orange boxes). Four genes showed higher expression at 24h and 72h after infection (purple box). Samples for 16h are clustered in green rectangles, for 24h in black rectangles and for 72h in red rectangles.

Validation of an identified gene, encoding a serine hydroxymethyltransferase (SHMT), as a promising target for antifungal drug development.

To demonstrate the value of the NanoString technology in identifying genes relevant for *A. fumigatus* virulence, and aiming to validate a novel drug target, we decided to further investigate the serine hydroxymethyltransferase encoding gene (*shmB,* AFUA_3G09320), which was consistently expressed at higher levels *in vivo* compared with *in vitro* conditions (Fig. 1B). *A. fumigatus* genome encodes two paralogues of serine hydroxymethyltransferase, ShmA (annotated as AFUA_2G07810, not included in the NanoString analysis) and ShmB. The ShmB protein contains 471 amino acids (aa) and ShmA is 537 aa in length, and both share an identity of 61.9% and similarity of 79.9% (Fig. 4A). Interestingly, ShmA has a long N-terminus region that is absent in ShmB (Fig. 4A), which prompted us to investigate the subcellular localisation of the proteins using three different public softwares: DeepMito [36], PProwler [37], and TargetP [38]. In contrast to the current annotation in FungiDB and NCBI databases, all three algorithms predicted a mitochondrial localisation for ShmA and a cytosolic for ShmB. Indeed, when we aligned the *A. fumigatus* proteins with their *Saccharomyces cerevisiae* orthologues, we observed that the yeast mitochondrial enzyme SHM1 also had a longer N-terminus in comparison to the cytosolic proteins (Fig. S2A).

Therefore, aiming to clarify the subcellular localisation of ShmA and ShmB in *A. fumigatus* we tagged

the proteins at their C-terminus with the fluorescent YFP-derivative Citrine to express them under the

control of their native promoters. In line with our predictions, we found that the ShmB-Citrine protein

localises throughout the cytoplasm, with a similar pattern to a control strain that expresses Citrine

under the control of the strong *gdpA* promoter (Fig. 4B). Regrettably, ShmA-Citrine could not be

detected, possibly reflecting very low or no expression of the encoding gene under steady-state

conditions, which would be supported by its dispensability under standard laboratory conditions (see

below).

To investigate the potential relevance of ShmA and ShmB for fungal virulence, we initially attempted

to construct single null mutants deleted for each encoding gene. We successfully constructed a *ΔshmA*

mutant, which showed no phenotype on regular laboratory conditions, growing perfectly on rich

Sabouraud and defined AMM media (Fig. S2B). In contrast, we were unable to generate a *ΔshmB*

mutant, suggesting that this might be an essential gene in *A. fumigatus*. We therefore constructed

two mutant strains in which we placed the *shmB* coding region under the control of the *tetOFF* module

[39, 40], one in the wild-type background and one in the *ΔshmA* background. The *tetOFF* system allows

downregulation of transcription of the gene of interest, in this case *shmB*, upon supplementation of

doxycycline (Dox) [39]. Addition of Dox inhibited growth of both strains (*shmB_tetOFF* and *ΔshmA-*

shmB_tetOFF) on AMM solid plates (Fig. 4C), proving that *shmB* is essential for viability independently

of its paralogue *shmA*. Gene essentiality is a conditional phenomenon, whereby particular

growth/environmental conditions may overcome the disturbances derived from the gene deficiency

[40-43]. Therefore, we attempted to reconstitute growth of the *shmB_tetOFF* strain in the presence

of Dox by supplementing compounds that could potentially be depleted in the absence of ShmB (Fig.

4D). We have previously observed that disturbances in the folate cycle causes a need of amino acid

supplementation [44], therefore, we assayed all phenotypic conditions on AMM without nitrogen

(AMM-N, to diversify and increase the presence of permeases in the membrane that ensure uptake

of amino acids and other compounds) supplemented with 1 mg/mL of all proteinogenic amino acids.

Additionally, we tried to reconstitute growth by supplementing the media with 5,10-

methylenetetrahydrofolate (CH_2-THF) and glycine (separately and combined), or with folic acid and

serine (substrates/products of the bidirectional enzymatic reaction), or with adenine (as blockade of

the folate cycle might impair purine biosynthesis [45]). In addition, we tested adding ATP to all those

supplements, as we previously showed that impairing the folate cycle *via* blocking the activity of

methionine synthase impacts cellular energetics in *A. fumigatus* [40]. The only condition that could

partly reconstitute growth, albeit not to wild-type levels, was supplementation with adenine and ATP

(Fig. 4D). None of the supplement combinations reconstituted growth of the *shmB_tetOFF* strain

under restrictive conditions (Fig. 4D). Therefore, either *shmB* is a truly essential gene, or the conditions

to overcome its essentiality are complex and yet unknown. Yet, based on these results *shmB*

constitutes a promising drug target candidate, and consequently we decided to test its value in an *in*

vivo infection model.

We have recently proposed that potential drug targets should be validated in models of established

infection, which is particularly relevant for essential genes (to confirm that essentiality cannot be

overcome during active growth in the host tissues) [40]. Therefore, we assayed our optimised protocol

to investigate targets in established *A. fumigatus* invasive pulmonary infections with the *tetOFF*

system [40]. Initially, we confirmed that Dox addition very strongly turns down *shmB* expression in

growing mycelium (Fig. S2C). Next, we infected leukopenic mice with 2×10^5 conidia of the

shmB_tetOFF strain, started Dox treatment 16h after infection and measured fungal burden in the

lungs 72h after infection. As control we also infected mice with the previously validated strain

Δcyp51B-cyp51A-tetOFF [40], which expresses *cyp51A*, the gene that encodes the target of the azole

antifungals, under the control of the same *tetOFF* system. Therefore, this control strain serves to

corroborate that downregulating a validated target (azoles are the gold standard therapy for invasive

pulmonary aspergillosis) causes a significant reduction in fungal burden, as indeed we detected (Fig.

4E). Interestingly, we found that downregulation of *shmB* in the established infection model triggered

a significant reduction in fungal burden (Fig. 4E). Hence, ShmB appears to be a valid target for the

treatment of aspergillosis infections.

Fig. 4

Figure 4. The cytosolic serine hydroxymethyltransferase is essential for viability and virulence in *Aspergillus fumigatus*.

A) Alignment of the amino acid sequences of the two serine hydroxymethyltransferase enzymes encoded in *A. fumigatus* genome. The proteins have 61.9% identity and 79.9% similarity. ShmA has a long N-terminus region that is absent in ShmB and which after further analysis (see text) suggests that ShmA is mitochondrial. **B)** A ShmB-Citrine tagged proteins localised diffused throughout the cytoplasm, similarly as the control P*gdpA*-Citrine strain. **C)** A Phenotypic assay with two independent clones of the *shmB_tetOFF* and *ΔshmA shmB_tetOFF* mutant strains. In the presence of doxycycline (Dox) the mutants did not grow, demonstrating that *shmB* is essential in *A. fumigatus*, and that ShmA cannot cover its function. Plates were incubated for 48 h at 37 °C. the experiment was repeated twice independently. **D)** Attempting to reconstitute *shmB_tetOFF* growth in restrictive conditions (+Dox), media was supplemented with various metabolites which could potentially be limited in the absence of serine hydroxymethyltransferase activity. The only condition that could partly reconstitute growth, not at wild-type level, was supplementation with ATP and adenine (Ade). Plates were incubated for 48 h at 37 °C. The experiment was repeated twice independently. **E)** Fungal burden in the lungs of leukopenic mice treated or untreated with Dox. Downregulation of *cyp51A* (encoding the target of the azoles) expression caused a significant reduction in burden (p=0.0159 in a nonparametric Mann Whitney test and p=0.0476 in an unpaired *t*-test with Welch's correction). Similarly, downregulation of *shmB* triggered a significant decrease in fungal burden (p=0.0159 in a nonparametric Mann Whitney test and p=0.0288 in an unpaired *t*-test with Welch's correction). Therefore, ShmB appears to be a valid antifungal target.Each point represents an independent mouse (biological replicate) and the graph shows the mean and the standard of the mean (SEM).

Structural-based analysis of ShmB druggability

The human serine hydroxymethyltransferase (SHMT) is considered a promising target for anti-cancer drug development [46] and accordingly various inhibitors have been developed (see for instance [47, 48]). We tested the capacity of two existing inhibitors of the human SHMT, Hit-1 [49] and SHIN-1 [48], to hamper *A. fumigatus* growth. However, these chemicals did not inhibit *A. fumigatus* growth at all at high concentrations (30 µg/mL = 65 µM of Hit1 or 16 µg/mL =16 µM of SHIN-1), and the fungus was able to grow even at very high concentrations (650 µM Hit-1 and 160 µM of SHIN-1) (Fig. S3). As our previous results with the *tetOFF* promoter indicated that ShmB is essential for *A. fumigatus* viability, we hypothesised that the inhibitors designed against the human enzyme may not be effective against the fungal enzyme, which would imply that fungal specific inhibitors that do not have cross-toxicity

could also be developed. To substantiate this hypothesis our initial step involved identifying the main

druggable binding pocket of the human SHMT based on its crystal structure as retrieved from the

Protein Data Bank (accession number PDB ID 1bj4), using a target-based *in silico* approach. We

conducted two independent analyses employing the DrugRep [50] and the PockDrug [51] engines. The

PockDrug analysis yielded ten possible binding pockets, with only six of them possessing a Druggability

Probability score exceeding 0.5 (Fig. 5A). Cross-referencing with the results from DrugRep revealed a

single common pocket from both predictions: pocket A (Fig. 5A) (Gln44 Ala383 Ile51 Glu374 Phe56

Ser381 Ala379 Glu69 Val46 Arg43 Cys384 Ala407 Gly47 Glu49 Ile382 Arg465 Glu378 Glu54 Asn55

Leu48). Next, we investigated whether this predicted binding pocket is preserved in the *A. fumigatus*

orthologue. To this purpose, we constructed a homological model of the fungal protein using the *in*

silico protein modelling AI AlphaFold [52]. Although the similarity between the sequences is

reasonably high (74.9%), indicating the protein is well preserved crossed the species, there are some

mutations in the amino acids within the key areas associated to the hypothetical binding pocket

(highlighted in red in Figure 5B). A close comparison revealed that such mutations likely altered the

conformation of the neighbouring amino acid residues hence resulting in a different topology of

pocket A compared to human SHMT, making it much shallower and smaller (Figure 5C). In particular,

the alteration in the position of Arg263 (Arg255 in *A. fumigatus* SHMT) has formed a barrier resulting

in the fungal pocket being almost half the size of the human pocket A.

Subsequently, we performed *in silico* docking using the SwissDock engine [53, 54] employing the Hit-

1 inhibitor (IC50 = 0.53 μM) [49] against pocket A of the human protein. This resulted in several

promising binding motifs occupying the entire pocket, as visually represented in Figure 5D. For

instance, in one of the top binding positions, HIT1.242 (ΔG = -7.57 kcal/mol; FullFitness = -2433.57

kcal/mol), Hit-1 nestled itself deep into the pocket, demonstrating hydrophobic interactions with the

pocket (Fig. 5E). Additionally, it also formed four hydrogen bonds distributed through the molecule

backbone (Ser`41, Gln`44, Arg`43 and Tip3`579; Fig. 5G), anchoring the ligand neatly into the pocket

(Fig. 5E). Subsequent *in silico* docking between Hit-1 and the fungal orthologue pocket A suggested

that Hit-1 could not bind as effectively. Only two binding motifs were suggested, both restricted to

the smaller *A. fumigatus* pocket (Figure 5D). Upon a closer inspection of the top binding motif, HIT1.28

(ΔG = -6.87 kcal/mol; FullFitness = -2143.87 kcal/mol; Fig. 5G), it is clear that the ligand is not

embedding deep into the pocket, indicating less hydrophobic interactions with the target. There is

only one hydrogen bond suggested between Hit-1 and the target (ASP`55; Fig. 5F), indicating a less

favourable binding compared to the human version of the protein. It also appeared that the altered

Arg255 indeed acted as a barrier, blocking Hit-1 from extending to the region proximal to Arg43 (Arg35

in *A. fumigatus* SHMT) and forming any additional hydrogen bonds (Fig. 5F). Therefore, it appears that

Hit-1 is specific to human SHMT as it cannot bind properly to the fungal orthologue, which would agree

with the observed absence of growth inhibition on *A. fumigatus* (Fig. S3). Similarly, we believe that

inhibitors that bind to the fungal, but not the human, enzyme pocket can be designed, and thus we

propose that the development of novel fungal-specific inhibitor agents against the promising SmhB

antifungal target is feasible.

Fig. 5

Pockets	Vol. Hull	Hydroph. Kyte	Polar Res	Aromtica Res	Drugg Prob	Standard Deviation
P1	667.66	0.18	0.47	0.20	0.89	0.03
P2	557.96	-0.03	0.50	0.06	0.68	0.04
P6	429.67	-0.03	0.42	0.08	0.70	0.03
P7	439.19	0.10	0.33	0.11	0.79	0.01
P8	407.80	-0.44	0.50	0.21	0.69	0.06
P9	473.76	0.22	0.40	0.00	0.73	0.07

Hydrogen bond pairs between Hit-1 and SHMT				
Distance	2.7	2.5	2.5	2.4	2.2
Hit-1	HIT1.242/O5	HIT1.242/H16	HIT1.242/O2	HIT1.242/O2	HIT1.28/ H21
Human	SER`41/HG1	GLN`44/OE1	ARG`43/HH21	TIP3`579/H1	
A. fumigatus					ASP`55/OD1
The unit of the distance measuring is Å.					

Figure 5. An inhibitor of the human SHMT does not bind properly to the fungal enzyme's ligand binding pocket

A) Binding pockets with draggability score above 0.5 predicated by DrupRep highlighted in human SHMT1 1bj4, with 'Pocket A' Zoomed in (left corner). **B)** Alignment of the amino acid sequences (partial) of the *A. fumigatus* (ShmB) and human (SHMT1) cytosolic SHMT enzymes. Different amino acids (highlighted in red rectangles) were detected surrounding the key residues Arg263 in SHMT1 (Arg255 in ShmB, highlighted in blue rectangle); and Arg43 (Arg35 in ShmB, highlighted in green rectangle). Orange arrows indicate SHMT1 residues forming hydrogen bonds with Hit-1 and purple arrow indicates ShmB residue forming hydrogen bond with Hit-1. **C)** Predicted binding pockets comparison between SHMT1 and ShmB with Arg263 (Arg255) highlighted in red. **D)** Hit-1 binding motifs suggested by SwissDock. **E)** Detailed interactions formed between Hit-1 and SHMT1. **F)** Detailed interactions formed between Hit-1 and ShmB. **G)** List of hydrogen bond pairs between Hit-1 and SHMT1 and ShmB.

DISCUSION

New antifungal agents are urgently needed, as resistance to the current drugs is widespread [55-57], posing a great threat to human health [24]. In the endeavour of developing novel and efficient chemotherapy, the selection of valid drug targets is crucial [58]. In the last few years, a variety of innovative approaches have been applied to identify suitable target candidates. For instance, comparative genomics [59], computer modelling [60], *in silico* analysis of protein interactions [61], screening of deletion libraries [62], screening of regulatable libraries [43, 63], construction of transposon insertion libraries [64, 65], or machine learning approaches [66] have been employed to identify antifungal drug targets. In this regard, *in vivo* transcriptomics has also been used frequently to understand fungal pathogenicity [34, 67-70], albeit its potential to identify suitable antifungal targets still encounters some scepticism. In a seminal study, McDonagh and colleagues [34] compared the transcriptome obtained from conidia germinating in murine lungs with germlings obtained from *in vitro* conditions expected to match the *in vivo* environment. This work could confirm that *A. fumigatus* encounters nutrient (particularly iron and nitrogen) limitation as well as alkaline and oxidative stress during initiation of infection, providing foundational cues about the conditions encountered by the fungal pathogen in the lungs. However, this study needed to use pools of 24 mice per sample, which limited the number of conditions that could be tested (for instance, only one model of immunosuppression was assayed), could only analysed the early initiation of infection (12-14 hours) and had to recover the fungal material from bronchoalveolar lavages. These limitations were mainly

due to the challenge of recovering sufficient fungal RNA material from the infected tissue for the

subsequent analysis. In a more recent study, NanoString analysis was employed to identify *A.*

fumigatus transcriptional factors which are highly expressed or strongly up-regulated during growth

in the murine lung tissues, and experimentally proved that two of them are relevant in the context of

infection [71]. Here we reasoned that by comparing the transcription levels of genes related to sulfur

metabolism in the tissues versus a variety of *in vitro* S-sources could reveal gene products that are

specifically upregulated *in vivo*, and may therefore be relevant for the infection process and constitute

promising antifungal targets.

In our analysis of genes related with sulfur metabolism, we have identified 15 genes that are

consistently expressed at higher levels in the murine lungs than in all tested *in vitro* conditions. At

least 5 of those genes have been previously proven to be important for *A. fumigatus* virulence, and

additionally 6 genes have been suspected to be relevant as well (Table 1). We previously showed that

AFUA_5G02180, encoding cysteine synthase (CysB), is important for *A. fumigatus* virulence, as a

cysteine auxotroph had reduced virulence in a murine model of infection [14]. We have also

demonstrated that AFUA_8G04340 encodes cystathionine-γ-lyase (MecB), the major persulfidating

enzyme in *A. fumigatus*, which is relevant for fungal virulence [33]. The genes AFUA_2G07680,

encoding L-ornithine-*N* ^{5}-monooxygenase (SidA), which catalyses the first committed step of

siderophore biosynthesis, and AFUA_5G03920, encoding the master transcriptional regulator for

adaptation to iron limitation (HapX), have been shown to be crucial for virulence [28, 29]. Finally, the

gene AFUA_8G04340, encoding the transcriptional activator of the cross-pathway control system of

amino acid biosynthesis (CpcA), was also reported to contribute to *A. fumigatus* virulence [30].

Therefore, we propose that the remaining genes are also likely to be important for *A. fumigatus*

virulence and thus constitute good candidates to validate experimentally. It is noteworthy that genes

that we have characterized and shown to be dispensable for virulence (AFUA_1G05020 -*sB*- and

AFUA_2G15590 -*sF*- [14]) were not detected in this analysis, further supporting the hypothesis that

the genes identified may indeed be relevant. Nevertheless, it also needs to be mentioned that several

genes that have been described as relevant for virulence in other studies were not identified as

universally important in our analysis (for instance, AFUA_6G04700 -HisB- , AFUA_4G06530 -MetR- or

AFUA_2G14210 - Ilv3A/IlvC- [13, 26, 72]), implying that our approach is not able to detect all genes

relevant for infection. Yet, our analysis has identified some of these genes as potential disease

initiation or progression factors, which opens the question of whether their relevance for infection

might be time-dependent.

In fact, our temporal analysis has revealed various potential disease initiation and progression factors.

These are defined as products that are required for the initiation of infection or for the persistence

and progression of disease [35]. Differentiating these actions is paramount for drug discovery, as it

would not be adequate to develop a drug to treat established infections that targets an initiation

factor, and similarly, it would not be appropriate using a drug that targets a progressing factor as

prophylaxis. Therefore, discerning disease initiation and progression factors would be important to

better understand the infective capacity of *A. fumigatus* and to expand our reservoir of potential

antifungal targets. In fact, progression factors, which are usually not identified in classic virulence

studies, are regarded as valuable targets, as antimicrobials are often employed to treat established

infection [35]. In this respect, we previously negated a role of *sFα* (AFUA_6G08920) and *sB*

(AFUA_1G05020) in *A. fumigatus* virulence by infecting leukopenic mice with null mutants [14], but

our current results suggest that they might be disease progression factors specifically relevant in the

steroid model (Fig. 3C).

It was unfortunate that the samples from two leukopenic mice did not run properly and had to be

excluded from the analyses. We acknowledge that this limits the value of some of our results,

particularly the comparisons among time-points and the analysis of temporal expression of genes in

leukopenic mice. However, as all the infections for the *in vivo* samples of the analysis were carried out

simultaneously, we reasoned that repeating just two mice would not provide fully consistent results.

Indeed, we tried to include the data obtained from the two leukopenic mice of the preliminary

experiment to validate the linear amplification protocol (Fig. 1A), but found that these samples

clustered completely separated from all other *in vivo* samples in the study (Fig S4), and therefore could

not be used for the model and temporal analyses. Importantly, even if they clustered separately, the

overall profile of expression relative to *in vitro* remained highly similar, such that the 15 genes

previously detected to be consistently upregulated *in vivo* VS *in vitro* were maintained with the

inclusion of these mice (Fig. S4), which shows the strength of the results. In addition, with these two

additional mice one more gene was included in the list of candidates, AFUA_5G07210. This gene

encodes a homoserine O-acetyltransferase, which has been described as important for virulence in

the fungal pathogen *Cryptococcus neoformans* [73].

One of the uncharacterised genes in our list of 15 potentially relevant genes encodes a serine

hydroxymethyltransferase (SHMT). SHMT is a pyridoxal phosphate-dependent enzyme of the folate

cycle that catalyses the reversible conversion of serine and tetrahydrofolate (THF) into glycine and

5,10-methylenetetrahydrofolate (CH_2-THF). Fungi encode two isoforms in the nuclear genome, one

cytosolic and one mitochondrial [45]. In *S. cerevisiae* both genes were disrupted singly and in

combination, generating auxotrophic viable strains [74]. In contrast to our results, the deletant of the

cytosolic SHMT encoding gene (*SHM2*) grew well in the presence of the substrates glycine and

formate. In addition, the cytosolic and mitochondrial SHMT encoding genes could also be deleted in

the ascomycete *Ashbya gossypii* [75], where supplementation with adenine or glycine could reconstitute growth of the cytosolic *SHMT* (*SHM2*) knock out. Therefore, essentiality of the cytosolic ShmB in *A. fumigatus* suggests that the metabolic implications of disrupting the one carbon-cycle are more complex in this pathogen than in other fungi. Indeed, we previously showed that in contrast to other fungi, the absence of methionine synthase, which forms a junction between the one carbon cycle and the trans-sulfuration pathway, causes an imbalance in cell energetics, possibly initiated by a cellular sensing of purine depletion, that halts fungal growth [40]. Here we found that adenine could partly reconstitute growth of the *shmB_tetOFF* strain in restrictive conditions, which resembles the phenotype we observed with *metH_tetOFF* [40]; but in contrast to the absence of MetH, ATP could not fully reconstitute growth in the absence of ShmB, reflecting that we do not understand the metabolic implications of eliminating the cytosolic serine hydroxymethyltransferase activity. Given its essentiality, we hypothesised that it could constitute a good target for the development of novel antifungals, and indeed we have validated it in a murine model of established infection using the *tetOFF* system. To our knowledge the relevance of serine hydroxymethyltransferases for fungal pathogenicity had not been investigated before. In contrast, this enzyme has been shown to be a promising antimalarial target [76] and implicated in a variety of virulence related traits in bacteria. For instance, a null mutant in *Staphylococcus aureus* showed impaired survival inside macrophages and reduced virulence in a *Galleria mellonella* infection model [77]. In *Helicobacter pylori*, deletion of the SHMT encoding gene *glyA* caused reduced growth rate [78], and in *Pseudomonas aeruginosa* controls rugose colony morphology, increases biofilm formation, abolishes swarming and relates to the redox status of the cells and the control of iron acquisition [79]. Therefore, targeting microbial SMHT enzymes seems to be a promising antimicrobial strategy. Nevertheless, as the current SHMT inhibitors have been developed against the human enzyme, specific inhibitors against the microbial proteins need to be designed. By comparing the pocket landscape of the *A. fumigatus* and human enzymes and performing a docking assay with a known inhibitor of the human SHMT, here we show that there are sufficient differences to pursue the development of a fungal specific agent. A comparable situation occurs in the case of efforts to inhibit the promising antifungal target calcineurin. The human and fungal enzymes are highly similar, such that a drug developed against the human protein, FK506, also inhibits the fungal enzyme and it is a potent antifungal agent. Current research aims to develop FK506 derivatives that are fungal specific, and much of this work is fuelled by structural insight [80-82].

In conclusion, in this study we show that *in vivo* transcriptomics is a valid strategy to identify virulence traits in *A. fumigatus*. Using the NanoString technology we have detected 15 genes as potentially relevant for pathogenicity, five of which have already been proven experimentally. We further investigate one novel gene, encoding the cytosolic serine hydroxymethyltransferase, thereby

demonstrating that this is an essential gene product in *A. fumigatus* and that it seems possible to
specifically inhibit the enzyme's activity. Therefore, we have validated cSHMT as a promising
antifungal target for future research efforts.

MATERIAL AND METHODS

Fungal strains and culture media

The *Aspergillus fumigatus* strain ATCC46645 [83] was used for the transcriptomic analysis, and the
MFIG001 isolate (*ΔKU80*) for genetic manipulation [84]. The strains were routinely grown on
Sabouraud solid medium (Oxoid) for 3-5 days at 37 °C for spore harvesting.

For phenotyping analysis, the isolates were inoculated on solid *Aspergillus* Minimal Medium (AMM,
1% glucose, 5 mM ammonium tartrate, 7 mM KCl, 11 mM KH$_2$PO$_4$, 0.25 mM MgSO$_4$, 1× Hutner's trace
elements solution [pH 5.5], 1.5% agar), supplemented as indicated in each section, and incubated at
37 °C for 2-3 days. Phenotypic experiments were performed at least twice independently.

For selection in the presence of resistance markers 50 µg ml^{-1} of hygromycin B or 100 µg ml^{-1} of
pyrithiamine (InvivoGen) were added to the AMM in the growth plates.

Mutant construction

To generate the *ΔshmA* mutant and attempt to construct the *ΔshmB* mutant, we followed a previously
optimised protocol [85]. Briefly, upstream and downstream fragments of the genes were amplified by
HIFI-PCR and fused to the Hygromycin-B (*hph*) selective cassette by fusion PCR. This deletant cassette
was used to replace the target gene by homologous recombination in the *ΔKU80* strain MFIG001.

To construct the *shmB_tetOFF* isolate, we followed CRISPR-Cas9 mediated recombination, as
previously described for *A. fumigatus* [86]. Briefly, a repair template containing the *tetOFF* module
and a pyrithiamine resistance marker (PtrA), was amplified from pSK606 [39] using Phusion green hot
start II high-fidelity PCR master mix (ThermoFisher Scientific). The repair template had 50 bp flanking
regions homologous to the upstream and downstream sequences at the ATG start codon, placing the
tetOFF module immediately 5'- of the *shmB* ORF. The sequence of the targeted gRNA (crRNA) was
(GTCCAAATGAAAGAAAACAA|TGG) and was designed using EuPaGDT [87].

To construct the fluorescent SHMT strains, we inserted the correspondent repair template, containing
Citrine and the pyrithiamine resistance marker, at the 3'-UTR of each gene using CRISPR-Cas9 as

previously descried [86]. Each repair template carried flanking homology regions to the targeted gene.

The targeted gRNAs used were (5'-GACAGGCAGGGGGTAGGTGC-3') and (5'-

GGGCACTTTCAGCCTTCCGT-3') for *shmB* and *shmA*, respectively.

All primers used in the study can be found in Table S3.

Localisation studies of ShmB-Citrine

A. fumigatus conidia were inoculated in 200 µL of filtered autoclaved AMM on 8 well chambers slides

(IBIDI) and incubated at 30 °C for 16h. Imagining was performed using a Leica SP8x laser scanning

confocal microscope equipped with a 63x (NA 1.4) HC PLAN APO CS2 oil immersion objective. The

pinhole was set to one Airy unit and the gain was set to 200. To visualise the localisation of labelled

proteins, fluorescence imaging was performed using an excitation of 516 nm and 529 nm emission

using a white light laser. Image format captures was 1024x1024 at 400 hz scan speed. Z-stacks of

germlings were taken at 1.0 microns and slices were projected as average intensity using ImageJ no

adjustments to brightness or contrast were made. Scale bars are shown.

Murine Infections

Mouse infections experiments were performed under United Kingdom Home Office project license

PDF8402B7 and approved by the University of Manchester Ethics Committee and by the Biological

Services Facility at the Faculty of Biology, Medicine and Health, University of Manchester.

Outbred CD1 male mice (22 to 26 g) were purchased from Charles Rivers and left to rest for at least

one week before the experiment. Mice were allowed access *ad libitum* to water and food throughout

the experiment.

For the leukopenic model mice were immunosuppressed with 150 mg/kg of cyclophosphamide

(Baxter) on days −3 and −1 plus 1 subcutaneous injection of 250 mg/kg hydrocortisone 21-Acetate

(Sigma-Aldrich) on day −1. For the corticosteroid model, mice were injected with a single dose of 40

mg/kg of triamcinolone (Bristol Myers Squibb) on day −1. Bacterial infections were prevented by

adding 2 g/L neomycin to the drinking water. Mice were anesthetized by exposure to 2% to 3%

isoflurane (Sigma-Aldrich) for 5 to 10 minutes and intranasally infected with 40 µL of a suspension

containing 10^5 spores for the leukopenic model and 10^6 spores for the corticosteroid model. For both

models, 3 mice per time point post infection (16, 24 or 72h) were used. At the designated time-points,

mice were sacrificed by a lethal injection of pentobarbital and the lungs were harvested to be immediately snap frozen for RNA isolation.

For the chronic model, we followed a published protocol [25, 88]. Briefly, an immunocompetent mouse was anesthetised by intraperitoneal injection of a ketamine (1%)/xylazine (0,2%) solution and intratracheally inoculated with 50 μL of a suspension of agar beads containing 5×10^7 conidia/mL. Infection was left to progress for 21 days before the mouse was sacrificed for further processing.

To monitor virulence of the *shmB_tetOFF* strain, we followed our previously optimised protocol [40]. Briefly, groups of 10 mice were infected with 40 μL of a suspension containing 2×10^5 of the *shmB_tetOFF* or the control *Δcyp51B-cyp51A-tetOFF* strain (inocula were confirmed by plating dilutions of the used suspensions). Five mice per strain received doxycycline (Dox) treatment and five PBS vehicle. Dox treatment started 16h after infection by an intraperitoneal injection (50 mg/kg) and change to Dox-containing food (Envingo Safe-Diet U8200 version 0115 A03 0.625 g/kg Doxycycline Hyclate pellets). Treatment was maintained by subcutaneous 50 mg/kg Dox injections every 12h post infection until the end of the experiment after 72h.

NanoString analysis: Preparation of samples, hybridisation and data processing

RNA from *in vitro* conditions was isolated as previously described [13]. Briefly, mycelia were shifted from a culture grown overnight in AMM to new flasks containing the defined sources described in Table S2. Mycelia were incubated for 4h in the new conditions, then filtrated trough sterile Miracloth (Merck Millipore) and snap frozen in liquid nitrogen. Mycelia were ground in the constant presence of liquid nitrogen and RNA was isolated using the RNeasy Plant Mini Kit (Qiagen) following the manufacturer's instructions and including the on column DNAse digestion step. RNA was assessed by a NanoDrop 2000 spectrophotometer (Thermo Fisher Scientific) to determine quality and concentration.

RNA from murine lungs was isolated as follows: Lungs were lyophilized for 48h in a CoolSafe ScanVac freeze drier connected to a VacuuBrand pump and subsequently ground in the presence of liquid nitrogen. The resulting powder was resuspended in TRIzol reagent (Invitrogen – Thermofisher) and processed following the manufacturer's instructions. RNA was cleaned with Phenol:Chloroform:Isoamyl (25:24:1) (Sigma) and extracted with Chloroform (Invitrogen). Subsequently, RNA was purified using the RNeasy Minikit (Qiagen), following the clean-up protocol, and including the on column DNAse digestion step.

RNA isolated from murine lungs was subjected to the NanoString nCounter Single Cell Gene Expression

protocol for linear amplification of target genes. Briefly, RNA was converted to cDNA using SuperScript

VILO MasterMix (Life Technologies). To confirm that target RNA was present in the samples, and that

cDNA has been properly generated, a Real-Time PCR was performed using the gene AFUA_4G07360

(*metH*) as target, as a RT-PCR for this transcript had been validated before [40]. Genes of interest were

linearly amplified using the Multiplex Target Enrichment (MTE) primers (Supplementary File 1) and a

PCR protocol consisting of 16 MTE amplification cycles of 94°C 15 seconds and 60°C 4 minutes, with

the TaqMan PreAmp Master Mix (Life Technologies).

cDNA samples were incubated for 2 minutes at 94°C and then snap cool on ice prior to addition to the

hybridization reaction. Hybridization was performed according to the Elements Tagset Hybridisation

setup, as detailed by Nanostring and using the NanoString custom designed probes A and B

(Supplementary File 1). The cartridge was read using the flex system, consisting of nCounter Prep

Station and the Digital Analyser.

The saw data file (Supplementary File S2) was initially subjected to a Quality Control check. This

consisted of the monitorization of the counts of the internal positive control (positive performance

for hybridization verification) and the total counts detected for each of the conditions (Fig. S1).

Raw data (Supplementary File 2) was normalised using the nSolver 4.0 software (NanoString). Data

was first subjected to Background Subtraction using the geometric mean of the negative control

counts. Subsequently, data was normalized using both the geometric mean of positive control count

and the geometric mean of the housekeeping genes (Supplementary File 3).

Normalised data was analysed using the Agglomerative Cluster – Heat map option of the nSolver 4.0

software (NanoString). Spearman Correlation was used as distance metric and Average was used as

linkage method.

Computational structure analyses

The crystal structure of the serine hydroxymethyltransferase (SHMT) was obtained from Research

Collaboratory for Structural Bioinformatics Protein Data Bank (RCSB PDB, http://www.rcsb.org) in PDB

format, referenced by its protein data bank ID 1bj4. The structure of the fungal SHMT was predicted

using AlphaFold2 [52]. The amino acid sequence was retrieved from FungiDatabase [89] and utilised

as input, with the highest scoring model (pLDDT= 96.9, pTMscore= 0.943) selected for downstream *in*

silico analysis.

For the human SHMT, DrugRep [50] and PockDrug [51] were employed for target based binding pocket

prediction. DrugRep is a receptor-based screening tool utilising CB-Dock to identify docking pockets

of the protein receptor. PockDrug utilizes the fpocket estimation method, based on holo or apo

proteins, for predicted pocket estimation. This method involves preliminary detection of cavities

capable of binding a ligand of sufficient size, though without ligand proximity information. Pockets

with a predicted druggability score above 0.5 from DrugRep were recorded and cross-referenced with

suggestions from PockDoc to determine the 'optimal' binding pocket for subsequent *in-silico* docking.

Ligand Hit-1 was drawn using Marvin JS (https://marvinjs-demo.chemaxon.com/latest/demo.html

[90]). The binding mode of Hit-1 to optimal binding pocket in SHMT was predicted using the EADock

DS based SwissDock (http://www.swissdock.ch/docking# [54]). The binding modes with the most

favourable energies were evaluated with FACTS (Fast Analytical Continuum Treatment of Solvation

[91]) and then output as clusters. The best docking pose, exhibiting maximal hydrophobicity pockets

coverage and hydrogen bonding, was manually selected. Outputs from all programmes were

visualised in PyMOL 2.5 (Schrödinger, LLC).

ACKOWLEDGEMENTS

We would like to specially thank Prof Sven Krappmann, who guided JA to discover the NanoString

technology some years ago and so inspired this work. In addition, we would like to thank Emilia

Mellado and Sven Krappmann for critical reading of the manuscript. We acknowledge the use of the

Genomic Technologies Facility (Faculty of Biology Medicine and Health, University of Manchester) for

the NanoString hybridisation and data read. Help and support from members of the Manchester

Fungal Infection Group (MFIG) and the Spanish National Centre for Mycology (LRIM) is greatly

appreciated.

This work was funded by the MRC Career Development Award (MR/N008707/1)

REFERENCES

1. Tekaia, F. and J.P. Latge, *Aspergillus fumigatus: saprophyte or pathogen?* Curr Opin Microbiol, 2005. **8**(4): p. 385-92.
2. Hospenthal, D.R., K.J. Kwon-Chung, and J.E. Bennett, *Concentrations of airborne Aspergillus compared to the incidence of invasive aspergillosis: lack of correlation.* Med Mycol, 1998. **36**(3): p. 165-8.
3. Latge, J.P., *Aspergillus fumigatus and aspergillosis.* Clin Microbiol Rev, 1999. **12**(2): p. 310-50.
4. Park, S.J. and B. Mehrad, *Innate Immunity to Aspergillus Species.* Clinical Microbiology Reviews, 2009. **22**(4): p. 535-551 DOI: 10.1128/cmr.00014-09.
5. Kosmidis, C. and D.W. Denning, *The clinical spectrum of pulmonary aspergillosis.* Thorax, 2015. **70**(3): p. 270-7 DOI: 10.1136/thoraxjnl-2014-206291.
6. Lowes, D., et al., *Predictors of mortality in chronic pulmonary aspergillosis.* Eur Respir J, 2017. **49**(2) DOI: 10.1183/13993003.01062-2016.
7. Brown, G.D., et al., *Hidden killers: human fungal infections.* Sci Transl Med, 2012. **4**(165): p. 165rv13 DOI: 10.1126/scitranslmed.3004404.
8. Lestrade, P.P., et al., *Voriconazole Resistance and Mortality in Invasive Aspergillosis: A Multicenter Retrospective Cohort Study.* Clin Infect Dis, 2019. **68**(9): p. 1463-1471 DOI: 10.1093/cid/ciy859.
9. Meis, J.F., et al., *Clinical implications of globally emerging azole resistance in Aspergillus fumigatus.* Philos Trans R Soc Lond B Biol Sci, 2016. **371**(1709) DOI: 10.1098/rstb.2015.0460.
10. Scott, J. and J. Amich, *Primary Metabolism of Human Pathogenic Fungi, Importance for Virulence and Potential for Drug Development*, in *Encyclopedia of Infection and Immunity*. 2022. p. 377-407.
11. Ene, I.V., et al., *Metabolism in Fungal Pathogenesis.* Cold Spring Harbor Perspectives in Medicine, 2014. **4**(12): p. a019695-a019695 DOI: 10.1101/cshperspect.a019695.
12. Amich, J., S. Krappmann, and A.K. Bachhawat, *Editorial: Sulphur Metabolism of Fungi - Implications for Virulence and Opportunities for Therapy.* Frontiers in Microbiology, 2020. **11** DOI: 10.3389/fmicb.2020.583689.
13. Amich, J., et al., *Regulation of sulphur assimilation is essential for virulence and affects iron homeostasis of the human-pathogenic mould Aspergillus fumigatus.* PLoS Pathog, 2013. **9**(8): p. e1003573 DOI: 10.1371/journal.ppat.1003573.
14. Amich, J., et al., *Exploration of Sulfur Assimilation of Aspergillus fumigatus Reveals Biosynthesis of Sulfur-Containing Amino Acids as a Virulence Determinant.* Infect Immun, 2016. **84**(4): p. 917-29 DOI: 10.1128/IAI.01124-15.
15. Scott, J., et al., *Targeting Methionine Synthase in a Fungal Pathogen Causes a Metabolic Imbalance That Impacts Cell Energetics, Growth, and Virulence.* mBio, 2020. **11**(5) DOI: 10.1128/mBio.01985-20.
16. Traynor, A.M., et al., *Involvement of Sulfur in the Biosynthesis of Essential Metabolites in Pathogenic Fungi of Animals, Particularly Aspergillus spp.: Molecular and Therapeutic Implications.* Frontiers in Microbiology, 2019. **10** DOI: 10.3389/fmicb.2019.02859.
17. Bachhawat, A.K. and A.K. Yadav, *Metabolic Pathways as Drug Targets: Targeting the Sulphur Assimilatory Pathways of Yeast and Fungi for Novel Drug Discovery*, in *Combating Fungal Infections*. 2010. p. 327-346.
18. Kuplińska, A. and K. Rząd, *Molecular targets for antifungals in amino acid and protein biosynthetic pathways.* Amino Acids, 2021. **53**(7): p. 961-991 DOI: 10.1007/s00726-021-03007-6.
19. Xu, W., et al., *Pathogen Gene Expression Profiling During Infection Using a Nanostring nCounter Platform*, in *Yeast Functional Genomics*. 2016. p. 57-65.

20.   Bourré, L., *Genome-Wide RNAseq and Array-Based NanoString Transcriptomic Technologies:*
Which To Use and When? - Crown Bioscience. 2020: https://blog.crownbio.com/genome-
wide-rnaseq-and-array-based-nanostring-transcriptomic-technologies.
21.   Veldman-Jones, M.H., et al., *Evaluating Robustness and Sensitivity of the NanoString*
Technologies nCounter Platform to Enable Multiplexed Gene Expression Analysis of Clinical
Samples. Cancer Research, 2015. **75**(13): p. 2587-2593 DOI: 10.1158/0008-5472.Can-15-0262.
22.   Geiss, G.K., et al., *Direct multiplexed measurement of gene expression with color-coded probe*
pairs. Nature Biotechnology, 2008. **26**(3): p. 317-325 DOI: 10.1038/nbt1385.
23.   Fisher, M.C., et al., *Tackling the emerging threat of antifungal resistance to human health.*
Nature Reviews Microbiology, 2022. **20**(9): p. 557-571 DOI: 10.1038/s41579-022-00720-1.
24.   Fisher, M.C., et al., *Worldwide emergence of resistance to antifungal drugs challenges human*
health and food security. Science, 2018. **360**(6390): p. 739-742 DOI: 10.1126/science.aap7999.
25.   Urb, M., et al., *Evolution of the Immune Response to Chronic Airway Colonization with*
Aspergillus fumigatus Hyphae. Infection and Immunity, 2015. **83**(9): p. 3590-3600 DOI:
10.1128/iai.00359-15.
26.   Dietl, A.-M., et al., *Histidine biosynthesis plays a crucial role in metal homeostasis and virulence*
of Aspergillus fumigatus. Virulence, 2016. **7**(4): p. 465-476 DOI:
10.1080/21505594.2016.1146848.
27.   Wangsanut, T. and M. Pongpom, *The Role of the Glutathione System in Stress Adaptation,*
Morphogenesis and Virulence of Pathogenic Fungi. International Journal of Molecular
Sciences, 2022. **23**(18) DOI: 10.3390/ijms231810645.
28.   Schrettl, M., et al., *HapX-mediated adaption to iron starvation is crucial for virulence of*
Aspergillus fumigatus. PLoS Pathog, 2010. **6**(9): p. e1001124.
29.   Schrettl, M., et al., *Siderophore biosynthesis but not reductive iron assimilation is essential for*
Aspergillus fumigatus virulence. J Exp Med, 2004. **200**(9): p. 1213-9.
30.   Krappmann, S., et al., *The Aspergillus fumigatus transcriptional activator CpcA contributes*
significantly to the virulence of this fungal pathogen. Mol Microbiol, 2004. **52**(3): p. 785-99.
31.   Liu, T.-B. and C. Xue, *The Ubiquitin-Proteasome System and F-box Proteins in Pathogenic Fungi.*
Mycobiology, 2018. **39**(4): p. 243-248 DOI: 10.5941/myco.2011.39.4.243.
32.   Jöhnk, B., et al., *SCF Ubiquitin Ligase F-box Protein Fbx15 Controls Nuclear Co-repressor*
Localization, Stress Response and Virulence of the Human Pathogen Aspergillus fumigatus.
PLOS Pathogens, 2016. **12**(9) DOI: 10.1371/journal.ppat.1005899.
33.   Sueiro-Olivares, M., et al., *Fungal and host protein persulfidation are functionally correlated*
and modulate both virulence and antifungal response. PLOS Biology, 2021. **19**(6) DOI:
10.1371/journal.pbio.3001247.
34.   McDonagh, A., et al., *Sub-telomere directed gene expression during initiation of invasive*
aspergillosis. PLoS Pathog, 2008. **4**(9): p. e1000154 DOI: 10.1371/journal.ppat.1000154.
35.   Cramer, R.A. and C.H. Kowalski, *Is It Time To Kill the Survival Curve? A Case for Disease*
Progression Factors in Microbial Pathogenesis and Host Defense Research. mBio, 2021. **12**(1)
DOI: 10.1128/mBio.03483-20.
36.   Savojardo, C., et al., *DeepMito: accurate prediction of protein sub-mitochondrial localization*
using convolutional neural networks. Bioinformatics, 2020. **36**(1): p. 56-64 DOI:
10.1093/bioinformatics/btz512.
37.   Hawkins, J. and M. BodÉN, *Detecting and Sorting Targeting Peptides with Neural Networks*
and Support Vector Machines. Journal of Bioinformatics and Computational Biology, 2011.
04(01): p. 1-18 DOI: 10.1142/s0219720006001771.
38.   Almagro Armenteros, J.J., et al., *Detecting sequence signals in targeting peptides using deep*
learning. Life Science Alliance, 2019. **2**(5) DOI: 10.26508/lsa.201900429.
39.   Wanka, F., et al., *Tet-on, or Tet-off, that is the question: Advanced conditional gene expression*
in Aspergillus. Fungal Genet Biol, 2016. **89**: p. 72-83 DOI: 10.1016/j.fgb.2015.11.003.

40. Scott, J., et al., *Targeting Methionine Synthase in a Fungal Pathogen Causes a Metabolic*
Imbalance That Impacts Cell Energetics, Growth, and Virulence. mBio, 2020. **11**(5) DOI:
10.1128/mBio.01985-20.
41. Beattie, S.R., et al., *Filamentous fungal carbon catabolite repression supports metabolic*
plasticity and stress responses essential for disease progression. PLoS Pathog, 2017. **13**(4): p.
e1006340 DOI: 10.1371/journal.ppat.1006340.
42. Cramer, R.A., *In vivo veritas: Aspergillus fumigatus proliferation and pathogenesis--*
conditionally speaking. Virulence, 2016. **7**(1): p. 7-10 DOI: 10.1080/21505594.2015.1134074.
43. Hu, W., et al., *Essential Gene Identification and Drug Target Prioritization in Aspergillus*
fumigatus. PLoS Pathogens, 2007. **3**(3) DOI: 10.1371/journal.ppat.0030024.
44. Scott, J., et al., *Pseudomonas aeruginosa-Derived Volatile Sulfur Compounds Promote Distal*
Aspergillus fumigatus Growth and a Synergistic Pathogen-Pathogen Interaction That Increases
Pathogenicity in Co-infection. Front Microbiol, 2019. **10**: p. 2311 DOI:
10.3389/fmicb.2019.02311.
45. Cossins, E.A. and L. Chen, *Folates and one-carbon metabolism in plants and fungi.*
Phytochemistry, 1997. **45**(3): p. 437-452 DOI: 10.1016/s0031-9422(96)00833-3.
46. Scaletti, E., et al., *Structural basis of inhibition of the human serine hydroxymethyltransferase*
SHMT2 by antifolate drugs. FEBS Letters, 2019. **593**(14): p. 1863-1873 DOI: 10.1002/1873-
3468.13455.
47. Geeraerts, S.L., et al., *The ins and outs of serine and glycine metabolism in cancer.* Nature
Metabolism, 2021. **3**(2): p. 131-141 DOI: 10.1038/s42255-020-00329-9.
48. Ducker, G.S., et al., *Human SHMT inhibitors reveal defective glycine import as a targetable*
metabolic vulnerability of diffuse large B-cell lymphoma. Proceedings of the National Academy
of Sciences, 2017. **114**(43): p. 11404-11409 DOI: 10.1073/pnas.1706617114.
49. Nonaka, H., et al., *Design strategy for serine hydroxymethyltransferase probes based on retro-*
aldol-type reaction. Nature Communications, 2019. **10**(1) DOI: 10.1038/s41467-019-08833-7.
50. Gan, J.-h., et al., *DrugRep: an automatic virtual screening server for drug repurposing.* Acta
Pharmacologica Sinica, 2022. **44**(4): p. 888-896 DOI: 10.1038/s41401-022-00996-2.
51. Hussein, H.A., et al., *PockDrug-Server: a new web server for predicting pocket druggability on*
holo and apo proteins. Nucleic Acids Research, 2015. **43**(W1): p. W436-W442 DOI:
10.1093/nar/gkv462.
52. Jumper, J., et al., *Highly accurate protein structure prediction with AlphaFold.* Nature, 2021.
596(7873): p. 583-589 DOI: 10.1038/s41586-021-03819-2.
53. Grosdidier, A., V. Zoete, and O. Michielin, *Fast docking using the CHARMM force field with*
EADock DSS. J Comput Chem, 2011. **32**(10): p. 2149-59 DOI: 10.1002/jcc.21797.
54. Grosdidier, A., V. Zoete, and O. Michielin, *SwissDock, a protein-small molecule docking web*
service based on EADock DSS. Nucleic Acids Res, 2011. **39**(Web Server issue): p. W270-7 DOI:
10.1093/nar/gkr366.
55. Lestrade, P.P.A., et al., *Triazole resistance in Aspergillus fumigatus: recent insights and*
challenges for patient management. Clinical Microbiology and Infection, 2019. **25**(7): p. 799-
806 DOI: 10.1016/j.cmi.2018.11.027.
56. Pristov, K.E. and M.A. Ghannoum, *Resistance of Candida to azoles and echinocandins*
worldwide. Clinical Microbiology and Infection, 2019. **25**(7): p. 792-798 DOI:
10.1016/j.cmi.2019.03.028.
57. Zafar, H., et al., *A titanic drug resistance threat in Cryptococcus neoformans.* Current Opinion
in Microbiology, 2019. **52**: p. 158-164 DOI: 10.1016/j.mib.2019.11.001.
58. Mota Fernandes, C., et al., *The Future of Antifungal Drug Therapy: Novel Compounds and*
Targets. Antimicrobial Agents and Chemotherapy, 2021. **65**(2) DOI: 10.1128/aac.01719-20.
59. Abadio, A.K.R., et al., *Comparative genomics allowed the identification of drug targets against*
human fungal pathogens. BMC Genomics, 2011. **12**(1) DOI: 10.1186/1471-2164-12-75.

60.   Bencurova, E., et al., *Identification of Antifungal Targets Based on Computer Modeling.* Journal
of Fungi, 2018. **4**(3) DOI: 10.3390/jof4030081.

61.   Chaudhary, R., et al., *In Silico Protein Interaction Network Analysis of Virulence Proteins*
Associated with Invasive Aspergillosis for Drug Discovery. Current Topics in Medicinal
Chemistry, 2019. **19**(2): p. 146-155 DOI: 10.2174/1568026619666181120150633.

62.   Noble, S.M., et al., *Systematic screens of a Candida albicans homozygous deletion library*
decouple morphogenetic switching and pathogenicity. Nature Genetics, 2010. **42**(7): p. 590-
598 DOI: 10.1038/ng.605.

63.   Becker, J.M., et al., *Pathway analysis of Candida albicans survival and virulence determinants*
in a murine infection model. Proceedings of the National Academy of Sciences, 2010. **107**(51):
p. 22044-22049 DOI: 10.1073/pnas.1009845107.

64.   Ianiri, G., et al., *Essential Gene Discovery in the Basidiomycete Cryptococcus neoformans for*
Antifungal Drug Target Prioritization. mBio, 2015. **6**(2) DOI: 10.1128/mBio.02334-14.

65.   Segal, E.S., et al., *Gene Essentiality Analyzed by In Vivo Transposon Mutagenesis and Machine*
Learning in a Stable Haploid Isolate of Candida albicans. MBio, 2018. **9**(5) DOI:
10.1128/mBio.02048-18.

66.   Fu, C., et al., *Leveraging machine learning essentiality predictions and chemogenomic*
interactions to identify antifungal targets. Nature Communications, 2021. **12**(1) DOI:
10.1038/s41467-021-26850-3.

67.   Chung, D., et al., *ChIP-seq and In Vivo Transcriptome Analyses of the Aspergillus fumigatus*
SREBP SrbA Reveals a New Regulator of the Fungal Hypoxia Response and Virulence. PLoS
Pathogens, 2014. **10**(11) DOI: 10.1371/journal.ppat.1004487.

68.   Cheng, S., et al., *Profiling of Candida albicans Gene Expression During Intra-abdominal*
Candidiasis Identifies Biologic Processes Involved in Pathogenesis. The Journal of Infectious
Diseases, 2013. **208**(9): p. 1529-1537 DOI: 10.1093/infdis/jit335.

69.   Xu, W., et al., *Activation and Alliance of Regulatory Pathways in C. albicans during Mammalian*
Infection. PLOS Biology, 2015. **13**(2) DOI: 10.1371/journal.pbio.1002076.

70.   Amorim-Vaz, S., et al., *RNA Enrichment Method for Quantitative Transcriptional Analysis of*
Pathogens In Vivo Applied to the Fungus Candida albicans. mBio, 2015. **6**(5) DOI:
10.1128/mBio.00942-15.

71.   Liu, H., et al., *Determining Aspergillus fumigatus transcription factor expression and function*
during invasion of the mammalian lung. PLOS Pathogens, 2021. **17**(3) DOI:
10.1371/journal.ppat.1009235.

72.   Nielsen, K., et al., *The Aspergillus fumigatus Dihydroxyacid Dehydratase Ilv3A/IlvC Is Required*
for Full Virulence. PLoS ONE, 2012. **7**(9) DOI: 10.1371/journal.pone.0043559.

73.   Nazi, I., et al., *Role of homoserine transacetylase as a new target for antifungal agents.*
Antimicrobial Agents and Chemotherapy, 2007. **51**(5): p. 1731-1736 DOI: 10.1128/Aac.01400-
06.

74.   Kastanos, E.K., Y.Y. Woldman, and D.R. Appling, *Role of Mitochondrial and Cytoplasmic Serine*
Hydroxymethyltransferase Isozymes in de Novo Purine Synthesis in Saccharomyces cerevisiae.
Biochemistry, 1997. **36**(48): p. 14956-14964 DOI: 10.1021/bi971610n.

75.   Schlüpen, C., et al., *Disruption of the SHM2 gene, encoding one of two serine*
hydroxymethyltransferase isoenzymes, reduces the flux from glycine to serine in Ashbya
gossypii. Biochemical Journal, 2003. **369**(2): p. 263-273 DOI: 10.1042/bj20021224.

76.   Schwertz, G., et al., *Antimalarial Inhibitors Targeting Serine Hydroxymethyltransferase (SHMT)*
with in Vivo Efficacy and Analysis of their Binding Mode Based on X-ray Cocrystal Structures.
Journal of Medicinal Chemistry, 2017. **60**(12): p. 4840-4860 DOI:
10.1021/acs.jmedchem.7b00008.

77.   Batool, N., et al., *Functional Identification of Serine Hydroxymethyltransferase as a Key Gene*
Involved in Lysostaphin Resistance and Virulence Potential of Staphylococcus aureus Strains.
International Journal of Molecular Sciences, 2020. **21**(23) DOI: 10.3390/ijms21239135.

78.   Antunes, L.C.M., et al., *Structural and functional insight into serine hydroxymethyltransferase*
917   *from Helicobacter pylori.* Plos One, 2018. **13**(12) DOI: 10.1371/journal.pone.0208850.
918   79.   Pu, M., et al., *Serine Hydroxymethyltransferase ShrA (PA2444) Controls Rugose Small-Colony*
919   *Variant Formation in Pseudomonas aeruginosa.* Frontiers in Microbiology, 2018. **9** DOI:
920   10.3389/fmicb.2018.00315.
921   80.   Juvvadi, P.R., et al., *Harnessing calcineurin-FK506-FKBP12 crystal structures from invasive*
922   *fungal pathogens to develop antifungal agents.* Nature Communications, 2019. **10**(1) DOI:
923   10.1038/s41467-019-12199-1.
924   81.   Gobeil, S.M.C., et al., *Leveraging Fungal and Human Calcineurin-Inhibitor Structures,*
925   *Biophysical Data, and Dynamics To Design Selective and Nonimmunosuppressive FK506*
926   *Analogs.* mBio, 2021. **12**(6) DOI: 10.1128/mBio.03000-21.
927   82.   Lee, Y., et al., *In Vitro and In Vivo Assessment of FK506 Analogs as Novel Antifungal Drug*
928   *Candidates.* Antimicrobial Agents and Chemotherapy, 2018. **62**(11) DOI: 10.1128/aac.01627-
929   18.
930   83.   Hearn, V.M. and D.W. Mackenzie, *Mycelial antigens from two strains of Aspergillus fumigatus:*
931   *an analysis by two-dimensional immunoelectrophoresis.* Mykosen, 1980. **23**(10): p. 549-62.
932   84.   Bertuzzi, M., et al., *On the lineage of Aspergillus fumigatus isolates in common laboratory use.*
933   Medical Mycology, 2021. **59**(1): p. 7-13 DOI: 10.1093/mmy/myaa075.
934   85.   Zhao, C., et al., *High-Throughput Gene Replacement in Aspergillus fumigatus.* Current
935   Protocols in Microbiology, 2019. **54**(1) DOI: 10.1002/cpmc.88.
936   86.   van Rhijn, N., et al., *Development of a marker-free mutagenesis system using CRISPR-Cas9 in*
937   *the pathogenic mould Aspergillus fumigatus.* Fungal Genetics and Biology, 2020. **145** DOI:
938   10.1016/j.fgb.2020.103479.
939   87.   Peng, D. and R. Tarleton, *EuPaGDT: a web tool tailored to design CRISPR guide RNAs for*
940   *eukaryotic pathogens.* Microbial Genomics, 2015. **1**(4) DOI: 10.1099/mgen.0.000033.
941   88.   Ralph, B. and D.C. Sheppard, *A Murine Model for Chronic A. fumigatus Airway Infections*, in
942   *Host-Fungal Interactions*. 2021. p. 215-224.
943   89.   Amos, B., et al., *VEuPathDB: the eukaryotic pathogen, vector and host bioinformatics resource*
944   *center.* Nucleic Acids Research, 2022. **50**(D1): p. D898-D911 DOI: 10.1093/nar/gkab929.
945   90.   Cherinka, B., et al., *Marvin: A Tool Kit for Streamlined Access and Visualization of the SDSS-IV*
946   *MaNGA Data Set.* The Astronomical Journal, 2019. **158**(2) DOI: 10.3847/1538-3881/ab2634.
947   91.   Haberthür, U. and A. Caflisch, *FACTS: Fast analytical continuum treatment of solvation.* Journal
948   of Computational Chemistry, 2007. **29**(5): p. 701-715 DOI: 10.1002/jcc.20832.
949
950

SUPPLEMENTARY FILES

Table S1: Genes included in the NanoString custom codeset.

Table S2: *In vitro* and *in vivo* conditions assayed for the NanoString run.

Table S3: Primers employed in this study.

Supplementary File 1: NanoString designed Probes A and B for hybridisation and MTE primers for enrichment.

Supplementary File 2: Raw data from digital analysis of hybridised codesets.

Supplementary File 3: Normalized data used for the analyses.