The molecules lost

NEW COMPOUNDS BINDING AT THE HISTAMINE SITE

OF THE NMDA RECEPTOR (NMDA$_{(HA)}$R)

V. ARMAND

Université de Paris, SPPIN - Saints-Pères Paris Institute for the Neurosciences, CNRS, Paris F-75006, France 45, rue des Saints Pères 75006 Paris, France

Corresponding author: *Vincent ARMAND, Université de Paris, SPPIN - Saints-Pères Paris Institute for the Neurosciences, CNRS, Paris F-75006, France* Descartes 45, rue des Saints Pères 75006 Paris, France

Tel: (+33) 1 42 86 42 21 FAX: (+33) 1 42 86 3830

e-mail: vincent.armand@parisdescartes.fr

Abstract

NMDA receptor ligands have been the target of intensive research for the treatment of psychotic diseases and central nervous system diseases. Our group published the characterization of the NMDA receptor histamine site. We developed modulators for this site, about 500 drugs were tested and we finally have a partial agonist FUBn293 with a nanomolar affinity,and also an antagonist ST-579 with a nanomolar affinity. We suggest that agonists at the (NMDA$_{(HA)}$R) constitute an innovative class of antipsychotics for the treatment of schizophrenia and other neurological or psychiatric disorders.

In this article, I will relate more than 20 years of research on a series of molecules that would have been forgotten without this short article summarizing the activity structure of this one. This work was done by many researchers, and I will list them here in the hope that I will not forget one, here they are:

V. Armand, J.-M. Arrang, C. Bayard, A. Burban, R. Faucard, C. R. Ganellin, S. Graßmann, P. P. Griffin, H. Kubas, I. Nuss, U. Reichert, W. Schunack, J.-C. Schwartz and H. Stark.

During this period, several science theses were written:

Для the pharmacological results

For the pharmacological results

in 1999 Cécile Bayard in Paris

Caractérisation du site de liaison de l'histamine du recepteur NMDA

in 2004 Raphaêl Faucard in Paris

Caractérisation pharmacologique et fonctionnelle du site histamine, modulateur du récepteur NMDA

in 2009 Aude Burban in Paris

Modulation du récepteur NMDA du glutamate par l'histamine : intérêt dans la schizophrénie

For the compounds formula and method for preparing

in 2000 Sven. Graßmann in Berlin

Synthese und Pharmakologie von Liganden der Histaminbindungsstelle des N-Methyl-D-aspartat-Rezeptors

in 2000 Ulrich Reichert in Berlin

Heterocyclische Alkanamine als Modulatoren der Histamin-Bindungsstelle des NMDA-Rezeptors : Synthese, Analytik und Struktur-Wirkungsbeziehungen

i

 in 2004 Isabelle Nuss in Berlin,

Heterobicyclische Alkanamine als Liganden einer modulatorischen Bindungsstelle des N-Methyl-D-aspartat-Rezeptors : Synthese, Analytik und Struktur-Wirkungsbeziehungen

in 2005 Perry Paige. Griffin in Frankfurt am Main

Neue Liganden einer modulatorischen Bindungsstelle an NMDA-Rezeptoren : Synthese, Analytik und Struktur-Wirkungsbeziehungen

in 2007 Holger Kubas in Frankfurt am Main

Substituierte Benzothiazole als allosterische Modulatoren an NMDA-Rezeptoren

There have also been communications on the subject;

 U. Reichert, S. Graßmann, C. Bayard, J.-M. Arrang, J.-C. Schwartz, H. Stark und W.Schunack.

Heterocyclische Alkanamine als positive Modulatoren des NMDA-Rezeptors.Deutsche Pharmazeutische Gesellschaft: Landesgruppe Berlin-Brandenburg: Der wissenschaftliche Nachwuchs stellt sich vor. Berlin (5. Juli 1999), Abstractbuch

S. Graßmann, U. Reichert, C. Bayard, J.-M. Arrang, J.-C. Schwartz, H. Stark

und W.Schunack.

Neue Leitstrukturen für Agonisten einer modulatorischen Bindungsstelle des NMDA-Rezeptors.

Deutsche Pharmazeutische Gesellschaft: Landesgruppe Berlin-Brandenburg: Der wissenschaftliche Nachwuchs stellt sich vor. Berlin (5. Juli 1999), Abstractbuch

I Nuss, U. Reichert, S. Graßmann, R. Faucard, C. Bayard, J.-M. Arrang, J.-C. Schwartz, H.Stark, and W. Schunack.

Synthesis and Pharmacology of Novel Potent Ligands at the Histamine Binding Site of the NMDA Receptor. 3rd European Graduate Student Meeting of the German Pharmaceutical Society (DPhG),Frankfurt am Main (23.-25. Februar 2001). Arch. Pharm. Pharm. Med. Chem. 2001, 334 (Suppl. 1), 23 (A-65).

I. Nuss, S. Graßmann, R. Faucard, J.-M. Arrang, J.-C. Schwartz, H. Stark und W. Schunack.

Synthese und Pharmakologie von neuen potenten Liganden der Histamin-Bindungsstelle des NMDA-Rezeptors. Deutsche Pharmazeutische Gesellschaft: Landesgruppe Berlin-Brandenburg: Der wissenschaftliche Nachwuchs stellt sich vor. Berlin (12. Juli 2001), Abstractbuch.

S. Graßmann, C. Bayard, J.-M. Arrang, C. R. Ganellin, J.-C. Schwartz, H. Stark, and W.Schunack.

In Vitro Investigation on Novel Potent Antagonists of a Modulatory Binding Site of NMDA Receptors. Jahrestagung der Deutschen Pharmazeutischen Gesellschaft

(DPhG) „Die wissenschaftliche Pharmazie im neuen Jahrtausend – Trends, Entwicklungen, Highlights", Halle (Saale) (10.- 13. Oktober 2001). Arch. Pharm. Pharm. Med. Chem. 2001, 334 (Suppl. 2), 18 (V1-17).

P. P. Griffin, S. Graßmann, C. Bayard, J.-M. Arrang, J.-C. Schwartz, W. Schunack, and H.Stark.

Histamine Derivatives as Potential NMDA Receptor Ligands. Deutsche Pharmazeutische Gesellschaft (DPhG) – Jahrestagung 2002, Berlin (9.-12.

Oktober 2002), Poster.Arch. Pharm. Pharm. Med. Chem. 2002, 335 (Suppl. 1), 81 (P:C36)

I. Nuss, S. Graßmann, R. Faucard, J.-C. Schwartz, J.-M. Arrang, H. Stark, and W. Schunack.

Bicyclic Alkanamines as Highly Potent Ligands of a Modulatory Binding Site of NMDA

Receptors. Deutsche Pharmazeutische Gesellschaft (DPhG) – Jahrestagung 2002, Berlin (9.-12.Oktober 2002), Koautor. Arch. Pharm. Pharm. Med. Chem. 2002, 335 (Suppl. 1), 48 (V:C16).

P. P. Griffin, S. Graßmann, C. Bayard, J.-M. Arrang, J.-C. Schwartz, W. Schunack, and H.Stark

Cytoprotective Histamine Derivatives as NMDA Receptor Modulators. ZAFES Kick-Off Symposium "Lipid Signaling", Frankfurt am Main (14. Oktober 2004)

The project was stopped more than 10 years ago because the patent on these molecules entitled; **NMDA RECEPTOR HISTAMINE SITE MODULATORS FOR USE IN THE TREATMENT OF CENTRAL NERVOUS SYSTEM DISEASES** was not accepted, some of the data had been released to the public, confer to the symposium.

So these are the molecules I think it's a shame to forget them for good.

Introduction

The effects of histamine are mediated by four G protein-coupled receptors (H_1, H_2, H_3 and H_4). In the brain, histamine also binds to the histamine site ($NMDA_{(HA)}R$) of the N-methyl-D-aspartate receptor (NMDAR) (1). Histamine potentiates NMDA currents in isolated (2), and cultured (3) hippocampal neurons and this effect requires NMDARs containing NR1 variants lacking exon 5 with NR2B subunits (4, 5). This potentiation is inversely related to the concentration of glycine (2, 6) and is reproduced by tele-methylhistamine (tele-MeHA), the catabolite of histamine in the brain (3, 4-6). Histamine also binds to $NMDA_{(HA)}R$ to potentiate NMDA-induced [^{3}H]noradrenaline release from hippocampal synaptosomes(6). Histamine potentiates N-methyl-D-aspartate receptors by interacting with an allosteric site distinct from the polyamine binding site (6)

Materials and Methods

[^{3}H] noradrenaline release from hippocampal synaptosomes.

A crude synaptosomal fraction was prepared as described previously with minor modifications (7).Adult male Wistar rats (200-250g) were killed by decapitation. The hippocampus wa rapidly removed and homogenized (Potter Elvehjem glass; eight up-down strokes) in 40 volumes of 0,32 M sucrose. The homogenate was first cecntrifuged (100g for 10 min) to remove nuclei and cellular debris. Synaptosomes

were isolated from the supernatrant by a second centrifugation (12,000 g for 20 min). The synaptosomal pellet was then suspended in modified Krebs-Ringer bicarbonate medium of the following composition : NaCl 120 mM ; KCl 0.8 mM; KH2PO4,1.2 mM ; CaCl2 1.3 mM ; MgSO4 1,2 mM; NaHCO3 27.5 mM ; glucose, 10 mM; ascorbic acid 0.06 mM ; EDTA 0.03 mM; gassed with 95% O2 and 5% CO2); pH 7.4.

Synaptosomes are then suspended in this same medium in the require volume of assay buffer abd are then incubated for 1 hour at 37°C in a rotary water bath, in an atmosphere of 95% 02 and 5% CO2, with [3H]noradrenaline (final concentration 30 nM, GE Healthcare, Buckingamshire, UK). During this incubation, synaptosomes are loaded with the labelled neurotransmitter. Then, the labelling of synaptosomes is followed by 4 washes with a Mg2+-free medium prewarmed at 37°C.

Synaptosomes are distributed in identical aliquots (200 µg of protein) in a final volume of 500 µL and incubated with NMDA (200µM), glycine (1 µM) and drugs to test in the presence of thioperamide, an H3 receptor antagonist in a saturating concentration (1 µM) to prevent the action of the heteroreceptor H3 modulating [3H] noradrenaline release in the system.(13Schlicker 1989) After 3 minutes of incubation at 37°C, reaction is stopped by immersion of tubes in ice-cold water, immediately followed by a centrifugation (14 000 × g, 10 sec). The amount of radioactivity released into each supernatant is finally determined by liquid scintillation using a β counter.

Results

We have investigated new drugs, ligands of the histamine site of the NMDA receptor by using the model of the NMDA-mediated [3H]noradrenaline release from hippocampal synaptosomes.

Over the years, we have tested more than 500 molecules and identified agonists and antagonists of the histamine binding site on the NMDA receptor, we present the most remarkable molecules with structure activity relationships in two series of tables in supplementary datas with the agonist (SD1) and the antagonist (SD 2)

On this response we identify some 2-aminobenzothiazole derivatives as potent

agonists of the NMDA$_{(HA)}$R .The FUB$_n$7, is the lead compound, it was a full agonist with a micromolar potency. Optimisation of this lead was obtained with substitution of the aliphatic amino group, that led to agonists with a nanomolar agonist potency such as FUB$_n$293 (8).

The FUB$_n$7,as lead coupound was first tested in various brain regions and the hippocampus give the best answer on [^{3}H]noradrenaline release (Fig 1B), then we used several NMDAR antagonist on the FUB$_n$7 and we observed a non competive inhibition (Fig 1C,D,E).

FUB$_n$7, the first lead compound obtained in this series, behaved as a full agonist with a micromolar potency (EC$_{50}$ = 2.1 $\pm$ 0.1 µM). Its effect was antagonized by the NMDAR blockers MK-801 (Figure 1C) and by ifenprodil, the NR2B antagonist (Figure 1E).

As we previously reported (8), FUB$_n$293 also potentiated NMDA-induced [^{3}H]noradrenaline release from hippocampal synaptosomes (EC$_{50}$ = 2,8 $\pm$ 1.8 nM). FUB$_n$293 displayed a nanomolar agonist potency on NMDA-induced [^{3}H]noradrenaline release from hippocampal synaptosomes. It was 50 fold more potent than FUB$_n$7 and 25,000 fold more potent than histamine, but its maximal effect was 49 $\pm$ 6% that of FUB$_n$7, suggesting that it behaved as a partial agonist on this response (Fig 2B).

In agreement, FUB$_n$293 decreased in a concentration-dependent manner the sub-maximal effect of FUB$_n$7. The maximal antagonism reached at the highest concentrations tested led to the same plateau as its maximal agonist effect (-50.2 $\pm$ 3.1% vs +50.9 $\pm$ 4.1% of FUB$_n$7-induced release), and its K_i assuming a competitive antagonism of FUB$_n$7 was in the same nanomolar range as its agonist potency (3.7 $\pm$ 1.4 nM) (Figure. 2B). A similar pattern and K_i value (7.6 $\pm$ 1.9 nM) was obtained when FUB$_n$293 was opposed to tele-MeHA (Figure. 2C), which confirmed that FUB$_n$7 and FUB$_n$293 bind at the histamine site i.e. the NMDA$_{(HA)}$R.

We obtain also antagonist the best was the ST-579

ST-579 was able to inhibite the potentiation of [³H]noradrenaline release induced by FUB$_n$7 (Fig 3B) and FUB$_n$293 (Fig 3C)

Activity Structure

The activity structure of a part of the compounds is presented in tables.

THE AGONIST

The tables give the intrinsic activity of the drugs compared to the FUB 7 and the EC 50 of the drugs in μM.

The first series describes agonists, in which the molecules are derived from the general formula around the lead compound FUBn 7 (Table 1 to Table 3)

The second series describes agonists, in which the molecules are derivated from the general formula around the lead compound FUB 16 (derivated from R-1,3-benzothiazol-2-amine) and lead to a first best compound FUB 247 and then to the FUBn293 the most potent drug realized (Table 4 to Table 9)

THE ANTAGONIST

The tables give the IC 50 of the drugs in μM.

The first series describes antagonists, in which the molecules are derived from the general formula the first compound FUB 88 (derivated of R-(1H-imidazol-5-yl)ethan-1-amine) to the first interesting compound of this series FUB 114 (Table 10

to Table 12).

The second series describes the antagonist, in which the molecules are derived from the general formula around the lead compound FUB 16 and lead to ST-579 the most potent drug realized (Table 13 and Table 14).

CONCLUSION

In conclusion, these data confirm the existence of a histamine site, distinct from other allosteric sites, of the NMDAR. Since histamine also activates the human NMDAR (9), agonists of the $NMDA_{(HA)}R$ may be helpful in therapeutics. We suggest that agonists at the $NMDA_{(HA)}R$ constitute an innovative class of antipsychotics for the treatment of schizophrenia and other neurological or psychiatric disorders.

REFERENCES

1.   H. L. Haas , O.A. Sergeeva, O. Selbach, Histamine in the nervous system. *Physiol Rev* **88**, 1183-1241 (2008).

2.   V. S. Vorobjev, I. N. Sharonova, I. B. Walsh, H.L. Haas, Histamine potentiates N-methyl-D-aspartate responses in acutely isolated hippocampal neurons. *Neuron* **11**,

837-844 (1993).

3.   J. M.Bekkers,   Enhancement by histamine of NMDA-mediated synaptic transmission in the hippocampus. *Science* **261**, 104-106 (1993).

4.   K.Williams, Subunit-specific potentiation of recombinant N-methyl-D-aspartate receptors by histamine. *Mol Pharmacol* **46**, 531-541 (1994).

5.   K.Williams, Pharmacological properties of recombinant N-methyl-D-aspartate (NMDA) receptors containing the epsilon 4 (NR2D) subunit. *Neurosci Lett* **184**, 181-184 (1995).

6.   A. Burban, R. Faucard, V. Armand, C. Bayard, V. Vorobjev, J.-M. Arrang. Histamine potentiates N-Methyl-D-Aspartate receptors by interacting with an allosteric site distinct from the polyamine-binding site. *J Pharmacol Exp Ther* **332(3)**:912-921 (2010)

7.   The isolation of nerve endings from brain: An electron microscopic study of cell fragments derived by homogenization and centrifugation E. G. Gray, V. P. WhittakerJ Anat. Jan; 96: 79–88.(1962).

8.   I. Nuss, U. Reichert, S. Graßmann, R. Faucard, C. Bayard, J.-M. Arrang, J.-C. Schwartz, H. Stark, W. Schunack. Synthesis and pharmacology of novel potent ligands at the histamine binding site of the NMDA receptor. 3rd European Graduate Student Meeting of the German Pharmaceutical Society (DPhG), Frankfurt am Main (23.-25. February 2001). Arch. Pharm. Pharm. Med. Chem.2001, **334** (Suppl. 1), 23 (A-65).

9.   M. Munir, et *al.* Pharmacological and immunological characterization of N-methyl-D-aspartate receptors in human NT2-N neurons. *J Pharmacol Exp Ther* **276**, 819-828 (1996).9.

10.   S. Graßmann, C. Bayard, J.-M. Arrang, C. R. Ganellin, J.-C. Schwartz, H.

Stark, W. Schunack. In Vitro Investigation on Novel Potent Antagonists of a Modulatory Binding Site of NMDA Receptors. Jahrestagung der Deutschen Pharmazeutischen Gesellschaft (DPhG) „Die wissenschaftliche Pharmazie im neuen Jahrtausend – Trends, Entwicklungen, Highlights, Halle (Saale) (10.-13. October 2001). Arch. Pharm. Pharm. Med. Chem.2001, 334 (Suppl. 2), 18 (V1-17).

11. 11. R.Faucard, V. Armand, A. Héron, V. Cochois, J-C Schwartz, J-M. Arrang. N-methyl-D-aspartate receptor antagonists enhance histamine neuron activity in rodent brain. J Neurochem 98, 1487-1496 (2006).

12. Inhibition of noradrenaline release in the rat brain cortex via presynaptic H3 receptors. Schlicker E, Fink K, Hinterthaner M, Göthert M. Naunyn Schmiedebergs Arch Pharmacol. 340:633-8. (1989)

This study was supported by INSERM, the French Ministère de la Recherche and the Fondation pour la Recherche Médicale.

Fig 1   **Effects of FUBn7 NMDA-induced [3H]noradrenaline release.**
A, Chemical structure of FUBn7..
B, Effect of FUBn7 on NMDA-induced [3H]noradrenaline release from synaptosomes of various rat brain regions. **C, D, E**, Effect of MK-801 (1-100 µM) (**C**), Mg2+ (1.2 mM) (**D**) and ifenprodil (0.03-1 µM) (**E**) on NMDA-induced [3H]noradrenaline release from hippocampal synaptosomes. Results are expressed as dpm/µg protein over [3H]noradrenaline release induced by NMDA (200µM) and glycine (1µM).
Each point represents the mean ± SEM of values obtained in 3-8 separate experiments.

Fig 2

Effects of FUBn293 NMDA-induced [3H]noradrenaline release. A, Chemical structure of FUBn293. **B, C,** Effect of FUBn293 tested alone, or against FUBn7 (**B**), or tele-MeHA (**C**), on NMDA-induced [3H]noradrenaline release from hippocampal synaptosomes. Results are expressed as per cent of the effect of FUBn7 (**B**), or tele-MeHA (**C**) (means of 6-16 determinations from 3-8 separate experiments).

AGONIST

TABLE 1

Name			Intrinsec activity %	Agonist EC 50 µM
Fub 7	n=3	$R^1 = 0$	**full**	**2,1**
Fub 169	n=4	$R^1 = 0$	full	1,6
Fub 216	n=5	$R^1 = 0$	20	0,12
Fub 239	n=6	$R^1 = 0$	23	0,12
Fub 282	n=7	$R^1 = 0$	?	>100
Fub 283	n=8	$R^1 = 0$	?	>100
Fub 241	n=3	$R^1 = CH3$	53	1,5
Fub 302	n=4	$R^1 = CH3$	29	0,2
Fub 310	n=5	$R^1 = CH$	?	>100

TABLE 2

Name		Intrinsec activity %	Agonist EC 50 µM
Fub 242		75	3,1
Fub 206		40	0,068
Fub 235		38	26
Fub 210		33	0,024
Fub 132	n=2	full	4,4
Fub 144	n=3	35	11
Fub 218	n=4	81	1
Fub 295	n=5	41	0,052
Fub 296	n=6	41	1,1

TABLE 3

Name		Intrinsec activity %	Agonist EC 50 µM
Fub 170	n=3	full	2,2
Fub 217	n=4	52	0,19
Fub 238	n=5	50	4,5
Fub 240	n=2	82	6,3
Fub 301	n=3	61	0,9
Fub 309	n=4	40	9

TABLE 4

Name		Intrinsec activity %	Agonist EC 50 µM
Fub 16		30	0,39
Fub 223		34	5,2
Fub 247		**48**	**0,024**
Fub 251		?	>100
Fub 252		?	>100
Fub 253		?	>10
Fub 300		45	0,037

TABLE 5

Name		Intrinsec activity%	Agonist EC 50 µM
Fub 266		48	1
Fub 281	n=2	30	0,19
Fub 292	n=3	32	0,034
Fub 293	**N=4**	**39**	**0,0028**
Fub 299	n=1	36	0,5
Fub 306	n=5	33	0,014
Fub 307	n=6	33	0,047
Fub 297		32	0,09
Fub 298		19	0,014

TABLE 6

Name		Intrinsec activity %	Agonist EC 50 µM
	H₂N—[benzothiazole]—CH₂CH₂—R1		
Fub 275	NH—(CH2)n—CH3, n=1	15	0,32
Fub 276	n=2	15	0,98
Fub 286	n=3	31	0,49
Fub 287	n=4	30	0,33
Fub 288	n=5	35	0,020
Fub 303	n=6	33	0,010
Fub 304	n=7	33	0,010
Fub 305	n=8	30	0,062
Fub 289	NH—C(CH3)(CH3)(H3C)	23	3,20
Fub 290	NH—C(CH3)(H3C)—CH2—CH3	27	0,11
Fub 291	NH—C(CH3)(H3C)—CH2—CH2—CH3	32	0,44
Fub 294	NH—CH2—CH2—CH2—O—CH3	34	0,60

TABLE 7

		Intrinsec activity %	Agonist EC 50 µM
ST 544		26	0,018
ST 545		31	0,025
ST 562		20	0,005
ST 563		50	0,007
ST 575		40	0,3
ST 576		38	0,20
ST 577		25	0,07
ST 578		30	0,60
ST 580		?	>10

TABLE 8

Name		Intrinsec activity %	Agonist EC 50 µM
ST 581		50	12
ST 582		45	0,013
ST			
ST 590		30	0,030
ST 591		45	0,097
ST 592		30	0,005
ST			
ST 593		25	0,010
ST 594		30	0,007
ST 595		35	0,03

TABLE 9

ST 615		43	0,009
ST 617		64	1,2
ST 619		?	>10

ANTAGONIST
TABLE 10

name		Antagonist IC 50 µM
Fub 88		0,37
Fub 98	n=0	52
Fub 99	n=1	16
Fub 100	n=2	1,8
Fub166	n=3	0,65
Fub 113	n=4	1,09
Fub 114	**n=5**	**0,29**
Fub 122	n=6	2,1
Fub 164	n=7	0,17
Fub 183	n=8	0,61
Fub 184	n=9	0,91

TABLE 11

Name		Antagonist IC 50 µM
Fub 120		3,65
Fub 125	n=3	0,79
Fub 126	n=4	2,46
Fub137	n=5	1,37
Fub 212		0,13

TABLE 12

Name		Antagonist IC 50 µM
Fub 130		0,2
Fub 131		0,69
Fub 133		>100
Fub 134		0,21
Fub 135		0,52
Fub167		0,5
Fub 168		0,2

Fub 171	H₃C ... R1 structure	0,5
Fub 172	CH2n, H₃C—R1 structure n=7	0,1
Fub 177	n=3	128
Fub 214	n=8	0,36

TABLE 13

Name		Antagonist IC 50 µM
Fub 243	 n=1	26
Fub 249	n=3	2,1
Fub 255	n=5	1,7
Fub 248		29
Fub 250	R1= Cl	13
Fub 256		30
Fub 258		0,8
Fub 259		0,23
Fub 260		1,1

TABLE 14

Name		Antagonist IC 50 µM
Fub 262		1,2
Fub 263		0,28
Fub 264		0,1
Fub 265		1,1
Fub 278		5,9

		Antagonist IC 50 μM
Fub 270		3
Fub 277		1,39
Fub 308		>100
Name		Antagonist IC 50 μM
St 541		0,077
ST 542		0,44
ST 543		0,1
ST 579		**0,038**

ST 587		0,5
ST 588		5
ST 589		0,9
		1,2
ST 616		0,042
ST 618		0,064