Resolution of SARS-CoV-2 infection in human lung tissues is driven by

extravascular CD163+ monocytes

Devin Kenney[1,2], Aoife K. O'Connell[1,2,4], Anna E. Tseng[1,2,4], Jacquelyn Turcinovic[1,2,3], Meagan L.

Sheehan[7,12], Adam D. Nitido[7,12], Paige Montanaro[2,4], Hans P. Gertje[2,4], Maria Ericsson[6], John H.

Connor[1,2], Vladimir Vrbanac[7], Nicholas A. Crossland[1,2,4], Christelle Harly[8,9,12]*, Alejandro B.

Balazs[7,12]*, and Florian Douam[1,2,12,13] *

[1]Department of Virology, Immunology, and Microbiology, Boston University Chobanian &

Avedisian School of Medicine, Boston, MA, USA.

[2] National Emerging Infectious Diseases Laboratories, Boston University, Boston, MA, USA.

[3] Bioinformatics Program, Boston University, Boston, MA, USA.

[4] Department of Pathology and Laboratory Medicine, Boston University Chobanian & Avedisian

School of Medicine, Boston, MA, USA.

[6] Electron Microscopy Core Facility, Harvard Medical School, Boston, MA, USA.

[7] Ragon Institute of MGH, MIT and Harvard, Cambridge, MA, USA.

[8] Université de Nantes, INSERM, CNRS, CRCINA, Nantes, France

[9] LabEx IGO 'Immunotherapy, Graft, Oncology', Nantes, France.

[12] These authors contributed equally to the work.

[13] Lead contact

*Correspondence: fdouam@bu.edu; abalazs@mgh.harvard.edu; christelle.harly@inserm.fr.

ABSTRACT

The lung-resident immune mechanisms driving resolution of SARS-CoV-2 infection in humans remain elusive. Using mice co-engrafted with a genetically matched human immune system and fetal lung xenograft (fLX), we mapped the immunological events defining resolution of SARS-CoV-2 infection in human lung tissues. Viral infection is rapidly cleared from fLX following a peak of viral replication. Acute replication results in the emergence of cell subsets enriched in viral RNA, including extravascular inflammatory monocytes (iMO) and macrophage-like T-cells, which dissipate upon infection resolution. iMO display robust antiviral responses, are transcriptomically unique among myeloid lineages, and their emergence associates with the recruitment of circulating CD4+ monocytes. Consistently, mice depleted for human CD4+ cells but not CD3+ T-cells failed to robustly clear infectious viruses and displayed signatures of chronic infection. Our findings uncover the transient differentiation of extravascular iMO from CD4+ monocytes as a major hallmark of SARS-CoV-2 infection resolution and open avenues for unravelling viral and host adaptations defining persistently active SARS-CoV-2 infection.

INTRODUCTION

Coronavirus disease 2019 (COVID-19) is a respiratory disease that has swept the world since its emergence in the Wuhan province of China in late 2019. The etiologic agent of COVID19, the Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) is a plus-sense, enveloped RNA virus that targets the epithelium of the respiratory tract. Infection results in varying severities of COVID-19, with the majority of cases being mild to asymptomatic. The onset of severe disease is associated with aberrant immune responses (e.g. increased pulmonary infiltration of myeloid lineages, exacerbated inflammation, lymphopenia, and "cytokine storm") and severe lung injury (e.g. lung consolidation, diffuse alveolar damage (DAD), and thrombosis) (Boumaza et al., 2021a; Delorey et al., 2021b; Grant et al., 2021; Gubernatorova et al., 2020; Kosyreva et al., 2021; Schultze and Aschenbrenner, 2021; Zhang et al., 2020).

Extensive studies, using both human samples and animal models, have been instrumental in unraveling cellular and molecular processes driving severe COVID19 disease (Edahiro et al., 2023; Fan et al., 2022; Ren et al., 2021; Stephenson et al., 2021). However, the immunological mechanisms defining successful resolution of SARS-CoV-2 infection remain elusive. While a recent study identified HLA-B*15 expression and previous coronavirus exposure as a determinant of rapid SARS-CoV-2 clearance and asymptomatic infection, this involved only 17-25% of asymptomatic patients (Augusto et al., 2023). The ability of SARS-CoV-2 to downregulate HLA-B suggests that other important immune mechanisms enabling SARS-CoV-2 clearance may be at play in the human lung (Moriyama et al., 2023; Yoo et al., 2021).

Our limited understanding of the human mechanisms driving effective viral clearance largely stems from the lack of appropriate experimental platforms. Only peripheral blood and bronchiolar lavage samples (which prevent access to the lung interstitial compartment) can be accessed from human patients with asymptomatic or mild COVID-19. There are also logistical challenges associated with the capture of early pulmonary immune responses in infected individuals, and/or in the validation of purely correlative findings, as this requires the need to conduct experimental

human infection. Though large and small animal models of COVID-19 have been developed, the

high-cost and limited reagent availability associated with non-human primate models, and the

large divergence between rodent and human immune systems further underscore the need for

additional models capable of recapitulating human protective immune responses to SARS-CoV-

2 infection.

Mice co-engrafted with human fetal lung xenograft (fLX) and a human immune system (HIS)

represent a robust platform to uncover human immunological determinants of SARS-CoV-2

infection resolution and lung tissue protection from excessive inflammation (Di et al., 2022;

Kenney et al., 2022; Sun et al., 2022a). We recently demonstrated that while mice engrafted with

fLX are highly susceptible to SARS-CoV-2 infection and lung tissue damage, co-engraftment of

fLX and a HIS in a xenorecipient strain supporting enhanced myelopoiesis (i.e., HNFL mouse

model) resulted in complete lung tissue protection from both infection and histopathology (Kenney

et al., 2022). Infection was cleared as early as day 2 post-virus inoculation, preventing widespread

viral replication in tissues. Macrophage antiviral responses were the dominant correlates of rapid

viral clearance and limited tissue damage. However, how the human immune system mobilizes

and resolves infection upon extensive viral replication within human lung tissues, a context that

more likely describes mild cases of SARS-CoV-2 infection, has remained elusive.

In this study, we leverage immunodeficient mice engrafted with a human fetal lung xenograft

(fLX) and a genetically matched human immune system, fetal liver and thymus (BLT-L mice) to

identify human immune determinants and mechanisms, or lack thereof, that drive SARS-CoV-2

resolution or persistence in human lung tissues. Consistent with previous reports (Di et al., 2022;

Sun et al., 2022a), BLT-L mice were permissive to SARS-CoV-2 infection following direct viral

inoculation into the fLX, with infection swiftly resolving following an early viral replication peak at

2-days post-inoculation (dpi). Single cell transcriptomic approaches identified extravascular

CD163+ inflammatory monocyte (iMO), which were highly enriched in viral RNA, as the primary

mediator of antiviral responses preceding viral clearance. Proteomic and cell depletion assays

supported a model in which recruitment of circulating CD4+ monocytes into infected fLX, and their

subsequent differentiation as iMO, is a defining feature of infection resolution. Abrogation of

infection clearance via systemic depletion of CD4+ cells resulted in persistent SARS-CoV-2

replication and a remodeling of lung antiviral responses consistent with chronic viral infection.

Collectively, our work sheds light on key immunological mechanisms driving SARS-CoV-2

resolution in human lung tissues, and provides experimental evidence that monocyte

differentiation into extravascular iMO is a critical hallmark of this process. Our study also

underscores the potential of the BLT-L mouse model to inform the development of

immunotherapies against viral respiratory infections and examine the cause and consequences

of chronic SARS-CoV-2 infection on both the virus and the host.

RESULTS

BLT-L mice effectively clear SARS-CoV-2 infection following acute viral replication

Previous work from our laboratory (Kenney et al., 2022) and others (Di et al., 2022; Sun et al.,

2022a; Wahl et al., 2019) have shown that immunodeficient mice can successfully be engrafted

with human fetal lung xenograft (fLX) independently, or in combination with a human immune

system. In this study, we leveraged a previously reported model co-engrafted with fetal liver and

thymus as well as with human hematopoietic stem cells (HSC) and fLX (BLT-L mice) (Wahl et al.,

2019). Fetal liver and thymus were engrafted under the renal capsule of adult mice (12-16 weeks

old) prior to intravenous HSC injection, while a piece of fLX was surgically introduced

subcutaneously on one flank of each animal, as described previously (Wahl et al., 2019). To

determine the susceptibility of BLT-L mice to SARS-CoV-2 infection and their ability to effectively

clear infection, BLT-L mice were inoculated with SARS-CoV-2 (2019-nCoV/USA_WA1/2020) via

intra-graft injection. We used a viral dose (10^6 PFU) that we previously established to drive robust

fLX infection in an immunodeficient mouse strain not engrafted with a HIS (Kenney et al., 2022).

Throughout the course of infection, mice did not display any weight loss (**Fig. 1 A)** or clinical

signs of disease such as lethargy or lack of responsiveness (data not shown). fLX were collected

at 2, 6, and 12 dpi to assess lung histopathology and viral titers. Immunohistochemistry (IHC) for

SARS-CoV-2 S protein revealed evidence of epithelial cell infection in fLX at 2 dpi, while the

majority of viral antigen was cleared by 6 dpi and became undetectable by 12 dpi (**Fig.1 B**[a-d]).

SARS-CoV-2 S antigen was primarily detected in the cytoplasm of alveolar epithelium

morphologically consistent with type II pneumocytes (AT2 cells) and macrophages within alveolar

spaces admixed with necrotic cellular debris and bronchiole epithelium, consistent with the

primary cell targets of SARS-CoV-2 (**Fig.1 B**[b]). Transmission electron microscopy (TEM) imaging

of fLX at 2 dpi substantiated evidence of productive fLX infection, as evidenced by the presence

of viral particles in the cytosol of epithelial cells and budding/invagination of virions (**Fig. S1 A-D**).

Viral titer quantification in fLX also supported evidence of acute infection at 2 dpi prior to effective

resolution (**Fig.1 C**). A significant amount of infectious viral particles could be recovered from fLX

at 2 dpi (3.20 ± 2.52 log([PFU/mg of tissue]), followed by a significant and incremental reduction

at 6 dpi (0.281 ± 0.688 log[PFU/mg of tissue]) and 12 dpi (0.111 ± 0.293 log[PFU/mg of tissue])

(**Fig. 1 C**). No serum neutralizing activity against SARS-CoV-2 was detected at any time point

(**Fig. S1 E,F**). The absence of neutralizing humoral responses contributing to infection resolution

is consistent with the rapid clearance of infectious viral particles in fLX.

Next, we examined histopathological phenotypes associated with active and resolved fLX

infection. Interpretation of hematoxylin and eosin (H&E) staining illustrated denuding of

pneumocytes, neutrophil infiltration, edema, hemorrhage, thrombosis, and pneumocyte necrosis

at 2 dpi (**Fig. 1 B**[e-h]**; S1 G-I**). No major disease phenotypes were observed at 6 and 12 dpi

compared to naïve fLX, underscoring that fLX can mount repair mechanisms upon resolution of

infection. A previously described semi-quantitative histopathological scoring system (Kenney et

al., 2022) provided statistical confirmation for significant lung pathology at 2 dpi, which was no

longer apparent at day 6 dpi (**Fig. 1 D**). Collectively, BLT-L fLX recapitulates important hallmarks

of acute SARS-CoV-2 infection, including viral replication and histopathological manifestations of

disease, prior to effective viral clearance and lung tissue repair.

SARS-CoV-2 induces cellular remodeling of fLX upon infection

We performed bulk-RNA sequencing on fLX to identify gene signatures associated with viral

clearance. Strikingly, transcriptomic responses at both 2 and 12 dpi were limited and mostly

defined by transcript downregulation relative to naïve fLX mRNA expression (150 and 333 genes

at 2 dpi and 12 dpi, respectively) (**Fig. S1 J-N**) which is consistent with the ability of SARS-CoV-

2 to accelerate the degradation of cytosolic cellular mRNAs (Finkel et al., 2021). At both time

points a smaller subset of genes were upregulated (91 and 21 genes at 2 and 12 dpi,

respectively); these upregulated genes primarily associated with B cell response/activation

(FCRL1, FCRL5, FCRLA, CR2, BLK, PAX5 and MZB1) and immunoglobin recombination

(*IGHV3-11, IGHV3-51, IGHD, IGHM, IGHJ3, IGHJ4, IGLV1-36, and IGLC5*) (**Fig. S2 K; Data S1**);

albeit no neutralizing antibody responses could be detected in infected BLT-L mice at any time

point (**Fig. S1 K**). Low to moderate upregulation of genes associated with NF-kb

activation/signaling (*IL1A*, *IL1RL1, TLR9, FABP4*) and monocyte/macrophage activation (*FGR*,

CD101, SLC11A1, HBEGF, TREM1, VSIG4, CCL24) were also observed (**Data S1**).

Given that bulk transcriptomic data provided limited insight into significant gene signatures

associated with infection resolution, we hypothesized that antiviral and clearance mechanisms

are likely to be compartmentalized within specific immune effectors whose contribution cannot be

comprehensively captured using bulk approaches. As we previously reported that effective control

of SARS-CoV-2 infection in fLX is dependent on the engraftment of a functional human immune

system (Kenney et al., 2022), we performed single-cell RNA sequencing (scRNA-seq) on naïve

and infected fLX of BLT-L mice to identify critical immune effectors of viral clearance. While the

majority of cells detected by scRNA-seq were human, a minor population of mouse cells were

detected in fLX and excluded from downstream analysis (**Fig. S2 A-C**). Analysis of the human

compartment revealed diverse immune subsets (T cells/Innate lymphoid cells (ILC), B cells,

myeloid cells and mast cells) and non-immune cell types (ciliated and non-ciliated epithelial,

endothelial, mesenchymal cells and chondrocytes) in both naïve and infected fLX (**Fig. 2 A,B; S2**

D). Important to note, because our tissue digestion process favors hematopoietic cell recovery

compared to non-immune compartments, the relative representation of the different cell types in

fLX is likely biased, as asserted by the limited size of the epithelial compartment. Regardless, the

relative dynamics of each cell type captured could be examined upon infection.

T-cell/ILC frequency dramatically decreased in fLX upon infection (naïve, 66.8%; 2 dpi, 35.3%;

12 dpi, 21.8%), which is consistent with evidence that COVID19 can induce lymphoid depletion

(**Fig. 2 B,C**). Notably, lymphopenia was not observed in the peripheral blood of BLT-L mice

suggesting lymphocyte depletion in fLX was not directly attributed to declining circulating

peripheral lymphocytes (**Fig. S2 E**). In contrast, myeloid (naïve, 9.94%; 2 dpi, 25.3%; 12 dpi,

22.2%) and B-cell subsets (naïve, 3.64%; 2 dpi, 5.38%; 12 dpi, 5.31%) relatively expanded upon

infection (**Fig. 2 B,C**). The epithelial, endothelial and mesenchymal cell frequencies also

increased upon infection, while mast cell, ciliated cell and chondrocyte frequencies decreased

(**Fig. 2 B,C**). Interestingly, within clusters, some cell populations (mesenchymal, myeloid, and T

cells/ILCs) showed clear segregation by time point (**Fig. 2 B, S2 F**; 2 dpi: red clusters and 12 dpi:

blue clusters), suggesting that infection alters the transcriptional state of some cell types or drives

development of novel cell subsets. This temporal segregation suggests a two-step tissue

remodeling process upon infection likely involving i) early responses to acute viral replication

followed by ii) tissue repair mechanisms and potential immune imprinting.

Next, we explored the distribution of viral RNA throughout clusters to determine if temporal-

specific clusters associated with the presence of viral RNA. Similar to clearance of infectious viral

particles (**Fig. 1C**), viral RNA was not detected in fLX at 12 dpi but was detected in several cell

clusters at 2 dpi (**Fig. 2D**). While minor yet detectable expression of viral reads were observed in

epithelial, endothelial, chondrocyte and B cell lineages (**Fig. 2 D**), mesenchymal subsets exhibited

higher levels of viral reads. However, reads were limited and distributed sporadically without

significant enrichment within a given cluster (**Fig. 4 E and Fig S3 A**), (including the 2 dpi-specific

mesenchymal cluster defined in group B; **Fig. 2 B**), underscoring a lack of association between

viral RNA and mesenchymal activation upon SARS-CoV-2 challenge. In contrast, 2 dpi-specific

T-cell and myeloid cell groupings (A and C; **Fig. 2B,E**) were strongly enriched in viral RNA,

suggesting a contribution of these subpopulations to the resolution of acute infection.

Even though minimal viral RNA was detected in the epithelial compartment, we wished to

examine if there were transcriptional indicators of current or prior infection of the epithelium. Initial

sub-clustering of the epithelial compartment revealed populations of airway basal and secretory

cells, alveolar type 1 (AT1) and type 2 (AT2) cells, and serous cells (**Fig. 2F**). Consistent with

hallmarks of COVID-19 in humans, we noted a relative reduction of the AT2 compartment at 2 dpi

(25.6%) compared to naïve mice (73.1%), which was partially restored at 12 dpi (49.1%) (**Fig. 2**

F,G). AT2 cells were not enriched in viral RNA and did not exhibit any robust antiviral responses,

consistent with evidence that SARS-CoV-2 infection inhibits type I IFN signaling (Chen et al.,

2021). However, they displayed previously reported COVID-19 signatures in human patients

including downregulation of MHC genes (*HLA-C, HLA-B, and HLA-DPB1)* and *CCL5* (a cytokine

associated with immune recruitment during respiratory infection). Elevated expression of

mitochondrial transcripts and transcripts associated with ROS-mediated mitochondrial

dysfunctions (i.e., LARS2) were observed upon infection and particularly at 12 dpi, suggestive of

major cellular and metabolic remodeling in AT2 cells. (**Fig. 2H**). Consistently, increased levels of

SFTPA1 transcripts, a gene associated with AT2 to AT1 transition upon lung damage, was noted

at 2 dpi (**Fig. 2H**). An increase in AT1 and airway basal cell frequencies at 2 dpi (9.89% and

60.4%, respectively) compared to naïve fLX (5.00% and 19.4%, respectively) suggest the

activation of tissue repair processes of the lung epithelium following infection (**Fig. 2G**).

Collectively, these data demonstrate that scRNA-seq has the resolution to capture both

immune and non-immune cellular remodeling of fLX during acute SARS-CoV-2 infection and can

provide insight into the dynamics of immune-mediated viral clearance. Additionally, SARS-CoV-

2 infection within fLX of BLT-L mice mimic many hallmarks of human lung infection.

Macrophage-like T-cells are enriched in viral RNA and mount robust antiviral signatures

Next, we aimed to identify the specific T-cell subpopulation that was enriched in viral RNA through

sub-clustering of the main T cell/ILC clusters. A CD3+ subpopulation clustered very distinctively

from other CD3+ T-cell lineages and was highly enriched in viral RNA (Cluster 6; **Fig. 3 A-C**).

Cells included in this subpopulation displayed low levels of both *CD4* and *CD8* transcripts (**Fig. 3**

E), as well as elevated levels of mitochondrial transcripts **(Fig. S3 B,C**). Most notably, this T-cell

subset was uniquely defined by the expression of several transcripts known to be enriched in

macrophages (*MARCO*, *TIMP1*, *LYZ*), fibroblasts (*MGP*, *CALD1*, *COL1A*), or both (*A2M*, *IFITM3*)

but not in T-cells (**Fig. 3 E and S3 D**). Cells included in this subset also exhibited evidence of

cytotoxic function through the expression of *GZMA*, *GZMB* and *GNLY* transcripts, albeit

expression was also detected in CD8+ T-cells and activated tissue resident memory T cells (TRM)

(Clusters 1,3,4, **Fig. 3 E and S3 D**). Given the proximity of this cluster with myeloid lineages (**Fig.**

2A and E) and evidence of expression of macrophage-enriched transcripts, we referred to this

unique cluster as Macrophage-like T-cells (MTC). MTC also exhibited upregulation of key gene

pathways related to COVID-19, SARS-CoV-2 cell-intrinsic immune responses, cellular cytotoxicity

(e.g., degranulation) and cell death, consistently with a role for this subset to serve as a robust

primary responder to viral infection. (**Fig. 3 D**). Collectively, our findings suggest a potential role

for this previously uncharacterized subset in contributing to early control of SARS-CoV-2 infection

in lung tissues.

Extravascular CD163+ inflammatory monocytes are the dominant antiviral mediator among

myeloid lineages

Prior work has shown that the regulation of myeloid cells and their responses play a critical role

in the progression of SARS-CoV-2 related disease (Delorey et al., 2021a; Junqueira et al., 2022;

Rendeiro et al., 2021; Wauters et al., 2021); therefore, we investigated how myeloid lineages

respond to SARS-CoV-2 infection in fLX. Sub-clustering of myeloid lineages unveiled diverse

subpopulations, including alveolar and interstitial macrophages, various monocytes subsets, and

one conventional dendritic cell subset (cDC) (**Fig.4 A; Fig. S5**). Notably, we found the 2 dpi-

specific, viral RNA-enriched myeloid subset, earlier referred to as Cluster C, (see **Fig.2 B,E**) to

bear a signature of inflammatory monocytes (iMO) (**Fig.4 A-C**). iMO exhibited elevated

expression of *VCAN, S100A8, CD14, CD163* and absence/minimal expression of *CD4, MARCO*

and CD206 (MRC1) (**Fig. 4 D,E,F; Fig S4**). This gene expression profile was consistent with

those of human lung monocytes previously reported in human-immune system engrafted mice

(Evren et al., 2021) and human FCN1-monocytes recovered from the BAL of COVID-19 patients

with acute respiratory disease syndrome (Wendisch et al., 2021), collectively underscoring the

non-classical, extravascular monocyte nature of iMO and their human relevance in COVID-19.

Upon acute infection, only iMO and alveolar macrophages increased in frequencies, whereas

cDC and interstitial macrophage frequencies decreased (**Fig. 4 G**). iMO were the most abundant

lineage at 2 dpi (33.5%; **Fig. 4 G**) and were the leading mediators of antiviral responses, as

exemplified by the robust upregulated expression of interferon stimulated genes (ISGs) and

inflammatory cytokines (*CCL8, CXCL10, CCL2, ISG15, DEFB1, IL1RN, IFIT1, ISG20, IFIT3,* and

MX1), as well as inflammatory markers such as *CD163* (**Fig. 4 D,H**). While some ISGs (e.g.,

ISG15, IFIT3, and MX1) were expressed in other myeloid lineages at 2 dpi, their expression was

markedly lower compared to iMO, suggesting a strong association between viral RNA and

potentiated antiviral responses. Notably, upregulation of *CCL8*, *CXCL11* and *DEFB1* transcripts

were the most exclusive to iMO (**Fig. 4 H**). Decreased *CD4* expression was a defining feature of

iMO among monocyte sublineages (**Fig. 4 E**), suggesting that iMO differentiation is associate with

loss of CD4 expression concomitantly with extravasation.

In addition to iMO, a large population of CD4+ CD163- CD206(MRC1)- patrolling monocyte

was detected at all time points (Cluster 0; **Fig. 4 A,D,E**). This subset was however dominant at

12 dpi over other monocyte subsets (50.3%; **Fig. 4 G**), underscoring tissue repair and anti-

inflammatory mechanisms following viral clearance. Notably, a recent study provided evidence

for circulating patrolling monocyte to give rise to a non-classical, extravascular monocyte

population in the mouse lung that expand upon stimuli (Schyns et al., 2019), suggesting that

recruitment of CD4+ patrolling monocyte in BLT-L mice upon infection may be a pre-requisite for

extensive iMO differentiation and extravasation.

Multiplex immunohistochemistry (mIHC) analysis was consistent with our scRNAseq data.

While naïve fLX displayed high numbers of CD4+ cells and detectable CD163+ interstitial cells

(likely macrophages), acute infection at 2 dpi showed an increase in CD163+ cells both in the

extravascular interstitium and alveolar spaces and a dramatic reduction in CD4+ cells (**Fig. 4 I**).

These dynamics mirrored the 2 dpi-specific expansion of CD163+ iMO and alveolar macrophages

uncovered through scRNAseq. CD3+ T-cell lymphoid depletion was also apparent at 2 dpi (**Fig.**

4 I). At 12 dpi, the number of CD4+ cells had resolved reflecting the expansion of CD4+ patrolling

monocytes while the amount of CD163+ was reduced compared to 2 dpi (**Fig. 4 I**).

We also examined whether viral RNA-associated iMOs (CoV-iMOs) displayed specific

transcriptomic signature compared to iMOs (noCoV-iMOs) with undetectable viral RNA. Notably,

only two genes correlated with the presence of viral RNA in iMOs: *FGL2 and C15orf48* (**Fig. 4 J**).

While *FGL2* was down-regulated in CoV-iMO compared to noCoV-iMOs, *C15orf48* was

upregulated. Soluble FGL2 is known to exert immunosuppressive functions (notably through

inhibition of the NF-kB pathway) (Fu et al., 2023) whereas C15orf48 is a positive regulator of

inflammation previously associated with severe COVID-19 (Clayton et al., 2021). Collectively,

these findings suggest that productive or abortive SARS-CoV-2 infection could further potentiate

iMO inflammation through a specific set of genes.

Systemic depletion of CD4+ cells abrogates viral clearance in fLX.

To test the hypothesis that CD4+ monocyte recruitment is critical for SARS-CoV-2 clearance in fLX, we performed systemic depletion of CD4+ cells through intraperitoneal injection of an anti-CD4 antibody, both, prior to and after infection (**Fig. 5 A**). In parallel, we conducted depletion of CD3+ or CD8+ cells in separate animal cohorts and in a similar manner to deconvolute the role of T-cells and CD4+ non-T-cells in driving SARS-CoV-2 clearance. Effective systemic depletion was confirmed in the blood by flow cytometry (**Fig. S5 A**), and all animals were euthanized at 12 dpi to probe for persistent SARS-CoV-2 infection in fLX. fLX isolated from CD4-depleted animals exhibited robust persistence of productive SARS-CoV-2 infection at 12 dpi (mean Log PFU/mg tissue=2.37), and to a more significant extent than CD3- and CD8-depleted animals (mean Log PFU/mg tissue=0.96 and 0.83, respectively) when compared to IgG isotype treated mice (mean Log PFU/mg tissue=0) (**Fig. 5 B**). Notably, persistent infection in CD4-depleted animals was associated with significant downregulation of MHC class I (**Fig. 5 C**), a phenomenon we similarly observed in acutely infected fLX (2 dpi) (**Fig. 5 C**) and within our scRNAseq dataset (**Fig. 2 H**) and that has been previously reported in cells actively replicating SARS-CoV-2 (Arshad et al., 2023; Yoo et al., 2021), further emphasizing that depletion of CD4+ cells is associated with persistent viral replication.

Using mIHC, we validated the absence of CD4+ cells in fLX at 12 dpi. CD163+ cells were also consistently reduced in the fLX of CD4-depleted animals compared to the fLX of animals treated with IgG isotype control antibodies (**Fig. 5 D**[a,d,e]). In CD4-depleted mice, the remaining CD163+ cells co-localized with PPARγ+ cells, indicating that lung alveolar macrophages are not effectively depleted by systemically-delivered antibodies (**Fig. 5 D**[d-e]). Furthermore, some fLX from CD4-depleted mice appeared more enriched in CD3+ CD4- cells compared to isotype-treated mice (**Fig. 5 D**[e]), suggesting that defective monocyte recruitment may lead to enhanced CD8+ T-cell recruitment. CD3- and CD8-depleted mice that successfully cleared infection exhibited a higher presence of CD163+ cells in fLX at 12 dpi compared to isotype-treated mice, indicating that the

absence of CD8+ T-cells may lead to increased monocyte recruitment for effective viral clearance

(**Fig. 5 D**[b-c,f-g]).

Collectively, our findings suggest that the recruitment of CD4+ monocyte into infected fLX is

a crucial mechanism associated with effective viral clearance.

Depletion of CD4+ cells in BLT-L mice recapitulates features of chronic viral infection

We then investigated whether persistent SARS-CoV-2 infection at 12 dpi in CD4-depleted mice

displayed similarities with acute infection (i.e., at 2 dpi), or bore hallmarks of chronic viral infection.

Both acutely infected mice (2 dpi) and CD4-depleted mice (12 dpi) displayed similarly elevated

serum levels of CCL2 and CCL3 (**Fig. 5 E,F**) which contrasted with protein levels found in naïve

mice and those that resolved infection. Given the importance of CCL2 and CCL3 as monocyte

chemoattractants, these findings further support the hypothesis that acute SARS-CoV-2

replication in fLX is associated with effective recruitment of circulating monocytes. To identify the

potential source of CCL2 and CCL3 expression in fLX upon acute infection, we returned to our

scRNAseq data. While *CCL2* and *CCL3* transcripts were detectable in several fLX human

lineages at naïve state (mesenchymal and endothelial for *CCL2*, and T-cell and myeloid for *CCL3*)

(**Fig. S5 B,C**), three major sources of CCL2 transcripts emerged at 2 dpi, namely iMO, MTC and

interstitial macrophages (**Fig. S5 B**). No significant transcriptomic upregulation of *CCL3* was

detected in fLX consistently with previously reported mechanisms of post-transcription regulation

(Kang et al., 2011; Mao et al., 2019). Unlike CCL2 and CCL3, serum levels of CXCL10 were only

elevated in acutely infected mice (2 dpi) but not in CD4-depleted mice at 12 dpi (**Fig. 5 G**). This

finding is consistent with evidence that iMO are major source of CXCL10 (**Fig. 4 E**), and that iMO-

mediated inflammation is associated with effective viral clearance.

Notably, persistently infected mice can be distinguished from acutely infected (2 dpi) and

isotype-treated (12 dpi) mice by elevated levels of CCL19 in serum (**Fig. 5 H**). CCL19 is a pro-

inflammatory cytokine that has been linked with persistent viral replication and inflammation such

as in the context of HIV-1 infection (Damas et al., 2009), underlining that persistent SARS-CoV-

2 infection in BLTL-L mice recapitulate key immunological features of chronic viral infection.

Unlike acutely infected fLX, persistently infected fLX did not display any significant

histopathological manifestations of disease (**Fig. 5 I**). While the lack of CD4+ cells and reduced

inflammation could promote such an outcome, we have previously reported that fLX exhibit

significant histopathological manifestations of disease upon SARS-CoV-2 infection even in the

absence of engraftment of a human immune system (Kenney et al., 2022). These findings suggest

that persistent SARS-CoV-2 infection remodels the lung tissue environment to better tolerate

sustained viral replication.

Lastly, it has been suggested that persistent SARS-CoV-2 infection can give rise to stable

mutations and eventually, to novel variants (Machkovech et al., 2024; Maponga et al., 2023).

While we failed to acquire viral reads of acceptable quality from persistently infected fLX at 12

dpi, we successfully recovered viral genome sequences from acutely infected fLX at 2 dpi, prior

to infection resolution. 75% of 2 dpi-associated viral sequences exhibited the selection of two

stable mutations not observed in our inoculum and located in the Spike N-terminal domain (NTD):

an insertion (216KLRS) and a non-synonymous mutation (R245H) (**Fig. 5J,K**; **Table S1**.

Interestingly, these two mutations were found together in 100% of the viral sequences, suggesting

potential co-evolution (**Fig. 5L**), and have been reported as positively selected in the context of

sub-optimal neutralizing antibody concentration (216KLRS) (PMDA, 2021) or in the context of

cross-species adaptation (216KLRS and R245H) (Fagre et al., 2021). The ability of SARS-CoV-

2 proteins to undergo rapid mutational selection with reported functional relevance strongly

support the potential of BLT-L mice to predict viral adaptation associated with persistent infection

and the potential emergence of novel variants.

Non-hematopoietic responses to infection likely contribute to effective resolution.

Finally, we aimed to further leverage our scRNAseq datasets to examine the contribution of

the mesenchymal and endothelial compartment in mediating effective infection resolution.

Relative expansion of the mesenchymal compartment upon infection was mainly driven by an

increase in fibroblast populations (**Fig. 2 C; Fig. 6 A-C**). An increase in alveolar fibroblasts was

observed at 12 dpi (49.6%; naïve=28.1%) which was preceded by the emergence of a 2 dpi-

specific cluster of activated fibroblasts (26.5%; naïve=8.9%), consistent with effective tissue repair

in response to lung damage (**Fig. 6 A-C**). Activated fibroblasts were the dominant mesenchymal

population during antiviral responses, as displayed by robust upregulation of interferon stimulated

genes (*LY6E*, *BST2*, *ISG15* and *IFI44L*; (**Fig. 6 D**) and cytokine signatures, which included major

pro-inflammatory mediators such as *CXCL10* and *TIMP1* (**Fig. 6 E-G**). Elevated expression of

mitochondrial encoded genes at 12 dpi was also observed, which we hypothesize to be reflective

of high metabolic activity.

The endothelial compartment is an important modulator of immune recruitment through

cytokine/chemokine signaling. Infected fLX showed the emergence of an activated, 2 dpi-specific

endothelial cell cluster (**Fig. 6 H,I**), which was identified by increased expression of several

transcripts involved in myeloid chemotaxis (**Fig. 6 J, Fig. S5 D**), including *CXCR4* and *MCAM*

(Gupta et al., 1998; Leroyer et al., 2019). In addition, transcripts of INSR, which can

downregulates CXCR4 expression through binding with insulin (Payankaulam et al., 2019;

Rathjen et al., 2022), were also found to be upregulated (**Fig. 6 J, Fig. S5 E,F**).

Venous endothelial cells displayed significant upregulation of a panel of interferon-stimulated

gene transcripts (*XAF1, IFIT1, MX1, IFI44L, RSAD2)* at 2 dpi (**Fig. 6 L**) underscoring an active

antiviral response. Downregulation of several transcripts coding for activation markers (*CD69*),

MHC genes (*HLA-DPB1, HLA-C*) and major proteins regulating cellular metabolism and

transcription (*HSP90AA1, RPL13A*) was reflective of endothelial dysfunction and stress (**Fig. 6**

L); consistent with human patient reports and animal studies (Kadiyska et al., 2022; Otifi and

Adiga, 2022; Qin et al., 2021; Xu et al., 2023). At 12 dpi, venous endothelial cells upregulated

expression of mitochondrial genes, which was similar to the transcriptomic pattern of activated

fibroblasts. Increases in mitochondrial transcripts were also observed in other endothelial clusters

(lymphatic endothelial cells and capillary cells) at 12 dpi (**Fig. 6 L**). This consistent finding across

multiple cell types in the endothelial compartment highlights the elevated metabolic activity

associated with tissue repair following infection resolution.

Collectively, BTL-L mice recapitulate a wide panel of cellular and transcriptomic hallmarks of

SARS-CoV-2 pulmonary infection observed in COVID-19 patients, and support a contribution of

the endothelium in driving infection resolution by enhancing monocyte recruitment.

DISCUSSION

As our appreciation of the immunological differences between mice and humans continue to grow,

humanized mouse models are increasingly standing out as robust platforms to understand how

viral pathogens interact with human tissues, and with the human immune system. These models

are especially valuable when investigating tissue and mucosal immunity, since such

investigations remain impractical in human patients.

The SARS-CoV-2 pandemic has emphasized the need to increase our understanding of an

effective, or failing, pulmonary immune responses during coronavirus infection. Mice engrafted

with human immune systems and genetically matched human lung tissues are valuable

alternatives to investigate lung-resident human immune responses to coronavirus infection (Di et

al., 2022; Kenney et al., 2022; Sun et al., 2022a; Wahl et al., 2019), bridging the limitations of

conventional animal models and the challenges associated with human studies. Using the HNFL

mouse model, we previously reported that such models can be leveraged to capture

immunological signatures defining effective control of SARS-CoV-2 infection (Kenney et al.,

2022). However, the enhanced myeloid reconstitution of the HNFL model rapidly inhibits

extensive viral replication in fLX, precluding our ability to study protective immunological

mechanisms that are at play upon acute, and potentially symptomatic, infection.

The goal of this study was to fill in this knowledge gap and uncover the array of lung-resident

immune mechanisms promoting effective control of acute SARS-CoV-2 infection. In contrast to

the HNFL mouse model, we postulated that the lesser myeloid engraftment and the more

prominent lymphoid reconstitution of BLT-L mice would enable us to more effectively mimic acute

infection and subsequent resolution in a human immunocompetent context. Consistently, BTL-L

mice were effectively able to clear both infectious viral particles and viral RNA following an early

peak of viral replication in fLX. Infection resolution was associated with a rapid mobilization of the

human immune cells into fLX upon viral inoculation, which aligns with previous evidence that

immunodeficient mice only engrafted with fLX are unable to clear infection (Kenney et al., 2022;

Wahl et al., 2021).

Here we identified two major hallmarks of protective lung-resident human immune responses

to SARS-CoV-2 infection. The first was the transient differentiation of circulating monocytes as

extravascular iMO and their role as primary mediators of antiviral responses. To the best of our

knowledge, this study represents the initial evidence of a direct role played by human lung

extravascular inflammatory monocytes in the resolution of respiratory viral infections. Several

lines of evidence support the monocyte nature of these cells. This include their close

transcriptomic proximity with patrolling monocytes which have been reported to give rise to

transient, extravascular (including alveolar) monocytes in the mouse lung (Schyns et al., 2019),

as well as their resemblance to monocytes reported in the BAL of COVID-19 patients (Wendisch

et al., 2021) and from the lung of human-immune system engrafted mice at steady state (Evren

et al., 2021).

The upregulation of myeloid recruitment signals and chemokines, including CCL2 and CCL3,

suggested a strong association between infiltration of circulating monocytes and iMO

differentiation. Abrogation of effective viral clearance following systemic depletion of CD4+

monocytes, but not CD3+ T-cells, was consistent with this model. Elevated serum levels of CCL2

and CCL3 persisted at 12 dpi in CD4-depleted animals, unlike mice that cleared infection at a

similar time point, further underscoring a connection between active viral replication and

monocyte recruitment signals in fLX. While iMO were identified as a substantial source of *CCL2*

transcripts at 2 dpi, CCL2 serum concentration was similarly elevated between CD4-depleted

animals (12 dpi) and non-depleted acutely infected animals (2 dpi) suggesting that MTC may

compensate for limited CCL2 production in absence of defective iMO differentiation during

infection.

The second hallmark of effective control of SARS-CoV-2 infection in fLX that we identified was

the detection of MTC, a T-cell population highly enriched in viral RNA. MTC exhibited expression

of key macrophage-defining genes and elevated cytotoxic activity. While this population presents

some similarity with innate-like T-cells (Molgora and Colonna, 2022), which harbor cytotoxic

signatures and whose roles in cancer have been investigated, MTC appear to be unique given

their combination of lymphoid, myeloid and mesenchymal markers and their short-lived, apoptotic

nature suggested by their complete absence at 12 dpi. MTC also share transcriptomic similarities

with γδ T-cell subsets previously described as bearing myeloid and cytotoxic functions (Lebedev

et al., 2021; Muto et al., 2015). However, their expression of conventional macrophage makers

emphasizes their uniqueness. Additional investigation is required to better our understanding of

the identity and functions of this potentially novel cell subset. However, it is tempting to speculate

that MTC may uniquely synergize CD8+ cytotoxic T-cell functions with viral particle phagocytosis

to effectively control infection prior to undergoing rapid cell death.

Severe COVID-19 has been associated with elevated macrophage infiltration, inflammatory

responses and profibrotic phenotypes (Delorey et al., 2021a; Rendeiro et al., 2021; Wauters et

al., 2021; Wendisch et al., 2021), supporting an overall picture in which excessive myeloid

inflammation and fibrotic response -despite promoting infection resolution- drive potentially fatal

acute respiratory distress syndrome (ARDS) and fibrosis. SARS-CoV-2 can trigger abortive

infection of both monocytes and macrophages (Boumaza et al., 2021b; Junqueira et al., 2022;

Yang et al., 2020; Zheng et al., 2021), and viral RNA replication and protein production in these

cells results in inflammasome activation and exacerbated inflammation (Junqueira et al., 2022;

Munnur et al., 2021; Sefik et al., 2022). However, we observed that viral RNA in iMO was

associated with tissue protection, transient and balanced inflammation (with no inflammasome

activation). By 12 dpi and infection resolution, fLX were nearly devoid of iMO but enriched in

patrolling monocytes. In mice, patrolling monocytes can give rise to lung extravascular

monocytes, which exhibit a dynamic transition state and can differentiate into CD206- interstitial

macrophages (Schyns et al., 2019). Our findings, in conjunction with reports indicating that severe

COVID-19 correlates with the persistence of monocyte-derived profibrotic macrophages

(Wendisch et al., 2021), suggest that the rapid reversion of iMO toward a patrolling phenotype

following encounter with viral RNA, rather than differentiation into interstitial macrophages,

represents a hallmark of protection from COVID19 disease. The limited antiviral responses and

major population contraction of interstitial macrophages at 2 dpi (**Fig. 4 B, D**) supported the notion

that circulating monocytes and/or iMO are not extensively differentiating into interstitial

macrophages upon infection.

The contribution of T-cells in infection resolution in our model remains elusive, particularly in light

of the inability of CD3-depleted mice to fully clear infection (50% of animals cleared infection). It

is worth noting that even when viral infection persisted in CD3-depleted animals, viral titers in fLX

remained significantly lower than in the fLX of persistently infected CD4-depleted animals (**Fig.**

S5 G). This suggests that CD4+ cells, mainly monocytes rather than T-cells, are critical to viral

clearance. Our mIHC data supports this hypothesis by revealing greater numbers of CD163+ cells

in fLX of CD3-depleted mice that were able to effectively clear infection at 12 dpi compared to

those that remained persistently infected. Therefore, while T-cells may contribute to viral

clearance, their absence could be compensated by increased iMO responses of various

magnitudes, ultimately resulting in complete or partial viral clearance. Given the significant

lymphoid depletion observed in fLX at 2 dpi and the limited antiviral responses mounted by CD4+

and CD8+ T-cells, one could then postulate that CD3+ T-cell depletion may interfere with the

differentiation and/or effective function of MTC in infected fLX, and that the lack of MTC may be

totally or partially compensated by a more extensive iMO response. How CD3+ depletion may

impact MTC, and how MTC and iMO functions may synergize to effectively control infection and

prevent extensive inflammation-induced tissue damage will need to be investigated.

Our data suggest a unique immunological model that defines cellular mechanisms leading to

effective control of SARS-CoV-2 infection in the human lung (**Fig. 7**). Upon SARS-CoV-2 infection

of fLX, initial epithelial infection triggers the activation of fibroblasts and endothelial cells. While

fibroblasts potentiate initial myeloid inflammatory responses, endothelial activation and CCL2

secretion by MTC drives the recruitment of patrolling monocytes, which subsequently differentiate

into iMO. iMO then exert a bimodal function: they "sponge" viral particles and components, and

exert tissue-wide antiviral responses protecting the epithelial compartment from *de novo* infection.

In parallel, MTC independently complement the "sponging" functions of iMO by adsorbing viral

particles. A defining hallmark of mild disease is the ability of patrolling monocyte to transiently

differentiate into iMO prior to reverting back into a patrolling monocyte state, instead of upholding

an inflammatory phenotype and further differentiating into inflammatory interstitial macrophages.

Several aspects of this model will need to be further investigated. Understanding the cellular

and molecular drivers defining the fate of infiltrating monocytes upon exposure to SARS-CoV-2,

and how key regulatory cross-roads in this cell type define the course of disease, is of particular

interest. Compounding immune dysregulations, such as elevated inflammatory baseline and/or

specific myeloid imprinting, may contribute to monocyte's inability to adequately regulate their

activation in a timely manner upon encountering viral materials, leading to rapid differentiation

into macrophages and exacerbated inflammation. Investigating the role of fibroblast and

endothelial activation in driving effective clearance mechanisms as well as these cells'

contribution to iMO differentiation is also of particular significance. It remains unclear as to how

endothelial and fibroblasts are activated during infection as neither cell population displayed signs

of viral RNA enrichment. Whether activation of these cells is triggered indirectly by early epithelial

and/or myeloid activation, or via direct exposure to viral components such as the nucleocapsid

protein (Qian et al., 2021) could be investigated using the BLT-L mouse model.

Patients suffering from immunodeficiencies can experience persistent SARS-CoV-2 replication in

the respiratory tract for over 100 days (Abe et al., 2021; Hettle et al., 2022; Sepulcri et al., 2021).

Persistent infection in immunodeficient individuals has a strong public heath significance, since

extended periods of replication in a single host has been postulated to favor the emergence of

variants of concern (Hettle et al., 2022; Shrestha et al., 2022). The ability of the BLT-L mouse

model to recapitulate productive and persistent SARS-CoV-2 infection in human lung tissues

through induction of a specific immunodeficiency opens avenues to dissect viral and host

adaptations favoring viral persistence in a human tissue context. Further, this could shed light into

how these adaptations can contribute to viral evolution and antigenic divergence. Persistent

SARS-CoV-2 infection in BLT-L mice associated with three major features: (1) persistent CCL2

and CCL3 production, (2) absence of lung histopathological manifestation of disease and (3)

elevated serum concentration of CCL19.

CCL19 can potentiate antiviral immune responses by promoting dendritic cell and T-cell

migration into lymph nodes through CCR7 binding (Yan et al., 2019). However, the CCR7-CCL19

axis can also contribute to viral pathogenesis as elevated serum levels of CCL19 correlate with

severe COVID-19 (Tveita et al., 2022) and persistent inflammation in untreated HIV-1 patients

(Damas et al., 2009). For instance, CCL19 mediated inflammation can enhance immune tolerance

by promoting CCR7 internalization (Kohout et al., 2004) and anti-inflammatory signaling (Serrano-

Lopez et al., 2021).

Our current knowledge on CCL19 functionality during viral infection suggests that persistent

infection in BLT-L mice drives tissue remodeling processes typical of chronic inflammation and

potentiates mechanisms of immune tolerance. Histopathological manifestations of disease in fLX

upon acute SARS-CoV-2 infection are persistent in mice engrafted solely with fLX (Kenney et al.,

2022; Wahl et al., 2020) but are transient in BTL-L mice, underscoring that such cytopathic

manifestations are not driven by human hematopoietic cells but rather by extensive viral

replication. Thereby, the absence of histopathological manifestations of disease observed in fLX

upon persistent SARS-CoV-2 replication (**Fig. 6 I**) support evidence of a tissue-wide remodeling

toward viral tolerance. The remarkable ability of BLT-L mice to support the rapid emergence of

adaptive mutations in the viral genome previously reported to confer increased fitness *in vitro* and

in vivo, emphasizes the strong potential of this mouse model to recapitulate not only host, but

also viral adaptations underlying persistent SARS-CoV-2 infection and to predict potential

evolutionary courses.

It is important to note that there are inherent limitations associated with the BLT-L mouse model.

First, direct viral inoculation into fLX potentially bypasses key immune checkpoints in the upper

respiratory tract, which are not recapitulated in our model. Second, the hematopoietic

reconstitution and functions of the BLT-L mouse model remain imperfect, notably through the

underrepresentation of specific hematopoietic subsets, including dendritic cells and granulocytes,

and limited B-cell responses – which may ultimately bias the dynamics of viral clearance. Despite

these limitations, BLT-L mice recapitulated human lung responses to SARS-CoV-2 infection

through AT2 loss of programming, fibroblast and endothelial activation, effective clearance of

infection through robust hematopoietic responses. This mouse model also supported viral

adaptation processes with public health relevance, and recapitulated signatures of persistent viral

replication that are consistent with patients suffering from chronic infection. Together, our work

unravels the potential of the BLT-L mouse model to uncover critical human mechanisms

governing effective resolution of coronavirus infection. It highlights the value of this mouse model

to identify novel immunotherapeutic strategies against coronaviral diseases and enhance our

comprehension of the host and viral mechanisms underlying SARS-CoV-2 persistent infection

and evolution.

MATERIALS AND METHODS

Institutional approvals

All animal experiments described in this study were performed in accordance with protocols that

were reviewed and approved by the Institutional Animal Care and Use and Committee of the

Ragon Institution and Boston University (PROTO202000020). All mice were maintained in

facilities accredited by the Association for the Assessment and Accreditation of Laboratory Animal

Care (AAALAC). All replication-competent SARS-CoV-2 experiments were performed in a

biosafety level 3 laboratory (BSL-3) at the Boston University National Emerging Infectious

Diseases Laboratories (NEIDL).

Resources

Mouse strains. Female NOD.Cg.-$Prkdc^{Scid}Il2rg^{tm1Wjl}$/SzJ (NSG) mice were obtained from the

Jackson Laboratory, catalog number 005557. NSG mice were maintained by the Ragon Institute

Human Immune System Mouse core prior to engraftment and shipment to the NEIDL, Boston

University.

Human fetal tissues. De-identified human fetal tissues were procured from Advanced Bioscience

Resources (Alameda, CA, USA).

Cell lines. VeroE6 cells were grown in Dulbecco's modified Eagle's medium (DMEM)

supplemented with 10% heat inactivated fetal bovine serum (Bio-Techne, R&D systems,

Minneapolis, MN, USA) and 1% (v/v) Penicillin Streptomycin (Thermo Scientific, Waltham, MA,

USA).

Antibodies. The anti-mouse antibody CD45-PE-Dazzle5 clone 30-F11 was purchased from

BioLegend (San Diego, CA, USA) and used for flow cytometry. The following anti-human

antibodies were used for flow cytometry and were all acquired from BD Biosciences (San Jose,

CA, USA): HLA-DR-BV510 clone L243, CD3-BV605 clone UCHT1, CD20-BV650 clone 2H7,

CD16 (FcγRIII)-BV117 clone B73.1, CD45-BV785 clone H130, CD8-FITC clone RPA-T8, CD33-

PE clone WM53, CD14-PerCP-Cy5.5 clone HCD14, CD45RA-PE-Cy7 clone HI100, CD56-

allophycocyanin clone QA17A16, CD4-Alexa Fluor 700 clone SK3. The following anti-human

antibodies were used for *in vivo* depletions: From Bio-X Cell (Lebanon, NH, USA): anti-human

CD3 clone OKT3, anti-human CD4 clone OKT4, anti-human CD8 clone OKT8; From

Thermofisher: mouse IgG2a isotype. The anti-SARS-CoV-2 (2019-nCoV) Spike RBD-mFc clone

#57 used as a control for neutralization assays was acquired from Sino Biologicals.

Experimental methods

Generation of BLT-L mice. BLT mice were generated via irradiation of female NOD.Cg.-

Prkdc^{Scid}Il2rg^{tm1Wjl}/SzJ mice (NSG mice; Jackson Laboratory #005557) with 200 rads from an X-

ray irradiator. Mice then underwent surgery to implant 1mm^3 of human fetal thymic and fetal liver

tissue (Advanced Bioscience Resources) under the murine kidney capsula. After the blunt

resection of the thoracal subcutis, through the same surgical opening, two pieces of homologous

human fetal lung tissue into the subcutaneous dorsal area of the 8-15 week old females. Post-

implantation, mice intravenously received 1x10^5 homologous CD34+ cells. Human immune

reconstitution was determined by flow cytometry at weeks post implantation. Three distinct human

donors were used for this study.

Generation of SARS-CoV-2 isolate and recombinant SARS-CoV-2 NanoLuc viral stocks. All replication-competent SARS-CoV-2 experiments were performed in a BSL-3 facility at the Boston University National Emerging Infectious Diseases Laboratories. The clinical isolate named 2019-nCoV/USA-WA1/2020 strain (NCBI accession number: MN985325) of SARS-CoV-2 was obtained from BEI Resources (Manassas, VA, USA). Recombinant SARS-CoV-2 expressing NanoLuc Luciferase (rSARS-CoV-2 NL) (Xie et al., 2020) was graciously provided by the Laboratory of Pei-Yong Shei. Viral stocks were grown and purified as previously described (reference HNFL manuscript). Briefly, 1×10^7 Vero E6 cells were plated in a T-175 flask the day prior to virus generation. The next day, cells were infected with virus diluted in 10 mL of Opti-MEM (ThermoFisher Scientific, Waltham, MA, USA) and incubated for 1 h at 37°C to allow for virus adsorption. After 1 h, 15 mL of DMEM containing 10% FBS and 1% penicillin/streptomycin was added to cells and incubated overnight. The next day, media was removed, cells were rinsed with 1X PBS, pH 7.5 (ThermoFisher Scientific) and 25 mL of fresh DMEM containing 2% FBS was added. Cells were assessed for cytopathic effect (CPE), media was collected, filtered with a 0.22 µm filter, and concentrated by sucrose gradient. Concentrated virus was suspended in sterile 1X PBS, pH 7.5, aliquoted and stored at –80C.

Viral quantification by plaque assay. The titer of our viral stocks was determined by plaque assay. Vero E6 cells were seeded into a 12-well plate at a density of 2×10^5 cells per well. The next day, cells were infected with 10-fold serial dilutions of the viral stocks and incubated for 1 hour at 37°C to allow for viral adsorption. After 1 hour incubation, 1 mL of overlay media (1.2% Avicel (DuPont, Wilmington, DE, USA; RC-581) in DMEM supplemented with 2% FBS and 1% Pen/Strep) was added to each well. Three days later, the overlay media was removed, and cells were fixed for 1 hour at room temperature with 10% neutral buffered formalin (ThermoFisher Scientific). Formalin was removed after fixation and cells were stained for 30 min at room temperature with 0.1% crystal violet (Sigma-Aldrich) prepared in 10% ethanol/water. Crystal violet

was removed, cells were washed with water and plaque forming unites were counted to determine

viral titers.

Mouse inoculation with SARS-CoV-2. BLT-L mice were anesthetized with 1–3% isoflurane prior

to inoculation via subcutaneous, intra-fetal lung xenograft (intra-fLX) injection with 10^6 plaque

forming unites (PFU) of SARS-CoV-2 WA-1 isolate in 50 µL of sterile 1X PBS. Mice were

euthanized at 2-, 6-, and 12-days post inoculation.

In vivo antibody depletion. BLT-L mice were administered 200 mg of anti-CD3e (OKT3)

(BioxCell; cat. # BE0001-2), anti-CD4 (OKT4) (BioxCell; cat. # BE0003-2), anti-CD8 (OKT8)

(BioxCell; cat. # BE0004-2), or isotype IgG2a (Thermofisher; cat # 02-6200) antibody 3-, 2-, and

1-day prior to inoculation and 4- and 8-days post inoculation with 1x10^6 PFU SARS-CoV-2 WA-

1.

Generation of single cell suspension from mouse blood. Blood processing was performed as

previously described (Kenney et al (2022) *Cell Reports*). Briefly, blood was collected via

submandibular bleeding or heart puncture and transferred into EDTA capillary collection tubes

(Microvette 600 K3E; Sarstedt, Nümbrecht, Germany). Blood was centrifuged at 3,500 RPM for

10 minutes at room temperature, serum was removed, and whole blood was subject to ACK lysing

(ThermoFisher Scientific; #A1049201) for 10 minutes at room temperature. Lysing was quenched

with 10% (v/v) FBS DMEM media and cells were washed twice with a 1% (v/v) FBS-PBS solution

(FACS Buffer) before downstream processing.

Generation of single cell suspension from fLX. Fetal lung xenografts were processed as

previously described (Kenney et al., 2022). Briefly, lung tissues were placed in a 60 mm dish,

minced using a disposable scalpel, then transferred into a 15 mL conical tube containing 3 mL of

digestion buffer (HBSS minus Ca2+, Mg2+, and phenol red, 0.5 mg/mL Liberase TM, 1 mg/mL

DNase I) and incubated at 37°C for 30 min with agitation every 10 min. After digestion, tissue was

passed through a 70 µm strainer on a 50 mL tube, mashed using the plunger of a 3 mL syringe

and washed twice with 1 mL FACS buffer. Cell suspension was centrifuged at 300 x g for 5

minutes at 4°C, suspended in 1 mL of ACK lysing buffer, and incubated for 2 min at room

temperature. After incubation, lysis was quenched with 12 mL of FACS buffer, cell suspension

was centrifuged at 300 x g for 5 min at 4°C, and the cell pellet was suspended in 1 mL of FACS

buffer prior to downstream processing.

Viral RNA extraction from serum. Viral RNA was extracted from serum using a Zymo Viral RNA

extraction kit (Zymo Research, Irvine, CA, USA: #R1035) following the manufacturers protocol.

Briefly, serum was mixed with RNA/DNA shield (Zymo) at a 1:1 ratio. RNA buffer was then added

to the serum (2:1 ratio) and passed through a column by centrifugation at 13,000 x g. The column

was then washed twice, and RNA was eluted with 15 µL of RNase/DNase free water.

RNA extraction from fLX. During necropsy a piece of fLX tissue was placed in 600 µL of

RNAlater (MilliporeSigma: #R0901500ML) and stored at −80°C until analysis. For RNA extraction,

20–30 mg of tissue was weighed out, placed into a 2 mL tube with 600 µL of RLT buffer with 1%

β–mercaptoethanol and a 5 mm stainless steel bead (Qiagen, Hilden, Germany: #69989), then

dissociated using a Qiagen TissueLyser II (Qiagen) with the following cycle: two min dissociation

at 1800 oscillations/min, one min rest, two min dissociation at 1800 oscillations/min. Dissociated

tissues were centrifuged at 13,000 rpm for 10 min at room temperature and cleared supernatant

was transferred to a new 1.5 mL eppendorf tube. RNA extractions were performed using a Qiagen

RNeasy Plus Mini Kit (Qiagen: #74134), according to the manufacturer's instructions (Qiagen:

#79256). RNA was eluted in 30 µL of RNase/DNase free water and quantified by Nanodrop.

fLX processing for viral titer quantification. Tissue pieces stored in RNALater were thawed at

room temperature and then 22-40 mg of tissue were weighed out and placed into 600 uL of

OptiMEM (ThermoFisher Scientific). Tissues were homogenized using a Qiagen TissueLyser II

with two dissociation cycles, centrifuged and supernatant was transferred to a new 1.5 mL

Eppendorf tube (see "*RNA extraction from fLX*" section). Tissue homogenates were then serial

diluted (10^{-1} – 10^{-6}) and 300 uL of supernatant was plated on VeroE6 cells in 12-well plates (2×10^5

cells/well). Supernatant was incubated for 1 hour at 37°C to allow for viral adsorption then 1 mL

of a 1:1 mixture of 2X DMEM 4% FBS 2% penicillin/streptomycin and 2.4% Avicel (Dupont) was

overlaid onto each well. Plates were incubated for 72 hours at 37°C with 5% CO_2. Avicel was then

removed, cells were fixed in 10% buffered formalin (ThermoFisher Scientific) for 1 hour, then

stained with 0.1% crystal violet in 10% ethanol/H_2O for 30 minutes before washing and

quantification.

Serum neutralization assay. One day prior to the experiment, 1×10^4 Vero E6 cells were plated

into a 96-well plate. Serum was decomplemented at 56 °C for 30 min and an initial dilution of 1:10

was prepared in OptiMEM. Two-fold dilutions were subsequently prepared and mixed with

rSARS-CoV-2 NL virus (MOI = 1) for 1 h at room temperature and then plated onto cells. After a

1-h incubation at 37 °C inoculum was removed and 200 µL of fresh DMEM containing 2% FBS

and 1% penicillin/streptomycin was added. After a 24 h incubation at 37 °C with 5% CO2 media

was removed and cells were fixed with 10% formalin for 1 h. A SARS-CoV-2 spike neutralizing

antibody (Sino Biological Inc., Beijing, China; 2 µg/µL; Clone #57) was used as a positive control.

Cells were washed with 1× PBS and 20 µM furimazine (MedChem Express, Monmouth Junction,

NJ, USA) luciferin substrate was added onto cells. Cells were then imaged using an IVIS spectrum

imager (PerkinElmer) and analyzed using LivingImage software (PerkinElmer). Titers were

determined as the reciprocal of the highest dilution with >50% reduction of cytopathic effect.

Quantification of peripheral human chimerism in BLT-L mice. 2–4x10^6 PBMCs of human or murine origin were isolated as described above and stained for 1 hour at 4°C in the dark with an antibody cocktail targeting human(h)CD45, mouse CD45, hCD3, hCD4, hCD8, hCD16, hCD19, hCD11c, hCD56 and hCD14. Following washing with FACS Buffer, cells were fixed with fixation buffer (1% (v/v) FBS, 4% (w/v) PFA in PBS) for 30 min at 4°C in the dark. Chimerism of all humanized mice was assessed by quantifying the following human populations: Human CD45+, human CD45+ murine CD45-; T-cells, CD45+ CD3+; CD4+ T cells, CD45+ CD3+ CD4+; CD8+ T cells, CD45+ CD3+ CD8+; CD45+ CD16+ leukocytes; B-cells, CD45+ CD19+; conventional dendritic cells, CD45+ CD11c+; NK/NKT cells, CD45+ CD56+; Monocytes, CD45+ CD14+.

Antibody staining. After generation of single cell suspension, 5x10^5 - 1x10^6 cells were used for flow cytometry staining. Cells were centrifuged at 300 X *g* for 5 min at 4°C. The cell pellet was resuspended in a mix of 22.5 µL FACS buffer and 2.5 µL of FcX (Biolegend; #422302) and incubated for 10 min at room temperature. After blocking, 25 µL of antibody cocktail targeting hCD3, hCD20, hCD16, hHLA-DR, hCD45, hCD8, hCD4, hCD33, hCD45RA, hCD56, hCD14, mCD45, and containing a LIVE/DEAD viability dye (ThermoFisher Scientific) was added to each sample and incubated in the dark for 30 min at 4°C. After staining, 1 mL of FACS buffer was added to each sample, samples were centrifuged at 300 x g for 5 min, washed with 1 mL FACS buffer, centrifuged at 300 x g for 5 min, and then fixed in 200 µL 4% PFA for 30 min. After fixation cells were washed twice with 1 mL FACS buffer, resuspended in FACS buffer, and stored protected from light at 4°C until analysis. Human immune cell subsets were gated as follows: human CD45+, hCD45+ mCD45-; human CD3+, hCD45+ hCD3+; human CD4+, hCD45+ hCD3+ hCD4+; human CD8+, hCD45+ hCD3+ hCD8+.

Flow cytometry analysis. For all flow cytometry experiments, flow cytometric analysis was performed using an LSRII Flow Cytometer (BD Biosciences) or Cytek Aurora instrument (Cytek

Biosciences). Flow cytometry fluorophore compensation for antibodies was performed using an

AbC™ Anti-Mouse Bead Kit (ThermoFisher Scientific). Flow cytometry data were analyzed using

FlowJo software (TreeStar, Ashland, OR, USA).

Histologic processing and analysis. Tissue samples were fixed for 72 h in 10% neutral buffered

formalin, processed in a Tissue-Tek VIP-5 automated vacuum infiltration processor (Sakura

Finetek USA, Torrance, CA, USA), followed by paraffin embedding using a HistoCore Arcadia

paraffin embedding station (Leica, Wetzlar, Germany). Generated formalin-fixed, paraffin-

embedded (FFPE) blocks were sectioned to 5 µm using a RM2255 rotary microtome (Leica),

transferred to positively charged slides, deparaffinized in xylene, and dehydrated in graded

ethanol. Tissue sections were stained with hematoxylin and eosin for histologic examination, while

unstained slides were used for immunohistochemistry. Qualitative and semi-quantitative

histomorphological analyses were performed by a single board-certified veterinary pathologist

(N.A.C.). An ordinal scoring system was developed to capture the heterogeneity of histologic

findings in fLX. Individual histologic findings that were scored included: syncytial cells, hyaline

membranes, intra-airspace neutrophils and necrosis, hemorrhage, edema, denuded

pneumocytes, capillary fibrin thrombi, intermediate vessel fibrin thrombi and coagulative necrosis.

The entire fLX was examined at 200x with a DM2500 light microscope (Leica) using the following

criteria: 0 = not present, 1 = found in <5% of fields, 2 = found in >5% but <25% of fields, or 3 =

found in >25% of fields. Several criteria were also restricted to 'not observed' (0) or 'observed'

(1). Scores were combined to generate a cumulative lung injury score.

Histological Antibodies. The following primary antibodies from Cell Signaling Technology

(Danvers, MA, USA) were used: rabbit anti-human/mouse PAPRg (clone C26H12), rabbit anti-

human CD4 (clone EP204), rabbit anti-human CD163 (clone D6U1J), rabbit anti-human CD3e

(clone D7A6E), mouse anti-MHC class I (clone EMR8-5), and mouse anti-SARS-CoV-2 Spike (S)

(clone E7U6O/2B3E5]). The secondary antibody used in this study included HRP Goat anti-Rabbit

IgG (H&L) (Vector Laboratories, Burligame, CA, USA). For mouse derived primary antibodies, a

linker antibody (Abcam) was used prior to application of the secondary antibody to prevent non-

specific binding. DAB chromogen (Roche) and chromogens used for TSA-conjugated Opal 480,

520, 570, and 620 fluorophores (Akoya Biosciences, Marlborough, MA, USA) were utilized to

develop immunohistochemical assays. The following anti-SARS-CoV-2 antibodies were used for

immunohistochemistry: rabbit polyclonal anti-SARS-CoV Nucleoprotein (Novus Biological,

Littleton, CO, USA), mouse monoclonal anti-SARS-CoV-2 Spike clone 2B3E5 (This antibody was

used in this study as clone E7U60, which was the pre-production clone ID of clone 2B3E5; Cell

Signaling Technology).

Multispectral fluorescent imaging. Fluorescently labeled slides were imaged using either a

Mantra 2.0TMor Vectra Polaris TM Quantitative Pathology Imaging System (Akoya Biosciences).

To maximize signal-to-noise ratios, fluorescently acquired images were spectrally unmixed using

a synthetic library specific for the Opal fluorophores used in each assay plus DAPI. An unstained

fLX section was used to create an autofluorescence signature that was subsequently removed

from multispectral images using InForm software version 2.4.8 (Akoya Biosciences).

Transmission electron microscopy. Tissue samples were fixed for 72 h in a mixture of 2.5%

glutaraldehyde and 2% formaldehyde in 0.1 M sodium cacodylate buffer (pH 7.4). Samples were

then washed in 0.1 M cacodylate buffer and postfixed with 1% osmiumtetroxide (OsO4)/1.5%

potassiumferrocyanide (KFeCN6) for 1 h at room temperature. After washes in water and 50 mM

maleate buffer pH 5.15 (MB), the samples were incubated in 1% uranyl acetate in MB for 1 h,

washed in MB and water, and dehydrated in grades of alcohol (10 min each: 50%, 70%, 90%,

2x10 min 100%). The tissue samples were then put in propyleneoxide for 1 h and infiltrated

overnight in a 1:1 mixture of propyleneoxide and TAAB Epon. The following day the samples were

embedded in fresh TAAB Epon and polymerized at 60°C for 48 h. Semi-thin (0.5 µm) and ultrathin

sections (50-80 nm) were cut on a Reichert Ultracut-S microtome (Leica). Semi-thin sections were

picked up on glass slides and stained with toluidine blue for examination by light microscopy to

find affected areas in the tissue. Ultrathin sections from those areas were picked up onto

formvar/carbon coated copper grids, stained with 0.2% lead citrate and examined in a JEOL

1200EX transmission electron microscope (JOEL, Akishima, Tokyo, Japan). Images were

recorded with an AMT 2k CCD camera.

Bulk RNA sequencing. Total RNA was processed from fLX as described above, and sent to BGI

genomics (Hong Kong, China) for library preparation and sequencing (Pair-ends, 100 bp, 20M

reads per sample). Raw FASTQ files were quality-checked with FastQC v0.11.7. Reads were

found to be excellent quality and were aligned to the combined human (GRCh38, Ensembl 101)

and mouse (GRCm38, Ensembl 101) genomes with STAR v2.7.1a (Dobin et al., 2013) followed

by quantification with featureCounts v1.6.2 (Liao et al., 2014). Quality was checked with MultiQC

v1.6. All samples passed quality thresholds of >75% sequences aligned and >15 million aligned

reads per sample. Significantly up- and downregulated genes were identified with DESeq2

v1.23.10 (Love et al., 2014) in R v3.6.0. Each post-infection timepoint was compared to naïve: 2

dpi vs. naïve, and 12 dpi vs. naïve. P-values were FDR-adjusted, and log2 fold change was shrunk

with the apeglm method. Significance was determined by an FDR-adjusted $p<0.01$ and a

shrunken log2 fold change>2 or <-2. DESeq2 result were imported into Ingenuity Pathway

Analysis (IPA; Service curated by Qiagen; Access provided through the Boston University

Genome Science Institute) (Kramer et al., 2014), and a canonical pathway enrichment analysis

was performed using the default settings and the same differential expression thresholds as

before (shrunken log2 fold change>2 or <-2 and FDR-adjusted p-value<0.01).

Single cell barcoding and sequencing. Following fLX processing into single cell suspension as

described above, cells were frozen down in a 90% FBS (Bio-Techne, R&D systems) 10% DMSO

solution (ThermoFisher Scientific) and kept at -80°C. Four to five days following freezing, cells

were thawed, and viability was assessed using Trypan blue (Fisher Scientific). Samples with

≥90% viability were then processed using the Chromium Next GEM Single Cell 3' GEM, Library

& Gel Bead Kit v3.1, as per manufacturer instructions, and single cell barcoding was performed

using a Chromium instrument (10x genomics) located in the NEIDL BSL-3. Reverse transcription

of RNAs was performed in the BSL-3 using a thermocycler (Applied Biosciences), and cDNA was

then removed from containment. Full-length, barcoded cDNAs were then amplified by PCR to

generate sufficient mass for library construction. Enzyme fragmentation, A tailing, adaptor ligation

and PCR were then performed at the Boston University single-cell sequencing core to obtain final

libraries containing P5 and P7 primers used in Illumina bridge amplification. Size distribution and

molarity of resulting cDNA libraries were assessed via Bioanalyzer High Sensitivity DNA Assay

(Agilent Technologies, USA). All cDNA libraries were sequenced on an Illumina NextSeq 500

instrument at the Boston University microarray and sequencing core according to Illumina and

10x Genomics guidelines with 1.4-1.8pM input and 1% PhiX control library spike-in (Illumina,

USA).

Single-cell RNA sequencing analysis.

Sequencing data was processed via 10X Genomics Cell Ranger 6.0.2. A custom reference was

built from the combined human (GRCh38, Ensembl 101) and mouse (GRCm38, Ensembl 101)

genomes. FASTQ files were generated from raw Illumina BCL files using *cellranger mkfastq*.

FASTQ files were aligned to the custom reference and quantified with *cellranger count*.

Quantification outputs were loaded and analyzed in R with the Seurat package (4.0.2, (Hao et al.,

2021)). Cells with nFeature_RNA between 200 and 7500, and MT.percent<= 50 were analyzed.

The dataset was normalized and scaled, and cell cycle scores (S+G2M) and nCount_RNA were

regressed. Seurat's RunPCA function was used to determine the number of PCA used for

subsequent analysis. FindClusters function was used to cluster cells. RunUMAP function was

used for UMAP visualization. A human score was calculated using all human genes, to

discriminate between mouse and human cells (**Fig. S3A-C**). Human cells were kept for

subsequent analysis. The following ("dimension";"resolution") parameters were used to cluster

the whole human dataset (15;0.1), epithelial cells (10;0.3), mesenchymal (10;0.4), endothelial

(10;0.5), T cell/ILC (12;0.4), and Myeloid cells (10;0.2). B cell, endothelial cell, epithelial cell,

mesenchymal cell, and mast cell clusters were identified by scoring using gene lists defined for

lung subsets in the COVID-19 atlas of ToppCell (**Fig. S3D**) (Jin et al., 2021). For chondrocytes,

which are not present in this dataset, we used the following gene list *ACAN, ITGA10, CYTL1,*

HAPLN1, SNORC, COL2A1, COL9A2, COL11A2, SOX9, SOX6, ZBTB16, FOXC1, and GATA2.

For T cells and ILC, a manually curated gene list based on ToppCell was used. Consensus

signature genes from LGEA LungMAP (Sun et al., 2022b) were used to validate each cluster.

Differentially expressed genes between clusters and conditions were identified using

FindMarkers. Only genes showing expression in at least 10% of cells in one of the groups

compared, p_val_adj < 0.05, and abs(fold change)>2 were considered.

Sequencing of SARS-CoV-2 virus from fLX. Viral sequences were amplified using a modified

version of the ARTIC3 tiled amplicon sequencing protocol (Quick 2020, Tyson 2020) and

sequenced on an Illumina MiSeq V3 600 cycle kit. Briefly, viral RNA was extracted from serum

using X protocol (I don't know the details). cDNA was synthesized from the viral RNA using a mix

of random hexamer and oligo-dT primers (IDT) and SSIV reverse transcriptase (Thermo Fisher).

Tiled amplicons were amplified using Q5 Hot Start polymerase (NEB) and the pooled primers

described in ARTIC3 (Quick 2020, Tyson 2020). Illumina sequencing adaptors were ligated onto

the amplicons using the NEBNext Adaptor kit for Illumina (NEB). The amplicons were then pooled,

cleaned up, and size selected using AMPureXP beads (Beckman Coulter). The pooled library

was quantified using Tape Station (Agilent) and Qbit (Thermo Fisher) kits. The library was

sequenced on a MiSeq with a V3 600 cycle kit (Illumina). Sequencing reads were filtered, trimmed,

and mapped to the D-WA reference sequence (Geneious). Variants were called using the

Geneious variant caller, with a minimum frequency threshold of 25%. Each positive variant was

visually inspected.

Quantification and statistical Analysis. For histopathological score, viral loads/titers

comparison and Spike quantification, Kruskal-Wallis, Kolmogorov-Smirnov non-parametric t-test,

or a two-way ANOVA with Benjamini, Krieger, and Yekutieli correction for multiple comparisons

were applied given the non-continuous nature of the data (i.e., viral inoculation in areas displaying

differential stage of tissue development, and/or differences in fLX engraftment, and/or

donor/gestational age differences). A two-way ANOVA with Benjamini, Krieger, and Yekutieli

correction for multiple comparisons was also used to generate the ISG p.value heatmap given

the non-parametric nature of these data. For quantitative analysis of hematopoietic cell infiltration

(i.e. multiplex IHC and flowcytometry data), Welch's t-test or one-way ANOVA tests were used.

For comparison of viral gene count between macrophage clusters and other clusters, a Mann-

Whitney t-test was used. All statistical tests and graphical depictions of results were performed

using GraphPad Prism version 9.0.1 software (GraphPad Software, La Jolla, CA). For all tests,

$p \leq 0.05$ was considered statistically significant. Statistical significance on figures and

supplemental figures is labelled as follow: *$p \leq 0.05$, **$p \leq 0.01$, ***$p \leq 0.001$, ****$p \leq 0.0001$.

Data availability. The accession number for the raw data of the bulk and single-cell RNA-Seq is

GSE255200.

ACKNOWLEDGMENTS

This work was supported in part by a start-up fund and Peter Paul Career Development Professorship from Boston University (to F.D.), grants from the National Institutes of Health (K22 AI144050 to F.D.), Clinical and Translational Science Awards (grant UL1 TR001430) from the National Center for Advancing Translational Sciences of the National Institutes of Health (to F.D., N.A.C. and A.E.), and an Evergrande MassCPR award (to R.B.). A.B.B. was supported by R01AI174875, R01AI174276, DP2DA040254, as well as CDC subcontract 200-2016-91773-T.O.2. and a Massachusetts Consortium on Pathogenesis Readiness (MassCPR) grant. We also thank the Evans Center for Interdisciplinary Biomedical Research at Boston University Chobanian and Avedisian School of Medicine for their support of the Affinity Research Collaborative on 'Respiratory Viruses: A Focus on COVID19'. This work utilized a Ventana Discovery Ultra and Akoya PhenoImager HT that were purchased with funding from National Institutes of Health SIG grants (S10 OD026983 & S10OD030269). We thank Ronald B. Corley, NEIDL director at the time of this study and MassCPR award recipient, for his constant encouragement and support for this study. We thank the Boston University Animal Science Center, the Ragon Institute Human Immune System Mouse core, the single cell sequencing core and microarray and sequencing core, and the flow cytometry core at Boston University Chobanian and Avedisian School of Medicine, and all the NEIDL animal core staff for their outstanding support. We also thank all the Douam, Balazs, Connor and Crossland lab members, NEIDL members, and members of the department of virology, immunology, and microbiology and pathology at Boston University for their constant support and advice. D.K. is supported by a T32 training grant in immunology (T32AI007309). The authors have no conflicting financial interests.

Author contributions: D.K., A.B.B. and F.D. conceptualized the overall study. C.H. conceptualized the computational component of this study and oversaw the analysis of the single-cell RNA sequencing data. D.K., A.B.B., N.A.C. and F.D. designed the experiments. D.K., A.K.O., M.S, A.N, H.P.G., M.E., V.V., A.B.B. and F.D. performed experiments. D.K., A.K.O., A.T., J.T.,

M.S, A.N, P.M., M.E., J.H.C., V.V., N.A.C., C.H., A.B.B. and F.D. analyzed the data. J.T and C.H.

carried out computational analysis. M.E. carried out electron microscopy analysis. A.B.B. and V.V.

provided access to key resources. D.K. and F.D. wrote the manuscript with contributions from all

authors.

FIGURE LEGENDS

Figure 1. BLT-L mice effectively resolve SARS-CoV-2 infection and repair lung tissue. Fetal

lung xenografts (fLX) of BLT-L mice were infected with 1×10^6 PFU of SARS-CoV-2 WA-1 isolate

via subcutaneous intra-graft injection. **(A)** BLT-L mice were monitored for weight change over the

course of 12-days post infection (dpi). **(B)** Immunohistochemistry for SARS-CoV-2 S protein **(a-**

d) and hematoxylin and eosin staining **(e-f)** were performed on naïve fLX **(a,e)** and fLX at 2 dpi

(b,f), 6 dpi **(c,g),** and 12 dpi **(d,h)**. Scale bar = 100 μM **(C)** Viral titer (log(PFU/mg)) in fLX at 2, 6,

and 12 dpi as determined by plaque assay. **(D)** Histopathological scoring of naïve fLX and fLX at

2, 6, and 12 dpi. Data are representative of two or three independent experiments. n = 6 -10 per

timepoint. Error bars represent mean ± standard error of the mean (SEM). *One-way ANOVA*

analysis was performed. p-values are indicated on graphs and only significant comparisons are

shown.

Figure 2. **Cellular remodeling and viral dynamics in fLX upon SARS-CoV-2 infection**

Single cell RNA sequencing analysis was performed on single cell suspensions of naïve fLX and

fLX at 2 and 12 dpi. Naïve n=3 fLX (9,605 cells), 2 dpi n=2 fLX (4,405 cells), and 12 dpi n = 2 fLX

(5,857 cells). **(A)** UMAP plot clustering of the human cell compartment of naïve fLX and fLX at 2

and 12 dpi. **(B)** Temporal annotation of the human clusters on the UMAP plot: naïve (black), 2 dpi

(red), 12 dpi (blue). Clusters unique to 2 dpi are annotated by A (T cell compartment), B

(mesenchymal compartment), or C (myeloid compartment). **(C)** Frequency of each cell

compartment determined by single cell RNA-sequencing. **(D)** Dot plots displaying the expression

of SARS-CoV-2 viral RNA transcripts identified in scRNA-seq. **(E)** UMAP plot showing the

distribution of SARS-CoV-2 viral RNA transcripts across all human cell clusters and time points.

(F) UMAP plot sub-clustering of the human epithelial compartment of fLX across all time points.

(G) Temporal annotation of the human epithelial sub-clusters on the UMAP plot: naïve (black), 2

dpi (red), 12 dpi (blue). The inset graph indicates the change in frequency of each compartment

(AT2; black, Basal; pink, Secretory; teal, and AT1; purple) within the epithelial compartment. Red

line indicates the change in the AT2 compartment over time. **(H)** Heatmap displaying the relative

expression (from -1 to 1) of the top differentially regulated genes over the course of infection within

the AT2 compartment.

Figure 3. A unique T-cell subset is enriched for viral RNA and specific for 2 dpi. (A) UMAP

plot showing the sub-clustering of the human T-cell compartment of fLX across all time points.

(B) Temporal annotation of the human T-cell sub-clusters on the UMAP plot: naïve (black), 2 dpi

(red), 12 dpi (blue). **(C)** UMAP plot (all time points) showing the distribution of SARS-CoV-2 viral

RNA transcripts in the T-cell compartment. **(D)** Go-term analysis showing the most highly

upregulated signaling pathways in the macrophage-like T-cell cluster (cluster 6). **(E)** Relative

expression of T-cells, macrophage, mesenchymal and antiviral markers within each of the T-cell

sub-clusters (From 0 to 8; as labeled in panel A.).

1002

Figure 4. CD163+ inflammatory monocytes arise during acute viral infection and are

predominant antiviral mediators. (A) UMAP plot showing sub-clustering of the human myeloid

compartment of fLX across all time points. **(B)** Temporal annotation of the myeloid sub-clusters

on the UMAP plot: naïve (black), 2 dpi (red), 12 dpi (blue). iMO cluster is indicated with a blue

circle. **(C)** UMAP plot (all time points) showing the distribution of SARS-CoV-2 viral RNA

transcripts in the myeloid compartment. **(D-F)** UMAP plot showing the expression of CD*163* **(D),**

CD4 **(E)**, and *MRC1* **(F)** across myeloid sub-sets. **(G)** Frequency of each myeloid sub-cluster by

timepoint. **(H)** Relative expression of highly upregulated ISGs and inflammatory markers in each of the myeloid sub-clusters. **(I)** Multiplex fluorescent immunohistochemistry of naïve fLX and fLX at 2 and 12 dpi. CD163: teal, CD4: yellow, CD3: orange, PPARγ: green, Dapi: gray. Two representative images. Scale bar = 100 μM **(J)** Violin plots showing expression level of differentially expressed genes between SARS-CoV-2 positive and negative iMo (cluster 3).

Figure 5. Loss of CD4+ monocytes results in persistent infection. (A) BLT-L mice were administered 200 μg of anti-CD3e (OKT3), anti-CD4 (OKT4), anti-CD8 (OKT8) or IgG2a isotype. **(B)** Viral titer (log(PFU/mg)) in fLX extracted from BLT-L mice treated with isotype, OKT3, OKT4, or OKT8 antibody at 12 dpi. **(C)** MHC-I staining of fLX tissue sections (anti-MHC class I (EMR8-5) CST 88274) extracted from naïve mice, OKT4-treated mice (12 dpi), or from infected fLX at 2 and 12 dpi. Scale bar = 100 μM **(D^{a-g})** Multiplex immunohistochemistry of fLX tissue sections extracted from Isotype (a), OKT3 (b,c), OKT4 (d,e), or OKT8 (f,g)-treated BLT-L mice at 12 dpi. CD163: teal, CD4: yellow, CD3: orange, PPARγ: green, Dapi: gray. Viral titer (PFU/mg) of the respective fLX is inlaid in white. Scale bar = 100 μM **(E-H)** Quantification of selected cytokines (CCL2, CCL3, CXCL10 and CCL19) in the serum of naïve and infected BLT-L mice (2 dpi or 12 dpi), and of OKT4-treated BLT-L mice (12 dpi). **(I)** Histopathological scoring of fLX extracted from isotype, OKT3, OKT4, or OKT8-treated BLT-L mice. One-*way ANOVA. p-values are indicated on graphs. Error bars indicate mean ± Standard error of the mean.*

Figure 6. The non-hematopoietic compartment of BLT-L fLX respond to infection and promote immune recruitment and tissue repair. (A) UMAP plot sub-clustering of the human mesenchymal compartment of fLX across all time points. **(B)** Temporal annotation of the mesenchymal sub-clusters on the UMAP plot: naïve (black), 2 dpi (red), 12 dpi (blue). **(C)** Frequency of each mesenchymal sub-cluster within the entire mesenchymal compartment over

the course of infection. Each cluster ID is labelled as in Panel A. **(D)** Heatmap displaying the

relative expression of top differentially regulated genes over the course of infection within the

activated fibroblast sub-cluster (Cluster 3). **(E)** Go-term analysis showing the most highly

upregulated signaling pathways in the activated fibroblast sub-cluster (Cluster 3). **(F-G)** UMAP

plot showing the expression of *CXCL10* **(F)** and *TIMP1* **(G)** in the mesenchymal compartment and

across all time points. **(H)** UMAP plot sub-clustering of the human endothelial compartment of fLX

across all time points. **(I)** Temporal annotation of the endothelial sub-clusters on the UMAP plot:

naïve (black), 2 dpi (red), 12 dpi (blue). **(J)** List of upregulated (red) and downregulated (blue)

genes specific to the activated endothelial sub-cluster (Cluster 2) over the other endothelial sub-

clusters (*p.adj*≤0.05). **(K)** Heatmap displaying the relative expression of the top differentially

regulated genes over the course of infection within the venous endothelial sub-cluster (Cluster 0).

Figure 7. Immunological events defining the successful resolution of SARS-CoV-2 infection in human lung tissues.

SUPPLEMENTAL FIGURE LEGENDS

Figure S1. BLT-L mice are susceptible to SARS-CoV-2 infection and display with transcriptomic changes at 2 and 12 dpi. (A-D) Transmission electron microscopy (TEM) of fLX

tissue sections extracted from BLT-L mice at 2 dpi illustrating virus particles in AT2 cells as evident

by presence of lamellar bodies. Scale bars indicated on image. **(E)** Neutralizing efficacy (ND50

values) of serum extracted from naïve or infected mice BLT-L mice (2 and 12 dpi). Assay was

performed using VeroE6 cells and a recombinant WA-1 SARS-CoV-2 virus expressing NanoLuc.

(F) Titration of the neutralizing activity of an anti-RBD antibody (serving as positive control)

against WA-1 SARS-CoV-2 virus expressing NanoLuc. **(G-I)** SARS-Cov-2 attributable

histopathology in BLT-L fLX at 2 dpi. **(G)** Bronchiolar attenuation and denuding (blue arrows)

correlates directly with the presence of SARS-CoV-2 Spike protein. **(H)** Alveolar spaces are

multifocally filled with necrotic debris admixed with neutrophils and edema (blue hashes). Type II

pneumocytes (ATII) are often denuded or attenuated in areas of inflammation that correlates

directly with the presence of SARS-CoV-2 Spike protein. Fibrin thrombi are routinely observed in

neighboring parenchyma (black asterisks). **(I)** Coagulative necrosis as evidenced by loss of

cellular detail and generalized eosinophilia (above the black hashed line) with numerous regional

fibrin thrombi (black asterisks). Although SARS-CoV-2 Spike protein is not observed in the area

of coagulative necrosis viral antigen is located within adjacent tissue. **(J-N)** Bulk RNA-sequencing

analysis was performed on total RNA extracted from fLX of naïve (n=4) or infected BLT-L mice at

2 (n=4) and 12 dpi (n=3). **(J)** Sequencing reads were aligned to a combined mouse (white), human

(black) and SARS-CoV-2 viral (red) genome. Samples containing greater than 60% human reads

(red dotted line) were included for analysis. **(K)** PCA plot showing clustering of naïve (black), 2-

(green) and 12 dpi (purple) samples. **(L)** Significantly differentially expressed genes (*p.adj*≤0.05;

upregulated: red, downregulated: blue) at 2- and 12 dpi. **(M)** Venn diagram showing the overlap

of upregulated (left) and downregulated (right) genes between 2- and 12 dpi. **(N)** Volcano plot

showing significantly downregulated (blue) and upregulated (red) genes at 2- and 12 dpi.

1076

Figure S2. Single-cell RNA sequencing analysis of fLX upon SARS-CoV-2 infection. (A-E)

Single cell RNA sequencing was performed on naïve fLX and infected fLX (at 2- and 12 dpi).

Sequencing reads were aligned to a combined human, mouse and SARS-CoV-2 viral genome.

(A) UMAP plot clustering on all cell (human and mouse) populations detected. **(B)** Expression of

mouse (top) and human *GAPDH* (bottom) in all clusters. **(C)** A human score was applied to each

cluster to identify human clusters (above dotted line) and mouse clusters (below dotted line; gray

zone). Mouse clusters were removed from downstream analysis. **(D)** Cell scoring system applied

to human cell clusters to classify each cluster. **(E)** Flow cytometric analysis showing frequency of

hCD45+, hCD3+, hCD4+, and hCD8+ cells among PBMCs extracted from the blood of naïve and

infected BLT-L mice at 2-, 6-, and 12 dpi. **(F)** UMAP clustering of human cells separated by time

point (naïve: left, 2 dpi: center, and 12 dpi: right).

1088

1089   **Figure S3. Gene signature of macrophage-like T cells. (A)** UMAP showing viral RNA reads

1090   across all human clusters. The mesenchymal cluster is enlarged. **(B)** Mitochondrial gene

1091   expression among the different T cell sub-clusters. **(C)** Expression of T-cell associated genes

1092   among the different T-cell sub-clusters. **(D)** Expression of myeloid/macrophage-associated genes

1093   among the different T-cell sub-clusters.

1094

1095   **Figure S4. Gene signature of myeloid and inflammatory monocytes sub-clusters. (A-C)**

1096   Violin plot displaying monocyte **(A)**, macrophage **(B)** and dendritic cell **(C)** gene signature score

1097   for each myeloid sub-cluster. **(D)** UMAP plots representing the expression of several myeloid

1098   markers that were employed to further define the identity of the different myeloid sub-clusters.

1099

Figure S5. Hematopoietic subset depletion in BLT-L mice and myeloid chemotaxis in

infected fLX. (A) Flow cytometric analysis showing the frequency of hCD45+, hCD3+, hCD3+

hCD4+, and hCD3+ hCD8+ cells among PBMCs extracted from the blood of BLT-L mice treated

with an isotype antibody, OKT3, OKT4 or OKT8 antibody. *One-way ANOVA. P-value indicated*

on graph. Error bars represent mean ± SEM. **(B-C)** UMAP plots showing the

compartmentalization of *CCL2* **(B)** and *CCL3* **(C)** expression within all human cell populations in

naïve or infected (2 and 12 dpi) fLX. **(D)** List of additional downregulated genes specific to the

activated endothelial sub-cluster (Cluster 2) over the other endothelial sub-clusters (*p.adj*≤0.05).

(E,F) UMAP plot showing the expression of *INSR* **(E)** and *CXCR4* **(F)** among all human

endothelial sub-clusters. **(H)** Viral titer (log(PFU/mg)) of CD3 depleted (OKT3) and CD4 depleted

(OKT4) fLX that were positive for infection. *Unpaired, non-parametric T-test. P-value indicated on*

graph.

SUPPLEMENTAL MATERIALS

Table S1. SARS-CoV-2 mutations identified among viral reads isolated from fLX at 2 dpi and from

our concentrated viral stock (i.e., WA1 positive stock used to inoculate fLX), as compared to the

Wuhan-1 consensus sequence.

Data S1. List of up- and down-regulated human genes in SARS-CoV-2-infected fLX at 2 and 12

dpi using bulk RNA sequencing analysis.

REFERENCES

Abe, T., T. Ikeda, Y. Tokuda, J. Ito, Y. Suzuki, C. Narahara, H. Iriyama, and K. Sato. 2021. A patient infected with SARS-CoV-2 over 100 days. *QJM*. 114:47-49.

Arshad, N., M. Laurent-Rolle, W.S. Ahmed, J.C. Hsu, S.M. Mitchell, J. Pawlak, D. Sengupta, K.H. Biswas, and P. Cresswell. 2023. SARS-CoV-2 accessory proteins ORF7a and ORF3a use distinct mechanisms to down-regulate MHC-I surface expression. *Proc Natl Acad Sci U S A*. 120:e2208525120.

Augusto, D.G., L.D. Murdolo, D.S.M. Chatzileontiadou, J.J. Sabatino, Jr., T. Yusufali, N.D. Peyser, X. Butcher, K. Kizer, K. Guthrie, V.W. Murray, V. Pae, S. Sarvadhavabhatla, F. Beltran, G.S. Gill, K.L. Lynch, C. Yun, C.T. Maguire, M.J. Peluso, R. Hoh, T.J. Henrich, S.G. Deeks, M. Davidson, S. Lu, S.A. Goldberg, J.D. Kelly, J.N. Martin, C.A. Vierra-Green, S.R. Spellman, D.J. Langton, M.J. Dewar-Oldis, C. Smith, P.J. Barnard, S. Lee, G.M. Marcus, J.E. Olgin, M.J. Pletcher, M. Maiers, S. Gras, and J.A. Hollenbach. 2023. A common allele of HLA is associated with asymptomatic SARS-CoV-2 infection. *Nature*. 620:128-136.

Boumaza, A., L. Gay, S. Mezouar, E. Bestion, A.B. Diallo, M. Michel, B. Desnues, D. Raoult, B. La Scola, P. Halfon, J. Vitte, D. Olive, and J.-L. Mege. 2021a. Monocytes and macrophages, targets of SARS-CoV-2: the clue for Covid-19 immunoparalysis. *The Journal of Infectious Diseases*.

Boumaza, A., L. Gay, S. Mezouar, E. Bestion, A.B. Diallo, M. Michel, B. Desnues, D. Raoult, B. La Scola, P. Halfon, J. Vitte, D. Olive, and J.L. Mege. 2021b. Monocytes and Macrophages, Targets of Severe Acute Respiratory Syndrome Coronavirus 2: The Clue for Coronavirus Disease 2019 Immunoparalysis. *J Infect Dis*. 224:395-406.

Chen, D.Y., N. Khan, B.J. Close, R.K. Goel, B. Blum, A.H. Tavares, D. Kenney, H.L. Conway, J.K. Ewoldt, V.C. Chitalia, N.A. Crossland, C.S. Chen, D.N. Kotton, S.C. Baker, S.Y. Fuchs, J.H. Connor, F. Douam, A. Emili, and M. Saeed. 2021. SARS-CoV-2 disrupts proximal elements in the JAK-STAT pathway. *J Virol*:JVI0086221.

Clayton, S.A., K.K. Daley, L. MacDonald, E. Fernandez-Vizarra, G. Bottegoni, J.D. O'Neil, T. Major, D. Griffin, Q. Zhuang, A.B. Adewoye, K. Woolcock, S.W. Jones, C. Goodyear, A. Elmesmari, A. Filer, D.A. Tennant, S. Alivernini, C.D. Buckley, R.D.S. Pitceathly, M. Kurowska-Stolarska, and A.R. Clark. 2021. Inflammation causes remodeling of mitochondrial cytochrome c oxidase mediated by the bifunctional gene C15orf48. *Sci Adv*. 7:eabl5182.

Damas, J.K., L. Landro, B. Fevang, L. Heggelund, G.E. Tjonnfjord, Y. Floisand, B. Halvorsen, S.S. Froland, and P. Aukrust. 2009. Homeostatic chemokines CCL19 and CCL21 promote

inflammation in human immunodeficiency virus-infected patients with ongoing viral replication. *Clin Exp Immunol*. 157:400-407.

Delorey, T.M., C.G.K. Ziegler, G. Heimberg, R. Normand, Y. Yang, A. Segerstolpe, D. Abbondanza, S.J. Fleming, A. Subramanian, D.T. Montoro, K.A. Jagadeesh, K.K. Dey, P. Sen, M. Slyper, Y.H. Pita-Juarez, D. Phillips, J. Biermann, Z. Bloom-Ackermann, N. Barkas, A. Ganna, J. Gomez, J.C. Melms, I. Katsyv, E. Normandin, P. Naderi, Y.V. Popov, S.S. Raju, S. Niezen, L.T. Tsai, K.J. Siddle, M. Sud, V.M. Tran, S.K. Vellarikkal, Y. Wang, L. Amir-Zilberstein, D.S. Atri, J. Beechem, O.R. Brook, J. Chen, P. Divakar, P. Dorceus, J.M. Engreitz, A. Essene, D.M. Fitzgerald, R. Fropf, S. Gazal, J. Gould, J. Grzyb, T. Harvey, J. Hecht, T. Hether, J. Jane-Valbuena, M. Leney-Greene, H. Ma, C. McCabe, D.E. McLoughlin, E.M. Miller, C. Muus, M. Niemi, R. Padera, L. Pan, D. Pant, C. Pe'er, J. Pfiffner-Borges, C.J. Pinto, J. Plaisted, J. Reeves, M. Ross, M. Rudy, E.H. Rueckert, M. Siciliano, A. Sturm, E. Todres, A. Waghray, S. Warren, S. Zhang, D.R. Zollinger, L. Cosimi, R.M. Gupta, N. Hacohen, H. Hibshoosh, W. Hide, A.L. Price, J. Rajagopal, P.R. Tata, S. Riedel, G. Szabo, T.L. Tickle, P.T. Ellinor, D. Hung, P.C. Sabeti, R. Novak, R. Rogers, D.E. Ingber, Z.G. Jiang, D. Juric, M. Babadi, S.L. Farhi, B. Izar, J.R. Stone, et al. 2021a. COVID-19 tissue atlases reveal SARS-CoV-2 pathology and cellular targets. *Nature*.

Delorey, T.M., C.G.K. Ziegler, G. Heimberg, R. Normand, Y. Yang, A. Segerstolpe, D. Abbondanza, S.J. Fleming, A. Subramanian, D.T. Montoro, K.A. Jagadeesh, K.K. Dey, P. Sen, M. Slyper, Y.H. Pita-Juárez, D. Phillips, Z. Bloom-Ackerman, N. Barkas, A. Ganna, J. Gomez, E. Normandin, P. Naderi, Y.V. Popov, S.S. Raju, S. Niezen, L.T.Y. Tsai, K.J. Siddle, M. Sud, V.M. Tran, S.K. Vellarikkal, L. Amir-Zilberstein, D.S. Atri, J. Beechem, O.R. Brook, J. Chen, P. Divakar, P. Dorceus, J.M. Engreitz, A. Essene, D.M. Fitzgerald, R. Fropf, S. Gazal, J. Gould, J. Grzyb, T. Harvey, J. Hecht, T. Hether, J. Jane-Valbuena, M. Leney-Greene, H. Ma, C. McCabe, D.E. McLoughlin, E.M. Miller, C. Muus, M. Niemi, R. Padera, L. Pan, D. Pant, C. Pe'er, J. Pfiffner-Borges, C.J. Pinto, J. Plaisted, J. Reeves, M. Ross, M. Rudy, E.H. Rueckert, M. Siciliano, A. Sturm, E. Todres, A. Waghray, S. Warren, S. Zhang, D.R. Zollinger, L. Cosimi, R.M. Gupta, N. Hacohen, W. Hide, A.L. Price, J. Rajagopal, P.R. Tata, S. Riedel, G. Szabo, T.L. Tickle, D. Hung, P.C. Sabeti, R. Novak, R. Rogers, D.E. Ingber, Z. Gordon Jiang, D. Juric, M. Babadi, S.L. Farhi, J.R. Stone, I.S. Vlachos, I.H. Solomon, O. Ashenberg, C.B.M. Porter, B. Li, A.K. Shalek, A.-C. Villani, et al. 2021b. A single-cell and spatial atlas of autopsy tissues reveals pathology and cellular targets of SARS-CoV-2. *bioRxiv*:2021.2002.2025.430130.

Di, Y., J. Lew, U. Goncin, A. Radomska, S.S. Rout, B.E.T. Gray, S. Machtaler, D. Falzarano, and K.J. Lavender. 2022. SARS-CoV-2 Variant-Specific Infectivity and Immune Profiles Are Detectable in a Humanized Lung Mouse Model. *Viruses*. 14.

Dobin, A., C.A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, S. Jha, P. Batut, M. Chaisson, and T.R. Gingeras. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*. 29:15-21.

Edahiro, R., Y. Shirai, Y. Takeshima, S. Sakakibara, Y. Yamaguchi, T. Murakami, T. Morita, Y. Kato, Y.C. Liu, D. Motooka, Y. Naito, A. Takuwa, F. Sugihara, K. Tanaka, J.B. Wing, K. Sonehara, Y. Tomofuji, C.-T.F. Japan, H. Namkoong, H. Tanaka, H. Lee, K. Fukunaga, H. Hirata, Y. Takeda, D. Okuzaki, A. Kumanogoh, and Y. Okada. 2023. Single-cell analyses and host genetics highlight the role of innate immune cells in COVID-19 severity. *Nat Genet*. 55:753-767.

Evren, E., E. Ringqvist, K.P. Tripathi, N. Sleiers, I.C. Rives, A. Alisjahbana, Y. Gao, D. Sarhan, T. Halle, C. Sorini, R. Lepzien, N. Marquardt, J. Michaelsson, A. Smed-Sorensen, J. Botling, M.C.I. Karlsson, E.J. Villablanca, and T. Willinger. 2021. Distinct developmental pathways from blood monocytes generate human lung macrophage diversity. *Immunity*. 54:259-275 e257.

Fagre, A., J. Lewis, M. Eckley, S. Zhan, S.M. Rocha, N.R. Sexton, B. Burke, B. Geiss, O. Peersen,
T. Bass, R. Kading, J. Rovnak, G.D. Ebel, R.B. Tjalkens, T. Aboellail, and T. Schountz.
2021. SARS-CoV-2 infection, neuropathogenesis and transmission among deer mice:
Implications for spillback to New World rodents. *PLoS Pathog*. 17:e1009585.
Fan, C., Y. Wu, X. Rui, Y. Yang, C. Ling, S. Liu, S. Liu, and Y. Wang. 2022. Animal models for
COVID-19: advances, gaps and perspectives. *Signal transduction and targeted therapy*.
7:220.
Finkel, Y., A. Gluck, A. Nachshon, R. Winkler, T. Fisher, B. Rozman, O. Mizrahi, Y. Lubelsky, B.
Zuckerman, B. Slobodin, Y. Yahalom-Ronen, H. Tamir, I. Ulitsky, T. Israely, N. Paran, M.
Schwartz, and N. Stern-Ginossar. 2021. SARS-CoV-2 uses a multipronged strategy to
impede host protein synthesis. *Nature*. 594:240-245.
Fu, L., Z. Liu, and Y. Liu. 2023. Fibrinogen-like protein 2 in inflammatory diseases: A future
therapeutic target. *Int Immunopharmacol*. 116:109799.
Grant, R.A., L. Morales-Nebreda, N.S. Markov, S. Swaminathan, M. Querrey, E.R. Guzman, D.A.
Abbott, H.K. Donnelly, A. Donayre, I.A. Goldberg, Z.M. Klug, N. Borkowski, Z. Lu, H.
Kihshen, Y. Politanska, L. Sichizya, M. Kang, A. Shilatifard, C. Qi, J.W. Lomasney, A.C.
Argento, J.M. Kruser, E.S. Malsin, C.O. Pickens, S.B. Smith, J.M. Walter, A.E. Pawlowski,
D. Schneider, P. Nannapaneni, H. Abdala-Valencia, A. Bharat, C.J. Gottardi, G.R.S.
Budinger, A.V. Misharin, B.D. Singer, R.G. Wunderink, R.A. Grant, L. Morales-Nebreda,
N.S. Markov, S. Swaminathan, M. Querrey, E.R. Guzman, D.A. Abbott, H.K. Donnelly, A.
Donayre, I.A. Goldberg, Z.M. Klug, N. Borkowski, Z. Lu, H. Kihshen, Y. Politanska, L.
Sichizya, M. Kang, A. Shilatifard, C. Qi, J.W. Lomasney, A.C. Argento, J.M. Kruser, E.S.
Malsin, C.O. Pickens, S.B. Smith, J.M. Walter, A.E. Pawlowski, D. Schneider, P.
Nannapaneni, H. Abdala-Valencia, A. Bharat, C.J. Gottardi, G.R.S. Budinger, A.V.
Misharin, B.D. Singer, R.G. Wunderink, A.A. Wagh, A.R. Hauser, A.R. Wolfe, A. Thakrar,
A.V. Yeldandi, A.A. Wang, A.R. Levenson, A.M. Joudi, B. Tran, C.A. Gao, C. Kurihara,
C.J. Schroedl, C.M. Horvath, D. Meza, D.D. Odell, D.W. Kamp, D.R. Winter, E.A. Ozer,
E.D. Shanes, E.T. Bartom, E.J. Rendleman, E.M. Leibenguth, F. Wehbe, G.Y. Liu, G.T.
Gadhvi, H.T. Navarro, J.I. Sznajder, J.E. Dematte, et al. 2021. Circuits between infected
macrophages and T cells in SARS-CoV-2 pneumonia. *Nature*. 590:635-641.
Gubernatorova, E.O., E.A. Gorshkova, A.I. Polinova, and M.S. Drutskaya. 2020. IL-6: Relevance
for immunopathology of SARS-CoV-2. *Cytokine Growth Factor Rev*. 53:13-24.
Gupta, S.K., P.G. Lysko, K. Pillarisetti, E. Ohlstein, and J.M. Stadel. 1998. Chemokine receptors
in human endothelial cells. Functional expression of CXCR4 and its transcriptional
regulation by inflammatory cytokines. *J Biol Chem*. 273:4282-4287.
Hao, Y., S. Hao, E. Andersen-Nissen, W.M. Mauck, 3rd, S. Zheng, A. Butler, M.J. Lee, A.J. Wilk,
C. Darby, M. Zager, P. Hoffman, M. Stoeckius, E. Papalexi, E.P. Mimitou, J. Jain, A.
Srivastava, T. Stuart, L.M. Fleming, B. Yeung, A.J. Rogers, J.M. McElrath, C.A. Blish, R.
Gottardo, P. Smibert, and R. Satija. 2021. Integrated analysis of multimodal single-cell
data. *Cell*. 184:3573-3587 e3529.
Hettle, D., S. Hutchings, P. Muir, E. Moran, and C.-G.U. consortium. 2022. Persistent SARS-CoV-
2 infection in immunocompromised patients facilitates rapid viral evolution: Retrospective
cohort study and literature review. *Clin Infect Pract*. 16:100210.
Jin, K., E.E. Bardes, A. Mitelpunkt, J.Y. Wang, S. Bhatnagar, S. Sengupta, D.P. Krummel, M.E.
Rothenberg, and B.J. Aronow. 2021. Implicating Gene and Cell Networks Responsible for
Differential COVID-19 Host Responses via an Interactive Single Cell Web Portal.
bioRxiv:2021.2006.2007.447287.
Junqueira, C., A. Crespo, S. Ranjbar, L.B. de Lacerda, M. Lewandrowski, J. Ingber, B. Parry, S.
Ravid, S. Clark, M.R. Schrimpf, F. Ho, C. Beakes, J. Margolin, N. Russell, K. Kays, J.
Boucau, U. Das Adhikari, S.M. Vora, V. Leger, L. Gehrke, L.A. Henderson, E. Janssen,
D. Kwon, C. Sander, J. Abraham, M.B. Goldberg, H. Wu, G. Mehta, S. Bell, A.E. Goldfeld,

M.R. Filbin, and J. Lieberman. 2022. FcgammaR-mediated SARS-CoV-2 infection of
monocytes activates inflammation. *Nature*. 606:576-584.
Kadiyska, T., I. Tourtourikov, K. Dabchev, R. Cherneva, N. Stoynev, R. Hadjiolova, V. Mitev, D.A.
Spandidos, M. Adamaki, and V. Zoumpourlis. 2022. Role of endothelial dysfunction in the
severity of COVID-19 infection (Review). *Molecular medicine reports*. 26.
Kang, J.G., M.J. Amar, A.T. Remaley, J. Kwon, P.J. Blackshear, P.Y. Wang, and P.M. Hwang.
2011. Zinc finger protein tristetraprolin interacts with CCL3 mRNA and regulates tissue
inflammation. *J Immunol*. 187:2696-2701.
Kenney, D.J., A.K. O'Connell, J. Turcinovic, P. Montanaro, R.M. Hekman, T. Tamura, A.R.
Berneshawi, T.R. Cafiero, S. Al Abdullatif, B. Blum, S.I. Goldstein, B.L. Heller, H.P. Gertje,
E. Bullitt, A.J. Trachtenberg, E. Chavez, E.T. Nono, C. Morrison, A.E. Tseng, A. Sheikh,
S. Kurnick, K. Grosz, M. Bosmann, M. Ericsson, B.R. Huber, M. Saeed, A.B. Balazs, K.P.
Francis, A. Klose, N. Paragas, J.D. Campbell, J.H. Connor, A. Emili, N.A. Crossland, A.
Ploss, and F. Douam. 2022. Humanized mice reveal a macrophage-enriched gene
signature defining human lung tissue protection during SARS-CoV-2 infection. *Cell Rep*.
39:110714.
Kohout, T.A., S.L. Nicholas, S.J. Perry, G. Reinhart, S. Junger, and R.S. Struthers. 2004.
Differential desensitization, receptor phosphorylation, beta-arrestin recruitment, and
ERK1/2 activation by the two endogenous ligands for the CC chemokine receptor 7. *J Biol*
Chem. 279:23214-23222.
Kosyreva, A., D. Dzhalilova, A. Lokhonina, P. Vishnyakova, and T. Fatkhudinov. 2021. The Role
of Macrophages in the Pathogenesis of SARS-CoV-2-Associated Acute Respiratory
Distress Syndrome. *Frontiers in Immunology*. 12.
Lebedev, M., B. Faburay, J.A. Richt, and A. Young. 2021. Myeloid-like gammadelta T cell subset
in the immune response to an experimental Rift Valley fever vaccine in sheep. *Vet*
Immunol Immunopathol. 233:110184.
Leroyer, A.S., M.G. Blin, R. Bachelier, N. Bardin, M. Blot-Chabaud, and F. Dignat-George. 2019.
CD146 (Cluster of Differentiation 146). *Arterioscler Thromb Vasc Biol*. 39:1026-1033.
Liao, Y., G.K. Smyth, and W. Shi. 2014. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics*. 30:923-930.
Love, M.I., W. Huber, and S. Anders. 2014. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome biology*. 15:550.
Machkovech, H.M., A.M. Hahn, J. Garonzik Wang, N.D. Grubaugh, P.J. Halfmann, M.C. Johnson,
J.E. Lemieux, D.H. O'Connor, A. Piantadosi, W. Wei, and T.C. Friedrich. 2024. Persistent
SARS-CoV-2 infection: significance and implications. *Lancet Infect Dis*.
Mao, G., Y. Kang, R. Lin, S. Hu, Z. Zhang, H. Li, W. Liao, and Z. Zhang. 2019. Long Non-coding
RNA HOTTIP Promotes CCL3 Expression and Induces Cartilage Degradation by
Sponging miR-455-3p. *Front Cell Dev Biol*. 7:161.
Maponga, T.G., M. Jeffries, H. Tegally, A. Sutherland, E. Wilkinson, R.J. Lessells, N. Msomi, G.
van Zyl, T. de Oliveira, and W. Preiser. 2023. Persistent Severe Acute Respiratory
Syndrome Coronavirus 2 Infection With accumulation of mutations in a patient with poorly
controlled Human Immunodeficiency Virus infection. *Clin Infect Dis*. 76:e522-e525.
Molgora, M., and M. Colonna. 2022. Innate-like T cells: A promising asset in anti-cancer immunity.
Cancer Cell. 40:714-716.
Moriyama, M., C. Lucas, V.S. Monteiro, S.-C.-G.S.I. Yale, and A. Iwasaki. 2023. Enhanced
inhibition of MHC-I expression by SARS-CoV-2 Omicron subvariants. *Proc Natl Acad Sci*
U S A. 120:e2221652120.
Munnur, D., Q. Teo, D. Eggermont, H.H.Y. Lee, F. Thery, J. Ho, S.W. van Leur, W.W.S. Ng, L.Y.L.
Siu, A. Beling, H. Ploegh, A. Pinto-Fernandez, A. Damianou, B. Kessler, F. Impens, C.K.P.
Mok, and S. Sanyal. 2021. Altered ISGylation drives aberrant macrophage-dependent
immune responses during SARS-CoV-2 infection. *Nat Immunol*. 22:1416-1427.

Muto, M., M. Baghdadi, R. Maekawa, H. Wada, and K. Seino. 2015. Myeloid molecular
characteristics of human gammadelta T cells support their acquisition of tumor antigen-
presenting capacity. *Cancer Immunol Immunother*. 64:941-949.
Otifi, H.M., and B.K. Adiga. 2022. Endothelial Dysfunction in Covid-19 Infection. *Am J Med Sci*.
363:281-287.
Payankaulam, S., A.M. Raicu, and D.N. Arnosti. 2019. Transcriptional Regulation of INSR, the
Insulin Receptor Gene. *Genes (Basel)*. 10.
PMDA. 2021. Xevudy for Intravenous Infusion 500 mg_GlaxoSmithKline K.K._Report on Special
Approval for Emergency; https://www.pmda.go.jp/files/000245006.pdf.
Qian, Y., T. Lei, P.S. Patel, C.H. Lee, P. Monaghan-Nichols, H.B. Xin, J. Qiu, and M. Fu. 2021.
Direct Activation of Endothelial Cells by SARS-CoV-2 Nucleocapsid Protein Is Blocked by
Simvastatin. *J Virol*. 95:e0139621.
Qin, Z., F. Liu, R. Blair, C. Wang, H. Yang, J. Mudd, J.M. Currey, N. Iwanaga, J. He, R. Mi, K.
Han, C.C. Midkiff, M.A. Alam, B.H. Aktas, R.S.V. Heide, R. Veazey, G. Piedimonte, N.J.
Maness, S. Ergun, F. Mauvais-Jarvis, J. Rappaport, J.K. Kolls, and X. Qin. 2021.
Endothelial cell infection and dysfunction, immune activation in severe COVID-19.
Theranostics. 11:8076-8091.
Rathjen, T., B. Kunkemoeller, C.T. Cederquist, X. Wang, S.M. Lockhart, J.C. Patti, H. Willenbrock,
G.S. Olsen, G.K. Povlsen, H.C. Beck, L.M. Rasmussen, Q. Li, K. Park, G.L. King, and C.
Rask-Madsen. 2022. Endothelial Cell Insulin Signaling Regulates CXCR4 (C-X-C Motif
Chemokine Receptor 4) and Limits Leukocyte Adhesion to Endothelium. *Arterioscler*
Thromb Vasc Biol. 42:e217-e227.
Ren, X., W. Wen, X. Fan, W. Hou, B. Su, P. Cai, J. Li, Y. Liu, F. Tang, F. Zhang, Y. Yang, J. He,
W. Ma, J. He, P. Wang, Q. Cao, F. Chen, Y. Chen, X. Cheng, G. Deng, X. Deng, W. Ding,
Y. Feng, R. Gan, C. Guo, W. Guo, S. He, C. Jiang, J. Liang, Y.M. Li, J. Lin, Y. Ling, H.
Liu, J. Liu, N. Liu, S.Q. Liu, M. Luo, Q. Ma, Q. Song, W. Sun, G. Wang, F. Wang, Y. Wang,
X. Wen, Q. Wu, G. Xu, X. Xie, X. Xiong, X. Xing, H. Xu, C. Yin, D. Yu, K. Yu, J. Yuan, B.
Zhang, P. Zhang, T. Zhang, J. Zhao, P. Zhao, J. Zhou, W. Zhou, S. Zhong, X. Zhong, S.
Zhang, L. Zhu, P. Zhu, B. Zou, J. Zou, Z. Zuo, F. Bai, X. Huang, P. Zhou, Q. Jiang, Z.
Huang, J.X. Bei, L. Wei, X.W. Bian, X. Liu, T. Cheng, X. Li, P. Zhao, F.S. Wang, H. Wang,
B. Su, Z. Zhang, K. Qu, X. Wang, J. Chen, R. Jin, and Z. Zhang. 2021. COVID-19 immune
features revealed by a large-scale single-cell transcriptome atlas. *Cell*. 184:1895-1913
e1819.
Rendeiro, A.F., H. Ravichandran, Y. Bram, V. Chandar, J. Kim, C. Meydan, J. Park, J. Foox, T.
Hether, S. Warren, Y. Kim, J. Reeves, S. Salvatore, C.E. Mason, E.C. Swanson, A.C.
Borczuk, O. Elemento, and R.E. Schwartz. 2021. The spatial landscape of lung pathology
during COVID-19 progression. *Nature*. 593:564-569.
Schultze, J.L., and A.C. Aschenbrenner. 2021. COVID-19 and the human innate immune system.
Cell. 184:1671-1692.
Schyns, J., Q. Bai, C. Ruscitti, C. Radermecker, S. De Schepper, S. Chakarov, F. Farnir, D.
Pirottin, F. Ginhoux, G. Boeckxstaens, F. Bureau, and T. Marichal. 2019. Non-classical
tissue monocytes and two functionally distinct populations of interstitial macrophages
populate the mouse lung. *Nat Commun*. 10:3964.
Sefik, E., R. Qu, C. Junqueira, E. Kaffe, H. Mirza, J. Zhao, J.R. Brewer, A. Han, H.R. Steach, B.
Israelow, H.N. Blackburn, S.E. Velazquez, Y.G. Chen, S. Halene, A. Iwasaki, E. Meffre,
M. Nussenzweig, J. Lieberman, C.B. Wilen, Y. Kluger, and R.A. Flavell. 2022.
Inflammasome activation in infected macrophages drives COVID-19 pathology. *Nature*.
606:585-593.
Sepulcri, C., C. Dentone, M. Mikulska, B. Bruzzone, A. Lai, D. Fenoglio, F. Bozzano, A. Bergna,
A. Parodi, T. Altosole, E. Delfino, G. Bartalucci, A. Orsi, A. Di Biagio, G. Zehender, F.
Ballerini, S. Bonora, A. Sette, R. De Palma, G. Silvestri, A. De Maria, and M. Bassetti.

2021. The Longest Persistence of Viable SARS-CoV-2 With Recurrence of Viremia and
Relapsing Symptomatic COVID-19 in an Immunocompromised Patient-A Case Study.
Open Forum Infect Dis. 8:ofab217.
Serrano-Lopez, J., S. Hegde, S. Kumar, J. Serrano, J. Fang, A.M. Wellendorf, P.A. Roche, Y.
Rangel, L.J. Carrington, H. Geiger, H.L. Grimes, S. Luther, I. Maillard, J. Sanchez-Garcia,
D.T. Starczynowski, and J.A. Cancelas. 2021. Inflammation rapidly recruits mammalian
GMP and MDP from bone marrow into regional lymphatics. *Elife*. 10.
Shrestha, L.B., C. Foster, W. Rawlinson, N. Tedla, and R.A. Bull. 2022. Evolution of the SARS-
CoV-2 omicron variants BA.1 to BA.5: Implications for immune escape and transmission.
Rev Med Virol. 32:e2381.
Stephenson, E., G. Reynolds, R.A. Botting, F.J. Calero-Nieto, M.D. Morgan, Z.K. Tuong, K. Bach,
W. Sungnak, K.B. Worlock, M. Yoshida, N. Kumasaka, K. Kania, J. Engelbert, B. Olabi,
J.S. Spegarova, N.K. Wilson, N. Mende, L. Jardine, L.C.S. Gardner, I. Goh, D. Horsfall, J.
McGrath, S. Webb, M.W. Mather, R.G.H. Lindeboom, E. Dann, N. Huang, K. Polanski, E.
Prigmore, F. Gothe, J. Scott, R.P. Payne, K.F. Baker, A.T. Hanrath, I.C.D. Schim van der
Loeff, A.S. Barr, A. Sanchez-Gonzalez, L. Bergamaschi, F. Mescia, J.L. Barnes, E. Kilich,
A. de Wilton, A. Saigal, A. Saleh, S.M. Janes, C.M. Smith, N. Gopee, C. Wilson, P.
Coupland, J.M. Coxhead, V.Y. Kiselev, S. van Dongen, J. Bacardit, H.W. King, I.
Cambridge Institute of Therapeutic, C.-B.C. Infectious Disease-National Institute of Health
Research, A.J. Rostron, A.J. Simpson, S. Hambleton, E. Laurenti, P.A. Lyons, K.B. Meyer,
M.Z. Nikolic, C.J.A. Duncan, K.G.C. Smith, S.A. Teichmann, M.R. Clatworthy, J.C.
Marioni, B. Gottgens, and M. Haniffa. 2021. Single-cell multi-omics analysis of the immune
response in COVID-19. *Nat Med*. 27:904-916.
Sun, R., Z. Zhao, C. Fu, Y. Wang, Z. Guo, C. Zhang, L. Liu, C. Zhang, C. Shu, J. He, Y.G. Yang,
S. Hua, Y. Gao, and Z. Hu. 2022a. Humanized mice for investigating SARS-CoV-2 lung
infection and associated human immune responses. *Eur J Immunol*. 52:1640-1647.
Sun, X., A.K. Perl, R. Li, S.M. Bell, E. Sajti, V.V. Kalinichenko, T.V. Kalin, R.S. Misra, H.
Deshmukh, G. Clair, J. Kyle, L.E. Crotty Alexander, J.A. Masso-Silva, J.A. Kitzmiller, K.A.
Wikenheiser-Brokamp, G. Deutsch, M. Guo, Y. Du, M.P. Morley, M.J. Valdez, H.V. Yu, K.
Jin, E.E. Bardes, J.A. Zepp, T. Neithamer, M.C. Basil, W.J. Zacharias, J. Verheyden, R.
Young, G. Bandyopadhyay, S. Lin, C. Ansong, J. Adkins, N. Salomonis, B.J. Aronow, Y.
Xu, G. Pryhuber, J. Whitsett, E.E. Morrisey, and N.L. Consortium. 2022b. A census of the
lung: CellCards from LungMAP. *Dev Cell*. 57:112-145 e112.
Tveita, A., S.L. Murphy, J.C. Holter, A.B. Kildal, A.E. Michelsen, T.V. Lerum, M. Kaarbo, L.
Heggelund, A.R. Holten, A.K. Finbraten, K.E. Muller, A. Mathiessen, S. Boe, B. Fevang,
B.K. Granerud, K. Tonby, A. Lind, S.G. Dudman, K.N. Henriksen, F. Muller, O.H.
Skjonsberg, M. Troseid, A. Barratt-Due, A.M. Dyrhol-Riise, P. Aukrust, B. Halvorsen, T.B.
Dahl, T. Ueland, N.-S. Consortium, and S.-C.-S.G.I. the Norwegian. 2022. High Circulating
Levels of the Homeostatic Chemokines CCL19 and CCL21 Predict Mortality and Disease
Severity in COVID-19. *J Infect Dis*. 226:2150-2160.
Wahl, A., C. De, M. Abad Fernandez, E.M. Lenarcic, Y. Xu, A.S. Cockrell, R.A. Cleary, C.E.
Johnson, N.J. Schramm, L.M. Rank, I.G. Newsome, H.A. Vincent, W. Sanders, C.R.
Aguilera-Sandoval, A. Boone, W.H. Hildebrand, P.A. Dayton, R.S. Baric, R.J. Pickles, M.
Braunstein, N.J. Moorman, N. Goonetilleke, and J. Victor Garcia. 2019. Precision mouse
models with expanded tropism for human pathogens. *Nat Biotechnol*. 37:1163-1173.
Wahl, A., L. Gralinski, C. Johnson, W. Yao, M. Kovarova, K. Dinnon, H. Liu, V. Madden, H.
Krzystek, C. De, K. White, A. Schafer, T. Zaman, S. Leist, P. Grant, K. Gully, F. Askin, E.
Browne, C. Jones, R. Pickles, R. Baric, and J.V. Garcia. 2020. Acute SARS-CoV-2
Infection is Highly Cytopathic, Elicits a Robust Innate Immune Response and is Efficiently
Prevented by EIDD-2801. *Res Sq*.

Wahl, A., L.E. Gralinski, C.E. Johnson, W. Yao, M. Kovarova, K.H. Dinnon, 3rd, H. Liu, V.J.
Madden, H.M. Krzystek, C. De, K.K. White, K. Gully, A. Schafer, T. Zaman, S.R. Leist,
P.O. Grant, G.R. Bluemling, A.A. Kolykhalov, M.G. Natchus, F.B. Askin, G. Painter, E.P.
Browne, C.D. Jones, R.J. Pickles, R.S. Baric, and J.V. Garcia. 2021. SARS-CoV-2
infection is effectively treated and prevented by EIDD-2801. *Nature*. 591:451-457.
Wauters, E., P. Van Mol, A.D. Garg, S. Jansen, Y. Van Herck, L. Vanderbeke, A. Bassez, B.
Boeckx, B. Malengier-Devlies, A. Timmerman, T. Van Brussel, T. Van Buyten, R.
Schepers, E. Heylen, D. Dauwe, C. Dooms, J. Gunst, G. Hermans, P. Meersseman, D.
Testelmans, J. Yserbyt, S. Tejpar, W. De Wever, P. Matthys, J. Neyts, J. Wauters, J. Qian,
and D. Lambrechts. 2021. Discriminating mild from critical COVID-19 by innate and
adaptive immune single-cell profiling of bronchoalveolar lavages. *Cell Res*. 31:272-290.
Wendisch, D., O. Dietrich, T. Mari, S. von Stillfried, I.L. Ibarra, M. Mittermaier, C. Mache, R.L.
Chua, R. Knoll, S. Timm, S. Brumhard, T. Krammer, H. Zauber, A.L. Hiller, A. Pascual-
Reguant, R. Mothes, R.D. Bulow, J. Schulze, A.M. Leipold, S. Djudjaj, F. Erhard, R.
Geffers, F. Pott, J. Kazmierski, J. Radke, P. Pergantis, K. Bassler, C. Conrad, A.C.
Aschenbrenner, B. Sawitzki, M. Landthaler, E. Wyler, D. Horst, C.-O.I. Deutsche, S.
Hippenstiel, A. Hocke, F.L. Heppner, A. Uhrig, C. Garcia, F. Machleidt, S. Herold, S.
Elezkurtaj, C. Thibeault, M. Witzenrath, C. Cochain, N. Suttorp, C. Drosten, C. Goffinet,
F. Kurth, J.L. Schultze, H. Radbruch, M. Ochs, R. Eils, H. Muller-Redetzky, A.E. Hauser,
M.D. Luecken, F.J. Theis, C. Conrad, T. Wolff, P. Boor, M. Selbach, A.E. Saliba, and L.E.
Sander. 2021. SARS-CoV-2 infection triggers profibrotic macrophage responses and lung
fibrosis. *Cell*. 184:6243-6261 e6227.
Xu, S.W., I. Ilyas, and J.P. Weng. 2023. Endothelial dysfunction in COVID-19: an overview of
evidence, biomarkers, mechanisms and potential therapies. *Acta Pharmacol Sin*. 44:695-
709.
Yan, Y., R. Chen, X. Wang, K. Hu, L. Huang, M. Lu, and Q. Hu. 2019. CCL19 and CCR7
Expression, Signaling Pathways, and Adjuvant Functions in Viral Infection and Prevention.
Front Cell Dev Biol. 7:212.
Yang, D., H. Chu, Y. Hou, Y. Chai, H. Shuai, A.C. Lee, X. Zhang, Y. Wang, B. Hu, X. Huang, T.T.
Yuen, J.P. Cai, J. Zhou, S. Yuan, A.J. Zhang, J.F. Chan, and K.Y. Yuen. 2020. Attenuated
Interferon and Proinflammatory Response in SARS-CoV-2-Infected Human Dendritic
Cells Is Associated With Viral Antagonism of STAT1 Phosphorylation. *J Infect Dis*.
222:734-745.
Yoo, J.S., M. Sasaki, S.X. Cho, Y. Kasuga, B. Zhu, R. Ouda, Y. Orba, P. de Figueiredo, H. Sawa,
and K.S. Kobayashi. 2021. SARS-CoV-2 inhibits induction of the MHC class I pathway by
targeting the STAT1-IRF1-NLRC5 axis. *Nat Commun*. 12:6602.
Zhang, Y., Y. Chen, and Z. Meng. 2020. Immunomodulation for Severe COVID-19 Pneumonia:
The State of the Art. *Frontiers in Immunology*. 11.
Zheng, J., Y. Wang, K. Li, D.K. Meyerholz, C. Allamargot, and S. Perlman. 2021. Severe Acute
Respiratory Syndrome Coronavirus 2-Induced Immune Activation and Death of Monocyte-
Derived Human Macrophages and Dendritic Cells. *J Infect Dis*. 223:785-795.

1448

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7